Correlation Functions of Classical and Quantum Artin System defined on Labachevsky Plane and Scrambling Time

Hrachya Babujian, Rubik Poghossian and George Savvidy

Institute of Nuclear and Particle Physics
Demokritos National Research Centre, Athens, Greece
Yerevan Physics Institute,
Alikhanian Br. 2, AM-0036 Yerevan, Armenia
E-mail: babujian@yerphi.am, poghos@yerphi.am,
savvidy@inp.demokritos.gr

ABSTRACT: We consider the quantisation of the Artin dynamical system defined on the fundamental region of the Lobachevsky plane. This fundamental region of the modular group has finite volume and infinite extension in the vertical axis that correspond to a cusp. In classical regime the geodesic flow in this fundamental region represents one of the most chaotic dynamical systems, has mixing of all orders, Lebesgue spectrum and non-zero Kolmogorov entropy. The classical correlation functions decay exponentially with an exponent proportional to the entropy. Here we calculated the quantum mechanical correlation functions in the approximation when the oscillation modes in the horizontal direction are neglected. The effective influence of these modes is taken into account to some extent by using the exact expression for the reflection amplitude. By performing a numerical integration we observed that a two-point correlation function decays exponentially and that a four-point function demonstrates tendency to decay with a lower pace. With the numerical data available to us it is impossible to estimate scrambling time t^* or to confirm its existence in the hyperbolic system considered above, but qualitatively we observe a short time exponential decay of the out-of-time correlation function to almost zero value and then an essential increase with the subsequent large fluctuations.

KEYWORDS: Artin billiard, Chaotic dynamical systems, Anosov systems, Kolmogorov systems, Modular invariance, Non holomorphic Automorphic functions, Quantum and Classical correlation functions, Scrambling time.

Contents

1	Introduction	1
2	Lobachevsky plane and its isometry group	4
3	Geodesic flow in Hamiltonian gauge	4
4	Quantization	6
5	Continuous spectrum and the reflection amplitude	8
6	Correlation functions	13
	6.1 Two-point correlation function	13
	6.2 Four-point correlation function	15
7	Acknowledgment	17

1 Introduction

The hyperbolic Anosov C-systems have exponential instability of their trajectories and as such represent the most natural chaotic dynamical systems [7]. Of special interest are C-systems which are defined on closed surfaces of the Lobachevsky plane of constant negative curvature. An example of such system has been introduced in a brilliant article published in 1924 by the mathematician Emil Artin [1]. The dynamical system is defined on the fundamental region of the Lobachevsky plane which is obtained by the identification of points congruent with respect to the modular group SL(2, Z), a discrete subgroup of the Lobachevsky plane isometries [2–4]. The fundamental region \mathcal{F} in this case is a hyperbolic triangle on Fig.1. The geodesic trajectories are bounded to propagate on the fundamental hyperbolic triangle. The geodesic flow in this fundamental region represents one of the most chaotic dynamical systems with exponential instability of its trajectories, has mixing of all orders, Lebesgue spectrum and non-zero Kolmogorov entropy [5–16, 18–20, 22, 23, 32–37, 40].

There is a great interest to consider quantisation of the hyperbolic dynamical systems and investigate their quantum-mechanical properties [22, 23]. Our main interest in this article is a study of the behaviour of the correlation functions of the Artin hyperbolic dynamical system in its classical and quantum regimes.

In classical regime the correlation functions are defined as an integral over a pair of functions/observables A and B in which the first one is stationary and the second one evolves with the geodesic flow g_t :

$$\mathcal{D}_t(A,B) = \int_{\mathcal{M}} A(g) \overline{B(gg_t)} d\mu(g). \tag{1.1}$$

The earlier investigation of the classical correlation functions of the geodesic flows was performed in [16–20] by using different approaches including Fourier series for the SL(2,R) group, zeta function for the geodesic flows, relating the poles of the Fourier transform of the correlation functions to the spectrum of an associated Ruelle operator, the methods of unitary representation theory, spectral properties of the corresponding Laplacian and other approaches. In recent articles [38, 39] the authors demonstrated exponential decay of the correlation functions with time on the classical phase space. The result was derived by using the differential geometry, group-theoretical methods of Gelfand and Fomin, the time evolution equations and the properties of automorphic functions on \mathcal{F} . The exponential decay rate was expressed in terms of the entropy $h(\mathcal{F})$ of the system

$$|\mathcal{D}_t(A,B)| \le M e^{-Kh|t|} , \qquad (1.2)$$

where M and K are constants depending on the smoothness of the functions. In classical regime the exponential divergency of the geodesic trajectories resulted into the universal exponential decay of its classical correlation functions [38, 39].

In order to investigate the behaviour of the correlation functions in quantum-mechanical regime it is necessary to know the spectrum of the system and the corresponding wave functions. In the case of the modular group the energy spectrum has continuous part, which is originating from the asymptotically free motion inside an infinitely long "y -channel" extended in the vertical direction of the fundamental region, as well as infinitely many discrete energy states corresponding to a bounded motion at the "bottom" of the fundamental triangle. The spectral problem has deep number-theoretical origin and was partially solved in a series of pioneering articles [24–27]. It was solved partially because the discrete spectrum and the corresponding wave functions are not known analytically. The general properties of the discrete spectrum have been derived using Selberg trace formula [26–31]. Numerical calculation of the discrete energy levels were performed for many energy states [42–44].

In the next three sections we shall review the geometry of Lobachevsky hyperbolic plane and of the fundamental region which corresponds to the modular group $SL(2, \mathbb{Z})$, the geodesic flow on that region and the quantisation of the system. The derivation of the Maass wave functions [24] for the continuous spectrum will be reviewed in details. We shall use the Poincaré representation for Maass non-holomorphic automorphic wave functions. We introduce a natural physical variable \tilde{y} for the distance in the vertical direction on the fundamental triangle

 $\int dy/y = \ln y = \tilde{y}$ and the corresponding momentum p_y in order to represent the Maass wave functions (5.22) in the form which is appealing to the physical intuition

$$\psi_{p_y}(x, \tilde{y}) = e^{-ip_y \tilde{y}} + \frac{\theta(\frac{1}{2} + ip_y)}{\theta(\frac{1}{2} - ip_y)} e^{ip_y \tilde{y}} + 4 \sum_{l=1}^{\infty} \tau_{ip_y}(l) K_{ip_y}(2\pi l e^{\tilde{y}}) \cos(2\pi l x).$$

Indeed, the first two terms describe the incoming and outgoing plane waves. The plane wave $e^{-ip_y\tilde{y}}$ incoming from infinity of the y axis on Fig. 1 (the vertex \mathcal{D}) elastically scatters on the boundary ACB of the fundamental triangle \mathcal{F} on Fig. 1. The reflection amplitude is a pure phase and is given by the expression in front of the outgoing plane wave $e^{ip_y\tilde{y}}$

$$\frac{\theta(\frac{1}{2} + ip_y)}{\theta(\frac{1}{2} - ip_y)} = \exp\left[i\,\varphi(p_y)\right]. \tag{1.3}$$

The rest of the wave function describes the standing waves $\cos(2\pi lx)$ in the x direction between boundaries $x = \pm 1/2$ with the amplitudes $K_{ip_y}(2\pi ly)$, which are exponentially decreasing with index l Fig.2. The continuous energy spectrum is given by the formula

$$E = p_y^2 + \frac{1}{4}.$$

The wave functions of the discrete spectrum have the form [24–27, 42–44]

$$\psi_n(z) = \sum_{l=1}^{\infty} c_l(n) K_{iu_n}(2\pi l e^{\tilde{y}}) \{ \frac{\cos(2\pi l x)}{\sin(2\pi l x)},$$
 (1.4)

where the spectrum $E_n = \frac{1}{4} + u_n^2$ and the coefficients $c_l(n)$ are not known analytically, but have been computed numerically for many values of n [42–44]. Having explicit expressions of the wave functions one can analyse a quantum-mechanical behaviour of the correlation functions [36], which we investigate in the sixth section. Here we calculated the quantum-mechanical correlation functions in the approximation when the oscillation modes in the horizontal direction are neglected. The effective influence of these modes is taken into account to some extent by using the exact expression for the reflection amplitude. By performing a numerical integration we observed that a two-point correlation function decays exponentially, Fig.3 and that a four-point function demonstrates tendency to decay with lower pace, Fig.4. With the numerical data available to us it is impossible to estimate scrambling time t^* [36] or to confirm its existence in the hyperbolic system considered above, but qualitatively we observe a short time exponential decay of the out-of-time correlation function and of the commutator to almost zero value and then an essential increase with the subsequent large fluctuations on Fig.4,5 .

2 Lobachevsky plane and its isometry group

Let us start with Poincare model of the Lobachevsky plane, i.e. the upper half of the complex plane: $H = \{z \in \mathbb{C}, \Im z > 0\}$ supplied with the metric (we set z = x + iy)

$$dl^2 = \frac{dx^2 + dy^2}{y^2} \tag{2.1}$$

with the Ricci scalar R = -2. Isometries of this space are given by $SL(2, \mathbb{R})$ transformations. The $SL(2, \mathbb{R})$ matrix (a,b,c,d) are real and ad-bc=1

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

acts on a point z by linear fractional substitutions

$$z \to \frac{az+b}{cz+d} \ . \tag{2.2}$$

Note also that g and -g give the same transformation, hence the effective group is $SL(2,\mathbb{R})/\mathbb{Z}_2$. We'll be interested in the space of orbits of a discrete subgroup $G \subset SL(2,\mathbb{R})$ in H. Our main example will be the modular group $G = SL(2,\mathbb{Z})$. A nice choice of the fundamental region \mathcal{F} of $SL(2,\mathbb{Z})$ is displayed in Fig.1. The fundamental region \mathcal{F} of the modular group consisting of those points between the lines $x = -\frac{1}{2}$ and $x = +\frac{1}{2}$ which lie outside the unit circle in Fig.1. The modular triangle \mathcal{F} has two equal angles $\alpha = \beta = \frac{\pi}{3}$ and with the third one equal to zero, $\gamma = 0$, thus $\alpha + \beta + \gamma = 2\pi/3 < \pi$. The area of the fundamental region is finite and equals to $\frac{\pi}{3}$ and gets a topology of sphere by "gluing" the opposite edges of the triangle. The invariant area element on the Lobachevsky plane is proportional to the square root of the determinant of the matric (2.1):

$$d\mu(z) = \frac{dxdy}{y^2} \,. \tag{2.3}$$

thus

Area(
$$\mathcal{F}$$
) = $\int_{-\frac{1}{2}}^{\frac{1}{2}} dx \int_{\sqrt{1-x^2}}^{\infty} \frac{dy}{y^2} = \frac{\pi}{3}$.

3 Geodesic flow in Hamiltonian gauge

Consider geodesic flow on \mathcal{F} , which is conveniently described by the least action principle $\delta S = 0$, where (cf. with (2.1))

$$S = \int Ldt = \int \frac{\sqrt{\dot{x}^2 + \dot{y}^2}}{y} dt . {(3.1)}$$




Figure 1. The non-compact fundamental region \mathcal{F} of a finite area is represented by the hyperbolic triangle ABD. The vertex D is at infinity of the y axis and corresponds to a cusp. The edges of the triangle are the arc AB, the rays AD and BD. The points on the edges AD and BD and the points of the arks AC with CB should be identified by the transformations w = z + 1 and w = -1/z in order to form a closed non-compact surface $\bar{\mathcal{F}}$ by "gluing" the opposite edges of the modular triangle together. The hyperbolic triangle OAB can be considered equally well as the fundamental region. The modular transformations of the fundamental region \mathcal{F} create a regular tessellation of the whole Lobachevsky plane by congruent hyperbolic triangles. K is the geodesic trajectory passing through the point (x, y) of \mathcal{F} in the \vec{v} direction.

By varying the action, we immediately get the equations of motion

$$\frac{d}{dt} \frac{\dot{x}}{y\sqrt{\dot{x}^2 + \dot{y}^2}} = 0,$$

$$\frac{d}{dt} \frac{\dot{y}}{y\sqrt{\dot{x}^2 + \dot{y}^2}} + \frac{\sqrt{\dot{x}^2 + \dot{y}^2}}{y^2} = 0.$$
(3.2)

Notice the invariance of the action and of the equations under time reparametrizations $t \to t(\tau)$. Presence of a local ("gauge") symmetry indicates that we have a constrained dynamical system. One particularly convenient choice of gauge fixing specifying the time parameter t proportional to the proper time, is archived by imposing the condition

$$\frac{\dot{x}^2 + \dot{y}^2}{y^2} = 2H\,, (3.3)$$

where H is a constant. In this gauge the equations (3.2) will take the form [21]

$$\frac{d}{dt} \left(\frac{\dot{x}}{y^2} \right) = 0$$

$$\frac{d}{dt} \left(\frac{\dot{y}}{y^2} \right) + \frac{2H}{y} = 0.$$
(3.4)




Figure 2. The incoming and outgoing plane waves. The plane wave $e^{-ip_y\tilde{y}}$ incoming from infinity of the y axis on Fig. 1 (the vertex \mathcal{D}) elastically scatters on the boundary ACB of the fundamental triangle \mathcal{F} on Fig. 1. The reflection amplitude is a pure phase and is given by the expression in front of the outgoing plane wave $e^{ip_y\tilde{y}}$. The rest of the wave function describes the standing waves in the x direction between boundaries $x = \pm 1/2$ with the amplitudes, which are exponentially decreasing.

Defining the canonical momenta p_x , p_y , conjugate to the coordinates x, y as

$$p_x = \frac{\dot{x}}{y^2}, \quad p_y = \frac{\dot{y}}{y^2},$$
 (3.5)

we shall get the geodesic equations (3.4) in the Hamiltonian form:

$$\dot{p_x} = 0, \quad \dot{p_y} = -\frac{2H}{y}.$$
 (3.6)

Indeed, after defining the Hamiltonian as

$$H = \frac{1}{2}y^2(p_x^2 + p_y^2) \tag{3.7}$$

the corresponding equations will take the form:

$$\dot{x} = \frac{\partial H}{\partial p_x} = y^2 p_x, \qquad \dot{y} = \frac{\partial H}{\partial p_y} = y^2 p_y$$

$$\dot{p_x} = -\frac{\partial H}{\partial x} = 0, \qquad \dot{p_y} = -\frac{\partial H}{\partial y} = -y(p_x^2 + p_y^2) = -\frac{2H}{y}$$
(3.8)

and coincide with (3.5) and (3.6). The advantage of the gauge (3.3) is that the Hamiltonian (3.7) coincides with the constraint.

4 Quantization

Now it is fairly standard to quantize this Hamiltonian system. We simply replace in (3.7)

$$p_x = -i\frac{\partial}{\partial x}, \qquad p_y = -i\frac{\partial}{\partial y}$$

and consider the (time independent) Schrödinger equation

$$H\psi = E\psi$$
.

The resulting equation explicitly reads:

$$-y^2(\partial_x^2 + \partial_y^2)\psi = E\psi. \tag{4.1}$$

On the lhs one easily recognises the Laplace operator (with an extra minus sign) in Poincare metric (2.1). It is easy to see that the Hamiltonian is positive semi-definite Hermitian operator. Indeed, for any quadratically integrable function $\psi(x, y)$

$$-\int \psi^*(x,y) \, y^2 (\partial_x^2 + \partial_y^2) \, \psi(x,y) \frac{dxdy}{y^2} = \int (|\partial_x \psi(x,y)|^2 + |\partial_y \psi(x,y)|^2) dxdy \ge 0.(4.2)$$

It is convenient to introduce parametrization of the energy E = s(1 - s) and to rewrite this equation as

$$-y^{2}(\partial_{x}^{2} + \partial_{y}^{2}) \ \psi(x, y) = s(1 - s) \ \psi(x, y). \tag{4.3}$$

As far as E is real and semi-positive and parametrisation is symmetric with respect to $s \leftrightarrow 1-s$ it follows that the parameter s should be chosen within the range

$$s \in [1/2, 1] \text{ or } s = 1/2 + iu, \quad u \in [0, \infty].$$
 (4.4)

One should impose the "periodic" boundary condition on the wave function with respect to the modular group

$$\psi(\frac{az+b}{cz+d}) = \psi(z), \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$$
(4.5)

in order to have the wave function which is properly defined on the fundamental region $\bar{\mathcal{F}}$ shown in Fig. 1. Taking into account that the transformation $T:z\to z+1$ belongs to SL(2,Z), one has to impose the periodicity condition $\psi(z)=\psi(z+1)$. Thus we have a Fourier expansion

$$\psi(x,y) = \sum_{n=-\infty}^{\infty} f_n(y) \exp(2\pi i n x). \tag{4.6}$$

Inserting this into Eq. (4.3), for the Fourier component $f_n(y)$ we get

$$\frac{d^2 f_n(y)}{dy^2} + (s(1-s) - 4\pi^2 n^2) f_n(y) = 0.$$
 (4.7)

For the case $n \neq 0$ the solution which exponentially decays at large y reads

$$f_n(y) = \sqrt{y} K_{s-\frac{1}{2}}(2\pi n|y|) \tag{4.8}$$

and for n = 0 one simply gets

$$f_0(y) = c_0 y^s + c_0' y^{1-s}. (4.9)$$

Thus the solution can be represented in the form

$$\psi(x,y) = c_0 y^s + c_0' y^{1-s} + \sqrt{y} \sum_{\substack{n=-\infty\\n\neq 0}}^{\infty} c_n K_{s-\frac{1}{2}}(2\pi n|y|) \exp(2\pi i n x), \tag{4.10}$$

where the coefficients c_0, c'_0, c_n should be defined such that the wave function will fulfil the boundary conditions (4.5). Thus one should impose also the invariance with respect to the second generator of the modular group SL(2, Z), that is, with respect to the transformation $S: z \to -1/z$:

$$\psi(z) = \psi(-1/z). \tag{4.11}$$

This functional equation defines the coefficients c_0, c'_0, c_n . We have found that it is much easier to resolve it using the full group of SL(2, Z) transformations acting on a particular solution (4.9). The wave function generated in this way will be invariant with respect to the SL(2, Z) transformations. We shall follow this approach in the next section.

5 Continuous spectrum and the reflection amplitude

As we just mentioned above in order to get SL(2, Z) invariant solutions, one should define the coefficients c_0, c'_0 and c_n in (4.10). Another option is to start from a particular solution and perform summation over all nonequivalent shifts of the argument by the elements of SL(2, Z), that is using the Poincaré series representation [2, 3]. We'll demonstrate this strategy using the simplest solution (4.9), (4.10) with $c_0 = 1, c'_0 = 0$:

$$\psi(z) = y^s = (\Im z)^s.$$

Let us denote by Γ_{∞} the subgroup of $\Gamma = SL(2, \mathbb{Z})$, generating shifts $z \to z + n$, $n \in \mathbb{Z}$. Explicitly the elements of Γ_{∞} are given by 2×2 matrices:

$$g_n = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \tag{5.1}$$

Since y^s is already invariant with respect to Γ_{∞} , we should perform summation over the conjugacy classes $\Gamma_{\infty}\backslash\Gamma$. Let us define these conjugacy classes. If two $SL(2, \mathbb{Z})$ matrices

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \qquad \text{and} \qquad \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$$

belong to the same class, then by definition for some $n \in \mathbb{Z}$

$$\begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

so that c'=c, d'=d, a'-a=nc and b'-b=nd. Since ad-bc=1 it follows that a and c do not have a common devisor. In fact, the opposite is also true. Given a pair of mutually prime integers (c,d) it is always possible to find a pair of integers (a,b) such that ad-bc=1. For any other pair (a',b') satisfying the same condition a'd-b'c=1, the relations a'-a=nc and b'-b=nd are satisfied for some integer n. Thus we established a bijection between the set of mutually prime pairs (c,d) with $(c,d) \neq (0,0)$ and the set of conjugacy classes $\Gamma_{\infty} \backslash \Gamma$. The fact that the integers (c,d) are mutually prime integers means that their greatest common divisor (gcd) is equal to one: gcd(c,d)=1. As a result, it is defined by the classical Poincaré series representation [2,3] and for the sum of our interest we get [2,3]

$$\psi_s(z) \equiv \frac{1}{2} \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} (\Im(\gamma z))^s = \frac{1}{2} \sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ \gcd(c,d)=1}} \frac{y^s}{((cx+d)^2 + c^2 y^2)^s} , \qquad (5.2)$$

where, as explained above, the sum on r.h.s. is taken over all mutually prime pairs (c,d). The series (5.2) is convergent when $\Re s > 1$. We have used also the simple relation

$$\Im \gamma z \equiv \Im \frac{az+b}{cz+d} = \frac{y}{(cx+d)^2 + c^2 y^2} .$$

To simplify further the sum let us multiply both sides of the eq. (5.2) by [24]

$$\sum_{n=1}^{\infty} \frac{1}{n^{2s}} \equiv \zeta(2s)$$

so that we shall get

$$\zeta(2s)\,\psi_s(z) = \frac{1}{2} \sum_{n=1}^{\infty} \sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ ard(c,d)=1}} \frac{y^s}{((ncx+nd)^2 + (nc)^2 y^2)^s}.$$
 (5.3)

It is easy now to get convinced that the set of all pairs (nc, nd) with n a positive integer and (c, d) - mutually prime, coincides with the set of all pairs of integers (m, k) which are not simultaneously zero. Indeed, given a pair (m, k) we can factor out the greatest common divisor n and represent it as (nc, nd) with mutually prime (c, d). Thus we arrive at the Eisenstein series representation of the wave function

$$\zeta(2s)\,\psi_s(z) = \frac{1}{2} \sum_{\substack{(m,k) \in \mathbb{Z}^2\\(m,k) \neq (0,0)}} \frac{y^s}{((mx+k)^2 + m^2 y^2)^s}.$$
 (5.4)

¹ The factor 1/2 below is introduced to remove the double degeneracy due to the fact that $SL(2, \mathbb{Z})$ elements $\pm \gamma$ both act on z in the same way.

Since the r.h.s. of this equation is periodic in x with period 1, we can expand it in Fourier series. Our next goal is to find the coefficients of this expansion

$$c_l(y) = \frac{1}{2} \sum_{\substack{(m,k) \in \mathbb{Z}^2 \\ (m,k) \neq (0,0)}} \int_0^1 \frac{y^s e^{-2\pi i l x} dx}{((mx+k)^2 + m^2 y^2)^s}.$$
 (5.5)

First let us handle the term with m = 0:

$$\frac{1}{2} \sum_{\substack{k \in \mathbb{Z} \\ k \neq 0}} \int_0^1 \frac{y^s e^{-2\pi i l x} dx}{k^{2s}} = \delta_{l,0} \zeta(2s) y^s.$$
 (5.6)

For the sum over non-zero m's let's notice that we may drop the factor 1/2 and sum over $m \ge 1$. Indeed, the sum over negative m's can be reverted to a sum over positive ones through redefinition $k \to -k$. For fixed positive m it is instructive to represent k as k = nm + s thus splitting the initial sum over $k \in \mathbb{Z}$ into double sum over $r = 0, 1, \ldots, m-1$ and $n \in \mathbb{Z}$. In this way after few simple manipulations we get

$$\sum_{m=1}^{\infty} \sum_{r=0}^{m-1} \sum_{n \in \mathbb{Z}} \int_{0}^{1} \frac{y^{s} e^{-2\pi i l x} dx}{((m(x+n)+r)^{2}+m^{2}y^{2})^{s}} = \sum_{m=1}^{\infty} \sum_{r=0}^{m-1} \int_{-\infty}^{\infty} \frac{y^{s} e^{-2\pi i l x} dx}{((mx+r)^{2}+m^{2}y^{2})^{s}}$$

$$= \sum_{m=1}^{\infty} \sum_{r=0}^{m-1} m^{-2s} y^{1-s} e^{\frac{2\pi i l r}{m}} \int_{-\infty}^{\infty} \frac{\cos(2\pi |l| yx) dx}{(x^{2}+1)^{s}}.$$
(5.7)

The last integral is expressed in terms of modified Bessel's K function:

$$\int_{-\infty}^{\infty} \frac{\cos(2\pi|l|yx)dx}{(x^2+1)^s} = \begin{cases} \frac{2\pi^s}{\Gamma(s)} |ly|^{s-\frac{1}{2}} K_{s-\frac{1}{2}}(2\pi|l|y), & \text{if } l \neq 0\\ \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}, & \text{if } l = 0. \end{cases}$$
(5.8)

A useful alternative representation of modified Bessel's K function which makes its properties more transparent is given by

$$K_{iu}(y) = \frac{1}{2} \int_{-\infty}^{\infty} e^{-y \cosh t} e^{iut} dt.$$
 (5.9)

This expression allows analytical continuation of the wave function from the region $\Re s > 1$ in (5.2) into the whole complex plane s because the Bessel's $K_s(y)$ functions are well defined for any s. Besides, an easy examination shows that the finite sum

$$\sum_{r=0}^{m-1} e^{\frac{2\pi i l r}{m}} = \begin{cases} m & \text{if } m \text{ divides } l \\ 0 & \text{otherwise} \end{cases}$$
 (5.10)

To summarise, for the Fourier coefficients (5.5) we shall get

$$c_l(y) = \frac{2\pi^s}{\Gamma(s)} \tau_{s-\frac{1}{2}}(|l|) \sqrt{y} K_{s-\frac{1}{2}}(2\pi|l|y), \quad \text{if } l \neq 0$$
 (5.11)

where

$$\tau_{\nu}(n) = \sum_{a \cdot b = n} \left(\frac{a}{b}\right)^{\nu},\tag{5.12}$$

while for l = 0:

$$c_0(y) = \frac{\sqrt{\pi}\Gamma(s - \frac{1}{2})\zeta(2s - 1)}{\Gamma(s)} y^{1-s}.$$
 (5.13)

Thus we recovered the second solution y^{1-s} in (4.10) and calculated the coefficient c'_0 in front of it. Thus the invariant solution (5.4) takes the following form:

$$\zeta(2s) \,\psi_s(x,y) = \zeta(2s)y^s + \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})\zeta(2s-1)}{\Gamma(s)} y^{1-s} +$$

$$+ \sqrt{y} \frac{4\pi^s}{\Gamma(s)} \sum_{l=1}^{\infty} \tau_{s-\frac{1}{2}}(l) K_{s-\frac{1}{2}}(2\pi ly) \cos(2\pi lx).$$
(5.14)

Using Riemann's reflection relation

$$\zeta(s) = \frac{\pi^{s - \frac{1}{2}} \Gamma\left(\frac{1 - s}{2}\right)}{\Gamma\left(\frac{s}{2}\right)} \zeta(1 - s)$$
(5.15)

and introducing the notation

$$\theta(s) = \pi^{-s} \zeta(2s) \Gamma(s) \tag{5.16}$$

we arrive at the elegant final expression for the energy eigenfunctions obtained by Maas [24]:

$$\theta(s)\psi_s(z) = \theta(s)y^s + \theta(1-s)y^{1-s} + 4\sqrt{y}\sum_{l=1}^{\infty} \tau_{s-\frac{1}{2}}(l)K_{s-\frac{1}{2}}(2\pi ly)\cos(2\pi lx), \quad (5.17)$$

This wave function is well defined in the complex s plane and has a simple pole at s=1. The physical continuous spectrum was defined in (4.4), where $s=\frac{1}{2}+iu$, $u \in [0,\infty]$ so that

$$E = s(1 - s) = \frac{1}{4} + u^2. \tag{5.18}$$

The continuous spectrum wave functions $\psi_s(x,y)$ are delta function normalisable [24–29]. The wave function (5.17) can be conveniently represented also in the form

$$\psi_{\frac{1}{2}+iu}(z) = y^{\frac{1}{2}+iu} + \frac{\theta(\frac{1}{2}-iu)}{\theta(\frac{1}{2}+iu)}y^{\frac{1}{2}-iu} + 4\sqrt{y}\sum_{l=1}^{\infty}\tau_{iu}(l)K_{iu}(2\pi ly)\cos(2\pi lx), (5.19)$$

where

$$K_{-iu}(y) = K_{iu}(y), \qquad \tau_{-iu}(l) = \tau_{iu}(l) .$$
 (5.20)

The physical interpretation of the wave function becomes more transparent when we introduce the new variables

$$\tilde{y} = \ln y, \quad p_y = -u, \quad E = p_y^2 + \frac{1}{4}.$$
 (5.21)

as well as alternative normalisation of the wave function

$$\psi_{p_y}(x,\tilde{y}) \equiv y^{-\frac{1}{2}} \psi_{\frac{1}{2}+iu}(z) =$$

$$= e^{-ip_y\tilde{y}} + \frac{\theta(\frac{1}{2}+ip_y)}{\theta(\frac{1}{2}-ip_y)} e^{+ip_y\tilde{y}} + 4 \sum_{l=1}^{\infty} \tau_{ip_y}(l) K_{ip_y}(2\pi l e^{\tilde{y}}) \cos(2\pi l x).$$
(5.22)

Indeed, the first two terms describe the incoming and outgoing plane waves. The plane wave $e^{-ip_y\tilde{y}}$ incoming from infinity of the y axis on Fig.1-2 (the vertex \mathcal{D}) elastically scatters on the boundary ACB of the fundamental region \mathcal{F} on Fig. 1. The reflection amplitude is a pure phase and is given by the expression in front of the outgoing plane wave $e^{ip_y\tilde{y}}$

$$\frac{\theta(\frac{1}{2} + ip_y)}{\theta(\frac{1}{2} - ip_y)} = \exp\left[i\,\varphi(p_y)\right]. \tag{5.23}$$

The rest of the wave function describes the standing waves $\cos(2\pi lx)$ in the x direction between boundaries $x = \pm 1/2$ with the amplitudes $K_{ip_y}(2\pi ly)$, which are exponentially decreasing with index l.

In addition to the continuous spectrum the system (4.3) has a discrete spectrum [24–29]. The number of discrete states is infinite: $E_0 = 0 < E_1 < E_2 < \rightarrow \infty$, the spectrum is extended to infinity - unbounded from above - and lacks any accumulation point except infinity. Let us denote the wave functions of the discrete spectrum by $\psi_n(z)$ so that the expansion into the full set of basis vectors will take the form

$$f(x,\tilde{y}) = \sum_{n>0} a_n \,\psi_n(x,\tilde{y}) + \frac{1}{2\pi} \int_0^\infty a_{p_y} \,\psi_{p_y}(x,\tilde{y}) dp_y \tag{5.24}$$

and the Parseval identity will be

$$||f(z)||^2 = \sum_{n>0} |a_n|^2 + \frac{1}{2\pi} \int_0^\infty |a_{p_y}|^2 dp_y,$$
 (5.25)

and

$$\sum_{n\geq 0} \psi_n(x,\tilde{y}) \psi_n^*(x_1,\tilde{y}_1) + \frac{1}{2\pi} \int_0^\infty \psi_{p_y}(x,\tilde{y}) \psi_{-p_y}(x_1,\tilde{y}_1) dp_y = \delta^{(2)}(z-z_1).$$
(5.26)

The wave functions of the discrete spectrum have the form [24-27, 42-44]

$$\psi_n(z) = \sum_{l=1}^{\infty} c_l(n) \sqrt{y} K_{iu_n}(2\pi l y) \{ \frac{\cos(2\pi l x)}{\sin(2\pi l x)},$$
 (5.27)

where the spectrum $E_n = \frac{1}{4} + u_n^2$ and the coefficients $c_l(n)$ are not known analytically, but were computed numerically for many values of n [42–44]. Having explicit expressions of the wave functions one can analyse the quantum-mechanical behaviour of the correlation functions, which we shall investigate in the next sections.

6 Correlation functions

6.1 Two-point correlation function

First let us calculate the two-point correlation function

$$\mathcal{D}_{2}(\beta, t) = \langle A(t)B(0)e^{-\beta H} \rangle = \sum_{n} \langle n|e^{iHt}A(0)e^{-iHt}B(0)e^{-\beta H}|n> =$$

$$= \sum_{n,m} e^{i(E_{n}-E_{m})t-\beta E_{n}} \langle n|A(0)|m> \langle m|B(0)|n>.$$
(6.1)

The energy eigenvalues (5.18) are parametrised by $n \sim \frac{1}{2} + iu$ and $m \sim \frac{1}{2} + iv$, thus

$$\mathcal{D}_{2}(\beta,t) = \int_{0}^{+\infty} \int_{0}^{+\infty} du \, dv \, e^{i(u^{2}-v^{2})t-\beta(\frac{1}{4}+u^{2})} \int_{\mathcal{F}} \psi_{\frac{1}{2}-iu}(z) \, A \, \psi_{\frac{1}{2}+iv}(z) \, d\mu(z)$$

$$\int_{\mathcal{F}} \psi_{\frac{1}{2}-iv}(w) \, B \, \psi_{\frac{1}{2}+iu}(w) \, d\mu(w) ,$$
(6.2)

where the complex conjugate function is $\psi_{\frac{1}{2}+iu}^*(z) = \psi_{\frac{1}{2}-iu}(z)$. Defining the basic matrix element as

$$A_{uv} = \int_{\mathcal{F}} \psi_{\frac{1}{2} - iu}(z) A \psi_{\frac{1}{2} + iv}(z) d\mu(z) = \int_{-1/2}^{1/2} dx \int_{\sqrt{1 - x^2}}^{\infty} \frac{dy}{y^2} \psi_{\frac{1}{2} - iu}(z) A \psi_{\frac{1}{2} + iv}(z)$$
 (6.3)

for the two-point correlation function we shall get

$$\mathcal{D}_2(\beta, t) = \int_{-\infty}^{+\infty} e^{i(u^2 - v^2)t - \beta(\frac{1}{4} + u^2)} A_{uv} B_{vu} du dv .$$
 (6.4)

In terms of the new variables (5.21) the basic matrix element (6.3) will take the form

$$A_{pq} = \int_{\mathcal{F}} \psi_p^*(x, y) \left(e^{-\frac{1}{2}y} A e^{\frac{1}{2}y} \right) \psi_q(x, y) dx dy =$$

$$= \int_{-1/2}^{1/2} dx \int_{\frac{1}{2} \log(1-x^2)}^{\infty} dy \ \psi_p^*(x, y) \left(e^{-\frac{1}{2}y} A e^{\frac{1}{2}y} \right) \psi_q(x, y) . \tag{6.5}$$

The matrix element (6.3) plays a fundamental role in the investigation of the correlation functions because all correlations can be expressed through it. One should choose also appropriate observables A and B. The operator y^{-2} seems very appropriate for two reasons. First, the convergence of the integrals over the fundamental region \mathcal{F} will be much stronger². This operator is reminiscent of the exponentiated

Louiville field since $y^{-2} = e^{-2\tilde{y}}$. Thus we have to calculate the matrix element (6.5) and for the observable $A = e^{-2y}$ we shall get³

$$A_{pq} = \int_{-1/2}^{1/2} dx \int_{\frac{1}{2}\log(1-x^2)}^{\infty} dy \ \psi_p^*(x,y) e^{-2y} \psi_q(x,y) \ . \tag{6.6}$$

In all the above formulas one should also include the contribution coming from the discrete spectrum, and we always mean that they are present. But as far as the region $\Im z \gg 1$, $\Im w \gg 1$ is concerned the contribution of the sum over Bessel's functions in (5.22) and the contribution of the discrete spectrum (5.27) we shall neglect. Indeed, the wave functions of the discrete spectrum (5.27) are entirely expressed in terms of the Bessel's functions and also will be neglected in our approximation because of the their exponential decay at large $\Im z \gg 1$ as one can observe in (5.9). In this approximation of the wave function (5.22) we shall have

$$A_{pq}^{(0)} = \frac{1}{2\pi} \int_{-1/2}^{1/2} dx \int_{\frac{1}{2}\log(1-x^2)}^{\infty} dy \left(e^{-ipy} + \frac{\theta(\frac{1}{2}+ip)}{\theta(\frac{1}{2}-ip)} e^{ipy} \right) e^{-2y} \left(e^{iqy} + \frac{\theta(\frac{1}{2}-iq)}{\theta(\frac{1}{2}+iq)} e^{-iqy} \right) =$$

$$= \frac{1}{2\pi} \int_{-1/2}^{1/2} dx \int_{\frac{1}{2}\log(1-x^2)}^{\infty} dy \left(e^{-ipy} + e^{ipy-i\varphi(p)} \right) e^{-2y} \left(e^{iqy} + e^{-iqy+i\varphi(q)} \right) =$$

$$= \frac{1}{4\pi} \int_{-1/2}^{1/2} dx \left(\frac{(1-x^2)^{-1+\frac{p-q}{2i}}}{1-\frac{p-q}{2i}} + \frac{(1-x^2)^{-1+\frac{p+q}{2i}} e^{i\varphi(q)}}{1-\frac{p+q}{2i}} \right)$$

$$+ \frac{(1-x^2)^{-1-\frac{p+q}{2i}} e^{-i\varphi(p)}}{1+\frac{p+q}{2i}} + \frac{(1-x^2)^{-1-\frac{p-q}{2i}} e^{-i(\varphi(p)-\varphi(q))}}{1+\frac{p-q}{2i}} \right), \tag{6.7}$$

where we used also (5.23). Integration over x can be performed exactly with the result for the basic matrix element

$$8\pi A_{pq}^{(0)} = \frac{{}_{2}F_{1}\left(\frac{1}{2}, 1 - \frac{p-q}{2i}; \frac{3}{2}; \frac{1}{4}\right)}{1 - \frac{p-q}{2i}} + \frac{{}_{2}F_{1}\left(\frac{1}{2}, 1 - \frac{p+q}{2i}; \frac{3}{2}; \frac{1}{4}\right)}{1 - \frac{p+q}{2i}} e^{i\varphi(q)} + \frac{{}_{2}F_{1}\left(\frac{1}{2}, 1 + \frac{p+q}{2i}; \frac{3}{2}; \frac{1}{4}\right)}{1 + \frac{p+q}{2i}} e^{-i\varphi(p)} + \frac{{}_{2}F_{1}\left(\frac{1}{2}, 1 + \frac{p-q}{2i}; \frac{3}{2}; \frac{1}{4}\right)}{1 + \frac{p-q}{2i}} e^{-i(\varphi(p)-\varphi(q))}.$$
(6.8)

In this approximation for the two-point correlation function we shall get

$$\mathcal{D}_{2}^{(0)}(\beta,t) = \int_{-\infty}^{+\infty} e^{i(p^{2}-q^{2})t-\beta(\frac{1}{4}+p^{2})} A_{pq}^{(0)} A_{qp}^{(0)} dpdq . \tag{6.9}$$

This expression is very convenient for the numerical calculation. On Fig.3 one can see exponential decay of the two-point correlation function with time at different temperatures. It is expected that the two-point correlation function should decay as [36]

$$\mathcal{D}_2(\beta, t) \sim e^{-t/t_d},\tag{6.10}$$

where $t_d \sim \beta$.

³The other interesting observable is $A = \cos(2\pi Nx), N = 1, 2, ...$




Figure 3. The two-point correlation function at $\beta = 1, 2, 3$.

6.2 Four-point correlation function

The out-of-time four-point correlation function of interest is [36]

$$\mathcal{D}_4(\beta, t) = \langle A(t)B(0)A(t)B(0)e^{-\beta H} \rangle =$$

$$= \sum_{n,m,l,r} e^{i(E_n - E_m + E_l - E_r)t - \beta E_n} \langle n|A(0)|m \rangle \langle m|B(0)|l \rangle \langle l|A(0)|r \rangle \langle r|B(0)|n \rangle$$
(6.11)

as well as the behaviour of the commutator $<[A(t),B(0)]^2e^{-\beta H}>$. In order to compute the square of the commutator one should consider also the following four-point correlation functions

$$\mathcal{D}_{4}'(\beta,t) = \langle A(t)B(0)B(0)A(t)e^{-\beta H} \rangle =$$

$$= \sum_{n,m,l,r} e^{i(-E_{m}+E_{r})t-\beta E_{n}} \langle n|A(0)|m \rangle \langle m|B(0)|l \rangle \langle l|B(0)|r \rangle \langle r|A(0)|n \rangle$$
(6.12)

$$\mathcal{D}_{4}''(\beta,t) = \langle B(0)A(t)A(t)B(0)e^{-\beta H} \rangle =$$

$$= \sum_{n,m,l,r} e^{i(E_{m}-E_{r})t-\beta E_{n}} \langle n|B(0)|m \rangle \langle m|A(0)|l \rangle \langle l|A(0)|r \rangle \langle r|B(0)|n \rangle$$
(6.13)

$$\mathcal{D}_{4}'''(\beta,t) = \langle B(0)A(t)B(0)A(t)e^{-\beta H} \rangle =$$

$$= \sum_{n,m,l,r} e^{i(-E_n + E_m - E_l + E_r)t - \beta E_n} \langle n|B(0)|m \rangle \langle m|A(0)|l \rangle \langle l|B(0)|r \rangle \langle r|A(0)|n \rangle$$
(6.14)

The corresponding energy eigenvalues we shall parametrise as $n \sim \frac{1}{2} + iu$, $m \sim \frac{1}{2} + iv$, $l \sim \frac{1}{2} + il$ and $l \sim \frac{1}{2} + ir$, thus

$$\mathcal{D}_{4}(\beta,t) = \int_{-\infty}^{+\infty} e^{i(u^{2}-v^{2}+l^{2}-r^{2})t-\beta(\frac{1}{4}+u^{2})} A_{uv} B_{vl} A_{lr} B_{ru} du dv dl dr =$$

$$= \int_{-\infty}^{+\infty} e^{i(u^{2}+l^{2})t-\beta(\frac{1}{4}+u^{2})} (AB)_{ul}(t) (AB)_{lu}(t) du dl , \qquad (6.15)$$

where

$$(AB)_{ul}(t) = \int_{\mathcal{F}} A_{uv} e^{-iv^2 t} B_{vl} dv . \qquad (6.16)$$




Figure 4. The four-point correlation function $D_4^{(0)}(t,\beta)$ at $\beta=1$.

In our plane wave approximation (analogue of so called mini-superspace approximation in Liouville theory) the four-point correlation functions will take the form

$$\mathcal{D}_{4}^{(0)}(\beta,t) = \int_{-\infty}^{+\infty} e^{i(p^{2}-q^{2}+l^{2}-r^{2})t-\beta(\frac{1}{4}+p^{2})} A_{pq}^{(0)} A_{ql}^{(0)} A_{lr}^{(0)} A_{rp}^{(0)} dpdqdldr$$

$$(6.17)$$

On Fig.4 one can see the behaviour of the four-point correlation function as a function of time at different temperatures.

It was speculated that the most important correlation functions indicating the traces of the classical chaotic dynamics in quantum regime is [36]

$$C(t,\beta) = -\langle [A(t), B(0)]^2 e^{-\beta H} \rangle = -\mathcal{D}_4(\beta, t) + \mathcal{D}_4'(\beta, t) + \mathcal{D}_4''(\beta, t) - \mathcal{D}_4'''(\beta, t)$$
(6.18)

and that the parts of the above commutation relation include the correlation functions which have the following behaviour:

$$\mathcal{D}_4(\beta, t) \sim 1 - f_0 e^{\frac{2\pi}{\beta}t} . \tag{6.19}$$

These formulas and the explicit form of the wave functions allow to calculate the above correlation functions at least using numerical integration. In our plane wave approximation we calculated the commutator

$$C^{(0)}(t,\beta) = -\langle [A(t),A(0)]^{2}e^{-\beta H} \rangle = (-\mathcal{D}_{4}^{(0)} + \mathcal{D}_{4}^{'(0)} + \mathcal{D}_{4}^{''(0)} - \mathcal{D}_{4}^{'''(0)})(\beta,t) = -\int_{-\infty}^{+\infty} e^{-\beta(\frac{1}{4}+p^{2})} \{\cos(p^{2}-q^{2}+l^{2}-r^{2})t - \cos(q^{2}-r^{2})t\} A_{pq}^{(0)} A_{ql}^{(0)} A_{lr}^{(0)} A_{rp}^{(0)} dpdqdldr .$$

$$(6.20)$$

The results of the numerical integration are presented on the Fig.5. Thus by performing numerical integration we observed that a two-point correlation function decays exponentially and that a four-point function demonstrates tendency to decay with a lower pace. With the numerical data available to us it is impossible to estimate




Figure 5. Time evolution of the correlation function C(t) (6.20) at the first and the following thirty "seconds" at $\beta = 1$. With the numerical data available to us it is impossible to estimate scrambling time t^* [36] or to confirm it existence in hyperbolic system considered above, but qualitatively one can observe short time exponential decay of the commutator to almost zero value and then essential increase with the subsequent large fluctuations.

scrambling time t^* [36] or to confirm its existence in the hyperbolic system considered above, but qualitatively one can observe short time exponential decay of the commutator to almost zero value and then essential increase with the subsequent large fluctuations.

7 Acknowledgment

R.P. is thankful to Demokritos Centre for hospitality during June-July 2018 when this work was completed . H.B. would like to thank Demokritos Centre for kind hospitality and generous financial support of the MIXMAX project. G.S. would like to thank Luis Alvarez-Gaume for stimulating discussions of SL(2,Z) automorphic functions, the references [28, 31] and kind hospitality at the Simon Center for Geometry and Physics where part of this work was completed. This project received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skĺodowska-Curie grant agreement No 644121.

References

- [1] Emil Artin, Ein mechanisches system mit quasiergodischen bahnen, E. Abh. Math. Semin. Univ. Hambg. (1924) 3: 170.
- [2] Henri Poincaré *Théorie des Groupes Fuchsiennes*, Acta Mathematica, **1** (1882) 1-62.
- [3] Henri Poincaré Mémoire sur les Fonctions Fuchsiennes, Acta Mathematica, 1 (1882) 193-294.

- [4] Lazarus Fuchs, Ueber eine Klasse von Funktionen mehrerer Variablen, welche durch Umkehrung der Integrale von Lösungen der linearen Differentialgleichungen mit rationalen Coeffizienten entstehen, J. Reine Angew. Math., 89 (1880) 151-169
- [5] J. Hadamard, Les surfaces á courbures opposées et leur linges geodesiques,
 Liouville, Journal de Mathématique, 4 (1898) 27
- [6] G.Hedlund, The dynamics of geodesic flow, Bull.Am.Math.Soc. 45 (1939) 241-246
- [7] D. V. Anosov, Geodesic flows on closed Riemannian manifolds with negative curvature, Trudy Mat. Inst. Steklov., Vol. **90** (1967) 3 210.
- [8] R.Bowen, Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms. (Lecture Notes in Mathematics, no. 470: A. Dold and B. Eckmann, editors). Springer-Verlag (Heidelberg, 1975), 108 pp.
- [9] A.N. Kolmogorov, New metrical invariant of transitive dynamical systems and automorphisms of Lebesgue spaces, Dokl. Acad. Nauk SSSR, 119 (1958) 861-865
- [10] A.N. Kolmogorov, On the entropy per unit time as a metrical invariant of automorphism, Dokl. Acad. Nauk SSSR, 124 (1959) 754-755
- [11] Ya.G. Sinai, On the Notion of Entropy of a Dynamical System, Doklady of Russian Academy of Sciences, **124** (1959) 768-771.
- [12] D. Ruelle, Thermodynamic Formalism, Addison-Wesley, Reading, Mass., 1978
- [13] E.Hopf. Statistik der Lösungen geodätischer Probleme vom unstabilen Typus. II. Math.Ann. 117 (1940) 590-608
- [14] E. Hopf, Ergodic theory and the geodesic flow on surfaces of constant negative curvature, Bull. Amer. Math. Soc, 77 (1971) 863-877.
- [15] I.M.Gelfand and S.V. Fomin, Geodesic flows on manifolds of constant negative curvature, Uspekhi Mat. Nauk, 7 (1952) 118-137; Amer. Math. Soc.Translation 1 (1965) 49-65.
- [16] P. Collet, H. Epstein and G. Gallavotti, Perturbations of Geodesic Flows on Surfaces of Constant Negative Curvature and Their Mixing Properties, Commun. Math. Phys. 95 (1984) 61-112
- [17] M. Pollicott, On the rate of mixing of Axiom A flows, Invent. math. 81 (1985) 413-426.
- [18] C. C. Moore, Exponential decay of correlation coefficients for geodesic flows, Group representations, ergodic theory, operator algebras, and mathematical physics (Berkeley, Calif., 1984), Math. Sci. Res. Inst. Publ., vol. 6, Springer, New York, 1987, pp. 163 181.
- [19] D. Dolgopyat, On Decay of Correlations in Anosov Flows, Annals of Mathematics Second Series, 147 (1998) 357-390
- [20] N. I. Chernov, Markov Approximations and Decay of Correlations for Anosov Flows, Annals of Mathematics Second Series. 147 (1998) 269-324

- [21] L. D. Faddeev, Feynman integral for singular Lagrangians, Theor. Math. Phys. 1 (1969) 1 [Teor. Mat. Fiz. 1 (1969) 3]. doi:10.1007/BF01028566
- [22] G.Savvidy, The Yang-Mills mechanics as a Kolmogorov K-system, Phys.Lett.B 130 (1983) 303
- [23] G. Savvidy, Classical and Quantum Mechanics of non-Abelian Gauge Fields, Nucl. Phys. B 246 (1984) 302.
- [24] H. Maass, Über eine neue Art von nichtanalytischen automorphen Funktionen, Math. Ann. 121, No 2 (1949), 141-183.
- [25] W. Roelcke, Über die Wellengleichung bei Grenzkreisgruppen erster Art, Sitzungsber. Heidelberg. Acad. Wiss. 4 Abh. (1953/1956), 161-267.
- [26] A.Selberg, Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series, Indian Journ. Math. Soc. 20 (1956) 47-87.
- [27] A.Selberg, *Discontinuous groups and harmonic analysis*, Proceedings of Stockholm Mathematical Congress (1962).
- [28] D. Bump, Automorphic Forms and Representations, Cambridge Studies in Advance Mathematics: 55, Cambridge University Press, 1998.
- [29] L.D.Faddeev, Expansion in eigenfunctions of the Laplace operator on the fundamental domain of a discrete group on the Lobachevsky plane, Trans. Moscow Math. Soc., 17 (1967) 357-386.
- [30] L.D.Faddeev, A. B. Venkov and V. L. Kalinin A non-arithmetic derivation of the Selberg trace formula, J. Soviet Math., 8 2 (1977) 171-199.
- [31] D.A.Hejhal, The Selberg Trace Formula for PSL(2, R), Lecture Notes in Mathematics 548, Springer-Verlag Vol. 1 1976.
- [32] G. Savvidy, Anosov C-systems and random number generators, Theor. Math. Phys. 188 (2016) 1155; doi:10.1134/S004057791608002X [arXiv:1507.06348 [hep-th]].
- [33] G. Savvidy and N. Ter-Arutyunyan-Savvidy, On the Monte Carlo simulation of physical systems, J.Comput.Phys. 97 (1991) 566; Preprint EFI-865-16-86-YEREVAN, Jan. 1986. 13pp.
- [34] K.Savvidy, The MIXMAX random number generator, Comput.Phys.Commun. 196 (2015) 161-165. (http://dx.doi.org/10.1016/j.cpc.2015.06.003); arXiv:1404.5355
- [35] K. Savvidy and G. Savvidy, Spectrum and Entropy of C-systems. MIXMAX random number generator, Chaos Solitons Fractals **91** (2016) 33 doi:10.1016/j.chaos.2016.05.003 [arXiv:1510.06274 [math.DS]].
- [36] J. Maldacena, S. H. Shenker and D. Stanford, A bound on chaos, JHEP 1608 (2016) 106 doi:10.1007/JHEP08(2016)106 [arXiv:1503.01409 [hep-th]].
- [37] G. Gur-Ari, M. Hanada and S. H. Shenker, Chaos in Classical D0-Brane

- Mechanics, JHEP **1602** (2016) 091 doi:10.1007/JHEP02(2016)091 [arXiv:1512.00019 [hep-th]].
- [38] G. Savvidy and K. Savvidy, Exponential decay of correlations functions in MIXMAX generator of pseudorandom numbers, Chaos Solitons and Fractals, 107 (2018) 244. https://doi.org/10.1016/j.chaos.2018.01.007
- [39] H. Poghosyan, H. Babujian and G. Savvidy, Artin Billiard Exponential Decay of Correlation Functions, arXiv:1802.04543 [nlin.CD].
- [40] M. Hanada, H. Shimada and M. Tezuka, Universality in Chaos: Lyapunov Spectrum and Random Matrix Theory, Phys. Rev. E 97 (2018) no.2, 022224 doi:10.1103/PhysRevE.97.022224 [arXiv:1702.06935 [hep-th]].
- [41] T. Anous and C. Cogburn, Mini-BFSS in Silico, arXiv:1701.07511 [hep-th].
- [42] A.Winkler, Cusp forms and Hecke groups, J. Reine Angew. Math. 386 (1988) 187
- [43] D.A.Hejhal, Eigenvalues of the Laplacian for PSL(2,Z): some new results and computational techniques, in International Symposium in Memory of Hua Loo-Keng (ed. by Gong, Lu, Wang, Yang), Science Press and Springer-Verlag 1 (1991) 59-102.
- [44] D.A.Hejhal and B.Berg, Some new results concerning eigenvalues of the non-Euclidean Laplacian for PSL(2,Z), Univ. of Minn. Math. Report No. 82-172 (1982) 7pp.