

Large N twisted partition functions in 3d-3d correspondence and Holography

Dongmin Gang

Center for Theoretical Physics, Seoul National University, Seoul 08826, Korea

Nakwoo Kim

Department of Physics and Research Institute of Basic Science, Kyung Hee University, Seoul 02447, Korea

School of Physics, Korea Institute for Advanced Study, Seoul 02445, Korea

(Dated: August 14, 2018)

We study the large N limit of twisted partition functions on $\mathcal{M}_{g,p}$, the S^1 bundle of degree p over a Riemann surface of genus g , for 3D $\mathcal{N} = 2$ superconformal field theories arising as low-energy limit of wrapped N M5-branes on hyperbolic 3-manifold M . We study contributions from two Bethe vacua which correspond to two canonical irreducible $SL(N, \mathbb{C})$ flat connections on M via 3D-3D correspondence. Using mathematical results on perturbative Chern-Simons invariants around the flat connections, we find universal expressions for the large N twisted partition functions contributed from the two Bethe vacua in term of the hyperbolic volume of M . The two large N partition functions perfectly match the on-shell actions for two Boltzmann-type solutions in the holographic dual AdS_4 gravity respectively.

INTRODUCTION AND SUMMARY

As a consistent theory of quantum gravity, string/M-theory is expected to provide microscopic understandings of quantum aspects of black holes (BHs). In a celebrated work [1], this hope was realized for extremal black holes in Minkowski spacetime, by reproducing the Bekenstein-Hawking entropy through counting D-brane bound states. Recently, this success was extended to black holes in asymptotically anti-de-Sitter spacetime. In particular, the entropy of magnetically charged supersymmetric black holes in AdS_4 supergravity can be explained using the holographic principle [2–11]. The AdS/CFT correspondence [12] says quantum gravity in asymptotically AdS_4 spacetime should be dual to a conformal field theory (CFT) on the 3-dimensional (3D) boundary. The field theories of interest have $\mathcal{N} \geq 2$ supersymmetry, and the black hole entropy on the gravity side turns out to be related to the so-called *topologically twisted indices* [13–16] in the dual field theory. They are the partition functions (ptns) on supersymmetric curved backgrounds $\mathcal{M}_{g,p=0} := \Sigma_g \times S^1$, with an appropriately chosen background magnetic flux coupled to R-symmetry current. The magnetic flux is turned on along the Riemann surface Σ_g of genus g . The twisted indices can be then computed using the supersymmetric localization technique. Recently, the realm of localizable 3D manifolds has been further extended in [17], and we now have formulae for twisted partition functions $Z_{g,p}$, i.e. the ptn on degree- p S^1 -bundles $\mathcal{M}_{g,p}$ (see (4)) over Σ_g .

In this letter, we study holographic duality for a large class of 3D $\mathcal{N} = 2$ SCFTs $T_N[M]$, defined in (2) (see also (3)), arising from wrapped M5-branes on closed hyperbolic 3-manifolds M . The 3D theory is characterized by N , number of M5-branes, and the choice of a 3-manifold M . For each M ,

there is an associated AdS_4/CFT_3 correspondence. The holographic dual of the wrapped M5-brane 3D SCFT was studied in [18, 19]. Since there are infinitely many such 3-manifolds [20], the wrapped M5-branes system provides a huge set of AdS_4/CFT_3 examples. We probe the holography using the twisted ptns. For $p = 0$, the ptn becomes a twisted index which counts ground states of M5-branes on $\Sigma_g \times M$. The counting is holographically dual to the microstates counting for a supersymmetric BH solution interpolating the asymptotic AdS_4 and its near-horizon limit $AdS_2 \times \Sigma_g$ [11].

The 3D-3D correspondence [21–23] provides a novel way of analyzing the 3D $\mathcal{N} = 2$ SCFTs. Schematically, the correspondence says

$$\begin{aligned} & (\text{supersymmetric ptns of 3D } T_N[M] \text{ theory}) \\ & = (SL(N, \mathbb{C}) \text{ Chern-Simons theory invariants on } M) . \end{aligned} \quad (1)$$

Refer to [24–31] for more details on 3D-3D dictionary. Via the correspondence, some supersymmetric quantities of 3D $T_N[M]$ theory can be evaluated without relying on a field theoretic description of the 3D SCFT. For example, as summarized in Table I, the twisted partition function $Z_{g,p}$ can be written in terms of basic perturbative invariants of the complex Chern-Simons (CS) theory. These invariants are mathematically well-defined and have been extensively studied in math literature.

Combining the 3D-3D dictionary with mathematical results, we obtain the large N behavior of the twisted partition functions contributed from two distinguished Bethe vacua in the $T_N[M]$ theory. The Bethe vacua of our interest correspond to two irreducible $SL(N, \mathbb{C})$ flat connections on M in 3D-3D dual complex CS theory. These two flat connections also have a natural interpretation in terms of hyperbolic geometry, see eq (18). They also give global minimum and maximum of

the absolute value of the fibering operator \mathcal{F} appearing in the ptn computation (12), see (20). We confirm that the large N twisted ptns from the two Bethe-vacua nicely match the on-shell action for the two Bolt-type solutions [10] in the gravity dual respectively. The comparison is summarized in Table II, which is the main point of this letter.

3D $T_N[M]$ THEORY

A large class of 3D $\mathcal{N} = 2$ SCFTs can be engineered through a twisted compactification of 6-dimensional SCFTs. They are labelled by the internal 3-manifolds M :

$$T_N[M] := (\text{Effective 3D } \mathcal{N} = 2 \text{ SCFT obtained from a twisted compactification of 6D } A_{N-1} (2, 0) \text{ theory on a 3-manifold } M) . \quad (2)$$

For simplicity, we assume M is a closed (compact) hyperbolic 3-manifold without boundary. To preserve supersymmetry, we perform a partial topological twisting along the internal 3-manifold using $SO(3)$ vector subgroup of $SO(5)$ R-symmetry of the 6D theory. The twisting preserves a quarter of supersymmetries and the resulting 3D theory becomes a 3D $\mathcal{N} = 2$ SCFT with 4 supercharges. The 6D theory describes the low-energy effective world-volume theory of N coincident M5-branes in M-theory, and the 3D theory can be considered as an effective world-volume theory of N coincident M5-branes wrapped on the compact 3-manifold M , i.e.

$$\begin{aligned} & N \text{ coincident M5-branes on } \mathbb{R}^3 \times M \\ & \text{in M-theory on } \mathbb{R}^3 \times (T^*M) \times \mathbb{R}^2 \\ & \xrightarrow{\text{IR world-volume theory of M5-branes}} T_N[M] \text{ on } \mathbb{R}^3 . \end{aligned} \quad (3)$$

Here T^*M is the cotangent bundle of M , which is a local Calabi-Yau.

Let us comment on a subtle point in the setup. As emphasized in [32], in taking the twisted compactification we need to choose a connected subset of the vacuum moduli-space of the theory defined on \mathbb{R}^3 , in order to have a genuine 3D SCFT. For a hyperbolic M , there is a natural choice (which is actually a single point) which is expected to become a discrete set of vacua when the theory is put on $\mathbb{R}^2 \times S^1$. This discrete set of vacua corresponds to a subset of irreducible $SL(N, \mathbb{C})$ flat connections on M . A field theoretic construction of the effective 3D gauge theory is proposed in [32], extending the beautiful construction in [23, 33] for cusped 3-manifolds with at least one torus boundary component, by incorporating gauge theoretical operations corresponding to Dehn filling (removing torus boundaries) operations on 3-manifold.

$T_N[M]$ ON $\mathcal{M}_{g,p}$ FOR EVEN p

We now turn to the case where $T_N[M]$ is put on a large class of nontrivial backgrounds $\mathcal{M}_{g,p}$ [17]:

$$\begin{aligned} & \mathcal{M}_{g,p} := (S^1\text{-bundle of degree } p \\ & \text{over a Riemann surface } \Sigma_g \text{ of genus } g) , \quad (4) \\ & \text{i.e. } S^1 \xrightarrow{p} \mathcal{M}_{g,p} \rightarrow \Sigma_g . \end{aligned}$$

The metric can be written as

$$ds^2 = \beta^2(d\psi - pa(z, \bar{z}))^2 + 2g_{z\bar{z}}dzd\bar{z} , \quad (5)$$

where z, \bar{z} are local coordinates on the Riemann surface and $\psi \sim \psi + 2\pi$ parameterizes the S^1 -fiber of length β . a is a 1-form on Σ_g whose curvature $F_a := da$ is normalized as

$$\frac{1}{2\pi} \int_{\Sigma_g} da = 1 . \quad (6)$$

To preserve some supersymmetry, we turn on the following background gauge field coupled to $U(1)$ R-symmetry.

$$A^R = \beta\nu_R(d\psi - pa) + n_R(\pi^*a) , \quad (7)$$

with proper quantization conditions for (ν_R, n_R) [10]. Here π^*a is a 1-form on $\mathcal{M}_{g,p}$ given as the pull-back of a using the projection map $\pi : \mathcal{M}_{g,p} \rightarrow \Sigma_g$.

For later comparison with the bolt-type solutions in the supergravity, we follow [10] and choose

$$\nu_R = \frac{1}{2} , \quad n_R = \frac{p}{2} + g - 1 , \quad p \in 2\mathbb{Z} . \quad (8)$$

Throughout the letter, we restrict our attention to the choice in (8) and some formulae below may not work for other cases. For example, the large N computation in Table II give incorrect answer for the usual round S^3 case which is $\mathcal{M}_{g=0,p=1}$.

For small N the effective 3d theory $T_N[M]$ might witness emergent symmetries in addition to R-symmetry, as pointed out in [32, 34]. When N is large enough, on the other hand, there is no accidental symmetry and the $U(1)$ R-symmetry in the IR should be simply inherited from the compact $SO(2)$ subgroup of $SO(5)$ R-symmetry in the 6D theory. It implies that the $U(1)$ R-charge, R , should be properly quantized

$$R(\mathcal{O}) \in \mathbb{Z} , \text{ for any state } \mathcal{O} \text{ of } T_N[M] \text{ on } \Sigma_g . \quad (9)$$

The Dirac quantization conditions for the $U(1)$ R-symmetry flux on Σ_g are

$$R(\mathcal{O}) \times n_R = R(\mathcal{O}) \times \left(\frac{p}{2} + g - 1\right) \in \mathbb{Z} , \quad (10)$$

for any state \mathcal{O} of $T_N[M]$ on Σ_g .

From (9), we see that the Dirac quantizations are always satisfied for even p . In summary, for large enough N we can put the 3D $T_N[M]$ theory on any $\mathcal{M}_{g \in \mathbb{Z}_{\geq 0}, p \in 2\mathbb{Z}}$ with supersymmetry preserving background gauge field, given in (7) and (8), coupled to the R-symmetry in the IR.

HOLOGRAPHIC DUAL OF $T_N[M]$

The gravity dual description is given by the uplift of a certain magnetically charged AdS_4 solution in the maximally supersymmetric $D = 7$ gauged supergravity [18, 19]. Schematically, the $D = 11$ solution is a product of AdS_4 , hyperbolic 3-manifold M , and a squashed 4-sphere \tilde{S}^4 . Consistency of the truncation from $D = 11$ down to minimal $\mathcal{N} = 2$, $D = 4$ gauged supergravity is established in [35] and it is guaranteed that we may replace the AdS_4 part with any nontrivial $D = 4$ solution and we still have an exact $D = 11$ solution.

The computation of holographic free energy can be also first done in $D = 4$ setup, and substitute the Newton constant with [36]

$$G_4 = \frac{3\pi^2}{2N^3 \text{vol}(M)}. \quad (11)$$

Here, the hyperbolic volume is defined as

$$\begin{aligned} \text{vol}(M) &= (\text{hyperbolic volume of } M) \\ &:= (\text{volume measured in the unique hyperbolic metric}). \end{aligned}$$

The hyperbolic metric is normalized as $R_{\mu\nu} = -2g_{\mu\nu}$. The Mostow's rigidity theorem [37] guarantees the uniqueness of the hyperbolic metric and thus the volume is actually a topological invariant.

As gravity duals of the boundary theory put on $\mathcal{M}_{g,p}$, we utilize the supersymmetric AdS-Taub-NUT and bolt solutions constructed in [38]. Since these solutions have non-vanishing Maxwell field, which in $D = 11$ uplift appears as a twisting of the R-symmetry angle in \tilde{S}^4 , one might worry about a conflict with the quantization condition g, p . But it turns out, since the R-symmetry angle is part of \tilde{S}^4 and we have a standard periodicity of 2π , the regularity condition for $D = 4$ NUT/Bolt is enough. This is in line with the field theory side discussion, in particular (9). A comment is in order here, in comparison with the uplifts involving Sasaki-Einstein 7-manifolds. In that case, the periodicity of the R-symmetry angle from the regularity of Bolt solution should be compatible with the periodicity condition due to collapsing cycles in the Kähler-Einstein base manifold of the Sasaki-Einstein space. The readers are referred to [10] for more details, where the authors considered an explicit example of Sasaki-Einstein manifolds such as $V^{5,2} = SO(5)/SO(3)$.

TWISTED PARTITION FUNCTIONS OF $T_N[M]$ IN 3D-3D CORRESPONDENCE

The twisted partition function $Z_{g,p}$ on the $\mathcal{M}_{g,p}$ for general 3D $\mathcal{N} = 2$ SCFTs is given as the following finite sum [17, 29]

$$Z_{g,p} = \sum_{\alpha} Z_{g,p}^{\alpha} := \sum_{\alpha} (\mathcal{H}^{\alpha})^{g-1} (\mathcal{F}^{\alpha})^p. \quad (12)$$

Here α labels the so-called Bethe vacua [39] of the 3D theory. It is obtained by extremizing the effective 2d twisted superpotential in the compactification on $\mathbb{R}^2 \times S^1$. The number of vacua is equal to the Witten index [40, 41] of the 3D SCFT. \mathcal{H} and \mathcal{F} are called *handle-gluing* and *fibering* operators respectively. The explicit forms of \mathcal{H} and \mathcal{F} for any given ultraviolet (UV) Lagrangian are available in [17]. Let us emphasize that, the formula (12) applied to the case of S^3 partition function which corresponds to $(g, p) = (0, 1)$ is apparently *different* from the more familiar Coulomb branch integral expression [42]. But their equivalence is illustrated for a number of examples in [17].

3D $T_N[M]$ theory	$SL(N, \mathbb{C})$ CS theory
Bethe vacuum α	Irreducible flat connection \mathcal{A}^{α}
Handle gluing operator \mathcal{H}^{α}	$\exp(-2S_1^{\alpha})$
Fibering operator \mathcal{F}^{α}	$\exp(iS_0^{\alpha}/(2\pi))$

TABLE I: 3d-3d dictionaries for basic ingredients in twisted partition function computation. $S_{n=0,1}^{\alpha}$ are perturbative invariants of the complex Chern-Simons theory around a flat-connection \mathcal{A}^{α} , see eq (15).

Now let us specialize to the $T_N[M]$ theories in (2). The twisted ptns for these theories can be analyzed using the 3D-3D dictionaries summarized in Table I. Twisted ptns in 3D-3D correspondence were studied in [29, 30]. In the table, the $\{S_n^{\alpha}\}_{n=0}^{\infty}$ represent terms in the loop expansion of the complex CS partition function around a flat-connection \mathcal{A}^{α} [43–46]:

$$\begin{aligned} Z_{\text{CS}}^{\alpha} \text{ pert} &:= \int \frac{D(\delta\mathcal{A})}{(\text{gauge})} e^{-\frac{1}{2\hbar} CS[\mathcal{A}^{\alpha} + \delta\mathcal{A}; M]} \\ &\xrightarrow{\text{as } \hbar \text{ goes to } 0} \exp\left(\frac{1}{\hbar} S_0^{\alpha} + S_1^{\alpha} + \dots + \hbar^{n-1} S_n^{\alpha} + \dots\right). \end{aligned} \quad (13)$$

The Chern-Simons functional is

$$CS[\mathcal{A}, M] := \int_M \text{Tr}(\mathcal{A} \wedge d\mathcal{A} + \frac{2}{3} \mathcal{A}^3). \quad (14)$$

Note that the counterpart of \mathcal{F} and \mathcal{H} are simply *tree level* and *one-loop* contributions in perturbation theory! More explicitly, the perturbative coefficients are given as

$$\begin{aligned} S_0^{\alpha} &= -\frac{1}{2} CS[\mathcal{A}^{\alpha}, M], \\ S_1^{\alpha} &:= \frac{1}{2} \log \text{Tor}_{R=\text{adjoint}}[\mathcal{A}^{\alpha}, M]. \end{aligned} \quad (15)$$

$\text{Tor}_R[\mathcal{A}^{\alpha}, M]$ is the Ray-Singer torsion of an associated vector bundle in a representation $R \in \text{Hom}[SL(N, \mathbb{C}) \rightarrow GL(V_R)]$ twisted by a flat connection \mathcal{A}^{α} . Here V_R is the vector space for representation R and $GL(V_R)$ is the general linear group

on the V_R . The analytic torsion is defined as follows [45, 47, 48]

$$\text{Tor}_R[\mathcal{A}^\alpha, M] := \frac{[\det' \Delta_0(R, \mathcal{A}^\alpha)]^{3/2}}{[\det' \Delta_1(R, \mathcal{A}^\alpha)]^{1/2}}. \quad (16)$$

Here $\Delta_n(R, \mathcal{A}^\alpha)$ is a Laplacian acting on V_R -valued n -form twisted by a flat connection \mathcal{A}^α . $\det' \Delta$ denotes the zeta function regularized determinant of the Laplacian Δ . For the one-loop part, the denominator comes from gauge field fluctuations $\delta \mathcal{A}$ while the numerator comes from the ghosts associated to a gauge fixing [49].

The 3D-3D dictionary in Table I can be derived combining several known results in literatures. The Bethe-vacua (vacua on $\mathbb{R}^2 \times S^1$) of 3D $T_N[M]$ theory are in one-to-one correspondence to a subset of irreducible flat-connections on M [21, 32]. According to a dictionary of 3D-3D relation, the asymptotic expansion $Z_{\text{CS pert}}^\alpha$ in (14) is equal to the perturbative expansion of holomorphic block $B^\alpha(q)$ [21, 27] associated to the Bethe-vacuum α in the limit $q \rightarrow 1$,

$$Z_{\text{CS pert}}^\alpha(\hbar) \simeq B^\alpha(q := e^\hbar), \quad (17)$$

as an asymptotic expansion in $\hbar \rightarrow 0$.

For general 3D $\mathcal{N} = 2$ theory, the asymptotic expansion coefficients S_0 and S_1 of holomorphic block are related to the operators \mathcal{F} and \mathcal{H} as given in Table I [50, 51].

LARGE N TWISTED PARTITION FUNCTIONS AROUND TWO BETHE-VACUA AND ITS HOLOGRAPHIC DUAL

For every hyperbolic 3-manifold M , there are two characteristic irreducible $SL(N, \mathbb{C})$ flat connections $\mathcal{A}_N^{\text{geom}}$ and $\overline{\mathcal{A}}_N^{\text{geom}}$ which can be constructed from the hyperbolic structure on M ,

$$\mathcal{A}_N^{\text{geom}} := \rho_N \cdot (\omega + ie), \quad \overline{\mathcal{A}}_N^{\text{geom}} := \rho_N \cdot (\omega - ie). \quad (18)$$

Here ω and e are respectively the spin-connection and vielbein of the unique hyperbolic metric on M . They are both locally $so(3)$ -valued 1-forms and the complex combinations $\omega \pm ie$ form an $SL(2, \mathbb{C})$ flat-connections on M . ρ_N is the N -dimensional irreducible representation of $sl(2, \mathbb{C}) = su(2)_\mathbb{C}$, and obviously $\rho_N \cdot (\omega \pm ie)$ become also irreducible $SL(N, \mathbb{C})$ flat connections. A crucial property of these two flat-connections is that they take the minimum (maximum) value of $\text{Im}[S_0]$ among all $SL(N, \mathbb{C})$ flat connections. Namely,

$$\text{Im}[S_0^{\overline{\text{geom}}}] < \text{Im}[S_0^\alpha] < \text{Im}[S_0^{\text{geom}}], \quad (19)$$

for any flat connection \mathcal{A}^α which is neither $\overline{\mathcal{A}}^{\text{geom}}$ nor $\mathcal{A}^{\text{geom}}$. Combined with the 3D-3D dictionary in Table I, it is implied

$$|\mathcal{F}^{\text{geom}}| < |\mathcal{F}^\alpha| < |\overline{\mathcal{F}}^{\text{geom}}|, \quad (20)$$

for any Bethe-vacuum α which is neither $(\overline{\text{geom}})$ nor (geom) .

Classical actions S_0^α for the two connections above can be computed as follows

$$\begin{aligned} \text{Im}[S_0^{\text{geom}}] &= -\frac{1}{2} \text{Im}[CS(\mathcal{A}_N^{\text{geom}})] = -\frac{1}{2} \text{Im}[CS(\rho_N \cdot (\omega + ie))] \\ &= -\frac{1}{2} \frac{\text{Tr}[\rho_N \cdot (T^a) \rho_N \cdot (T^b)]}{\text{Tr}[T^a T^b]} \text{Im}[CS(\omega + ie)] \\ &= -\frac{1}{2} \frac{N^3 - N}{6} \text{Im}[CS(\omega + ie)] = \frac{N^3 - N}{6} \text{vol}(M), \quad \text{and} \\ \text{Im}[S_0^{\overline{\text{geom}}}] &= -\text{Im}[S_0^{\text{geom}}] = -\frac{N^3 - N}{6} \text{vol}(M). \end{aligned} \quad (21)$$

In the second line, T^a ($a = 1, 2, 3$) are Pauli matrices and $\rho_N \cdot (T^a)$ are generators in the N -dimensional irreducible representation. From a simple group theoretical fact

$$\frac{\text{Tr}[\rho_N \cdot (T^a) \rho_N \cdot (T^b)]}{\text{Tr}[T^a T^b]} = \frac{N^3 - N}{6}, \quad (22)$$

the expected N^3 -scaling of $T_N[M]$ theory follows. In the third line of (21), we use the fact that the imaginary part of Chern-Simons functional of $\mathcal{A} = \omega + ie$ is equal to the Einstein-Hilbert action with unit negative cosmology constant up to an overall numerical factor [52]. The action for the unique hyperbolic metric is twice of the hyperbolic volume of 3-manifold with a minus sign.

The large N asymptotic behavior of the 1-loop coefficients, S_1^{geom} and $S_1^{\overline{\text{geom}}}$, can be analyzed using a following mathematical theorem [53],

$$\begin{aligned} \log |\text{Tor}_{R=\rho_{2m+1}}[\mathcal{A}_{N=2}^{\text{geom}}, M]| \\ \xrightarrow{\text{as } m \text{ goes to } \infty} -\frac{1}{\pi} m^2 \text{vol}(M) + o(m). \end{aligned} \quad (23)$$

Here ρ_{2m+1} is the $(2m+1)$ -dimensional irreducible representation of $sl(2, \mathbb{C}) = su(2)_\mathbb{C}$. Combining the theorem with the following branching rule,

$$(\text{adjoint of } sl(N, \mathbb{C})) = \bigoplus_{m=1}^{N-1} \rho_{2m+1} \text{ of } sl(2, \mathbb{C}), \quad (24)$$

when the $sl(2, \mathbb{C})$ is embedded into $sl(N, \mathbb{C})$ via ρ_N .

we have following large N behavior of the 1-loop coefficients

$$\begin{aligned} \text{Re}[S_1^{\text{geom}}] &= \frac{1}{2} \log |\text{Tor}_{R=\text{adjoint}}[\mathcal{A}_N^{\text{geom}}, M]| \\ &= \frac{1}{2} \sum_{m=1}^{N-1} \log |\text{Tor}_{R=\rho_{2m+1}}[\mathcal{A}_{N=2}^{\text{geom}}, M]| \\ &= -\frac{1}{2\pi} \text{vol}(M) \sum_{m=1}^{N-1} m^2 + o(m) = -\frac{N^3 + o(N^2)}{6\pi} \text{vol}(M), \end{aligned}$$

and

$$\text{Re}[S_1^{\overline{\text{geom}}}]] = \text{Re}[S_1^{\text{geom}}] = -\frac{N^3 + o(N^2)}{6\pi} \text{vol}(M). \quad (25)$$

Combining the 3D-3D dictionaries in Table I with the large N analysis in (21) and (25), we finally obtain following universal large N behavior of the twisted ptns (12)

$$\begin{aligned} F_{g,p}^{\text{geom}} &:= -\log |Z_{g,p}^{\text{geom}}(T_N[M])| \\ &= \frac{4(1-g)N^3 + pN^3}{12\pi} \text{vol}(M) + o(N^2), \\ \overline{F}_{g,p}^{\text{geom}} &:= -\log |\overline{Z}_{g,p}^{\text{geom}}(T_N[M])| \\ &= \frac{4(1-g)N^3 - pN^3}{12\pi} \text{vol}(M) + o(N^2). \end{aligned} \quad (26)$$

They nicely match the on-shell actions $I_{g,p}^{\text{Bolt}\pm}$ of $\text{Bolt}\pm$ solution in [10]. The large N computations are summarized in Table II.

M-theory on $AdS_4 \times M \times S^4$	$SL(N, \mathbb{C})$ CS theory on M
Bolt_+ solution $I_{g,p}^{\text{Bolt}_+} = \frac{\pi(4(1-g)-p)}{8G_4}$	Flat connection $\mathcal{A}_N^{\text{geom}}$ $F_{g,p}^{\text{geom}} = \frac{4(1-g)N^3 - pN^3}{12\pi} \text{vol}(M)$
Bolt_- solution $I_{g,p}^{\text{Bolt}_-} = \frac{\pi(4(1-g)+p)}{8G_4}$	Flat connection $\mathcal{A}_N^{\text{geom}}$ $\overline{F}_{g,p}^{\text{geom}} = \frac{4(1-g)N^3 + pN^3}{12\pi} \text{vol}(M)$

TABLE II: The M-theory is holographic dual to 3D $T_N[M]$ theory while the complex Chern-Simons theory is 3D-3D dual to the $T_N[M]$ theory. The 4d Newton constant G_4 is given in (11).

COMPARISON WITH LARGE N S_b^3 -PTN

The prescription to compute the twisted partition function $Z_{g,p}$ for $T_N[M]$ through 3D-3D correspondence naturally shares several ingredients with the corresponding computation of a squashed 3-sphere partition function $Z_b(T_N[M])$ studied in [36, 54]. The squashed 3-sphere S_b^3 of our interest is a supersymmetric curved background introduced in [55], defined as

$$S_b^3 = \{(z, w) \in \mathbb{C}^2 : b^2|z|^2 + \frac{1}{b^2}|w|^2 = 1\}. \quad (27)$$

Setting $b = 1$ gives the usual round 3-sphere. According to the 3D-3D relation [22, 23], the extreme squashing limit $b \in \mathbb{R} \rightarrow 0$ corresponds to a weakly coupled limit of the Chern-Simons theory. More concretely Z_b is determined by the perturbative invariants S_n^{geom} in (14) around the flat connection $\mathcal{A}_N^{\text{geom}}$ [31, 56, 57],

$$\begin{aligned} F_b(T_N[M]) &:= -\log |Z_b(T_N[M])| \\ &\xrightarrow{\text{as } \hbar := 2\pi i b^2 \text{ goes to } 0} \\ &= -\text{Re} \left[\left(\frac{1}{\hbar} S_0^{\text{geom}} + \dots + \hbar^{n-1} S_n^{\text{geom}} + \dots \right) \right]. \end{aligned} \quad (28)$$

Combined with the large N behaviors of S_n^{geom} for $n = 1, 2$ in eqs. (21) and (25), we see that the asymptotic expansion is

compatible with the gravity dual side of free-energy I_b^{gravity} [54],

$$\begin{aligned} I_b^{\text{gravity}} &= \frac{\pi(b + b^{-1})^2}{8G_4} = \frac{N^3}{12\pi} (b + b^{-1})^2 \text{vol}(M) \\ &\xrightarrow{\text{as } \hbar := 2\pi i b^2 \text{ goes to } 0} \\ &= \frac{iN^3 \text{vol}(M)}{6\hbar} + \frac{N^3 \text{vol}(M)}{6\pi} - \frac{iN^3 \text{vol}(M)}{24\pi^2} \hbar, \end{aligned} \quad (29)$$

up to $o(\hbar^0)$. Motivated from the comparison, it was further conjectured that [54]

$$\lim_{N \rightarrow \infty} \frac{1}{N^3} S_2^{\text{geom}} = \frac{i \text{vol}(M)}{24\pi^2}, \quad \lim_{N \rightarrow \infty} \frac{1}{N^3} S_{n \geq 3}^{\text{geom}} = 0. \quad (30)$$

This conjecture was checked numerically for a number of concrete examples.

Now we compare the two large N analysis and see

$$\begin{aligned} &\lim_{N \rightarrow \infty} \frac{1}{N^3} F_b(T_N[M]) \\ &= \lim_{N \rightarrow \infty} \frac{1}{N^3} \left(-\frac{\text{Im}[S_0^{\text{geom}}]}{2\pi b^2} - \text{Re}[S_1^{\text{geom}}] + (2\pi b^2) \text{Im}[S_2^{\text{geom}}] \right), \end{aligned}$$

and

$$\begin{aligned} &\lim_{N \rightarrow \infty} \frac{1}{N^3} F_{g,p}^{\text{geom}}(T_N[M]) \\ &= \lim_{N \rightarrow \infty} \frac{1}{N^3} \left(2(g-1) \text{Re}[S_1^{\text{geom}}] + \frac{p}{2\pi} \text{Im}[S_0^{\text{geom}}] \right). \end{aligned} \quad (31)$$

From the comparison, we obtain a general relation of the following form, between the twisted and the squashed S^3 partition functions in the large N limit

$$\lim_{N \rightarrow \infty} \frac{F_{g,p}^{\text{geom}}}{F_{b=1}} = (1-g) - \frac{p}{4}. \quad (32)$$

which holds for every 3D $T_N[M]$ theory. To arrive the conclusion, we use following universal relation between perturbative invariants

$$\lim_{N \rightarrow \infty} \frac{\pi \text{Re}[S_1^{\text{geom}}]}{\text{Im}[S_0^{\text{geom}}]} = \lim_{N \rightarrow \infty} \frac{-4\pi^2 \text{Im}[S_2^{\text{geom}}]}{\text{Im}[S_0^{\text{geom}}]} = 1, \quad (33)$$

which follows from the fact [58]

$$\begin{aligned} &\lim_{N \rightarrow \infty} \frac{F_b}{F_{b=1}} \\ &= \lim_{N \rightarrow \infty} \frac{-\frac{1}{2\pi} \text{Im}[S_0^{\text{geom}}] b^{-2} - \text{Re}[S_1^{\text{geom}}] + (2\pi) \text{Im}[S_2^{\text{geom}}] b^2}{-\frac{1}{2\pi} \text{Im}[S_0^{\text{geom}}] - \text{Re}[S_1^{\text{geom}}] + (2\pi) \text{Im}[S_2^{\text{geom}}]} \\ &= \frac{1}{4} (b^{-2} + 2 + b^2). \end{aligned}$$

The same universal relation (32) for $p = 0$ was observed in [4, 11] for different class of 3D $\mathcal{N} = 2$ SCFTs.

CONCLUSION

In this letter, we probe a large class of $\text{AdS}_4/\text{CFT}_3$ associated to M5-branes wrapped on 3-manifolds by computing large N twisted ptns. For M2 and D2-branes and their Chern-Simons-matter theories, the large N computations have been performed already in [2–11]. A nicer feature of our analysis is that we map the large N analysis to a mathematical problem via 3D-3D correspondence which can be solved from known mathematical results, such as (19), (21) and (25). The results hold for any closed hyperbolic 3-manifold M and one does not need to perform the large N analysis for individual $\text{AdS}_4/\text{CFT}_3$ model associated to each M . We can also give a simple explanation why there is a universal relation (see (32)) between the twisted ptns and S^3 -ptn in the large N limit. Both types of ptns are related to the same perturbative invariants of a complex CS theory through the 3D-3D correspondence in the large N limit. We hope that the improvement made in our analysis may provide a better way of understanding the sub-leading corrections to the large N twisted ptns which might be related to quantum corrections to the Bekenstein-Hawking entropy.

ACKNOWLEDGMENTS

We would like to thank Seok Kim, Sunjin Choi, Ching Hwang, Jaewon Song, Masahito Yamazaki and Victor Mikhaylov for interesting discussions on related works. This work was initiated while the authors were visiting APCTP, Pohang for a workshop “Strings, Branes and Gauge theories”, 16-25 July 2018. We thank APCTP for hospitality. The work of DG was supported by Samsung Science and Technology Foundation under Project Number SSTBA1402-08. The research of NK was supported by NRF grant 2015R1D1A1A09059301.

-
- [1] A. Strominger and C. Vafa, “Microscopic origin of the Bekenstein-Hawking entropy,” *Phys. Lett.* **B379** (1996) 99–104, arXiv:hep-th/9601029.
- [2] F. Benini, K. Hristov, and A. Zaffaroni, “Black hole microstates in AdS_4 from supersymmetric localization,” *JHEP* **05** (2016) 054, arXiv:1511.04085 [hep-th].
- [3] F. Benini, K. Hristov, and A. Zaffaroni, “Exact microstate counting for dyonic black holes in AdS_4 ,” *Phys. Lett.* **B771** (2017) 462–466, arXiv:1608.07294 [hep-th].
- [4] S. M. Hosseini and A. Zaffaroni, “Large N matrix models for $3d \mathcal{N} = 2$ theories: twisted index, free energy and black

- holes,” *JHEP* **08** (2016) 064, arXiv:1604.03122 [hep-th].
- [5] S. M. Hosseini and N. Mekareeya, “Large N topologically twisted index: necklace quivers, dualities, and Sasaki-Einstein spaces,” *JHEP* **08** (2016) 089, arXiv:1604.03397 [hep-th].
- [6] A. Cabo-Bizet, V. I. Giraldo-Rivera, and L. A. Pando Zayas, “Microstate counting of AdS_4 hyperbolic black hole entropy via the topologically twisted index,” *JHEP* **08** (2017) 023, arXiv:1701.07893 [hep-th].
- [7] F. Azzurli, N. Bobev, P. M. Cricigno, V. S. Min, and A. Zaffaroni, “A universal counting of black hole microstates in AdS_4 ,” *JHEP* **02** (2018) 054, arXiv:1707.04257 [hep-th].
- [8] S. M. Hosseini, K. Hristov, and A. Passias, “Holographic microstate counting for AdS_4 black holes in massive IIA supergravity,” *JHEP* **10** (2017) 190, arXiv:1707.06884 [hep-th].
- [9] F. Benini, H. Khachatryan, and P. Milan, “Black hole entropy in massive Type IIA,” *Class. Quant. Grav.* **35** no. 3, (2018) 035004, arXiv:1707.06886 [hep-th].
- [10] C. Toldo and B. Willett, “Partition functions on 3d circle bundles and their gravity duals,” *JHEP* **05** (2018) 116, arXiv:1712.08861 [hep-th].
- [11] N. Bobev and P. M. Cricigno, “Universal RG Flows Across Dimensions and Holography,” *JHEP* **12** (2017) 065, arXiv:1708.05052 [hep-th].
- [12] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Int. J. Theor. Phys.* **38** (1999) 1113–1133, arXiv:hep-th/9711200. [Adv. Theor. Math. Phys.2,231(1998)].
- [13] S. Gukov and D. Pei, “Equivariant Verlinde formula from fivebranes and vortices,” *Commun. Math. Phys.* **355** no. 1, (2017) 1–50, arXiv:1501.01310 [hep-th].
- [14] F. Benini and A. Zaffaroni, “A topologically twisted index for three-dimensional supersymmetric theories,” *JHEP* **07** (2015) 127, arXiv:1504.03698 [hep-th].
- [15] F. Benini and A. Zaffaroni, “Supersymmetric partition functions on Riemann surfaces,” *Proc. Symp. Pure Math.* **96** (2017) 13–46, arXiv:1605.06120 [hep-th].
- [16] C. Closset and H. Kim, “Comments on twisted indices in 3d supersymmetric gauge theories,” *JHEP* **08** (2016) 059, arXiv:1605.06531 [hep-th].
- [17] C. Closset, H. Kim, and B. Willett, “Supersymmetric partition functions and the three-dimensional A-twist,” *JHEP* **03** (2017) 074, arXiv:1701.03171 [hep-th].
- [18] M. Pernici, “Spontaneous compactification of seven-dimensional supergravity theories,” *Classical and Quantum Gravity* **2** no. 5, (1985) 673.
- [19] J. P. Gauntlett, N. Kim, and D. Waldram, “M Five-branes wrapped on supersymmetric cycles,” *Phys. Rev.* **D63** (2001) 126001, arXiv:hep-th/0012195.
- [20] W. P. Thurston, *The geometry and topology of three-manifolds*. Princeton University Princeton, NJ, 1979.

- [21] T. Dimofte, S. Gukov, and L. Hollands, “Vortex Counting and Lagrangian 3-manifolds,” *Lett. Math. Phys.* **98** (2011) 225–287, arXiv:1006.0977 [hep-th].
- [22] Y. Terashima and M. Yamazaki, “SL(2,R) Chern-Simons, Liouville, and Gauge Theory on Duality Walls,” *JHEP* **08** (2011) 135, arXiv:1103.5748 [hep-th].
- [23] T. Dimofte, D. Gaiotto, and S. Gukov, “Gauge Theories Labelled by Three-Manifolds,” *Commun. Math. Phys.* **325** (2014) 367–419, arXiv:1108.4389 [hep-th].
- [24] J. Yagi, “3d TQFT from 6d SCFT,” *JHEP* **08** (2013) 017, arXiv:1305.0291 [hep-th].
- [25] S. Lee and M. Yamazaki, “3d Chern-Simons Theory from M5-branes,” *JHEP* **12** (2013) 035, arXiv:1305.2429 [hep-th].
- [26] C. Cordova and D. L. Jafferis, “Complex Chern-Simons from M5-branes on the Squashed Three-Sphere,” *JHEP* **11** (2017) 119, arXiv:1305.2891 [hep-th].
- [27] C. Beem, T. Dimofte, and S. Pasquetti, “Holomorphic Blocks in Three Dimensions,” *JHEP* **12** (2014) 177, arXiv:1211.1986 [hep-th].
- [28] T. Dimofte, “Complex Chern-Simons Theory at Level k via the 3d-3d Correspondence,” *Commun. Math. Phys.* **339** no. 2, (2015) 619–662, arXiv:1409.0857 [hep-th].
- [29] S. Gukov, P. Putrov, and C. Vafa, “Fivebranes and 3-manifold homology,” *JHEP* **07** (2017) 071, arXiv:1602.05302 [hep-th].
- [30] S. Gukov, D. Pei, P. Putrov, and C. Vafa, “BPS spectra and 3-manifold invariants,” arXiv:1701.06567 [hep-th].
- [31] V. Mikhaylov, “Teichmüller TQFT vs Chern-Simons Theory,” *JHEP* **04** (2018) 085, arXiv:1710.04354 [hep-th].
- [32] D. Gang and K. Yonekura, “Symmetry enhancement and closing of knots in 3d/3d correspondence,” arXiv:1803.04009 [hep-th].
- [33] T. Dimofte, M. Gabella, and A. B. Goncharov, “K-Decompositions and 3d Gauge Theories,” *JHEP* **11** (2016) 151, arXiv:1301.0192 [hep-th].
- [34] D. Gang, Y. Tachikawa, and K. Yonekura, “Smallest 3d hyperbolic manifolds via simple 3d theories,” *Phys. Rev.* **D96** no. 6, (2017) 061701, arXiv:1706.06292 [hep-th].
- [35] A. Donos, J. P. Gauntlett, N. Kim, and O. Varela, “Wrapped M5-branes, consistent truncations and AdS/CMT,” *JHEP* **12** (2010) 003, arXiv:1009.3805 [hep-th].
- [36] D. Gang, N. Kim, and S. Lee, “Holography of 3d-3d correspondence at Large N ,” *JHEP* **04** (2015) 091, arXiv:1409.6206 [hep-th].
- [37] G. D. Mostow, “Quasi-conformal mappings inn-space and the rigidity of hyperbolic space forms,” *Publications Mathématiques de l’IHÉS* **34** no. 1, (1968) 53–104.
- [38] D. Martelli, A. Passias, and J. Sparks, “The supersymmetric NUTs and bolts of holography,” *Nucl. Phys.* **B876** (2013) 810–870, arXiv:1212.4618 [hep-th].
- [39] N. A. Nekrasov and S. L. Shatashvili, “Bethe/Gauge correspondence on curved spaces,” *JHEP* **01** (2015) 100, arXiv:1405.6046 [hep-th].
- [40] H.-C. Kim and S. Kim, “Supersymmetric vacua of mass-deformed M2-brane theory,” *Nucl. Phys.* **B839** (2010) 96–111, arXiv:1001.3153 [hep-th].
- [41] K. Intriligator and N. Seiberg, “Aspects of 3d $N=2$ Chern-Simons-Matter Theories,” *JHEP* **07** (2013) 079, arXiv:1305.1633 [hep-th].
- [42] A. Kapustin, B. Willett, and I. Yaakov, “Exact Results for Wilson Loops in Superconformal Chern-Simons Theories with Matter,” *JHEP* **03** (2010) 089, arXiv:0909.4559 [hep-th].
- [43] S. Gukov, “Three-dimensional quantum gravity, Chern-Simons theory, and the A polynomial,” *Commun. Math. Phys.* **255** (2005) 577–627, arXiv:hep-th/0306165.
- [44] T. Dimofte, S. Gukov, J. Lenells, and D. Zagier, “Exact Results for Perturbative Chern-Simons Theory with Complex Gauge Group,” *Commun. Num. Theor. Phys.* **3** (2009) 363–443, arXiv:0903.2472 [hep-th].
- [45] S. Gukov and H. Murakami, “SL(2,C) Chern-Simons theory and the asymptotic behavior of the colored Jones polynomial,” *Lett. Math. Phys.* **86** (2008) 79–98, arXiv:math/0608324 [math.GT].
- [46] D. Gang, M. Romo, and M. Yamazaki, “All-Order Volume Conjecture for Closed 3-Manifolds from Complex Chern-Simons Theory,” arXiv:1704.00918 [hep-th].
- [47] D. B. Ray and I. M. Singer, “R-torsion and the laplacian on riemannian manifolds,” *Advances in Mathematics* **7** no. 2, (1971) 145–210.
- [48] S. Gukov and P. Sulkowski, “A-polynomial, B-model, and Quantization,” *JHEP* **02** (2012) 070, arXiv:1108.0002 [hep-th].
- [49] E. Witten, “Quantum field theory and the jones polynomial,” *Communications in Mathematical Physics* **121** no. 3, (1989) 351–399.
- [50] C. Closset, H. Kim, and B. Willett, “Seifert fibering operators in 3d $\mathcal{N} = 2$ theories,” arXiv:1807.02328 [hep-th].
- [51] D. Gang, V. Mikhaylov, and M. Yamazaki, “To appear;”
- [52] E. Witten, “2+ 1 dimensional gravity as an exactly soluble system,” *Nuclear Physics B* **311** no. 1, (1988) 46–78.
- [53] W. Müller, “The asymptotics of the ray-singer analytic torsion of hyperbolic 3-manifolds,” in *Metric and differential geometry*, pp. 317–352. Springer, 2012.
- [54] D. Gang, N. Kim, and S. Lee, “Holography of wrapped M5-branes and Chern–Simons theory,” *Phys. Lett.* **B733** (2014) 316–319, arXiv:1401.3595 [hep-th].
- [55] N. Hama, K. Hosomichi, and S. Lee, “SUSY Gauge Theories on Squashed Three-Spheres,” *JHEP* **05** (2011) 014, arXiv:1102.4716 [hep-th].
- [56] J.-B. Bae, D. Gang, and J. Lee, “3D $\mathcal{N} = 2$ Minimal SCFTs from Wrapped M5-Branes,” arXiv:1610.09259 [hep-th].
- [57] D. Gang and Y. Hatsuda, “S-duality resurgence in SL(2) Chern-Simons theory,” arXiv:1710.09994 [hep-th].
- [58] D. Martelli, A. Passias, and J. Sparks, “The gravity dual of supersymmetric gauge theories on a squashed three-sphere,”

Nucl. Phys. **B864** (2012) 840–868,
arXiv:1110.6400 [hep-th].