

ESTIMATES FOR ELLIPTIC SYSTEMS IN A NARROW REGION ARISING FROM COMPOSITE MATERIALS

HONGJIE JU, HAIGANG LI, AND LONGJUAN XU

ABSTRACT. In this paper, we establish the pointwise upper and lower bounds of the gradients of solutions to a class of elliptic systems, including linear systems of elasticity, in a general narrow region and in all dimensions. This problem arises from the study of damage analysis of high-contrast composite materials. Our results show that the damage may initiate from the narrowest place.

1. INTRODUCTION AND MAIN RESULTS

From the structure of fiber-reinforced composite, there are a relatively large number of fibers which are touching or nearly touching. The maximal strains can be strongly influenced by the distances between fibers. Especially, in high-contrast fiber-reinforced composites high concentration of extreme electric field or mechanical loads will occur in the narrow regions between two adjacent fibers. The purpose of this paper is to establish gradient estimates for solutions to a class of elliptic systems, including linear systems of elasticity, in such narrow regions.

A composite medium would be represented by a bounded domain Ω , divided into a finite number of subdomains. A simple two-dimensional example, which very well illustrates the main feature of our estimates, would have the domain $\Omega \subset \mathbb{R}^2$ model the cross-section of a fiber-reinforced composite, with $D_1 \cup D_2 \subset \Omega$ representing the cross-section of the fibers, the remaining subdomain $\Omega \setminus \overline{D_1 \cup D_2}$ representing the matrix surrounding the fibers. It is well known that for the scalar case, the anti-plane shear model is inconsistent with the two-dimensional conductivity model. Thus, the blow-up analysis for electric field has a valuable meaning in relation to the damage analysis of composite material. The most important quantities from an engineering point of view are $|\nabla u|$, representing the electric field in the conductivity problem or the stresses in the anti-plane shear model. Therefore, stimulated by the well-known work on damage analysis of fiber composites [9, 22, 29], there have been a number of papers, starting from [17, 26, 27], on gradient estimates for solutions of elliptic equations and systems with piecewise smooth coefficients which are relevant in such studies. See [3–8, 10, 11, 13–16, 20, 24, 28, 31, 32] and the references therein.

In order to investigate the high concentration phenomenon of high-contrast composites when $\text{dist}(D_1, D_2)$ is small, it is important to study the gradient estimate for the limiting case of a class of elliptic equations and systems with partially degenerated coefficients, that is, the coefficients in D_1 and D_2 degenerate to ∞ . In a

Date: August 13, 2018.

H.J. Ju was partially supported by NSFC (11301034) (11471050).

H.G. Li was partially supported by NSFC (11571042) (11371060) (11631002), Fok Ying Tung Education Foundation (151003). Corresponding author.




FIGURE 1. Two closely spaced balls.




FIGURE 2. A narrow region Ω_1 .

recent paper [5], some gradient estimates were obtained concerning the conductivity problem where the conductivity is allowed to be ∞ (perfect conductor).

Theorem A ([5]). Let B_1 and B_2 be two balls in \mathbb{R}^3 with radius R and centered at $(0, 0, \pm R \pm \frac{\varepsilon}{2})$, respectively, (See Figure 1). Let H be a harmonic function in \mathbb{R}^3 such that $H(0) = 0$. Define u to be the solutions of

$$\begin{cases} \Delta u = 0, & \text{in } \mathbb{R}^3 \setminus \overline{B_1 \cup B_2}, \\ u = 0, & \text{on } \partial B_1 \cup \partial B_2, \\ u(x) - H(x) = O(|x|^{-1}), & \text{as } |x| \rightarrow +\infty. \end{cases}$$

Then there exists a constant C independent of ε such that

$$\|\nabla(u - H)\|_{L^\infty(\mathbb{R}^3 \setminus \overline{B_1 \cup B_2})} \leq C.$$

Theorem A shows that when the boundary values are both zero on ∂B_1 and ∂B_2 , $|\nabla u|$ is bounded, so no concentration occurs. Theorem A was extended to the elliptic systems in [24], and was improved that $|\nabla u|$ decays exponentially fast near the origin. Later, another proof for scalar case was given in [21]. However, it is much more interesting to study the case when the boundary data are different on ∂B_1 and ∂B_2 , which more frequently appears in practical engineering applications, see [18, 22, 23, 29], where it is shown that in dimension two $|\nabla u|$ may blow up in the narrow region between B_1 and B_2 , as $\varepsilon \rightarrow 0$.

Contrary to the scalar equation, less is known on such blow-up phenomenon for the linear elasticity case. Therefore, our effort is focussed on the narrow region (see Figure 2) to investigate the gradient estimate for a class of general elliptic systems, including linear systems of elasticity.

Before stating our results, we first fix our domain. To be precise, we define a more general narrow region in all dimensions as follows: for $r \leq 1$,

$$\Omega_r := \left\{ x = (x', x_n) \in \mathbb{R}^n \mid -\frac{\varepsilon}{2} + h_2(x') < x_n < \frac{\varepsilon}{2} + h_1(x'), x' \in B_r(0') \right\},$$

where $B_r(0') := \{x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1} \mid |x'| < r\}$ is a ball in \mathbb{R}^{n-1} , centered at the origin $0'$ of radius r , h_1 and h_2 are smooth functions in $B_1(0')$ satisfying

$$\begin{aligned} -\frac{\varepsilon}{2} + h_2(x') < \frac{\varepsilon}{2} + h_1(x'), & \quad \text{for } |x'| \leq 1, \\ h_1(0') = h_2(0') = 0, \quad \nabla h_1(0') = \nabla h_2(0') = 0, & \quad (1.1) \end{aligned}$$

$$\nabla^2(h_1 - h_2)(0') \geq \kappa_0 I_{n-1} \quad (1.2)$$

and

$$\|h_1\|_{C^2(B_1(0'))} + \|h_2\|_{C^2(B_1(0'))} \leq \kappa_1, \quad (1.3)$$

where I_{n-1} is the $(n-1) \times (n-1)$ identity matrix, κ_0, κ_1 are some positive constants. We denote the top and bottom boundaries of Ω_r as

$$\Gamma_r^+ = \{x \in \mathbb{R}^n \mid x_n = \frac{\varepsilon}{2} + h_1(x'), |x'| \leq r\}, \quad \Gamma_r^- = \{x \in \mathbb{R}^n \mid x_n = -\frac{\varepsilon}{2} + h_2(x'), |x'| \leq r\},$$

respectively.

Let $u = (u^1, \dots, u^N)$ be a vector-valued function. Consider the following boundary value problem:

$$\begin{cases} \partial_\alpha \left(A_{ij}^{\alpha\beta}(x) \partial_\beta u^j + B_{ij}^\alpha(x) u^j \right) + C_{ij}^\beta(x) \partial_\beta u^j + D_{ij}(x) u^j = 0, & \text{in } \Omega_1, \\ u = \varphi(x), & \text{on } \Gamma_1^+, \\ u = \psi(x), & \text{on } \Gamma_1^-, \end{cases} \quad (1.4)$$

where $\varphi(x) = (\varphi^1(x), \varphi^2(x), \dots, \varphi^N(x)) \in C^2(\Gamma_1^+; \mathbb{R}^N)$, $\psi(x) = (\psi^1(x), \psi^2(x), \dots, \psi^N(x)) \in C^2(\Gamma_1^-; \mathbb{R}^N)$ are given vector-valued functions. Here the usual summation convention is used: α and β are summed from 1 to n , while i and j are summed from 1 to N .

The coefficients $A_{ij}^{\alpha\beta}(x)$ are measurable, bounded, that is,

$$|A_{ij}^{\alpha\beta}(x)| \leq \Lambda \quad (1.5)$$

for some constant $\Lambda > 0$ and satisfy the rather weak ellipticity condition, that is, there exists a constant $0 < \lambda < \infty$ such that

$$\int_{\Omega_1} A_{ij}^{\alpha\beta}(x) \partial_\alpha \xi^i \partial_\beta \xi^j dx \geq \lambda \int_{\Omega_1} |\nabla \xi|^2 dx, \quad \forall \xi \in H_0^1(\Omega_1; \mathbb{R}^N). \quad (1.6)$$

Recall that a system is called a system of elasticity if $N = n$, the coefficients satisfy

$$A_{ij}^{\alpha\beta}(x) = A_{ji}^{\beta\alpha}(x) = A_{\alpha j}^{i\beta}(x),$$

and for all $n \times n$ symmetric matrices ξ_α^i ,

$$\lambda |\xi| \leq A_{ij}^{\alpha\beta}(x) \xi_\alpha^i \xi_\beta^j \leq \Lambda |\xi|^2.$$

It is clear that hypotheses (1.5) and (1.6) are satisfied by the linear systems of elasticity, especially by the Lamé system, see [30],

$$\lambda \Delta u + (\lambda + \mu) \nabla(\nabla \cdot u) = 0.$$

Furthermore, we assume that

$$\|A\|_{C^2(\Omega_1)} + \|B\|_{C^2(\Omega_1)} + \|C\|_{C^2(\Omega_1)} + \|D\|_{C^2(\Omega_1)} \leq \kappa_2, \quad (1.7)$$

for some positive constant κ_2 . Throughout the paper, unless otherwise stated, we use C to denote some positive constant, whose values may vary from line to line, which depend only on $n, N, \lambda, \Lambda, \kappa_0, \kappa_1, \kappa_2$, but not on ε . Also, we call a constant having such dependence a *universal constant*.

In the paper, the main result concerns local piecewise gradient estimates of weak solutions u of problem (1.4); that is, $u \in H^1(\Omega_1; \mathbb{R}^N)$, and satisfies

$$\int_{\Omega_1} \left(A_{ij}^{\alpha\beta}(x) \partial_\beta u^j + B_{ij}^\alpha(x) u^j \right) \partial_\alpha \zeta^i - C_{ij}^\beta(x) \partial_\beta u^j \zeta^i - D_{ij}(x) u^j \zeta^i dx = 0,$$

for every vector-valued function $\zeta = (\zeta^1, \dots, \zeta^N) \in C_c^\infty(\Omega_1; \mathbb{R}^N)$, and hence for every $\zeta \in H_0^1(\Omega_1; \mathbb{R}^N)$.

Theorem 1.1. *Assume that hypotheses (1.1)–(1.3) and (1.5)–(1.7) are satisfied, and let $u \in H^1(\Omega_1; \mathbb{R}^N)$ be a weak solution of problem (1.4). Then, for $x \in \Omega_{1/2}$,*

$$\begin{aligned} |\nabla u(x', x_n)| &\leq \frac{C}{\varepsilon + |x'|^2} \left| \varphi(x', \varepsilon/2 + h_1(x')) - \psi(x', -\varepsilon/2 + h_2(x')) \right| \\ &\quad + C \left(\|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)} + \|u\|_{L^2(\Omega_1)} \right). \end{aligned} \quad (1.8)$$

Moreover, if $\varphi^l(0', \varepsilon/2) \neq \psi^l(0', -\varepsilon/2)$ for some integer l , then

$$|\nabla u(0', x_n)| \geq \frac{|\varphi^l(0', \varepsilon/2) - \psi^l(0', -\varepsilon/2)|}{C\varepsilon}, \quad \forall x_n \in \left(-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right). \quad (1.9)$$

Remark 1.2. Some remarks on Theorem 1.1 are in order.

(i) When $\varphi(x', \varepsilon/2 + h_1(x')) \equiv \psi(x', -\varepsilon/2 + h_2(x'))$, we first observe from (1.8) that $|\nabla u| \leq C$ and there is no blow-up occurring, which is consistent with the main result of [24].

(ii) On the other hand, it is easy to see that if $\varphi = a$, $\psi = b$ for two different constant vectors a and b , then $|\nabla u|$ will blow up, especially with rate ε^{-1} at the origin.

(iii) When $\varphi(0', \varepsilon/2) = \psi(0', -\varepsilon/2)$, by the Taylor expansions of φ and ψ , we have

$$\begin{aligned} |\nabla u(x', x_n)| &\leq \frac{C}{\varepsilon + |x'|^2} \left| (\nabla_{x'} \varphi(0', \varepsilon/2) - \nabla_{x'} \psi(0', -\varepsilon/2)) \cdot x' + O(|x'|^2) \right| \\ &\quad + C \left(\|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)} + \|u\|_{L^2(\Omega_1)} \right), \quad x \in \Omega_{1/2}. \end{aligned}$$

Clearly, if $\nabla_{x'} \varphi(0', \varepsilon/2) \neq \nabla_{x'} \psi(0', -\varepsilon/2)$, then $|\nabla u| \leq C$ on Ω_ε , where C is independent of ε . If $\nabla_{x'} \varphi(0', \varepsilon/2) = \nabla_{x'} \psi(0', -\varepsilon/2)$, $|\nabla u|$ is uniformly bounded on $\Omega_{1/2}$. Consequently, in the case that $\varphi(0', \varepsilon/2) = \psi(0', -\varepsilon/2)$, there is no blow-up occurring at the origin.

Theorem 1.1 gives more information about the dependence of $|\nabla u|$, which will play an important role in the study of the perfect conductivity problem (e.g. [10, 11, 25]) and Lamé system with partially infinite coefficients (e.g. [12–14]), where the coefficients in the inclusions are allowed to be ∞ .

For the convenience of further applications, we list the analog result for the conductivity problem in the narrow region as a consequence. For the boundary value problem of Laplace equation

$$\begin{cases} \Delta u = 0, & \text{in } \Omega_1, \\ u = \varphi(x), & \text{on } \Gamma_1^+, \\ u = \psi(x), & \text{on } \Gamma_1^-, \end{cases} \quad (1.10)$$

we have

Corollary 1.3. *Assume that $u \in H^1(\Omega_1)$ is a weak solution of (1.10), $\varphi(x) \in C^2(\Gamma_1^+)$, $\psi(x) \in C^2(\Gamma_1^-)$ are given functions. Then, for $x \in \Omega_{1/2}$,*

$$\begin{aligned} |\nabla u(x', x_n)| &\leq \frac{C}{\varepsilon + |x'|^2} \left| \varphi(x', \varepsilon/2 + h_1(x')) - \psi(x', -\varepsilon/2 + h_2(x')) \right| \\ &\quad + C \left(\|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)} + \|u\|_{L^2(\Omega_1)} \right). \end{aligned} \quad (1.11)$$

If $\varphi(0', \varepsilon/2) \neq \psi(0', -\varepsilon/2)$, then

$$|\nabla u(0', x_n)| \geq \frac{|\varphi(0', \varepsilon/2) - \psi(0', -\varepsilon/2)|}{C\varepsilon}, \quad \forall x_n \in \left(-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right).$$

Remark 1.2 is also true for the problem (1.10).

The paper is organized as follows. In Section 2, we use energy method and an adaptive version of Bao-Li-Li's iteration technique [13] to prove Theorem 1.1. The main differences of the proof of Corollary 1.3 with that of Theorem 1.1 are given in Section 3.

2. PROOF OF THEOREM 1.1

We decompose the solution of (1.4) as follows:

$$u = v_1 + v_2 + \cdots + v_N,$$

where $v_l = (v_l^1, v_l^2, \dots, v_l^N)$, $l = 1, 2, \dots, N$, with $v_l^j = 0$ for $j \neq l$, and v_l satisfies the following boundary value problem

$$\begin{cases} \partial_\alpha \left(A_{ij}^{\alpha\beta}(x) \partial_\beta v_l^j + B_{ij}^\alpha(x) v_l^j \right) + C_{ij}^\beta(x) \partial_\beta v_l^j + D_{ij}(x) v_l^j = 0, & \text{in } \Omega_1, \\ v_l = (0, \dots, 0, \varphi^l, 0, \dots, 0), & \text{on } \Gamma_1^+, \\ v_l = (0, \dots, 0, \psi^l, 0, \dots, 0), & \text{on } \Gamma_1^-. \end{cases} \quad (2.1)$$

Then

$$\nabla u = \sum_{l=1}^N \nabla v_l.$$

In order to estimate $|\nabla v_l|$, $l = 1, \dots, N$, we introduce a scalar function $\bar{u} \in C^2(\mathbb{R}^n)$ such that $\bar{u} = 1$ on Γ_1^+ , $\bar{u} = 0$ on Γ_1^- and

$$\bar{u}(x) = \frac{x_n - h_2(x') + \frac{\varepsilon}{2}}{\varepsilon + h_1(x') - h_2(x')}, \quad \text{in } \Omega_1. \quad (2.2)$$

By a direct calculation, we obtain that

$$|\partial_\alpha \bar{u}(x)| \leq \frac{C|x'|}{\varepsilon + |x'|^2}, \quad \alpha = 1, \dots, n-1, \quad \frac{1}{C(\varepsilon + |x'|^2)} \leq |\partial_n \bar{u}(x)| \leq \frac{C}{\varepsilon + |x'|^2}, \quad (2.3)$$

and for $\alpha, \beta = 1, \dots, n-1$,

$$|\partial_{\alpha\beta} \bar{u}(x)| \leq \frac{C}{\varepsilon + |x'|^2}, \quad |\partial_{\alpha n} \bar{u}(x)| \leq \frac{C|x'|}{(\varepsilon + |x'|^2)^2}, \quad \partial_{nn} \bar{u}(x) = 0. \quad (2.4)$$

For $l = 1, 2, \dots, N$, we define

$$\tilde{u}_l(x) = (0, \dots, 0, \varphi^l(x', \varepsilon/2 + h_1(x')) \bar{u}(x) + \psi^l(x', -\varepsilon/2 + h_2(x')) (1 - \bar{u}(x)), 0, \dots, 0). \quad (2.5)$$

Thus, in view of (2.3) and (2.4),

$$\begin{aligned} |\nabla_{x'} \tilde{u}_l(x)| &\leq \frac{C|x'|}{\varepsilon + |x'|^2} |\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))| \\ &\quad + C(\|\nabla\varphi^l\|_{L^\infty} + \|\nabla\psi^l\|_{L^\infty}), \end{aligned} \quad (2.6)$$

$$\begin{aligned} \frac{|\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))|}{C(\varepsilon + |x'|^2)} &\leq \\ |\partial_n \tilde{u}_l(x)| &\leq \frac{C|\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))|}{\varepsilon + |x'|^2}, \end{aligned} \quad (2.7)$$

and by using (2.3) and (2.4), for $\alpha, \beta = 1, \dots, n-1$,

$$\begin{aligned} |\partial_{\alpha\beta} \tilde{u}_l(x)| &\leq \frac{C}{\varepsilon + |x'|^2} |\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))| \\ &\quad + C \left(\frac{|x'|}{\varepsilon + |x'|^2} + 1 \right) (\|\nabla\varphi^l\|_{L^\infty} + \|\nabla\psi^l\|_{L^\infty}) \\ &\quad + C(\|\nabla^2\varphi^l\|_{L^\infty} + \|\nabla^2\psi^l\|_{L^\infty}), \end{aligned} \quad (2.8)$$

$$\begin{aligned} |\partial_{\alpha n} \tilde{u}_l(x)| &\leq \frac{C|x'|}{(\varepsilon + |x'|^2)^2} |\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))| \\ &\quad + \frac{C}{\varepsilon + |x'|^2} (\|\nabla\varphi^l\|_{L^\infty} + \|\nabla\psi^l\|_{L^\infty}), \end{aligned} \quad (2.9)$$

$$\partial_{nn} \tilde{u}_l(x) = 0. \quad (2.10)$$

Here and throughout the paper, for simplicity we use $\|\nabla\varphi\|_{L^\infty}$, $\|\nabla\psi\|_{L^\infty}$, $\|\nabla^2\varphi\|_{L^\infty}$ and $\|\nabla^2\psi\|_{L^\infty}$ to denote $\|\nabla\varphi\|_{L^\infty(\Gamma_1^+)}$, $\|\nabla\psi\|_{L^\infty(\Gamma_1^-)}$, $\|\nabla^2\varphi\|_{L^\infty(\Gamma_1^+)}$ and $\|\nabla^2\psi\|_{L^\infty(\Gamma_1^-)}$, respectively.

Let

$$w_l = v_l - \tilde{u}_l, \quad l = 1, \dots, N.$$

Then w satisfies

$$\begin{cases} \partial_\alpha \left(A_{ij}^{\alpha\beta}(x) \partial_\beta w^j + B_{ij}^\alpha(x) w^j \right) + C_{ij}^\beta(x) \partial_\beta w^j + D_{ij}(x) w^j = \tilde{f}^i, & \text{in } \Omega_1, \\ w = 0, & \text{on } \Gamma_1^\pm, \end{cases} \quad (2.11)$$

where

$$\begin{aligned} \tilde{f}^i &= -\partial_\alpha \left(A_{ij}^{\alpha\beta}(x) \partial_\beta \tilde{u}^j + B_{ij}^\alpha(x) \tilde{u}^j + C_{ij}^\alpha(x) \tilde{u}^j \right) \\ &\quad + \partial_\beta (C_{ij}^\beta(x) \tilde{u}^j) - D_{ij}(x) \tilde{u}^j. \end{aligned}$$

Let $\tilde{f} := (\tilde{f}^1, \dots, \tilde{f}^N)$, then it follows from (1.7) and (2.5)–(2.10) that for $(x', x_n) \in \Omega_1$,

$$\begin{aligned} |\tilde{f}(x', x_n)| &\leq C|\nabla^2 \tilde{u}(x', x_n)| + C|\nabla \tilde{u}(x', x_n)| + C|\tilde{u}(x', x_n)| \\ &\leq \left(\frac{C}{\varepsilon + |x'|^2} + \frac{C|x'|}{(\varepsilon + |x'|^2)^2} \right) |\varphi(x', \varepsilon/2 + h_1(x')) - \psi(x', -\varepsilon/2 + h_2(x'))| \\ &\quad + \left(\frac{C}{\varepsilon + |x'|^2} + \frac{C|x'|}{\varepsilon + |x'|^2} \right) (\|\nabla\varphi\|_{L^\infty} + \|\nabla\psi\|_{L^\infty}) \\ &\quad + C(\|\nabla^2\varphi\|_{L^\infty} + \|\nabla^2\psi\|_{L^\infty}), \end{aligned} \quad (2.12)$$

where C is independent of ε .

Lemma 2.1. *Let $v_l \in H^1(\Omega_1; \mathbb{R}^N)$ be a weak solution of (2.1), then*

$$\int_{\Omega_{1/2}} |\nabla w_l|^2 dx \leq C \left(\|w_l\|_{L^2(\Omega_1)}^2 + \|\varphi^l\|_{C^2(\Gamma_1^+)}^2 + \|\psi^l\|_{C^2(\Gamma_1^-)}^2 \right), \quad l = 1, \dots, N, \quad (2.13)$$

where C depends on $n, \lambda, \kappa_0, \kappa_1$ and κ_2 .

Proof. For simplicity, we assume that $\psi \equiv 0$. We only prove the case when $l = 1$ for instance. The other cases are the same. Denote

$$w := w_1, \quad \tilde{u} := \tilde{u}_1 \quad \text{and} \quad \varphi := \varphi^1.$$

Then,

$$\begin{aligned} \tilde{f}^i &= -\partial_\alpha \left(A_{i1}^{\alpha\beta}(x) \partial_\beta \tilde{u}^1 + B_{i1}^\alpha(x) \tilde{u}^1 + C_{i1}^\alpha(x) \tilde{u}^1 \right) \\ &\quad + \partial_\beta (C_{i1}^\beta(x) \tilde{u}^1 - D_{i1}(x) \tilde{u}^1), \end{aligned}$$

and it follows from (1.7) and (2.5)–(2.10) that

$$\begin{aligned} |\tilde{f}^i(x)| &\leq C (|\nabla^2 \tilde{u}^1(x)| + |\nabla \tilde{u}^1(x)| + |\tilde{u}^1(x)|) \\ &\leq C \|\varphi\|_{C^2(\Gamma_1^+)}, \quad x \in \Omega_1 \setminus \overline{\Omega_{1/4}}. \end{aligned} \quad (2.14)$$

Multiplying the equation in (2.11) by w and applying integration by parts in $\Omega_{1/2}$, we have

$$\begin{aligned} &\int_{\Omega_{1/2}} A_{ij}^{\alpha\beta}(x) \partial_\beta w^j \partial_\alpha w^i dx \\ &= - \int_{\Omega_{1/2}} B_{ij}^\alpha(x) w^j \partial_\alpha w^i dx + \int_{\Omega_{1/2}} C_{ij}^\beta(x) \partial_\beta w^j w^i dx + \int_{\Omega_{1/2}} D_{ij}(x) w^j w^i dx \\ &\quad - \int_{\Omega_{1/2}} \tilde{f}^i w^i dx + \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x') < x_n < \frac{\varepsilon}{2}+h_1(x')}} \left(A_{ij}^{\alpha\beta}(x) \partial_\beta w^j + B_{ij}^\alpha(x) w^j \right) w^i \frac{x_\alpha}{r} ds. \end{aligned}$$

Using the weak ellipticity condition and the Cauchy inequality, we obtain

$$\begin{aligned} \lambda \int_{\Omega_{1/2}} |\nabla w|^2 dx &\leq \int_{\Omega_{1/2}} A_{ij}^{\alpha\beta}(x) \partial_\beta w^j \partial_\alpha w^i dx \\ &\leq \frac{\lambda}{4} \int_{\Omega_{1/2}} |\nabla w|^2 dx + C \int_{\Omega_{1/2}} |w|^2 dx + \left| \int_{\Omega_{1/2}} \tilde{f}^i w^i dx \right| \\ &\quad + C \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x') < x_n < \frac{\varepsilon}{2}+h_1(x')}} (|\nabla w|^2 + |w|^2) ds, \end{aligned} \quad (2.15)$$

Note that $w = 0$ on Γ_1^\pm and $\overline{\Omega_{2/3}} \setminus \Omega_{1/3} \subset \left((\Omega_1 \setminus \overline{\Omega_{1/4}}) \cup (\Gamma_1^\pm \setminus \Gamma_{1/4}^\pm) \right)$, by using the Sobolev embedding theorem and classical $W^{2,p}$ estimates for elliptic systems,

we have, for some $p > n$,

$$\begin{aligned} \|\nabla w\|_{L^\infty(\Omega_{2/3} \setminus \overline{\Omega_{1/3}})} &\leq C \|w\|_{W^{2,p}(\Omega_{2/3} \setminus \overline{\Omega_{1/3}})} \\ &\leq C \left(\|w\|_{L^2(\Omega_1 \setminus \overline{\Omega_{1/4}})} + \|\tilde{f}\|_{L^\infty(\Omega_1 \setminus \overline{\Omega_{1/4}})} \right) \\ &\leq C \left(\|w\|_{L^2(\Omega_1)} + \|\varphi\|_{C^2(\Gamma_1^+)} \right), \end{aligned}$$

and for $x = (x', x_n) \in \Omega_{2/3} \setminus \overline{\Omega_{1/3}}$,

$$\begin{aligned} |w(x', x_n)| &= |w(x', x_n) - w(x', \frac{\varepsilon}{2} + h_1(x'))| \\ &\leq C(\varepsilon + |x'|^2) \|\nabla w\|_{L^\infty(\Omega_{2/3} \setminus \overline{\Omega_{1/3}})} \\ &\leq C \left(\|w\|_{L^2(\Omega_1)} + \|\varphi\|_{C^2(\Gamma_1^+)} \right). \end{aligned}$$

In particular, this implies that

$$\int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x')<x_n<\frac{\varepsilon}{2}+h_1(x')}} (|w|^2 + |\nabla w|^2) ds \leq C \left[\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2 \right], \quad (2.16)$$

where C depends only on n , λ and κ_0 .

Obviously,

$$\int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x')<x_n<\frac{\varepsilon}{2}+h_1(x')}} |\nabla_{x'} \tilde{u}|^2 ds \leq C \|\varphi\|_{C^1(\Gamma_1^+)}^2, \quad (2.17)$$

and

$$\begin{aligned} &\int_{\Omega_{1/2}} |\nabla_{x'} \tilde{u}|^2 dx \\ &\leq C \int_{|x'|<\frac{1}{2}} (\varepsilon + h_1(x') - h_2(x')) \left(\frac{|x'|^2 |\varphi|^2}{(\varepsilon + |x'|^2)^2} + \|\nabla \varphi\|_{L^\infty}^2 \right) dx' \\ &\leq C \|\varphi\|_{C^1(\Gamma_1^+)}^2. \end{aligned} \quad (2.18)$$

Applying integration by parts and making use of (2.10) and (2.16)–(2.18), we have

$$\begin{aligned} &\left| \int_{\Omega_{1/2}} \tilde{f}^i w^i dx \right| \\ &\leq \left| \int_{\Omega_{1/2}} \sum_{\alpha+\beta<2n} A_{i1}^{\alpha\beta} w^i \partial_{\alpha\beta} \tilde{u}^1 dx \right| + \left| \int_{\Omega_{1/2}} \partial_\alpha A_{i1}^{\alpha\beta} w^i \partial_\beta \tilde{u}^1 dx \right| + \left| \int_{\Omega_{1/2}} (B_{i1}^\alpha + C_{i1}^\alpha) \tilde{u}^1 \partial_\alpha w^i dx \right| \\ &\quad + \left| \int_{\Omega_{1/2}} (\partial_\beta (C_{i1}^\beta(x)) - D_{i1}(x)) \tilde{u}^1 w^i dx \right| + \left| \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x')<x_n<\frac{\varepsilon}{2}+h_1(x')}} (B_{i1}^\alpha + C_{i1}^\alpha) \tilde{u}^1 w^i \frac{x_\alpha}{r} ds \right| \end{aligned}$$

$$\begin{aligned}
&\leq C \int_{\Omega_{1/2}} |\nabla_{x'} \tilde{u}| |\nabla w| dx + C \int_{\Omega_{1/2}} |\nabla_{x'} \tilde{u}| |w| dx + C \int_{\Omega_{1/2}} |\tilde{u}| |\nabla w| dx \\
&\quad + C \int_{\Omega_{1/2}} |\tilde{u}| |w| dx + C \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x')<x_n<\frac{\varepsilon}{2}+h_1(x')}} (|\nabla_{x'} \tilde{u}| |w| + |\tilde{u}| |w|) ds \\
&\leq C \left(\int_{\Omega_{1/2}} |\nabla_{x'} \tilde{u}|^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega_{1/2}} |\nabla w|^2 dx \right)^{\frac{1}{2}} + C \left(\int_{\Omega_{1/2}} |\nabla_{x'} \tilde{u}|^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega_{1/2}} |w|^2 dx \right)^{\frac{1}{2}} \\
&\quad + C \left(\int_{\Omega_{1/2}} |\tilde{u}|^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega_{1/2}} |\nabla w|^2 dx \right)^{\frac{1}{2}} + C \left(\int_{\Omega_{1/2}} |\tilde{u}|^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega_{1/2}} |w|^2 dx \right)^{\frac{1}{2}} \\
&\quad + C \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x')<x_n<\frac{\varepsilon}{2}+h_1(x')}} (|\nabla_{x'} \tilde{u}|^2 + |u|^2) ds + C \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x')<x_n<\frac{\varepsilon}{2}+h_1(x')}} |w|^2 ds \\
&\leq C \|\varphi\|_{C^1(\Gamma_1^+)} \left(\int_{\Omega_{1/2}} |\nabla w|^2 dx \right)^{\frac{1}{2}} + C (\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2) \\
&\leq \frac{\lambda}{4} \int_{\Omega_{1/2}} |\nabla w|^2 dx + C (\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2). \tag{2.19}
\end{aligned}$$

Inserting (2.16) and (2.19) to (2.15), we obtain that

$$\int_{\Omega_{1/2}} |\nabla w|^2 dx \leq C (\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2).$$

Lemma 2.1 is established. \square

Denote

$$\delta(x') := \varepsilon + h_1(x') - h_2(x').$$

By (1.2) and (1.3), we have

$$\frac{1}{C} (\varepsilon + |x'|^2) \leq \delta(x') \leq C (\varepsilon + |x'|^2). \tag{2.20}$$

For $x_0 \in \Omega_{1/2}$, we set

$$\widehat{\Omega}_s(x_0) := \{ x \in \Omega_{1/2} \mid |x' - x'_0| < s \}, \quad \forall 0 \leq s \leq 1/2. \tag{2.21}$$

Lemma 2.2. For $0 \leq |x'_0| \leq \sqrt{\varepsilon}$,

$$\begin{aligned}
\int_{\widehat{\Omega}_\delta(x_0)} |\nabla w_l|^2 dx &\leq C \varepsilon^{n-1} [|\varphi^l(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi^l(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \\
&\quad + \varepsilon (\|\varphi^l\|_{C^2(\Gamma_1^+)}^2 + \|\psi^l\|_{C^2(\Gamma_1^-)}^2 + \|w_l\|_{L^2(\Omega_1)}^2)]; \tag{2.22}
\end{aligned}$$

and for $\sqrt{\varepsilon} < |x'_0| < \frac{1}{2}$,

$$\begin{aligned}
\int_{\widehat{\Omega}_\delta(x_0)} |\nabla w_l|^2 dx &\leq C |x'_0|^{2(n-1)} [|\varphi^l(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi^l(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \\
&\quad + |x'_0|^2 (\|\varphi^l\|_{C^2(\Gamma_1^+)}^2 + \|\psi^l\|_{C^2(\Gamma_1^-)}^2 + \|w_l\|_{L^2(\Omega_1)}^2)], \tag{2.23}
\end{aligned}$$

where $\delta = \delta(x'_0)$, $l = 1, \dots, N$.

Proof. We still assume that $\psi \equiv 0$ and only prove the case when $l = 1$ for instance, and denote $w := w_1$, $\tilde{u} := \tilde{u}_1$ and $\varphi := \varphi^1$.

For $0 < t < s < 1$, let $\eta(x')$ be a smooth function satisfying $0 \leq \eta(x') \leq 1$, $\eta(x') = 1$ if $|x' - x'_0| < t$, $\eta(x') = 0$ if $|x' - x'_0| > s$ and $|\nabla\eta(x')| \leq \frac{2}{s-t}$. Multiplying $\eta^2 w$ on both sides of the equation in (2.11) and applying integration by parts, we have

$$\begin{aligned} & - \int_{\widehat{\Omega}_s(x_0)} (A_{ij}^{\alpha\beta}(x) \partial_\beta w^j + B_{ij}^\alpha(x) w^j) \partial_\alpha (\eta^2 w^i) dx \\ & + \int_{\widehat{\Omega}_s(x_0)} C_{ij}^\beta(x) \partial_\beta w^j \eta^2 w^i dx + \int_{\widehat{\Omega}_s(x_0)} D_{ij}(x) w^j \eta^2 w^i dx = \int_{\widehat{\Omega}_s(x_0)} \tilde{f}^i \eta^2 w^i dx. \end{aligned}$$

Since

$$\begin{aligned} & \int_{\widehat{\Omega}_s(x_0)} (A_{ij}^{\alpha\beta}(x) \partial_\beta w^j + B_{ij}^\alpha(x) w^j) \partial_\alpha (\eta^2 w^i) dx \\ & = \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) \partial_\beta (\eta w^j) \partial_\alpha (\eta w^i) dx - \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) (\partial_\beta \eta w^j) \partial_\alpha (\eta w^i) dx \\ & + \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) \partial_\beta (\eta w^j) (\partial_\alpha \eta w^i) dx - \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) (\partial_\beta \eta w^j) (\partial_\alpha \eta w^i) dx, \\ & + \int_{\widehat{\Omega}_s(x_0)} B_{ij}^\alpha(x) (\eta w^j) \partial_\alpha (\eta w^i) dx + \int_{\widehat{\Omega}_s(x_0)} B_{ij}^\alpha(x) (\partial_\alpha \eta w^j) (\eta w^i) dx, \end{aligned}$$

and

$$\begin{aligned} & \int_{\widehat{\Omega}_s(x_0)} C_{ij}^\beta(x) \partial_\beta w^j \eta^2 w^i dx \\ & = \int_{\widehat{\Omega}_s(x_0)} C_{ij}^\beta(x) \partial_\beta (\eta w^j) (\eta w^i) dx - \int_{\widehat{\Omega}_s(x_0)} C_{ij}^\beta(x) (\partial_\beta \eta w^j) (\eta w^i) dx, \end{aligned}$$

by using the weak ellipticity condition (1.6) and the Cauchy inequality, we have

$$\begin{aligned} & \lambda \int_{\widehat{\Omega}_s(x_0)} |\nabla(\eta w)|^2 dx \leq \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) \partial_\beta (\eta w^j) \partial_\alpha (\eta w^i) dx \\ & = \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) (\partial_\beta \eta w^j) \partial_\alpha (\eta w^i) dx - \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) \partial_\beta (\eta w^j) (\partial_\alpha \eta w^i) dx \\ & + \int_{\widehat{\Omega}_s(x_0)} A_{ij}^{\alpha\beta}(x) (\partial_\beta \eta w^j) (\partial_\alpha \eta w^i) dx - \int_{\widehat{\Omega}_s(x_0)} B_{ij}^\alpha(x) (\eta w^j) \partial_\alpha (\eta w^i) dx \\ & - \int_{\widehat{\Omega}_s(x_0)} B_{ij}^\alpha(x) (\partial_\alpha \eta w^j) (\eta w^i) dx + \int_{\widehat{\Omega}_s(x_0)} C_{ij}^\beta(x) \partial_\beta (\eta w^j) (\eta w^i) dx \\ & - \int_{\widehat{\Omega}_s(x_0)} C_{ij}^\beta(x) (\partial_\beta \eta w^j) (\eta w^i) dx + \int_{\widehat{\Omega}_s(x_0)} D_{ij}(x) (\eta w^j) (\eta w^i) dx - \int_{\widehat{\Omega}_s(x_0)} \eta^2 \tilde{f}^i w^i dx \\ & \leq \frac{\lambda}{2} \int_{\widehat{\Omega}_s(x_0)} |\nabla(\eta w)|^2 dx + C \int_{\widehat{\Omega}_s(x_0)} |(\nabla\eta)w|^2 dx + \frac{C}{(s-t)^2} \int_{\widehat{\Omega}_s(x_0)} |w|^2 dx \\ & + (s-t)^2 \int_{\widehat{\Omega}_s(x_0)} |\tilde{f}|^2 dx \\ & \leq \frac{\lambda}{2} \int_{\widehat{\Omega}_s(x_0)} |\nabla(\eta w)|^2 dx + \frac{C}{(s-t)^2} \int_{\widehat{\Omega}_s(x_0)} |w|^2 dx + (s-t)^2 \int_{\widehat{\Omega}_s(x_0)} |\tilde{f}|^2 dx. \end{aligned}$$

Thus, we obtain

$$\int_{\widehat{\Omega}_t(x_0)} |\nabla w|^2 dx \leq \frac{C}{(s-t)^2} \int_{\widehat{\Omega}_s(x_0)} |w|^2 dx + C(s-t)^2 \int_{\widehat{\Omega}_s(x_0)} |\tilde{f}|^2 dx. \quad (2.24)$$

Note that $w = 0$ on Γ_1^- , by (1.3) and the Hölder inequality, we obtain

$$\begin{aligned} \int_{\widehat{\Omega}_s(x_0)} |w|^2 dx &= \int_{\widehat{\Omega}_s(x_0)} \left| \int_{-\frac{\varepsilon}{2}+h_2(x')}^{x_n} \partial_n w(x', x_n) dx_n \right|^2 dx \\ &\leq \int_{\widehat{\Omega}_s(x_0)} (\varepsilon + h_1(x') - h_2(x')) \int_{-\frac{\varepsilon}{2}+h_2(x')}^{\frac{\varepsilon}{2}+h_1(x')} |\nabla w|^2 dx_n dx \\ &\leq \int_{|x'-x'_0|<s} C(\varepsilon + |x'|)^2 \int_{-\frac{\varepsilon}{2}+h_2(x')}^{\frac{\varepsilon}{2}+h_1(x')} |\nabla w|^2 dx_n dx'. \end{aligned} \quad (2.25)$$

It follows from (2.12) and the mean value theorem that

$$\begin{aligned} &\int_{\widehat{\Omega}_s(x_0)} |\tilde{f}|^2 dx \\ &\leq |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 \int_{\widehat{\Omega}_s(x_0)} \left(\frac{C}{\varepsilon + |x'|^2} + \frac{C|x'|}{(\varepsilon + |x'|^2)^2} \right)^2 dx \\ &\quad + \|\nabla \varphi\|_{L^\infty}^2 \int_{\widehat{\Omega}_s(x_0)} \left(\frac{C}{\varepsilon + |x'|^2} + \frac{C|x'|}{(\varepsilon + |x'|^2)^2} \right)^2 |x' - x'_0|^2 dx \\ &\quad + \|\nabla \varphi\|_{L^\infty}^2 \int_{\widehat{\Omega}_s(x_0)} \left(\frac{C}{\varepsilon + |x'|^2} + \frac{C|x'|}{\varepsilon + |x'|^2} \right)^2 dx + Cs^{n-1} \|\nabla^2 \varphi\|_{L^\infty}^2 \\ &\leq C|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 \int_{|x'-x'_0|<s} \frac{1}{(\varepsilon + |x'|^2)^2} dx' \\ &\quad + C\|\nabla \varphi\|_{L^\infty}^2 \int_{|x'-x'_0|<s} \left(\frac{1}{\varepsilon + |x'|^2} + \frac{|x' - x'_0|^2}{(\varepsilon + |x'|^2)^2} \right) dx' + Cs^{n-1} \|\nabla^2 \varphi\|_{L^\infty}^2. \end{aligned} \quad (2.26)$$

Case 1. For $0 \leq |x'_0| \leq \sqrt{\varepsilon}$, $0 < t < s < \sqrt{\varepsilon}$, from (2.25) and (2.26), we have

$$\int_{\widehat{\Omega}_s(x_0)} |w|^2 dx \leq C\varepsilon^2 \int_{\widehat{\Omega}_s(x_0)} |\nabla w|^2 dx, \quad (2.27)$$

and

$$\begin{aligned} &\int_{\widehat{\Omega}_s(x_0)} |\tilde{f}|^2 dx \\ &\leq C|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 \frac{s^{n-1}}{\varepsilon^2} + C\|\nabla \varphi\|_{L^\infty}^2 \frac{s^{n-1}}{\varepsilon} + Cs^{n-1} \|\nabla^2 \varphi\|_{L^\infty}^2. \end{aligned} \quad (2.28)$$

Denote

$$F(t) := \int_{\widehat{\Omega}_t(x_0)} |\nabla w|^2 dx.$$

By (2.24), (2.27) and (2.28), for some universal constant $C_1 > 0$, we have for $0 < t < s < \sqrt{\varepsilon}$,

$$F(t) \leq \left(\frac{C_1 \varepsilon}{s-t} \right)^2 F(s) + C(s-t)^2 s^{n-1} \cdot \left(\frac{|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2}{\varepsilon^2} + \frac{\|\nabla \varphi\|_{L^\infty}^2}{\varepsilon} + \|\nabla^2 \varphi\|_{L^\infty}^2 \right). \quad (2.29)$$

Let $t_i = \delta + 2C_1 i \varepsilon$, $i = 0, 1, \dots$ and $k = \left\lceil \frac{1}{4C_1 \sqrt{\varepsilon}} \right\rceil + 1$, then

$$\frac{C_1 \varepsilon}{t_{i+1} - t_i} = \frac{1}{2}.$$

Using (2.29) with $s = t_{i+1}$ and $t = t_i$, we obtain that, for $i = 0, 1, 2, \dots, k$,

$$F(t_i) \leq \frac{1}{4} F(t_{i+1}) + C(i+1)^{n-1} \varepsilon^{n-1} \left(|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + \varepsilon \|\varphi\|_{C^2(\Gamma_1^+)}^2 \right).$$

After k iterations, making use of (2.13), we have, for sufficiently small ε ,

$$\begin{aligned} F(t_0) &\leq \left(\frac{1}{4} \right)^k F(t_k) + C \sum_{i=1}^k \left(\frac{1}{4} \right)^{i-1} i^{n-1} \varepsilon^{n-1} \left(|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + \varepsilon \|\varphi\|_{C^2(\Gamma_1^+)}^2 \right) \\ &\leq \left(\frac{1}{4} \right)^k F(\sqrt{\varepsilon}) + C \varepsilon^{n-1} \left(|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + \varepsilon \|\varphi\|_{C^2(\Gamma_1^+)}^2 \right) \\ &\leq C \varepsilon^{n-1} \left[|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + \varepsilon (\|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|w\|_{L^2(\Omega_1)}^2) \right], \end{aligned}$$

here we used that the first term in the last but one line decays exponentially, which implies that for $0 \leq |x'_0| \leq \sqrt{\varepsilon}$,

$$\|\nabla w\|_{L^2(\widehat{\Omega}_s(x_0))}^2 \leq C \varepsilon^{n-1} \left[|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + \varepsilon (\|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|w\|_{L^2(\Omega_1)}^2) \right].$$

Case 2. For $\sqrt{\varepsilon} \leq |x'_0| < \frac{1}{2}$, $0 < t < s < \frac{2|x'_0|}{3}$, by (2.25) and (2.26), we have

$$\int_{\widehat{\Omega}_s(x_0)} |w|^2 dx \leq C |x'_0|^4 \int_{\widehat{\Omega}_s(x_0)} |\nabla w|^2 dx,$$

and

$$\begin{aligned} &\int_{\widehat{\Omega}_s(x_0)} |\tilde{f}|^2 dx \\ &\leq C |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 \frac{s^{n-1}}{|x'_0|^4} + C \|\nabla \varphi\|_{L^\infty}^2 \frac{s^{n-1}}{|x'_0|^2} + C s^{n-1} \|\nabla^2 \varphi\|_{L^\infty}^2. \end{aligned}$$

Thus, we obtain that, for $0 < t < s < \frac{2|x'_0|}{3}$,

$$F(t) \leq \left(\frac{C_2 |x'_0|^2}{s-t} \right)^2 F(s) + C(s-t)^2 s^{n-1} \cdot \left(\frac{|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2}{|x'_0|^4} + \frac{\|\nabla \varphi\|_{L^\infty}^2}{|x'_0|^2} + \|\nabla^2 \varphi\|_{L^\infty}^2 \right), \quad (2.30)$$

for some universal constant $C_2 > 0$. Taking the same iteration procedure as Case 1, set $t_i = \delta + 2C_2 i |x'_0|^2$, $i = 0, 1, \dots$ and $k = \left\lceil \frac{1}{4C_2 |x'_0|} \right\rceil + 1$, by (2.30) with $s = t_{i+1}$

and $t = t_i$, we have, for $i = 0, 1, 2, \dots, k$,

$$F(t_i) \leq \frac{1}{4}F(t_{i+1}) + C(i+1)^{n-1}|x'_0|^{2(n-1)} \left(|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + |x'_0|^2 \|\varphi\|_{C^2(\Gamma_1^+)}^2 \right).$$

Similarly, after k iterations, we have

$$\begin{aligned} F(t_0) &\leq \left(\frac{1}{4}\right)^k F(t_k) + C \sum_{i=1}^k \left(\frac{1}{4}\right)^{i-1} i^{n-1} |x'_0|^{2(n-1)} \\ &\quad \left(|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + |x'_0|^2 \|\varphi\|_{C^2(\Gamma_1^+)}^2 \right) \\ &\leq \left(\frac{1}{4}\right)^k F(t_k) + C |x'_0|^{2(n-1)} \left(|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + |x'_0|^2 \|\varphi\|_{C^2(\Gamma_1^+)}^2 \right) \\ &\leq C |x'_0|^{2(n-1)} \left[|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + |x'_0|^2 (\|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|w\|_{L^2(\Omega_1)}^2) \right], \end{aligned}$$

which implies that, for $\sqrt{\varepsilon} \leq |x'_0| < \frac{1}{2}$,

$$\begin{aligned} &\|\nabla w\|_{L^2(\widehat{\Omega}_\delta(x_0))}^2 \\ &\leq C |x'_0|^{2(n-1)} \left[|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))|^2 + |x'_0|^2 (\|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|w\|_{L^2(\Omega_1)}^2) \right]. \end{aligned}$$

The proof of Lemma 2.2 is completed. \square

Lemma 2.3. For $l = 1, \dots, N$, if $|x'| \leq \sqrt{\varepsilon}$,

$$\begin{aligned} |\nabla w_l(x)| &\leq \frac{C|\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))|}{\sqrt{\varepsilon}} \\ &\quad + C \left(\|\varphi^l\|_{C^2(\Gamma_1^+)} + \|\psi^l\|_{C^2(\Gamma_1^-)} + \|w_l\|_{L^2(\Omega_1)} \right), \end{aligned} \quad (2.31)$$

and if $\sqrt{\varepsilon} < |x'| < R_0$,

$$\begin{aligned} |\nabla w_l(x)| &\leq \frac{C|\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))|}{|x'|} \\ &\quad + C \left(\|\varphi^l\|_{C^2(\Gamma_1^+)} + \|\psi^l\|_{C^2(\Gamma_1^-)} + \|w_l\|_{L^2(\Omega_1)} \right). \end{aligned} \quad (2.32)$$

Consequently, by (2.6) and (2.7), we have for sufficiently small ε and $x \in \Omega_{R_0}$,

$$\begin{aligned} |\nabla v_l(x)| &\leq \frac{C|\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))|}{\varepsilon + |x'|^2} \\ &\quad + C \left(\|\varphi^l\|_{C^2(\Gamma_1^+)} + \|\psi^l\|_{C^2(\Gamma_1^-)} + \|v_l\|_{L^2(\Omega_1)} \right). \end{aligned} \quad (2.33)$$

Moreover, if $\varphi^l(0', \frac{\varepsilon}{2}) \neq \psi^l(0', -\frac{\varepsilon}{2})$, then

$$|\nabla v_l(0', x_n)| \geq \frac{|\varphi^l(0', \frac{\varepsilon}{2}) - \psi^l(0', -\frac{\varepsilon}{2})|}{C\varepsilon}, \quad \forall x_n \in \left(-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right).$$

Proof. Take the case when $\psi \equiv 0$ and $l = 1$ for instance, and denote $v := v_1$, $w := w_1$, $\tilde{u} := \tilde{u}_1$ and $\varphi := \varphi^1$. Given $x_0 = (x'_0, x_{0n}) \in \Omega_{R_0}$, making a change of variables

$$\begin{cases} x' - x'_0 = \delta y', \\ x_n = \delta y_n. \end{cases} \quad (2.34)$$

Define

$$\hat{h}_1(y') := \frac{1}{\delta} \left(\frac{\varepsilon}{2} + h_1(\delta y' + x'_0) \right), \quad \hat{h}_2(y') := \frac{1}{\delta} \left(-\frac{\varepsilon}{2} + h_2(\delta y' + x'_0) \right).$$

Then, the region $\hat{\Omega}_\delta(x_0)$ becomes Q_1 , where

$$Q_r = \{y \in \mathbb{R}^n \mid \hat{h}_2(y') < y_n < \hat{h}_1(y'), \ |y'| < r\}, \quad 0 < r \leq 1,$$

and the top and bottom boundaries of Q_r become

$$\hat{\Gamma}_r^+ := \{y \in \mathbb{R}^n \mid y_n = \hat{h}_1(y'), \ |y'| \leq r\}$$

and

$$\hat{\Gamma}_r^- := \{y \in \mathbb{R}^n \mid y_n = \hat{h}_2(y'), \ |y'| \leq r\},$$

respectively. From (1.3) and the definition of \hat{h}_1 and \hat{h}_2 , we have

$$\hat{h}_1(0') - \hat{h}_2(0') = 1,$$

and for $|y'| < 1$,

$$|\nabla \hat{h}_1(y')| + |\nabla \hat{h}_2(y')| \leq C(\delta + |x'_0|), \quad |\nabla^2 \hat{h}_1(y')| + |\nabla^2 \hat{h}_2(y')| \leq C\delta.$$

Since R_0 is small, $\|\hat{h}_1\|_{C^{1,1}((-1,1))}$ and $\|\hat{h}_2\|_{C^{1,1}((-1,1))}$ are small and Q_1 is essentially a unit square as far as applications of Sobolev embedding theorems and classical L^p estimates for elliptic systems are concerned.

Let

$$\hat{u}(y', y_n) = \tilde{u}(\delta y' + x'_0, \delta y_n), \quad \hat{w}(y', y_n) = w(\delta y' + x'_0, \delta y_n).$$

Thus, $\hat{w}(y)$ satisfies

$$\begin{cases} \partial_\alpha \left(\hat{A}_{ij}^{\alpha\beta} \partial_\beta \hat{w}^j + \hat{B}_{ij}^\alpha \hat{w}^j \right) + \hat{C}_{ij}^\beta \partial_\beta \hat{w}^j + \hat{D}_{ij} \hat{w}^j = \hat{f}_i, & \text{in } Q_1, \\ \hat{w} = 0, & \text{on } \hat{\Gamma}_1^\pm, \end{cases}$$

where

$$\begin{aligned} \hat{A}(y) &= A(\delta y' + x'_0, \delta y_n), & \hat{B}(y) &= \delta B(\delta y' + x'_0, \delta y_n), \\ \hat{C}(y) &= \delta C(\delta y' + x'_0, \delta y_n), & \hat{D}(y) &= \delta^2 D(\delta y' + x'_0, \delta y_n), \end{aligned} \quad (2.35)$$

and $\hat{f}_i := -\partial_\alpha \left(\hat{A}_{ij}^{\alpha\beta} \partial_\beta \hat{w}^j + \hat{B}_{ij}^\alpha \hat{w}^j \right) - \hat{C}_{ij}^\beta \partial_\beta \hat{w}^j - \hat{D}_{ij} \hat{w}^j$.

In view of $\hat{w} = 0$ on the upper and lower boundaries of Q_1 , we have, by Poincaré inequality, that

$$\|\hat{w}\|_{H^1(Q_1)} \leq C \|\nabla \hat{w}\|_{L^2(Q_1)}.$$

Using the Sobolev embedding theorem and classical $W^{2,p}$ estimates for elliptic systems, we have, for some $p > n$,

$$\|\nabla \hat{w}\|_{L^\infty(Q_{1/2})} \leq C \|\hat{w}\|_{W^{2,p}(Q_{1/2})} \leq C \left(\|\nabla \hat{w}\|_{L^2(Q_1)} + \|\hat{f}\|_{L^\infty(Q_1)} \right).$$

Since

$$\|\nabla \hat{w}\|_{L^\infty(Q_{1/2})} = \delta \|\nabla w\|_{L^\infty(\hat{\Omega}_{\delta/2}(x_0))}, \quad \|\nabla \hat{w}\|_{L^2(Q_1)} = \delta^{1-\frac{n}{2}} \|\nabla w\|_{L^2(\hat{\Omega}_\delta(x_0))}$$

and

$$\|\hat{f}\|_{L^\infty(Q_1)} = \delta^2 \|\tilde{f}\|_{L^\infty(\hat{\Omega}_\delta(x_0))},$$

tracing back to w through the transforms, we have

$$\|\nabla w\|_{L^\infty(\hat{\Omega}_{\delta/2}(x_0))} \leq \frac{C}{\delta} \left(\delta^{1-\frac{n}{2}} \|\nabla w\|_{L^2(\hat{\Omega}_\delta(x_0))} + \delta^2 \|\tilde{f}\|_{L^\infty(\hat{\Omega}_\delta(x_0))} \right).$$

Case 1. For $0 \leq |x'_0| \leq \sqrt{\varepsilon}$.

By (2.12) and (2.22), we have

$$\begin{aligned} & \delta^{-\frac{n}{2}} \|\nabla w\|_{L^2(\widehat{\Omega}_\delta(x_0))} \\ & \leq \frac{C}{\sqrt{\varepsilon}} \left(\frac{\varepsilon}{\delta}\right)^{\frac{n}{2}} \left[|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + \sqrt{\varepsilon} (\|\varphi\|_{C^2(\Gamma_1^+)} + \|w\|_{L^2(\Omega_1)}) \right] \\ & \leq \frac{C}{\sqrt{\varepsilon}} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + C (\|\varphi\|_{C^2(\Gamma_1^+)} + \|w\|_{L^2(\Omega_1)}) \end{aligned}$$

and

$$\delta \|\tilde{f}\|_{L^\infty(\widehat{\Omega}_\delta(x_0))} \leq \frac{C}{\sqrt{\varepsilon}} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + C (\|\nabla \varphi\|_{L^\infty} + \|\nabla^2 \varphi\|_{L^\infty}).$$

Therefore,

$$\|\nabla w\|_{L^\infty(\widehat{\Omega}_{\delta/2}(x_0))} \leq \frac{C}{\sqrt{\varepsilon}} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + C (\|\varphi\|_{C^2(\Gamma_1^+)} + \|w\|_{L^2(\Omega_1)}).$$

(2.31) is proved.

Case 2. For $\sqrt{\varepsilon} \leq |x'_0| \leq R_0$.

Using (2.12) and (2.23), we obtain

$$\begin{aligned} & \delta^{-\frac{n}{2}} \|\nabla w\|_{L^2(\widehat{\Omega}_\delta(x_0))} \\ & \leq \frac{C}{|x'_0|} \left(\frac{|x'_0|^2}{\delta}\right)^{\frac{n}{2}} \left[|\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + |x'_0| (\|\varphi\|_{C^2(\Gamma_1^+)} + \|w\|_{L^2(\Omega_1)}) \right] \\ & \leq \frac{C}{|x'_0|} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + C (\|\varphi\|_{C^2(\Gamma_1^+)} + \|w\|_{L^2(\Omega_1)}), \end{aligned}$$

and

$$\delta \|\tilde{f}\|_{L^\infty(\widehat{\Omega}_\delta(x_0))} \leq \frac{C}{|x'_0|} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + C (\|\nabla \varphi\|_{L^\infty} + \|\nabla^2 \varphi\|_{L^\infty}).$$

Therefore,

$$\|\nabla w\|_{L^\infty(\widehat{\Omega}_{\delta/2}(x_0))} \leq \frac{C}{|x'_0|} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0))| + C (\|\varphi\|_{C^2(\Gamma_1^+)} + \|w\|_{L^2(\Omega_1)}).$$

(2.32) is proved.

Notice that $|\nabla v| \leq |\nabla w| + |\nabla \tilde{u}|$, by (2.6), (2.7), (2.31) and (2.32), we obtain (2.33). By the Taylor expansion and (1.1), we have

$$\varphi^l(x', \frac{\varepsilon}{2} + h_1(x')) = \varphi^l(0', \frac{\varepsilon}{2}) + \nabla_{x'} \varphi^l(0', \frac{\varepsilon}{2}) \cdot x' + O(|x'|^2). \quad (2.36)$$

It is clear that if $\varphi^l(0', \frac{\varepsilon}{2}) \neq 0$, then

$$|\nabla v_l(0', x_n)| \geq \frac{|\varphi^l(0', \frac{\varepsilon}{2})|}{C\varepsilon}, \quad \forall x_n \in \left(-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right).$$

The proof of Lemma 2.3 is finished. \square

Proof of Theorem 1.1. By Lemma 2.1–Lemma 2.3, we have, for $x \in \Omega_{R_0}$,

$$\begin{aligned} |\nabla u(x)| &\leq \sum_{l=1}^N |\nabla v_l| \\ &\leq \frac{C|\varphi^l(x', \varepsilon/2 + h_1(x')) - \psi^l(x', -\varepsilon/2 + h_2(x'))|}{\varepsilon + |x'|^2} \\ &\quad + C \left(\|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)} + \|u\|_{L^2(\Omega_1)} \right). \end{aligned}$$

Applying the standard elliptic theorem (see Agmon et al. [1] and [2]), we have

$$\|\nabla u\|_{L^\infty(\Omega_{1/2} \setminus \Omega_{R_0})} \leq C \left(\|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)} + \|u\|_{L^2(\Omega_1)} \right).$$

If $\varphi^l(0', \frac{\varepsilon}{2}) \neq \psi^l(0', -\frac{\varepsilon}{2})$ for some integer l , then by Lemma 2.3, we obtain

$$|\nabla u(0', x_n)| \geq \frac{|\varphi^l(0', \frac{\varepsilon}{2}) - \psi^l(0', -\frac{\varepsilon}{2})|}{C\varepsilon}, \quad \forall x_n \in \left(-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right).$$

The proof of Theorem 1.1 is completed. \square

3. PROOF OF COROLLARY 1.3

Since the problem (1.10) has much more applications in practice, such as conductivity problem, anti-plane shear model. We give a sketched proof of Corollary 1.3 and only list its main ingredients. We use the auxillary scalar function $\bar{u} \in C^2(\mathbb{R}^n)$ introduced in Section 2, which satisfying (2.2). Define

$$\tilde{u}(x) = \varphi(x', \frac{\varepsilon}{2} + h_1(x'))\bar{u} + \psi(x', -\frac{\varepsilon}{2} + h_2(x'))(1 - \bar{u}),$$

then

$$\begin{aligned} |\nabla_{x'} \tilde{u}(x)| &\leq \frac{C|x'|}{\varepsilon + |x'|^2} |\varphi(x', \varepsilon/2 + h_1(x')) - \psi(x', -\varepsilon/2 + h_2(x'))| \\ &\quad + C(\|\nabla \varphi\|_{L^\infty} + \|\nabla \psi\|_{L^\infty}), \end{aligned} \quad (3.1)$$

$$\begin{aligned} \frac{|\varphi(x', \varepsilon/2 + h_1(x')) - \psi(x', -\varepsilon/2 + h_2(x'))|}{C(\varepsilon + |x'|^2)} &\leq \\ |\partial_n \tilde{u}(x)| &\leq \frac{C|\varphi(x', \varepsilon/2 + h_1(x')) - \psi(x', -\varepsilon/2 + h_2(x'))|}{\varepsilon + |x'|^2}, \end{aligned} \quad (3.2)$$

and

$$\begin{aligned} |\partial_{\alpha\alpha} \tilde{u}(x)| &\leq \frac{C}{\varepsilon + |x'|^2} |\varphi(x', \varepsilon/2 + h_1(x')) - \psi(x', -\varepsilon/2 + h_2(x'))| \\ &\quad + C \left(\frac{|x'|}{\varepsilon + |x'|^2} + 1 \right) (\|\nabla \varphi\|_{L^\infty} + \|\nabla \psi\|_{L^\infty}) \\ &\quad + C(\|\nabla^2 \varphi\|_{L^\infty} + \|\nabla^2 \psi\|_{L^\infty}), \quad \alpha = 1, \dots, n-1, \end{aligned} \quad (3.3)$$

$$\partial_{nn} \tilde{u}(x) = 0. \quad (3.4)$$

Denote $w = u - \tilde{u}$, which satisfies the following boundary value problem

$$\begin{cases} \Delta w = -\Delta \tilde{u}, & \text{in } \Omega_1, \\ w = 0, & \text{on } \Gamma_1^\pm. \end{cases} \quad (3.5)$$

Step 1. Boundedness of $\int_{\Omega_{1/2}} |\nabla w|^2 dx$.

Multiplying the equation in (3.5) by w and applying integration by parts on $\Omega_{1/2}$, in view of $w = 0$ on Γ_1^\pm , we have

$$\begin{aligned}
& \int_{\Omega_{1/2}} |\nabla w|^2 dx \\
&= \int_{\Omega_{1/2}} \sum_{\alpha=1}^{n-1} w \partial_{\alpha\alpha} \tilde{u} dx + \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x') < x_n < \frac{\varepsilon}{2}+h_1(x')}} \sum_{\alpha=1}^{n-1} w \partial_{\alpha} w \frac{x_{\alpha}}{r} ds \\
&= - \int_{\Omega_{1/2}} \nabla_{x'} w \cdot \nabla_{x'} \tilde{u} dx + \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x') < x_n < \frac{\varepsilon}{2}+h_1(x')}} \sum_{\alpha=1}^{n-1} w (\partial_{\alpha} \tilde{u} + \partial_{\alpha} w) \frac{x_{\alpha}}{r} ds \\
&\leq \frac{1}{2} \int_{\Omega_{1/2}} |\nabla w|^2 dx + \frac{1}{2} \int_{\Omega_{1/2}} |\nabla_{x'} \tilde{u}|^2 dx \\
&\quad + \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x') < x_n < \frac{\varepsilon}{2}+h_1(x')}} C(|\nabla_{x'} \tilde{u}|^2 + |w|^2 + |\nabla w|^2) ds.
\end{aligned}$$

By using (3.1),

$$\begin{aligned}
& \int_{\Omega_{1/2}} |\nabla_{x'} \tilde{u}|^2 dx \leq C(\|\varphi\|_{C^1(\Gamma_1^+)}^2 + \|\psi\|_{C^1(\Gamma_1^-)}^2), \\
& \int_{\substack{|x'|=\frac{1}{2}, \\ -\frac{\varepsilon}{2}+h_2(x') < x_n < \frac{\varepsilon}{2}+h_1(x')}} |\nabla_{x'} \tilde{u}|^2 ds \leq C(\|\varphi\|_{C^1(\Gamma_1^+)}^2 + \|\psi\|_{C^1(\Gamma_1^-)}^2).
\end{aligned}$$

Using (3.3) and (3.4), for $x \in \Omega_1 \setminus \overline{\Omega_{1/4}}$,

$$|\Delta \tilde{u}(x)| \leq C(\|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)}).$$

In view of $w = 0$ on Γ_1^\pm , by a standard elliptic theorem for Poisson equation, we have

$$\begin{aligned}
\|w\|_{L^\infty(\Omega_{2/3} \setminus \overline{\Omega_{1/3}})}^2 + \|\nabla w\|_{L^\infty(\Omega_{2/3} \setminus \overline{\Omega_{1/3}})}^2 &\leq C(\|w\|_{L^2(\Omega_1 \setminus \overline{\Omega_{1/4}})} + \|\Delta \tilde{u}\|_{L^\infty(\Omega_1 \setminus \overline{\Omega_{1/4}})}) \\
&\leq C(\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2),
\end{aligned}$$

which, implies that

$$\int_{\Omega_{1/2}} |\nabla w|^2 dx \leq C(\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2), \quad (3.6)$$

where C depends on n , λ , κ_0 and κ_1 .

Step 2. Estimate of $\int_{\widehat{\Omega}_\delta(x_0)} |\nabla w|^2 dx$, where $x_0 \in \Omega_{1/2}$, $\delta = \delta(x'_0) := \varepsilon + h_1(x'_0) - h_2(x'_0)$.

Let $\eta(x')$ be the cut-off function in Lemma 2.2. Multiplying $\eta^2 w$ on both sides of the equation in (3.5) and applying integration by parts, we have

$$\int_{\widehat{\Omega}_\delta(x_0)} \nabla w \cdot \nabla(\eta^2 w) dx = \int_{\widehat{\Omega}_\delta(x_0)} \Delta \tilde{u} \eta^2 w dx.$$

Note that

$$\int_{\widehat{\Omega}_s(x_0)} \nabla w \cdot \nabla(\eta^2 w) dx = \int_{\widehat{\Omega}_s(x_0)} |\nabla(\eta w)|^2 dx - \int_{\widehat{\Omega}_s(x_0)} w^2 |\nabla \eta|^2 dx,$$

by the Cauchy inequality, we have

$$\begin{aligned} \int_{\widehat{\Omega}_t(x_0)} |\nabla w|^2 dx &\leq \int_{\widehat{\Omega}_s(x_0)} |\nabla(\eta w)|^2 dx \\ &= \int_{\widehat{\Omega}_s(x_0)} w^2 |\nabla \eta|^2 dx + \int_{\widehat{\Omega}_s(x_0)} \Delta \tilde{u} \eta^2 w dx \\ &\leq \frac{C}{(s-t)^2} \int_{\widehat{\Omega}_s(x_0)} |w|^2 dx + (s-t)^2 \int_{\widehat{\Omega}_s(x_0)} |\Delta \tilde{u}|^2 dx. \end{aligned} \quad (3.7)$$

It is easy to see that

$$\int_{\widehat{\Omega}_s(x_0)} |w|^2 dx \leq \int_{|x'-x'_0|<s} C(\varepsilon + |x'|^2)^2 \int_{-\frac{\varepsilon}{2}+h_2(x')}^{\frac{\varepsilon}{2}+h_1(x')} |\nabla w|^2 dx_n dx',$$

and by (3.3)-(3.4),

$$\begin{aligned} &\int_{\widehat{\Omega}_s(x_0)} |\Delta \tilde{u}|^2 dx \\ &\leq C |\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \int_{|x'-x'_0|<s} \frac{1}{\varepsilon + |x'|^2} dx' \\ &\quad + C(\|\nabla \varphi\|_{L^\infty}^2 + \|\nabla \psi\|_{L^\infty}^2) \int_{|x'-x'_0|<s} \left(\frac{|x-x'_0|^2}{\varepsilon + |x'|^2} + 1 \right) dx' \\ &\quad + Cs^{n-1}(\|\nabla^2 \varphi\|_{L^\infty}^2 + \|\nabla^2 \psi\|_{L^\infty}^2). \end{aligned}$$

Case 1. For $|x'_0| \leq \sqrt{\varepsilon}$, $0 < t < s < \sqrt{\varepsilon}$, we have

$$\int_{\widehat{\Omega}_s(x_0)} |w|^2 dx \leq C\varepsilon^2 \int_{\widehat{\Omega}_s(x_0)} |\nabla w|^2 dx, \quad (3.8)$$

and

$$\begin{aligned} \int_{\widehat{\Omega}_s(x_0)} |\Delta \tilde{u}|^2 dx &\leq \frac{Cs^{n-1}}{\varepsilon} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \\ &\quad + Cs^{n-1}(\|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2). \end{aligned} \quad (3.9)$$

Denote $F(t) := \int_{\widehat{\Omega}_t(x_0)} |\nabla w|^2 dx$. By (3.7)-(3.9), for some universal constant $\hat{C}_1 > 0$, we have for $0 < t < s < \sqrt{\varepsilon}$,

$$\begin{aligned} F(t) &\leq \left(\frac{\hat{C}_1 \varepsilon}{s-t} \right)^2 F(s) \\ &\quad + C(s-t)^2 s^{n-1} \left(\frac{1}{\varepsilon} |\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \right. \\ &\quad \left. + \|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2 \right). \end{aligned} \quad (3.10)$$

Let $t_i = \delta + 2i\hat{C}_1\varepsilon$, $i = 0, 1, \dots$ and $k = \left\lceil \frac{1}{4\hat{C}_1\sqrt{\varepsilon}} \right\rceil + 1$, then

$$\frac{\hat{C}_1 \varepsilon}{t_{i+1} - t_i} = \frac{1}{2}.$$

Using (3.10) with $s = t_{i+1}$ and $t = t_i$, we obtain that, for $i = 0, 1, 2, \dots, k$,

$$\begin{aligned} F(t_i) &\leq \frac{1}{4}F(t_{i+1}) \\ &\quad + C(i+1)^{n-1}\varepsilon^n [|\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \\ &\quad + \varepsilon(\|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2)]. \end{aligned}$$

After k iterations, making use of (3.6), we have, for sufficiently small ε ,

$$\begin{aligned} F(t_0) &\leq C\varepsilon^n [|\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \\ &\quad + \varepsilon(\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2)], \end{aligned}$$

here we used that the first term in the last but one line decays exponentially, which implies that for $0 \leq |x'_0| \leq \sqrt{\varepsilon}$,

$$\begin{aligned} \|\nabla w\|_{L^2(\widehat{\Omega}_\delta(x_0))}^2 &\leq C\varepsilon^n [|\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \\ &\quad + \varepsilon(\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2)]. \end{aligned} \quad (3.11)$$

Case 2. Similarly, for $\sqrt{\varepsilon} < |x'_0| < \frac{1}{2}$, $0 < t < s < \frac{2|x'_0|}{3}$, we have

$$\begin{aligned} \|\nabla w\|_{L^2(\widehat{\Omega}_\delta(x_0))}^2 &\leq C|x'_0|^{2n} [|\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))|^2 \\ &\quad + |x'_0|^2(\|w\|_{L^2(\Omega_1)}^2 + \|\varphi\|_{C^2(\Gamma_1^+)}^2 + \|\psi\|_{C^2(\Gamma_1^-)}^2)]. \end{aligned} \quad (3.12)$$

Step 3. Estimate of $|\nabla w(x)|$ for $x \in \Omega_{R_0}$, for some small $R_0 \in (0, \frac{1}{2})$.

By using a similar argument in Lemma 2.3, it follows from (3.11) and (3.12) that, for $0 \leq |x'_0| \leq \sqrt{\varepsilon}$,

$$\begin{aligned} \|\nabla w\|_{L^\infty(\widehat{\Omega}_{\delta/2}(x_0))} &\leq C|\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))| \\ &\quad + C\sqrt{\varepsilon}(\|w\|_{L^2(\Omega_1)} + \|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)}). \end{aligned}$$

and for $\sqrt{\varepsilon} \leq |x'_0| \leq R_0$,

$$\begin{aligned} \|\nabla w\|_{L^\infty(\widehat{\Omega}_{\delta/2}(x_0))} &\leq C|\varphi(x'_0, \varepsilon/2 + h_1(x'_0)) - \psi(x'_0, -\varepsilon/2 + h_2(x'_0))| \\ &\quad + C|x'_0|(\|w\|_{L^2(\Omega_1)} + \|\varphi\|_{C^2(\Gamma_1^+)} + \|\psi\|_{C^2(\Gamma_1^-)}). \end{aligned}$$

By using (3.1)-(3.2), we obtain (1.11).

Finally, if $\varphi(0', \frac{\varepsilon}{2}) \neq \psi(0', -\frac{\varepsilon}{2})$, then by the Taylor expansion, we obtain

$$|\nabla u(0', x_n)| \geq \frac{|\varphi(0', \frac{\varepsilon}{2}) - \psi(0', -\frac{\varepsilon}{2})|}{C\varepsilon}, \quad \forall x_n \in \left(-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right).$$

The proof of Corollary 1.3 is completed.

REFERENCES

- [1] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I. *Comm. Pure Appl. Math.* 12 (1959), 623-727.
- [2] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II. *Comm. Pure Appl. Math.* 17 (1964), 35-92.
- [3] H. Ammari, E. Bonnetier, F. Triki and M. Vogelius, Elliptic estimates in composite media with smooth inclusions: an integral equation approach. *Ann. Sci. Éc. Norm. Supér.* (4) 48 (2015), no. 2, 453-495.

- [4] H. Ammari, G. Ciraolo, H. Kang, H. Lee and K. Yun, Spectral analysis of the Neumann-Poincaré operator and characterization of the stress concentration in anti-plane elasticity. *Arch. Ration. Mech. Anal.* 208 (2013), 275-304.
- [5] H. Ammari, G. Dassios, H. Kang and M. Lim, Estimates for the electric field in the presence of adjacent perfectly conducting spheres, *Quart. Appl. Math.* 65 (2007), no. 2, 339-355.
- [6] H. Ammari, H. Kang and M. Lim, Gradient estimates for solutions to the conductivity problem. *Math. Ann.* 332 (2005), no. 2, 277-286.
- [7] H. Ammari, H. Kang, H. Lee, J. Lee and H.Zribi, Optimal estimates for the electric field in two dimensions. *J. Math. Pures Appl.* (9) 88 (2007), no. 4, 307-324.
- [8] H. Ammari, H. Kang, H. Lee, M. Lim and H.Zribi, Decomposition theorems and fine estimates for electrical fields in the presence of closely located circular inclusions. *J. Differential Equations* 247 (2009), no. 11, 2897-2912.
- [9] I. Babuška, B. Andersson, P. Smith and K. Levin, Damage analysis of fiber composites. I. Statistical analysis on fiber scale. *Comput. Methods Appl. Mech. Engrg.* 172 (1999), no 1-4, 27-77.
- [10] E.S. Bao, Y.Y. Li and B. Yin, Gradient estimates for the perfect conductivity problem, *Arch. Ration. Mech. Anal.* 193 (2009), no. 1, 195-226.
- [11] E.S. Bao, Y.Y. Li and B. Yin, Gradient estimates for the perfect and insulated conductivity problems with multiple inclusions. *Comm. Partial Differential Equations* 35 (2010), no. 11, 1982-2006.
- [12] J.G. Bao, H.J. Ju and H.G. Li, Optimal boundary gradient estimates for Lam systems with partially infinite coefficients. *Adv. Math.* 314 (2017), 583-629.
- [13] J.G. Bao, H.G. Li and Y.Y. Li, Gradient estimates for solutions of the Lamé system with partially infinite coefficients. *Arch. Ration. Mech. Anal.* 215 (2015), 307-351.
- [14] J.G. Bao, H.G. Li and Y.Y. Li, Gradient estimates for solutions of the Lamé system with partially infinite coefficients in dimensions greater than two. *Adv. Math.* 305 (2017), 298-338.
- [15] E. Bonnetier and F. Triki, Asymptotics in the presence of inclusions of small volume for a conduction equation: a case with a non-smooth reference potential. *Imaging microstructures, Contemp. Math.* vol. 494, Amer.Math. Soc., Providence, RI, 2009, 95-111.
- [16] E. Bonnetier and F. Triki, On the spectrum of the Poincaré variational problem for two close-to-touching inclusions in 2D. *Arch. Ration. Mech. Anal.* 209 (2013), no. 2, 541-567.
- [17] E. Bonnetier and M. Vogelius, An elliptic regularity result for a composite medium with "touching" fibers of circular cross-section. *SIAM J. Math. Anal.* 31 (2000), no. 3, 651-677.
- [18] B. Budiansky and G.F. Carrier, High shear stresses in stiff fiber composites. *J. Appl. Mech.* 51 (1984), 733-735.
- [19] M. Giaquinta, Multiple integrals in the calculus of variations and nonlinear elliptic systems. *Annals of Mathematics Studies*, 105, Princeton University, Princeton, N.J., 1983.
- [20] H. Kang, M. Lim and K. Yun, Asymptotics and computation of the solution to the conductivity equation in the presence of adjacent inclusions with extreme conductivities. *J. Math. Pures Appl.* (9) 99 (2013), no. 2, 234-249.
- [21] H. Kang, H. Lee and K. Yun, Optimal estimates and asymptotics for the stress concentration between closely located stiff inclusions. *Math. Ann.* 363 (2015), no. 3-4, 1281-1306.
- [22] J.B. Keller, Stresses in narrow regions. *Trans. ASME J. Appl. Mech.* 60 (1993), 1054-1056.
- [23] J.B. Keller, Conductivity of a medium containing a dense array of perfectly conducting spheres or cylinders or nonconducting cylinders. *J. Appl. Phys.* 3 (1963), 991-993.
- [24] H.G. Li, Y.Y. Li, E.S. Bao and B. Yin, Derivative estimates of solutions of elliptic systems in narrow regions. *Quart. Appl. Math.* 71 (2014), no. 3, 589-596.
- [25] H.G. Li and L.J. Xu, Optimal estimates for the perfect conductivity problem with inclusions close to the boundary. *SIAM J. Math. Anal.* 49 (2017), no. 4, 3125-3142.
- [26] Y.Y. Li and L. Nirenberg, Estimates for elliptic systems from composite material. Dedicated to the memory of Jürgen K. Moser, *Comm. Pure Appl. Math.* 56 (2003), no. 7, 892-925.
- [27] Y.Y. Li and M. Vogelius, Gradient estimates for solutions to divergence form elliptic equations with discontinuous coefficients. *Arch. Ration. Mech. Anal.* 153 (2000), no. 2, 91-151.
- [28] M. Lim and K. Yun, Blow-up of electric fields between closely spaced spherical perfect conductors, *Comm. Partial Differential Equations* 34 (2009), no. 10-12, 1287-1315.
- [29] X. Markenscoff, Stress amplification in vanishingly small geometries, *Comput. Mech.* 19 (1996), 77-83.

- [30] O. Oleinik, A. Shamaev and G. Yosifian, Mathematical problems in elasticity and homogenization. Studies in Mathematics and its Applications, vol. 26, North-Holland Publishing Co., Amsterdam, 1992.
- [31] K. Yun, Estimates for electric fields blown up between closely adjacent conductors with arbitrary shape. SIAM J. Appl. Math. 67 (2007), no. 3, 714-730.
- [32] K. Yun, Optimal bound on high stresses occurring between stiff fibers with arbitrary shaped cross-sections. J. Math. Anal. Appl. 350 (2009), no. 1, 306-312.

(H.J. Ju) SCHOOL OF SCIENCES, BEIJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS,
BEIJING 100876, CHINA
E-mail address: `hjju@bupt.edu.cn`

(H.G. Li) SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, LABORATORY
OF MATHEMATICS AND COMPLEX SYSTEMS, MINISTRY OF EDUCATION, BEIJING 100875, CHINA.
E-mail address: `hgli@bnu.edu.cn`

(L.J. Xu) SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, LABORATORY
OF MATHEMATICS AND COMPLEX SYSTEMS, MINISTRY OF EDUCATION, BEIJING 100875, CHINA.