

Two-field axion inflation and the swampland constraint in the flux-scaling scenario

Cesar Damian^a and Oscar Loaiza-Brito^b

^b*Departamento de Ingeniería Mecánica, Universidad de Guanajuato,
Carretera Salamanca-Valle de Santiago Km 3.5+1.8 Comunidad de Palo Blanco, Salamanca, Mexico*

^c*Departamento de Física, Universidad de Guanajuato,
Loma del Bosque No. 103 Col. Lomas del Campestre C.P 37150 Leon, Guanajuato, Mexico.*

^b*E-mail:* cesaredas@fisica.ugto.mx, oloaiza@fisica.ugto.mx

ABSTRACT: Based on the flux-scaling scenario we study a model consisting on Type IIB string theory compactified on a Calabi-Yau manifold with a frozen complex structure in the presence of generic fluxes. The model contains (meta)stable Minkowski and de Sitter vacua as well as inflationary directions driven by two independent linear combination of axions. Due to a numerical control by fluxes, we show that cosmological parameters as the spectral index, the scalar-to tensor ratio and non-Gaussianities can be kept within observed bounds while preserving the desired hierarchies on physical scales. Moreover we compute the deviation of the inflationary trajectories from geodesics on field space in terms of the fluxes showing that they fulfill the recent proposed swampland criterion for multi-field scenarios.

KEYWORDS: Flux scaling scenario, swampland, multifield inflation.

Contents

1	Introduction	1
2	String compactification with non-geometric fluxes	4
2.1	Uplifting to metastable dS by D-terms	5
2.2	Uplifting the flat direction	7
3	Effective model of inflation	8
3.1	Hierarchy of masses	9
3.2	Spectral index	9
3.3	Tensor to scalar ratio	11
3.4	Non-Gaussianities	11
4	The swampland criterion	13
4.1	Multi-field inflation and the swampland constraint	13
4.2	The swampland criterion for two-field inflationary model in the flux scaling scenario	14
5	Final Remarks	16
A	Specifics of the model	18
A.1	Frozen complex structure modulus	18
B	Cosmological Observables and Slow-Roll Multi-field Inflation	19

1 Introduction

Since its conception, the inflationary mechanism [1] has become one of the most promising models to describe the very early stage of the universe by generically explaining some theoretical and observational puzzles of present measurements. Additionally, observational evidence is confident with a nearly scale-invariance spectrum and a tensor to scalar ratio lower than 0.1 [2] implying a Universe with a gaussian CMB profile. Although single field inflation matches with several cosmological observables, multi-field inflation is not completely ruled out [2]. In particular, in single field inflation the primordial non-gaussianities are suppressed by powers of the slow-roll parameters [3, 4]. Thus, it is expected that the non-linear interaction with other fields shall produce observable non-gaussianities within observational data [5]. Naturally, the non-gaussianities are expected to be large in multifield inflationary scenarios [6–10]. It is our goal to show that non-gaussianities can be kept under control, this is, with values

below the present observational bounds, within the context of the recently proposed flux-scale scenario [11].

The use of string theory as a quantum gravity theory to model an effective and consistent inflationary setup follows from the fact that inflation is very sensitive to UV corrections, mainly due to the high energy scale at which it is expected to occur. In this context, the inflationary potential is identified with one or more directions on the scalar potential (constructed at the level of supergravity) which in turn depends on scalar fields arising from a string compactification. Such dependence is achieved by turning on string fluxes supported on non-contractible cycles of the internal space, leading to the well-known moduli stabilization scenario. Stabilization of moduli has opened up a window not only for effectively reproducing inflation but also for giving a consistent setup on which UV corrections of the scalar potential are under control as a consequence of shift symmetries on the ten-dimensional theory and their inheritance on the corresponding four-dimensional axions. In this way, one of the best motivated and supported models of inflation from string theory is F-term axion monodromy inflation [12, 13].

Besides the above, F-term axion monodromy inflation also allows us to expect polynomial suppressed ratios of different scales by considering a tree-level superpotential depending on all moduli, including Kähler moduli, which generically have been stabilized by non-perturbative effects [14, 15]. The perturbative dependence on the Kähler moduli is accomplished by turning on non-geometric fluxes [16–18] whose presence in string theory is supported by recent studies in generalized geometry and double field theory [19–23]. Since their incorporation to string modeling, a lot of work has been done, proving that dS vacua are possible at tree level [11, 24–35].

Moreover, in the F-term axion inflation scenario, one expects that inflationary directions are driven by axions with a Kähler potential depending solely on the corresponding saxions. This scheme guarantees that inflationary directions do not get quantum corrections from the Kähler potential, allowing the possible construction of inflationary models with large values of r . However, in order to have a physical consistent frame, it is necessary to have some hierarchies on the different scales as¹

$$M_s > M_{KK} > M_{\text{inf}} > M_{i, \text{mod}} > H_{\text{inf}} > M_\theta, \quad (1.1)$$

where θ is the inflaton. The flux-scaling scenario [11, 33, 38, 39] establishes a mechanism to accomplish all the above requirements by having a parametrical control of the different scales and mass of the moduli by the presence of different kind of fluxes which precisely generates

¹A relation between mass hierarchies and the presence of inflationary directions was pointed out in Appendix A in [30], while the role played by the conifold on the existence of hierarchies and inflation was studied in [36, 37].

a perturbative tree-level superpotential on moduli. Roughly speaking, it is required a model with a symmetric superpotential on n complex fields in the presence of $2n - 1$ real fluxes. This leads to a scalar potential with a flat direction on a linear combination of axions, while the saxions, appearing on the Kähler potential are stabilized at a non-supersymmetric minimum. The flat direction is uplifted by parametrically breaking the symmetry of the superpotential on the complex moduli by adding an extra flux [13]. This addition brings a new term in the superpotential which describes an interaction between the above mentioned moduli. Under this scheme, it was shown that it is possible to find non-supersymmetric vacua free of tachyons by considering p -fluxes, while in [33] it was explored a specific scenario in which a non-supersymmetric AdS vacuum is uplifted to dS by turning on non-geometric fluxes which contribute to D-terms in the scalar potential.

Flat directions on the scalar potential driven by one or more moduli are of particular interest to construct inflationary scenarios corresponding to effective field theories with scalar fields coupled to gravity. However, it has been difficult to distinguish whether the models obtained from string theory correspond to those models belonging to consistent quantum theories of gravity at high energies. The set of those models not connected to consistent theories of quantum gravity has been called the “string swampland”. With the purpose to identify such regions it was recently proposed a criterion which seems to rule out the possibility to reproduce single-field inflation on string scenarios [40, 41]. This criterion was studied in the context of multi-field inflation in [42] where it was shown that it is possible to have inflation driven by two or more scalar fields while fulfilling the swampland criterion. Therefore it is interesting to study the possibility to have inflation on the scale-flux scenario and whether this construction allows us to construct inflationary directions driven by more than two moduli while satisfying the swampland criterion on the slow-roll conditions.

Summarizing, we are interested in a model in which we can have a hierarchy of physical scales under a parametrical control by fluxes, with a stable minimum and inflationary directions. A minimalistic model consisting on an effective theory with two-scalar fields seems to be a simple scenario to look for all the above conditions. In this work, by selecting a particular model presented in [11], we study a concrete example in which inflationary directions are driven by two scalar fields corresponding to linear combinations of axions. Since in a multi-field scenario it is possible that non-Gaussianities are present, we show that under numerical control by fluxes, non-Gaussianities can also be bounded below observed values. Moreover, we show that for some inflationary trajectories around a Minkowski minimum, the swampland criterion is fulfilled in the context of multi-field inflation described in [42]. For that we follow the flux-scaling scenario by identifying the inflationary trajectories to those corresponding to unstabilized moduli which are slightly uplifted by the addition of p -fluxes. In the approximation of small p we show that the curvature parameter of inflationary trajectories can be written as function of fluxes allowing us to get large values of it by a parametrical control of fluxes. The ratio between this parameter and the Hubble constant, which determines whether

or not the swampland criterion is satisfied, is also larger than one on different zones of the trajectories. However, this ratio cannot be parametrically controlled by fluxes, at least for constant values. It is also important to mention that for all our analysis it is necessary to relax the integer quantization requirement on fluxes. We comment on this issue in our final remarks.

Our work is organized as follows: In section II we review the model studied in [11] and present our two-field inflationary model. We also compute the ratio of the different scales parametrically controlled by 3 different fluxes. Section III is devoted to compute cosmological parameters as the tensor to scalar ratio, mass hierarchy, the spectral index and non-Gaussianities. We show that all these values can be expressed in terms of a ratio of fluxes. However, as mentioned, integer quantization of fluxes seems to bring numerical values beyond observations. This issue must be understood and carefully studied. In section IV we explore the values of the inflationary swampland criterion for the above inflationary trajectories. Finally, we present our conclusions and a couple of Appendices are devoted to review some useful notation.

2 String compactification with non-geometric fluxes

We shall select a model studied in detailed in [11] (for details see Appendix A) consisting on Type IIB superstring compactified on Calabi–Yau manifolds in the presence of orientifold 3–planes with non-vanishing fluxes, including the common NS-NS and RR fluxes from the closed string sector and the so called non-geometric fluxes Q and R [16–18]. Specifically we study the case for a CY manifold with a frozen complex structure and a single Kähler modulus, this is $h_-^{2,1} = 0$ and $h_+^{1,1} = 1$. We have also limited our model to manifolds for which $h_-^{1,1} = 0$ meaning that the closed string potential form called the geometric moduli $G_2 = Sb_2 + c_2$ is absent in the superpotential, and with $h_+^{2,1} = 1$ allowing the presence of D-terms. For this case, the Kähler potential reads

$$K = -3\log(T + \bar{T}) - \log(S + \bar{S}), \quad (2.1)$$

with a perturbative dependence on the complex moduli fields $S = e^{-\phi} + ic$ and $T = \tau + i\rho$. The superpotential depends on the coupling between RR and NS-NS fluxes with the complex structure, the geometric moduli with the odd Kähler moduli (which we consider absent) and the non-geometric fluxes Q with the Kähler moduli. Notice that the R flux is not present in the superpotential. Therefore, by turning on 3 real fluxes f, h and q the superpotential is (see Appendix A for notation)

$$W = if + ihS + iqT, \quad (2.2)$$

showing a linear dependence on S and T , where f, h and q are *in principle* integer quantized RR, NS-NS and Q fluxes respectively.

This model presents an explicitly manifestation of the scenario proposed in [11] in which we have 3 fluxes and 4 real moduli. It is then expected that 3 out of 4 real moduli get a vacuum expectation value while a flat direction would be present. In fact, as shown in [11], the F-term potential generated by (2.2) has an interesting non-supersymmetric and non-tachyonic AdS extremum, with a scalar potential depending solely on τ , s and θ where

$$\theta = q\rho - hc, \quad (2.3)$$

and the minimum is obtained at $(s, \tau, \theta) = (-f/h, -6f/5q, 0)$ with

$$V_{0F} = -\frac{M_{\text{Pl}}^4}{2^4 \pi} \frac{5^2 q^3 h}{3^3 f^2}, \quad (2.4)$$

It is important to mention that for $|f/h| \gg 1$ and $|f/q| \gg 1$, this models exhibits a weak string coupling and large radius implying that higher-order corrections of the scalar potential can be safely ignored. These fluxes are subject to the Bianchi identities and tadpole cancellation conditions given in [11], however, Bianchi identities are trivially fulfilled by the selected fluxes.

2.1 Uplifting to metastable dS by D-terms

We now want to analyze the effect of adding the D-term potential allowed for $h_+^{2,1} = 1$ closely following [33] to uplift to Minkowski or dS minimum. For that it is considered the presence of extra non-geometric fluxes f and R which survive the orientifold projection and appear in the D-term of the scalar potential. Such a term was computed in [33] and by turning on just two fluxes, r and g respectively, it reads

$$V_D = \frac{\delta}{v\rho^2} \left(g - \frac{r\rho}{3s} \right)^2, \quad (2.5)$$

where δ is a positive constant, and $s = e^{-\phi}$. Observe that this term depends on all the saxions in the model. The fluxes entering in V_D are related to the action of the twisted differential \mathcal{D} on the even $(2,1)$ forms. Such fluxes do not enter at all in the superpotential W that determines the F-term potential. However, there are Bianchi identities that mix r and g with NS-NS and Q -fluxes appearing in W . In our case there is only one non trivial BI constraint given by

$$rh + gq = 0. \quad (2.6)$$

We find that the model admits an uplift that can be either to Minkowski or dS depending on the value of δ . Due to the complexity of the solution, we limit ourselves to present the values of stabilized moduli as Taylor series expansions to linear order in $V_0 = \Lambda$. For an arbitrary value of Λ , there is still an extremum at $\theta = 0$ whereas the saxions are stabilized at

$$s = -\frac{1}{2^3} \frac{f}{h} - \frac{3^4}{2^7} \frac{f^3}{q^3 h^2} \Lambda + \mathcal{O}(\Lambda^2), \quad \tau = -\frac{3^2}{2^3} \frac{f}{q} - \frac{11 \cdot 3^4}{2^8} \frac{f^3}{q^4 h} \Lambda + \mathcal{O}(\Lambda^2). \quad (2.7)$$

The constant δ in the D-term potential is given by

$$\delta = \frac{1}{2^4} \frac{qh}{g^2} + \frac{3^4}{2^{10}} \frac{f^2}{g^2 q^2} \Lambda + \mathcal{O}(\Lambda^2). \quad (2.8)$$

We then conclude that for small $\Lambda > 0$ and $q, h > 0$, we stay in the physical region for $f < 0$ which also guarantees that $\delta > 0$. To attain the perturbative regime we take $|f| > q, h$. We remark that the above series converge for small enough Λ . Clearly for $\Lambda = 0$ we have a Minkowski extremum.

Hence, at the minimum the moduli acquire the following masses:

$$M_{\text{mod},i}^2 = \left(\mu_i \frac{q^3 h}{f^2} + \tilde{\mu}_i \Lambda + \mathcal{O}(\Lambda^2) \right) \frac{M_{\text{Pl}}^2}{4\pi}, \quad (2.9)$$

with the coefficients given by

$$\mu_i = \left\{ \frac{2^6(17 + \sqrt{181})}{3^6}, \frac{2^6(17 - \sqrt{181})}{3^6}; \frac{2^6 \cdot 7}{3^6}, 0 \right\}, \quad (2.10)$$

and

$$\tilde{\mu}_i = \left\{ -\frac{19 \cdot 43}{2 \cdot 3^3} - \frac{5591}{2 \cdot 3^3 \cdot 181^{1/2}}, -\frac{19 \cdot 43}{2 \cdot 3^3} + \frac{5591}{2 \cdot 3^3 \cdot 181^{1/2}}; \frac{7 \cdot 19}{3^3}, 0 \right\}. \quad (2.11)$$

The first (last) two entries are related to linear combination of saxions (axions).

Notice that the scalar potential $V = V_F + V_D$ still has a flat direction on an orthogonal direction to θ . However the mass of moduli is altered by the value of Λ . We observe that for small enough Λ there are no tachyons but as Λ increases, the first normalized mass eigenvalue becomes negative. Therefore, in order to have a stable minimum, it is required a small value for Λ .

Finally, the string and KK mass are given by

$$\begin{aligned} M_s^2 &= \frac{2^{9/2} \pi q^{3/2} h^{1/2}}{3^3 f^2} M_{\text{Pl}}^2 - \frac{17\pi}{2^{3/2}} \frac{\Lambda}{q^{3/2} h^{1/2}} M_{\text{Pl}}^2 - \mathcal{O}(\Lambda^2), \\ M_{\text{KK}}^2 &= \frac{2^4 q^2}{3^4 f^2} \frac{M_{\text{Pl}}^2}{4\pi} - \frac{11}{3^2} \frac{\Lambda}{qh} \frac{M_{\text{Pl}}^2}{4\pi} - \mathcal{O}(\Lambda^2), \end{aligned} \quad (2.12)$$

from which we observe that the net contribution of a positive cosmological constant diminishes the values of M_s^2 and M_{KK}^2 at subleading order. The ratio of KK to string scale at linear order in Λ is

$$\frac{M_{\text{KK}}^2}{M_s^2} = \frac{1}{3 \cdot 2^{5/2} \pi^2} \frac{q^{1/2}}{h^{1/2}} + \frac{3 \cdot 7}{2^{17/2} \pi^2} \frac{f^2 \Lambda}{q^{5/2} h^{3/2}} + \mathcal{O}(\Lambda^2). \quad (2.13)$$

Thus, we have parametrical control of the KK mass over the string mass for $q < h$, as pointed out in [11]. Indeed, the ratio of the moduli to KK scale has the same behavior as in the

non-supersymmetric non-tachyonic AdS extremum in [11], which for sake of completeness we present here:

$$\frac{M_{\text{mod}}^2}{M_{\text{KK}}^2} = \frac{3^4}{2^4} h q \mu_i + \left(\frac{11 \cdot 3^6}{2^8} \frac{f^2 \mu_i}{q^2} - \frac{3^4}{2^4} \frac{f^2 \mu_i}{2^4 q^2} \right) \Lambda + \mathcal{O}(\Lambda^2), \quad (2.14)$$

where we have used the following expressions for m_s^2 and m_{KK}^2 [43]

$$m_s^2 = \frac{\pi}{s^{1/2} \mathcal{V}^{1/2}}, \quad (2.15)$$

and

$$m_{\text{KK}}^2 = \frac{1}{4\pi \mathcal{V}^2}. \quad (2.16)$$

with $s = e^{-\phi}$ and \mathcal{V} is the volume modulus of the CY in the Einstein frame in Planck units.

2.2 Uplifting the flat direction

In order to unflatten the orthogonal direction to θ , we follow the same receipt proposed in [11] and consider to turn on a p-flux, with a superpotential

$$W = \lambda W_0 + p S T, \quad (2.17)$$

where W_0 is original superpotential with a flat direction and λ is a scaling parameter. The lightest modes are two orthogonal axions, given by θ and

$$\sigma = -\frac{q}{s^2} c - 3 \frac{h}{\tau^2} \rho. \quad (2.18)$$

In terms of canonical terms, the corresponding normalized fields read

$$\tilde{\theta}^2 = \frac{3}{12h^2 s^2 + 4q^2 \tau^2} \quad \text{and} \quad \tilde{\sigma}^2 = \frac{s\tau}{12h^2 s^2 + 4q^2 \tau^2}, \quad (2.19)$$

where the saxionic fields stay fixed at the minima. This condition is expected to be valid due to the mass hierarchy.

In order to break this structure without destabilizing the former vacua, we choose the S-dual flux $p = -\tilde{\theta}$, and to preserve parametrical control of the masses, we choose $\lambda = 1/50$. In this way there is a Minkowsky minima with no flat directions. The masses now satisfies the proposed hierarchy:

$$m_s^2 > m_\rho^2 > m_\sigma^2 \sim m_\theta^2, \quad (2.20)$$

where the orthogonal axionic masses are of the same order. Thus, it is natural to identify the two orthogonal axions as the scalar fields driving inflation. It is interesting to note that the

lightest scalar is in the sgoldstino direction.

Since we are dealing with two scalar fields, it is necessary to look for inflationary trajectories in the context of multi-field inflation. In such a framework the slow-roll parameters for single field inflation are generalized according to the review shown in Appendix B. We then proceed to compute the existence of three different inflationary trajectories² which are shown in Figure 1 in which we show the scalar potential as function of the axions $\tilde{\theta}$ and $\tilde{\sigma}$ for a specific choice of fluxes. It is important to stress out the fractional values given to the fluxes. We have found that for integer values it is not possible neither to fulfill the hierarchy of mass nor to find inflationary trajectories.




Figure 1. Plot of a) scalar potential and b) stable inflationary trajectories for $f = 1/40$, $h = 1/50$, $q = -1/17$ and $\lambda = 1/50$.

3 Effective model of inflation

With the purpose of computing the number of e-folds that guarantees a required minimal amount of inflation, the initial points are selected in such a way that they satisfy the inflationary conditions at the beginning of the trajectory while the pivot scales are chosen once the inflationary conditions are violated by reaching $\eta = 1$. The three initial values as well the number of e-folds at the end of inflation are shown in Table 1.

(σ_0, θ_0)	$e - folds$
$(-3.8, 20)$	93
$(16, 13)$	60
$(16, -16)$	58

Table 1. Initial value and number of e-folds obtained for selected trajectories.

²Their selection also follows from the number of e-folds they produce as shown later on.

The inflationary trajectories are obtained by a numerical solution of Eqs. (B.2) and (B.1). As usual we choose arbitrary points in the field space as the initial value and we solve numerically the FRW equations until the minima is reached. At the end of inflation ($\epsilon \sim 1$) we compute the number of e-folds. We propose three trajectories where at least 40–60 e-folds are obtained. As we shall see, the expected hierarchy of scales is still preserved pointing out the possible presence of two-field inflation.

3.1 Hierarchy of masses

In this section we present the numerical values for the scales obtained in our model. The fluxes are selected in such a way that the saxions becomes heavier than its axion partners at least by one order of magnitude, and the two axions have a similar mass, this is

$$m_s > m_{\text{KK}} > m_{\text{inf}} \sim m_{\text{mod}} \sim H_{\text{inf}} < m_{\tilde{\theta}} \sim m_{\tilde{\sigma}}. \quad (3.1)$$

As is shown in Eq. 2.15 and 2.16 mass of scales depend only on the saxions expectation values. Thus, for the present model the KK scale as well the string scale are fixed during inflation due to the hierarchy of saxions. Table II shows the corresponding values in Planck units.

Table 2. Mass hierarchy for the selected model.

Mass	
m_s	5.64
m_{KK}	$5.25 \cdot 10^{-2}$
m_{inf}	$1.257 \cdot 10^{-2}$
m_{saxions}	$\{9.99 \cdot 10^{-3}, 2.23 \cdot 10^{-2}\}$
H_{inf}	$2.00 \cdot 10^{-2}$
m_{axions}	$\{1.75 \cdot 10^{-3}, 2.37 \cdot 10^{-4}\}$

Tadpole cancellation condition for the selected models implies

$$\mathfrak{f}h = N_{O3} - N_{D3}, \quad \mathfrak{f}q = N_{O7} - N_{D7}, \quad (3.2)$$

which for the selected fluxes requires $N_{O3} > N_{D3}$ and $N_{O7} < N_{D7}$. Notice that in our numerical example all fluxes have values less than 1. Therefore their contribution to tadpoles is of order $\mathcal{O}(10^{-4})$ in Planck units. Although fractional amounts of fluxes are not allowed by Dirac quantization, the fluxes in the superpotential could have perturbative and non-perturbative corrections as shown in [44]. In such case, no tadpole exists in the model.

3.2 Spectral index

In Figure 2 it is shown the evolution of the slow-roll parameters as a function of the e-folding. As observed for all the trajectories, at the beginning of inflation there exists a momentarily fast-rolling region which does not last more than a couple of e-folds. Besides that we find

that there exist a range in moduli space in which inflation occurs. A particularly interesting issue appears for trajectory III which contains a region in the field space where there is a slightly accelerated expansion while fulfilling the slow-roll conditions. As we shall see, this region has important contributions to all cosmological observables.




Figure 2. Slow-roll parameters ϵ (solid line) and η (dashed line) for a) trajectory I, b) trajectory II and c) trajectory III.

The spectral index n_s is calculated from the relation shown in Eq. B.9 and it is shown in Figure 3 for our three trajectories. The filled lines represent the allowed value of n_s as reported by Planck. The results shown that all trajectories meet the observable bounds at different times during the slow-roll dynamics. For the case of trajectory I, the bounds are matched at the end of inflation. For trajectories I and III the horizon exit is expected to occur at the end of inflation in order to agree with observations. However, for trajectory II, observational bounds are matched at the beginning of inflation. As is pointed out in the previous paragraph, there is a region in the field space where the spectral index varies in an oscillatory manner. This region is related to the variation of the ϵ parameter. This behavior implies an acceleration during the slow-roll regime.




Figure 3. Plot of spectral index n_s for a) trajectory I, b) trajectory II and c) trajectory III.

3.3 Tensor to scalar ratio

The tensor to scalar ratio r is calculated through the Eq. B.10 which is defined as the ratio of power spectra of tensor and scalar perturbations. Since the claim of the possibility to have scenarios with a non-negligible value of r , this cosmological observable has attracted considerable attention in the last years. In Figure 4 we present the tensor to scalar ratio for the three selected trajectories. As observed the trajectory III presents a large ratio at the end of inflation which does not meet the current observational bounds of $r < 0.05$ as reported by Planck. However, trajectory I and II, give a small value of r which are in good agreement with the reported values.

3.4 Non-Gaussianities

The local non-Gaussianities f_{NL} are determined through Eq. B.12 as a position space expansion around Gaussian perturbations. Under this ansatz, non-Gaussianities are generated independently at different spatial points [45]. Translating this requirement into the context of inflation, typically implies that non-Gaussianities are generated on super-Hubble scales. Therefore,

$$\xi(x) = \xi_g(x) + \frac{3}{5} f_{NL} \xi^2(x) + \dots \quad (3.3)$$

where $\xi_g(x)$ is a perturbation variable that satisfies the Gaussian statistics. Thus a positive value of local non-Gaussianities enhance the value of the power spectra, providing more hot spots on the CMB, whereas a negative value shall be related to colder spots. Thus, a




Figure 4. Plot of the tensor to scalar ratio r for a) trajectory I, b) trajectory II and c) trajectory III.

positive value of f_{NL} shall be related to more large structure formation of Galaxies. Figure




Figure 5. Plot of non-Gaussianities for a) trajectory I, b) trajectory II and c) trajectory III.

5 shows non-Gaussianities obtained in the presented model for our three trajectories. As observed, along trajectory II we find $|f_{NL}| < 0.1$ giving rise to a nearly Gaussian spectrum,

which is compatible with the current observation. However, for trajectories I and III, the model predicts a very large amount of non-Gaussianities and in consequence a large amount of hot(blues) spots in the CMB which are beyond the current observational bounds.

4 The swampland criterion

Construction of realistic scenarios from string compactifications has faced a variety of obstacles with efforts to overcome them. It is difficult to distinguish effective field theories obtained from string theory by some type of compactification to those which apparently are not compatible with a quantum theory of gravity. The landscape and the swampland, respectively, are two regions in which stringy models may belong to. Recently, a criterion for that was presented in [40, 41] establishing a bound from below to the gradient of the scalar potential in case it has a positive value. The proposal has received a lot of attention specially for its implications on cosmology [46–54] which essentially states that single-field slow-roll inflation is ruled out. However, for the case of a multi-field inflationary scenario it was shown in [42] that the above criterion and inflation can in fact coexist. Therefore, we want to study whether our model is compatible with the swampland criterion while preserving inflationary trajectories. For that we start with a very brief review on the swampland proposal in a multi-field scenario closely following [42].

4.1 Multi-field inflation and the swampland constraint

The inflationary swampland criterion about the bounding on the slope of the scalar potential is given by

$$\frac{|\nabla V|}{V} \leq c \sim \mathcal{O}(1), \quad (4.1)$$

which for a single field effective theory implies that the slow-roll parameter $\epsilon \sim \mathcal{O}(1)$ ruling out inflation. However, as shown in [42], the above criterion must be generalized for the case of a multi-field scenario in which the bound is now given by ϵ_V , with

$$\epsilon_V = \epsilon \left(1 + \frac{\Omega^2}{9H^2} \right). \quad (4.2)$$

The parameter Ω measures the bending of the trajectory with respect to a geodesic in the moduli field space. Essentially, is given by the modulus of the covariant derivative on time of the tangent component of the trajectory. Therefore, if the ratio $\Omega/3H$ is large enough to produce an equally large value of ϵ_V the swampland criterion is fulfilled and inflation is also present on such trajectory if ϵ is still small. It was then argued in [42] that for non-geodesic trajectories it is possible to have inflation with a large ϵ_V . In this section we shall show that this is indeed the case for our model.

4.2 The swampland criterion for two-field inflationary model in the flux scaling scenario

In Figure 6 we compare ϵ and ϵ_V for the three selected models studied in the previous section. As is shown in [42], there is a deviation from ϵ due to the fact that we have a two-field inflation scenario. This is, whereas ϵ remains small during all the trajectory, there exists variations for ϵ_V due to angular velocity on the field trajectory. It is however, important to mention that these variations do not fulfill the bound proposed in [42] about having $\epsilon_V > 180H$.




Figure 6. Plot of inflationary parameters ϵ and ϵ_V for a) trajectory I, b) trajectory II and c) trajectory III.

For sake of completeness let us try to construct an explicit form for Ω . For that we consider the superpotential related to a Minkowski vacuum, this is, we shall take $\Lambda = 0$. Once we have the stabilized values for $\tilde{\theta}$ and $\tilde{\sigma}$, the scalar potential can be written as a function of s and τ and also of the fluxes h, f and q . Now we take the contribution to the scalar potential by turning on the flux p by which we obtain the vev's for s and τ given by

$$\begin{aligned}
 s &= \frac{1}{2^3} \frac{f}{h} + \frac{47}{2^5} \frac{pf}{qh} c + \frac{2}{3} \frac{p\theta}{qf} c^2 - \frac{7}{2^7} \frac{pf\theta}{qh}, \\
 \tau &= -\frac{3^2}{2^3} \frac{f}{q} - \frac{303}{2^5} \frac{pf}{qh} c + \frac{59}{2^7} \frac{pf\theta}{q^2 h} - \frac{2 \cdot 7}{3} \frac{ph\theta}{q^2 f} c^2.
 \end{aligned} \tag{4.3}$$

For small values³ of p and $\tilde{\theta}$ the scalar potential is written at first order on p as

$$V = \frac{2^2}{\sqrt{7}} \frac{pq^2}{f} \left(\frac{13}{3^3} \tilde{\theta} - \frac{1}{3^5} \tilde{\theta}^3 - \frac{2}{\sqrt{3}} \tilde{\sigma} + \frac{2}{3^3 \cdot \sqrt{3}} \tilde{\sigma} \tilde{\theta}^2 - \frac{1}{3^2} \tilde{\sigma}^2 \tilde{\theta} \right). \quad (4.4)$$

In order to completely geometrize the swampland criteria, let us determine the dependence of the Ω parameter in the slow-roll approximation. First, we notice that Ω is related to the dynamics of the scalar fields as

$$\Omega = \frac{\dot{\tilde{\sigma}} \ddot{\tilde{\theta}} - \tilde{\theta} \ddot{\tilde{\sigma}}}{\dot{\tilde{\theta}}^2 + \dot{\tilde{\sigma}}^2} \quad (4.5)$$

and the time derivative of the Hubble scale is determined by

$$\dot{H} = -\frac{1}{2} \dot{\phi}_0^2, \quad (4.6)$$

for $\phi_0 = \sqrt{\delta_{ij} \dot{\phi}^i \dot{\phi}^j}$, thus we have that the time variation of the Hubble scale is related to the kinetic energy of the scalar fields as

$$\dot{H} = -\frac{1}{2} (\dot{\tilde{\theta}}^2 + \dot{\tilde{\sigma}}^2). \quad (4.7)$$

The denominator of Ω is similarly related to the time derivative of the Hubble scale as

$$\Omega = -2 \frac{\dot{\tilde{\sigma}} \ddot{\tilde{\theta}} - \tilde{\theta} \ddot{\tilde{\sigma}}}{\dot{H}}, \quad (4.8)$$

Now, in the slow-roll approximation, the second derivative with respect to time of the scalar fields shall be related to the gradients of the scalar potential. This is

$$\ddot{\phi}_1^a = -\frac{1}{3} \left(\frac{\dot{V}^a}{H} - \frac{V^a \dot{H}}{H^2} \right), \quad (4.9)$$

with $V^a = \partial_a V$. We realize that the second term is suppressed by the slow-roll parameter as

$$\ddot{\phi}_1^a = -\frac{1}{3} \left(\frac{\dot{V}^a}{H} + \epsilon V^a \right) \quad (4.10)$$

then, the numerator of Ω can be written as

$$\dot{\tilde{\sigma}} \ddot{\tilde{\theta}} - \tilde{\theta} \ddot{\tilde{\sigma}} = \frac{1}{3^2} \left(\frac{\dot{V}^1 V^2 - \dot{V}^2 V^1}{H^2} \right) \quad (4.11)$$

where the upper index, indicates derivative with respect to the scalar fields $\{1, 2\} = \{\tilde{\theta}, \tilde{\sigma}\}$. Let us define the vectors $\hat{V} = \langle V^1, V^2 \rangle$ and $\hat{V}_O = \langle V^2, -V^1 \rangle$ (notice that $\hat{V} \cdot \hat{V}_O = 0$), thus

$$\dot{\tilde{\sigma}} \ddot{\tilde{\theta}} - \tilde{\theta} \ddot{\tilde{\sigma}} = \frac{1}{3^2} \left(\frac{\hat{V}^a \cdot d_t \hat{V}^a}{H^2} \right) \quad (4.12)$$

³We comment on this issue in our final remarks.

the Ω parameter is written as

$$\Omega = \frac{2}{3^2} \left(\frac{\hat{V}_O^a \cdot d_t \hat{V}^a}{\dot{H} H^2} \right) \quad (4.13)$$

where in order to simplify notation we have used $d_t \hat{V}^a = \dot{\hat{V}}^a$. Thus in the slow-roll approximation, where $H^2 = \frac{1}{3}V$, the time derivative of the Hubble scale is given by

$$\dot{H} = \frac{1}{3^{1/2} \cdot 2} \frac{\dot{V}}{V^{1/2}}, \quad (4.14)$$

from which we get that

$$\Omega = \frac{2^2}{3^{1/2}} \left(\frac{\hat{V}_O^a \cdot d_t \hat{V}^a}{\dot{V} V^{1/2}} \right) \quad (4.15)$$

Thus, for constant fluxes any dependence of the form $V = \alpha F(\tilde{\theta}, \tilde{\phi})$, with α a function on the fluxes, shall provide a parametric control of the fluxes over the angular velocity Ω of the form $\alpha^{1/2}$. However, the its ratio

$$\frac{\Omega}{H} = 2^2 \left(\frac{\hat{V}_O^a \cdot d_t \hat{V}^a}{\dot{V} V} \right), \quad (4.16)$$

does not provide parametrical control from a potential of this kind. One way to rescue a parametrical control by fluxes on the ratio Ω/H is to allow the presence on non-constant fluxes [55] which would give rise to terms of the form α^a . This would also allows us to control wether this model belongs or not to the swampland.

5 Final Remarks

Based on the flux-string scenario we have constructed a simple model giving rise to a two-field inflationary scenario in which besides the hierarchy of masses of the involved scales as well as the presence of inflationary trajectories, stable vacua and moduli stabilization at tree-level, we also have numerical control by fluxes of cosmological parameters as the spectral index, scalar-to-tensor ratio and non-gaussianities. Moreover we show that by the incursion of non-geometric fluxes it is possible to uplift the stable non-supersymmetric AdS minimum to Minkowski or de Sitter. The inflationary trajectories correspond to those generated by uplifting flat directions by the presence of p -fluxes, which in turn breaks down the symmetry of the superpotential on the complex moduli S and T . However, as recently proposed, it is important to see wether inflation can coexist with an inflationary swampland criterion which establishes that effective inflationary models might be consistent to a theory of quantum gravity if the ratio Ω/H is of order one or larger.

For single-field scenarios this constraint rules out inflation, but not for multi-field inflation since Ω parametrizes the departure of inflationary trajectories to geodesics in the moduli

space. We have shown that for our model this is indeed the case where two-field inflation and the above mentioned swampland criterion can in fact coexist. It is important to mention that the inflaton corresponds to a linear combination of axions, which in the context of F-theory inflation monodromy, implies the absence of possible quantum corrections to the scalar potential.

A drawback in our approach is reflected on the impossibility to parametrically control the ratio Ω/H by fluxes. This follows from the approximation we are considering by selecting small values of p -fluxes. One way to circumvent this result is to consider instead non-constant fluxes but fluxes depending on moduli.

However, as followed from the last assertion, there is a subtle issue that must be stressed out. The existence of hierarchies, inflationary trajectories, and their subsequent parametrical control is based on the assumption of non-integer values for the fluxes. Therefore, this model could be considered an effective realistic model if there is a way to enforce the fluxes to violate Dirac quantization. Mirror symmetry contributions can be responsible of this, but we leave such study for future work.

Acknowledgments

We thank Nana Cabo, Anamaría Font, Xin Gao, Daniela Herschmann, Damián Mayorga, Pramod Shukla, Rui Sun, and Ivonne Zavala for useful discussions. We specially thanks Ralph Blumenhagen for explanations and suggestions. The work of C.D. is supported by a DAIP-Universidad de Guanajuato project No. CIIC 233/2018. O. L.-B. is supported by CONACyT project No. 258982 and by DAIP-Universidad de Guanajuato project No. CIIC 230/2018.

A Specifics of the model

Here we review the model used in our work which was studied in detail in [11] and [33]. We consider a Type IIB compactification on a threefold Calabi-Yau manifold in the presence of RR and NS-NS fluxes, non-geometric fluxes f, Q and R and $O3$ -planes. For generic values of Betti numbers we have a perturbative tree-level superpotential given by

$$W = -P_1(U) + iSP_2(U) - iG^a P_{3,a}(U) + iT_\alpha P_4^\alpha(U), \quad (\text{A.1})$$

where the complex polynomial $P_i(U)$ are given by

$$\begin{aligned} P_1(U) &= f_\lambda X^\lambda - \tilde{f}^\lambda F_\lambda, & \lambda = 0, \dots, h_-^{2,1}, \\ P_2(U) &= h_\lambda X^\lambda - \tilde{h}^\lambda F_\lambda, \\ P_{3,a}(U) &= f_{\lambda a} X^\lambda - \tilde{f}_a^\lambda F_\lambda, & a = 1, \dots, h_-^{1,1}, \\ P_4^\alpha(U) &= q_\lambda^\alpha X^\lambda - \tilde{q}^{\lambda\alpha} F_\lambda, & \alpha = 1, \dots, h_+^{1,1}. \end{aligned} \quad (\text{A.2})$$

The moduli fields are $S = s + ic$, with $s = e^{-i\phi}$ and c the integrated RR zero-form, and $T_\alpha = \rho_\alpha + i\tau_\alpha$. The orientifold surviving fluxes appearing in the superpotential are written in terms of the symplectic basis $(\alpha_\Lambda, \beta^\Lambda)$ ($\Lambda = 0, \dots, h^{2,1}$) as

$$\begin{aligned} F_3 &= -\tilde{f}^\lambda \alpha_\lambda + f_\lambda \beta^\lambda, & : \text{RR} \\ H_3 &= -\tilde{h}^\lambda \alpha_\lambda + h_\lambda \beta^\lambda, & : \text{NS-NS} \\ f \cdot \omega_a &= -f_{\lambda a} X^\lambda + \tilde{f}_a^\lambda F_\lambda, & : \text{geometric flux } f \\ Q \cdot \tilde{\omega}^\alpha &= -q_\lambda^\alpha X^\lambda + \tilde{q}^{\lambda\alpha} F_\lambda, & : \text{non-geometric flux } Q. \end{aligned} \quad (\text{A.3})$$

These fluxes contribute to D-brane tadpole conditions as follows

$$\begin{aligned} N_{O3} - N_{D3} &= -f_\lambda \tilde{h}^\lambda + \tilde{f}^\lambda h_\lambda, \\ (N_{O5} - N_{D5})_a &= f_\lambda \tilde{f}_a^\lambda - \tilde{f}^\lambda f_{\lambda a}, \\ (N_{O7} - N_{D7})_\alpha &= -f_\lambda \tilde{q}^{\lambda\alpha} + \tilde{f}^\lambda q_\lambda^\alpha. \end{aligned} \quad (\text{A.4})$$

A.1 Frozen complex structure modulus

By considering a frozen complex structure modulus, $X^0 = 1$ and $F_0 = i$ and by turning on only those fluxes which contribute to the superpotential linearly on the moduli, we get

$$W = i\tilde{f}^0 + iSh_0 + iT_\alpha q_0^\alpha, \quad (\text{A.5})$$

which for an isotropic six-dimensional torus ($T_1 = T_2 = T_3 = T$) reduces by turning on only a single q -flux to

$$W = i\tilde{f}^0 + iSh_0 + iTq_0^1. \quad (\text{A.6})$$

By re-labeling the fluxes as $\tilde{f}^0 = \mathfrak{f}$, $h_0 = h$ and $q_0^1 = q$ we recover the superpotential in Eq.2.2. Notice at this point that $D5$ contributions to tadpole vanish by our selection of fluxes, but $D3$ and $D7$ -branes contribution do not.

Therefore, since the scalar potential is given by

$$V = \frac{M_{Pl}^4}{4\pi} e^K \left(K^{I\bar{J}} D_I W D_{\bar{J}} \bar{W} - 3|W|^2 \right) \quad (\text{A.7})$$

with K the Kähler potential given in Eq. 2.1. we have that [11]

$$V = \frac{M_{Pl}^4}{64\pi} \left(\frac{(hs - \mathfrak{f})^2}{s\tau^3} - \frac{2q(3hs + \mathfrak{f})}{s\tau^2} - \frac{5q^2}{3s\tau} + \frac{(hc + q\rho)^2}{s\tau^3} \right) \quad (\text{A.8})$$

with a minimum at $s = \mathfrak{f}/h$, $\tau = -6\mathfrak{f}/5q$ and $\theta = 0$ where $\theta = hc + q\rho$ a linear combination of axions. It follows that there is a flat direction on the orthogonal linear combinations of such axions. Notice that in order to stay on a physical region, fluxes at the minimum must satisfy that $s, \tau > 0$.

B Cosmological Observables and Slow-Roll Multi-field Inflation

In the multifield scenario, for a homogenous and isotropic space-time the scalar fields shall satisfy the Einstein-Friedman equations, which in the N-fold formalism are given by

$$H^2 = \frac{1}{3M_{pl}^2} \left[V(\phi^i) + \frac{1}{2} H^2 \mathcal{G}_{ij} \frac{\partial \phi^i}{\partial N} \frac{\partial \phi^j}{\partial N} \right], \quad (\text{B.1})$$

and

$$\frac{d^2 \phi^i}{dN^2} + \left(3 + \frac{1}{H} \frac{dH}{dN} \right) \frac{d\phi^i}{dN} + \frac{1}{H^2} \mathcal{G}^{ij} \partial_j V = 0, \quad (\text{B.2})$$

where ϕ^i represent the i-th field, $dN = Hdt$ and $\partial_i V = \partial V / \partial \phi_i$. In the present work we shall specialize in a constant non-canonical kinetic terms for the scalar fields determined by the metric \mathcal{G}_{ab} . This condition is obtained since the lightest fields are a linear combination of axions. As usual, inflation requires that the parameters

$$\epsilon = -\frac{\dot{H}}{H^2}, \quad (\text{B.3})$$

$$\eta = \frac{\dot{\epsilon}}{\epsilon H}, \quad (\text{B.4})$$

shall satisfy $\eta \ll 1$ and $\epsilon \ll 1$ during inflation. Thus, the number of e-folds is determined by

$$dN_e = Hdt, \quad (\text{B.5})$$

where its duration shall be around 40–60 in order to reproduce the observable universe. The explicit form of cosmological observables in the E-fold formalism are related to the scalar power spectrum [56] given by

$$P_S^{(\chi, \zeta)} = \frac{H^2}{4\pi^2} (A^{AB} N_A N_B)_{\phi=\phi(e)}, \quad (\text{B.6})$$

where N is given by the set of differential equations

$$\frac{dN_A}{dt} = -N_B P_A^B, \quad (\text{B.7})$$

subject to the final condition

$$N_A^F = - \left(\frac{H_A}{H_B F^B} \right). \quad (\text{B.8})$$

Thus, the spectral index is determined by the standard expression

$$n_S^{(\chi, \zeta)} - 1 = -2\epsilon - 2 \frac{A^{AB} N_A P_B^C N_C}{A^{AB} N_A N_B} + \frac{d_N A^{AB} N_A N_B}{A^{AB} N_A N_B}. \quad (\text{B.9})$$

Similarly the tensor to scalar ratio r is given by

$$r^{(\chi, \zeta)} = 8 \frac{1 - (1 + \alpha)\epsilon}{N^A N_A}, \quad (\text{B.10})$$

where $N^A = A^{AB} N_B$ and $\alpha = 0.7296$.

Non-Gaussianities coming from the three point correlation functions related to the bi-spectrum are parametrized by f_{NL} . Using the N-fold formalism [4, 9, 57] which relates the curvature perturbations to the difference of e-foldings of two constant time hypersurfaces, we have that

$$\delta\phi^A(\lambda, N_e) = \phi^A(\lambda + \delta\lambda, N_e) - \phi^A(\lambda, N_e) \quad (\text{B.11})$$

where λ is an integration constant and ϕ^A stands for a scalar field. Thus, the curvature perturbations can be expressed as variations of the number of e-foldings. In particular, the f_{NL} parameters is expressed generically as

$$f_{NL} = \frac{5}{6} \frac{N^A N^B N_{AB}}{(N^C N_C)^2}, \quad (\text{B.12})$$

where N_{AB} is the solution of the differential equations

$$\frac{dN_{AB}}{dt} = -N_{AC} P_B^C - N_{BC} P_A^C - N_C Q_{AB}^C, \quad (\text{B.13})$$

subject to the final condition

$$N_{AB}^F = - \left(\frac{U_{AB}}{H_C F^C} \right)_{\phi=\phi(e)}. \quad (\text{B.14})$$

Thus for two scalar fields we require to solve 16 differential equations. The explicitly form of A^{AB} , P_A^b , U_{AB} , H_A and F^A can be found in [56].

References

- [1] Alan H. Guth. The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems. *Phys. Rev.*, D23:347–356, 1981.
- [2] PAR Ade, N Aghanim, M Arnaud, F Arroja, M Ashdown, J Aumont, C Baccigalupi, M Ballardini, AJ Banday, RB Barreiro, et al. Planck 2015 results-xx. constraints on inflation. *Astronomy & Astrophysics*, 594:A20, 2016.
- [3] TJ Allen, B Grinstein, and MB Wise. Nongaussian density perturbations in inflationary cosmologies 1987 phys. *Lett. B*, 197:66.
- [4] Juan Martin Maldacena. Non-Gaussian features of primordial fluctuations in single field inflationary models. *JHEP*, 05:013, 2003.
- [5] P. A. R. Ade et al. Planck 2015 results. XVII. Constraints on primordial non-Gaussianity. *Astron. Astrophys.*, 594:A17, 2016.
- [6] D. S. Salopek. Cold dark matter cosmology with nonGaussian fluctuations from inflation. *Phys. Rev.*, D45:1139–1157, 1992.
- [7] Andrei D. Linde and Viatcheslav F. Mukhanov. Nongaussian isocurvature perturbations from inflation. *Phys. Rev.*, D56:R535–R539, 1997.
- [8] N. Bartolo, S. Matarrese, and A. Riotto. Nongaussianity from inflation. *Phys. Rev.*, D65:103505, 2002.
- [9] Francis Bernardeau and Jean-Philippe Uzan. NonGaussianity in multifield inflation. *Phys. Rev.*, D66:103506, 2002.
- [10] Thorsten Battefeld and Richard Easther. Non-Gaussianities in Multi-field Inflation. *JCAP*, 0703:020, 2007.
- [11] Ralph Blumenhagen, Anamaria Font, Michael Fuchs, Daniela Herschmann, Erik Plauschinn, Yuta Sekiguchi, and Florian Wolf. A Flux-Scaling Scenario for High-Scale Moduli Stabilization in String Theory. *Nucl. Phys.*, B897:500–554, 2015.
- [12] Fernando Marchesano, Gary Shiu, and Angel M. Uranga. F-term Axion Monodromy Inflation. *JHEP*, 09:184, 2014.
- [13] Ralph Blumenhagen, Daniela Herschmann, and Erik Plauschinn. The Challenge of Realizing F-term Axion Monodromy Inflation in String Theory. *JHEP*, 01:007, 2015.
- [14] Vijay Balasubramanian and Per Berglund. Stringy corrections to Kahler potentials, SUSY breaking, and the cosmological constant problem. *JHEP*, 11:085, 2004.
- [15] Vijay Balasubramanian, Per Berglund, Joseph P. Conlon, and Fernando Quevedo. Systematics of moduli stabilisation in Calabi-Yau flux compactifications. *JHEP*, 03:007, 2005.
- [16] Jessie Shelton, Washington Taylor, and Brian Wecht. Nongeometric flux compactifications. *JHEP*, 10:085, 2005.
- [17] Jessie Shelton, Washington Taylor, and Brian Wecht. Generalized Flux Vacua. *JHEP*, 02:095, 2007.
- [18] Brian Wecht. Lectures on Nongeometric Flux Compactifications. *Class. Quant. Grav.*, 24:S773–S794, 2007.

- [19] Gerardo Aldazabal, Diego Marques, and Carmen Nunez. Double Field Theory: A Pedagogical Review. *Class. Quant. Grav.*, 30:163001, 2013.
- [20] David S. Berman and Daniel C. Thompson. Duality Symmetric String and M-Theory. *Phys. Rept.*, 566:1–60, 2014.
- [21] David Andriot, Magdalena Larfors, Dieter Lust, and Peter Patalong. A ten-dimensional action for non-geometric fluxes. *JHEP*, 09:134, 2011.
- [22] David Geissbuhler, Diego Marques, Carmen Nunez, and Victor Penas. Exploring Double Field Theory. *JHEP*, 06:101, 2013.
- [23] Ralph Blumenhagen, Xin Gao, Daniela Herschmann, and Pramod Shukla. Dimensional Oxidation of Non-geometric Fluxes in Type II Orientifolds. *JHEP*, 10:201, 2013.
- [24] Anamaria Font, Adolfo Guarino, and Jesus M. Moreno. Algebras and non-geometric flux vacua. *JHEP*, 12:050, 2008.
- [25] Adolfo Guarino and George James Weatherill. Non-geometric flux vacua, S-duality and algebraic geometry. *JHEP*, 02:042, 2009.
- [26] Beatriz de Carlos, Adolfo Guarino, and Jesus M. Moreno. Complete classification of Minkowski vacua in generalised flux models. *JHEP*, 02:076, 2010.
- [27] Gerardo Aldazabal, Diego Marques, Carmen Nunez, and Jose A. Rosabal. On Type IIB moduli stabilization and $N = 4, 8$ supergravities. *Nucl. Phys.*, B849:80–111, 2011.
- [28] Giuseppe Dibitetto, Adolfo Guarino, and Diederik Roest. Charting the landscape of $N=4$ flux compactifications. *JHEP*, 03:137, 2011.
- [29] Johan Blaback, Ulf Danielsson, and Giuseppe Dibitetto. Fully stable dS vacua from generalised fluxes. *JHEP*, 08:054, 2013.
- [30] Cesar Damian, Luis R Díaz-Barrón, Oscar Loaiza-Brito, and Miguel Sabido. Slow-roll inflation in non-geometric flux compactification. *Journal of High Energy Physics*, 2013(6):109, 2013.
- [31] Cesar Damian and Oscar Loaiza-Brito. More stable de Sitter vacua from S-dual nongeometric fluxes. *Phys. Rev.*, D88(4):046008, 2013.
- [32] Falk Hassler, Dieter Lust, and Stefano Massai. On Inflation and de Sitter in Non-Geometric String Backgrounds. *Fortsch. Phys.*, 65(10-11):1700062, 2017.
- [33] Ralph Blumenhagen, Cesar Damian, Anamaria Font, Daniela Herschmann, and Rui Sun. The Flux-Scaling Scenario: De Sitter Uplift and Axion Inflation. *Fortsch. Phys.*, 64(6-7):536–550, 2016.
- [34] J. Blaback, U. H. Danielsson, G. Dibitetto, and S. C. Vargas. Universal dS vacua in STU-models. *JHEP*, 10:069, 2015.
- [35] Daniela Herschmann. *Large field inflation and moduli stabilisation in type IIB string theory*. PhD thesis, LMU Munich (main), 2017-05-17.
- [36] Nana Cabo Bizet, Oscar Loaiza-Brito, and Ivonne Zavala. Mirror quintic vacua: hierarchies and inflation. *JHEP*, 10:082, 2016.
- [37] Ralph Blumenhagen, Daniela Herschmann, and Florian Wolf. String Moduli Stabilization at the Conifold. *JHEP*, 08:110, 2016.

- [38] Anamaria Font. A flux-scaling scenario for moduli stabilization and axion inflation in string theory. In *Proceedings, 14th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Astrophysics, and Relativistic Field Theories (MG14) (In 4 Volumes): Rome, Italy, July 12-18, 2015*, volume 4, pages 4249–4255, 2017.
- [39] Ralph Blumenhagen, Anamaria Font, Michael Fuchs, Daniela Herschmann, and Erik Plauschinn. Large field inflation and string moduli stabilization. *PoS, PLANCK2015:021*, 2015.
- [40] Georges Obied, Hirosi Ooguri, Lev Spodyneiko, and Cumrun Vafa. De Sitter Space and the Swampland. 2018.
- [41] Prateek Agrawal, Georges Obied, Paul J. Steinhardt, and Cumrun Vafa. On the Cosmological Implications of the String Swampland. 2018.
- [42] Ana Achúcarro and Gonzalo A. Palma. The string swampland constraints require multi-field inflation. 2018.
- [43] Wilfried Buchmuller, Clemens Wieck, and Martin Wolfgang Winkler. Supersymmetric Moduli Stabilization and High-Scale Inflation. *Phys. Lett.*, B736:237–240, 2014.
- [44] S. Hosono, A. Klemm, and S. Theisen. Lectures on mirror symmetry. *Lect. Notes Phys.*, 436:235–280, 1994.
- [45] Yi Wang. Inflation, Cosmic Perturbations and Non-Gaussianities. *Commun. Theor. Phys.*, 62:109–166, 2014.
- [46] David Andriot. On the de Sitter swampland criterion. 2018.
- [47] Gia Dvali and Cesar Gomez. On Exclusion of Positive Cosmological Constant. 2018.
- [48] Sumit K. Garg and Chethan Krishnan. Bounds on Slow Roll and the de Sitter Swampland. 2018.
- [49] Alex Kehagias and Antonio Riotto. A note on Inflation and the Swampland. 2018.
- [50] Frederik Denef, Arthur Hebecker, and Timm Wrase. The dS swampland conjecture and the Higgs potential. 2018.
- [51] Christoph Roupec and Timm Wrase. de Sitter extrema and the swampland. 2018.
- [52] David Andriot. New constraints on classical de Sitter: flirting with the swampland. 2018.
- [53] Hiroki Matsui and Fuminobu Takahashi. Eternal Inflation and Swampland Conjectures. 2018.
- [54] Ido Ben-Dayan. Draining the Swampland. 2018.
- [55] Cesar Damian and Oscar Loaiza-Brito. Meromorphic Flux Compactification. *JHEP*, 04:141, 2017.
- [56] Xin Gao, Tianjun Li, and Pramod Shukla. Cosmological observables in multi-field inflation with a non-flat field space. *JCAP*, 1410(10):008, 2014.
- [57] Leonardo Senatore and Matias Zaldarriaga. The Effective Field Theory of Multifield Inflation. *JHEP*, 04:024, 2012.