

GLOBAL AND LOCAL PROPERTIES OF FINITE GROUPS WITH ONLY FINITELY MANY CENTRAL UNITS IN THEIR INTEGRAL GROUP RING

ANDREAS BÄCHLE, MAURICIO CAICEDO, ERIC JESPEERS, AND SUGANDHA MAHESHWARY

ABSTRACT. The aim of this article is to explore global and local properties of finite groups whose integral group rings have only trivial central units, so-called cut groups. We prove that for such groups, the number of rational valued irreducible characters coincides with the number of rational valued conjugacy classes. Further, we provide a natural criterion for nilpotent groups of class 2 to be cut and give a complete list of simple cut groups. Also, the impact of the cut property on Sylow 3-subgroups is discussed. We also collect substantial data on groups which indicates that the class of cut groups is surprisingly large.

1. INTRODUCTION

Let G be a finite group and let $\mathcal{Z}(\mathcal{U}(\mathbb{Z}G))$ denote the group of central units of the integral group ring $\mathbb{Z}G$. Clearly, $\mathcal{Z}(\mathcal{U}(\mathbb{Z}G))$ contains $\pm\mathcal{Z}(G)$, where $\mathcal{Z}(G)$ denotes the center of G . However, if $\mathcal{Z}(\mathcal{U}(\mathbb{Z}G)) = \pm\mathcal{Z}(G)$, i.e., all central units of $\mathbb{Z}G$ are trivial, then G is called a *cut group* or a group with the *cut property* [BMP17]. The question of classifying such groups, was seemingly first put up by Goodaire and Parmenter [GP86] and is included in the list of problems in Sehgal's book [Seh93, Problem 26]. In their pioneering article [RS90], Ritter and Sehgal could reformulate the requirement of the cut property on the center of $\mathcal{U}(\mathbb{Z}G)$ into a group theoretic property of G , also termed as RS-property (see [BMP18]). A remarkable consequence of Ritter and Sehgal's work is the fact that rational groups are cut groups, in particular symmetric groups S_n and Weyl group of complex Lie algebras are examples of cut groups. Over the last three decades, the study of cut groups has been taken up by several authors under different names and exhibit a highly interesting interplay between group theory, representation theory, algebraic number theory and even K -theory (see e.g. [MP18], Section 3 for a complete survey).

The original reason for studying cut groups is that for an arbitrary finite group G the unit group $\mathcal{U}(\mathbb{Z}G)$ has a subgroup of finite index that is generated by the central units and the units of reduced norm one. For many finite groups G the latter group is generated by very specific unipotent units (called bicyclic units). These, together with the Bass units (a natural generalization of cyclotomic units) “determine” finitely many generators of the center; note that the Bass units are in general not central units (for details we refer to [JdR15, Chapter 11]). It thus is a natural question to determine when the central units can be avoided to determine finitely many generators of a large subgroup of $\mathcal{U}(\mathbb{Z}G)$, i.e., characterize the cut groups.

Various interesting properties of cut groups are known. For instance, it is apparent from Higman's work [Hig40] that an abelian group G is cut if and only if the exponent of G divides 4 or 6. The work of Ritter and Sehgal [RS90] yields that the cut property is quotient closed. It is observed by Bakshi, Maheshwary and Passi [BMP17] that the center of a cut group is again a cut group, yet the cut property is not preserved under taking direct products. They also proved that every cut group has order divisible by 2 or 3. Furthermore, in recent explorations, it has been observed that the primes dividing the order of cut groups in certain classes are restricted [CD10, BMP17, Mah18, Bäck18], for instance, in the case of solvable cut groups to 2, 3, 5 and 7. It has also interested these authors to give a description of cut groups in various classes of finite solvable groups.

In the present work, we continue to examine the class of finite cut groups. After setting up the necessary background in Section 2, we begin by observing the impact of the cut property of a group

2010 *Mathematics Subject Classification.* 16S34, 16U60, 20E25.

Key words and phrases. central units, cut groups, cut property, inverse semi-rational, rational groups.

The first and the second author are postdoctoral researchers of the FWO (Research Foundation Flanders). The third author is supported in part by Onderzoeksraad of Vrije Universiteit Brussel and FWO (Research Foundation Flanders). The research of the fourth author is supported by DST (Department of Science and Technology), India (INSPIRE/04/2017/000897).

on its character table, see Section 3: We prove that for a cut group G , the number of rational valued irreducible characters of G equals the number of rational valued conjugacy classes of G by exhibiting a nice symmetry between the rows and columns of the character tables of such groups. An element-wise criterion is known for a nilpotent group to be a cut group [Mah18]. In Section 4, we provide a necessary and sufficient condition for a nilpotent group of class 2 to be a cut group. Recently, it has been observed that an infinite simple group is always a cut group [BMP18]. However, this is not true for finite simple groups. We provide a complete list of finite simple cut groups in Section 5. Further, in Section 6, we explore some local properties of cut groups: the impact of the cut property on group elements and the Sylow subgroups. In particular, we prove that Sylow 3-subgroups of cut groups are again cut, for several classes of groups, including all supersolvable groups. Finally, in Section 7, we put up the surprising data on existence of cut groups. For instance, 86.62% of all groups up to order 512 are cut.

2. PRELIMINARIES

Throughout the article, all groups considered are finite, unless otherwise stated explicitly. Let G be a group and $x \in G$ be an element of G . The order of G is denoted by $|G|$. The order of x is denoted by $o(x)$ and $C_G(x)$ denotes the centralizer of x in G . Let $y \in G$. Then, by $x \sim y$ we mean that “ x is conjugate to y in G ”, i.e., $x^g := g^{-1}xg = y$ for some $g \in G$ and $[x, y]$ denotes the commutator $x^{-1}y^{-1}xy = x^{-1}x^y$. By $H \leq G$ ($H \trianglelefteq G$), we indicate that H is a subgroup (normal subgroup). For $H \leq G$, $N_G(H)$ and $[G : H]$ respectively denote the normalizer and the index of H in G . If p is a prime, then $\text{Syl}_p(G)$ is the set of Sylow p -subgroups of G and the commutator subgroup of G is denoted by G' .

By $\text{Irr}(G)$ we denote the set of irreducible complex characters of G and for $\chi \in \text{Irr}(G)$, by $\mathbb{Q}(\chi)$ we denote the field extension of the rationals generated by the values of the character χ . Likewise, $\mathbb{Q}(x)$ denotes the field extension of the rationals generated by the values of all the character values at the element $x \in G$.

Recall that an element $x \in G$ is said to be *real*, if $x \sim x^{-1}$, equivalently $\mathbb{Q}(x) \subseteq \mathbb{R}$. Similarly, a character χ is called *real* if $\mathbb{Q}(\chi) \subseteq \mathbb{R}$. A group G is called *real* if all its elements are real; equivalently all its characters are real. It is common to call an element $x \in G$ *rational in G* , if $x^j \sim x$ for all j coprime to $o(x)$ and in this case $\mathbb{Q}(x) = \mathbb{Q}$. Likewise, a group G is called *rational*, if every element $x \in G$ is rational in G , which is equivalent to saying that every character χ of G is rational i.e., $\mathbb{Q}(\chi) = \mathbb{Q}$.

We briefly mention certain classes of groups that are related to cut groups. An element $x \in G$ is called *quadratic* if $[\mathbb{Q}(x) : \mathbb{Q}] = 2$, and, dually a character χ is called *quadratic* if $[\mathbb{Q}(\chi) : \mathbb{Q}] = 2$. If every irreducible character of G is either rational or quadratic, then it is said to be *quadratic rational*. Following [CD10], an element $x \in G$ is called *semi-rational in G* , if there exists an integer m such that $x^j \sim x$ or $x^j \sim x^m$ for every $j \in \mathbb{Z}$ coprime to $o(x)$ and x is called *inverse semi-rational in G* , if $m = -1$, i.e., $x^j \sim x$ or $x^j \sim x^{-1}$ for every such j . The group G is called *(inverse) semi-rational* if every element of G is (inverse) semi-rational in G .

We begin by stating the following equivalent criteria for a cut group that are essential for this article.

Proposition 2.1. (see [Bäc18, Proposition 2.2], [MP18, Theorem 5]) *For a group G , the following statements are equivalent:*

- (1) G is a cut group.
- (2) For every $x \in G$ and $j \in \mathbb{Z}$ with j coprime to $o(x)$, $x^j \sim x$ or $x^j \sim x^{-1}$, i.e., x is inverse semi-rational in G .
- (3) If $\mathbb{Q}G \simeq \bigoplus_{k=1}^m M_{n_k}(D_k)$ is the Wedderburn decomposition ($m, n_k \in \mathbb{Z}_{\geq 1}$, D_k division algebras), then for each k ,

$$\mathcal{Z}(D_k) \simeq \mathbb{Q}(\sqrt{-d})$$

for some $d = d(k) \in \mathbb{Z}_{\geq 0}$.

- (4) For each $\chi \in \text{Irr}(G)$, $\mathbb{Q}(\chi) = \mathbb{Q}(\sqrt{-d})$ for some $d = d(\chi) \in \mathbb{Z}_{\geq 0}$.

Hence the notion of a cut group and an inverse semi-rational group are the same. To avoid confusion, we always use the term inverse semi-rational in the local sense i.e., only for elements of the group and the term cut for a group. Note that the above proposition implies that every element

of order 1, 2, 3, 4 or 6 is inverse semi-rational in every group it is contained in; in particular all groups of exponent a divisor of 4 or 6 are cut.

There are other known equivalent characterizations of cut groups (for example a group G is cut if and only if the Whitehead group $K_1(\mathbb{Z}G)$ is finite), cf. [MP18, Theorem 5]. We will give yet another one in Lemma 3.2 below. It follows from Proposition 2.1 that the rational groups are cut groups. The cut groups are in the intersection of quadratic rational and semi-rational groups. Observe that the dihedral group of order 16 is both, quadratic rational and semi-rational, yet it is not cut. However, we shall see in Section 7 that cut groups constitute a big part of this intersection.

3. RATIONALITY

It is well-known that the number of real conjugacy classes of a group agrees with the number of its real irreducible characters, where the conjugacy class of x is called real if all its elements are real. Yet the analogous statement is not true if the field of reals is replaced by the field of rationals; there are groups as small as order 32 such that the number of rational conjugacy classes exceeds the number of rational irreducible characters and vice versa (see Example 3.4 below). It is an old theme in representation theory to detect classes of groups for which the number of rational conjugacy classes coincides with the number of rational irreducible characters. The numbers, of course, coincide in the case of rational groups. Navarro and Tiep proved that this is also true for groups with only one or two rational irreducible characters [NT10, Corollary 9.7, Theorem A] (using the Classification of Finite Simple Groups). Also for groups with cyclic Sylow 2-subgroups the number of rational conjugacy classes and rational characters coincide, as shown by Navarro and Tent [NT10]. A group with a cyclic Sylow 2-subgroup P can only be cut if $|P| \leq 4$, by Cayley's 2-complement theorem. An old result of Broshi shows that there is the same quantity of rational classes and characters if all Sylow subgroups are abelian [Bro71]. We show that the numbers in question behave well for cut groups.

Theorem 3.1. *Let G be a cut group. Then the number of rational irreducible characters of G equals the number of rational conjugacy classes of G .*

We begin by proving the following lemma.

Lemma 3.2. *Let G be a group. The following statements are equivalent:*

- (1) G is a cut group.
- (2) For each $x \in G$, $\mathbb{Q}(x) = \mathbb{Q}(\sqrt{-d})$ for some $d = d(x) \in \mathbb{Z}_{\geq 0}$.

Proof. Assume first that G is a cut group. Let $x \in G$, so that x is inverse semi-rational in G and in particular semi-rational in G . It thus follows from [Ten12, Lemma 1], that $[\mathbb{Q}(x) : \mathbb{Q}] \leq 2$. Now, if x is not rational, then for some $\chi \in \text{Irr}(G)$, $\mathbb{Q} \neq \mathbb{Q}(\chi(x)) \subseteq \mathbb{Q}(x)$ and thus degree considerations yield $\mathbb{Q}(\chi(x)) = \mathbb{Q}(x)$. Furthermore, $\mathbb{Q} \neq \mathbb{Q}(\chi(x)) \subseteq \mathbb{Q}(\chi)$ and by Proposition 2.1, $\mathbb{Q}(\chi) = \mathbb{Q}(\sqrt{-d})$, for some $d > 0$. Consequently, (2) follows.

Conversely, let $x \in G$. Since $\mathbb{Q}(x) = \mathbb{Q}(\sqrt{-d})$, $d \geq 0$, again by [Ten12, Lemma 1], we have that x is semi-rational in G , so that if j is a positive integer coprime to $o(x)$, then

$$x^j \sim x \text{ or } x^j \sim x^m, \text{ for some } m. \quad (\text{C})$$

Note that if x is real, then in view of the assumption $\mathbb{Q}(x) = \mathbb{Q}(\sqrt{-d})$, we obtain that $\mathbb{Q}(x) = \mathbb{Q}$, i.e., x is rational, so that $x^j \sim x$. Further, if x is not real, i.e., x is not conjugate to x^{-1} , then in view of (C), x^{-1} must be conjugate to x^m . Hence, (C) is equivalent to saying that for every positive integer j coprime to $o(x)$, $x^j \sim x$ or $x^j \sim x^{-1}$. Hence, G is a cut group. (Proposition 2.1). \square

The following well-known fact is a consequence of a result of R. Brauer on actions of groups on character tables [Isa76, (6.32)].

Proposition 3.3. *Let G be a group. The number of real irreducible characters of G and the number of real conjugacy classes of G coincide.*

Proof of Theorem 3.1. Observe firstly that in view of Lemma 3.2 and Proposition 2.1, there is a nice symmetry between the rows and the columns of the character table of a cut group as we have the following equivalences:

- (1) G is a cut group.
- (2) For each $\chi \in \text{Irr}(G)$, $\mathbb{Q}(\chi) = \mathbb{Q}(\sqrt{-d})$ for some $d = d(\chi) \in \mathbb{Z}_{\geq 0}$.
- (3) For each $x \in G$, $\mathbb{Q}(x) = \mathbb{Q}(\sqrt{-d})$ for some $d = d(x) \in \mathbb{Z}_{\geq 0}$.

Therefore, if G is a cut group, then an irreducible character of G is real if and only if it is rational, and similarly for the conjugacy classes. Hence, the result follows immediately from Proposition 3.3. \square

Note that Theorem 3.1 cannot be extended to the slightly larger class of semi-rational groups nor to the class of quadratic rational groups. In this case the nice symmetry between the rows and the columns of the character table may not hold true.

Indeed, by [Ten12, Lemma 1] a group is semi-rational if and only if all the fields $\mathbb{Q}(x)$, $x \in G$, have degree at most 2 over the rationals. However there are examples of semi-rational groups of order 384 (e.g. with SmallGroupID [384, 3283] in GAP [GAP18]) with an irreducible character χ such that $\mathbb{Q}(\chi) = \mathbb{Q}(\zeta_{24})$, where ζ_n denotes a primitive complex n^{th} root of unity, i.e., $[\mathbb{Q}(\chi) : \mathbb{Q}] = 2^3$. The group G of order 32 in Example 3.4 below also has an irreducible character with field of character values $\mathbb{Q}(\zeta_8)$ over the rationals.

Example 3.4. In [Ten12, Section 6], J. Tent gave the following two groups:

$$G = \langle a, b, c \mid a^2 = b^2 = c^8 = 1, b^c = b, b^a = bc^4, c^a = c^3 \rangle$$

and

$$H = \langle a, b, c \mid a^2 = b^2 = c^8 = 1, b^c = b, b^a = b, c^a = bc^3 \rangle$$

(having GAP SmallGroupIDs [32, 42] and [32, 9], respectively; the first one being a central extension of C_4 by a dihedral group of order 8, the latter of a Klein four group by a dihedral group of order 8). If we denote by $\text{Irr}_{\mathbb{Q}}(G)$ and $\text{ccl}_{\mathbb{Q}}(G)$ the set of rational irreducible characters and rational conjugacy classes of G , respectively, then we have $|\text{Irr}_{\mathbb{Q}}(G)| = 10 > |\text{ccl}_{\mathbb{Q}}(G)| = 8$ and $|\text{Irr}_{\mathbb{Q}}(H)| = 6 < |\text{ccl}_{\mathbb{Q}}(H)| = 8$. These are examples of smallest possible order for which the number of rational irreducible characters and the number of rational conjugacy classes differ. It may be noted that G is semi-rational and H is quadratic rational.

4. NILPOTENT CUT GROUPS

As mentioned earlier, it is well known that an abelian group is cut, if and only if its exponent divides 4 or 6. In this section, we provide a characterization on the quotients of G , for G to be a cut group for nilpotent groups of class 2. For an element wise criterion for a nilpotent group to be a cut group, see [BMP17, Mah18]. As the cut property is quotient closed and the center of a cut group is again a cut group, we necessarily have for a cut group G :

$$\text{for all } N \trianglelefteq G : \quad \exp(\mathcal{Z}(G/N)) \mid 4 \quad \text{or} \quad \exp(\mathcal{Z}(G/N)) \mid 6, \quad (\text{N})$$

where $\exp(H)$ denotes the exponent of a group H . However, rarely condition (N) is sufficient for G to be a cut group. For example (N) holds for all simple groups, but not all simple groups are cut (see Section 5). Also, dihedral groups of order $2p$, p a prime, satisfy (N), but these groups are not cut for $p \geq 5$ (and are of derived length 2). We prove that in case of nilpotent groups of class 2 the condition is sufficient for G to be a cut group.

Lemma 4.1. *Let G be a nilpotent group of class 2 satisfying one of the following properties:*

- (1) $\exp(\mathcal{Z}(G)) = 4$ and $\exp(G/G') \mid 4$ or
- (2) $\exp(\mathcal{Z}(G)) = 3$ and $\exp(G/G') = 3$.

If G' is cyclic, then G has the cut-property.

Proof. Assume (1). Since G is a 2-group, it is enough to show that for every $x \in G$, $x^4 \in [x, G] := \{[x, g] \mid g \in G\}$ ([Mah18], Corollary 3). For this, let $x \in G$. Clearly, if $x \in \mathcal{Z}(G)$ then $x^4 = 1 \in [x, G]$. So, assume that $x \notin \mathcal{Z}(G)$. Since $G' \leq \mathcal{Z}(G)$, we have the following two cases:

- (a) $|G'| = 2$. In this case, $1 \neq [x, G] \leq G'$, and therefore, $[x, G] = G'$ which implies $x^4 \in [x, G] = G'$, as $\exp(G/G') \mid 4$.
- (b) $|G'| = 4$. Now, if $[x, G] = G'$ then $\exp(G/G') \mid 4$ implies that $x^4 \in [x, G]$ and if $[x, G]$ is a proper subgroup of G' , then the order of x must divide 8. Indeed, let $g \in G$ such that $[x, g] \neq 1$, so that $[x, g]^2 = [x^2, g] = 1$. Hence $x^2 \in \mathcal{Z}(G)$, which implies, $x^8 = 1$. As $x^4 \in G'$ we have either $x^4 = 1$ or x^4 lies in the unique subgroup of order 2 of G' . Therefore, $x^4 \in [x, G]$ in both cases.

The proof of (2) is similar and we omit the details. \square

Lemma 4.2. *Let G be a p -group of nilpotency class 2. Then G is cut if and only if one of the following holds:*

- (1) $p = 2$ and, $\exp(\mathcal{Z}(G/N)) \mid 4$, for all $N \trianglelefteq G$.
- (2) $p = 3$ and, $\exp(\mathcal{Z}(G/N)) \mid 3$, for all $N \trianglelefteq G$.

Furthermore, G is rational if and only if $\exp(\mathcal{Z}(G/N)) \mid 2$, for all $N \trianglelefteq G$.

Proof. Clearly, if G is p -group which is cut, (1) or (2) holds. Furthermore, if G is rational, then G is a 2-group and $\mathcal{Z}(G/N)$ is an abelian rational group for every normal subgroup N of G . Since the only abelian rational groups are elementary abelian 2-groups, we have $\exp(\mathcal{Z}(G/N)) \mid 2$ for all $N \trianglelefteq G$.

We now prove the converse. Suppose first that G is a 2-group of class 2 such that $\exp(\mathcal{Z}(G/N)) \mid 4$, for all $N \trianglelefteq G$. We begin by observing that for such a group G ,

$$\exp(\mathcal{Z}(G)) \mid 2 \implies \exp(G) \mid 4. \quad (*)$$

For this, let $x \in G$. Then, for any $g \in G$, $[x, g] \in G' \leq \mathcal{Z}(G)$ and hence, $[x, g]^2 = [x^2, g] = 1$, which implies that $x^2 \in \mathcal{Z}(G)$, i.e., $x^4 = 1$. Now, let $M_n(D)$ be a Wedderburn component of $\mathbb{Q}G$. Let $K \trianglelefteq G$ be the kernel of the corresponding representation. Then G/K is faithfully embedded in $M_n(D)$, which is also a Wedderburn component of $\mathbb{Q}(G/K)$. We distinguish two cases.

- (a) $\exp(\mathcal{Z}(G/K)) \mid 2$: In this case, by statement, we have that (*), $\exp(G/K) \mid 4$, which implies that G/K is a cut group and hence $\mathcal{Z}(D)$ is $\mathbb{Q}(\sqrt{-d})$ with $d \in \mathbb{Z}_{\geq 0}$ (Proposition 2.1).
- (b) $\exp(\mathcal{Z}(G/K)) = 4$: Here, $\exp((G/K)/(G/K)') = \exp(G/G'K)$ divides 4, by hypothesis. Since G/K has nilpotency class at most two, $(G/K)' \leq \mathcal{Z}(G/K)$. Moreover, $(G/K)' \leq \mathcal{Z}(G/K)$ is cyclic as G/K is embedded in $M_n(D)$. Therefore, by Lemma 4.1, G/K is cut. Hence, the center of every Wedderburn component of $\mathbb{Q}G$ is $\mathbb{Q}(\sqrt{-d})$ with $d \in \mathbb{Z}_{\geq 0}$.

In case $p = 3$, the second part of Lemma 4.1 can be used to get the corresponding statement. The proof of the last statement also follows using similar arguments. \square

Theorem 4.3. *Let G be a nilpotent group of class at most 2. Then G is cut if, and only if, the exponent of $\mathcal{Z}(G/N)$ divides 4 or 6, for all $N \trianglelefteq G$.*

Proof. Clearly, the condition is necessary. For sufficiency, assume that G is a finite nilpotent group of class at most 2 and the exponent of $\mathcal{Z}(G/N)$ divides 4 or 6, for all $N \trianglelefteq G$. Observe that since $\exp(\mathcal{Z}(G))$ divides 4 or 6, the nilpotency of G yields that G is a $\{2, 3\}$ -group and $G = P_2 \times P_3$, where P_2 and P_3 denote its Sylow 2- and 3-subgroups, respectively. In view of ([Mah18], Theorem 3; [BMP17], Theorems 2 & 3), G is a cut group if, and only if both P_2 and P_3 are cut groups; and P_2 is rational, if P_3 is non-trivial. We first check that P_2 is a cut group. This follows from Lemma 4.2, as $\exp(\mathcal{Z}(P_2/N_2))$ divides 4, for every $N_2 \trianglelefteq P_2$. For if $N \trianglelefteq G$, then $N = N_2 \times N_3$, where $N_2 \trianglelefteq P_2$ and $N_3 \trianglelefteq P_3$; thus yielding, $\exp(\mathcal{Z}(P_2/N_2)) = \exp(\mathcal{Z}(G/N_2 \times P_3))$ which divides 4. Similarly, P_3 is also a cut group. Hence, it only remains to show that if P_3 is non-trivial, then P_2 is rational. Observe that $\exp(\mathcal{Z}(P_2/N_2)) = 2$ for every $N_2 \trianglelefteq P_2$, for if $\mathcal{Z}(P_2/N_2)$ contains an element x of order 4, then xy , where $y \in \mathcal{Z}(P_3) \leq \mathcal{Z}(G)$ is an element of order 3, is an element of order 12 in $\mathcal{Z}(G/N_2 \times 1)$, a contradiction. But then P_2 is rational by Lemma 4.2. \square

Remark 4.4. Note that for a nilpotent group G of nilpotency class exceeding 2, $\exp(\mathcal{Z}(G/N)) \mid 4$ or $\exp(\mathcal{Z}(G/N)) \mid 6$ for all $N \trianglelefteq G$ is no more sufficient to conclude that G is cut. For instance, if $G = D_{16}$, the dihedral group of order 16, we have $\exp(\mathcal{Z}(G/N)) \mid 2$ for every normal subgroup, yet the group is not cut. Similarly, if

$$G = \langle a, b, c \mid a^9 = b^3 = c^3 = 1, b^a = b, a^c = ab, b^c = a^3b \rangle \simeq (C_9 \times C_3) \rtimes C_3,$$

all centers of quotients of G are of exponent 3, but this group is again not cut (G has SmallGroupID [81, 8]).

5. SIMPLE CUT GROUPS

So far, the properties of solvable cut groups have been explored. A complete classification of finite metacyclic cut groups is given in [BMP17, Theorem 5]. A description of Frobenius cut groups can be found in [Bäc18, Theorem 1.3] (it turns out that those groups are always solvable). In this section, we give a complete classification of finite simple cut groups.

Theorem 5.1. *Let G be a simple group. Then G is cut if and only if it is isomorphic to one of the following groups:*

- (1) $C_2, C_3,$
- (2) $A_7, A_8, A_9, A_{12},$
- (3) $L_2(7), U_3(3), U_3(5), U_4(3), U_5(2), U_6(2), S_4(3), S_6(2), O_8^+(2),$
- (4) $M_{11}, M_{12}, M_{22}, M_{23}, M_{24}, Co_1, Co_2, Co_3, HS, McL, Th, M.$

Proof. The simple semi-rational groups are classified in [AD17, Theorem 1.1] (using the Classification of Finite Simple Groups); disregarding the alternating groups this is a finite list. The alternating groups that are cut are described in [Fer04] and [AKS08]. An inspection of the character tables of the remaining 45 groups (for example in ATLAS [CCN⁺85] or GAP) using Proposition 2.1 reveals that of those exactly the groups listed above are cut groups. \square

Note that in contrast to the case of finite simple groups, every infinite simple group I is a cut group i.e., has the property that $\mathcal{Z}(\mathcal{U}(ZI)) = \pm\mathcal{Z}(I)$, see [BMP18, Examples following Theorem 5].

6. LOCAL PROPERTIES OF CUT GROUPS

It was conjectured for a long time that being rational for 2-elements is governed by the Sylow 2-subgroup of a group. More precisely, already in Kletzing's book from 1984 [Kle84, p. 13] it is referred to as a "long standing conjecture" that the Sylow 2-subgroup of a rational group is again rational (recall that every non-trivial rational group is of even order and hence has a non-trivial Sylow 2-subgroup). Eventually in 2012, Isaacs and Navarro provided rational groups of order $2^9 \cdot 3$ with Sylow 2-subgroups that are not rational [IN12]. However, they also proved the conjecture for solvable groups with Sylow 2-subgroups of nilpotency class at most 2 [IN12, Theorem A].

Recall that every non-trivial cut group has order divisible by 2 or 3 [BMP17, Theorem 1] and one might wonder which properties are determined locally (i.e., in p -subgroups or their normalizers). It is not hard to find cut groups that have Sylow 2-subgroups that are not cut (e.g. the groups with SmallGroupID [384, 18033] and [384, 18040] of order $384 = 2^7 \cdot 3$). However, for the prime 3 things seem to behave differently. The following lemma shows that being inverse semi-rational for 3-elements is indeed a somehow local property.

Lemma 6.1. *Let G be a group and let $x \in G$ be a 3-element. Then x is inverse semi-rational in G if and only if x is inverse semi-rational in P for some $P \in \text{Syl}_3(G)$.*

Proof. Let y be an element of a group Y . Then $B_Y(y) := N_Y(\langle y \rangle)/C_Y(y)$ can naturally be identified with a subgroup of $\text{Aut}(\langle y \rangle)$, the automorphism group of $\langle y \rangle$. By [CD10, Lemma 5], the element y is inverse semi-rational in Y if and only if $B_Y(y)\langle \tau \rangle = \text{Aut}(\langle y \rangle)$, where $\tau: \langle y \rangle \rightarrow \langle y \rangle$ ($w \mapsto w^{-1}$) denotes the inversion automorphism of $\langle y \rangle$. Recall that the automorphism group of the cyclic group C_{3^f} of order 3^f is cyclic ($\text{Aut}(C_{3^f}) \simeq C_2 \times C_{3^f-1}$), and 2 is a primitive root modulo 3^f , i.e., 2 generates the unit group of the ring of integers modulo 3^f . Hence, the 3-element x is inverse semi-rational in an ambient group if and only if x is conjugate to x^4 in that group. Assume that the 3-element $x \in G$ of order 3^f is inverse semi-rational in G . Then there exists $g \in N_G(\langle x \rangle)$ such that $x^g = x^4$. Since the automorphism $w \mapsto w^4$ of $\langle x \rangle$ is of 3-power order, we may replace g by its 3-part, if necessary, and can assume that g also is a 3-element. Now, since g normalizes $\langle x \rangle$, the subgroup $S = \langle x, g \rangle$ is a 3-subgroup of G . By Sylow's theorem, S is contained in a Sylow 3-subgroup P of G and x is inverse semi-rational in P . The other implication is clear. \square

The above lemma is clearly false for p -elements, $p \geq 5$.

It may be observed that if G is a cut group and $P \in \text{Syl}_3(G)$ then P is cut if and only if for all $x \in P$ and for all $S \in \text{Syl}_3(G)$ containing x , x is inverse semi-rational in S . Lemma 6.1 asserts that each 3-element of a cut group is inverse semi-rational in at least one Sylow 3-subgroup. From the next lemma one can deduce that it is inverse semi-rational in all Sylow 3-subgroups, in certain cases.

Lemma 6.2. *Let G be a cut group and $P \in \text{Syl}_p(G)$ for some prime p . Then $\exp \mathcal{Z}(P) \mid p$, if p is odd and $\exp \mathcal{Z}(P) \mid 4$, if $p = 2$.*

Proof. Let $x \in \mathcal{Z}(P)$ be an element of order p^f . Then $P \leq C_G(x)$ and hence $B_G(x) = N_G(\langle x \rangle)/C_G(x)$ has an order not divisible by p . Note that $\text{Aut}(\langle x \rangle) \simeq C_{p^f-1} \times C_{p-1}$, in case p odd and $\text{Aut}(\langle x \rangle) \simeq$

$C_{2f-2} \times C_2$, for $p = 2$. Further, by [CD10, Lemma 5], we have that $[\text{Aut}(\langle x \rangle) : B_G(x)] \leq 2$ and thus the result follows. \square

Let p be a prime. For a group X denote by $O_p(X)$ and $O_{p'}(X)$ the maximal normal p -subgroup of X and maximal normal p' -subgroup of X , respectively. For a group G we can define the upper p -series by

$$1 \leq O_{p'}(G) \leq O_{p',p}(G) \leq O_{p',p,p'}(G) \leq \dots,$$

where $O_{p',p}(G)/O_{p'}(G) = O_p(G/O_{p'}(G))$, $O_{p',p,p'}(G)/O_{p',p}(G) = O_{p'}(G/O_{p',p}(G))$ and so on (alternating between p and p'). If this series terminates in G , then G is called p -solvable. In this case, the p -length of G is defined to be the number of occurrences of the symbol p in the subscript of the first group in the upper p -series of G that equals G . For example, G has p -length at most 1 if and only if it has normal subgroups M and N , $M \subseteq N$, such that M and G/N are p' -groups and N/M is a p -group. We prove the following:

Proposition 6.3. *Let G be a 3-solvable cut group of 3-length at most 1, $P \in \text{Syl}_3(G)$. Then P is cut.*

Proof. Since G has 3-length 1, G has normal subgroups $1 \leq M \leq N \leq G$ such that M and G/N are 3'-groups and N/M is a 3-group. Let $x \in P$. Then x is inverse semi-rational in G and hence also in some Sylow 3-subgroup Q of G by Lemma 6.1. Since N is a normal subgroup of 3'-index, Q is contained in N and hence x is inverse semi-rational in N . Note that M is a normal complement for P in N , that is, $N = MP$ is the semi-direct product of the normal subgroup M with P and the restriction to P of the projection map $\pi: N = MP \rightarrow P$ is the identity map on P . As x is inverse semi-rational in N , there is $n \in N$ such that $x^n = x^4$. Write $n = my$, $m \in M$, $y \in P$. Then $x^y = x^4$, since $\pi(n) = y$. Hence, x is inverse semi-rational in P , implying that P is cut. \square

The following proposition follows from Lemma 6.2 and Proposition 6.3:

Proposition 6.4. *Let G be a cut group and $P \in \text{Syl}_3(G)$. Then P is also cut, provided one of the following holds:*

- (1) P is abelian.
- (2) P is a normal subgroup of G .

In particular, a unique Sylow 3-subgroup of a cut group is always cut.

Remark 6.5. Using Cayley's 2-complement theorem together with the previous proposition we obtain: Let G be a cut group with order only divisible by the primes 2 and 3 and let $P \in \text{Syl}_3(G)$. If a Sylow 2-subgroup of G is cyclic, then P is cut.

Theorem 6.6. *Let G be a cut group and $P \in \text{Syl}_3(G)$. Then P is also cut, provided one of the following holds:*

- (1) G is supersolvable,
- (2) G is a Frobenius group,
- (3) G is simple,
- (4) G is of odd order and $O_3(G)$ is abelian.

Proof.

- (1) Supersolvable groups have a Sylow tower [Hup67, VI, Satz 9.1], so in particular p -length at most 1 for all primes p . The claim thus follows from Proposition 6.3.
- (2) Assume that G is a Frobenius group with Frobenius kernel F and Frobenius complement K . Since F and K have coprime order, a Sylow 3-subgroup P of G is isomorphic to a Sylow 3-subgroup of F or to a Sylow 3-subgroup of K . Assume that 3 divides $|F|$. Since F is nilpotent and characteristic in G , P is normal in G . Hence it is cut by Proposition 6.4. If 3 divides $|K|$, then P is cyclic by [Pas68, Theorem 18.1 (iv)] and it follows from Lemma 6.2 that it is also cut. (Of course, we could also have used the description of Frobenius cut groups in [Bäc18, Theorem 1.3].)
- (3) In Theorem 5.1 the simple cut groups were determined. They are given along with the nilpotency class of their Sylow 3-subgroup P in Table 1, showing that their nilpotency class can exceed 2. Using, for example GAP, one can check that these Sylow 3-subgroups are again cut.

TABLE 1. Simple cut groups G with the nilpotency class of their Sylow 3-subgroup P

G	C_2	C_3	A_7	A_8	A_9	A_{12}	$L_2(7)$	$U_3(3)$	$U_3(5)$
$\text{cl}(P)$	0	1	1	1	3	3	1	2	1

G	$U_4(3)$	$U_5(2)$	$U_6(2)$	$S_4(3)$	$S_6(2)$	$O_8^+(2)$	M_{11}	M_{12}	M_{22}
$\text{cl}(P)$	3	3	3	3	3	3	1	2	1

G	M_{23}	M_{24}	Co_1	Co_2	Co_3	HS	McL	Th	M
$\text{cl}(P)$	1	2	5	3	3	1	3	7	9

- (4) In case a group has odd order, then inverse semi-rational is the same as semi-rational by [CD10, Remark 13]. These groups have been described in [CD10, Theorem 3]. They come in three families: 3-groups, certain Frobenius groups and certain groups that are occasionally called 2-Frobenius or double Frobenius groups. Only in the last case we still need to verify the claim. By [CD10, Theorem 3 (2)] they have the following structure: $|G| = 7 \cdot 3^b$ and G contains a normal Frobenius subgroup of index 3 with $O_3(G)$ the Frobenius kernel. Moreover $O_3(G)T \in \text{Syl}_3(G)$ for some subgroup $T = \langle t \rangle$ of order 3 and $G/O_3(G)$ is the non-abelian group of order 21.

Assume that $O_3(G)$ is abelian. We aim to show that every element of the Sylow 3-subgroup $P = O_3(G)\langle t \rangle$ is inverse semi-rational in P . Firstly, we claim that in this setting $O_3(G)$ has exponent 3. Assume there is $y \in C_{O_3(G)}(t)$ of order 9. Then $[t, y] = 1$ and hence $t \in C_G(y)$. However, $C_G(y) = O_3(G)$: clearly $O_3(G) \leq C_G(y)$, since the former is abelian, but also since $O_3(G)F$ is a Frobenius group for every $F \in \text{Syl}_7(G)$, $C_G(y)$ cannot contain elements of order 7 and since y is inverse semi-rational in G , $[N_G(\langle y \rangle) : C_G(y)] = 3$, so indeed $C_G(y) = O_3(G)$. Summing up, $t \in C_G(y) = O_3(G)$. This yields a contradiction since $t \notin O_3(G)$, by the form of the Sylow 3-subgroup. So there is no element of order 9 in $C_{O_3(G)}(t)$ and $\exp(C_{O_3(G)}(t)) = 3$. By [KMS14, Lemma 2.4], $O_3(G) = \langle C_{O_3(G)}(t)^f \mid f \in F \rangle$ for some $F \in \text{Syl}_7(G)$, so also $\exp(O_3(G)) = 3$, since the latter is abelian. Thus every element of $O_3(G)$ is inverse semi-rational in $O_3(G)$ and hence also in $P \geq O_3(G)$.

Now assume that $x \in P \setminus O_3(G)$. Then P is the unique Sylow 3-subgroup of G containing x , because $O_3(G) < \langle O_3(G), x \rangle \leq P$ and $[P : O_3(G)] = 3$. Since x is inverse semi-rational in G , Lemma 6.1 asserts that x is inverse semi-rational in a Sylow 3-subgroup that contains it. Hence, x is inverse semi-rational in P . Consequently, every element of P is inverse semi-rational in P , i.e., P is cut. \square

Note that the class of groups in (4) contains groups of 3-length 2. We do not know of any example of a cut group of odd order to which the above theorem does not apply. Theorem 6.6 could also have been proved using the dual characterization of cut groups using characters.

Recall that a group G is called a *Camina group*, if $G' \neq G$ and, for every $g \notin G'$, the coset gG' is a conjugacy class. As Camina groups are either Frobenius or p -groups [DS96], we immediately have the following corollary from Theorem 6.6:

Corollary 6.7. *A Sylow 3-subgroup of a Camina cut group is again a cut group.*

Using the above results and GAP with the `grpconst` package [BE18], we have verified that the Sylow 3-subgroups of all cut groups of order at most 2000, as well as for order $2^2 \cdot 3^6$, $2^3 \cdot 3^6$ and $2^2 \cdot 3^7$, are again cut. Naturally, the following question arises:

Question 6.8. **Are Sylow 3-subgroups of cut groups again cut?**

Question 6.8 essentially asks whether we can always replace the existence assertion for the Sylow 3-subgroup in Lemma 6.1 by a “for all” statement.

Remark 6.9. Recall that (normal) subgroups of cut groups need not to be cut groups. But for classes for which Question 6.8 has a positive answer, the Sylow 3-subgroups have this property. In view of Proposition 2.1, this implies that for such groups, the centers of the Wedderburn components of the rational group algebra of cut groups also influences those of its Sylow 3-subgroups. For, if the centers

of the Wedderburn components of $\mathbb{Q}G$ are all rational or quadratic imaginary, then the same holds true for $\mathbb{Q}P$ for $P \in \text{Syl}_3(G)$.

7. EXISTENCE OF CUT GROUPS

In this section, we will give some indications that the class of cut groups is surprisingly large in all finite groups. Recall that a p -group can only be a cut group if $p \in \{2, 3\}$ [BMP17, Theorem 1]. We show that in these cases, the ratio of cut groups tends to one in the logarithmic sense.

Proposition 7.1. *Let $c(r)$ denote the number of cut groups of order r and $f(r)$ the number of all groups of order r . Then*


$$\lim_{n \rightarrow \infty} \frac{\ln c(p^n)}{\ln f(p^n)} = 1, \quad \text{for } p \in \{2, 3\}. \quad (**)$$

Proof. In their work on the asymptotic behavior of the number of p -groups, Higman, Sims, Newman and Seeley showed that for each prime p ,

$$p^{\frac{2}{27}m^2(m-6)} \leq f(p^m) \leq p^{\frac{2}{27}m^3 + O(m^{\frac{5}{2}})},$$

see [BNV07, Theorem 4.5, Theorem 5.7]. Noting that all the groups constructed by Higman to obtain the lower bound are actually cut groups in case $p = 2$ (as they are extensions of elementary abelian 2-groups by elementary abelian 2-groups), see [BNV07, Section 4], shows that $(**)$ holds for $p = 2$. By [BNV07, Theorem 19.3] it follows that the number of 3-groups of exponent 3 (and nilpotency class 2) has the same asymptotic behavior in the leading term, so $(**)$ also holds for $p = 3$. \square

FIGURE 1. Rational and cut groups of order at most 1023



Remark 7.2. Though the above proposition shows that there are many cut groups in the class of p -groups, it says nothing about the actual percentage. For groups of small order we now also support this with numerical data which we obtained using GAP and the SgIPPOw package [EVL14]. Table 2 lists the number and percentage of rational as well as cut 2-groups up to order 2^9 . Observe that for the 2-groups, the percentage of cut groups increases again from order 2^8 . The number and percentage of cut 3-groups up to order 3^8 are enlisted in Table 3 (note that the only rational group of odd order is the trivial one). Also, in Table 4, we list the number and percentage of rational and cut groups of selected mixed orders. Figure 1 presents a chart showing the percentage of groups that are rational and cut, up to order 1023. The horizontal and the vertical axes indicate the order and the percentage respectively. The dashed graph gives the percentage of rational groups up to that order whereas the solid graph indicates the percentages of cut groups up to that order. Note that the upward bumps for the percentage of cut groups appear at 2-powers, whereas the (visible) bumps downwards for this percentage happen at orders of the form $2^a \cdot 3$. Yet also for these orders the cut groups are still surprisingly numerous, cf. Table 4. Also, when considering all groups up to a certain small order, cut groups are not negligible. For instance, about 86.62% of the groups of order at most 512 and 78.55% of groups of order at most 1023 are cut groups, whereas 0.57% of the groups of order at most 512 and 0.52% of groups of order at most 1023 are rational.

TABLE 2. Rational and cut groups in small order 2-groups

2^n	number of all groups of order 2^n	number of rational groups of order 2^n <i>percentage of rational groups of order 2^n</i>	number of cut groups of order 2^n <i>percentage of cut groups of order 2^n</i>
$2^2 = 4$	2	1 50%	2 100%
$2^3 = 8$	5	3 60%	4 80%
$2^4 = 16$	14	3 21.43%	10 71.43%
$2^5 = 32$	51	10 19.61%	33 64.71%
$2^6 = 64$	267	30 11.24%	161 60.30%
$2^7 = 128$	2 328	124 5.33%	1 349 57.95%
$2^8 = 256$	56 092	748 1.34%	37 593 67.02%
$2^9 = 512$	10 494 213	59 514 0.57%	9 127 858 86.98%

TABLE 3. Cut groups in small order 3-groups

3^n	number of all groups of order 3^n	number of cut groups of order 3^n <i>percentage of cut groups of order 3^n</i>
$3^2 = 9$	2	1 50%
$3^3 = 27$	5	3 60%
$3^4 = 81$	15	4 26.67%
$3^5 = 243$	67	14 20.90%
$3^6 = 729$	504	96 19.05%
$3^7 = 2187$	9 310	595 6.39%
$3^8 = 6561$	1 396 077	66 312 5.06%

TABLE 4. Rational and cut groups for some selected mixed orders

m	number of all groups of order m	number of rational groups of order m <i>percentage of rational groups of order m</i>	number of cut groups of order m <i>percentage of cut groups of order m</i>
$2^2 \cdot 5 = 20$	5	0 0%	1 20%
$2 \cdot 3 \cdot 7 = 42$	6	0 0%	2 33.33%
$2^6 \cdot 3 = 192$	1 543	18 1.17%	318 20.61%
$2^7 \cdot 3 = 384$	20 169	317 0.32%	2 279 11.30%
$2^3 \cdot 5^2 = 400$	221	1 0.45%	12 5.43%
$2^6 \cdot 3^2 = 576$	8 681	45 0.52%	1 074 12.37%
$2^3 \cdot 3 \cdot 7^2 = 588$	66	0 0%	4 6.06%
$2^3 \cdot 3^4 = 648$	757	11 1.45%	151 19.95%
$2^8 \cdot 3 = 768$	1 090 235	304 0.03%	64 765 5.94%
$2 \cdot 3^6 = 1458$	1 798	1 0.06%	387 21.52%
$2^3 \cdot 3^2 \cdot 5^2 = 1800$	749	0 0%	14 1.87%
$2^3 \cdot 3^5 = 1944$	3 973	17 0.43%	525 13.21%

Acknowledgements:

We are grateful to Urban Jezernik, Gabriel Navarro, Inder Bir Singh Passi and Ángel del Río for very interesting discussions. We want to thank Jürgen Müller for providing us with the Sylow 3-subgroups of the sporadic simple groups. The fourth author is thankful to Eric Jespers and FWO for the local hospitality provided for stay at VUB which played a vital role in the outcome of this paper.

REFERENCES

- [AD17] S.H. Alavi and A. Daneshkhah, *On semi-rational finite simple groups*, *Monatsh. Math.* **184** (2017), no. 2, 175–184.
- [AKS08] R. Zh. Aleev, A. V. Kargapolov, and V. V. Sokolov, *The ranks of central unit groups of integral group rings of alternating groups*, *Fundam. Prikl. Mat.* **14** (2008), no. 7, 15–21.
- [Bäc18] A. Bächle, *Integral group rings of solvable groups with trivial central units*, *Forum Math.* **30** (2018), no. 4, 845–855.
- [BE18] H.U. Besche and B. Eick, *Grpconst*, <https://www.gap-system.org/Packages/grpconst.html>, 2018, GAP package, Version 2.6.
- [BMP17] G. K. Bakshi, S. Maheshwary, and I. B. S. Passi, *Integral group rings with all central units trivial*, *J. Pure Appl. Algebra* **221** (2017), no. 8, 1955–1965.
- [BMP18] ———, *Group rings and the RS-property*, to appear in *Communications in Algebra* (2018).
- [BNV07] S.R. Blackburn, P.M. Neumann, and G. Venkataraman, *Enumeration of finite groups*, *Cambridge Tracts in Mathematics*, vol. 173, Cambridge University Press, Cambridge, 2007.
- [Bro71] A.M. Broshi, *Galois correspondences between the irreducible characters and the conjugacy classes of finite groups*, *J. Algebra* **19** (1971), 441–451. MR 0291308
- [CCN+85] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of finite groups*, Oxford University Press, Eynsham, 1985, Maximal subgroups and ordinary characters for simple groups, With computational assistance from J. G. Thackray.
- [CD10] D. Chillag and S. Dolfi, *Semi-rational solvable groups*, *J. Group Theory* **13** (2010), no. 4, 535–548.
- [DS96] R. Dark and C. M. Scoppola, *On Camina groups of prime power order*, *J. Algebra* **181** (1996), no. 3, 787–802.
- [EVL14] B. Eick and M. Vaughan-Lee, *SglPPow*, <https://gap-packages.github.io/sglppow/>, 2014, GAP package, Version 1.1.
- [Fer04] R. A. Ferraz, *Simple components and central units in group algebras*, *J. Algebra* **279** (2004), no. 1, 191–203.
- [GAP18] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.9.1*, 2018, <https://www.gap-system.org>.
- [GP86] E. G. Goodaire and M. M. Parmenter, *Units in alternative loop rings*, *Israel J. Math.* **53** (1986), no. 2, 209–216.
- [Hig40] G. Higman, *The units of group-rings*, *Proc. London Math. Soc. (2)* **46** (1940), 231–248.
- [Hup67] B. Huppert, *Endliche Gruppen. I*, *Die Grundlehren der Mathematischen Wissenschaften, Band 134*, Springer-Verlag, Berlin-New York, 1967.
- [IN12] I. M. Isaacs and G. Navarro, *Sylow 2-subgroups of rational solvable groups*, *Math. Z.* **272** (2012), no. 3-4, 937–945.
- [Isa76] I.M. Isaacs, *Character theory of finite groups*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1976, Pure and Applied Mathematics, No. 69.
- [JdR15] E. Jespers and Á. del Río, *Group Ring Groups*, Volume 1: Orders and Generic Constructions of Units, De Gruyter, Berlin-Boston, 2015.
- [Kle84] D. Kletzing, *Structure and representations of \mathbf{Q} -groups*, *Lecture Notes in Mathematics*, vol. 1084, Springer-Verlag, Berlin, 1984.
- [KMS14] E. Khukhro, N. Makarenko, and P. Shumyatsky, *Frobenius groups of automorphisms and their fixed points*, *Forum Math.* **26** (2014), no. 1, 73–112.
- [Mah18] S. Maheshwary, *Integral Group Rings With All Central Units Trivial: Solvable Groups*, *Indian J. Pure Appl. Math.* **49** (2018), no. 1, 169–175.
- [MP18] S. Maheshwary and I. B. S. Passi, *The upper central series of the unit groups of integral group rings: a survey*, 2018, to appear in *Indian Statistical Institute Series*, Springer.
- [NT10] G. Navarro and J. Tent, *Rationality and Sylow 2-subgroups*, *Proc. Edinb. Math. Soc. (2)* **53** (2010), no. 3, 787–798.
- [Pas68] D. Passman, *Permutation groups*, W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [RS90] J. Ritter and S. K. Sehgal, *Integral group rings with trivial central units*, *Proc. Amer. Math. Soc.* **108** (1990), no. 2, 327–329.
- [Seh93] S. K. Sehgal, *Units in integral group rings*, *Pitman Monographs and Surveys in Pure and Applied Mathematics*, vol. 69, Longman Scientific & Technical, Harlow, 1993, With an appendix by Al Weiss.
- [Ten12] J. Tent, *Quadratic rational solvable groups*, *J. Algebra* **363** (2012), 73–82.

(ANDREAS BÄCHLE) VAKGROEP WISKUNDE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSELS, BELGIUM
E-mail address: abachle@vub.be

(MAURICIO CAICEDO) VAKGROEP WISKUNDE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSELS, BELGIUM
E-mail address: mcaicedo@vub.be

(ERIC JESPERS) VAKGROEP WISKUNDE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSELS, BELGIUM
E-mail address: eric.jespers@vub.be

(SUGANDHA MAHESHWARY) INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, MOHALI, SECTOR 81, MOHALI (PUNJAB)-140306, INDIA
E-mail address: sugandha@iisermohali.ac.in