# Infinitely many solutions for a Hénon-type system in hyperbolic space

Patrícia Leal da Cunha \*

Departamento de Tecnologia e Ciência de Dados, FGV-SP, Brazil,

Flávio Almeida Lemos<sup>†</sup>

Departamento de Matemática, UFOP-MG, Brazil.

ABSTRACT. This paper is devoted to study the semilinear elliptic system of Hénon-type

$$\begin{cases} -\Delta_{\mathbb{B}^N} u = K(d(x))Q_u(u,v), \\ -\Delta_{\mathbb{B}^N} v = K(d(x))Q_v(u,v), \\ u, v \in H^1_r(\mathbb{B}^N), N \ge 3 \end{cases}$$

in the hyperbolic space  $\mathbb{B}^N$ , where  $H^1_r(\mathbb{B}^N) = \{u \in H^1(\mathbb{B}^N) : u \text{ is radial}\}$  and  $-\Delta_{\mathbb{B}^N}$  denotes the Laplace-Beltrami operator on  $\mathbb{B}^N$ ,  $d(x) = d_{\mathbb{B}^N}(0, x)$ ,  $Q \in C^1(\mathbb{R} \times \mathbb{R}, \mathbb{R})$  is p-homogeneous function and  $K \geq 0$  is a continuous function. We prove a compactness result and together with the Clark's theorem we establish the existence of infinitely many solutions.

Keywords. Hyperbolic space, Hénon equation Mathematics Subject Classification 2010. 58J05; 35J60; 58E30; 35J20; 35J75

## 1 Introduction and main result

This article concerns the existence of infinitely many solutions for the following semilinear elliptic system of Hénon type in hyperbolic space

$$\begin{cases} -\Delta_{\mathbb{B}^N} u = K(d(x))Q_u(u,v) \\ -\Delta_{\mathbb{B}^N} v = K(d(x))Q_v(u,v) \\ u,v \in H^1_r(\mathbb{B}^N), N \ge 3 \end{cases}$$
(\mathcal{H})

where  $\mathbb{B}^N$  is the Poincaré ball model for the hyperbolic space,  $H^1_r(\mathbb{B}^N)$  denotes the sobolev space of radial  $H^1(\mathbb{B}^N)$  function,  $r = d(x) = d_{\mathbb{B}^N}(0, x)$ ,  $\Delta_{\mathbb{B}^N}$  is the Laplace-Beltrami type operator on  $\mathbb{B}^N$ .

We assume the following hypothesis on  ${\cal K}$  and  ${\cal Q}$ 

 $(K_1)$   $K \ge 0$  is a continuous function with K(0) = 0 and  $K \ne 0$  in  $\mathbb{B}^N \setminus \{0\}$ .

<sup>\*</sup>E-mail addresses: patcunha80@gmail.com

 $<sup>^{\</sup>dagger}\textit{E-mail addresses: flavioal@gmail.com}$ 

- $(K_2) \ K = O(r^{\beta}) \text{ as } r \to 0 \text{ and } K = O(r^{\beta}) \text{ as } r \to \infty, \text{ for some } \beta > 0.$
- (Q<sub>1</sub>)  $Q \in C^1(\mathbb{R} \times \mathbb{R}, \mathbb{R})$  is such that  $Q(-s, t) = Q(s, -t) = Q(s, t), Q(\lambda s, \lambda t) = \lambda^p Q(s, t)$ (Q is p - homogeneous),  $\forall \lambda \in \mathbb{R}$  and  $p \in (2, \delta)$ , where

$$\delta = \begin{cases} \frac{2N+2\beta}{N-2} & \text{if } N-2 > 0\\ \infty & \text{if otherwise} \end{cases}$$

- (Q<sub>2</sub>) There exist  $C, C_1, C_2 > 0$  such that  $Q(s,t) \leq C(s^p + t^p), Q_s(s,t) \leq C_1 s^{p-1}$  and  $Q_t(s,t) \leq C_2 t^{p-1}, \ \forall s,t \geq 0.$
- (Q<sub>3</sub>) There exists  $C_3 > 0$  such that  $C_3(|s|^p + |t|^p) \le Q(s,t)$  with  $p \in (2, \delta)$ .

In the past few years the prototype problem

$$-\Delta_{\mathbb{B}^N} u = d(x)^{\alpha} |u|^{p-2} u, \quad u \in H^1_r(\mathbb{B}^N)$$

has been attracted attention. Unlike the corresponding problem in the Euclidean space  $\mathbb{R}^N$ , He in [10] proved the existence of a positive solution to the above problem over the range  $p \in (2, \frac{2N+2\alpha}{N-2})$  in the hyperbolic space. More precisely, she explored the Strauss radial estimate for hyperbolic space together with the Mountain Pass Theorem. In a subsequent paper [11], she proved the existence of at least one non-trivial positive solution for the critical Hénon equation

$$-\Delta_{\mathbb{B}^N} u = d(x)^{\alpha} |u|^{2^* - 2} u + \lambda u, \quad u \ge 0, \quad u \in H^1_0(\Omega'),$$

provided that  $\alpha \to 0^+$  and for a suitable value of  $\lambda$ , where  $\Omega'$  is a bounded domain in hyperbolic space  $\mathbb{B}^N$ . Finally, by working in the hole hyperbolic space  $\mathbb{H}^N$ , He [12] considered the following Hardy-Hénon type system

$$\begin{cases} -\Delta_{\mathbf{H}^N} u = d_b(x)^{\alpha} |v|^{p-1} v \\ -\Delta_{\mathbf{H}^N} v = d_b(x)^{\beta} |u|^{q-1} u, \end{cases}$$

for  $\alpha, \beta \in \mathbb{R}, N > 4$  and obtained infinitely many non-trivial radial solutions.

We would like to mention the paper of Carrião, Faria and Miyagaki [3] where they extended He's result by considering a general nonlinearity

$$\begin{cases} -\Delta_{\mathbb{B}^N}^{\alpha} u = K(d(x))f(u) \\ u \in H_r^1(\mathbb{B}^N). \end{cases}$$
(1)

They were able to prove the existence of at least one positive solution through a compact Sobolev embedding with the Mountain Pass Theorem.

In this paper we investigate the existence of infinitely many solutions by considering a gradient system that generalizes problem (1). We cite [2, 4, 7, 8, 9, 14, 15] for related gradient systems problems. In order to obtain our result, we applied the Clark's theorem [5, 6] and get inspiration on the nonlinearities condition employed by Morais Filho and Souto [13] in a p-laplacian system defined on a bounded domain in  $\mathbb{R}^N$ . As regarding the difficulties, many technical difficulties arise when working on  $\mathbb{B}^N$ , which is a non compact manifold. This means that the the embedding  $H^1(\mathbb{B}^N) \hookrightarrow L^p(\mathbb{B}^N)$  is not compact for  $2 \leq p \leq \frac{2N}{N-2}$  and the functional related to the system  $\mathcal{H}$  cannot satisfy the  $(PS)_c$  condition for all c > 0.

We also point out that since the weight function d(x) depends on the Riemannian distance r from a pole o, we have some difficulties in proving that

$$\int_{\mathbb{B}^N} d(x)^{\beta} \left( |u(x)|^p + |v(x)|^p \right) dV_{\mathbb{B}^N} < \infty, \quad \forall (u,v) \in H^1(\mathbb{B}^N) \times H^1(\mathbb{B}^N)$$

leading to a great effort in proving that the Euler-Lagrange functional associated is well defined.

To overcome these difficulties we restrict ourselves to the radial functions.

Our result is

**Theorem 1.1.** Under hypotheses  $(K_1)$ - $(K_2)$  and  $(Q_1)$ - $(Q_3)$ , the problem  $(\mathcal{H})$  has infinitely many solutions.

# 2 Preliminaries

Throughout this paper, C is a positive constant which may change from line to line.

The Poincaré ball for the hyperbolic space is

$$\mathbb{B}^N = \{ x \in \mathbb{R}^N | |x| < 1 \}$$

endowed with Riemannian metric g given by  $g_{i,j} = (p(x))^2 \delta_{i,j}$  where  $p(x) = \frac{2}{1-|x|^2}$ . We denote the hyperbolic volume by  $dV_{\mathbb{B}^N} = (p(x))^N dx$ . The geodesic distance from the origin to  $x \in \mathbb{B}^N$  is given by

$$d(x) := d_{\mathbb{B}^N}(0, x) = \int_0^{|x|} \frac{2}{1 - s^2} ds = \log\left(\frac{1 + |x|}{1 - |x|}\right).$$

The hyperbolic gradient and the Laplace-Beltrami operator are

$$-\Delta_{\mathbb{B}^N} u = -(p(x))^{-N} div(p(x)^{N-2} \nabla u), \quad \nabla_{\mathbb{B}^N} u = \frac{\nabla u}{p(x)}$$

where  $H^1(\mathbb{B}^N)$  denotes the Sobolev space on  $\mathbb{B}^N$  with the metric g.  $\nabla$  and div denote the Euclidean gradient and divergence in  $\mathbb{R}^N$ , respectively.

Let  $H^1_r(\mathbb{B}^N) = \{ u \in H^1(\mathbb{B}^N) : u \text{ is radial} \}.$ 

We shall find weak-solutions of problem  $(\mathcal{H})$  in the space

$$H = H^1_r(\mathbb{B}^N) \times H^1_r(\mathbb{B}^N)$$

endowed with the norm

$$\|(u,v)\|^{2} = \int_{\mathbb{B}^{N}} \left( \|\nabla_{\mathbb{B}^{N}} u\|_{\mathbb{B}^{N}}^{2} + \|\nabla_{\mathbb{B}^{N}} v\|_{\mathbb{B}^{N}}^{2} \right) dV_{\mathbb{B}^{N}}.$$

One can observe that system  $(\mathcal{H})$  is formally derived as the Euler-Lagrange equation for the functional

$$I(u,v) = \frac{1}{2} \int_{\mathbb{B}^N} (\|\nabla_{\mathbb{B}^N} u\|_{\mathbb{B}^N}^2 + \|\nabla_{\mathbb{B}^N} v\|_{\mathbb{B}^N}^2) \, dV_{\mathbb{B}^N} - \int_{\mathbb{B}^N} K(d(x))Q(u,v) \, dV_{\mathbb{B}^N}.$$

We endowed the norm for  $L^p(\mathbb{B}^N) \times L^p(\mathbb{B}^N)$  as

$$||(u,v)||_p^p = \int_{\mathbb{B}^N} (|u|^p + |v|^p) \, dV_{\mathbb{B}^N}.$$

To solve this problem we need the following lemmas.

**Lemma 2.1.** The map  $(u,v) \mapsto (d(x)^m u, d(x)^m v)$  from  $H = H^1_r(\mathbb{B}^N) \times H^1_r(\mathbb{B}^N)$  to  $L^p(\mathbb{B}^N) \times L^p(\mathbb{B}^N)$  is continuous for  $p \in (2, \tilde{m})$ , where m > 0 and

$$\tilde{m} = \begin{cases} \frac{2N}{N-2-2m} & \text{if } m < \frac{N-2}{2}, \\ \infty & \text{if otherwise.} \end{cases}$$

PROOF. In [10, Lemma 2.2] it has been proved that the map  $u \mapsto d(x)^m u$  from  $H^1_r(\mathbb{B}^N)$  to  $L^p(\mathbb{B}^N)$  is continuous for  $p \in (2, \tilde{m})$ . Therefore  $\|d(x)^m u\|_p \leq C \|u\|_{H^1_r}$  and  $\|d(x)^m v\|_p \leq C \|v\|_{H^1_r}$ . Hence,

$$\left( \|d(x)^m u\|_p^2 + \|d(x)^m v\|_p^2 \right)^{\frac{1}{2}} \le C \|(u,v)\|.$$

Now observe that

$$\begin{aligned} \|d(x)^{m}u\|_{p} + \|(d(x)^{m}v)\|_{p} &= \left[ \left( \|d(x)^{m}u\|_{p} + \|d(x)^{m}v\|_{p} \right)^{2} \right]^{\frac{1}{2}} \\ &= \left( \|d(x)^{m}u\|_{p}^{2} + 2 \|d(x)^{m}u\|_{p} \|d(x)^{m}v\|_{p} + \|d(x)^{m}v\|_{p} \right)^{\frac{1}{2}}. \end{aligned}$$

Applying the Cauchy's inequality  $ab \leq \frac{a^2+b^2}{2}$ , we get

$$\|d(x)^{m}u\|_{p} + \|d(x)^{m}v\|_{p} \le \sqrt{2} \Big( \|d(x)^{m}u\|_{p}^{2} + \|d(x)^{m}v\|_{p}^{2} \Big)^{\frac{1}{2}}$$

By the subaditivity,

$$\left( \left\| d(x)^m u \right\|_p^p + \left\| d(x)^m v \right\|_p^p \right)^{\frac{1}{p}} \le \left\| d(x)^m u \right\|_p + \left\| d(x)^m v \right\|_p$$

Therefore,

$$\|(d(x)^m u, d(x)^m v)\|_p \le C \|(u, v)\|$$

and the lemma holds.

**Remark 2.1.** From the previous lemma, there exists a positive constant C > 0, such that

$$\begin{split} \int_{\mathbb{B}^N} d(x)^{\beta} |u(x)|^p dV_{\mathbb{B}^N} + \int_{\mathbb{B}^N} d(x)^{\beta} |v(x)|^p dV_{\mathbb{B}^N} \\ &\leq C \left( \int_{\mathbb{B}^N} \|\nabla_{\mathbb{B}^N} u\|_{\mathbb{B}^N}^2 dV_{\mathbb{B}^N} + \int_{\mathbb{B}^N} \|\nabla_{\mathbb{B}^N} v\|_{\mathbb{B}^N}^2 dV_{\mathbb{B}^N} \right)^{\frac{p}{2}} \\ & \text{where } m = \frac{\beta}{p} \text{ and } 2$$

**Lemma 2.2.** The map  $(u,v) \mapsto (d(x)^m u, d(x)^m v)$  from  $H = H^1_r(\mathbb{B}^N) \times H^1_r(\mathbb{B}^N)$  to  $L^p(\mathbb{B}^N) \times L^p(\mathbb{B}^N)$  is compact for  $p \in (2, \tilde{m})$ , where m > 0 and

$$\tilde{m} = \begin{cases} \frac{2N}{N-2-2m} & \text{if } m < \frac{N-2}{2} \\ \infty & \text{if otherwise} \end{cases}$$

PROOF. Let  $(u_n, v_n) \in H$  be a bounded sequence. Then up to a subsequence, if necessary, we may assume that

$$(u_n, v_n) \rightharpoonup (u, v)$$

It is easy to see that  $u_n \rightharpoonup u$  and  $v_n \rightharpoonup v$  in  $H^1_r(\mathbb{B}^N)$ .

We will use the same calculus used by Haiyang He [10] (page 26). We want to show that

$$\lim_{n \to \infty} \int_{\mathbb{B}^N} d(x)^{mq} |u_n(x)|^p dV_{\mathbb{B}^N} = \int_{\mathbb{B}^N} d(x)^{mp} |u(x)|^p dV_{\mathbb{B}^N},$$
$$\lim_{n \to \infty} \int_{\mathbb{B}^N} d(x)^{mq} |v_n(x)|^p dV_{\mathbb{B}^N} = \int_{\mathbb{B}^N} d(x)^{mp} |v(x)|^p dV_{\mathbb{B}^N}.$$

Let  $u \in H^1_r(\mathbb{B}^N)$ , then by Haiyang He [10] we have

$$\begin{aligned} |u_n(x)| &\leq \frac{1}{\sqrt{\omega_{n-1}(N-2)}} \left(\frac{1-|x|^2}{2}\right)^{\frac{N-2}{2}} \frac{1}{|x|^{\frac{N-2}{2}}} \|u_n\|_{H^1_r(\mathbb{B}^N)},\\ |u_n(x)| &\leq \frac{1}{\sqrt{\omega_{n-1}}} \left(\frac{1-|x|^2}{2}\right)^{\frac{N-1}{2}} \frac{1}{|x|^{\frac{N}{2}}} \|u_n\|_{H^1_r(\mathbb{B}^N)}. \end{aligned}$$

Since  $\{|x| \leq \frac{1}{2}\}$ ,  $\ln \frac{1+|x|}{1-|x|} \leq \frac{2r}{1-r^2}$  and 2 , we have

$$d(x)^{mp}|u|^p \leq C\left(\ln\frac{1+|x|}{1-|x|}\right)^{mp}\left(\frac{1-|x|^2}{2}\right)^{p\frac{N-2}{2}}\left(\frac{1}{|x|^{\frac{N-2}{2}}}\right)^p \leq C\left(\frac{2|x|}{1-|x|^2}\right)^{mp}\left(\frac{1-|x|^2}{2}\right)^{p\frac{N-2}{2}}\left(\frac{1}{|x|^{\frac{N-2}{2}}}\right)^p \equiv h_1.$$

Set

$$g_1(x) = \begin{cases} h_1(x) & \text{if } 0 \le |x| < \frac{1}{2} \\ 0 & \text{if } \frac{1}{2} \le |x| < 1 \end{cases}$$

then

$$\begin{split} \int_{\mathbb{B}^N} g_1 \, dV_{\mathbb{B}^N} &= \int_0^{\frac{1}{2}} \left(\frac{2r}{1-r^2}\right)^{mp} \left(\frac{1-r^2}{2}\right)^{p\frac{N-2}{2}} \left(\frac{1}{r^{\frac{N-2}{2}}}\right)^p r^{N-1} \left(\frac{2}{1-r^2}\right)^N dr \\ &\leq C \int_0^{\frac{1}{2}} \left(\frac{2r}{1-r^2}\right)^{mp} \left(\frac{1-r^2}{2}\right)^{\left(p\frac{N-2}{2}-N\right)} r^{N-1-p\frac{N-2}{2}} dr \\ &\leq C \int_0^{\frac{1}{2}} r^{mp+N-1-p\frac{N-2}{2}} dr < \infty. \end{split}$$

Since  $\{|x| > \frac{1}{2}\}$  and 2 , we have

$$d(x)^{mp}|u|^p \leq C \left(\ln\frac{1+|x|}{1-|x|}\right)^{mp} \left(\frac{1-|x|^2}{2}\right)^{p\frac{N-1}{2}} \left(\frac{1}{|x|^{\frac{N}{2}}}\right)^p \leq C \left(\ln\frac{1+|x|}{1-|x|}\right)^{mp} \left(\frac{1-|x|^2}{2}\right)^{p\frac{N-1}{2}} \left(\frac{1}{|x|^{\frac{N}{2}}}\right)^p \equiv h_2.$$

Set

$$g_2(x) = \begin{cases} 0 & \text{if } 0 \le |x| < \frac{1}{2} \\ h_2(x) & \text{if } \frac{1}{2} \le |x| < 1, \end{cases}$$

then

$$\begin{split} \int_{\mathbb{B}^N} g_2 \, dV_{\mathbb{B}^N} &= \int_{\frac{1}{2}}^1 \left( \ln \frac{1+r}{1-r} \right)^{mp} \left( \frac{1-r^2}{2} \right)^{p\frac{N-1}{2}} \left( \frac{1}{r^{\frac{N}{2}}} \right)^p r^{N-1} \left( \frac{2}{1-r^2} \right)^N dr \\ &\leq C \int_{\frac{1}{2}}^1 \left( \ln \frac{1+r}{1-r} \right)^{mp} \left( \frac{1-r^2}{2} \right)^{(p\frac{N-1}{2}-N)} r^{N-1-\frac{N}{2}p} dr \\ &\leq \int_{\ln 3}^\infty s^{mp} \left( \frac{2e^s}{(e^s+1)^2} \right)^{\frac{N-1}{2}p-N+1} ds < \infty. \end{split}$$

Hence, we have

$$|d(x)^{mq}u_n(x)^q| \le g_1(x) + g_2(x).$$

By the Dominated convergence theorem, we obtain

$$\lim_{n \to \infty} \int_{\mathbb{B}^N} d(x)^{mq} u_n^p(x) dV_{\mathbb{B}^N} = \int_{\mathbb{B}^N} d(x)^{mp} u^p(x) dV_{\mathbb{B}^N}.$$

In the same way we conclude that

$$\lim_{n \to \infty} \int_{\mathbb{B}^N} d(x)^{mq} v_n^p(x) dV_{\mathbb{B}^N} = \int_{\mathbb{B}^N} d(x)^{mp} v^p(x) dV_{\mathbb{B}^N}$$

and the Lemma holds.

3 Proof of Theorem 1.1

The Clark's theorem is one of the most important results in critical point theory (see [5]). It was successfully applied to sublinear elliptic problems with symmetry and the existence of infinitely many solutions around the 0 was shown.

In order to state the Clark's theorem, we need some terminologies.

Let  $(X, \|\cdot\|_X)$  be a Banach Space and  $\mathcal{I} \in C^1(X, \mathbb{R})$ .

- (i) For  $c \in \mathbb{R}$  we say that  $\mathcal{I}(u)$  satisfies  $(PS)_c$  condition if any sequence  $(u_j)_{j=1}^{\infty} \subset X$  such that  $\mathcal{I}(u_j) \to c$  and  $\|\mathcal{I}'(u_j)\| \to 0$  has a convergent subsequence.
- (ii) Let S be a symmetric and closed set family in  $X \setminus \{0\}$ . For  $A \in S$ , the genus  $\gamma(A) = \min\{n \in \mathbb{N} : \phi \in C(A, \mathbb{R}^n\{0\}) \text{ is odd}\}$ . If there is no such natural, we set  $\gamma(A) = \infty$ .

(iii) Let  $\Omega$  be a open and bounded set,  $0 \in \Omega$  in  $\mathbb{R}^n$ . If  $A \in S$  is such that there exists a odd homeomorphism function from A to  $\partial\Omega$ , then  $\gamma(A) = n$ .

**Theorem 3.1** (Clark's Theorem). Let  $\mathcal{I} \in C(X, \mathbb{R})$  be an even function, bounded from below, with  $\mathcal{I}(0) = 0$  and there exists a compact, symmetric set  $K \in S$  such that  $\gamma(K) = k$  and  $\sup_K \mathcal{I} < 0$ . Then I has least k distinct pairs of critical points.

The proof of Theorem 1.1 is made by using Theorem 3.1.

The  $(\mathcal{H})$  system are the Euler-Lagrange equations related to the functional

$$I(u,v) = \frac{1}{2} \int_{\mathbb{B}^N} (\|\nabla_{\mathbb{B}^N} u\|_{\mathbb{B}^N}^2 + \|\nabla_{\mathbb{B}^N} v\|_{\mathbb{B}^N}^2) dV_{\mathbb{B}^N} - \int_{\mathbb{B}^N} K(d(x))Q(u,v)dV_{\mathbb{B}^N}$$
(2)

which is  $C^1$  on H.

The functional I is not bounded from below, therefore, we can't apply the Clark's technique for this functional.

In order to overcome this difficulty we consider the auxiliary functional

$$J(u,v) = \left(\int_{\mathbb{B}^N} \left(\|\nabla_{\mathbb{B}^N} u\|_{\mathbb{B}^N}^2 + \|\nabla_{\mathbb{B}^N} v\|_{\mathbb{B}^N}^2\right) dV_{\mathbb{B}^N}\right)^{p-1} - \int_{\mathbb{B}^N} K(d(x))Q(u,v) dV_{\mathbb{B}^N}, \quad (3)$$

where  $p \in (2, \delta)$ , while for J' we have,  $\forall (\phi, \psi) \in H$ 

$$J'(u,v)(\phi,\psi) = (2p-2) ||(u,v)||^{2p-4} \int_{\mathbb{B}^N} (\langle \nabla_{\mathbb{B}^N} u, \nabla_{\mathbb{B}^N} \phi \rangle_{\mathbb{B}^N} + \langle \nabla_{\mathbb{B}^N} v, \nabla_{\mathbb{B}^N} \psi \rangle_{\mathbb{B}^N}) dV_{\mathbb{B}^N} - \int_{\mathbb{B}^N} K(d(x))(\phi Q_u(u,v) + \psi Q_v(u,v)) dV_{\mathbb{B}^N}$$

$$(4)$$

We will show that the set of critical points of J is related to a set of critical points of I and J satisfies the conditions of Theorem 3.1.

The proof of Theorem 1.1 is divided into several lemmas.

**Lemma 3.1.** If  $(u, v) \in H$ ,  $(u, v) \neq (0, 0)$  is a critical point for J, then

$$(w,z) = \left(\frac{u}{\left[(2p-2) \|(u,v)\|^{2p-4}\right]^{\frac{1}{p-2}}}, \frac{v}{\left[(2p-2) \|(u,v)\|^{2p-4}\right]^{\frac{1}{p-2}}}\right)$$

is a critical point for I.

PROOF. Note that  $(u, v) \neq (0, 0)$  is a critical point for J if, and only if, (u, v) is a weak solution to the problem

$$\begin{cases} -(2p-2) \|(u,v)\|^{2p-4} \Delta_{\mathbb{B}^N} u = K(d(x))Q_u(u,v) \\ -(2p-2) \|(u,v)\|^{2p-4} \Delta_{\mathbb{B}^N} v = K(d(x))Q_v(u,v) \\ u,v \in H^1_r(\mathbb{B}^N), N \ge 3 \end{cases}$$
(S)

Define  $\lambda(\|(u,v)\|) = \left[(2p-2)\|(u,v)\|^{2p-4}\right]^{\frac{-1}{p-2}}$ , then  $(w,z) = \lambda(\|(u,v)\|)(u,v)$ . Using the p-1-homogeneity condition of  $Q_u(u,v)$  and  $Q_u(u,v)$ , observe that

$$\begin{aligned} &-\Delta_{\mathbb{B}^N} w - K(d(x))Q_u(w,z) \\ &= -\lambda(\|(u,v)\|)\Delta_{\mathbb{B}^N} u - (\lambda(\|(u,v)\|))^{p-1}K(d(x))Q_u(u,v) \\ &= -\lambda(\|(u,v)\|)K(d(x))Q_u(u,v)\left((\lambda(\|(u,v)\|))^{p-2} - \frac{1}{(2p-2)\|(u,v)\|^{2p-4}}\right) \end{aligned}$$

and

$$\begin{aligned} &-\Delta_{\mathbb{B}^N} z - K(d(x))Q_v(w,z) \\ &= -\lambda(\|(u,v)\|)\Delta_{\mathbb{B}^N} v - (\lambda(\|(u,v)\|))^{p-1}K(d(x))Q_v(u,v) \\ &= -\lambda(\|(u,v)\|)K(d(x))Q_v(u,v)\left((\lambda(\|(u,v)\|))^{p-2} - \frac{1}{(2p-2)\|(u,v)\|^{2p-4}}\right) \end{aligned}$$

Hence (w, z) is a weak solution for problem  $(\mathcal{H})$  and so, a critical point for I. Lemma 3.2. J(u, v) is bounded from below and satisfies the  $(PS)_c$  condition. PROOF. From  $(K_1)$ - $(K_2)$ ,  $(Q_2)$ - $(Q_3)$  and Remark 2.1

$$J(u,v) = \left( \int_{\mathbb{B}^{N}} \|\nabla_{\mathbb{B}^{N}} u\|_{\mathbb{B}^{N}}^{2} + \|\nabla_{\mathbb{B}^{N}} v\|_{\mathbb{B}^{N}}^{2} dV_{\mathbb{B}^{N}} \right)^{p-1} - \int_{\mathbb{B}^{N}} K(d(x))Q(u,v) dV_{\mathbb{B}^{N}}$$
  

$$\geq \|(u,v)\|^{2p-2} - C \int_{\mathbb{B}^{N}} d(x)^{\beta} (|u|^{p} + |v|^{p}) dV_{\mathbb{B}^{N}}$$
  

$$\geq \|(u,v)\|^{2p-2} - \|(u,v)\|^{p},$$

so that J(u, v) is bounded from below.

Let  $(u_n.v_n) \in H$  be such that  $|J(u_n,v_n)| \leq C$  with  $C \in \mathbb{R}^+$ ,  $J'(u_n,v_n) \to 0$ . Since

$$C \ge J(u_n, v_n) \ge ||(u_n, v_n)||^{2p-2} - ||(u_n, v_n)||^p$$

we conclude that  $||(u_n, v_n)||$  is bounded. So, there exists  $(u, v) \in H$  such that, passing to a subsequence if necessary,

$$(u_n, v_n) \rightharpoonup (u, v), \text{ as } n \to \infty.$$

From the Embedding Lemma 2.2,

$$\int_{\mathbb{B}^N} (d(x))^{\beta} (|u_n|^p + |v_n|^p) \, dV_{\mathbb{B}^N} \longrightarrow \int_{\mathbb{B}^N} (d(x))^{\beta} (|u|^p + |v|^p) \, dV_{\mathbb{B}^N}.$$

and by  $(K_2) - (Q_2)$ , we infer that

$$|K(d(x)(u_nQ_u(u_n, v_n) + v_nQ_v(u_n, v_n))| \le C(d(x))^{\beta}(|u_n|^p + v_n|^p).$$

Therefore, by the Lebesgue Dominated Convergence Theorem,

$$\begin{split} \int_{\mathbb{B}^N} K(d(x))(Q_u(u_n,v_n)u_n + Q_v(u_n,v_n)v_n) \, dV_{\mathbb{B}^N} \to \int_{\mathbb{B}^N} K(d(x))(Q_u(u,v)u + Q_v(u,v)v) \, dV_{\mathbb{B}^N}.\\ \text{Since } J'(u,v)(u,v) &= 0 \text{ and } J'(u_n,v_n)(u_n,v_n) = o_n(1) \text{ as } n \to \infty, \text{ we have} \end{split}$$

$$(2p-2)\Big(\|(u_n,v_n)\|^{2p-2} - \|(u,v)\|^{2p-2}\Big) = J'(u_n,v_n)(u_n,v_n) - J'(u,v)(u,v) + \\ + \int_{\mathbb{B}^N} K(d(x))\Big(Q_u(u_n,v_n)u_n + Q_v(u_n,v_n)v_n\Big) dV_{\mathbb{B}^N} + \\ - \int_{\mathbb{B}^N} K(d(x))\Big(Q_u(u,v)u + Q_v(u,v)v\Big) dV_{\mathbb{B}^N} = o_n(1)$$

then,  $||(u_n, v_n)|| \to ||(u, v)||$ . Therefore,

$$(u_n, v_n) \to (u, v), \text{ as } n \to \infty, \text{ in } H$$

The next lemma ends the proof of the Theorem 1.1.

**Lemma 3.3.** Given  $k \in \mathbb{N}$ , there exists a compact and symmetric set  $K \in H$  such that  $\gamma(K) = k$  and  $\sup_K J < 0$ .

**PROOF.** Let  $X_k \subset H$  be a subspace of dimension k. Consider the following norm in  $X_k$ 

$$\|(u,v)\|_{X_k} = \left(\int_{\mathbb{B}^N} K(d(x))(|u|^p + |v|^p) dV_{\mathbb{B}^N}\right)^{\frac{1}{p}}.$$

Since  $X_k \subset H$  has finite dimension, there exists a > 0 such that

$$a ||(u,v)||_{X_k} \le ||(u,v)|| \le \frac{1}{a} ||(u,v)||_{X_k}, \quad \forall (u,v) \in X_k.$$

Therefore, we obtain from  $(Q_3)$  that

$$J(u,v) \le \|(u,v)\|^{2p-2} - C \int_{\mathbb{B}^N} K(d(x))(|u|^p + |v|^p) dV_{\mathbb{B}^N} = \|(u,v)\|^{2p-2} - C \|(u,v)\|_{X_k}^p,$$

where  $C \in \mathbb{R}$  is a positive constant. We then conclude that

$$J(u,v) \le \|(u,v)\|_{X_k}^p \left(\frac{\|(u,v)\|_{X_k}^{p-2}}{a^{2p-2}} - C\right).$$

Let  $A = a^{\frac{2p-2}{p-2}}$  and consider the set  $K = \{(u, v) \in X_k : ||(u, v)||_{X_k} = \frac{A}{2}C^{\frac{1}{p-2}}\}$ , then

$$J(u,v) \le C \frac{A^p}{2^p} \left(\frac{1}{2^{q-2}} - 1\right) < 0, \quad \forall (u,v) \in K.$$

We get that  $\sup_K J < 0$ , where  $K \subset H$  is a compact and symmetric set such that  $\gamma(K) = k$ .

Finally, from Lemmas 3.2 and 3.3, Theorem 3.1 implies the existence of at least k distinct pairs of critical points for the functional J. Since k is arbitrary, we obtain infinitely many critical points for J.

In view of Lemma 3.1, we conclude that the functional J possesses, together with I, infinitely many critical points.

Finally, we point out that since H is a closed subspace of the Hilbert space  $H^1(\mathbb{B}^N) \times H^1(\mathbb{B}^N)$ , following some ideas in [1, 3], we can conclude that (u, v) is a critical point in  $H^1(\mathbb{B}^N) \times H^1(\mathbb{B}^N)$ .

#### 4 Further result

We can apply the same method that proved Theorem 1.1 to establish the existence of infinitely many solutions for the following semilinear elliptic equation

$$\begin{cases} -\Delta_{\mathbb{B}^N}^{\alpha} u = K(d(x)) |u|^{p-2} u, \\ u \in \mathbb{E} \subset H^1_r(\mathbb{B}^N), N \ge 3 \end{cases}$$
  $(\mathcal{H}*)$ 

where K satisfies  $(K_1) - (K_2)$ ,  $-\Delta^{\alpha}_{\mathbb{R}^N}$  is the Laplace-Beltrami type operator

$$-\Delta^{\alpha}_{\mathbb{B}^N} u = -(p(x))^{-N} div(p(x)^{N-2}(d(x))^{\alpha} \nabla u)$$

and

$$\mathbb{E} = \left\{ u \in H_r^1(\mathbb{B}^N) : \|u\|_{\mathbb{E}} = \left( \int_{\mathbb{B}^N} (d(x))^{\alpha} \|\nabla_{\mathbb{B}^N} u\|_{\mathbb{B}^N}^2 dV_{\mathbb{B}^N} \right)^{\frac{1}{2}} \right\}.$$

We obtain the following result

**Theorem 4.1.** Under hypotheses  $(K_1)$ - $(K_2)$ ,  $(\mathcal{H}^*)$  equation has infinitely many solutions.

The energy functional corresponding to  $(\mathcal{H}^*)$  is

$$I(u) = \frac{1}{2} \int_{\mathbb{B}^N} (d(x))^{\alpha} \left\| \nabla_{\mathbb{B}^N} u \right\|_{\mathbb{B}^N}^2 dV_{\mathbb{B}^N} - \frac{1}{q} \int_{\mathbb{B}^N} K(d(x)) |u|^q \, dV_{\mathbb{B}^N}$$
(5)

defined on  $H^1_r(\mathbb{B}^N)$ 

Problem  $(\mathcal{H}^*)$  is close related to one studied by Carrião, Faria and Miyagaki [3]. In [3], they proved that the map  $u \mapsto d(x)^m u$  from  $\mathbb{E}$  to  $L^q(\mathbb{B}^N)$  is compact for  $q \in (2, \tilde{m})$ , where

$$\tilde{m} = \begin{cases} \frac{2N}{N-2-2m+\alpha} & \text{if } m < \frac{N-2+\alpha}{2} \\ \infty & \text{if otherwise} \end{cases}$$

and then there exists a positive constant C > 0, such that

$$\int_{\mathbb{B}^N} d(x)^{\beta} |u(x)|^q dV_{\mathbb{B}^N} \le C \left( \int_{\mathbb{B}^N} (d(x))^{\alpha} \|\nabla_{\mathbb{B}^N} u\|_{\mathbb{B}^N} dV_{\mathbb{B}^N} \right)^{\frac{q}{2}},\tag{6}$$

by taking  $m = \frac{\beta}{q}$  with  $2 < q < \frac{2N}{N-2-2\frac{\beta}{q}+\alpha}$ .

Using  $(K_1)$ - $(K_2)$  together with inequality (6), we get that the functional I is well defined. This functional is not bounded from below, hence, we can't apply Clark's technique [5].

In order to overcome this difficulty we consider the auxiliary functional

$$\psi(u) = \left(\int_{\mathbb{B}^N} (d(x))^{\alpha} \|\nabla_{\mathbb{B}^N} u\|_{\mathbb{B}^N}^2 dV_{\mathbb{B}^N}\right)^{p-1} - \int_{\mathbb{B}^N} K(d(x)) |u|^p dV_{\mathbb{B}^N},\tag{7}$$

where  $p \in (2, 2^{\beta}_{\alpha}), u \in \mathbb{E}$  and

$$\psi'(u)v = (2p-2) \|u\|_{\mathbb{E}}^{2p-4} \int_{\mathbb{B}^N} (d(x))^{\alpha} \langle \nabla_{\mathbb{B}^N} u, \nabla_{\mathbb{B}^N} v \rangle_{\mathbb{B}^N} dV_{\mathbb{B}^N} + \int_{\mathbb{B}^N} K(d(x)) |u|^{p-2} uv \, dV_{\mathbb{B}^N}$$
(8)

We have the corresponding results of Lemmas 3.1, 3.2 and 3.3 for problem ( $\mathcal{H}*$ ). The set of critical points of  $\psi$  is related to a set of critical points of I and  $\psi$  satisfies the conditions of Theorem 3.1.

**Lemma 4.1.** If  $u \in \mathbb{E}$ ,  $u \neq 0$  is a critical point for  $\psi$ , then  $v = \frac{u}{[(2p-2) ||u||_{\mathbb{E}}^{2p-4}]^{\frac{1}{p-2}}}$  is a critical point for I,

**Lemma 4.2.**  $\psi(u)$  is bounded from below and satisfies the Palais-Smale condition (PS).

**Lemma 4.3.** Given  $k \in \mathbb{N}$ , there exists a compact and symmetric set  $K \in \mathbb{E}$  such that  $\gamma(K) = k$  and  $\sup_K \psi < 0$ .

From Lemmas 4.2 and 4.3 and Theorem 3.1, we conclude that the functional I possesses infinitely many critical points.

Acknowledgments. This paper was carried out while the first author was visiting the Mathematics Department of UFOP and she would like to thank the members of DEMAT-UFOP for their hospitality. The authors are greatful to professor Giovany Figueiredo for helpful comments.

## Referências

- G. Bianchi, J. Chabrowski, A. Szulkin, On symmetric solutions of an elliptic equation with nonlinearity involving critical Sobolev exponent, Nonlinear Anal., 25 (1995), 41-59.
- [2] L. Boccardo, D. G. de Figueiredo, Some remarks on a system of quasilinear elliptic equations Nonlinear Diff. Equa. Appl., 9 (2002), 309-323.
- [3] P. C. Carrião, L. F. O. Faria, O. H. Miyagaki, Semilinear elliptic equations of the Hénon-type in hyperbolic space, Comm. Contemp. Math., 18, No. 2 (2016), (13 pages).
- [4] M. Chipot, Handbook of Differential Equations Stationary Partial Differential Equations, Elsevier, 2008.
- [5] D. C. Clark, A variant of the Lusternik-Schnirelman theory, Indiana Univ. Math. J., 22 (1972), 65-74.
- [6] D. G. Costa, An Invitation to Variational Methods In Differential Equations, Birkhäuser Boston, 2007.
- [7] D. G. de Figueiredo, Nonlinear elliptic systems, Anais da Academia Brasileira de Ciências, vol. 72, 2000.
- [8] D. G. de Figueiredo, Y. H. Ding, Strongly indefinite functionals and multiple solutions of elliptic systems, Trans. Amer. Math. Soc., 355 No. 7 (2003), 2973-2989.
- D. G. de Figueiredo, P. L. Felmer, On superquadratic elliptic systems, Trans. Amer. Math. Soc., 343 (1994), 119-123.
- [10] H. He, The existence of solutions for Hénon equation in hyperbolic space, Proc. Japan Acad., 89, Ser. A (2013), 24-28.
- [11] H. He, J. Qiu, Existence of solutions for critical Hénon equations in hyperbolic spaces, Electr. J. Diff. Eq., 13 (2013), 1-11.
- [12] H. He, Infinitely many solutions for Hardy-Hénon type elliptic system in hyperbolic space, Ann. Acad. Sci. Fenn. Math, 40 (2015), 969-983.

- [13] D. C. de Morais Filho, M. A. S. Souto, Systems of p-Laplacian equations involving homogeneous nonlinearities with critical sobolev exponent degrees, Commun. in Partial Diff. Equations, 24 (1999), 1537-1553.
- [14] T. Bartsch, D. G. de Figueiredo Infinitely Many Solutions of Nonlinear Elliptic Systems, Progress in Nonlinar Differential Equations and Their Applicantions, (1999), Vol. 35.
- [15] J. Wang, J. Xu, F. Zhang, Existence and multiplicity of solutions for asymptotically Hamiltonian elliptic systems in R<sup>N</sup>, J. Math. Anal. Appl., 367 (2010), 193-203.