

Coordinates Adapted to Vector Fields III: Real Analyticity

Brian Street*

Abstract

Given a finite collection of C^1 vector fields on a C^2 manifold which span the tangent space at every point, we consider the question of when there is locally a coordinate system in which these vector fields are real analytic. We give necessary and sufficient, coordinate-free conditions for the existence of such a coordinate system. Moreover, we present a quantitative study of these coordinate charts. This is the third part in a three-part series of papers. The first part, joint with Stovall, lay the groundwork for the coordinate system we use in this paper and showed how such coordinate charts can be viewed as scaling maps for sub-Riemannian geometry. The second part dealt with the analogous questions with real analytic replaced by C^∞ and Zygmund spaces.

1 Introduction

Let X_1, \dots, X_q be C^1 vector fields on a C^2 manifold M , which span the tangent space at every point of M . In this paper, we investigate the following three closely related questions:

- (i) When is there a coordinate system near a fixed point $x_0 \in M$ such that the vector fields X_1, \dots, X_q are real analytic in this coordinate system?
- (ii) When is there a real analytic manifold structure on M , compatible with its C^2 structure, such that X_1, \dots, X_q are real analytic with respect to this structure? When such a structure exists, we will see it is unique.
- (iii) When there is a coordinate system as in (i), how can we pick it so that X_1, \dots, X_q are “normalized” in this coordinate system in a quantitative way which is useful for applying techniques from analysis?

We present necessary and sufficient conditions for (i) and (ii), and under these conditions give a quantitative answer to (iii). This is the third part in a three part series of papers. In the first two parts [SS18, Str21], the same questions were investigated where “real analytic” was replaced by Zygmund spaces.

The first paper in the series [SS18], joint with Stovall, was based on methods from ODEs, while the second paper [Str21] sharpened the results from the first paper using methods from PDEs. In this paper, we take the results from the first paper as a starting point, and use additional methods from ODEs to answer the above questions. Thus, this paper does not use any methods from PDEs.

The coordinate charts from (iii) can be viewed as scaling maps in sub-Riemannian geometry. When viewed in this light, these results can be seen as a continuation of results initiated by Nagel, Stein, and Wainger [NSW85] and C. Fefferman and Sánchez-Calle [FSC86], and furthered by Tao and Wright [TW03] and the author [Str11]. See Sections 1.1 and 5 for a description of this.

This paper is a continuation of the first part of the series [SS18]. That paper contains several applications and motivations for the types of results described in this paper. It also contains a more leisurely introduction to some of the definitions and results in this paper, though we include all the necessary definitions so that the statement of the results is self-contained.

*This material is partially based upon work supported by the National Science Foundation under Grant No. 1440140, while the author was in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the spring semester of 2017. The author was also partially supported by National Science Foundation Grant Nos. 1401671 and 1764265.

1.1 The role of real analyticity

Two important ways the main results of this paper can be used are:

- They give necessary and sufficient, coordinate free, conditions on a collection of C^1 vector fields, which span the tangent space, for there to exist a coordinate system in which these vector fields are real analytic.
- They give scaling maps adapted to real analytic sub-Riemannian geometries, which are useful for questions from harmonic analysis.

The first way seems to be new. The second way has a long history and similar results have been used in several areas of harmonic analysis. We now turn to describing some of this; see also Section 5.1.2.

Real analytic vector fields have important applications in several types of questions from harmonic analysis. Since the original work of Hörmander [Hör67], C^∞ vector fields satisfying Hörmander’s condition¹ have played a central role in several areas. Nagel, Stein, and Wainger [NSW85] developed a quantitative theory of the sub-Riemannian geometries induced by Hörmander vector fields. In particular, they introduced scaling maps adapted to Hörmander vector fields which allowed the use of many techniques from harmonic analysis to be generalized to the setting of sub-Riemannian manifolds. These ideas have been used in many different ways, including applications to partial differential equations defined by vector fields and singular Radon transforms. See the notes at the end of Chapter 2 of [Str14] for a history of some of these ideas.

A finite collection of real analytic vector fields does not necessarily satisfy Hörmander’s condition; however, it does satisfy a generalization of this condition: the C^∞ module generated by the vector fields and their commutators of all orders is locally finitely generated (as a C^∞ module). This was first noted by Lobry [Lob70] and is a simple consequence of the Weierstrass preparation theorem (see Section 9). Because of this, it is possible to generalize the quantitative theory of Nagel, Stein, and Wainger to a setting which applies to any finite collection of real analytic vector fields, whether or not they satisfy Hörmander’s condition. The techniques required for this generalization use ideas of Tao and Wright [TW03] and the author [Str11]. In the context of the quantitative theory of sub-Riemannian geometry applied to questions in analysis, this seems to have been first explicitly used by the author and Stein [SS12] to study singular Radon transforms.

Thus, real analytic vector fields hold a special place: the quantitative scaling techniques used to study Hörmander vector fields can often be applied to real analytic vector fields, whether or not they satisfy Hörmander’s condition. For many such applications, the scaling maps developed in [Str11] are sufficient; however, in the context of real analytic vector fields, the theory from that paper has several deficiencies which are fixed in this paper. One major deficiency is that if one starts with real analytic vector fields, the scaling theorems from [Str11] only guarantee quantitative bounds on the C^m norms of the rescaled vector fields and no estimates on their real analyticity. Thus, when one applies the results from [Str11] to a real analytic setting, the real analyticity is destroyed. In this paper, we show the rescaled vector fields are real analytic and give appropriate quantitative control of this fact. This is described in Section 5.1.2.

Remark 1.1. The results from this paper are useful even when considering some very classical settings. Indeed, suppose V_1, \dots, V_r are real analytic vector fields on an open set $\Omega \subseteq \mathbb{R}^n$. The classical Frobenius theorem applies to foliate Ω into leaves; each leaf is a real analytic, injectively immersed sub-manifold (see [Her63, Nag66, Lob70, Sus73]). This may be a singular foliation: the various leaves may have different dimension. Near a point where the dimension changes (a “singular point”), classical proofs do not give good quantitative control on the real analytic coordinate systems which define the leaves: classical proofs “blow up” near a singular point. Our methods give useful, uniform quantitative control near such a singular point and avoid this blow up, in a certain sense. See Remark 5.6.

Outline of the paper: In Section 2, we present the function spaces we use. There are two types of function spaces: the standard ones on Euclidean space described in Section 2.1, and analogs on a C^2 manifold endowed with a finite collection of C^1 vector fields described in Section 2.2. In Section 3 we describe the main results

¹A finite collection of vector fields satisfies “Hörmander’s condition” if the Lie algebra generated by the vector fields spans the tangent space at every point.

of this paper, starting with the qualitative results in Section 3.1 (i.e., (i) and (ii) from the beginning of the introduction), and then turning to the quantitative results in Section 3.2 (i.e., (iii)). In Section 4 we further study the function spaces introduced in Section 2. In Section 5 we describe how the quantitative results can be seen as scaling maps in sub-Riemannian type geometries. In Section 6 we describe the results we use from [SS18]. In Section 7 we prove the main results of this paper. In Section 8 we describe the special case of some of our results when working on Euclidean space with Lebesgue measure (which is the most common application). Finally, in Section 9 we prove the results concerning scaling from Section 5.

2 Function Spaces

Before we can state our main results, we need to introduce the function spaces we use. As described in [SS18], we make a distinction between function spaces on subsets of \mathbb{R}^n and function spaces on a C^2 manifold M . On \mathbb{R}^n we use the standard coordinate system to define the usual function spaces. On an abstract C^2 manifold M , we do not have access to any one natural coordinate system and so it does not make sense to discuss, for example, real analytic functions on M . However, if M is endowed with C^1 vector fields X_1, \dots, X_q , we are able to define what it means to be real analytic with respect to these vector fields, and that is how we shall proceed. The notion of a function being real analytic with respect to a finite collection of vector fields is a special case of a general notion due to Nelson [Nel59]. Throughout the paper, $B^n(\delta)$ denotes the ball of radius $\delta > 0$, centered at 0, in \mathbb{R}^n .

2.1 Function Spaces on Euclidean Space

Let $\Omega \subseteq \mathbb{R}^n$ be an open set. We have the usual Banach space of bounded, continuous functions on Ω :

$$C(\Omega) := \{f : \Omega \rightarrow \mathbb{C} \mid f \text{ is continuous and bounded}\}, \quad \|f\|_{C(\Omega)} := \sup_{x \in \Omega} |f(x)|.$$

We next define two closely related spaces of real analytic functions on \mathbb{R}^n . For $r > 0$ let $B^n(r)$ be the ball of radius r in \mathbb{R}^n , centered at 0. We define $\mathcal{A}^{n,r}$ to be the space of those $f \in C(B^n(r))$ such that $f(t) = \sum_{\alpha \in \mathbb{N}^n} \frac{c_\alpha}{\alpha!} t^\alpha$, $\forall t \in B^n(r)$, where

$$\|f\|_{\mathcal{A}^{n,r}} := \sum_{\alpha \in \mathbb{N}^n} \frac{|c_\alpha| r^{|\alpha|}}{\alpha!} < \infty.$$

For $\Omega \subseteq \mathbb{R}^n$ open, we let $f \in C^{\omega,r}(\Omega)$ consist of those $f \in C^\infty(\Omega)$ such that

$$\|f\|_{C^{\omega,r}(\Omega)} := \sum_{\alpha \in \mathbb{N}^n} \frac{\|\partial_x^\alpha f\|_{C(\Omega)} r^{|\alpha|}}{\alpha!} < \infty. \quad (2.1)$$

For the relationship between $\mathcal{A}^{n,r}$ and $C^{\omega,r}(\Omega)$ see Lemma 4.2. We set

$$C^\omega(\Omega) := \bigcup_{r>0} C^{\omega,r}(\Omega).$$

We say $f \in C_{\text{loc}}^\omega(\Omega)$ if for all $x \in \Omega$, there exists an open neighborhood $U \subseteq \Omega$ of x , with $f|_U \in C^\omega(U)$. It is easy to see that $C_{\text{loc}}^\omega(\Omega)$ is the usual space of real analytic functions on Ω .

If \mathcal{X} is a Banach space, we define the same spaces taking values in \mathcal{X} in the obvious way, and denote these spaces by $C(\Omega; \mathcal{X})$, $\mathcal{A}^{n,r}(\mathcal{X})$, $C^{\omega,r}(\Omega; \mathcal{X})$, and $C^\omega(\Omega; \mathcal{X})$. When we have a vector field X on Ω , we identify $X = \sum_{j=1}^n a_j(x) \frac{\partial}{\partial x_j}$ with the function $(a_1, \dots, a_n) : \Omega \rightarrow \mathbb{R}^n$. It therefore makes sense to consider quantities like $\|X\|_{C^{\omega,r}(\Omega; \mathbb{R}^n)}$.

2.2 Function Spaces on Manifolds

Let X_1, \dots, X_q be C^1 vector fields on a connected C^2 manifold M . Define the Carnot-Carathéodory ball associated to X_1, \dots, X_q , centered at $x \in M$, of radius $\delta > 0$ by

$$B_X(x, \delta) := \left\{ y \in M \mid \exists \gamma : [0, 1] \rightarrow M, \gamma(0) = x, \gamma(1) = y, \gamma'(t) = \sum_{j=1}^q a_j(t) \delta X_j(\gamma(t)), \right. \\ \left. a_j \in L^\infty([0, 1]), \left\| \sum_{j=1}^q |a_j|^2 \right\|_{L^\infty} < 1 \right\}, \quad (2.2)$$

and for $y \in M$ set

$$\rho(x, y) = \inf\{\delta > 0 : y \in B_X(x, \delta)\}. \quad (2.3)$$

ρ is called a sub-Riemannian distance. See Remark 2.3 for the precise definition of $\gamma'(t)$ used in (2.2).

We use ordered multi-index notation X^α . Here α denotes a list of elements of $\{1, \dots, q\}$ and $|\alpha|$ denotes the length of the list. For example, $X^{(2,1,3,1)} = X_2 X_1 X_3 X_1$ and $|(2, 1, 3, 1)| = 4$.

We have the usual Banach space of bounded continuous functions on M :

$$C(M) := \{f : M \rightarrow \mathbb{C} \mid f \text{ is continuous and bounded}\}, \quad \|f\|_{C(M)} := \sup_{x \in M} |f(x)|.$$

Next, we introduce what it means to be real analytic with respect to X_1, \dots, X_q . Following the setting in \mathbb{R}^n , we introduce two versions of this. Given $x_0 \in M$ and $r > 0$ we define $\mathcal{A}_X^{x_0, r}$ to be the space of those $f \in C(M)$ such that

$$h(t_1, \dots, t_q) := g(e^{t_1 X_1 + \dots + t_q X_q} x_0) \in \mathcal{A}^{q, r},$$

here we are assuming $e^{t_1 X_1 + \dots + t_q X_q} x_0$ exists for $(t_1, \dots, t_q) \in B^q(r)$ (see Definition 3.5). We define $\|f\|_{\mathcal{A}_X^{x_0, r}} := \|h\|_{\mathcal{A}^{q, r}}$. Note that $\|f\|_{\mathcal{A}_X^{x_0, r}}$ depends only on the values of $f(y)$ where $y = e^{t_1 X_1 + \dots + t_q X_q} x_0$ and $(t_1, \dots, t_q) \in B^q(r)$; thus this is merely a semi-norm.

For $r > 0$ we define $C_X^{\omega, r}(M)$ to be the space of those $f \in C(M)$ such that $X^\alpha f$ exists and is continuous for all ordered multi-indices α and such that

$$\|f\|_{C_X^{\omega, r}(M)} := \sum_{m=0}^{\infty} \frac{r^m}{m!} \sum_{|\alpha|=m} \|X^\alpha f\|_{C(M)} < \infty. \quad (2.4)$$

We set

$$C_X^\omega(M) := \bigcup_{r>0} C_X^{\omega, r}(M).$$

The norm $\|f\|_{C_X^{\omega, r}(M)}$ was originally introduced by Nelson [Nel59] in greater generality.

Remark 2.1. When we write Vf for a C^1 vector field V and $f : M \rightarrow \mathbb{R}$, we define this as $Vf(x) := \frac{d}{dt} \Big|_{t=0} f(e^{tX} x)$. When we say Vf exists, it means that this derivative exists in the classical sense, $\forall x$. If we have several C^1 vector fields V_1, \dots, V_L , we define $V_1 V_2 \dots V_L f := V_1(V_2(\dots V_L(f)))$ and to say that this exists means that at each stage the derivatives exist.

Note that if ∇ denotes the list of vector fields $\nabla = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \right)$ and $\Omega \subseteq \mathbb{R}^n$ is open, we have

$$\mathcal{A}_\nabla^{0, r} = \mathcal{A}^{n, r} \text{ and } C_\nabla^{\omega, r}(\Omega) = C^{\omega, r}(\Omega),$$

with equality of norms.² For more details on these spaces, see Section 4.

Throughout the paper if we claim $\|f\|_{C_X^{\omega, r}(M)} < \infty$ it means $f \in C_X^{\omega, r}(M)$, and similarly for any other function space. We refer the reader to [SS18] for a more detailed discussion of the above definitions.

An important property of the above spaces and norms is that they are invariant under diffeomorphisms.

²Notice that (2.1) uses regular multi-indices, while (2.4) uses ordered multi-indices. Once this is taken into account, proving the two norms are equal when $X = \nabla$ is straightforward.

Proposition 2.2. *Let N be another C^2 manifold, let $\Phi : M \rightarrow N$ be a C^2 diffeomorphism, and let Φ_*X denote the list of vector fields $\Phi_*X_1, \dots, \Phi_*X_q$. Then the map $f \mapsto f \circ \Phi$ is an isometric isomorphism between the following spaces: $C(N) \rightarrow C(M)$, $C_{\Phi_*X}^{\omega,r}(N) \rightarrow C_X^{\omega,r}(M)$, and $\mathcal{A}_{\Phi_*X}^{\Phi(x_0),r} \rightarrow \mathcal{A}_X^{x_0,r}$.*

Proof. This is immediate from the definitions. \square

Remark 2.3. In (2.2) (and in the rest of the paper), $\gamma'(t)$ is defined as follows. In the case that M is an open subset $\Omega \subseteq \mathbb{R}^n$ and $\gamma : [a, b] \rightarrow \Omega$, $\gamma'(t) = \sum_{j=1}^q a_j(t)X_j(\gamma(t))$ is defined to mean $\gamma(t) = \gamma(a) + \int_a^t \sum_j a_j(s)X_j(\gamma(s)) ds$; note that this definition is local in t (equivalently, we are requiring that γ be absolutely continuous and have the desired derivative almost everywhere). For an abstract C^2 manifold, this is interpreted locally. I.e., if $\gamma : [a, b] \rightarrow M$, we say $\gamma'(t) = \sum_{j=1}^q a_j(t)X_j(\gamma(t))$ if $\forall t_0 \in [a, b]$, there is an open neighborhood N of $\gamma(t_0)$ and a C^2 diffeomorphism $\Psi : N \rightarrow \Omega$, where $\Omega \subseteq \mathbb{R}^n$ is open, such that $(\Psi \circ \gamma)'(t) = \sum_{j=1}^q a_j(t)(\Psi_*X_j)(\Psi \circ \gamma(t))$ for t near t_0 ($t \in [a, b]$).

3 Results

We present the main results of the paper. We separate the results into the qualitative results (i.e., (i) and (ii) from the introduction) and quantitative results (i.e., (iii)). The qualitative results are a simple consequence of the quantitative results, and the quantitative results are useful for proving results in analysis (see Section 5).

3.1 Qualitative Results

Let X_1, \dots, X_q be C^1 vector fields on a C^2 manifold \mathfrak{M} . For $x, y \in \mathfrak{M}$, let $\rho(x, y)$ denote the sub-Riemannian distance associated to X_1, \dots, X_q on \mathfrak{M} defined by (2.3). Fix $x_0 \in M$ and let $Z := \{y \in \mathfrak{M} : \rho(x_0, y) < \infty\}$. ρ is a metric on Z , and we give Z the topology induced by ρ (this is finer³ than the topology as a subspace of \mathfrak{M} and may be strictly finer). Let $M \subseteq Z$ be a connected open subset of Z containing x_0 . We give M the topology of a subspace of Z . We begin with a classical result to set the stage.

Proposition 3.1. *Suppose $[X_i, X_j] = \sum_{k=1}^q c_{i,j}^k X_k$, where $c_{i,j}^k : M \rightarrow \mathbb{R}$ are locally bounded. Then, there is a C^2 manifold structure on M (compatible with its topology) such that:*

- *The inclusion $M \hookrightarrow \mathfrak{M}$ is a C^2 injective immersion.*
- *X_1, \dots, X_q are C^1 vector fields tangent to M .*
- *X_1, \dots, X_q span the tangent space at every point of M .*

Furthermore, this C^2 structure is unique in the sense that if M is given another C^2 structure (compatible with its topology) such that the inclusion map $M \hookrightarrow \mathfrak{M}$ is a C^2 injective immersion, then the identity map $M \rightarrow M$ is a C^2 diffeomorphism between these two structures.

For a proof of Proposition 3.1 (which is standard), see [SS18, Appendix A]. Henceforth, we assume the conditions of Proposition 3.1 so that M is a C^2 manifold and X_1, \dots, X_q are C^1 vector fields on M which span the tangent space at every point. We write $n := \dim \text{span}\{X_1(x_0), \dots, X_q(x_0)\}$ so that $\dim M = n$.

Remark 3.2. If $X_1(x_0), \dots, X_q(x_0)$ span $T_{x_0}\mathfrak{M}$, then M is an open submanifold of \mathfrak{M} . If X_1, \dots, X_q span the tangent space at every point of \mathfrak{M} and \mathfrak{M} is connected, one may take $M = \mathfrak{M}$.

Theorem 3.3 (The Local Theorem). *The following three conditions are equivalent:*

- (i) *There exists an open neighborhood $V \subseteq M$ of x_0 and a C^2 diffeomorphism $\Phi : U \rightarrow V$ where $U \subseteq \mathbb{R}^n$ is open, such that $\Phi_*X_1, \dots, \Phi_*X_q \in C^\omega(U; \mathbb{R}^n)$.*

³See [SS18, Lemma A.1] for a proof that this topology is finer than the subspace topology.

(ii) Reorder the vector fields so that $X_1(x_0), \dots, X_n(x_0)$ are linearly independent. There exists an open neighborhood $V \subseteq M$ of x_0 such that:

- $[X_i, X_j] = \sum_{k=1}^n \hat{c}_{i,j}^k X_k$, $1 \leq i, j \leq n$, where $\hat{c}_{i,j}^k \in C_X^\omega(V)$.
- For $n+1 \leq j \leq q$, $X_j = \sum_{k=1}^n b_j^k X_k$, where $b_j^k \in C_X^\omega(V)$.

(iii) There exists an open neighborhood $V \subseteq M$ of x_0 such that $[X_i, X_j] = \sum_{k=1}^q c_{i,j}^k X_k$, $1 \leq i, j \leq q$, where $c_{i,j}^k \in C_X^\omega(V)$.

Theorem 3.4 (The Global Theorem). *The following two conditions are equivalent:*

(i) *There is a real analytic atlas on M , compatible with its C^2 structure, such that X_1, \dots, X_q are real analytic with respect to this atlas.*

(ii) *For each $x_0 \in M$, any of the three equivalent conditions from Theorem 3.3 hold for this choice of x_0 .*

Furthermore, under these conditions, the real analytic manifold structure on M induced by the atlas in (i) is unique, in the sense that if there is another real analytic atlas on M , compatible with its C^2 structure and such that X_1, \dots, X_q are real analytic with respect to this second atlas, then the identity map $M \rightarrow M$ is a real analytic diffeomorphism between these two real analytic structures on M .

3.2 Quantitative Results

Theorem 3.3 gives necessary and sufficient conditions for a certain type of coordinate chart to exist. For applications in analysis, it is essential to have quantitative control of this coordinate chart. By using this quantitative control, these charts can be seen as generalized scaling maps in sub-Riemannian geometry—see Section 5 and [SS18, Section 7] for more details. We now turn to these quantitative results, which are the heart of this paper.

Let X_1, \dots, X_q be C^1 vector fields on a C^2 manifold \mathfrak{M} .

Definition 3.5. For $x \in \mathfrak{M}$, $\eta > 0$, and $U \subseteq \mathfrak{M}$, we say the list $X = X_1, \dots, X_q$ satisfies $\mathcal{C}(x_0, \eta, U)$ if for every $a \in B^q(\eta)$ the expression

$$e^{a_1 X_1 + \dots + a_q X_q} x_0$$

exists in U . More precisely, consider the differential equation

$$\frac{\partial}{\partial r} E(r) = a_1 X_1(E(r)) + \dots + a_q X_q(E(r)), \quad E(0) = x_0.$$

We assume that a solution to this differential equation exists up to $r = 1$, $E : [0, 1] \rightarrow U$. We have $E(r) = e^{ra_1 X_1 + \dots + ra_q X_q} x_0$.

For $1 \leq n \leq q$, we let

$$\mathcal{I}(n, q) := \{(i_1, i_2, \dots, i_n) : i_j \in \{1, \dots, q\}\} = \{1, \dots, q\}^n.$$

For $J = (j_1, \dots, j_n) \in \mathcal{I}(n, q)$ we write X_J for the list of vector fields X_{j_1}, \dots, X_{j_n} . We write $\bigwedge X_J := X_{j_1} \wedge X_{j_2} \wedge \dots \wedge X_{j_n}$.

Fix $x_0 \in \mathfrak{M}$, $\xi > 0$, $\zeta \in (0, 1]$, and set $n = \dim \text{span}\{X_1(x_0), \dots, X_q(x_0)\}$. We assume for $1 \leq j, k \leq q$,

$$[X_j, X_k] = \sum_{l=1}^q c_{j,k}^l X_l, \quad c_{j,k}^l \in C(B_X(x_0, \xi)),$$

where $B_X(x_0, \xi)$ is defined via (2.2) and is given the metric topology induced by ρ from (2.3). Proposition 3.1 applies to show that $B_X(x_0, \xi)$ is an n -dimensional, C^2 , injectively immersed submanifold of \mathfrak{M} . X_1, \dots, X_q

are C^1 vector fields on $B_X(x_0, \xi)$ and span the tangent space at every point. Henceforth, we treat X_1, \dots, X_q as vector fields on $B_X(x_0, \xi)$.

Let $J_0 \in \mathcal{I}(n, q)$ be such that $\bigwedge X_{J_0}(x_0) \neq 0$ and moreover

$$\max_{J \in \mathcal{I}(n, q)} \left| \frac{\bigwedge X_J(x_0)}{\bigwedge X_{J_0}(x_0)} \right| \leq \zeta^{-1}, \quad (3.1)$$

where $\frac{\bigwedge X_J(x_0)}{\bigwedge X_{J_0}(x_0)}$ is defined as follows. Let $\lambda : \bigwedge^n T_{x_0} B_X(x_0, \xi) \rightarrow \mathbb{R}$ be any nonzero linear functional; then

$$\frac{\bigwedge X_J(x_0)}{\bigwedge X_{J_0}(x_0)} := \frac{\lambda(\bigwedge X_J(x_0))}{\lambda(\bigwedge X_{J_0}(x_0))}. \quad (3.2)$$

Because $\bigwedge^n T_{x_0} B_X(x_0, \xi)$ is one dimensional, (3.2) is independent of the choice of λ ; see [SS18, Section 5] for more details. Notice that a $J_0 \in \mathcal{I}(n, q)$ satisfying (3.1) always exists—one can pick J_0 so that (3.1) holds with $\zeta = 1$; however it is important for some applications⁴ that we have the flexibility to choose $\zeta < 1$. Without loss of generality, reorder X_1, \dots, X_q so that $J_0 = (1, \dots, n)$.

- Let $\eta > 0$ be such that X_{J_0} satisfies $\mathcal{C}(x_0, \eta, \mathfrak{M})$.
- Let $\delta_0 > 0$ be such that for $\delta \in (0, \delta_0]$ the following holds: if $z \in B_{X_{J_0}}(x_0, \xi)$ is such that X_{J_0} satisfies $\mathcal{C}(z, \delta, B_{X_{J_0}}(x_0, \xi))$ and if $t \in B^n(\delta)$ is such that $e^{t_1 X_1 + \dots + t_n X_n} z = z$ and if $X_1(z), \dots, X_n(z)$ are linearly independent, then $t = 0$.

Remark 3.6. Because X_1, \dots, X_n are C^1 , such an $\eta > 0$ and $\delta_0 > 0$ always exist; see Lemma 7.22 and Remark 7.23. However, in general one can only guarantee that η, δ_0 are bounded below in terms of the C^1 norms of X_1, \dots, X_n in some coordinate system—and this is not a diffeomorphic invariant quantity. Thus, we state our results in terms of δ_0 and η to preserve the diffeomorphic invariance. See [SS18, Section 4.1] for a further discussion on η and δ_0 .

Key Assumption: We assume $c_{j,k}^l \in \mathcal{A}_{X_{J_0}}^{x_0, \eta}$, $1 \leq j, k, l \leq q$.

Remark 3.7. The assumption $c_{j,k}^l \in \mathcal{A}_{X_{J_0}}^{x_0, \eta}$ can be replaced with the stronger assumption⁵ $c_{j,k}^l \in C_{X_{J_0}}^{\omega, \eta}(B_{X_{J_0}}(x_0, \xi))$. Indeed, Lemma 4.3 shows $C_{X_{J_0}}^{\omega, \eta}(B_{X_{J_0}}(x_0, \xi)) \subseteq \mathcal{A}_{X_{J_0}}^{x_0, \min\{\xi, \eta\}}$.

Definition 3.8. We say C is a 0-admissible constant if C can be chosen to depend only on upper bounds for q, ζ^{-1}, ξ^{-1} , and $\|c_{j,k}^l\|_{C(B_{X_{J_0}}(x_0, \xi))}$, $1 \leq j, k, l \leq q$.

Definition 3.9. We say C is an admissible constant if C can be chosen to depend on anything a 0-admissible constant can depend on, and can also depend on upper bounds for η^{-1}, δ_0^{-1} , and $\|c_{j,k}^l\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta}}$ ($1 \leq j, k, l \leq q$).

We write $A \lesssim_0 B$ for $A \leq CB$ where C is a positive 0-admissible constant, and write $A \approx_0 B$ for $A \lesssim_0 B$ and $B \lesssim_0 A$. We write $A \lesssim B$ for $A \leq CB$ where C is a positive admissible constant, and write $A \approx B$ for $A \lesssim B$ and $B \lesssim A$.

For $t \in B^n(\eta)$ set

$$\Phi(t) = e^{t_1 X_1 + \dots + t_n X_n} x_0. \quad (3.3)$$

Let $\eta_0 := \min\{\eta, \xi\}$ so that $\Phi : B^n(\eta_0) \rightarrow B_{X_{J_0}}(x_0, \xi) \subseteq B_X(x_0, \xi)$.

Theorem 3.10 (The Quantitative Theorem). *Fix $x_0 \in \mathfrak{M}$ and let ξ, ζ, n, J_0, η , and δ_0 be as above, and suppose the Key Assumption is satisfied. Then, there exists a 0-admissible constant $\chi \in (0, \xi]$ such that:*

⁴For example, it will be essential that we may take $\zeta < 1$ in an upcoming work on similar questions with complex vector fields [Str20].

⁵A priori, $B_{X_{J_0}}(x_0, \xi)$ is not necessarily a manifold. Nevertheless, $C_{X_{J_0}}^{\omega, \eta}(B_{X_{J_0}}(x_0, \xi))$ can be defined with the same formulas. For further details on this, see [SS18, Section 2.2.1].

(a) $\forall y \in B_{X_{J_0}}(x_0, \chi), \bigwedge X_{J_0}(y) \neq 0.$

(b) $\forall y \in B_{X_{J_0}}(x_0, \chi),$

$$\sup_{J \in \mathcal{I}(n, q)} \left| \frac{\bigwedge X_J(y)}{\bigwedge X_{J_0}(y)} \right| \approx_0 1.$$

(c) $\forall \chi' \in (0, \chi], B_{X_{J_0}}(x_0, \chi')$ is an open subset of $B_X(x_0, \xi)$ and is therefore a submanifold.

There exist admissible constants $\eta_1, \xi_1, \xi_2 > 0$ such that:

(d) $\Phi(B^n(\eta_1))$ is an open subset of $B_{X_{J_0}}(x_0, \chi)$ and is therefore a submanifold of $B_X(x_0, \xi).$

(e) $\Phi : B^n(\eta_1) \rightarrow \Phi(B^n(\eta_1))$ is a C^2 diffeomorphism.

(f) $B_X(x_0, \xi_2) \subseteq B_{X_{J_0}}(x_0, \xi_1) \subseteq \Phi(B^n(\eta_1)) \subseteq B_{X_{J_0}}(x_0, \chi) \subseteq B_X(x_0, \xi).$

Let $Y_j = \Phi^* X_j$ and write $Y_{J_0} = (I + A)\nabla$, where Y_{J_0} denotes the column vector of vector fields $Y_{J_0} = [Y_1, Y_2, \dots, Y_n]^\top$, ∇ denotes the gradient in \mathbb{R}^n thought of as a column vector, and $A \in C(B^n(\eta_1); \mathbb{M}^{n \times n}).$ ⁶

(g) $A(0) = 0$ and $A \in \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ with $\|A\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \frac{1}{2}.$

(h) For $1 \leq j \leq q$, $Y_j \in \mathcal{A}^{n, \eta_1}(\mathbb{R}^n)$ and $\|Y_j\|_{\mathcal{A}^{n, \eta_1}(\mathbb{R}^n)} \lesssim 1.$

Remark 3.11. The main results of this paper (including Theorem 3.10) are invariant under arbitrary C^2 diffeomorphisms. This is true *quantitatively*—all of the estimates are unchanged when pushed forward under an arbitrary C^2 diffeomorphism; this is a consequence of Proposition 2.2. See [SS18] for more details.

3.2.1 Densities

As in [SS18, Str21], we describe how to study densities in the coordinate system given by Theorem 3.10. We refer the reader to Section 5 and [SS18, Section 7] for a further discussion of how these estimates can be used.

We take the same setting as in Theorem 3.10. Let $\chi \in (0, \xi]$ be as in that theorem and let ν be a C^1 density on $B_{X_{J_0}}(x_0, \chi).$ Suppose

$$\mathcal{L}_{X_j} \nu = f_j \nu, \quad 1 \leq j \leq n, \quad f_j \in C(B_{X_{J_0}}(x_0, \chi)),$$

where \mathcal{L}_{X_j} denotes the Lie derivative with respect to $X_j.$ We refer the reader to [Gui08] for a quick and easy to read introduction on the basics of densities (see also [Nic07] where densities are called 1-densities). We assume that there exists $r > 0$ such that $f_j \in \mathcal{A}_{X_{J_0}}^{x_0, r}.$

Definition 3.12. We say C is a $0; \nu$ -admissible constant if C is a 0-admissible constant which is also allowed to depend on upper bounds for $\|f_j\|_{C(B_{X_{J_0}}(x_0, \chi))}, 1 \leq j \leq n.$

Definition 3.13. We say C is a ν -admissible constant, if C is an admissible constant which is also allowed to depend on upper bounds for r^{-1} and $\|f_j\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}}, 1 \leq j \leq n.$

We write $A \lesssim_{0; \nu} B$ for $A \leq CB$ where C is a $0; \nu$ -admissible constant, and write $A \approx_{0; \nu} B$ for $A \lesssim_{0; \nu} B$ and $B \lesssim_{0; \nu} A.$ We similarly define \lesssim_ν and $\approx_\nu.$

Theorem 3.14. Define $h \in C^1(B^n(\eta_1))$ by $\Phi^* \nu = h \sigma_{\text{Leb}},$ where σ_{Leb} denotes the Lebesgue density on $\mathbb{R}^n.$ Then,

(a) $h(t) \approx_{0; \nu} \nu(X_1, \dots, X_n)(x_0), \forall t \in B^n(\eta_1).$ In particular, $h(t)$ always has the same sign, and is either never zero, or always zero.

⁶Here, and in the rest of the paper, $\mathbb{M}^{n \times n}$ denotes the Banach space of $n \times n$ real matrices endowed with the usual operator norm.

(b) Set $s := \min\{\eta_1, r\}$, where η_1 is as in Theorem 3.10. Then, $h \in \mathcal{A}^{n,s}$ and $\|h\|_{\mathcal{A}^{n,s}} \lesssim_\nu |\nu(X_1, \dots, X_n)(x_0)|$.

Corollary 3.15. *Let ξ_2 be as in Theorem 3.10. Then,*

$$\nu(B_{X_{j_0}}(x_0, \xi_2)) \approx_\nu \nu(B_X(x_0, \xi_2)) \approx_\nu \nu(X_1, \dots, X_n)(x_0), \quad (3.4)$$

and therefore,

$$|\nu(B_{X_{j_0}}(x_0, \xi_2))| \approx_\nu |\nu(B_X(x_0, \xi_2))| \approx_\nu |\nu(X_1, \dots, X_n)(x_0)| \approx_0 \max_{j_1, \dots, j_n \in \{1, \dots, q\}} |\nu(X_{j_1}, \dots, X_{j_n})(x_0)|. \quad (3.5)$$

4 Function Spaces Revisited

In this section, we present the basic results we need concerning the function spaces defined in Section 2. Let M be a C^2 manifold and let X_1, \dots, X_q be C^1 vector fields on M and let X denote the list $X = X_1, \dots, X_q$. Let $\Omega \subseteq \mathbb{R}^n$ be an open set.

Lemma 4.1. *The spaces $C^{\omega,r}(\Omega)$, $\mathcal{A}^{n,r}$, $C_X^{\omega,r}(M)$, and $\mathcal{A}_X^{x_0,r}$ are Banach algebras. In particular, if \mathcal{Y} denotes any one of these spaces and if $x, y \in \mathcal{Y}$, then $\|xy\|_{\mathcal{Y}} \leq \|x\|_{\mathcal{Y}} \|y\|_{\mathcal{Y}}$. More generally, the same holds for the analogous spaces of functions taking values in a Banach algebra.*

Proof. We prove only the result for functions taking values in \mathbb{C} ; the same proof proves the more general result for functions taking values in a Banach algebra.

We begin with proof for $\mathcal{A}^{n,r}$. Suppose $f, g \in \mathcal{A}^{n,r}$. Then if $f(t) = \sum_{\alpha \in \mathbb{N}^n} \frac{c_\alpha}{\alpha!} t^\alpha$ and $g(t) = \sum_{\alpha \in \mathbb{N}^n} \frac{d_\alpha}{\alpha!} t^\alpha$, we have $f(t)g(t) = \sum_{\alpha, \beta \in \mathbb{N}^n} \frac{c_\alpha d_\beta}{\alpha! \beta!} t^{\alpha+\beta}$, and therefore,

$$\|fg\|_{\mathcal{A}^{n,r}} \leq \sum_{\alpha, \beta \in \mathbb{N}^n} \frac{|c_\alpha d_\beta|}{\alpha! \beta!} r^{|\alpha|+|\beta|} \leq \left(\sum_{\alpha \in \mathbb{N}^n} \frac{|c_\alpha|}{\alpha!} r^{|\alpha|} \right) \left(\sum_{\beta \in \mathbb{N}^n} \frac{|d_\beta|}{\beta!} r^{|\beta|} \right) = \|f\|_{\mathcal{A}^{n,r}} \|g\|_{\mathcal{A}^{n,r}},$$

completing the proof for $\mathcal{A}^{n,r}$. The result for $\mathcal{A}_X^{x_0,r}$ follows immediately from the result for $\mathcal{A}^{n,r}$.

Next we consider $C_X^{\omega,r}(M)$. For this, we need some notation from [Nel59]—we refer the reader to that reference for more detailed information on these definitions. Let \mathcal{Y} be a Banach space and let $\mathcal{O}(\mathcal{Y})$ denote the set of all (bounded or unbounded) operators on \mathcal{Y} . For $A \in \mathcal{O}(\mathcal{Y})$, we write $|A|$ for the set consisting of A alone. Let $|\mathcal{O}(\mathcal{Y})|$ be the free abelian semigroup with the set of all $|A|$, $A \in \mathcal{O}(\mathcal{Y})$, as generators. Let $\alpha, \beta \in |\mathcal{O}(\mathcal{Y})|$ so that $\alpha = |A_1| + \dots + |A_l|$, $\beta = |B_1| + \dots + |B_m|$ (where these are formal sums). We define $\alpha\beta = \sum_{i=1}^l \sum_{j=1}^m |A_i B_j| \in |\mathcal{O}(\mathcal{Y})|$.

Let $A_1, \dots, A_l \in \mathcal{O}(\mathcal{Y})$, and set $\alpha = |A_1| + \dots + |A_l|$. For $y \in \mathcal{Y}$ in the domains of A_1, \dots, A_l , we define $\|\alpha y\| := \|A_1 y\|_{\mathcal{Y}} + \dots + \|A_l y\|_{\mathcal{Y}}$. For $f = (f_1, \dots, f_m) \in \mathcal{Y}^m$, with each f_j in the domains of A_1, \dots, A_l , define $\mathcal{A}f := (A_j f_k)_{1 \leq j \leq l, 1 \leq k \leq m} \in \mathcal{Y}^{ml}$. Note that $\|\alpha x\| = \|\mathcal{A}x\|_{\mathcal{Y}^{ml}}$, and more generally, $\|\alpha^m x\| = \|\mathcal{A}^m x\|_{\mathcal{Y}^{l^m m}}$. Here we are giving \mathcal{Y}^m the norm $\|f\|_{\mathcal{Y}^m} := \sum_{l=1}^m \|f_l\|_{\mathcal{Y}}$.

Now suppose \mathcal{Y} is a Banach algebra, and suppose $A_1, \dots, A_l \in \mathcal{O}(\mathcal{Y})$ satisfy $A_j(xy) = (A_j(x))y + x(A_j(y))$ (and the domains of A_1, \dots, A_l are algebras). For $f = (f_1, \dots, f_{m_1}) \in \mathcal{Y}^{m_1}$ and $g = (g_1, \dots, g_{m_2}) \in \mathcal{Y}^{m_2}$ set $fg = (f_j g_k)_{1 \leq j \leq m_1, 1 \leq k \leq m_2} \in \mathcal{Y}^{m_1 m_2}$. We then have, for $x, y \in \mathcal{Y}$ (and x and y in the domains of the appropriate operators),

$$\mathcal{A}^k(xy) = \sum_{k_1+k_2=k} \binom{k}{k_1} (\mathcal{A}^{k_1}x)(\mathcal{A}^{k_2}y),$$

and in particular

$$\|\alpha^k(xy)\| = \|\mathcal{A}^k(xy)\|_{\mathcal{Y}^{lk}} \leq \sum_{k_1+k_2=k} \binom{k}{k_1} \|\mathcal{A}^{k_1}x\|_{\mathcal{Y}^{lk_1}} \|\mathcal{A}^{k_2}y\|_{\mathcal{Y}^{lk_2}} = \sum_{k_1+k_2=k} \binom{k}{k_1} \|\alpha^{k_1}x\| \|\alpha^{k_2}y\|. \quad (4.1)$$

We now specialize to the case $\mathcal{B} = C(M)$ and $\alpha = |X_1| + |X_2| + \dots + |X_q|$. We have, using (4.1) and the definition of $\|\cdot\|_{C_X^{\omega,r}}$,

$$\begin{aligned} \|fg\|_{C_X^{\omega,r}(M)} &= \sum_{k=0}^{\infty} \frac{1}{k!} \|\alpha^k(fg)\|_{r^k} \leq \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{k_1+k_2=k} \binom{k}{k_1} \|\alpha^{k_1}f\| \|\alpha^{k_2}g\|_{r^{k_1}r^{k_2}} \\ &= \left(\sum_{k_1=0}^{\infty} \frac{1}{k_1!} \|\alpha^{k_1}f\|_{r^{k_1}} \right) \left(\sum_{k_2=0}^{\infty} \frac{1}{k_2!} \|\alpha^{k_2}g\|_{r^{k_2}} \right) = \|f\|_{C_X^{\omega,r}(M)} \|g\|_{C_X^{\omega,r}(M)}. \end{aligned}$$

This completes the proof for $C_X^{\omega,r}(M)$. Since $C^{\omega,r}(\Omega) = C_{\nabla}^{\omega,r}(\Omega)$ (with equality of norms), the result for $C^{\omega,r}(\Omega)$ follows as well. \square

The spaces $C_X^{\omega,r}(M)$, and $\mathcal{A}_X^{x_0,r}$ are closely related as the next three results show.

Lemma 4.2. (i) $C^{\omega,r}(B^n(r)) \subseteq \mathcal{A}^{n,r}$ and $\|f\|_{\mathcal{A}^{n,r}} \leq \|f\|_{C^{\omega,r}(B^n(r))}$.

(ii) $\mathcal{A}^{n,r} \subseteq C^{\omega,r/2}(B^n(r/2))$ and $\|f\|_{C^{\omega,r/2}(B^n(r/2))} \leq \|f\|_{\mathcal{A}^{n,r}}$.

Proof. (i) is a special case of Lemma 4.3, below, so we only prove (ii). We use the identity, for multi-indices $\alpha \in \mathbb{N}^n$,

$$\sum_{\beta \leq \alpha} \binom{\alpha}{\beta} = 2^{|\alpha|},$$

where the sum is taken over all $\beta \in \mathbb{N}^n$ with $\beta_j \leq \alpha_j$ for all j .

Suppose $f \in \mathcal{A}^{n,r}$. Then, $f(t) = \sum_{\alpha \in \mathbb{N}^n} \frac{c_\alpha}{\alpha!} t^\alpha$ with $\|f\|_{\mathcal{A}^{n,r}} = \sum_{\alpha \in \mathbb{N}^n} \frac{|c_\alpha|}{\alpha!} r^{|\alpha|} < \infty$. Set $r_1 = r/2$. We have

$$\begin{aligned} \|f\|_{C^{\omega,r_1}(B^n(r_1))} &= \sum_{\beta \in \mathbb{N}^n} \frac{r_1^{|\beta|}}{\beta!} \|\partial_t^\beta f\|_{C(B^n(r_1))} = \sum_{\beta \in \mathbb{N}^n} \frac{r_1^{|\beta|}}{\beta!} \sup_{t \in B^n(r_1)} \left| \sum_{\alpha \geq \beta} \frac{c_\alpha}{(\alpha - \beta)!} t^{\alpha - \beta} \right| \\ &\leq \sum_{\beta \in \mathbb{N}^n} \sum_{\alpha \geq \beta} \frac{r_1^{|\alpha|}}{\beta! (\alpha - \beta)!} |c_\alpha| = \sum_{\alpha \in \mathbb{N}^n} \frac{r_1^{|\alpha|}}{\alpha!} |c_\alpha| \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} = \sum_{\alpha \in \mathbb{N}^n} \frac{2^{|\alpha|} r_1^{|\alpha|}}{\alpha!} |c_\alpha| = \|f\|_{\mathcal{A}^{n,r}}, \end{aligned}$$

completing the proof. \square

Lemma 4.3. Suppose $X = X_1, \dots, X_q$ satisfies $\mathcal{C}(x_0, r, M)$. Then, $C_X^{\omega,r}(M) \subseteq \mathcal{A}_X^{x_0,r}$ and $\|f\|_{\mathcal{A}_X^{x_0,r}} \leq \|f\|_{C_X^{\omega,r}(M)}$.

Proof. We will show, for $f \in C_X^{\omega,r}(M)$ that

$$f(e^{t_1 X_1 + \dots + t_q X_q} x_0) = \sum_{m=0}^{\infty} \frac{((t_1 X_1 + \dots + t_q X_q)^m f)(x_0)}{m!} =: h(t), \quad t \in B^q(r). \quad (4.2)$$

The result will follow since the hypothesis $f \in C_X^{\omega,r}(M)$ implies that the sum in (4.2) converges absolutely for $|t| \leq r$, and $\|f\|_{\mathcal{A}_X^{x_0,r}} = \|h\|_{\mathcal{A}^{q,r}} \leq \|f\|_{C_X^{\omega,r}(M)}$.

Fix $t \in B^q(r)$. For $\delta > 0$ small (depending on t) and for $s_1, s_2 \in (-1 - \delta, 1 + \delta)$ define

$$g(s_1, s_2) = \sum_{m=0}^{\infty} \frac{s_1^m}{m!} ((t_1 X_1 + \dots + t_q X_q)^m f) \left(e^{s_2(t_1 X_1 + \dots + t_q X_q)} x_0 \right). \quad (4.3)$$

Since X satisfies $\mathcal{C}(x_0, r, M)$, for $s_2 \in (-1 - \delta, 1 + \delta)$ we have $e^{s_2(t_1 X_1 + \dots + t_q X_q)} x_0 \in M$, and therefore the sum in (4.3) converges absolutely by the hypothesis that $f \in C_X^{\omega,r}(M)$ (here we are taking δ small, depending on t). Hence $g(s_1, s_2)$ is defined for $s_1, s_2 \in (-1 - \delta, 1 + \delta)$.

We have,

$$\frac{\partial}{\partial s_1} g(s_1, s_2) = \sum_{m=0}^{\infty} \frac{s_1^m}{m!} \left((t_1 X_1 + \dots + t_q X_q)^{m+1} f \right) \left(e^{s_2(t_1 X_1 + \dots + t_q X_q)} x_0 \right) = \frac{\partial}{\partial s_2} g(s_1, s_2).$$

We conclude $g(s, 0) = g(0, s)$ for $s \in (-1 - \delta, 1 + \delta)$. In particular, $g(1, 0) = g(0, 1)$, which is exactly (4.2), completing the proof. \square

Unlike the Euclidean case in Lemma 4.2, the reverse containment to Lemma 4.3 is a more involved and requires more hypotheses. In fact, we see it as a corollary of Theorem 3.10.

Corollary 4.4. *We take all the same hypotheses and notation as in Theorem 3.10 and define admissible constants as in that theorem. Fix $r \in (0, \eta_1]$ (where η_1 is as in Theorem 3.10). Then, there is an admissible constant $s = s(r) > 0$ such that $\mathcal{A}_{X_{J_0}}^{x_0, r} \subseteq C_X^{\omega, s}(\Phi(B^n(r/2)))$. Moreover, there is an admissible constant $C = C(r)$ such that*

$$\|f\|_{C_X^{\omega, s}(\Phi(B^n(r/2)))} \leq C \|f\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}}. \quad (4.4)$$

Remark 4.5. Notice that $\|f\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}} \leq \|f\|_{\mathcal{A}_X^{x_0, r}}$ and so one may replace X_{J_0} with X throughout Corollary 4.4.

Proof of Corollary 4.4. Let $\Phi(t) = e^{t_1 X_1 + \dots + t_n X_n} x_0$ be as in Theorem 3.10. Suppose $f \in \mathcal{A}_{X_{J_0}}^{x_0, r}$; so that, by the definition of $\mathcal{A}_{X_{J_0}}^{x_0, r}$, $\Phi^* f \in \mathcal{A}^{n, r}$ with $\|\Phi^* f\|_{\mathcal{A}^{n, r}} = \|f\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}}$. By Lemma 4.2 (ii), $\Phi^* f \in C^{\omega, r/2}(B^n(r/2))$ with $\|\Phi^* f\|_{C^{\omega, r/2}(B^n(r/2))} \leq \|f\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}}$. Letting $Y_j = \Phi^* X_j$ as in Theorem 3.10, we have that $Y_1, \dots, Y_q \in \mathcal{A}^{n, \eta_1}(\mathbb{R}^n) \subseteq C^{\omega, \eta_1/2}(B^n(\eta_1/2); \mathbb{R}^n)$ with $\|Y_j\|_{C^{\omega, \eta_1/2}(B^n(\eta_1/2); \mathbb{R}^n)} \leq \|Y_j\|_{\mathcal{A}^{n, \eta_1}(\mathbb{R}^n)} \lesssim 1$ (where we have again used Lemma 4.2 (ii)).

Proposition 4.7 (below) shows that there exists an admissible $s = s(r) > 0$ such that $\Phi^* f \in C^{\omega, r/2}(B^n(r/2)) = C_Y^{\omega, s}(B^n(r/2))$ and

$$\|\Phi^* f\|_{C_Y^{\omega, s}(B^n(r/2))} \leq C \|\Phi^* f\|_{C^{\omega, r/2}(B^n(r/2))} \leq C \|f\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}},$$

where C is as in the statement of the corollary. Because $\Phi_* Y_j = X_j$, Proposition 2.2 implies $f \in C_X^{\omega, s}(\Phi(B^n(r/2)))$ with $\|f\|_{C_X^{\omega, s}(\Phi(B^n(r/2)))} = \|\Phi^* f\|_{C_Y^{\omega, s}(B^n(r/2))} \leq C \|f\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}}$, completing the proof. \square

Lemma 4.6. *For any $s \in (0, r)$, $X_j : C_X^{\omega, r}(M) \rightarrow C_X^{\omega, s}(M)$. Furthermore, for $f \in C_X^{\omega, r}(M)$, $\|X_j f\|_{C_X^{\omega, s}(M)} \leq (\sup_{m \in \mathbb{N}} (s/r)^m \frac{m+1}{r}) \|f\|_{C_X^{\omega, r}(M)}$.*

Proof. Let $f \in C_X^{\omega, r}(M)$, and consider

$$\begin{aligned} \|X_j f\|_{C_X^{\omega, s}(M)} &= \sum_{m=0}^{\infty} \frac{s^m}{m!} \sum_{|\alpha|=m} \|X^\alpha X_j f\|_{C(M)} \leq \sum_{m=0}^{\infty} (s/r)^m \frac{m+1}{r} \frac{r^{m+1}}{(m+1)!} \sum_{|\alpha|=m+1} \|X^\alpha f\|_{C(M)} \\ &\leq \left(\sup_{m \in \mathbb{N}} (s/r)^m \frac{m+1}{r} \right) \|f\|_{C_X^{\omega, r}(M)} < \infty, \end{aligned}$$

where in the last inequality, we have used $s < r$. The result follows. \square

4.1 Comparison with Euclidean function spaces and a result of Nelson

Let $\Omega \subseteq \mathbb{R}^n$ be an open set. If Y_1, \dots, Y_q are real analytic vector fields on Ω which span the tangent space at every point, it is a result of Nelson [Nel59, Theorem 2] that being real analytic with respect to Y_1, \dots, Y_q is the same as being real analytic in the classical sense. We state a quantitative version of this.

Proposition 4.7. *Fix $r > 0$, and let $Y_1, \dots, Y_q \in C^{\omega, r}(\Omega; \mathbb{R}^n)$.*

- (i) There exists $s > 0$ such that $C^{\omega,r}(\Omega) \subseteq C_Y^{\omega,s}(\Omega)$ and $\|f\|_{C_Y^{\omega,s}(\Omega)} \leq C\|f\|_{C^{\omega,r}(\Omega)}$, $\forall f \in C^{\omega,r}(\Omega)$, where s and C can be chosen to depend only on upper bounds for q , n , r^{-1} , and $\|Y_j\|_{C^{\omega,r}(\Omega;\mathbb{R}^n)}$ ($1 \leq j \leq q$).
- (ii) Suppose, in addition, that for $1 \leq j \leq n$, $1 \leq k \leq q$, there exists $b_j^k \in C^{\omega,r}(\Omega)$ such that $\frac{\partial}{\partial x_j} = \sum_{k=1}^q b_j^k Y_k$. For all $r_1 > 0$, there exists $s' > 0$ such that $C_Y^{\omega,r_1}(\Omega) \subseteq C^{\omega,s'}(\Omega)$ and $\|f\|_{C^{\omega,s'}(\Omega)} \leq C\|f\|_{C_Y^{\omega,r_1}(\Omega)}$, $\forall f \in C_Y^{\omega,r_1}(\Omega)$, where s' and C can be chosen to depend only on upper bounds for q , r^{-1} , r_1^{-1} , $\|Y_j\|_{C^{\omega,r}(\Omega;\mathbb{R}^n)}$ ($1 \leq j \leq q$), and $\|b_j^k\|_{C^{\omega,r}(\Omega)}$ ($1 \leq j \leq n$, $1 \leq k \leq q$).

Proof. This follows from a straightforward modification of the proof of [Nel59, Theorem 2] and we leave the details to the reader. \square

Remark 4.8. In the sequel, we only use (i) of Proposition 4.7.

5 Sub-Riemannian Geometry and Scaling

One of the main applications of results like Theorem 3.10 is as scaling maps for sub-Riemannian geometries. Such scaling maps were first introduced by Nagel, Stein, and Wainger [NSW85], and were further studied by many other authors including Tao and Wright [TW03], the author [Str11], Montanari and Morbidelli [MM12], and most recently in the first two parts of this series [SS18, Str21]. Since Nagel, Stein, and Wainger's results, these ideas have been used in a wide variety of problems. For a description of some of these applications, see the notes at the end of Chapter 2 of [Str14].

Nagel, Stein, and Wainger's results worked in the smooth category⁷. The later papers either worked in the smooth category, or with a finite level of smoothness. Thus, if one starts with a sub-Riemannian geometry based on real analytic vector fields, the results in these works do not yield appropriate quantitative control in the real analytic setting and therefore these results destroy the real analytic nature of the problem under consideration. Theorem 3.10 fixes this issue.

Furthermore, when dealing with real analytic vector fields, one does not need to assume Hörmander's condition on the vector fields. See Section 5.1.2.

We present such results, here.

5.1 Classical Real Analytic Sub-Riemannian Geometries

In this section, we describe a real analytic version of the foundational work of Nagel, Stein, and Wainger [NSW85], and see how it is a special case of Theorem 3.10. This is the simplest non-trivial setting where the results in this paper can be seen as providing scaling maps adapted to a sub-Riemannian geometry.

Let X_1, \dots, X_q be real analytic vector fields on an open set $\Omega \subseteq \mathbb{R}^n$; we assume X_1, \dots, X_q span the tangent space at every point of Ω . To each X_j assign a formal degree $d_j \in [1, \infty)$. We assume $\forall x \in \Omega$ there exists an open neighborhood $U_x \subseteq \Omega$ of x such that:

$$[X_j, X_k] = \sum_{d_l \leq d_j + d_k} c_{j,k}^{l,x} X_l, \quad c_{j,k}^{l,x} \in C^\omega(U_x). \quad (5.1)$$

We write (X, d) for the list $(X_1, d_1), \dots, (X_q, d_q)$ and for $\delta > 0$ write $\delta^d X$ for the list of vector fields $\delta^{d_1} X_1, \dots, \delta^{d_q} X_q$. The sub-Riemannian ball associated to (X, d) centered at $x_0 \in \Omega$, of radius $\delta > 0$ is defined by

$$B_{(X,d)}(x_0, \delta) := B_{\delta^d X}(x_0, 1),$$

where the later ball is defined by (2.2). $B_{(X,d)}(x_0, \delta)$ is an open subset of Ω . It is easy to see that the balls $B_{(X,d)}(x, \delta)$ are metric balls.

⁷More precisely, the quantitative estimates they proved involved C^m norms, for various $m \in \mathbb{N}$.

Define, for $x \in \Omega$, $\delta \in (0, 1]$,

$$\Lambda(x, \delta) := \max_{j_1, \dots, j_n \in \{1, \dots, q\}} |\det(\delta^{d_{k_1}} X_{k_1}(x) \cdots \delta^{d_{k_n}} X_{k_n}(x))|.$$

For each $x \in \Omega$, $\delta \in (0, 1]$, pick $j_1 = j_1(x, \delta), \dots, j_n = j_n(x, \delta)$ so that

$$|\det(\delta^{d_{j_1}} X_{j_1}(x) \cdots \delta^{d_{j_n}} X_{j_n}(x))| = \Lambda(x, \delta).$$

For this choice of $j_1 = j_1(x, \delta), \dots, j_n = j_n(x, \delta)$, define

$$\Phi_{x, \delta}(t_1, \dots, t_n) := \exp(t_1 \delta^{d_{j_1}} X_{j_1} + \cdots + t_n \delta^{d_{j_n}} X_{j_n})x.$$

We let σ_{Leb} denote the usual Lebesgue density on Ω .

Theorem 5.1. *Fix a compact set $\mathcal{K} \Subset \Omega$.⁸ In what follows, we write $A \lesssim B$ for $A \leq CB$ where C is a positive constant which may depend on \mathcal{K} , but does not depend on the particular points $x \in \mathcal{K}$ or $u \in \mathbb{R}^n$, or the scale $\delta \in (0, 1]$; we write $A \approx B$ for $A \lesssim B$ and $B \lesssim A$. Under the above described hypotheses, there exists $\eta_1, \xi_0 \approx 1$, such that $\forall x \in \mathcal{K}$,*

(i) $\sigma_{\text{Leb}}(B_{(X, d)}(x, \delta)) \approx \Lambda(x, \delta)$, $\forall \delta \in (0, \xi_0]$.

(ii) $\sigma_{\text{Leb}}(B_{(X, d)}(x, 2\delta)) \lesssim \sigma_{\text{Leb}}(B_{(X, d)}(x, \delta))$, $\forall \delta \in (0, \xi_0/2]$.

(iii) $\forall \delta \in (0, 1]$, $\Phi_{x, \delta}(B^n(\eta_1)) \subseteq \Omega$ is open and $\Phi_{x, \delta} : B^n(\eta_1) \rightarrow \Phi_{x, \delta}(B^n(\eta_1))$ is a real analytic diffeomorphism.

(iv) Define $h_{x, \delta}(t)$ by $h_{x, \delta} \sigma_{\text{Leb}} = \Phi_{x, \delta}^* \sigma_{\text{Leb}}$. Then, $h_{x, \delta}(t) \approx \Lambda(x, \delta)$, $\forall t \in B^n(\eta_1)$, and there exists $s \approx 1$ with $\|h_{x, \delta}\|_{\mathcal{A}^{n, s}} \lesssim \Lambda(x, \delta)$.

(v) $B_{(X, d)}(x, \xi_0 \delta) \subseteq \Phi_{x, \delta}(B^n(\eta_1)) \subseteq B_{(X, d)}(x, \delta)$, $\forall \delta \in (0, 1]$.

(vi) Let $Y_j^{x, \delta} := \Phi_{x, \delta}^* \delta^{d_j} X_j$, $1 \leq j \leq q$, so that $Y_j^{x, \delta}$ is a real analytic vector field on $B^n(\eta_1)$. We have

$$\|Y_j^{x, \delta}\|_{\mathcal{A}^{n, \eta_1}(\mathbb{R}^n)} \lesssim 1, \quad 1 \leq j \leq q. \quad (5.2)$$

Finally, $Y_1^{x, \delta}(u), \dots, Y_q^{x, \delta}(u)$ span $T_u B^n(\eta_1)$, uniformly in x, δ , and u , in the sense that

$$\max_{k_1, \dots, k_n \in \{1, \dots, q\}} \inf_{u \in B^n(\eta_1)} |\det(Y_{k_1}^{x, \delta}(u) \cdots Y_{k_n}^{x, \delta}(u))| \approx 1. \quad (5.3)$$

Remark 5.2. If real analytic was replaced in this entire section with C^∞ (and the estimates in (5.2) were replaced with appropriate estimates of C^m norms), then Theorem 5.1 is the main result, in a slightly different language, of Nagel, Stein, and Wainger's work [NSW85]—see [SS18, Section 7.1] for a further discussion. A main consequence of the results of this paper is that one can obtain good estimates on the real analyticity of the vector fields $Y_1^{x, \delta}, \dots, Y_q^{x, \delta}$ —see (5.2).

Proof. By a simple partition of unity argument, we may write

$$[X_j, X_k] = \sum_{d_l \leq d_j + d_k} c_{j, k}^l X_l, \quad c_{j, k}^l \in C_{\text{loc}}^\infty(\Omega).$$

Using this, most of Theorem 5.1 is contained in [SS18, Section 7.1]; the only parts which are not are those which relate to real analyticity. In particular, (i), (ii), (iii) (with real analytic replaced by C^∞), (v), and (5.3) are all explicitly stated in [SS18, Section 7.1]. Furthermore, since X_1, \dots, X_q are real analytic, $\Phi_{x, \delta}$

⁸We write $A \Subset B$ to mean A is a relatively compact subset of B .

is real analytic (by classical theorems), so (iii) follows. Thus, the new parts are (iv) and (5.2). These are simple consequences of Theorem 3.14 and Theorem 3.10, respectively, though we will see them as part of a more general theorem: Theorem 5.7, below.

Hence, to complete the proof, we show how Theorem 5.7 applies to this setting. Without loss of generality, we may shrink each U_x so that $U_x \Subset \Omega$. Set

$$X_j^\delta := \delta^{d_j} X_j, \quad \hat{c}_{j,k}^{l,x,\delta} := \begin{cases} \delta^{d_j+d_k-d_l} c_{j,k}^{l,x}, & d_l \leq d_j + d_k, \\ 0, & \text{otherwise,} \end{cases}$$

so that

$$[X_j^\delta, X_k^\delta] = \sum_{l=1}^q \hat{c}_{j,k}^{l,x,\delta} X_l^\delta, \quad \text{on } U_x.$$

We let X^δ denote the list $X_1^\delta, \dots, X_q^\delta$. Fix an open set Ω' with $\mathcal{K} \Subset \Omega' \Subset \Omega$ and set $\mathcal{K}_1 := \overline{\Omega'}$, so that $\mathcal{K}_1 \Subset \Omega$ is compact. Take $s_1 > 0$ so that $X_1, \dots, X_q \in C^{\omega, s_1}(\Omega'; \mathbb{R}^n)$. By the Phragmen-Lindelöf principle, we may take $\xi' > 0$ so small that $B_X(x, \xi') \subseteq \Omega' \subset \mathcal{K}_1$, $\forall x \in \mathcal{K}$. $\{U_x : x \in \mathcal{K}_1\}$ is an open cover for \mathcal{K}_1 and we extract a finite sub-cover U_{x_1}, \dots, U_{x_R} . The balls $B_X(x, \delta)$ are metric balls and the topology induced by these balls is the same as the usual topology on Ω . Let $\xi \in (0, \min\{\xi', 1\}]$ be less than or equal to the Lebesgue number for the cover U_{x_1}, \dots, U_{x_R} of \mathcal{K}_1 , with respect to the metric associated to the balls $B_X(x, \delta)$. Thus, since $\xi \leq \xi'$, $\forall x \in \mathcal{K}$, $\exists r \in \{1, \dots, R\}$ with $B_X(x, \xi) = B_X(x, \xi) \cap \mathcal{K}_1 \subseteq U_{x_r}$. For this choice of r , set $c_{j,k}^{l,x,\delta} := \hat{c}_{j,k}^{l,x_r,\delta}$. Take $s_2 > 0$ so that $\forall r \in \{1, \dots, R\}$, $c_{j,k}^{l,x_r} \in C^{\omega, s_2}(U_{x_r})$. By Proposition 4.7 (using $X_1, \dots, X_q \in C^{\omega, s_1}(\Omega'; \mathbb{R}^n)$), there exists $s \approx 1$ with $c_{j,k}^{l,x_r} \in C_X^{\omega, s}(U_{x_r})$, $1 \leq r \leq R$, $1 \leq j, k, l \leq q$, and

$$\|c_{j,k}^{l,x_r}\|_{C_X^{\omega, s}(U_{x_r})} \lesssim \|c_{j,k}^{l,x_r}\|_{C^{\omega, s_1}(U_{x_r})} \lesssim 1, \quad 1 \leq r \leq R, 1 \leq j, k, l \leq q.$$

Thus, by tracing through the definitions we have, for $x \in \mathcal{K}$, $\delta \in (0, 1]$, $c_{j,k}^{l,x,\delta} \in C_X^{\omega, s}(B_{X^\delta}(x, \xi))$ and

$$\sup_{\substack{x \in \mathcal{K} \\ \delta \in (0, 1]}} \|c_{j,k}^{l,x,\delta}\|_{C_X^{\omega, s}(B_{X^\delta}(x, \xi))} \leq \sup_{1 \leq r \leq R} \|c_{j,k}^{l,x_r}\|_{C_X^{\omega, s}(U_{x_r})} \lesssim 1.$$

Define f_j^δ by $f_j^\delta \sigma_{\text{Leb}} = \mathcal{L}_{X_j^\delta} \sigma_{\text{Leb}}$; i.e., $f_j^\delta = \delta^{d_j} f_j$, where $f_j := \nabla \cdot X_j$. By our hypotheses, we have $f_j \in C_{\text{loc}}^\omega(\Omega)$, $1 \leq j \leq q$. It follows that there exists $s_3 > 0$ with $f_j \in C^{\omega, s_3}(\Omega')$, $1 \leq j \leq q$. By Proposition 4.7 (using that $X_1, \dots, X_q \in C^{\omega, s_1}(\Omega'; \mathbb{R}^n)$), there exists $r \approx 1$ with $f_j \in C_X^{\omega, r}(\Omega')$. Directly from the definitions, we now have

$$\sup_{\delta \in (0, 1]} \|f_j^\delta\|_{C_X^{\omega, r}(\Omega')} \leq \|f_j\|_{C_X^{\omega, r}(\Omega')} \lesssim 1.$$

Using the above remarks, the result now follows directly from Theorem 5.7. \square

Remark 5.3. The most important part of Theorem 5.1 is (vi); which allows us to think of $\Phi_{x,\delta}$ as a scaling map. Indeed, one thinks of the vector fields $\delta^{d_1} X_1, \dots, \delta^{d_q} X_q$ as being “small” (for δ small). However, $\Phi_{x,\delta}$ gives a coordinate system in which these vector fields are unit size. Indeed, the vector fields $Y_1^{x,\delta}, \dots, Y_q^{x,\delta}$ are the vector fields $\delta^{d_1} X_1, \dots, \delta^{d_q} X_q$ written in the coordinate system given by $\Phi_{x,\delta}$. These vector fields are real analytic uniformly in x and δ (i.e., (5.2)) and span the tangent space uniformly in x and δ (i.e., (5.3)). Thus, we have “rescaled” the vector fields to be unit size.

5.1.1 Hörmander’s condition

The main way that Theorem 5.1 arises is via vector fields which satisfy Hörmander’s condition. Suppose V_1, \dots, V_r are real analytic vector fields on an open set $\Omega \subseteq \mathbb{R}^n$. We assume V_1, \dots, V_r satisfy Hörmander’s condition of order m on Ω . I.e., we assume that the finite list of vector fields

$$V_1, \dots, V_r, \dots, [V_i, V_j], \dots, [V_i, [V_j, V_k]], \dots, \dots, \text{commutators of order } m,$$

span the tangent space at every point of Ω .

To each V_1, \dots, V_r , we assign the formal degree 1. If Z has formal degree e , we assign to $[V_j, Z]$ the formal degree $e + 1$. Let $(X_1, d_1), \dots, (X_q, d_q)$ denote the finite list of vector fields with formal degree $d_j \leq m$. Hörmander's condition implies X_1, \dots, X_q span the tangent space at every point of Ω .

We claim (5.1) holds, and therefore Theorem 5.1 applies to $(X_1, d_1), \dots, (X_q, d_q)$. Indeed, if $d_j + d_k \leq m$ we have

$$[X_j, X_k] = \sum_{d_l = d_j + d_k} c_{j,k}^l X_l,$$

where $c_{j,k}^l$ are constants by the Jacobi identity. If $d_j + d_k > m$, then since X_1, \dots, X_q are real analytic and span the tangent space at every point, we have $\forall x \in \Omega$ there exists a neighborhood $U_x \subseteq \Omega$ of x such that

$$[X_j, X_k] = \sum_{l=1}^q c_{j,k}^{l,x} X_l = \sum_{d_l \leq d_j + d_k} c_{j,k}^{l,x} X_l, \quad c_{j,k}^{l,x} \in C^\omega(U_x).$$

Thus, (5.1) holds and Theorem 5.1 applies.

Let $\mathcal{K} \Subset \Omega$ be a compact set. Applying Theorem 5.1 for $\delta \in (0, 1]$, $x \in \mathcal{K}$, we obtain $\eta_1 > 0$ and $\Phi_{x,\delta} : B^n(\eta_1) \rightarrow B_{(X,d)}(x, \delta)$ as in that theorem. Set $V_j^{x,\delta} := \Phi_{x,\delta}^* \delta V_j$. If $d_k = l$, then

$$X_k = [V_{j_1}, [V_{j_2}, \dots, [V_{j_{l-1}}, V_{j_l}]]],$$

and so

$$\Phi_{x,\delta}^* \delta^{d_k} X_k = \Phi_{x,\delta}^* [\delta V_{j_1}, [\delta V_{j_2}, \dots, [\delta V_{j_{l-1}}, \delta V_{j_l}]]] = [V_{j_1}^{x,\delta}, [V_{j_2}^{x,\delta}, \dots, [V_{j_{l-1}}^{x,\delta}, V_{j_l}^{x,\delta}]]].$$

Theorem 5.1 implies the vector fields $\Phi_{x,\delta}^* \delta^{d_j} X_j$ are real analytic and span the tangent space, uniformly for $x \in \mathcal{K}$, $\delta \in (0, 1]$. We conclude that the vector fields $V_1^{x,\delta}, \dots, V_q^{x,\delta}$ are real analytic and satisfy Hörmander's condition, uniformly for $x \in \mathcal{K}$ and $\delta \in (0, 1]$. In short, the map $\Phi_{x,\delta}^*$ takes $\delta V_1, \dots, \delta V_r$ to $V_1^{x,\delta}, \dots, V_r^{x,\delta}$ which are real analytic and satisfy Hörmander's condition "uniformly"; i.e., it takes the case of δ small and rescales it to the case $\delta = 1$, while preserving real analyticity in a quantitative way.

5.1.2 Beyond Hörmander's condition

Let V_1, \dots, V_r be real analytic vector fields defined on an open set $\Omega \subseteq \mathbb{R}^n$. It turns out that the main conclusions of Section 5.1.1 hold without assuming V_1, \dots, V_r satisfy Hörmander's condition, so long as one is willing to work on an injectively immersed submanifold. We describe this here—many of these methods appeared in [SS12] and are based on an idea of Lobry [Lob70].

Fix a large integer m to be chosen later and a compact set $\mathcal{K} \Subset \Omega$. Assign to each V_1, \dots, V_r the formal degree 1. If Z has formal degree e , we assign to $[V_j, Z]$ the formal degree $e + 1$. Let $(X_1, d_1), \dots, (X_q, d_q)$ denote the finite list of vector fields with formal degree $d_j \leq m$. The results that follow are essentially independent of m , provided m is chosen sufficiently large; how large m needs to be depends on V_1, \dots, V_r and \mathcal{K} . As above, for $\delta \in (0, 1]$, we let $\delta^d X$ denote the list of vector fields $\delta^{d_1} X_1, \dots, \delta^{d_q} X_q$. We sometimes identify $\delta^d X$ with the $n \times q$ matrix $(\delta^{d_1} X_1 | \dots | \delta^{d_q} X_q)$. Set $B_{(X,d)}(x, \delta) := B_{\delta^d X}(x, 1)$, where the later ball is defined in (2.2).

The classical Frobenius theorem applies to the involutive distribution generated by V_1, \dots, V_r (see [Her63, Nag66, Lob70, Sus73]) to see that the ambient space is foliated into real analytic leaves⁹. Let L_x denote the leaf passing through x . V_1, \dots, V_r satisfy Hörmander's condition on each leaf. If m is sufficiently large and Ω' is an open set with $\mathcal{K} \Subset \Omega' \Subset \Omega$, then V_1, \dots, V_r satisfy Hörmander's condition of order at most m on $L_x \cap \Omega'$, $\forall x \in \mathcal{K}$. Therefore X_1, \dots, X_q span the tangent space at every point of $L_x \cap \Omega'$, $\forall x \in \mathcal{K}$, and $B_{(X,d)}(x, \delta)$ is an open subset of L_x . Let ν_x denote the induced Lebesgue density on the leaf passing through x . For an $n \times q$ matrix A , and for $n_0 \leq \min\{n, q\}$, let $\det_{n_0 \times n_0} A$ denote the vector consisting of the determinants of the $n_0 \times n_0$ submatrices of A —the order of the components does not matter.

⁹The various leaves may have different dimensions; i.e., this may be a singular foliation. See Remark 5.6 for further comments.

For each $x \in \Omega$ set $n_0(x) := \dim \text{span}\{X_1(x), \dots, X_q(x)\} = \dim L_x$. For each $x \in \Omega$, $\delta \in (0, 1]$, pick $j_1 = j_1(x, \delta), \dots, j_{n_0(x)} = j_{n_0(x)}(x, \delta)$ so that

$$\left| \det_{n_0(x) \times n_0(x)} \left(\delta^{d_{j_1}} X_{j_1}(x) | \dots | \delta^{d_{j_{n_0(x)}}} X_{j_{n_0(x)}}(x) \right) \right|_{\infty} = \left| \det_{n_0(x) \times n_0(x)} \delta^d X \right|_{\infty}.$$

For this choice of $j_1 = j_1(x, \delta), \dots, j_{n_0(x)} = j_{n_0(x)}(x, \delta)$ set (writing n_0 for $n_0(x)$):

$$\Phi_{x,\delta}(t_1, \dots, t_{n_0}) := \exp\left(t_1 \delta^{d_{j_1}} X_{j_1} + \dots + t_{n_0} \delta^{d_{j_{n_0}}} X_{j_{n_0}}\right)x. \quad (5.4)$$

Theorem 5.4. *Fix a compact set $\mathcal{K} \subseteq \Omega$ and $x \in \mathcal{K}$, take m sufficiently large (depending on \mathcal{K} and V_1, \dots, V_r), and define $(X_1, d_1), \dots, (X_q, d_q)$ as above. Define $n_0(x)$, ν_x , and $\Phi_{x,\delta}(t_1, \dots, t_{n_0(x)})$ as above. We write $A \lesssim B$ for $A \leq CB$ where C is a positive constant which may depend on \mathcal{K} , but does not depend on the particular points $x \in \mathcal{K}$ and $u \in \mathbb{R}^{n_0(x)}$ under consideration, or on the scale $\delta \in (0, 1]$; we write $A \approx B$ for $A \lesssim B$ and $B \lesssim A$. There exists $\eta_1, \xi_0 \approx 1$ such that $\forall x \in \mathcal{K}$,*

(i) $\nu_x(B_{(X,d)}(x, \delta)) \approx \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_{\infty}, \forall \delta \in (0, \xi_0]$.

(ii) $\nu_x(B_{(X,d)}(x, 2\delta)) \lesssim \nu_x(B_{(X,d)}(x, \delta)), \forall \delta \in (0, \xi_0/2]$.

(iii) $\forall \delta \in (0, 1]$, $\Phi_{x,\delta}(B^{n_0(x)}(\eta_1)) \subseteq L_x$ is open and $\Phi_{x,\delta} : B^n(\eta_1) \rightarrow \Phi_{x,\delta}(B^n(\eta_1))$ is a real analytic diffeomorphism.

(iv) For $\delta \in (0, 1]$, define $h_{x,\delta}(t)$ on $B^{n_0(x)}(\eta_1)$ by $h_{x,\delta} \sigma_{\text{Leb}} = \Phi_{x,\delta}^* \nu_x$. Then, $h_{x,\delta}(t) \approx \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_{\infty}, \forall t \in B^{n_0(x)}(\eta_1)$, and there exists $s \approx 1$ with $\|h_{x,\delta}\|_{\mathcal{A}^{n_0(x),s}} \lesssim \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_{\infty}$.

(v) $B_{(X,d)}(x, \xi_0 \delta) \subseteq \Phi_{x,\delta}(B^{n_0(x)}(\eta_1)) \subseteq B_{(X,d)}(x, \delta), \forall \delta \in (0, 1]$.

(vi) For $\delta \in (0, 1]$, $x \in \mathcal{K}$, let $Y_j^{x,\delta} := \Phi_{x,\delta}^* \delta^{d_j} X_j$, so that $Y_j^{x,\delta}$ is a real analytic vector field on $B^{n_0(x)}(\eta_1)$. We have

$$\|Y_j^{x,\delta}\|_{\mathcal{A}^{n_0(x),\eta_1}(\mathbb{R}^n)} \lesssim 1, \quad 1 \leq j \leq q.$$

Finally, $Y_1^{x,\delta}(u), \dots, Y_q^{x,\delta}(u)$ span $T_u B^{n_0(x)}(\eta_1)$, uniformly in x, δ , and u , in the sense that

$$\max_{k_1, \dots, k_{n_0(x)} \in \{1, \dots, q\}} \inf_{u \in B^{n_0(x)}(\eta_1)} \left| \det \left(Y_{k_1}^{x,\delta}(u) | \dots | Y_{k_{n_0(x)}}^{x,\delta}(u) \right) \right| \approx 1.$$

Remark 5.5. See Section 9.1 for a generalization of Theorem 5.4 to the ‘‘multi-parameter’’ setting.

For the proof of Theorem 5.4, see Section 9.

Remark 5.6. Theorem 5.4 is useful even when restricting attention to $\delta = 1$. Indeed, the leaves L_x are real analytic manifolds, and account for the foliation of Ω associated to the real analytic vector fields V_1, \dots, V_r . This may be a singular foliation: $\dim L_x$ may not be constant in x . Suppose $x_0 \in \mathcal{K}$ is a singular point; i.e., $\dim L_x$ is not constant on any neighborhood of x_0 . Then, the usual constructions of the real analytic coordinate systems on L_x ‘‘blow up’’ as x approaches x_0 ; one does not obtain a useful quantitative control of these charts. One can think of the map $\Phi_{x,1}$ from Theorem 5.4 as a real analytic coordinate system near the point x . The conclusions of Theorem 5.4 amount to a quantitative control of this chart, which is uniform in x . A similar sort of uniform control was an important ingredient in [SS12], and will likely be useful in other questions from analysis regarding real analytic vector fields.

5.2 Generalized Sub-Riemannian Geometries

The results described in Section 5.1 concern the classical setting of sub-Riemannian geometry; which arises in many questions, including in the study of “maximally hypoelliptic” differential equations (see [Str14, Chapter 2] for details). In [SS18, Section 7.3], this setting was generalized to account for certain situations which arise for some partial differential equations which are not maximally hypoelliptic. With the results of this paper in hand, the results from [SS18, Section 7.3] transfer to the real analytic setting. We present these results here (with a few slight modifications from the setting in [SS18, Section 7.3]). One important thing to note is that, in this section (and unlike the settings described above) we do not require that the vector fields be a priori real analytic. We only require them to be C^1 , along with certain estimates which allow us to construct a coordinate system in which they are real analytic.

Let M be a connected n dimensional C^2 manifold and for each $\delta \in (0, 1]$, let $X^\delta = X_1^\delta, \dots, X_q^\delta$ be a list of C^1 vector fields on Ω which span the tangent space at every point. For $x \in \Omega$, $\delta \in (0, 1]$ set $B(x, \delta) := B_{X^\delta}(x, 1)$, where $B_{X^\delta}(x, 1)$ is defined by (2.2). Let ν be a C^1 density on M . Our goal is to give conditions under which the balls $B(x, \delta)$ when paired with the density ν give a real analytic space of homogeneous type. Some of the conditions we give can be thought of as analogs of the axioms of a space of homogeneous type, while others can be thought of as endowing this space of homogeneous type with an adapted real analytic structure. In what follows, we write X^δ as the column vector of vector fields $[X_1^\delta, \dots, X_q^\delta]^\top$. Because of this, if we are given a matrix $A : M \rightarrow \mathbb{M}^{q \times q}$, it makes sense to consider $A(x)X^\delta$ which again gives a column vector of vector fields. It also makes sense to consider the space $L^\infty(M)$, which can be defined with any strictly positive density on M —all such densities give the same space and norm.

We assume the following, which are a real analytic version of the assumptions in [SS18, Section 7.3]:

- (I) $\forall \delta \in (0, 1]$, $x \in M$, we have $\text{span}\{X_1^\delta(x), \dots, X_q^\delta(x)\} = T_x M$.
- (II) $X_1^\delta, \dots, X_q^\delta$ are uniformly C^1 in the following sense. For every $x \in M$, there exists an open set $U \subseteq \mathbb{R}^n$ and a diffeomorphism $\Psi : U \rightarrow V$, where V is a neighborhood of x in M , such that

$$\sup_{\delta \in (0, 1]} \|\Psi^* X_j^\delta\|_{C^1(U; \mathbb{R}^n)} < \infty.$$

- (III) $X_j^\delta \rightarrow 0$ and $\delta \rightarrow 0$, uniformly on compact subsets of M . More precisely, for every $x \in M$, there exists an open set $U \subseteq \mathbb{R}^n$ and a diffeomorphism $\Psi : U \rightarrow V$, where V is a neighborhood of x in M , such that

$$\lim_{\delta \rightarrow 0} \|\Psi^* X_j^\delta\|_{C^0(U; \mathbb{R}^n)} = 0.$$

- (IV) $\forall 0 < \delta_1 \leq \delta_2 \leq 1$, $X^{\delta_1} = T_{\delta_1, \delta_2} X^{\delta_2}$, where $T_{\delta_1, \delta_2} \in L^\infty(M; \mathbb{M}^{q \times q})$ with $\|T_{\delta_1, \delta_2}\|_{L^\infty(M; \mathbb{M}^{q \times q})} \leq 1$.

- (V) $\exists B_1, B_2 \in (1, \infty)$, $b_1, b_2 \in (0, 1)$ such that $\forall \delta \in (0, 1/B_1]$, $\exists S_\delta \in L^\infty(M; \mathbb{M}^{q \times q})$ and $\forall \delta \in (0, 1/B_2]$, $\exists R_\delta \in L^\infty(M; \mathbb{M}^{q \times q})$ with $S_\delta X^{B_1 \delta} = X^\delta$, $R_\delta X^\delta = X^{B_2 \delta}$, and

$$\sup_{0 < \delta \leq 1/B_1} \|S_\delta\|_{L^\infty(M; \mathbb{M}^{q \times q})} \leq b_1, \quad \sup_{0 < \delta \leq 1/B_2} \|R_\delta\|_{L^\infty(M; \mathbb{M}^{q \times q})} \leq b_2^{-1}.$$

- (VI) For every compact set $\mathcal{K} \Subset M$, there exists $\xi \in (0, 1]$ and $s > 0$ such that $\forall \delta \in (0, 1]$ and $x \in \mathcal{K}$, $[X_j^\delta, X_k^\delta] = \sum_{l=1}^q c_{j,k}^{l,x,\delta} X_l^\delta$ on $B_{X^\delta}(x, \xi)$, where

$$\sup_{x \in \mathcal{K}, \delta \in (0, 1]} \left\| c_{j,k}^{l,x,\delta} \right\|_{C_X^{\omega,s}(B_{X^\delta}(x, \xi))} < \infty.$$

- (VII) We assume $\mathcal{L}_{X_j^\delta} \nu = f_j^\delta \nu$, for $1 \leq j \leq q$, $\delta \in (0, 1]$, where for every relatively compact open set $\Omega' \Subset M$, there exists $s > 0$ with

$$\sup_{\delta \in (0, 1]} \|f_j^\delta\|_{C_X^{\omega,s}(\Omega')} < \infty.$$

Fix $\zeta \in (0, 1]$ (we take $\zeta = 1$ for many applications). For each $x \in M$, $\delta \in (0, 1]$, pick $j_1 = j_1(x, \delta), \dots, j_n = j_n(x, \delta) \in \{1, \dots, q\}$ so that

$$\max_{k_1, \dots, k_n \in \{1, \dots, q\}} \frac{X_{k_1}^\delta(x) \wedge \dots \wedge X_{k_n}^\delta(x)}{X_{j_1}^\delta(x) \wedge \dots \wedge X_{j_n}^\delta(x)} \leq \zeta^{-1}.$$

For this choice of $j_1 = j_1(x, \delta), \dots, j_n = j_n(x, \delta)$, define

$$\Phi_{x, \delta}(t_1, \dots, t_n) := \exp(t_1 X_{j_1}^\delta + \dots + t_n X_{j_n}^\delta)x.$$

Define, for $x \in M$, $\delta \in (0, 1]$

$$\Lambda(x, \delta) := \max_{k_1, \dots, k_n \in \{1, \dots, q\}} |\nu(X_{k_1}, \dots, X_{k_n})(x)|.$$

Theorem 5.7. (i) $B(x, \delta_1) \subseteq B(x, \delta_2)$, $\forall x \in M$, $0 < \delta_1 \leq \delta_2 \leq 1$.

(ii) $\bigcap_{\delta \in (0, 1]} \overline{B(x, \delta)} = \{x\}$, $\forall x \in M$.

(iii) $B(x, \delta) \cap B(y, \delta) \neq \emptyset \Rightarrow B(y, \delta) \subseteq B(x, C\delta)$, $\forall \delta \in (0, 1/C]$, where $C = B_1^k$ and k is chosen so that $b_1^k \leq \frac{1}{3}$.

Fix a compact set $\mathcal{K} \Subset M$. In what follows, we write $A \lesssim B$ for $A \subseteq CB$ where C is a positive constant which may depend on \mathcal{K} , but does not depend on the particular point $x \in \mathcal{K}$ or on the scale $\delta \in (0, 1]$. We write $A \approx B$ for $A \lesssim B$ and $B \lesssim A$. There exist $\eta_1, \xi_0 \approx 1$ such that $\forall x \in \mathcal{K}$:

(iv) ν is either everywhere strictly positive, everywhere strictly negative, or everywhere 0.

(v) $\nu(B(x, \delta)) \approx \nu(X_{j_1(x, \delta)}^\delta, \dots, X_{j_n(x, \delta)}^\delta)(x)$ and $|\nu(B(x, \delta))| \approx \Lambda(x, \delta)$, $\forall \delta \in (0, \xi_0]$.

(vi) $|\nu(B(x, 2\delta))| \lesssim |\nu(B(x, \delta))|$, $\forall \delta \in (0, \xi_0/2]$.

(vii) $\forall \delta \in (0, 1]$, $\Phi_{x, \delta}(B^n(\eta_1)) \subseteq M$ is open and $\Phi_{x, \delta} : B^n(\eta_1) \rightarrow \Phi_{x, \delta}(B^n(\eta_1))$ is a C^2 diffeomorphism.

(viii) Define $h_{x, \delta} \in C(B^n(\eta_1))$ by $h_{x, \delta} \sigma_{\text{Leb}} = \Phi_{x, \delta}^* \nu$. Then,

$$h_{x, \delta}(t) \approx \nu(X_{j_1(x, \delta)}^\delta, \dots, X_{j_n(x, \delta)}^\delta)(x) \text{ and } |h_{x, \delta}(t)| \approx \Lambda(x, \delta), \quad \forall t \in B^n(\eta_1).$$

Also, $h_{x, \delta} \in \mathcal{A}^{n, \eta_1}$ and $\|h_{x, \delta}\|_{\mathcal{A}^{n, \eta_1}} \lesssim \Lambda(x, \delta)$.

(ix) $B(x, \xi_0 \delta) \subseteq \Phi_{x, \delta}(B^n(\eta_1)) \subseteq B(x, \delta)$, $\forall \delta \in (0, 1]$.

(x) Let $Y_j^{x, \delta} := \Phi_{x, \delta}^* X_j^\delta$, $1 \leq j \leq q$, so that $Y_j^{x, \delta}$ is a vector field on $B^n(\eta_1)$. Then, $Y_j^{x, \delta} \in \mathcal{A}^{n, \eta_1}$ and

$$\|Y_j^{x, \delta}\|_{\mathcal{A}^{n, \eta_1}(\mathbb{R}^n)} \lesssim 1, \quad 1 \leq j \leq q.$$

Furthermore, $Y_1^{x, \delta}(u), \dots, Y_q^{x, \delta}(u)$ span $T_u B^n(\eta_1)$, uniformly in x, δ , and u in the sense that

$$\max_{k_1, \dots, k_n \in \{1, \dots, q\}} \inf_{u \in B^n(\eta_1)} \left| \det \left(Y_{k_1}^{x, \delta}(u) \mid \dots \mid Y_{k_n}^{x, \delta}(u) \right) \right| \approx 1.$$

Proof. (i), (ii), and (iii) all follow just as in the corresponding results in [SS18, Theorem 7.6]. For the remaining parts, the goal is to apply Theorem 3.10, Theorem 3.14, and Corollary 3.15 to the vector fields $X_1^\delta, \dots, X_q^\delta$, for $\delta \in (0, 1]$ and for each base point $x_0 \in \mathcal{K}$ (uniformly for $\delta \in (0, 1]$, $x_0 \in \mathcal{K}$)—we use the choice of ξ and ν from above, and take $\zeta = 1$.

By the Picard–Lindelöf theorem, we make take $\eta \in (0, 1]$ depending on \mathcal{K} and the bounds from (II), so that $\forall x \in \mathcal{K}$, $\delta \in (0, 1]$, $X_1^\delta, \dots, X_q^\delta$ satisfy $\mathcal{C}(x, \eta, M)$. Take $\delta_0 > 0$ as in Lemma 7.22, when applied to

$X_1^\delta, \dots, X_q^\delta$. It can be seen from the proof of Lemma 7.22 (which can be found in [SS18, Proposition 4.14]) that δ_0 can be chosen independent of $\delta \in (0, 1]$ (this uses (II)). In light of (VI) (see, also, Remark 3.7), Theorem 3.10, Theorem 3.14, and Corollary 3.15 apply to the vector fields $X_1^\delta, \dots, X_q^\delta$ for each $\delta \in (0, 1]$ and each base point $x \in \mathcal{K}$ (with η replaced by $\min\{\eta, s\}$). Each constant which is 0-admissible, admissible, ν -admissible, or $0; \nu$ -admissible in these results can be chosen independent of $x \in \mathcal{K}$ and $\delta \in (0, 1]$. Let $\xi_1, \xi_2, \eta_1 > 0$ be as in Theorem 3.10, so that ξ_1, ξ_2 , and η_1 can be chosen independent of $x \in \mathcal{K}$ and $\delta \in (0, 1]$. The map $\Phi_{x, \delta}$ is precisely the map Φ from Theorem 3.10 when using the base point x and the vector fields $X_1^\delta, \dots, X_q^\delta$.

(vii) follows from Theorem 3.10 (d) and (e). (viii) follows from Theorem 3.14 and Corollary 3.15. (viii) implies that on a neighborhood of each point, ν is either strictly positive, strictly negative, or identically 0. Since M is connected, it follows that ν is either everywhere strictly positive, everywhere strictly negative, or everywhere 0; i.e., (iv) holds. By multiplying ν by ± 1 , we may henceforth assume (without loss of generality) that ν is everywhere non-negative—and is either identically 0 or everywhere strictly positive.

(ix): Theorem 3.10 gives $\xi_2 \approx 1$ ($\xi_2 < 1$) such that

$$B_{X^\delta}(x, \xi_2) \subseteq \Phi_{x, \delta}(B^n(\eta_1)) \subseteq B_{X^\delta}(x, \xi) \subseteq B_{X^\delta}(x, 1) = B(x, \delta).$$

Thus, to prove (ix), we wish to show $\exists \xi_0 \approx 1$ with

$$B(x, \xi_0 \delta) \subseteq B_{X^\delta}(x, \xi_2). \quad (5.5)$$

This follows just as in [SS18, Theorem 7.6], where it is shown that we may take $\xi_0 = B_1^{-k}$, where k is chosen so that $b_1^k \leq \xi_2$.

We claim, for $\delta_1 \leq \delta_2 \leq 1$,

$$\Lambda(x, \delta_1) \lesssim \Lambda(x, \delta_2), \quad (5.6)$$

where the implicit constant can be chosen to depend only on q . Indeed,

$$\begin{aligned} \Lambda(x, \delta_1) &= \max_{k_1, \dots, k_n \in \{1, \dots, q\}} \left| \nu(X_{k_1}^{\delta_1}, \dots, X_{k_n}^{\delta_1})(x) \right| \\ &= \max_{k_1, \dots, k_n \in \{1, \dots, q\}} \left| \nu((T_{\delta_1, \delta_2} X^{\delta_2})_{k_1}, \dots, (T_{\delta_1, \delta_2} X^{\delta_2})_{k_n})(x) \right|. \end{aligned}$$

Since $\|T_{\delta_1, \delta_2}\|_{L^\infty(M; \mathbb{M}^{q \times q})} \leq 1$, the right hand side is ν evaluated at a linear combination, with (variable) coefficients bounded by 1, of the vector fields $X_1^{\delta_2}, \dots, X_q^{\delta_2}$. Using the properties of densities, it follows that

$$\left| \nu((T_{\delta_1, \delta_2} X^{\delta_2})_{k_1}, \dots, (T_{\delta_1, \delta_2} X^{\delta_2})_{k_n})(x) \right| \lesssim \Lambda(x, \delta_2),$$

(5.6) follows.

Next we claim, for $c > 0$ fixed,

$$\Lambda(x, c\delta) \approx \Lambda(x, \delta), \quad \delta, c\delta \in (0, 1], \quad (5.7)$$

where the implicit constant depends on c , but not on x or δ . It suffices to prove (5.7) for $c < 1$. By (5.6), it suffices to prove (5.7) for $c = B_2^{-k}$ for some k . We have

$$\begin{aligned} \Lambda(x, \delta) &= \max_{k_1, \dots, k_n \in \{1, \dots, q\}} \left| \nu(X_{k_1}^\delta, \dots, X_{k_n}^\delta)(x) \right| \\ &= \max_{k_1, \dots, k_n \in \{1, \dots, q\}} \left| \nu((AX^{c\delta})_{k_1}, \dots, (AX^{c\delta})_{k_n})(x) \right|, \end{aligned} \quad (5.8)$$

where $A(x) = R_{B_2^{-1}\delta}(x)R_{B_2^{-2}\delta}(x) \cdots R_{B_2^{-k}\delta}(x)$. Since $\sup_{x \in M} \|A(x)\|_{\mathbb{M}^{q \times q}} \leq b_2^{-k} \lesssim 1$ (where the implicit constant depends on k), it follows that the right hand side of (5.8) is ν evaluated at linear combinations, with (variable) coefficients which have absolute value $\lesssim 1$, of the vectors $X_1^{c\delta}, \dots, X_q^{c\delta}$. It follows from the properties of densities that

$$\left| \nu((AX^{c\delta})_{k_1}, \dots, (AX^{c\delta})_{k_n})(x) \right| \lesssim \Lambda(x, c\delta).$$

We conclude $\Lambda(x, \delta) \lesssim \Lambda(x, c\delta)$. Combining this with (5.6) proves (5.7).

Corollary 3.15 and using that we have (without loss of generality) assumed ν is non-negative, shows

$$\nu(B_{X^\delta}(x, \xi_2)) \approx \Lambda(x, \delta). \quad (5.9)$$

Combining (5.9) with (5.7) and (5.5) shows

$$\nu(B(x, \xi_0\delta)) \leq \nu(B_{X^\delta}(x, \xi_2)) \approx \Lambda(x, \delta) \approx \Lambda(x, \xi_0\delta). \quad (5.10)$$

Conversely, using (5.9) again, we have

$$\Lambda(x, \delta) \approx \nu(B_{X^\delta}(x, \xi_2)) \leq \nu(B_{X^\delta}(x, 1)) = \nu(B(x, \delta)). \quad (5.11)$$

Combining (5.10) and (5.11) proves $|\nu(B(x, \delta))| \approx \Lambda(x, \delta)$, $\forall \delta \in (0, \xi_0]$. Since we have assumed ν is everywhere non-negative, (v) follows from this and Corollary 3.15. (vi) follows from (v) and (5.7). (x) follows from Theorem 3.10. \square

6 Results from the first paper

In this section, we describe the main results needed from [SS18]. We do not require as detailed information as is discussed in that paper, and so we instead state an immediate consequence of the results in that paper.

We take the same setup and hypotheses as in Theorem 3.10, and define 0-admissible constants and admissible constants as in that theorem. We take $\Phi(t)$ as in (3.3). We separate the results we need into two parts.

Proposition 6.1. *There exists a 0-admissible constant $\chi \in (0, \xi]$ such that:*

(a) $\forall y \in B_{X_{J_0}}(x_0, \chi)$, $\bigwedge X_{J_0}(y) \neq 0$.

(b) $\forall y \in B_{X_{J_0}}(x_0, \chi)$,

$$\sup_{J \in \mathcal{I}(n, q)} \left| \frac{\bigwedge X_J(y)}{\bigwedge X_{J_0}(y)} \right| \approx_0 1.$$

(c) $\forall \chi' \in (0, \chi]$, $B_{X_{J_0}}(x_0, \chi')$ is an open subset of $B_X(x_0, \xi)$ and is therefore a submanifold.

Proof. This is contained in [SS18, Theorem 4.7]. \square

Proposition 6.2. *In the special case $n = q$ (so that $X_{J_0} = X$), there exists an admissible constant $\hat{\eta} \in (0, \eta_0]$ such that:*

(a) $\Phi(B^n(\hat{\eta}))$ is an open subset of $B_X(x_0, \xi)$, and is therefore a submanifold.

(b) $\Phi : B^n(\hat{\eta}) \rightarrow \Phi(B^n(\hat{\eta}))$ is a C^2 diffeomorphism.

Let $Y_j = \Phi^* X_j$ and write $Y = (I + A)\nabla$, where Y denotes the column vector of vector fields $Y = [Y_1, \dots, Y_n]^\top$, ∇ denotes the gradient in \mathbb{R}^n thought of as a column vector, and $A \in C(B^n(\hat{\eta}); \mathbb{M}^{n \times n})$.

(c) $A(0) = 0$.

For $t \in B^n(\hat{\eta})$, let $C(t)$ denote the $n \times n$ matrix with j, k component given by $\sum_{i=1}^n t_i c_{j,i}^k(\Phi(t))$. For $t \in B^n(\hat{\eta})$ write t in polar coordinates $t = r\theta$.

(d) A satisfies the differential equation

$$\frac{\partial}{\partial r} rA(r\theta) = -A(r\theta)^2 - C(r\theta)A(r\theta) - C(r\theta). \quad (6.1)$$

Proof. All of the results except (d) are contained in [SS18, Proposition 9.26]. [SS18, Proposition 9.26] uses the notion of a $1'$ -admissible constant, which involves bounds on $\|c_{j,k}^l \circ \Phi\|_{C^1(B^n(\eta_0))}$. Our assumptions imply $\|c_{j,k}^l \circ \Phi\|_{\mathcal{A}^{n,\eta_0}} \lesssim 1$; this does not quite imply $c_{j,k}^l \circ \Phi \in C^1(B^n(\eta_0))$ (the problem is that while $c_{j,k}^l \circ \Phi$ is C^1 , its first derivatives might not be bounded on $B^n(\eta_0)$). Instead, we proceed as follows. By defining $\tilde{\eta} := \eta_0/2$, we do have $\|c_{j,k}^l \circ \Phi\|_{C^1(B^n(\tilde{\eta}))} \lesssim \|c_{j,k}^l \circ \Phi\|_{\mathcal{A}^{n,\eta_0}} \lesssim 1$. Applying [SS18, Proposition 9.26] with η replaced by $\tilde{\eta}$ yields all of the above except (d).

(d) is an immediate consequence of [SS18, Proposition 9.1]. \square

7 Proofs

In this section, we prove the main results of this paper; namely, Theorems 3.3, 3.4, 3.10, and 3.14. In Section 7.1 we describe the main way we show functions are real analytic. Namely, we prove the function in question satisfies an appropriate real analytic ODE, which forces it to be real analytic; Section 7.1 contains several quantitative instances of this. In Section 7.2 we prove the main quantitative theorem: Theorem 3.10. In Section 7.3 we study densities and prove Theorem 3.14. Finally, in Section 7.4, we prove the qualitative results (Theorems 3.3 and 3.4), which are simple consequences of Theorem 3.10.

7.1 Real Analytic Functions and ODEs

At various points, we will need to prove functions are real analytic. The way we will do this is by showing the functions satisfy a ODE which depends real analytically on the appropriate parameters. We begin with a simple and classical version of this where we deduce the solutions to a certain real analytic PDE are real analytic, by reducing it to a real analytic ODE.

Proposition 7.1. *Let $\Omega \subseteq \mathbb{R}^n$ be an open set and $f \in C^1(\Omega; \mathbb{R}^m)$ satisfy*

$$\frac{\partial}{\partial t_j} f(t) = F_j(t, f(t)), \quad \forall t \in \Omega, 1 \leq j \leq n,$$

where F_j is real analytic in both variables. Then, f is real analytic.

Proof Sketch. Fix $s \in \Omega$. We will show f is real analytic near s . Set $g(\epsilon, t) := f(\epsilon t + s)$. Then, we have

$$\frac{\partial}{\partial \epsilon} g(\epsilon, t) = \sum_{j=1}^n t_j \frac{\partial f}{\partial t_j}(\epsilon t + s) = \sum_{j=1}^n t_j F_j(\epsilon t + s, f(\epsilon t + s)) =: G_s(\epsilon, t, g(\epsilon, t)),$$

where G_s is analytic in all its variables. Also, $g(0, t) = f(s)$ (which is constant in t).

Hence, $g(\epsilon, t)$ satisfies a real analytic ODE, and classical results show g is real analytic for ϵ and t small. Since $g(\epsilon, t) = f(\epsilon t + s)$, this shows that f is real analytic near s , completing the proof. \square

7.1.1 A Particular ODE

Fix $D, \hat{\eta} > 0$ and for $1 \leq j \leq n$, let $C_j \in \mathcal{A}^{n, \hat{\eta}}(\mathbb{M}^{n \times n})$ with $\sum_{j=1}^n \|C_j\|_{\mathcal{A}^{n, \hat{\eta}}(\mathbb{M}^{n \times n})} \leq D$. Set $C(t) := \sum_{j=1}^n t_j C_j(t)$. For $t \in B^n(\hat{\eta})$, we write t in polar coordinates $t = r\theta$. We consider the differential equation, defined for functions $A(t)$ taking values in $\mathbb{M}^{n \times n}$, given by

$$\frac{\partial}{\partial r} r A(r\theta) = -A(r\theta)^2 - C(r\theta)A(r\theta) - C(r\theta), \quad A(0) = 0. \quad (7.1)$$

Proposition 7.2. *Let $\eta_1 \in (0, \min\{\hat{\eta}, 5/8D\}]$. There exists a solution $A \in \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ to (7.1). Moreover, this solution satisfies $\|A\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \frac{1}{2}$. Finally, this solution is unique in the sense that if $B(t) \in C(B^n(\delta); \mathbb{M}^{n \times n})$ is another solution to (7.1), then $A(t) = B(t)$ for $|t| < \min\{\delta, \eta_1\}$.*

The rest of this section is devoted to the proof of Proposition 7.2. Following the proof in [SS18, Proposition 9.4], we introduce the map $\mathcal{T} : \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n}) \rightarrow \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ given by

$$\mathcal{T}(A)(x) := - \int_0^1 A(sx)^2 + C(sx)A(sx) + C(sx) ds.$$

Using that $\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ is an algebra (Lemma 4.1) it is immediate to verify $\mathcal{T} : \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n}) \rightarrow \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$. A simple change of variables shows, for $r > 0$,

$$\mathcal{T}(A)(r\theta) = \frac{1}{r} \int_0^r -A(s\theta)^2 - C(s\theta)A(s\theta) - C(s\theta) ds.$$

Thus, A is a solution to (7.1) if and only if $\mathcal{T}(A) = A$ and $A(0) = 0$. We will prove the existence of such a fixed point by using the contraction mapping principle.

Lemma 7.3. *Let $A \in \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ satisfy $A(0) = 0$. For $s \in [0, 1]$ set $A_s(x) = A(sx)$. Then, $\|A_s\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq s\|A\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})}$.*

Proof. This is immediate from the definitions. □

Lemma 7.4. *For $s \in [0, 1]$, set $C_s(t) = C(st)$. Then, $\|C_s\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \eta_1 s D$.*

Proof. Let $C_j(t) = \sum_{\alpha \in \mathbb{N}^n} \frac{c_{\alpha, j}}{\alpha!} t^\alpha$. Then, $C_s(t) = \sum_{\alpha \in \mathbb{N}^n} \sum_{j=1}^n \frac{c_{\alpha, j}}{\alpha!} s^{|\alpha|+1} t_j^\alpha$. Thus,

$$\begin{aligned} \|C_s\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} &\leq \sum_{j=1}^n \sum_{\alpha \in \mathbb{N}^n} \frac{\|c_{\alpha, j}\|_{\mathbb{M}^{n \times n}}}{\alpha!} s^{|\alpha|+1} \eta_1^{|\alpha|+1} \leq s \eta_1 \sum_{j=1}^n \sum_{\alpha \in \mathbb{N}^n} \frac{\|c_{\alpha, j}\|_{\mathbb{M}^{n \times n}}}{\alpha!} \eta_1^{|\alpha|} \\ &= s \eta_1 \sum_{j=1}^n \|C_j\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq s \eta_1 D, \end{aligned}$$

completing the proof. □

Define

$$\mathcal{M} := \left\{ A \in \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n}) : A(0) = 0 \text{ and } \|A\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \frac{1}{2} \right\}.$$

We give \mathcal{M} the induced metric as a subset of $\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$. With this metric, \mathcal{M} is a complete metric space.

Lemma 7.5. *For $\eta_1 \in (0, \min\{\hat{\eta}, 5/8D\}]$, $\mathcal{T} : \mathcal{M} \rightarrow \mathcal{M}$ and is a strict contraction.*

Proof. Using that $\|B_1 B_2\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \|B_1\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \|B_2\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})}$ (Lemma 4.1) and Lemmas 7.3 and 7.4, we have, for $A \in \mathcal{M}$,

$$\begin{aligned} \|\mathcal{T}(A)\|_{\mathcal{A}^{n, \eta_1}} &\leq \int_0^1 \|A(s\cdot)^2\|_{\mathcal{A}^{n, \eta_1}} + \|C(s\cdot)A(s\cdot)\|_{\mathcal{A}^{n, \eta_1}} + \|C(s\cdot)\|_{\mathcal{A}^{n, \eta_1}} ds \\ &\leq \int_0^1 s^2 \|A\|_{\mathcal{A}^{n, \eta_1}}^2 + (D\eta_1 s^2) \|A\|_{\mathcal{A}^{n, \eta_1}} + D\eta_1 s ds \leq \frac{1}{3} \cdot \frac{1}{4} + \frac{D\eta_1}{3} \cdot \frac{1}{2} + \frac{D\eta_1}{2} \\ &= \frac{1}{12} + \frac{2}{3} D\eta_1 \leq \frac{1}{12} + \frac{2}{3} \cdot \frac{5}{8} = \frac{1}{2}. \end{aligned}$$

Clearly, since $A(0) = 0$ and $C(0) = 0$, we have $\mathcal{T}(A)(0) = 0$. We conclude $\mathcal{T} : \mathcal{M} \rightarrow \mathcal{M}$.

For $A, B \in \mathcal{M}$, we have using $A^2 - B^2 = \frac{1}{2}(A+B)(A-B) + \frac{1}{2}(A-B)(A+B)$,

$$\begin{aligned} \|\mathcal{T}(A) - \mathcal{T}(B)\|_{\mathcal{A}^{n, \eta_1}} &\leq \int_0^1 \|A(s\cdot) + B(s\cdot)\|_{\mathcal{A}^{n, \eta_1}} \|A(s\cdot) - B(s\cdot)\|_{\mathcal{A}^{n, \eta_1}} + \|C(s\cdot)\|_{\mathcal{A}^{n, \eta_1}} \|A(s\cdot) - B(s\cdot)\|_{\mathcal{A}^{n, \eta_1}} ds \\ &\leq \int_0^1 s^2 \|A - B\|_{\mathcal{A}^{n, \eta_1}} + D\eta_1 s^2 \|A - B\|_{\mathcal{A}^{n, \eta_1}} ds \leq \frac{1 + D\eta_1}{3} \|A - B\|_{\mathcal{A}^{n, \eta_1}} \leq \frac{13}{24} \|A - B\|_{\mathcal{A}^{n, \eta_1}}, \end{aligned}$$

completing the proof. □

Proof of Proposition 7.2. Uniqueness for (7.1) was established in [SS18, Proposition 9.4]; and the same proof yields the claimed uniqueness in Proposition 7.2. For existence, Lemma 7.5 shows that the contraction mapping principle applies to $\mathcal{T} : \mathcal{M} \rightarrow \mathcal{M}$ to show that there is a unique fixed point, $A \in \mathcal{M}$, of \mathcal{T} . As described above, this A is a solution to (7.1) and clearly satisfies $\|A\|_{\mathcal{A}^{n,n_1}(\mathbb{M}^{n \times n})} \leq \frac{1}{2}$ (since $A \in \mathcal{M}$). This completes the proof. \square

7.1.2 Identifying Real Analytic Functions I: Euclidean Space

In Proposition 7.1, we showed how to prove a function was real analytic by introducing a new variable and proving the function satisfied an ODE in this new variable. In this section, we present a quantitative version of a similar argument. We make no effort to state the result in the greatest generality, and instead focus on the setting needed for this paper.

Fix $n, N, L \in \mathbb{N}$ and $r > 0$. We consider functions $F(t) = (F_1(t), \dots, F_N(t)) \in C(B^n(r); \mathbb{R}^N)$ satisfying a certain ODE. For each $1 \leq l \leq N$, $1 \leq j \leq n$, and $\alpha \in \mathbb{N}^N$ with $|\alpha| \leq L$ (α a multi-index), let $a_{\alpha,j,l} \in \mathcal{A}^{n,r}$. For $1 \leq l \leq N$, fix $F_{l,0} \in \mathbb{R}$. We consider the following system of differential equations for $1 \leq l \leq N$:

$$\frac{\partial}{\partial \epsilon} F_l(\epsilon t) = \sum_{j=1}^n \sum_{\substack{\alpha \in \mathbb{N}^N \\ |\alpha| \leq L}} t_j a_{\alpha,j,l}(\epsilon t) F(\epsilon t)^\alpha, \quad F_l(0) = F_{l,0}. \quad (7.2)$$

Fix D such that

$$|F_{l,0}|, \|a_{\alpha,j,l}\|_{\mathcal{A}^{n,r}} \leq D, \quad 1 \leq j \leq n, 1 \leq l \leq N, |\alpha| \leq L.$$

Proposition 7.6. *Set $r' := \min\{r, D(n2^L(L+1)^N(\max\{1, D\})^{L+1})^{-1}, (n(L+1)^{N+1}(\max\{1, D\})^{L2^L})^{-1}\}$. Suppose $F \in C(B^n(r); \mathbb{R}^N)$ satisfies (7.2). Then, $F|_{B^n(r')} \in \mathcal{A}^{n,r'}(\mathbb{R}^N)$. Moreover, for each $1 \leq l \leq N$, $\|F_l\|_{\mathcal{A}^{n,r'}} \leq 2D$.*

To prove Proposition 7.6 we prove a more general auxiliary result where we separate ϵt into two variables (ϵ, t) . To this end, set

$$r' := \min\{r, D(n2^L(L+1)^N(\max\{1, D\})^{L+1})^{-1}, (n(L+1)^{N+1}(\max\{1, D\})^{L2^L})^{-1}\}. \quad (7.3)$$

We will consider functions $\tilde{F}(\epsilon, t) : [0, 1] \times B^n(r') \rightarrow \mathbb{R}^N$, and we will think of these as functions $\tilde{F}(\epsilon, t) \in \mathcal{A}^{n,r'}(C([0, 1]; \mathbb{R}^N))$. I.e.,

$$\tilde{F}(\epsilon, t) = \sum_{\beta \in \mathbb{N}^n} \frac{t^\beta}{\beta!} c_\beta(\epsilon),$$

where $c_\beta \in C([0, 1]; \mathbb{R}^N)$ and

$$\|\tilde{F}\|_{\mathcal{A}^{n,r'}(C([0,1]; \mathbb{R}^N))} = \sum_{\beta \in \mathbb{N}^n} \frac{(r')^{|\beta|}}{\beta!} \|c_\beta\|_{C([0,1]; \mathbb{R}^N)}.$$

For $1 \leq j \leq n$, $1 \leq l \leq N$, and $\alpha \in \mathbb{N}^N$ with $|\alpha| \leq L$, let $\tilde{a}_{\alpha,j,l}(\epsilon, t) \in \mathcal{A}^{n,r}(C([0, 1]))$. For $1 \leq l \leq N$, fix a constant $\tilde{F}_{l,0} \in \mathbb{R}$. We consider the following system of differential equations for $1 \leq l \leq N$:

$$\frac{\partial}{\partial \epsilon} \tilde{F}_l(\epsilon, t) = \sum_{j=1}^n \sum_{\substack{\alpha \in \mathbb{N}^N \\ |\alpha| \leq L}} t_j \tilde{a}_{\alpha,j,l}(\epsilon, t) \tilde{F}(\epsilon, t)^\alpha, \quad \tilde{F}_l(0, t) \equiv \tilde{F}_{l,0}. \quad (7.4)$$

We suppose:

$$|\tilde{F}_{l,0}|, \|\tilde{a}_{\alpha,j,l}\|_{\mathcal{A}^{n,r}(C([0,1]))} \leq D, \quad 1 \leq j \leq n, 1 \leq l \leq N, |\alpha| \leq L.$$

Proposition 7.7. *Let r' be given by (7.3). There exists $\tilde{F}(\epsilon, t) \in \mathcal{A}^{n, r'}(C([0, 1]; \mathbb{R}^N))$ satisfying (7.4). This solution satisfies*

$$\max_{1 \leq l \leq N} \|\tilde{F}_l\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \leq 2D. \quad (7.5)$$

Finally, this solution is unique in the sense that if for some $\delta > 0$, $\hat{F}(\epsilon, t) \in C([0, 1] \times B^n(\delta); \mathbb{R}^N)$ is another solution to (7.4), then $\tilde{F}(\epsilon, t) = \hat{F}(\epsilon, t)$ for $|t| < \min\{\delta, r'\}$ and $\epsilon \in [0, 1]$.

The rest of this section is devoted to the proofs of Propositions 7.6 and 7.7. We begin with Proposition 7.7, the existence portion of which we will prove using the contraction mapping principle.

Lemma 7.8. *For $f(\epsilon, t) \in \mathcal{A}^{n, r'}(C([0, 1]))$, set $g(\epsilon, t) := \int_0^\epsilon f(\epsilon', t) d\epsilon'$. Then, $g \in \mathcal{A}^{n, r'}(C([0, 1]))$ and*

$$\|g\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \leq \|f\|_{\mathcal{A}^{n, r'}(C([0, 1]))}.$$

Proof. This is immediate from the definitions. \square

For $\tilde{F}(\epsilon, t) = (\tilde{F}_1(\epsilon, t), \dots, \tilde{F}_N(\epsilon, t)) \in \mathcal{A}^{n, r'}(C([0, 1]; \mathbb{R}^N))$ and $1 \leq l \leq N$, set

$$\mathcal{T}_l(\tilde{F})(\epsilon, t) = \tilde{F}_{l,0} + \int_0^\epsilon \left(\sum_{j=1}^n \sum_{\substack{\alpha \in \mathbb{N}^N \\ |\alpha| \leq L}} t_j \tilde{a}_{\alpha, j, l}(\epsilon', t) \tilde{F}(\epsilon', t)^\alpha \right) d\epsilon',$$

and set $\mathcal{T}(\tilde{F}) := (\mathcal{T}_1(\tilde{F}), \dots, \mathcal{T}_N(\tilde{F}))$. Using that $\mathcal{A}^{n, r'}(C([0, 1]))$ is an algebra (Lemma 4.1), $r' \leq r$, and Lemma 7.8, we have $\mathcal{T} : \mathcal{A}^{n, r'}(C([0, 1]; \mathbb{R}^N)) \rightarrow \mathcal{A}^{n, r'}(C([0, 1]; \mathbb{R}^N))$. Furthermore, \tilde{F} solves (7.4) if and only if $\mathcal{T}(\tilde{F}) = \tilde{F}$.

Set

$$\mathcal{M} := \left\{ \tilde{F} = (\tilde{F}_1, \dots, \tilde{F}_N) \in \mathcal{A}^{n, r'}(C([0, 1]; \mathbb{R}^N)) : \|\tilde{F}_l\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \leq 2D, 1 \leq l \leq N \right\}.$$

We give \mathcal{M} the metric $\rho(\tilde{F}, \hat{F}) := \max_{1 \leq l \leq N} \|\tilde{F}_l - \hat{F}_l\|_{\mathcal{A}^{n, r'}(C([0, 1]))}$. With this metric, \mathcal{M} is a complete metric space.

Lemma 7.9. $\mathcal{T} : \mathcal{M} \rightarrow \mathcal{M}$ and is a strict contraction.

Proof. For $\tilde{F} \in \mathcal{M}$, $|\alpha| \leq L$, $1 \leq j \leq n$, and $1 \leq l \leq N$, we have by Lemma 4.1,

$$\|t_j \tilde{a}_{\alpha, j, l} \tilde{F}^\alpha\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \leq \|t_j\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \|\tilde{a}_{\alpha, j, l}\|_{\mathcal{A}^{n, r'}(C([0, 1]))} (2D)^{|\alpha|} \leq r' 2^{|\alpha|} D^{|\alpha|+1} \leq r' 2^L (\max\{1, D\})^{L+1}.$$

Thus, by Lemma 7.8,

$$\|\mathcal{T}_l(\tilde{F})\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \leq |\tilde{F}_{l,0}| + \sum_{j=1}^n \sum_{|\alpha| \leq L} r' 2^L (\max\{1, D\})^{L+1} \leq D + r' n 2^L (L+1)^N (\max\{1, D\})^{L+1} \leq 2D,$$

where the last inequality follows from the choice of r' . It follows that $\mathcal{T} : \mathcal{M} \rightarrow \mathcal{M}$.

Again using Lemma 4.1, we have for $\tilde{F}, \hat{F} \in \mathcal{M}$, $|\alpha| \leq L$, $1 \leq j \leq n$, and $1 \leq l \leq N$,

$$\begin{aligned} \|t_j \tilde{a}_{\alpha, j, l} (\tilde{F}^\alpha - \hat{F}^\alpha)\|_{\mathcal{A}^{n, r'}(C([0, 1]))} &\leq \|t_j\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \|\tilde{a}_{\alpha, j, l}\|_{\mathcal{A}^{n, r'}(C([0, 1]))} |\alpha| (2D)^{|\alpha|-1} \max_{1 \leq k \leq N} \|\tilde{F}_k - \hat{F}_k\|_{\mathcal{A}^{n, r'}(C([0, 1]))} \\ &\leq r' 2^{|\alpha|-1} D^{|\alpha|} |\alpha| \rho(\tilde{F}, \hat{F}) \leq r' 2^{L-1} (\max\{1, D\})^L L \rho(\tilde{F}, \hat{F}). \end{aligned}$$

Thus, by Lemma 7.8,

$$\begin{aligned} \|\mathcal{T}_l(\tilde{F}) - \mathcal{T}_l(\hat{F})\|_{\mathcal{A}^{n, r'}(C([0, 1]))} &\leq \sum_{j=1}^n \sum_{|\alpha| \leq L} r' 2^{L-1} (\max\{1, D\})^L L \rho(\tilde{F}, \hat{F}) \\ &\leq n(L+1)^{N+1} r' 2^{L-1} (\max\{1, D\})^L \rho(\tilde{F}, \hat{F}) \leq \frac{1}{2} \rho(\tilde{F}, \hat{F}), \end{aligned}$$

where the last inequality follows from the choice of r' . It follows that $\rho(\mathcal{T}(\tilde{F}), \mathcal{T}(\hat{F})) \leq \frac{1}{2}\rho(\tilde{F}, \hat{F})$, and therefore $\mathcal{T} : \mathcal{M} \rightarrow \mathcal{M}$ is a strict contraction. \square

Proof of Proposition 7.7. Lemma 7.9 shows that the contraction mapping principle applies to $\mathcal{T} : \mathcal{M} \rightarrow \mathcal{M}$ to yield a unique fixed point $\tilde{F} \in \mathcal{M}$ of \mathcal{T} . This fixed point is the desired solution to (7.4). Since $\tilde{F} \in \mathcal{M}$, (7.5) follows. Finally, since (7.4) is a standard ODE, standard uniqueness theorems (using, for example, Grönwall's inequality) give the claimed uniqueness. \square

Proof of Proposition 7.6. Suppose $F \in C(B^n(r); \mathbb{R}^N)$ satisfies (7.2). Set $\tilde{F}(\epsilon, t) = F(\epsilon t)$, $\tilde{a}_{\alpha, j, l}(\epsilon, t) = a_{\alpha, j, l}(\epsilon t)$, and $\tilde{F}_{l, 0} := F_{l, 0}$. Then, \tilde{F} satisfies (7.4). The uniqueness from Proposition 7.7 shows that \tilde{F} is the solution described in that result, and therefore $\tilde{F}(\epsilon, t) \in \mathcal{A}^{n, r'}(C([0, 1]; \mathbb{R}^N))$ and (7.5) holds. Since $F(t) = \tilde{F}(1, t)$, the result follows. \square

7.1.3 Identifying Real Analytic Functions II: Manifolds

Let X_1, \dots, X_q be C^1 vector fields on a C^2 manifold \mathfrak{M} . Fix $x_0 \in \mathfrak{M}$ and suppose X_1, \dots, X_q satisfy $\mathcal{C}(x_0, \eta, \mathfrak{M})$ for some $\eta > 0$. Fix $N, L \in \mathbb{N}$, $\xi > 0$. For $1 \leq l \leq N$, $1 \leq j \leq q$, let $P_{l, j}$ be a polynomial of degree L , in N indeterminates, with coefficients in $\mathcal{A}_X^{x_0, \eta} \cap C(B_X(x_0, \xi))$:

$$P_{l, j}(x, y) = \sum_{\substack{\alpha \in \mathbb{N}^N \\ |\alpha| \leq L}} b_{\alpha, l, j}(x) y^\alpha,$$

where $b_{\alpha, l, j} \in \mathcal{A}_X^{x_0, \eta} \cap C(B_X(x_0, \xi))$. Fix $D > 0$ with $\|b_{\alpha, l, j}\|_{\mathcal{A}_X^{x_0, \eta}} \leq D$, $\forall \alpha, j, l$.

Proposition 7.10. *Suppose $G = (G_1, \dots, G_N) \in C(B_X(x_0, \xi); \mathbb{R}^N)$ satisfies, for $1 \leq j \leq q$, $1 \leq l \leq N$,*

$$X_j G_l(x) = P_{l, j}(x, G(x)).$$

Then, $\exists r' \in (0, \eta]$ with $G \in \mathcal{A}_X^{x_0, r'}(\mathbb{R}^N)$. Furthermore, $\|G\|_{\mathcal{A}_X^{x_0, r'}} \leq C$ where C and r' can be chosen to depend only on upper bounds for q , η^{-1} , ξ^{-1} , D , L , N , and $|G(x_0)|$.

Proof. Let $\Psi(t_1, \dots, t_q) := \exp(t_1 X_1 + \dots + t_q X_q)x_0$, and set $F(t) := G(\Psi(t))$. The goal is to show $F \in \mathcal{A}^{q, r'}(\mathbb{R}^N)$ with $\|F\|_{\mathcal{A}^{q, r'}(\mathbb{R}^N)} \leq C$, where r' and C are as in the statement of the result. Note that F satisfies

$$\frac{\partial}{\partial \epsilon} F_l(\epsilon t) = \sum_{j=1}^q t_j (X_j F_l)(\Psi(\epsilon t)) = \sum_{j=1}^q \sum_{\substack{\alpha \in \mathbb{N}^N \\ |\alpha| \leq L}} t_j b_{\alpha, l, j} \circ \Phi(\epsilon t) F(\epsilon t)^\alpha.$$

By hypothesis, $b_{\alpha, l, j} \circ \Psi \in \mathcal{A}^{q, \eta}$ with $\|b_{\alpha, l, j} \circ \Psi\|_{\mathcal{A}^{q, \eta}} \leq D$. From here, the result follows immediately from Proposition 7.6 (with $n = q$ and $r = \min\{\eta, \xi\}$). \square

7.2 The Quantitative Theorem

In this section, we prove Theorem 3.10 which is the main theorem of this paper. We separate the proof into two parts: when the vector fields are linearly independent at x_0 (i.e., when $n = q$) and more generally when the vector fields may be linearly dependent at x_0 (i.e., when $q \geq n$).

7.2.1 Linearly Independent

In this section we prove Theorem 3.10 in the special case when $n = q$; so that we have $X = X_{J_0}$. By Proposition 3.1, X_1, \dots, X_n span the tangent space at every point of $B_X(x_0, \xi)$. Since we know $B_X(x_0, \xi)$ is an n -dimensional manifold, we have that X_1, \dots, X_n form a basis for the tangent space at every point of $B_X(x_0, \xi)$. Taking $\chi = \xi$, Theorem 3.10 (a), (b), and (c) follow immediately.

We apply Proposition 6.2 to obtain $\hat{\eta} \in (0, \eta_0]$ as in that proposition; so that $\Phi : B^n(\hat{\eta}) \rightarrow B_X(x_0, \xi)$ and is a C^2 diffeomorphism onto its image. We let $Y_j = \Phi^* X_j$, and define A as in Theorem 3.10 by $Y = (I + A)\nabla$. Our main goal is to show that A is real analytic; this will imply that Y_1, \dots, Y_n are real analytic as well.

We have assumed $c_{j,k}^l \in \mathcal{A}_X^{x_0, \eta}$ with $\|c_{j,l}^l\|_{\mathcal{A}_X^{x_0, \eta}} \lesssim 1$. Thus, by the definition of $\mathcal{A}_X^{x_0, \eta}$, we have $c_{j,k}^l \circ \Phi \in \mathcal{A}^{n, \hat{\eta}}$ with $\|c_{j,k}^l \circ \Phi\|_{\mathcal{A}^{n, \hat{\eta}}} \lesssim 1$. We conclude $c_{j,k}^l \circ \Phi \in \mathcal{A}^{n, \hat{\eta}}$ with $\|c_{j,k}^l \circ \Phi\|_{\mathcal{A}^{n, \hat{\eta}}} \lesssim 1$.

For $1 \leq l \leq n$, define an $n \times n$ matrix $C_l(t)$ by letting the j, k component of $C_l(t)$ equal $c_{j,l}^k \circ \Phi$. Thus, $C_l \in \mathcal{A}^{n, \hat{\eta}}(\mathbb{M}^{n \times n})$ with $\|C_l\|_{\mathcal{A}^{n, \hat{\eta}}(\mathbb{M}^{n \times n})} \lesssim 1$. Set $C(t) := \sum_{l=1}^n t_l C_l(t)$. By Proposition 6.2 (c), $A(0) = 0$, and by Proposition 6.2 (d), A satisfies the differential equation (6.1).

Proposition 7.2 shows that there is an admissible constant $\eta_1 \in (0, \hat{\eta}]$ such that (6.1) has a solution $\hat{A} \in \mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ with $\|\hat{A}\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \frac{1}{2}$. By the uniqueness of this solution described in Proposition 7.2, $A|_{B^n(\eta_1)} = \hat{A}$. This establishes Theorem 3.10 (g). Theorem 3.10 (h) is an immediate consequence of (g) (since $Y = Y_{J_0}$ when $n = q$).

Proposition 6.2 (a) shows $\Phi(B^n(\hat{\eta}))$ is an open subset of $B_X(x_0, \xi)$ and Proposition 6.2 (b) shows $\Phi : B^n(\hat{\eta}) \rightarrow \Phi(B^n(\hat{\eta}))$ is a C^2 diffeomorphism. Since $\eta_1 \leq \hat{\eta}$, we see $\Phi(B^n(\eta_1))$ is an open subset of $B_X(x_0, \xi)$ and $\Phi : B^n(\eta_1) \rightarrow \Phi(B^n(\eta_1))$ is a C^2 diffeomorphism. This establishes Theorem 3.10 (d) and (e).

Finally, we prove Theorem 3.10 (f). We have already taken $\chi = \xi$, and we have $\Phi(B^n(\eta_1)) \subseteq B_X(x_0, \xi)$. Thus it suffices to prove the existence of ξ_1 and ξ_2 . Since $X_{J_0} = X$, we may take $\xi_1 = \xi_2$, and therefore we only need to prove the existence of ξ_1 . This follows just as in [SS18, Lemma 9.23].

7.2.2 Linearly Dependent

In this section we prove Theorem 3.10 in the general case, $q \geq n$. As in [SS18] the goal is to reduce the problem to the case $q = n$. Set

$$\mathcal{I}_0(n, q) := \{(i_1, \dots, i_n) \in \mathcal{I}(n, q) : 1 \leq i_1 < i_2 < \dots < i_n \leq q\}.$$

Lemma 7.11. *For $J \in \mathcal{I}(n, q)$, $1 \leq j \leq n$,*

$$\mathcal{L}_{X_j} \bigwedge X_J = \sum_{K \in \mathcal{I}_0(n, q)} g_{j,J}^K \bigwedge X_K, \text{ on } B_{X_{J_0}}(x_0, \xi),$$

where $\|g_{j,J}^K\|_{C(B_{X_{J_0}}(x_0, \xi))} \lesssim 0$ 1 and $\|g_{j,J}^K\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta}} \lesssim 1$. Here, \mathcal{L}_{X_j} denotes the Lie derivative with respect to X_j .

Proof. Let $J = (j_1, \dots, j_n)$. By the definition of \mathcal{L}_{X_j} (see [SS18, Section 5] for more details), we have

$$\mathcal{L}_{X_j} \bigwedge X_J = \mathcal{L}_{X_j} (X_{j_1} \wedge X_{j_2} \wedge \dots \wedge X_{j_n}) = \sum_{l=1}^n X_{j_1} \wedge X_{j_2} \wedge \dots \wedge X_{j_{l-1}} \wedge [X_j, X_{j_l}] \wedge X_{j_{l+1}} \wedge \dots \wedge X_{j_n}.$$

But, $[X_j, X_{j_l}] = \sum_{k=1}^q c_{j,j_l}^k X_k$, by assumption. Thus,

$$\mathcal{L}_{X_j} \bigwedge X_J = \sum_{l=1}^n \sum_{k=1}^q c_{j,j_l}^k X_{j_1} \wedge X_{j_2} \wedge \dots \wedge X_{j_{l-1}} \wedge X_k \wedge X_{j_{l+1}} \wedge \dots \wedge X_{j_n}.$$

The result now follows from the anti-commutativity of \wedge and the assumptions on $c_{i,j}^k$. \square

Take $\chi \in (0, \xi]$ to be the 0-admissible constant given by Proposition 6.1. With this choice of χ , Theorem 3.10 (a), (b), and (c) follow immediately. In particular, $\bigwedge X_{J_0}(y) \neq 0$ for $y \in B_X(x_0, \xi)$. It therefore makes sense to consider $\frac{\bigwedge X_J(y)}{\bigwedge X_{J_0}(y)}$ for any $J \in \mathcal{I}(n, q)$ and $y \in B_{X_{J_0}}(x_0, \chi)$.

Lemma 7.12. *For $J \in \mathcal{I}(n, q)$, $1 \leq j \leq n$,*

$$X_j \frac{\bigwedge X_J}{\bigwedge X_{J_0}} = \sum_{K \in \mathcal{I}_0(n, q)} g_{j,J}^K \frac{\bigwedge X_K}{\bigwedge X_{J_0}} - \sum_{K \in \mathcal{I}_0(n, q)} g_{j,J_0}^K \frac{\bigwedge X_J}{\bigwedge X_{J_0}} \frac{\bigwedge X_K}{\bigwedge X_{J_0}},$$

where $g_{j,J}^K$ are the functions from Lemma 7.11.

Proof. We use the identity

$$X_j \frac{\bigwedge X_J}{\bigwedge X_{J_0}} = \frac{\mathcal{L}_{X_j} \bigwedge X_J}{\bigwedge X_{J_0}} - \frac{\bigwedge X_J}{\bigwedge X_{J_0}} \frac{\mathcal{L}_{X_j} \bigwedge X_{J_0}}{\bigwedge X_{J_0}},$$

which is proved in [SS18, Lemma 5.1]. From here, the result follows immediately from Lemma 7.11. \square

Lemma 7.13. *For $J \in \mathcal{I}(n, q)$ we have $\frac{\bigwedge X_J}{\bigwedge X_{J_0}} \in C(B_{X_{J_0}}(x_0, \chi))$ and*

$$\left\| \frac{\bigwedge X_J}{\bigwedge X_{J_0}} \right\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_0 1.$$

Proof. This is just a restatement of Theorem 3.10 (b), which we have already shown. \square

Lemma 7.14. *There exists an admissible constant $\eta' \in (0, \eta]$ such that $\forall J \in \mathcal{I}(n, q)$, $\frac{\bigwedge X_J}{\bigwedge X_{J_0}} \in \mathcal{A}_{X_{J_0}}^{x_0, \eta'}$ and*

$$\left\| \frac{\bigwedge X_J}{\bigwedge X_{J_0}} \right\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \lesssim 1.$$

Proof. Set $N := |\mathcal{I}_0(n, q)|$ and let $G(x) \in C(B_{X_{J_0}}(x_0, \chi); \mathbb{R}^N)$ be given by $G_J(x) = \frac{\bigwedge X_J(x)}{\bigwedge X_{J_0}(x)}$, for $J \in \mathcal{I}_0(n, q)$. Lemma 7.12 shows that Proposition 7.10 applies to G (with $\xi = \chi$, $X = X_{J_0}$, $q = n$, $L = 2$, D an admissible constant, and $|G(x_0)| \leq \zeta^{-1}$). Here we taken η' to be r' from that result. The lemma follows. \square

Lemma 7.15. *For $1 \leq k \leq q$, $1 \leq l \leq n$, there exist $\tilde{b}_k^l \in C(B_{X_{J_0}}(x_0, \chi)) \cap \mathcal{A}_{X_{J_0}}^{x_0, \eta'}$ such that*

$$X_k = \sum_{l=1}^n \tilde{b}_k^l X_l. \quad (7.6)$$

Furthermore, $\|\tilde{b}_k^l\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_0 1$ and $\|\tilde{b}_k^l\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \lesssim 1$.

Proof. For $1 \leq k \leq q$, $1 \leq l \leq n$, set $J(l, k) = (1, 2, \dots, l-1, k, l+1, \dots, n) \in \mathcal{I}(n, q)$ and define

$$\tilde{b}_k^l := \frac{\bigwedge X_{J(l, k)}}{\bigwedge X_{J_0}}.$$

Cramer's rule shows (7.6) (see [SS18, Section 5] for more comments on this application of Cramer's rule). The desired estimates follow from Lemmas 7.13 and 7.14, completing the proof. \square

Lemma 7.16. *For $1 \leq i, j, k \leq n$, there exist $\hat{c}_{i,j}^k \in C(B_{X_{J_0}}(x_0, \chi)) \cap \mathcal{A}_{X_{J_0}}^{x_0, \eta'}$ such that*

$$[X_i, X_j] = \sum_{k=1}^n \hat{c}_{i,j}^k X_k. \quad (7.7)$$

Furthermore, $\|\hat{c}_{i,j}^k\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_0 1$ and $\|\hat{c}_{i,j}^k\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \lesssim 1$.

Proof. For $1 \leq i, j \leq n$ we have using Lemma 7.15

$$[X_i, X_j] = \sum_{k=1}^q c_{i,j}^k X_k = \sum_{l=1}^n \left(\sum_{k=1}^q c_{i,j}^k \tilde{b}_k^l \right) X_l.$$

Setting $\hat{c}_{i,j}^k := \sum_{k=1}^q c_{i,j}^k \tilde{b}_k^l$, (7.7) follows. We have $\|\hat{c}_{i,j}^k\|_{C(B_{X_{J_0}}(x_0, \xi))} \lesssim_0 1$ and $\|\hat{c}_{i,j}^k\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \lesssim 1$, by the definition of admissible constants. Combining this with Lemma 7.15, the fact that $\|fg\|_{C(B_{X_{J_0}}(x_0, \xi))} \leq \|f\|_{C(B_{X_{J_0}}(x_0, \xi))} \|g\|_{C(B_{X_{J_0}}(x_0, \xi))}$ (which is immediate from the definition), and $\|fg\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \leq \|f\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \|g\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}}$ (see Lemma 4.1), the desired estimates follow. \square

In light of Lemma 7.16, the case $n = q$ of Theorem 3.10 (which we proved in Section 7.2.1) applies to X_1, \dots, X_n with η replaced by η' and ξ replaced by χ ,¹⁰ yielding admissible constants $\eta_1 \in (0, \eta']$ and $\xi_1 > 0$ as in that theorem. This establishes (d), (e), (g), (f) (except the existence of ξ_2), and (h) for $1 \leq j \leq n$. Since we have already shown (a), (b), and (c), all that remains to show is the existence of ξ_2 as in (f) and (h) for $n+1 \leq j \leq q$. The existence of ξ_2 follows directly from [SS18, Lemma 9.35].

All that remains is (h) for $n+1 \leq j \leq q$. Lemma 7.15 shows $\|\tilde{b}_k^l\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \lesssim 1$. Thus, by the definition of $\mathcal{A}_{X_{J_0}}^{x_0, \eta'}$, we have $\|\tilde{b}_k^l \circ \Phi\|_{\mathcal{A}^{n, \eta_1}} \lesssim 1$. Pulling back (7.6) via Φ , we have for $n+1 \leq k \leq q$,

$$Y_k = \sum_{l=1}^n \left(\tilde{b}_k^l \circ \Phi \right) Y_l.$$

Since we already know $\|Y_l\|_{\mathcal{A}^{n, \eta_1}(\mathbb{R}^n)} \lesssim 1$ ($1 \leq l \leq n$) it follows that $\|Y_k\|_{\mathcal{A}^{n, \eta_1}(\mathbb{R}^n)} \lesssim 1$. Here, we have used that \mathcal{A}^{n, η_1} is a Banach algebra (Lemma 4.1). This completes the proof.

7.3 Densities

In this section, we prove Theorem 3.14 and Corollary 3.15. We take the same setting and notation as in that Section 3.2.1. In particular, we have all of the conclusions of Theorem 3.10, and take η_1 , χ , and Φ as in that theorem. In addition, we have a density ν on $B_{X_{J_0}}(x_0, \chi)$ (where χ is as in Theorem 3.10). Let f_j be as in Section 3.2.1. As in Theorem 3.10, we (without loss of generality) take $J_0 = (1, \dots, n)$.

Following [SS18], we introduce an auxiliary density on $B_{X_{J_0}}(x_0, \chi)$ given by

$$\nu_0(Z_1, \dots, Z_n) := \frac{|Z_1 \wedge Z_2 \wedge \dots \wedge Z_n|}{|X_1 \wedge X_2 \wedge \dots \wedge X_n|}. \quad (7.8)$$

Since $X_1 \wedge X_2 \wedge \dots \wedge X_n$ is never zero on $B_{X_{J_0}}(x_0, \chi)$ (Theorem 3.10 (a)), ν_0 is defined on $B_{X_{J_0}}(x_0, \chi)$ and is clearly a density. Note that $\nu_0(X_1, \dots, X_n) \equiv 1$, so that ν_0 is nonzero everywhere on $B_{X_{J_0}}(x_0, \chi)$.

Lemma 7.17. *For $1 \leq j \leq n$, $\mathcal{L}_{X_j} \nu_0 = f_j^0 \nu_0$, where $f_j^0 \in \mathcal{A}_{X_{J_0}}^{x_0, \eta_1} \cap C(B_{X_{J_0}}(x_0, \chi))$ and $\|f_j^0\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta_1}} \lesssim 1$, $\|f_j^0\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim 0$.*

Proof. The function f_j^0 was computed in the proof of [SS18, Lemma 9.38]. There it is shown

$$f_j^0 = - \sum_{K \in \mathcal{I}_0(n, q)} g_{j, J_0}^K \frac{\bigwedge X_K}{\bigwedge X_{J_0}},$$

where g_{j, J_0}^K is the function from Lemma 7.11. We have by Lemma 7.11,

$$\|g_{j, J_0}^K\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta_1}} \leq \|g_{j, J_0}^K\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta}} \lesssim 1, \quad \|g_{j, J_0}^K\|_{C(B_{X_{J_0}}(x_0, \chi))} \leq \|g_{j, J_0}^K\|_{C(B_{X_{J_0}}(x_0, \xi))} \lesssim 0.$$

Also, by Lemma 7.13 and Lemma 7.14,

$$\left\| \frac{\bigwedge X_K}{\bigwedge X_{J_0}} \right\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta_1}} \leq \left\| \frac{\bigwedge X_K}{\bigwedge X_{J_0}} \right\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta}} \lesssim 1, \quad \left\| \frac{\bigwedge X_J}{\bigwedge X_{J_0}} \right\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim 1.$$

where we have used that η' from the proof of Theorem 3.14 satisfies $\eta' \geq \eta_1$. Using the above, the result follows from Lemma 4.1. \square

Define $h_0 \in C(B^n(\eta_1))$ by $\Phi^* \nu_0 = h_0 \sigma_{\text{Leb}}$, where σ_{Leb} denotes the Lebesgue density on \mathbb{R}^n .

¹⁰When $n = q$, we proved Theorem 3.10 with $\chi = \xi$.

Lemma 7.18. $h_0(t) = \det(I + A(t))^{-1}$ where A is the matrix from Theorem 3.14. Furthermore, $h_0(t) \approx_0 1$, $\forall t \in B^n(\eta_1)$, and $h_0 \in \mathcal{A}^{n, \eta_1}$ with $\|h_0\|_{\mathcal{A}^{n, \eta_1}} \lesssim_0 1$.

Proof. Because $\sup_{t \in B^n(\eta_1)} \|A(t)\|_{\mathbb{M}^{n \times n}} \leq \|A\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \frac{1}{2}$ (by Theorem 3.10 (g)), we have $|\det(I + A(t))^{-1}| = \det(I + A(t))^{-1}$, for all $t \in B^n(\eta_1)$. Thus,

$$\begin{aligned} h_0(t) &= (\Phi^* \nu_0)(t) \left(\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_2}, \dots, \frac{\partial}{\partial t_n} \right) = (\Phi^* \nu_0)(t) ((I + A(t))^{-1} Y_1(t), \dots, (I + A(t))^{-1} Y_n(t)) \\ &= |\det(I + A(t))^{-1}| (\Phi^* \nu_0)(t) (Y_1(t), \dots, Y_n(t)) = \det(I + A(t))^{-1} \nu_0(\Phi(t)) (X_1(\Phi(t)), \dots, X_n(\Phi(t))) \\ &= \det(I + A(t))^{-1}. \end{aligned}$$

That $h_0(t) \approx_0 1$, $\forall t \in B^n(\eta_1)$, now follows from the above mentioned fact that $\sup_{t \in B^n(\eta_1)} \|A(t)\|_{\mathbb{M}^{n \times n}} \leq \frac{1}{2}$.

Since $\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ is a Banach algebra (Lemma 4.1) and since $\|A\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq \frac{1}{2}$, it follows that $I + A$ is invertible in $\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})$ with $\|(I + A)^{-1}\|_{\mathcal{A}^{n, \eta_1}(\mathbb{M}^{n \times n})} \leq 2$. We conclude $\|h_0\|_{\mathcal{A}^{n, \eta_1}} = \|\det(I + A)^{-1}\|_{\mathcal{A}^{n, \eta_1}} \lesssim_0 1$. \square

Since ν_0 is an everywhere nonzero density, there is a unique $g \in C(B_{X_{J_0}}(x_0, \chi))$ such that $\nu_0 = g\nu$.

Lemma 7.19. For $1 \leq j \leq n$, $X_j g = (f_j - f_j^0)g$, and $g(x) \approx_{0; \nu} g(x_0) = \nu(x_0)(X_1(x_0), \dots, X_n(x_0))$ for all $x \in B_{X_{J_0}}(x_0, \chi)$.

Proof. See [SS18, Section 9.4]. \square

Lemma 7.20. Set $s := \min\{\eta_1, r\}$. Then, $g \in \mathcal{A}_{X_{J_0}}^{x_0, s}$ and $\|g\|_{\mathcal{A}_{X_{J_0}}^{x_0, s}} \lesssim_{\nu} |\nu(X_1, \dots, X_n)(x_0)|$.

Proof. Set $B(t) := g \circ \Phi(t)$. The result can be rephrased as saying $B \in \mathcal{A}^{n, s}$ with $\|B\|_{\mathcal{A}^{n, s}} \lesssim_{\nu} |B(0)| = |g(x_0)| = |\nu(X_1, \dots, X_n)(x_0)|$. We have, using Lemma 7.19,

$$\begin{aligned} \frac{\partial}{\partial \epsilon} B(\epsilon t) &= ((t_1 X_1 + \dots + t_n X_n)g)(\Phi(\epsilon t)) = \sum_{j=1}^n t_j (f_j \circ \Phi(\epsilon t) - f_j^0 \circ \Phi(\epsilon t))g \circ \Phi(\epsilon t) \\ &= \sum_{j=1}^n t_j (f_j \circ \Phi(\epsilon t) - f_j^0 \circ \Phi(\epsilon t))B(\epsilon t). \end{aligned} \tag{7.9}$$

Solving the linear ODE (7.9) we have

$$B(t) = e^{\int_0^1 \sum_{j=1}^n t_j (f_j \circ \Phi(\epsilon t) - f_j^0 \circ \Phi(\epsilon t)) d\epsilon} B(0).$$

Since $\|f_j\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}} \lesssim_{\nu} 1$, by assumption, and $\|f_j^0\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta_1}} \lesssim_{\nu} 1$ by Lemma 7.17, we have $\|f_j \circ \Phi - f_j^0 \circ \Phi\|_{\mathcal{A}^{n, s}} \lesssim_{\nu} 1$.

By Lemma 7.8, if $F(t) = \int_0^1 \sum_{j=1}^n t_j (f_j \circ \Phi(\epsilon t) - f_j^0 \circ \Phi(\epsilon t)) d\epsilon$, then $\|F\|_{\mathcal{A}^{n, s}} \lesssim_{\nu} 1$. Finally, since $\mathcal{A}^{n, s}$ is a Banach algebra (by Lemma 4.1), $\|e^F\|_{\mathcal{A}^{n, s}} \leq e^{\|F\|_{\mathcal{A}^{n, s}}} \lesssim_{\nu} 1$. We conclude $\|B\|_{\mathcal{A}^{n, s}} \lesssim_{\nu} |B(0)|$, completing the proof. \square

Proof of Theorem 3.14. We have

$$h\sigma_{\text{Leb}} = \Phi^* \nu = \Phi^* g\nu_0 = (g \circ \Phi)\Phi^* \nu_0 = (g \circ \Phi)h_0\sigma_{\text{Leb}},$$

and therefore $h = (g \circ \Phi)h_0$. (a) follows by combining the fact that $g \approx_{0; \nu} \nu(X_1, \dots, X_n)(x_0)$ (Lemma 7.19) and $h_0 \approx 1$ (Lemma 7.18).

Since $\|g \circ \Phi\|_{\mathcal{A}^{n, s}} = \|g\|_{\mathcal{A}_{X_{J_0}}^{x_0, s}} \lesssim_{\nu} |\nu(X_1, \dots, X_n)(x_0)|$ (Lemma 7.20) and $\|h_0\|_{\mathcal{A}^{n, s}} \leq \|h_0\|_{\mathcal{A}^{n, \eta_1}} \lesssim_0 1$ (Lemma 7.18), (b) follows by the formula $h = (g \circ \Phi)h_0$ and Lemma 4.1. \square

Having proved Theorem 3.14 we turn to Corollary 3.15. To facilitate this, we introduce a corollary of Theorem 3.10.

Corollary 7.21. *Let $\eta_1, \xi_1, \xi_2 > 0$ be as in Theorem 3.10. Then, there exist admissible constants $\eta_2 \in (0, \eta_1]$, $0 < \xi_4 \leq \xi_3 \leq \xi_2$ such that*

$$\begin{aligned} B(x_0, \xi_4) &\subseteq B_{X_{J_0}}(x_0, \xi_3) \subseteq \Phi(B^n(\eta_2)) \subseteq B_{X_{J_0}}(x_0, \xi_2) \subseteq B_X(x_0, \xi_2) \\ &\subseteq B_{X_{J_0}}(x_0, \xi_1) \subseteq \Phi(B^n(\eta_2)) \subseteq B_{X_{J_0}}(x_0, \chi) \subseteq B_X(x_0, \xi). \end{aligned}$$

Proof. After obtaining η_1, ξ_1, ξ_2 from Theorem 3.10, apply Theorem 3.10 again with ξ replaced by ξ_2 to obtain η_2, ξ_3 , and ξ_4 as in the statement of the corollary. \square

Proof of Corollary 3.15. We have

$$\begin{aligned} \nu(B_{X_{J_0}}(x_0, \xi_2)) &= \int_{B_{X_{J_0}}(x_0, \xi_2)} \nu = \int_{\Phi^{-1}(B_{X_{J_0}}(x_0, \xi_2))} \Phi^* \nu \\ &= \int_{\Phi^{-1}(B_{X_{J_0}}(x_0, \xi_2))} h(t) dt \approx_{0; \nu} \sigma_{\text{Leb}}(\Phi^{-1}(B_{X_{J_0}}(x_0, \xi_2))) \nu(X_1, \dots, X_n)(x_0), \end{aligned} \quad (7.10)$$

where σ_{Leb} denotes Lebesgue measure, and we have used Theorem 3.14 (a). By Corollary 7.21 and the fact that $\eta_1, \eta_2 > 1$ are admissible constants, we have

$$1 \approx \sigma_{\text{Leb}}(B^n(\eta_2)) \leq \sigma_{\text{Leb}}(\Phi^{-1}(B_{X_{J_0}}(x_0, \xi_2))) \leq \sigma_{\text{Leb}}(B^n(\eta_1)) \approx 1. \quad (7.11)$$

Combining (7.10) and (7.11) proves $\nu(B_{X_{J_0}}(x_0, \xi_2)) \approx_{\nu} \nu(X_1, \dots, X_n)(x_0)$. The same proof works with $B_{X_{J_0}}(x_0, \xi_2)$ replaced by $B_X(x_0, \xi_2)$, which completes the proof of (3.4).

To prove (3.5), all that remains is to show

$$|\nu(X_1, \dots, X_n)(x_0)| \approx_0 \max_{j_1, \dots, j_n \in \{1, \dots, q\}} |\nu(X_{j_1}, \dots, X_{j_n})(x_0)| \quad (7.12)$$

Note that

$$1 = |\nu_0(X_1, \dots, X_n)(x_0)| \leq \max_{k_1, \dots, k_n \in \{1, \dots, q\}} |\nu_0(X_{k_1}, \dots, X_{k_n})(x_0)| \leq \zeta^{-1} \lesssim_0 1,$$

by the definition of ζ . We conclude

$$\max_{k_1, \dots, k_n \in \{1, \dots, q\}} |\nu_0(X_{k_1}, \dots, X_{k_n})(x_0)| \approx_0 1.$$

Thus,

$$\begin{aligned} |\nu(X_1, \dots, X_n)(x_0)| &= |g(x_0) \nu_0(X_1, \dots, X_n)(x_0)| = |g(x_0)| \\ &\approx_0 |g(x_0)| \max_{k_1, \dots, k_n \in \{1, \dots, q\}} |\nu_0(X_{k_1}, \dots, X_{k_n})(x_0)| = \max_{k_1, \dots, k_n \in \{1, \dots, q\}} |\nu(X_{k_1}, \dots, X_{k_n})(x_0)| \end{aligned}$$

This establishes (7.12) and completes the proof. \square

7.4 Qualitative Results

In this section, we prove the qualitative results; i.e., Theorems 3.3 and 3.4. These are simple consequences of Theorem 3.10. We begin with Theorem 3.3. For this we recall [SS18, Proposition 4.14].

Lemma 7.22 (Proposition 4.14 of [SS18]). *Let X_1, \dots, X_q be C^1 vector fields on a C^2 manifold \mathfrak{M} .*

(i) *Let $\mathcal{K} \Subset \mathfrak{M}$ be a compact set. Then $\exists \eta > 0$ such that $\forall x_0 \in \mathcal{K}$, X_1, \dots, X_n satisfy $\mathcal{C}(x_0, \eta, \mathfrak{M})$.*

(ii) *Let $\mathcal{K} \Subset \mathfrak{M}$ be a compact set. Then, there exists $\delta_0 > 0$ such that $\forall \theta \in S^{q-1}$ if $x \in \mathcal{K}$ is such that $\theta_1 X_1(x) + \dots + \theta_q X_q(x) \neq 0$, then $\forall r \in (0, \delta_0]$,*

$$e^{r\theta_1 X_1 + \dots + r\theta_q X_q} x \neq x.$$

Comments on the proof. In [SS18, Proposition 4.14], (i) was only stated for a fixed $x_0 \in \mathfrak{M}$ and not “uniformly on compact sets.” However, the same proof yields (i); indeed, it is an immediate consequence of the Picard-Lindelöf theorem. (ii) is stated directly in [SS18, Proposition 4.14]. \square

Remark 7.23. Lemma 7.22 shows that we always have η and δ_0 as in the assumptions of Theorem 3.10. Thus, if we wish to apply Theorem 3.10 to obtain a qualitative result, we do not need to verify the existence of η and δ_0 .

Lemma 7.24. *Let X_1, \dots, X_q be C^1 vector fields on an n -dimensional C^2 manifold M . Let $V \subseteq M$ be open, let $U \subseteq \mathbb{R}^n$ be an open neighborhood of $0 \in \mathbb{R}^n$, and let $\Phi : U \rightarrow V$ be a C^2 -diffeomorphism. Fix $r > 0$ so that $B^n(r) \subseteq U$ and set $x_0 = \Phi(0)$. Set $Y_j = \Phi^* X_j$ and suppose $Y_j \in \mathcal{A}^{n,r}(\mathbb{R}^n)$, $1 \leq j \leq q$, and for some j_0, k_0 $[Y_{j_0}, Y_{k_0}] = \sum_{l=1}^q \tilde{c}_{j_0, k_0}^l Y_l$, where $\tilde{c}_{j_0, k_0}^l \in \mathcal{A}^{n,r}$. Then, there exists a neighborhood V' of x_0 and $s > 0$ such that $[X_{j_0}, X_{k_0}] = \sum_{l=1}^q c_{j_0, k_0}^l X_l$ where $c_{j_0, k_0}^l \in C_X^{\omega, s}(V')$ and $c_{j_0, k_0}^l = \tilde{c}_{j_0, k_0}^l \circ \Phi^{-1}$.*

Proof. Since $\tilde{c}_{j_0, k_0}^l \in \mathcal{A}^{n,r}$ and $Y_j \in \mathcal{A}^{n,r}(\mathbb{R}^n)$ ($1 \leq j \leq q$), Lemma 4.2 (ii) shows $\tilde{c}_{j_0, k_0}^l \in C^{\omega, r/2}(B^n(r/2))$ and $Y_j \in C^{\omega, r/2}(B^n(r/2); \mathbb{R}^n)$. Proposition 4.7 shows that there exists $s > 0$ with $\tilde{c}_{j_0, k_0}^l \in C_Y^{\omega, s}(B^n(r/2))$, where Y denotes the list of vector fields Y_1, \dots, Y_q . Set $c_{j_0, k_0}^l := \tilde{c}_{j_0, k_0}^l \circ \Phi^{-1}$ and $V' := \Phi(B^n(r/2))$. Proposition 2.2 shows $c_{j_0, k_0}^l \in C_X^{\omega, s}(V')$ and pushing the formula $[Y_{j_0}, Y_{k_0}] = \sum_{l=1}^q \tilde{c}_{j_0, k_0}^l Y_l$ forward via Φ shows $[X_{j_0}, X_{k_0}] = \sum_{l=1}^q c_{j_0, k_0}^l X_l$. This completes the proof. \square

Proof of Theorem 3.3. (i) \Rightarrow (ii): Let U, V, Φ , and x_0 be as in (i). By the definition of $C^\omega(U; \mathbb{R}^n)$, there exists an $r > 0$ such that $\Phi^* X_1, \dots, \Phi^* X_q \in C^{\omega, r}(U; \mathbb{R}^n)$. Without loss of generality, assume $0 \in U$ and $\Phi(0) = x_0$. Reorder X_1, \dots, X_q so that $X_1(x_0), \dots, X_n(x_0)$ are linearly independent and let $Y_j := \Phi^* X_j$, so that $Y_j \in C^{\omega, r}(U; \mathbb{R}^n)$. Note that $Y_1(0), \dots, Y_n(0)$ span the tangent space $T_0 U$. Take $r_1 \in (0, r]$ so small $B^n(r_1) \subseteq U$. By Lemma 4.2 (i), $Y_j \in \mathcal{A}^{n, r_1}(\mathbb{R}^n)$, $1 \leq j \leq q$. Since Y_1, \dots, Y_n are real analytic and form a basis for the tangent space near 0, there exists $r_2 > 0$ such that

$$[Y_i, Y_j] = \sum_{k=1}^n \tilde{c}_{i,j}^k Y_k, \quad \tilde{c}_{i,j}^k \in \mathcal{A}^{n, r_2}.$$

Pushing this statement forward via Φ shows, for $1 \leq i, j \leq n$, $[X_i, X_j] = \sum_{k=1}^n \hat{c}_{i,j}^k X_k$, where $\hat{c}_{i,j}^k := \tilde{c}_{i,j}^k \circ \Phi^{-1}$. Lemma 7.24 shows there exists $s_1 > 0$ with $\hat{c}_{i,j}^k \in C_X^{\omega, s_1}(V') \subseteq C_X^\omega(V')$ for some neighborhood V' of x_0 .

Furthermore, since each Y_j ($1 \leq j \leq q$) is real analytic, and Y_1, \dots, Y_n form a basis of the tangent space near 0, there is $s_2 > 0$ such that, for $n+1 \leq j \leq q$,

$$Y_j = \sum_{k=1}^n \tilde{b}_j^k Y_k, \quad \tilde{b}_j^k \in \mathcal{A}^{n, s_2}. \quad (7.13)$$

By Lemma 4.2 (ii), $\tilde{b}_j^k \in C^{\omega, s_2/2}(B^n(s_2/2))$. Proposition 4.7 shows that there exists $s_3 > 0$ such that $\tilde{b}_j^k \in C_Y^{\omega, s_3}(B^n(s_2/2))$. Proposition 2.2 shows $b_j^k := \tilde{b}_j^k \circ \Phi^{-1} \in C_X^{\omega, s_3}(\Phi(B^n(s_2/2))) \subseteq C_X^\omega(\Phi(B^n(s_2/2)))$. Pushing (7.13) forward via Φ , we have $X_j = \sum_{k=1}^n b_j^k X_k$. Combining the above proves (ii).

(ii) \Rightarrow (iii): Suppose (ii) holds. Let V be as in (ii) and take $r > 0$ so that $\hat{c}_{i,j}^k, b_j^k \in C_X^{\omega, r}(V)$, $\forall i, j, k$. We wish to show for $1 \leq i, j \leq q$,

$$[X_i, X_j] = \sum_{k=1}^q \hat{c}_{i,j}^k X_k, \quad \hat{c}_{i,j}^k \in C_X^{\omega, r/2}(V). \quad (7.14)$$

For $1 \leq i, j \leq n$, (7.14) follows directly from the hypothesis (ii). We prove (7.14) for $n+1 \leq i, j \leq q$. The remaining cases ($1 \leq i \leq n$ and $n+1 \leq j \leq q$, or $n+1 \leq i \leq q$ and $1 \leq j \leq n$) are similar and easier. We have

$$[X_i, X_j] = \left[\sum_{k_1=1}^n b_i^{k_1} X_{k_1}, \sum_{k_2=1}^n b_j^{k_2} X_{k_2} \right] = \sum_{k_1, k_2=1}^n \left(b_i^{k_1} (X_{k_1} b_j^{k_2}) X_{k_2} - b_j^{k_2} (X_{k_2} b_i^{k_1}) X_{k_1} + \sum_{l=1}^n b_i^{k_1} b_j^{k_2} \hat{c}_{k_1, k_2}^l X_l \right).$$

We are given $b_j^k, \hat{c}_{k_1, j_2} \in C_X^{\omega, r}(V) \subseteq C_X^{\omega, r/2}(V)$ and Lemma 4.6 shows $X_l b_j^k \in C_X^{\omega, r/2}(V)$. Now the result follows from the fact that $C_X^{\omega, r/2}(V)$ is a Banach algebra (Lemma 4.1).

(iii) \Rightarrow (i): This is a consequence of Theorem 3.10. We make a few comments to this end. First of all, as discussed in Lemma 7.22 and Remark 7.23, there exist η and δ_0 as in the hypotheses of Theorem 3.10. In (iii) we are given $c_{i,j}^k \in C_X^{\omega, r}$ near x_0 for some $r > 0$, while in Theorem 3.10 it is assumed $c_{j,k}^l$ is continuous near x_0 and $c_{j,k}^k \in \mathcal{A}_{X_{J_0}}^{x_0, \eta}$ (where J_0 is chosen as in that theorem with $\zeta = 1$). However, $c_{i,j}^k \in C_X^{\omega, r}$ near x_0 clearly implies $c_{j,k}^l$ is continuous near x_0 , and Lemma 4.3 shows $C_X^{\omega, \eta} \subseteq \mathcal{A}_{X_{J_0}}^{x_0, \eta} \subseteq \mathcal{A}_{X_{J_0}}^{x_0, \eta}$, so by shrinking η to be $\leq r$, those hypotheses follow. With these remarks, Theorem 3.10 applies to yield the coordinate chart Φ as in that theorem. This coordinate chart has the properties given in (i), completing the proof. \square

We now turn to Theorem 3.4. The uniqueness of the real analytic structure described in that theorem follows from the next lemma.

Lemma 7.25. *Let M, N be two n -dimensional real analytic manifolds and suppose X_1, \dots, X_q are real analytic vector fields on M which span the tangent space at every point, and Z_1, \dots, Z_q are real analytic vector fields on N . Let $\Psi : M \rightarrow N$ be a C^2 diffeomorphism such that $\Psi_* X_j = Z_j$. Then Ψ is real analytic.*

Proof. Fix a point $x_0 \in M$. We will show Ψ is real analytic near x_0 . Reorder the vector fields X_1, \dots, X_q so that $X_1(x_0), \dots, X_n(x_0)$ are linearly independent; and therefore form a basis for the tangent space near x_0 . Reorder Z_1, \dots, Z_q in the corresponding way, so that we have $\Psi_* X_j = Z_j$. I.e., we have

$$d\Psi(x)X_j(x) = Z_j(\Psi(x)).$$

Pick a real analytic coordinate system, x_1, \dots, x_n , on M near x_0 . Since X_1, \dots, X_n span the tangent space near x_0 and are real analytic, and Z_1, \dots, Z_n are real analytic, we have, for x near x_0 ,

$$\frac{\partial \Psi}{\partial x_j}(x) = \sum_{l=1}^n a_{j,l}(x) F_{j,l}(\Psi(x)), \quad 1 \leq j \leq n,$$

where $a_{j,l}$ and $F_{j,l}$ are real analytic near x_0 and $\Psi(x_0)$, respectively. Proposition 7.1 applies to show Ψ is real analytic near x_0 , completing the proof. \square

Proof of Theorem 3.4. (i) \Rightarrow (ii) is obvious. For the converse, suppose (ii) holds. Thus, for each $x \in M$, there exist open sets $U_x \subseteq \mathbb{R}^n$, $V_x \subseteq M$, and a C^2 diffeomorphism $\Phi_x : U_x \rightarrow V_x$ such that if $Y_j^x = \Phi_x^* X_j$, then Y_j^x is real analytic on U_x . We wish to show that the collection $\{(\Phi_x^{-1}, V_x) : x \in M\}$ forms a real analytic atlas on M ; once this is shown, (i) will follow since then X_j will be real analytic with respect to this atlas by definition, and this atlas is clearly compatible with the C^2 structure on M . Hence, we need only verify that the transition functions are real analytic. Take $x_1, x_2 \in M$ such that $V_{x_1} \cap V_{x_2} \neq \emptyset$. Set $\Psi = \Phi_{x_2}^{-1} \circ \Phi_{x_1} : U_{x_1} \cap \Phi_{x_1}^{-1}(V_{x_2}) \rightarrow U_{x_2} \cap \Phi_{x_2}^{-1}(V_{x_1})$. We wish to show Ψ is a real analytic diffeomorphism. We already know Ψ is a C^2 diffeomorphism and $\Psi_* Y_j^{x_1} = Y_j^{x_2}$. That Ψ is real analytic now follows from Lemma 7.25, completing the proof of (i). As mentioned before, the uniqueness of the real analytic structure, as described in the theorem, follows from Lemma 7.25, completing the proof. \square

8 Densities in Euclidean Space

While the hypotheses in Section 3.2.1 concern densities on abstract manifolds, the most important special case which arises in applications is that of the induced Lebesgue density on real analytic submanifolds of Euclidean space. In this section, we describe how to apply Theorem 3.14 and Corollary 3.15 in such a setting.

Let $\Omega \subseteq \mathbb{R}^N$ be open and fix $x_0 \in \Omega$. Let $X_1, \dots, X_q \in C_{\text{loc}}^\omega(\Omega; \mathbb{R}^N)$ be real analytic vector fields on Ω . We suppose X_1, \dots, X_q satisfy the conditions of Theorem 3.10 with $\mathfrak{M} = \Omega$, and so we have admissible constants as in that theorem, and ξ, δ_0, η, J_0 , and ζ as in the hypotheses of Theorem 3.10, and we take χ as

in the conclusion of Theorem 3.10.¹¹ As in Theorem 3.10, we take $n := \dim \text{span}\{X_1(x_0), \dots, X_q(x_0)\}$. We also assume that ξ is chosen so that $B_X(x_0, \xi) \Subset \Omega$. Fix an open set Ω' with $B_X(x_0, \xi) \Subset \Omega' \Subset \Omega$.

Under these hypotheses, Proposition 3.1 applies to show that $B_X(x_0, \xi)$ is an n -dimensional, injectively immersed submanifold of Ω . Classical theorems show that $B_X(x_0, \xi)$ can be given the structure of a real analytic,¹² injectively immersed submanifold of Ω and X_1, \dots, X_q are real analytic vector fields on $B_X(x_0, \xi)$. Let ν denote the induced Lebesgue density on $B_X(x_0, \xi)$. The goal of this section is to describe how the hypotheses of Theorem 3.14 and Corollary 3.15 hold, for this choice of ν , in a quantitative way.¹³

As in Theorem 3.10 we, without loss of generality, reorder the vector fields so that $J_0 = (1, \dots, n)$.

Fix $\delta_1 > 0$ and $s_1 > 0$ so that

$$X_1, \dots, X_n \in C^{\omega, s_1}(B^N(x_0, \delta_1); \mathbb{R}^N),$$

where $B^N(x_0, \delta_1) = \{y \in \mathbb{R}^N : |x_0 - y| < \delta_1\}$.

Definition 8.1. We say C is an E-admissible constant¹⁴ if C can be chosen to depend only on anything an admissible constant may depend on (see Definition 3.9), and upper bounds for δ_1^{-1} , s_1^{-1} , N , $\|X_j\|_{C^1(\Omega')}$, $1 \leq j \leq n$, and $\|X_j\|_{C^{\omega, s_1}(B^N(x_0, \delta_1); \mathbb{R}^N)}$, $1 \leq j \leq n$. We write $A \lesssim_E B$ if $A \leq CB$, where C is a positive E-admissible constant. We write $A \approx_E B$ for $A \lesssim_E B$ and $B \lesssim_E A$.

The main result of this section is the following:

Proposition 8.2. *There exists an E-admissible constant $r > 0$ and functions $f_1, \dots, f_n \in \mathcal{A}_{X_{J_0}}^{x_0, r} \cap C(B_{X_{J_0}}(x_0, \chi))$ such that for $1 \leq j \leq n$, $\mathcal{L}_{X_j} \nu = f_j \nu$ and $\|f_j\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}} \lesssim_E 1$, $\|f_j\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_E 1$.*

Remark 8.3. It is an immediate consequence of Proposition 8.2 that Theorem 3.14 and Corollary 3.15 hold, for this choice of ν , where any constant which is ν -admissible in the sense of those results, is E-admissible in the sense of Definition 8.1.

The rest of this section is devoted to the proof of Proposition 8.2. By Lemma 7.16, for $1 \leq i, j, k \leq n$, there exists an admissible constant $\eta' > 0$ and $\hat{c}_{i,j}^k \in C(B_{X_{J_0}}(x_0, \chi)) \cap \mathcal{A}_{X_{J_0}}^{x_0, \eta'}$ such that

$$[X_i, X_j] = \sum_{k=1}^n \hat{c}_{i,j}^k X_k, \quad 1 \leq i, j \leq n, \quad (8.1)$$

where $\|\hat{c}_{i,j}^k\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_0 1$ and $\|\hat{c}_{i,j}^k\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \lesssim 1$.

We abuse notation and write X_{J_0} to both denote the list of vector fields X_1, \dots, X_n and the $N \times n$ matrix whose columns are given by X_1, \dots, X_n . For $K \in \mathcal{I}_0(n, N)$ we write X_{K, J_0} to denote the $n \times n$ submatrix of X_{J_0} given by taking the rows listed in K . We set $\det_{n \times n} X_{J_0} = (\det X_{K, J_0})_{K \in \mathcal{I}_0(n, N)}$, so that $\det_{n \times n} X_{J_0}$ is a vector (it is not important in which order the coordinates are arranged). Since $X_1 \wedge X_2 \wedge \dots \wedge X_n$ is never zero on $B_{X_{J_0}}(x_0, \chi)$ (by Theorem 3.10 (a)), we have $|\det_{n \times n} X_{J_0}| > 0$ on $B_{X_{J_0}}(x_0, \chi)$.

Lemma 8.4. *There exists an E-admissible constant $\eta'' > 0$ such that for $1 \leq j \leq n$ and $K \in \mathcal{I}_0(n, N)$,*

$$X_j \det X_{K, J_0} = \sum_{L \in \mathcal{I}_0(n, N)} \tilde{g}_{j, K}^L \det X_{L, J_0},$$

where $\tilde{g}_{j, K}^L \in \mathcal{A}_{X_{J_0}}^{x_0, \eta''} \cap C(B_{X_{J_0}}(x_0, \chi))$ and $\|\tilde{g}_{j, K}^L\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta''}} \lesssim_E 1$, $\|\tilde{g}_{j, K}^L\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_E 1$.

¹¹There is a sense in which Theorem 3.10 can always be applied to real analytic vector fields. This is the subject of Section 5.1.2 and Section 9. However, this section has a different thrust and so we assume the hypotheses of Theorem 3.10.

¹²It is not important for the results in this section that $B_X(x_0, \xi)$ can be given a real analytic structure.

¹³The quantitative estimates in this section do not follow from classical proofs. The main difficulty is that classical proofs break down near a singular point of the associated foliation. See Remarks 1.1 and 5.6.

¹⁴Here, E stands for ‘‘Euclidean’’.

Proof. For $K = (k_1, \dots, k_n) \in \mathcal{I}_0(n, N)$, set $\nu_K = dx_{k_1} \wedge dx_{k_2} \wedge \dots \wedge dx_{k_n}$, so that ν_K is an n -form on \mathbb{R}^N . Note that $\det X_{K, J_0} = \nu_K(\bigwedge X_{J_0})$. Using [Lee03, Proposition 18.9] we have for $1 \leq j \leq n$,

$$X_j \det X_{K, J_0} = \mathcal{L}_{X_j} \nu_K \left(\bigwedge X_{J_0} \right) = (\mathcal{L}_{X_j} \nu_K) \left(\bigwedge X_{J_0} \right) + \nu_K \left(\mathcal{L}_{X_j} \bigwedge X_{J_0} \right). \quad (8.2)$$

We address the two terms on the right hand side of (8.2) separately.

For the second term on the right hand side of (8.2), we have

$$\begin{aligned} \nu_K \left(\mathcal{L}_{X_j} \bigwedge X_{J_0} \right) &= \nu_K (\mathcal{L}_{X_j} (X_1 \wedge X_2 \wedge \dots \wedge X_n)) \\ &= \nu_K ([X_j, X_1] \wedge X_2 \wedge \dots \wedge X_n) + \nu_K (X_1 \wedge [X_j, X_2] \wedge X_3 \wedge \dots \wedge X_n) \\ &\quad + \dots + \nu_K (X_1 \wedge X_2 \wedge \dots \wedge X_{n-1} \wedge [X_j, X_n]) \end{aligned} \quad (8.3)$$

The terms on the right hand side of (8.3) are all similar, so we address only the first. We have, using (8.1),

$$\nu_K ([X_j, X_1] \wedge X_2 \wedge \dots \wedge X_n) = \sum_{l=1}^n \hat{c}_{j,1}^l \nu_K (X_l \wedge X_2 \wedge \dots \wedge X_n) = \hat{c}_{j,1}^1 \nu_K (X_1 \wedge X_2 \wedge \dots \wedge X_n) = \hat{c}_{j,1}^1 \det X_{K, J_0}.$$

Since $\hat{c}_{j,1}^1 \in \mathcal{A}_{X_{J_0}}^{x_0, \eta'} \cap C(B_{X_{J_0}}(x_0, \chi))$ with $\|\hat{c}_{j,1}^1\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'}} \lesssim 1$, $\|\hat{c}_{j,1}^1\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim 0$, this is of the desired form for any $\eta'' \leq \eta'$.

We now turn to the first term on the right hand side of (8.2). We have, for $K = (k_1, \dots, k_n)$,

$$\begin{aligned} \mathcal{L}_{X_j} \nu_K &= \mathcal{L}_{X_j} (dx_{k_1} \wedge dx_{k_2} \wedge \dots \wedge dx_{k_n}) \\ &= (\mathcal{L}_{X_j} dx_{k_1}) \wedge dx_{k_2} \wedge \dots \wedge dx_{k_n} + dx_{k_1} \wedge (\mathcal{L}_{X_j} dx_{k_2}) \wedge \dots \wedge dx_{k_n} + \dots + dx_{k_1} \wedge dx_{k_2} \wedge \dots \wedge (\mathcal{L}_{X_j} dx_{k_n}). \end{aligned} \quad (8.4)$$

Each term on the right hand side of (8.4) is similar, so we describe the first. For this, we write $X_j = \sum_{k=1}^N a_j^k \frac{\partial}{\partial x_k}$ ($1 \leq j \leq n$), where $a_j^k \in C^{\omega, s_1}(B^N(x_0, \delta_1)) \cap C^1(\Omega')$ with $\|a_j^k\|_{C^{\omega, s_1}(B^N(x_0, \delta_1))} \lesssim_{\mathbb{E}} 1$ and $\|a_j^k\|_{C^1(\Omega')} \lesssim_{\mathbb{E}} 1$. Then, $\mathcal{L}_{X_j} dx_{k_1} = da_j^{k_1} = \sum_{l=1}^N \frac{\partial a_j^{k_1}}{\partial x_l} dx_l$. Thus,

$$(\mathcal{L}_{X_j} dx_{k_1}) \wedge dx_{k_2} \wedge \dots \wedge dx_{k_n} = \sum_{l=1}^N \frac{\partial a_j^{k_1}}{\partial x_l} dx_l \wedge dx_{k_2} \wedge \dots \wedge dx_{k_n} = \sum_{l=1}^N \frac{\partial a_j^{k_1}}{\partial x_l} \epsilon_{K, l} \nu_{K_l},$$

where $\epsilon_{K, l} \in \{-1, 0, 1\}$ and $K_l \in \mathcal{I}_0(n, N)$ is obtained by reordering $(l, k_2, k_3, \dots, k_n)$ to be non-decreasing. Applying the same ideas to the other terms on the right hand side of (8.4), we see

$$(\mathcal{L}_{X_j} \nu_K) \left(\bigwedge X_{J_0} \right) = \sum_{L \in \mathcal{I}_0(n, N)} b_{j, K}^L \nu_L \left(\bigwedge X_{J_0} \right) = \sum_{L \in \mathcal{I}_0(n, N)} b_{j, K}^L \det X_{L, J_0}, \quad (8.5)$$

where each $b_{j, K}^L$ is a sum of terms of the form $\pm \frac{\partial a_j^k}{\partial x_l}$ where $1 \leq j \leq n$, $1 \leq k, l \leq N$ (and the number of terms in this sum is $\lesssim_{\mathbb{E}} 1$).

Since $a_j^k \in C^{\omega, s_1}(B^N(x_0, \delta_1))$ with $\|a_j^k\|_{C^{\omega, s_1}(B^N(x_0, \delta_1))} \lesssim_{\mathbb{E}} 1$, applying Lemma 4.6 (with $X = \nabla$) shows

$$\frac{\partial a_j^k}{\partial x_l} \in C^{\omega, s_1/2}(B^N(x_0, \delta_1)), \quad \left\| \frac{\partial a_j^k}{\partial x_l} \right\|_{C^{\omega, s_1/2}(B^N(x_0, \delta_1))} \lesssim_{\mathbb{E}} 1, \quad 1 \leq j \leq n, 1 \leq k, l \leq N.$$

We conclude $b_{j, K}^L \in C^{\omega, s_1/2}(B^N(x_0, \delta_1))$ with $\|b_{j, K}^L\|_{C^{\omega, s_1/2}(B^N(x_0, \delta_1))} \lesssim_{\mathbb{E}} 1$.

By Proposition 4.7 (i), there exists an E-admissible constant $s_2 > 0$ such that $b_{j, K}^L \in C_{X_{J_0}}^{\omega, s_2}(B^N(x_0, \delta_1))$ with $\|b_{j, K}^L\|_{C_{X_{J_0}}^{\omega, s_2}(B^N(x_0, \delta_1))} \lesssim_{\mathbb{E}} 1$. Also, it immediately follows from the properties of the a_j^k that $b_{j, K}^L \in C(B_{X_{J_0}}(x_0, \chi))$ with $\|b_{j, K}^L\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_{\mathbb{E}} 1$ (here we have used $\chi \leq \xi$ and therefore $B_{X_{J_0}}(x_0, \chi) \subseteq \Omega'$).

Because, for $1 \leq j \leq n$, $\|X_j\|_{C^1(B^N(x_0, \delta_1); \mathbb{R}^N)} \lesssim_E \|X_j\|_{C^{\omega, s_1}(B^N(x_0, \delta_1); \mathbb{R}^N)} \lesssim_E 1$, the Picard-Lindelöf theorem shows that we may take an E-admissible constant $\eta'' \in (0, \min\{s_2, \eta'\}]$ so small that X_{J_0} satisfies $\mathcal{C}(x_0, \eta'', B^N(x_0, \delta_1))$. Then, by Lemma 4.3 we have, for $1 \leq j \leq n$, $K, L \in \mathcal{I}_0(n, N)$,

$$\|b_{j,K}^L\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta''}} \leq \|b_{j,K}^L\|_{C_{X_{J_0}}^{\omega, s_2}(B^N(x_0, \delta_1))} \lesssim_E 1.$$

Combining this with (8.5) and the above mentioned fact that $\|b_{j,K}^L\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_E 1$, shows the first term on the right hand side of (8.2) is of the desired form, completing the proof. \square

Lemma 8.5. *Let $K \in \mathcal{I}_0(n, N)$, $1 \leq j \leq n$. Then,*

$$X_j \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} = \sum_{L \in \mathcal{I}_0(n, N)} \tilde{g}_{j,K}^L \frac{\det X_{L, J_0}}{|\det_{n \times n} X_{J_0}|} - \sum_{L_1, L_2 \in \mathcal{I}_0(n, N)} \tilde{g}_{j, L_1}^{L_2} \frac{(\det X_{K, J_0})(\det X_{L_1, J_0})(\det X_{L_2, J_0})}{|\det_{n \times n} X_{J_0}|^3},$$

where $\tilde{g}_{j, L_1}^{L_2}$ are the functions from Lemma 8.4.

Proof. We have, using Lemma 8.4, for $1 \leq j \leq n$, $K \in \mathcal{I}_0(n, N)$,

$$\begin{aligned} X_j \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} &= \frac{X_j \det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} - \frac{1}{2} \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|^3} X_j |\det_{n \times n} X_{J_0}|^2 \\ &= \sum_{L \in \mathcal{I}_0(n, N)} \tilde{g}_{j,K}^L \frac{\det X_{L, J_0}}{|\det_{n \times n} X_{J_0}|} - \sum_{L_1 \in \mathcal{I}_0(n, N)} \frac{(\det X_{K, J_0})(\det X_{L_1, J_0})(X_j \det X_{L_1, J_0})}{|\det_{n \times n} X_{J_0}|^3} \\ &= \sum_{L \in \mathcal{I}_0(n, N)} \tilde{g}_{j,K}^L \frac{\det X_{L, J_0}}{|\det_{n \times n} X_{J_0}|} - \sum_{L_1, L_2 \in \mathcal{I}_0(n, N)} \tilde{g}_{j, L_1}^{L_2} \frac{(\det X_{K, J_0})(\det X_{L_1, J_0})(\det X_{L_2, J_0})}{|\det_{n \times n} X_{J_0}|^3}, \end{aligned}$$

completing the proof. \square

Lemma 8.6. *There exists an E-admissible constant $\eta''' > 0$ such that $\forall K \in \mathcal{I}_0(n, N)$,*

$$\frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} \in \mathcal{A}_{X_{J_0}}^{x_0, \eta'''}, \quad \left\| \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} \right\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta'''}} \lesssim_E 1.$$

Proof. For $K \in \mathcal{I}_0(n, N)$, set $G_K := \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|}$. Lemma 8.5 shows, for $1 \leq j \leq n$,

$$X_j G_K = \sum_{L \in \mathcal{I}_0(n, N)} \tilde{g}_{j,K}^L G_L - \sum_{L_1, L_2 \in \mathcal{I}_0(n, N)} \tilde{g}_{j, L_1}^{L_2} G_K G_{L_1} G_{L_2}.$$

Using the estimates on the functions $\tilde{g}_{j, L_1}^{L_2}$ described in Lemma 8.4, the result follows from Proposition 7.10, with $q = n$, $\eta = \eta'''$, $\xi = \chi$, D an E-admissible constant, $L = 3$, $N = |\mathcal{I}_0(n, N)|$, and $|G(x_0)| = 1$. Here, we take η''' to be the r' guaranteed by that result. \square

Lemma 8.7.

$$\frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} \in C(B_{X_{J_0}}(x_0, \chi)), \quad \left\| \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} \right\|_{C(B_{X_{J_0}}(x_0, \chi))} \leq 1.$$

Proof. Since $\wedge X_{J_0}$ is never zero on $B_{X_{J_0}}(x_0, \chi)$ (by Theorem 3.10 (a)), the continuity of $\frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|}$ follows immediately. That it is bounded by 1 follows directly from the definition. \square

Lemma 8.8. *There exists an E-admissible constant $\eta_2 > 0$ and functions $h_j \in \mathcal{A}_{X_{J_0}}^{x_0, \eta_2} \cap C(B_{X_{J_0}}(x_0, \chi))$, $1 \leq j \leq n$ so that*

$$X_j \left| \det_{n \times n} X_{J_0} \right| = h_j \left| \det_{n \times n} X_{J_0} \right|, \quad 1 \leq j \leq n,$$

and $\|h_j\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta_2}} \lesssim_E 1$, $\|h_j\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_E 1$.

Proof. Using Lemma 8.4 we have

$$\begin{aligned}
X_j \left| \det_{n \times n} X_{J_0} \right| &= \frac{1}{2} \left| \det_{n \times n} X_{J_0} \right|^{-1} X_j \left| \det_{n \times n} X_{J_0} \right|^2 = \frac{1}{2} \left| \det_{n \times n} X_{J_0} \right|^{-1} \sum_{K \in \mathcal{I}_0(n, N)} X_j (\det X_{K, J_0})^2 \\
&= \left| \det_{n \times n} X_{J_0} \right|^{-1} \sum_{K, L \in \mathcal{I}_0(n, N)} \tilde{g}_{j, K}^L (\det X_{K, J_0}) (\det X_{L, J_0}) \\
&= \left| \det_{n \times n} X_{J_0} \right| \sum_{K, L \in \mathcal{I}_0(n, N)} \tilde{g}_{j, K}^L \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} \frac{\det X_{L, J_0}}{|\det_{n \times n} X_{J_0}|}.
\end{aligned}$$

We set $h_j := \sum_{K, L \in \mathcal{I}_0(n, N)} \tilde{g}_{j, K}^L \frac{\det X_{K, J_0}}{|\det_{n \times n} X_{J_0}|} \frac{\det X_{L, J_0}}{|\det_{n \times n} X_{J_0}|}$, and let $\eta_2 := \eta'' \wedge \eta'''$. Then, using the bounds on $\tilde{g}_{j, K}^L$ from Lemma 8.4, the bounds on $\frac{\det X_{L, J_0}}{|\det_{n \times n} X_{J_0}|}$ from Lemmas 8.6 and 8.7, and Lemma 4.1, we have $h_j \in \mathcal{A}_{X_{J_0}}^{x_0, \eta_2} \cap C(B_{X_{J_0}}(x_0, \chi))$ and $\|h_j\|_{\mathcal{A}_{X_{J_0}}^{x_0, \eta_2}} \lesssim_{\mathbb{E}} 1$, $\|h_j\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_{\mathbb{E}} 1$ completing the proof. \square

Lemma 8.9. $\nu = |\det_{n \times n} X_{J_0}| \nu_0$ on $B_{X_{J_0}}(x_0, \chi)$, where ν_0 is given by (7.8).

Proof. Since $\nu_0(\wedge X_{J_0}) \equiv 1$, ν_0 is a strictly positive density on $B_{X_{J_0}}(x_0, \chi)$. Thus, $\nu = f(x)\nu_0$ for some $f : B_{X_{J_0}}(x_0, \chi) \rightarrow \mathbb{R}$. To solve for f we evaluate this equation at $\wedge X_{J_0}$ and since $\nu_0(\wedge X_{J_0}) \equiv 1$, we have $f = \nu(\wedge X_{J_0})$. Since ν is the induced Lebesgue density on an n -dimensional, injectively immersed, submanifold of \mathbb{R}^N (to which X_1, \dots, X_q are tangent), we have $\nu(\wedge X_{J_0}) = |\det_{n \times n} X_{J_0}|$, completing the proof. \square

Proof of Proposition 8.2. For $1 \leq j \leq n$, let $f_j := f_j^0 + h_j$, where f_j^0 is described in Lemma 7.17 and h_j is described in Lemma 8.8. Then, if $r := \min\{\eta_1, \eta_2\} > 0$ we have that r is an E-admissible constant, $f_j \in \mathcal{A}_{X_{J_0}}^{x_0, r} \cap C(B_{X_{J_0}}(x_0, \chi))$, and $\|f_j\|_{\mathcal{A}_{X_{J_0}}^{x_0, r}} \lesssim_{\mathbb{E}} 1$, $\|f_j\|_{C(B_{X_{J_0}}(x_0, \chi))} \lesssim_{\mathbb{E}} 1$.

Using Lemmas 7.17, 8.8, and 8.9 we have, for $1 \leq j \leq n$,

$$\mathcal{L}_{X_j} \nu = \mathcal{L}_{X_j} \left| \det_{n \times n} X_{J_0} \right| \nu_0 = \left(X_j \left| \det_{n \times n} X_{J_0} \right| \right) \nu_0 + \left| \det_{n \times n} X_{J_0} \right| \mathcal{L}_{X_j} \nu_0 = h_j \left| \det_{n \times n} X_{J_0} \right| \nu_0 + f_j^0 \left| \det_{n \times n} X_{J_0} \right| \nu_0 = f_j \nu,$$

completing the proof. \square

9 Scaling and real analyticity: the proof of Theorem 5.4

In this section, we prove Theorem 5.4. Fix a compact set $\mathcal{K} \Subset \Omega$. The idea is that, if m is chosen sufficiently large (depending on \mathcal{K} and V_1, \dots, V_r), then a proof similar to that of Theorem 5.7 will work uniformly for $x \in \mathcal{K}$ with the manifold M from Theorem 5.7 replaced by L_x . As in Theorem 5.4, throughout this section we use $A \lesssim B$ to denote $A \leq CB$ where C can be chosen independent of $x \in \mathcal{K}$ and $\delta \in (0, 1]$.

We first show how the appropriate conditions hold uniformly, which uses the Weierstrass preparation theorem. This is based on an idea of Lobry [Lob70]¹⁵ (see also [Sus73, p. 188]), and was used in a similar context in [SS12] and [Str14, Section 2.15.5].

We take the same setting and notation as in Theorem 5.4. Thus, we have real analytic vector fields V_1, \dots, V_r on an open set $\Omega \subseteq \mathbb{R}^n$. We assign to each V_1, \dots, V_r the formal degree 1. If Z has formal degree e , we assign to $[V_j, Z]$ the formal degree $e + 1$. We let \mathcal{S} denote the (infinite) set of all such vector fields with formal degrees; thus each $(X, e) \in \mathcal{S}$ is a pair of a real analytic vector field X and $e \in \mathbb{N}$, where X is an iterated commutator of V_1, \dots, V_r .

An important ingredient is the next proposition:

¹⁵Lemmas 9.2 and 9.3 are classical, and it was Lobry [Lob70] who first used them to prove a result similar to Proposition 9.1. Unfortunately, there is a slight error in [Lob70]; see [Ste80]. Below, we begin with a *finite* collection of vector fields V_1, \dots, V_r , which allows us to avoid this issue.

Proposition 9.1. Fix $x \in \Omega$. Then, there exists an open neighborhood $U_x \subseteq \Omega$ of x and $m_x \in \mathbb{N}$ such that the following holds. Let $m \geq m_x$, and let $(X_1, d_1), \dots, (X_q, d_q)$ be an enumeration of those elements $(Z, e) \in \mathcal{S}$ with $e \leq m$. Then there exist real analytic functions $\hat{c}_{j,k}^{l,x} \in C^\omega(U_x)$ with

$$[X_j, X_k] = \sum_{d_l \leq d_j + d_k} \hat{c}_{j,k}^{l,x} X_l \text{ on } U_x.$$

We turn to the proof of Proposition 9.1. In the next few results, we (without loss of generality) relabel the fixed point $x \in \Omega$ from Proposition 9.1 to be 0. Thus, we work near the point $0 \in \mathbb{R}^n$. We write $f : \mathbb{R}_0^n \rightarrow \mathbb{R}^n$ to denote that f is a germ of a function defined near $0 \in \mathbb{R}^n$. Let

$$\mathcal{A}_n := \{f : \mathbb{R}_0^n \rightarrow \mathbb{R} \mid f \text{ is real analytic}\},$$

$$\mathcal{A}_n^m := \{f : \mathbb{R}_0^n \rightarrow \mathbb{R}^m \mid f \text{ is real analytic}\}.$$

Notice that \mathcal{A}_n^m can be identified with the m -fold Cartesian product of \mathcal{A}_n ; justifying our notation. \mathcal{A}_n^n can also be identified with the space of germs of real analytic vector fields near $0 \in \mathbb{R}^n$.

Lemma 9.2. The ring \mathcal{A}_n is Noetherian.

Comments on the proof. This is a simple consequence of the Weierstrass preparation theorem. See page 148 of [ZS75]. The proof in [ZS75] is for the formal power series ring, however, as mentioned on page 130 of [ZS75], the proof also works for the ring of convergent power series. This is exactly the ring \mathcal{A}_n . \square

Lemma 9.3. The module \mathcal{A}_n^m is a Noetherian \mathcal{A}_n module.

Comments on the proof. It is a standard fact that for any Noetherian ring R , the R -module R^m is Noetherian. \square

Lemma 9.4. Let $\mathcal{S} \subset \mathcal{A}_n^n \times \mathbb{N}$. Then, there exists a finite subset $\mathcal{F} \subseteq \mathcal{S}$ such that for every $(g, e) \in \mathcal{S}$,

$$g(x) = \sum_{\substack{(f,d) \in \mathcal{F} \\ d \leq e}} c_{(f,d)}(x) f(x),$$

with $c_{f,d} \in \mathcal{A}_n$.

Proof. Define a map $\iota : \mathcal{A}_n^n \times \mathbb{N} \rightarrow \mathcal{A}_{n+1}^n$ by $\iota(f, d) = t^d f(x)$, where $t \in \mathbb{R}$, $x \in \mathbb{R}^n$. Let \mathcal{M} be the submodule of \mathcal{A}_{n+1}^n generated by $\iota\mathcal{S}$. \mathcal{M} is finitely generated by Lemma 9.3. Let $\mathcal{F} \subseteq \mathcal{S}$ be a finite subset so that $\iota\mathcal{F}$ generates \mathcal{M} . We will show that \mathcal{F} satisfies the conclusions of the lemma.

Let $(g, e) \in \mathcal{S}$. Since $t^e g \in \mathcal{M}$, we have

$$t^e g(x) = \sum_{(f,d) \in \mathcal{F}} \hat{c}_{(f,d)}(t, x) t^d f(x), \quad \hat{c}_{(f,d)} \in \mathcal{A}_{n+1},$$

on a neighborhood of $(0, 0) \in \mathbb{R} \times \mathbb{R}^n$. Applying $\frac{1}{e!} \partial_t^e \big|_{t=0}$ to both sides and using that $\frac{1}{e!} \partial_t^e \big|_{t=0} t^d c_{(f,d)}(t, x) = 0$ if $d > e$, we have

$$g(x) = \sum_{\substack{(f,d) \in \mathcal{F} \\ d \leq e}} \left[\frac{1}{e!} \partial_t^e \big|_{t=0} t^d \hat{c}_{(f,d)}(t, x) \right] f(x).$$

The result follows. \square

We return to the setting at the start of this section. We let $\widehat{\mathcal{S}}$ denote the smallest collection of vector fields paired with formal degrees such that:

- $(V_1, 1), \dots, (V_r, 1) \in \widehat{\mathcal{S}}$.

- If $(Y, d), (Z, e) \in \widehat{\mathcal{S}}$, then $([Y, Z], d + e) \in \widehat{\mathcal{S}}$.

Note that $\widehat{\mathcal{S}}$ and \mathcal{S} (where \mathcal{S} is defined at the start of this section) are closely related. Indeed, $\mathcal{S} \subseteq \widehat{\mathcal{S}}$, and if $(Z, e) \in \widehat{\mathcal{S}}$ then Z is a finite linear combination (with constant coefficients) of vector fields Z' such that $(Z', e) \in \mathcal{S}$ (this follows from the Jacobi identity).

Lemma 9.5. *Fix $x \in \Omega$. There exists a finite set $\mathcal{F}_x \subseteq \mathcal{S}$ and an open neighborhood $U_x \Subset \Omega$ of x such that for every $(X, d) \in \widehat{\mathcal{S}}$,*

$$X = \sum_{\substack{(Z,e) \in \mathcal{F}_x \\ e \leq d}} c_{(X,d)}^{(Z,e)} Z, \quad c_{(X,d)}^{(Z,e)} \in C^\omega(U_x).$$

Proof. In this proof we relabel x to be 0. We apply Lemma 9.4 to $\widehat{\mathcal{S}}$ to obtain a finite set $\widehat{\mathcal{F}} \subseteq \widehat{\mathcal{S}}$ as in that lemma. Thus, every element $(X, d) \in \widehat{\mathcal{S}}$ can be written in the form:

$$X = \sum_{\substack{(Z,e) \in \widehat{\mathcal{F}} \\ e \leq d}} c_{(X,d)}^{(Z,e)} Z, \quad c_{(X,d)}^{(Z,e)} \in \mathcal{A}_n.$$

Since every vector field appearing in $\widehat{\mathcal{S}}$ is a finite linear combination (with constant coefficients) of vector fields with the same degree in \mathcal{S} , there is a finite set $\mathcal{F}_1 \subseteq \mathcal{S}$ such that every element $(X, d) \in \widehat{\mathcal{S}}$ can be written in the form:

$$X = \sum_{\substack{(Z,e) \in \mathcal{F}_1 \\ e \leq d}} c_{(X,d)}^{(Z,e)} Z, \quad c_{(X,d)}^{(Z,e)} \in \mathcal{A}_n. \quad (9.1)$$

The problem is that, a priori, the neighborhood of 0 on which $c_{(X,d)}^{(Z,e)}$ are defined might depend on (X, d) . Our goal is to find a common neighborhood of 0 which works for all $(X, d) \in \widehat{\mathcal{S}}$.

Let $m := \max\{d : \exists (X, d) \in \mathcal{F}_1\}$ and set $\mathcal{F} := \{(X, d) \in \mathcal{S} : d \leq m\}$. Note that $\mathcal{F} \subset \mathcal{S}$ is a finite set and $\mathcal{F}_1 \subseteq \mathcal{F}$. Furthermore, (9.1) holds with \mathcal{F}_1 replaced by \mathcal{F} (since $\mathcal{F}_1 \subseteq \mathcal{F}$). For each $(X_1, d_1), (X_2, d_2) \in \mathcal{F}$, $([X_1, X_2], d_1 + d_2) \in \widehat{\mathcal{S}}$ and therefore (9.1) holds with (X, d) replaced by $([X_1, X_2], d_1 + d_2)$ and \mathcal{F}_1 replaced by \mathcal{F} . Since there are only finitely many such vector fields, there is a common neighborhood U_1 of 0 such that for each $(X_1, d_1), (X_2, d_2) \in \mathcal{F}$ we may write

$$[X_1, X_2] = \sum_{\substack{(Z,e) \in \mathcal{F} \\ e \leq d_1 + d_2}} b_{(X_1, d_1), (X_2, d_2)}^{(Z,e)} Z, \quad b_{(X_1, d_1), (X_2, d_2)}^{(Z,e)} \in C_{\text{loc}}^\omega(U_1). \quad (9.2)$$

We claim, $\forall (X, d) \in \widehat{\mathcal{S}}$,

$$X = \sum_{\substack{(Z,e) \in \mathcal{F} \\ e \leq d}} c_{(X,d)}^{(Z,e)} Z, \quad c_{(X,d)}^{(Z,e)} \in C_{\text{loc}}^\omega(U_1). \quad (9.3)$$

We prove (9.3) by induction on d . Since $(V_1, 1), \dots, (V_r, 1) \in \mathcal{F}$, the base case, $d = 1$, is clear. Let $d_0 \geq 2$; we assume the result for all $d < d_0$ and prove the result for d_0 . If $(X, d_0) \in \widehat{\mathcal{S}}$ with $d_0 \geq 2$, then $X = [X_1, X_2]$ where $(X_1, d_1), (X_2, d_2) \in \widehat{\mathcal{S}}$ with $d_1 + d_2 = d_0$. In particular, $d_1, d_2 < d_0$ and so by the inductive hypothesis we may write for $j = 1, 2$,

$$X_j = \sum_{\substack{(Z,e) \in \mathcal{F} \\ e \leq d_j}} c_{(X_j, d_j)}^{(Z,e)} Z, \quad c_{(X_j, d_j)}^{(Z,e)} \in C_{\text{loc}}^\omega(U_1).$$

Thus, we have, using (9.2),

$$\begin{aligned}
X = [X_1, X_2] &= \left[\sum_{\substack{(Z_1, e_1) \in \mathcal{F} \\ e_1 \leq d_1}} c_{(X_1, d_1)}^{(Z_1, e_1)} Z_1, \sum_{\substack{(Z_2, e_2) \in \mathcal{F} \\ e_2 \leq d_2}} c_{(X_2, d_2)}^{(Z_2, e_2)} Z_2 \right] \\
&= \sum_{\substack{(Z_1, e_1) \in \mathcal{F} \\ e_1 \leq d_1}} \sum_{\substack{(Z_2, e_2) \in \mathcal{F} \\ e_2 \leq d_2}} \left(\left(Z_1 c_{(X_2, d_2)}^{(Z_2, e_2)} \right) Z_2 - \left(Z_2 c_{(X_1, d_1)}^{(Z_1, e_1)} \right) Z_1 + c_{(X_1, d_1)}^{(Z_1, e_1)} c_{(X_2, d_2)}^{(Z_2, e_2)} [Z_1, Z_2] \right) \\
&= \sum_{\substack{(Z_1, e_1) \in \mathcal{F} \\ e_1 \leq d_1}} \sum_{\substack{(Z_2, e_2) \in \mathcal{F} \\ e_2 \leq d_2}} \left(\left(Z_1 c_{(X_2, d_2)}^{(Z_2, e_2)} \right) Z_2 - \left(Z_2 c_{(X_1, d_1)}^{(Z_1, e_1)} \right) Z_1 + c_{(X_1, d_1)}^{(Z_1, e_1)} c_{(X_2, d_2)}^{(Z_2, e_2)} \sum_{\substack{(Z_3, e_3) \in \mathcal{F} \\ e_3 \leq e_1 + e_2}} b_{(Z_1, e_1), (Z_2, e_2)}^{(Z_3, e_3)} Z_3 \right).
\end{aligned}$$

Since $c_{(X_1, d_1)}^{(Z_1, e_1)}, c_{(X_2, d_2)}^{(Z_2, e_2)}, b_{(Z_1, e_1), (Z_2, e_2)}^{(Z_3, e_3)} \in C_{\text{loc}}^\omega(U_1)$, $\forall (Z_1, e_1), (Z_2, e_2), (Z_3, e_3) \in \mathcal{F}$, (9.3) follows for (X, d) , completing the proof of (9.3). Taking $\mathcal{F}_x := \mathcal{F}$ and $U \Subset U_1$ a relatively compact open set containing $0 = x$, the result follows. \square

Proof of Proposition 9.1. Let $\mathcal{F}_x \subset \mathcal{S}$ and $U_x \subseteq \Omega$ be as in Lemma 9.5. Set $m_x := \max\{e : \exists (Z, e) \in \mathcal{F}_x\}$. For $m \geq m_x$ let $(X_1, d_1), \dots, (X_q, d_q)$ be an enumeration of those elements $(Z, e) \in \mathcal{S}$ with $e \leq m$. Since $\mathcal{F}_x \subseteq \{(X_1, d_1), \dots, (X_q, d_q)\}$ and for each i, j , $([X_i, X_j], d_i + d_j) \in \widehat{\mathcal{S}}$, the result follows from Lemma 9.5. \square

Lemma 9.6. *There exists $m = m(\mathcal{K}) \in \mathbb{N}$ such that the following holds. Let $(X_1, d_1), \dots, (X_q, d_q)$ be the list of vector fields with formal degrees $\leq m$ as defined at the start of this section. Then, there exists $\xi \in (0, 1]$, $s > 0$ such that $\forall x \in \mathcal{K}$,*

$$[X_j, X_k] = \sum_{d_l \leq d_j + d_k} c_{j,k}^{l,x} X_l, \quad c_{j,k}^{l,x} \in C_X^{\omega,s}(B_X(x, \xi)), \quad (9.4)$$

where X denotes the list of vector fields X_1, \dots, X_q . Furthermore,

$$\sup_{x \in \mathcal{K}} \|c_{j,k}^{l,x}\|_{C_X^{\omega,s}(B_X(x, \xi))} < \infty.$$

Finally, $\bigcup_{x \in \mathcal{K}} B_X(x, \xi) \Subset \Omega$.

Proof. For each $x \in \Omega$, let $m_x \in \mathbb{N}$ and $U_x \Subset \Omega$ be as in Proposition 9.1. Fix an open set Ω' with $\mathcal{K} \Subset \Omega' \Subset \Omega$, and set $\mathcal{K}_1 := \overline{\Omega'} \Subset \Omega$. $\{U_x : x \in \mathcal{K}_1\}$ is an open cover for \mathcal{K}_1 and we extract a finite sub-cover U_{x_1}, \dots, U_{x_R} . Set $m := \max\{m_{x_k} : 1 \leq k \leq R\}$ and let $(X_1, d_1), \dots, (X_q, d_q) \in \mathcal{S}$ be an enumeration of all the vector fields in \mathcal{S} with degree $d_j \leq m$.

We claim that there exists $\xi \in (0, 1]$ such that $\forall x \in \mathcal{K}$, $\exists k \in \{1, \dots, R\}$ with $B_X(x, \xi) \subseteq U_{x_k}$. Consider the list of vector fields $W = X_1, \dots, X_q, \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}$. By the Phragmen-Lindelöf principle, we may take $\xi' > 0$ so small $B_W(x, \xi') \subseteq \Omega' \subset \mathcal{K}_1$, for all $x \in \mathcal{K}$. The balls $B_W(x, \delta)$ are metric balls and the topology induced by these balls on Ω is the usual topology. Let $\xi \in (0, \xi']$ be less than or equal to the Lebesgue number for the cover U_{x_1}, \dots, U_{x_R} of \mathcal{K}_1 , with respect to the metric associated to the balls $B_W(x, \delta)$. Then, $\forall x \in \mathcal{K}$, $\exists k \in \{1, \dots, R\}$ with $B_X(x, \xi) \subseteq B_W(x, \xi) = B_W(x, \xi) \cap \mathcal{K}_1 \subseteq U_{x_k}$, as desired. Also, $\bigcup_{x \in \mathcal{K}} B_X(x, \xi) \subseteq U_{x_1} \cup \dots \cup U_{x_R} \Subset \Omega$.

Let $\hat{c}_{j,k}^{l,x}$ be the functions from Proposition 9.1 (with this choice of m). Take $s_1 > 0$ so that $\forall r \in \{1, \dots, R\}$, $\hat{c}_{j,k}^{l,x_r} \in C^{\omega, s_1}(U_{x_r})$.

Take $x \in \mathcal{K}$ and $1 \leq r \leq R$ so that $B_X(x, \xi) \subseteq U_{x_r}$. Set $c_{j,k}^{l,x} := \hat{c}_{j,k}^{l,x_r}|_{B_X(x, \xi)}$. By Proposition 4.7, there exists $s \approx 1$ with $c_{j,k}^{l,x} \in C_X^{\omega, s}(B_X(x, \xi)) \subseteq C_X^{\omega, s}(U_{x_r}) \subseteq C^{\omega, s_1}(U_{x_r})$ and

$$\|c_{j,k}^{l,x}\|_{C_X^{\omega, s}(B_X(x, \xi))} \leq \|\hat{c}_{j,k}^{l,x_r}\|_{C_X^{\omega, s}(U_{x_r})} \lesssim \|\hat{c}_{j,k}^{l,x_r}\|_{C^{\omega, s_1}(U_{x_r})} \leq \max_{1 \leq r \leq R} \|\hat{c}_{j,k}^{l,x_r}\|_{C^{\omega, s_1}(U_{x_r})} \lesssim 1.$$

By the definition of $c_{j,k}^{l,x}$, we have $[X_j, X_k] = \sum_l c_{j,k}^{l,x} X_l$, for $x \in B_X(x, \xi)$, completing the proof. \square

Let $m, c_{j,k}^{l,x}, \xi, s$, and $(X_1, d_1), \dots, (X_q, d_q)$ be as in Lemma 9.6. We will prove Theorem 5.4 with this choice of m . For $\delta \in (0, 1]$, set $X_j^\delta := \delta^{d_j} X_j$, let X^δ denote the list $X_1^\delta, \dots, X_q^\delta$, and set

$$c_{j,k}^{l,x,\delta} := \begin{cases} \delta^{d_j+d_k-d_l} c_{j,k}^{l,x}, & \text{if } d_j + d_k \geq d_l, \\ 0, & \text{otherwise.} \end{cases}$$

Lemma 9.7. $[X_j^\delta, X_k^\delta] = \sum_l c_{j,k}^{l,x,\delta} X_l^\delta$ on $B_{X^\delta}(x, \xi)$ and

$$\sup_{x \in \mathcal{K}, \delta \in (0,1]} \left\| c_{j,k}^{l,x,\delta} \right\|_{C_{X^\delta}^{\omega,s}(B_{X^\delta}(x,\xi))} < \infty.$$

Proof. This follows immediately from Lemma 9.6, by multiplying (9.4) by $\delta^{d_j+d_k}$ and tracing through the definitions. \square

We pick $j_1 = j_1(x, \delta), \dots, j_{n_0(x)} = j_{n_0(x)}(x, \delta)$ as in Section 5.1.2 and we set

$$J_0 = J_0(x, \delta) := (j_1(x, \delta), \dots, j_{n_0(x)}(x, \delta)) \in \mathcal{I}(n_0(x), q) \in \mathcal{I}(n_0(x), q).$$

Let $X_{J_0}^\delta$ denote the list $X_{j_1(x,\delta)}^\delta, \dots, X_{j_{n_0(x)}(x,\delta)}^\delta$.

Lemma 9.8. *The conditions Theorem 3.10 hold uniformly when applied to $X_1^\delta, \dots, X_q^\delta$, at the base point x , for $x \in \mathcal{K}$ and $\delta \in (0, 1]$. Here we take ξ and $J_0 = J_0(x, \delta)$ as defined above. The corresponding map Φ from Theorem 3.10 is the map $\Phi_{x,\delta}$ defined in (5.4). In particular, any constant which is admissible in the sense of Theorem 3.10 can be taken independent of $x \in \mathcal{K}$ and $\delta \in (0, 1]$, when applied to $X_1^\delta, \dots, X_q^\delta$ at the base point x .*

Proof. We use $\xi > 0$ and $c_{j,k}^{l,x,\delta}$ from Lemma 9.7. The existence of δ_0 , independent of $x \in \mathcal{K}$ and $\delta \in (0, 1]$, as in the hypotheses of Theorem 3.10 follows directly from Lemma 7.22. The existence of $\eta > 0$, independent of $x \in \mathcal{K}$ and $\delta \in (0, 1]$, such that X^δ satisfies $\mathcal{C}(x, \eta, \Omega)$, $\forall x \in \mathcal{K}$ also follows from Lemma 7.22. With J_0 as above, we may take $\zeta \approx 1$ in Theorem 3.10. By Lemma 4.3 and Lemma 9.7, we have

$$\sup_{x \in \mathcal{K}, \delta \in (0,1]} \left\| c_{j,k}^{l,x,\delta} \right\|_{\mathcal{A}_{X_{J_0}^\delta(x,\delta)}^{x, \min\{s, \eta, \xi\}}} \leq \sup_{x \in \mathcal{K}, \delta \in (0,1]} \left\| c_{j,k}^{l,x,\delta} \right\|_{C_{X^\delta}^{\omega,s}(B_{X^\delta}(x,\xi))} < \infty.$$

Thus, we use $\min\{s, \eta, \xi\}$ in place of η in the hypotheses of Theorem 3.10. The result follows. \square

Lemma 9.8 shows that Theorem 3.10 applies to $X_1^\delta, \dots, X_q^\delta$ at the base point $x \in \mathcal{K}$, to yield constants $\chi, \eta_1, \xi_1, \xi_2 > 0$ as in that theorem (with $0 < \xi_2 \leq \xi_1 \leq \chi \leq \xi$), which are independent of $x \in \mathcal{K}$ and $\delta \in (0, 1]$.

Proof of Theorem 5.4 (vi). This follows directly from Theorem 3.10 (g) and (h). \square

Proof of Theorem 5.4 (v). By Theorem 3.10 we have $B_{X^\delta}(x, \xi_2) \subseteq \Phi_{x,\delta}(B^n(\eta_1)) \subseteq B_{X^\delta}(x, \xi)$. Taking $\xi_0 = \xi_2$, we have $B_{(X,d)}(x, \xi_0 \delta) \subseteq B_{X^\delta}(x, \xi_2)$ and since $\xi \leq 1$, $B_{X^\delta}(x, \xi) \subseteq B_{X^\delta}(x, 1) = B_{(X,d)}(x, \delta)$, completing the proof of (v). \square

Remark 9.9. In the proof of Theorem 5.4 (v), we chose ξ_0 so that $B_{(X,d)}(x, \xi_0 \delta) \subseteq B_{X^\delta}(x, \xi_2)$.

Proof of Theorem 5.4 (iii). This is contained in Theorem 3.10 (d) and (e), except that Theorem 3.10 (e) only guarantees that $\Phi_{x,\delta}$ is a C^2 diffeomorphism, not a real analytic diffeomorphism. However, using (vi) (which we have already proved), $Y_1^{x,\delta}, \dots, Y_q^{x,\delta}$ form a real analytic spanning set for $T_u B^{n_0(x)}(\eta_1)$ ($\forall u \in B^{n_0(x)}(\eta_1)$). Since $(\Phi_{x,\delta})_* Y_j^{x,\delta} = \delta^{d_j} X_j$ where X_1, \dots, X_q are real analytic on L_x , Lemma 7.25 shows $\Phi_{x,\delta}$ is real analytic. \square

Lemma 9.10. *There exists $r > 0$, independent of $x \in \mathcal{K}$, $\delta \in (0, 1]$, and functions $f_l^{x, \delta}$ ($1 \leq l \leq n_0(x)$) such that $\forall x \in \mathcal{K}$, $\delta \in (0, 1]$,*

$$\mathcal{L}_{X_{j_l(x, \delta)}^\delta} \nu_x = f_l^{x, \delta} \nu_x, \quad 1 \leq l \leq n_0(x),$$

where $f_l^{x, \delta} \in \mathcal{A}_{X_{j_0(x, \delta)}^\delta}^{x, r} \cap C(B_{X_{j_0(x, \delta)}^\delta}(x, \chi))$ with

$$\|f_l^{x, \delta}\|_{\mathcal{A}_{X_{j_0(x, \delta)}^\delta}^{x, r}}, \|f_l^{x, \delta}\|_{C(B_{X_{j_0(x, \delta)}^\delta}(x, \chi))} \lesssim 1, \quad 1 \leq l \leq n_0(x).$$

Proof. It is immediate to verify that the hypotheses of Proposition 8.2, when applied to the vector fields X^δ , hold uniformly for $x \in \mathcal{K}$, $\delta \in (0, 1]$. Thus, E-admissible constants in that theorem can be chosen independent of $x \in \mathcal{K}$, $\delta \in (0, 1]$. Because of this, the lemma is an immediate consequence of Proposition 8.2. \square

In light of Lemma 9.10, the assumptions of Theorem 3.14 and Corollary 3.15 hold, when applied to the vector fields X^δ and the density ν_x , at the base point x , uniformly for $x \in \mathcal{K}$ and $\delta \in (0, 1]$. Thus, any constant which is ν -admissible (or $0; \nu$ -admissible) in the sense of those results can be chosen independent of $x \in \mathcal{K}$, $\delta \in (0, 1]$.

Since ν_x is the induced Lebesgue density on an $n_0(x)$ -dimensional injectively immersed submanifold of \mathbb{R}^N , we have

$$\nu_x(Z_1 \wedge \cdots \wedge Z_{n_0(x)}) = \left| \det_{n_0(x) \times n_0(x)}(Z_1 | \cdots | Z_{n_0(x)}) \right|, \quad (9.5)$$

where $Z_1, \dots, Z_{n_0(x)}$ are vector fields tangent to L_x and $(Z_1 | \cdots | Z_{n_0(x)})$ denotes the $N \times n_0(x)$ matrix with columns given by $Z_1, \dots, Z_{n_0(x)}$.

Proof of Theorem 5.4 (iv). This follows from Theorem 3.14 and Corollary 3.15, using (9.5). \square

Proof of Theorem 5.4 (i). By Remark 9.9, Corollary 3.15, and (9.5), we have for $\delta \in (0, 1]$,

$$\begin{aligned} \nu_x(B_{(X, d)}(x, \xi_0 \delta)) &\leq \nu_x(B_{X^\delta}(x, \xi_2)) \approx \max_{k_1, \dots, k_{n_0(x)} \in \{1, \dots, q\}} \nu_x(\delta^{d_{k_1}} X_{k_1}, \dots, \delta^{d_{k_{n_0(x)}}} X_{k_{n_0(x)}}) \\ &\approx \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_\infty \approx \left| \det_{n_0(x) \times n_0(x)} (\xi_0 \delta)^d X(x) \right|_\infty, \end{aligned} \quad (9.6)$$

where the last \approx follows immediately from the formula for $|\det_{n_0(x) \times n_0(x)} \delta^d X(x)|_\infty$ and the fact that $\xi_0 \approx 1$. Similarly, since $\xi_2 \leq 1$, we have by Corollary 3.15 and (9.5),

$$\begin{aligned} \nu_x(B_{(X, d)}(x, \delta)) &\geq \nu_x(B_{X^\delta}(x, \xi_2)) \approx \max_{k_1, \dots, k_{n_0(x)} \in \{1, \dots, q\}} \nu_x(\delta^{d_{k_1}} X_{k_1}, \dots, \delta^{d_{k_{n_0(x)}}} X_{k_{n_0(x)}}) \\ &\approx \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_\infty \end{aligned} \quad (9.7)$$

Combining (9.6) and (9.7) completes the proof. \square

Proof of Theorem 5.4 (ii). This follows immediately from the formula given in Theorem 5.4 (i). \square

9.1 Multi-parameter geometries

The results in Section 5.1.2 concerned single-parameter sub-Riemannian geometries. Theorem 5.4 can be generalized to the setting of multi-parameter geometries with essentially the same proof. We outline these ideas in this section. Such multi-parameter geometries arise in applications: see [SS12, Str14].

Let V_1, \dots, V_r be real analytic vector fields defined on an open set $\Omega \subseteq \mathbb{R}^n$. Fix $\nu \in \mathbb{N}$ and to each V_j assign a formal degree $0 \neq e_j \in \mathbb{N}^\nu$. If Z has formal degree $e \in \mathbb{N}^\nu$, we assign to $[V_j, Z]$ the formal degree $e + e_j$. Fix $m \in \mathbb{N}$ a large integer and let $(X_1, d_1), \dots, (X_q, d_q)$ denote the finite list of vector fields with

ν -parameter formal degree $d_j \in \mathbb{N}^\nu$ with $|d_j|_\infty \leq m$. The results which follow are essentially independent of m , so long as m is chosen sufficiently large (depending on $(V_1, e_1), \dots, (V_r, e_r)$ and \mathcal{K}). For $\delta \in (0, 1]^\nu$, we let $\delta^d X$ denote the list of vector fields $\delta^{d_1} X_1, \dots, \delta^{d_q} X_q$. We sometimes identify $\delta^d X$ with the $n \times q$ matrix $(\delta^{d_1} X_1 | \dots | \delta^{d_q} X_q)$. As before, we set $B_{(X,d)}(x, \delta) := B_{\delta^d X}(x, 1)$, though now $\delta \in (0, 1]^\nu$.

As in Section 5.1.2, the involutive distribution generated by V_1, \dots, V_r foliates Ω into leaves, and we let L_x denote the leaf passing through x , and ν_x the induced Lebesgue density on L_x . $B_{(X,d)}(x, \delta)$ is an open subset of L_x .

For each $x \in \Omega$ set $n_0(x) := \dim \text{span}\{X_1(x), \dots, X_q(x)\}$. For each $x \in \Omega$, $\delta \in (0, 1]^\nu$, pick $j_1 = j_1(x, \delta), \dots, j_{n_0(x)} = j_{n_0(x)}(x, \delta)$ so that

$$\left| \det_{n_0(x) \times n_0(x)} \left(\delta^{d_{j_1}} X_{j_1}(x) | \dots | \delta^{d_{j_{n_0(x)}}} X_{j_{n_0(x)}}(x) \right) \right|_\infty = \left| \det_{n_0(x) \times n_0(x)} \delta^d X \right|_\infty.$$

For this choice of $j_1 = j_1(x, \delta), \dots, j_{n_0(x)} = j_{n_0(x)}(x, \delta)$ set (writing n_0 for $n_0(x)$):

$$\Phi_{x,\delta}(t_1, \dots, t_{n_0}) := \exp\left(t_1 \delta^{d_{j_1}} X_{j_1} + \dots + t_{n_0} \delta^{d_{j_{n_0}}} X_{j_{n_0}}\right)x.$$

Theorem 9.11. *Fix a compact set $\mathcal{K} \Subset \Omega$ and $x \in \mathcal{K}$, take m sufficiently large (depending on \mathcal{K} and $(V_1, e_1), \dots, (V_r, e_r)$), and define $(X_1, d_1), \dots, (X_q, d_q)$ as above. Define $n_0(x)$, ν_x , and $\Phi_{x,\delta}(t_1, \dots, t_{n_0})$ as above. We write $A \lesssim B$ for $A \leq CB$ where C is a positive constant which may depend on \mathcal{K} , but does not depend on the particular points $x \in \mathcal{K}$ and $u \in \mathbb{R}^{n_0(x)}$ under consideration, or on the scale $\delta \in (0, 1]^\nu$; we write $A \approx B$ for $A \lesssim B$ and $B \lesssim A$. There exist $\eta_0, \xi_0 \approx 1$ such that $\forall x \in \mathcal{K}$,*

(i) $\nu_x(B_{(X,d)}(x, \delta)) \approx \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_\infty, \forall \delta \in (0, 1]^\nu, |\delta| \leq \xi_0.$

(ii) $\nu_x(B_{(X,d)}(x, 2\delta)) \lesssim \nu_x(B_{(X,d)}(x, \delta)), \forall \delta \in (0, 1]^\nu, |\delta| \leq \xi_0/2.$

(iii) $\forall \delta \in (0, 1]^\nu, \Phi_{x,\delta}(B^{n_0(x)}(\eta_1)) \subseteq L_x$ is open and $\Phi_{x,\delta} : B^n(\eta_1) \rightarrow \Phi_{x,\delta}(B^n(\eta_1))$ is a real analytic diffeomorphism.

(iv) For $\delta \in (0, 1]^\nu$, define $h_{x,\delta}(t)$ on $B^{n_0(x)}(\eta_1)$ by $h_{x,\delta} \sigma_{\text{Leb}} = \Phi_{x,\delta}^* \nu_x$. Then, $h_{x,\delta}(t) \approx \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_\infty, \forall t \in B^{n_0(x)}(\eta_1)$, and there exists $s \approx 1$ with $\|h_{x,\delta}\|_{\mathcal{A}^{n_0(x),s}} \lesssim \left| \det_{n_0(x) \times n_0(x)} \delta^d X(x) \right|_\infty.$

(v) $B_{(X,d)}(x, \xi_0 \delta) \subseteq \Phi_{x,\delta}(B^{n_0(x)}(\eta_1)) \subseteq B_{(X,d)}(x, \delta), \forall \delta \in (0, 1]^\nu.$

(vi) For $\delta \in (0, 1]^\nu, x \in \mathcal{K}$, let $Y_j^{x,\delta} := \Phi_{x,\delta}^* \delta^{d_j} X_j$, so that $Y_j^{x,\delta}$ is a real analytic vector field on $B^{n_0(x)}(\eta_1)$. We have

$$\|Y_j^{x,\delta}\|_{\mathcal{A}^{n_0(x),\eta_1}(\mathbb{R}^n)} \lesssim 1, \quad 1 \leq j \leq q.$$

Finally, $Y_1^{x,\delta}(u), \dots, Y_q^{x,\delta}(u)$ span $T_u B^{n_0(x)}(\eta_1)$, uniformly in x, δ , and u , in the sense that

$$\max_{k_1, \dots, k_{n_0(x)} \in \{1, \dots, q\}} \inf_{u \in B^{n_0(x)}(\eta_1)} \left| \det \left(Y_{k_1}^{x,\delta}(u) | \dots | Y_{k_{n_0(x)}}^{x,\delta}(u) \right) \right| \approx 1.$$

Comments on the proof. The proof is nearly identical to the proof of Theorem 5.4; anywhere in the proof where one writes $d \leq e$, where now $d, e \in \mathbb{N}^\nu$, the inequality means $d_\mu \leq e_\mu, \forall 1 \leq \mu \leq \nu$. A main change needed is that the set \mathcal{S} consists of vector fields paired with formal degrees in \mathbb{N}^ν , instead of formal degrees in \mathbb{N} . To deal with this one needs to generalize Lemma 9.4 to deal with $\mathcal{S} \subset \mathcal{A}_n^n \times \mathbb{N}^\nu$; the same proof works by treating $t \in \mathbb{R}^\nu$ and using each degree as a multi-index. With these modifications it is straightforward to adapt the proof to yield this more general result. We leave the details to the interested reader. \square

References

- [FSC86] Charles L. Fefferman and Antonio Sánchez-Calle, *Fundamental solutions for second order subelliptic operators*, Ann. of Math. (2) **124** (1986), no. 2, 247–272. MR 855295
- [Gui08] Victor Guillemin, *Lecture notes*, <http://math.mit.edu/~arita/18.101/>, 2008, prepared by Ana Rita Pires, Accessed: 2017-05-02.
- [Her63] Robert Hermann, *On the accessibility problem in control theory*, Internat. Sympos. Nonlinear Differential Equations and Nonlinear Mechanics, Academic Press, New York, 1963, pp. 325–332. MR 0149402
- [Hör67] Lars Hörmander, *Hypoelliptic second order differential equations*, Acta Math. **119** (1967), 147–171. MR 0222474
- [Lee03] John M. Lee, *Introduction to smooth manifolds*, Graduate Texts in Mathematics, vol. 218, Springer-Verlag, New York, 2003. MR 1930091
- [Lob70] Claude Lobry, *Contrôlabilité des systèmes non linéaires*, SIAM J. Control **8** (1970), 573–605. MR 0271979
- [MM12] A. Montanari and D. Morbidelli, *Nonsmooth Hörmander vector fields and their control balls*, Trans. Amer. Math. Soc. **364** (2012), no. 5, 2339–2375. MR 2888209
- [Nag66] Tadashi Nagano, *Linear differential systems with singularities and an application to transitive Lie algebras*, J. Math. Soc. Japan **18** (1966), 398–404. MR 0199865
- [Nel59] Edward Nelson, *Analytic vectors*, Ann. of Math. (2) **70** (1959), 572–615. MR 0107176
- [Nic07] Liviu I. Nicolaescu, *Lectures on the geometry of manifolds*, second ed., World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007. MR 2363924
- [NSW85] Alexander Nagel, Elias M. Stein, and Stephen Wainger, *Balls and metrics defined by vector fields. I. Basic properties*, Acta Math. **155** (1985), no. 1-2, 103–147. MR 793239
- [SS12] Elias M. Stein and Brian Street, *Multi-parameter singular Radon transforms III: Real analytic surfaces*, Adv. Math. **229** (2012), no. 4, 2210–2238. MR 2880220
- [SS18] Betsy Stovall and Brian Street, *Coordinates adapted to vector fields: canonical coordinates*, Geom. Funct. Anal. **28** (2018), no. 6, 1780–1862. MR 3881835
- [Ste80] P. Stefan, *Integrability of systems of vector fields*, J. London Math. Soc. (2) **21** (1980), no. 3, 544–556. MR 577729
- [Str11] Brian Street, *Multi-parameter Carnot-Carathéodory balls and the theorem of Frobenius*, Rev. Mat. Iberoam. **27** (2011), no. 2, 645–732. MR 2848534
- [Str14] ———, *Multi-parameter Singular Integrals*, Annals of Mathematics Studies, vol. 189, Princeton University Press, Princeton, NJ, 2014.
- [Str20] ———, *Sub-Hermitian geometry and the quantitative Newlander-Nirenberg theorem*, Adv. Math. **368** (2020), 107137, 87. MR 4085140
- [Str21] ———, *Coordinates Adapted to Vector Fields II: Sharp Results*, in press., Amer. Jour. Math., arXiv:1808.04159, 2021.
- [Sus73] Héctor J. Sussmann, *Orbits of families of vector fields and integrability of distributions*, Trans. Amer. Math. Soc. **180** (1973), 171–188. MR 0321133

- [TW03] Terence Tao and James Wright, *L^p improving bounds for averages along curves*, J. Amer. Math. Soc. **16** (2003), no. 3, 605–638. MR 1969206
- [ZS75] Oscar Zariski and Pierre Samuel, *Commutative algebra. Vol. II*, Springer-Verlag, New York-Heidelberg, 1975, Reprint of the 1960 edition, Graduate Texts in Mathematics, Vol. 29. MR 0389876

University of Wisconsin-Madison, Department of Mathematics, 480 Lincoln Dr., Madison, WI, 53706

street@math.wisc.edu

MSC 2010: 58A30 (Primary), 32C05 and 53C17 (Secondary)