

ANISOTROPIC SOBOLEV EMBEDDINGS AND THE SPEED OF PROPAGATION FOR PARABOLIC EQUATIONS

FATMA GAMZE DÜZGÜN, SUNRA MOSCONI* AND VINCENZO VESPRI

ABSTRACT. We consider a quasilinear parabolic Cauchy problem with spatial anisotropy of orthotropic type and study the spatial localization of solutions. Assuming the initial datum is localized with respect to a coordinate having slow diffusion rate, we bound the corresponding directional velocity of the support along the flow. The expansion rate is shown to be optimal for large times.

CONTENTS

1.	Introduction	1
2.	Preliminaries	5
3.	Basic properties of solutions	11
4.	L^∞ -estimates	15
5.	Proof of the main Theorem	20
	Appendix A. An example á la Tikhonov	27
	References	29

1. INTRODUCTION

This paper deals with some properties of weak solutions to the Cauchy problem

$$(1.1) \quad \begin{cases} u_t - \operatorname{div} A(x, u, Du) = 0 & \text{in } S_T = \mathbb{R}^N \times]0, T[, \\ u(x, 0) = u_0. \end{cases}$$

The field $A : S_T \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ is only measurable and *anisotropic*, i.e.

$$(1.2) \quad \begin{cases} A_i(x, s, z) z_i \geq \Lambda^{-1} |z_i|^{p_i} \\ |A_i(x, s, z)| \leq \Lambda |z_i|^{p_i-1} \end{cases}$$

for some $\Lambda > 0$ and a suitable choice of $p_i > 1$, $i = 1, \dots, N$. In the case $p_i \equiv p$, $A_i(x, s, z) = |z_i|^{p-2} z_i$, the equation in (1.1) is known as the parabolic orthotropic p -Laplacian equation and, while its principal part is homogeneous, it fails to be isotropic.

Many materials, such as liquid crystals, wood or earth's crust usually present different diffusion rates along different directions. Moreover, in most of the physical phenomena involved in such media, finite speed of propagation of disturbances is a much more reasonable assumption than the usual infinite-speed one implied by linear equations. This effect can either be caused by additional absorption terms in the model equation or by the intrinsic diffusion rate of the medium. We are interested in this latter situation which, in the framework assumed here, consists in studying (1.1) when some of the p_i 's are greater than 2.

Let us discuss some features of equation (1.1) from the mathematical point of view.

2010 *Mathematics Subject Classification.* 35K15, 35K65, 35K92, 35R35.

Key words and phrases. Anisotropic equations, Finite speed of propagation, L^∞ -estimates, Non-uniqueness.

– *Regularity*

The anisotropy prescribed by (1.2) falls into the wider class of problems with *non-standard* growth condition. Even in the stationary case, a reasonable regularity theory for such equations requires a bound on the sparseness of the powers p_i . For elliptic equations driven by a principal part of the form (1.2), counterexamples to boundedness are given in the seminal papers [18, 20, 26]. Sufficient conditions for local boundedness are found in [7, 13, 17] and read

$$\bar{p} < N, \quad \max\{p_1, \dots, p_N\} < \bar{p}^*,$$

where

$$(1.3) \quad \frac{1}{\bar{p}} = \frac{1}{N} \sum_{i=1}^N \frac{1}{p_i}, \quad \bar{p}^* = \frac{N\bar{p}}{N - \bar{p}}$$

while in the limit case $\max\{p_1, \dots, p_N\} = \bar{p}^*$ it suffices to assume that $u \in L_{\text{loc}}^{\bar{p}^*}(\mathbb{R}^N)$ (which does not follow from the anisotropic Sobolev embedding, see [23]). With stronger conditions on the p_i 's and more smoothness assumptions on A , Hölder and Lipschitz regularity can be obtained and the literature on the elliptic case is huge. We refer to [27, Section 6] and [16] for a complete survey on the subject and related bibliography.

Regarding the parabolic degenerate (or singular) case of (1.1) much less is known. Indeed, the regularity theory going from linear to degenerate/singular equations in the parabolic framework is much more subtle than in the elliptic one. As far as we know, continuity of solutions to (1.1) is known only when the principal part is homogeneous, i.e. $p_i \equiv p$, thanks to the work of DiBenedetto and collaborators, see [12]. Nevertheless, some L^∞ -theory has been developed in the fully anisotropic case, see [9, 31] and the book [2].

– *Existence/Uniqueness*

In general, even for the heat equation, the Cauchy problem suffers heavy non-uniqueness phenomena, even for smooth and compactly supported initial data. This holds true also for the general isotropic model equation

$$(1.4) \quad \begin{cases} u_t = \operatorname{div}(|Du|^{p-2}Du) & \text{in } S_T, \\ u(x, 0) = u_0, \end{cases}$$

and examples can be found either in [6] or by coupling the results in [32] (see also [21]) for the porous media equation with [22]. In appendix A we will exhibit self-similar examples *à la* Tikhonov [30] (i.e., solutions $u \neq 0$ for $u_0 \equiv 0$) through a different and direct approach.

The main point of the theory is then to identify natural function classes ensuring unique solvability of the Cauchy problem. For linear parabolic equations this is a classical theme initiated by Widder [33] and developed by Aronsson [4], see the Introduction to [11, Ch XI] for a discussion. Roughly speaking, well-posedness holds for the class of functions (and initial data) obeying a suitable growth condition at infinity. Moreover, the growth condition is called optimal if any non-negative solution on S_T *a-priori* satisfies it.

The optimal well-posedness class was determined for (1.4) in [14, 15], following a number of related works on the porous medium equation. In the model case of the anisotropic parabolic equation a growth condition ensuring existence and uniqueness has been studied in [9, 10], however its optimality is still missing up to now. Indeed, the proofs of optimality usually involve some form of Harnack inequality in turn implying the prescribed growth, but in the anisotropic case no Harnack inequality is in general known.

A much smaller class where well-posedness holds true for (1.4) is the one of $L^p(S_T)$ solutions. Lions' method [25] indeed gives existence and uniqueness for the Cauchy problem with $L^2(\mathbb{R}^N)$ initial data, and eventually the method can be extended to problem (1.1) assuming monotonicity of $z \mapsto A(x, s, z)$ and $L^1(\mathbb{R}^N)$ initial data (or even u_0 being a finite measure). In this respect see [10] and the book [2]. However, at the level of generality dictated by the sole condition (1.2) in (1.1), no existence and/or uniqueness theory is available, whatever *a-priori* summability condition one imposes on the solution.

– *Description of the result and related problems*

We are interested in finding an optimal bounds for the directional speeds of propagation of the support of u_0 under the flow (1.1). Due to the changes in the diffusion coefficients with respect to various coordinates, we expect different velocities in each direction. It is well-known that, when the principal part is homogeneous, i.e. $p_i \equiv p$, finite energy solutions of (1.1) with initial condition $u_0 \in C_c^\infty(\mathbb{R}^N)$ preserve compactness of the support along the motion whenever $p > 2$ (slow-diffusion rate). In this case, an optimal bound from above to the speed of $\text{supp}(u(\cdot, t))$ has been determined in [1], [8] for a wide class of second order nonlinear parabolic equations, in [5] for higher order ones, while [28] deals with suitable parabolic systems. A typical result in this setting reads as follows: if $u \in L^p(S_T)$ solves (1.4) for some $p > 2$ and $\text{supp}(u_0) \subseteq B_{R_0}$, then for any $t \in]0, T]$

$$(1.5) \quad \text{supp}(u(\cdot, t)) \subseteq B_{2R_0+R(t)}, \quad R(t) = Ct^{\frac{1}{N(p-2)+p}} \|u_0\|_1^{\frac{p-2}{N(p-2)+p}},$$

where here and in the following we denote by $\|\cdot\|_p$ the $L^p(\mathbb{R}^N)$ norm, $p \in [1, \infty]$. In the anisotropic case, the model equation presents *locally* in space the same *qualitative* finite speed of propagation in each direction x_i of slow diffusion rate (i.e., for which $p_i > 2$). We refer to [2] and the literature therein for these kinds of results. What seems to be missing is a *global, quantitative* estimate of the aforementioned velocities, as prescribed e.g. by (1.5) in the homogeneous principal part scenario.

Since we will deal with local solutions to (1.1), the well-posedness issues described above suggest that we cannot expect finite speed of propagation to hold for generic local solutions, even for the parabolic p -Laplace equation (1.4). Indeed, in appendix A (see also Remark 2.2), self-similar solutions are constructed showing that finite speed of propagation fails at the global level. The main step consists in proving a non-uniqueness result for a merely Hölder continuous system of ODE with suitable initial data. In order to achieve this, we employ a sub-supersolution method which we think will be of some use for other problems, avoiding the full dynamical system analysis performed, e.g., in [21] or [6].

Despite the failure of finite speed of propagation in general, we will see that any local solution u of (1.1) actually possesses a so-called *branch* \tilde{u} (i.e., a solution of (1.1) coinciding with u on $\text{supp}(\tilde{u})$, see Remark 2.2 for some examples), exhibiting finite speed of propagation in each direction of slow diffusion. Finally, we will obtain a quantitative bound from above on the velocity, showing its optimality in a wide range of anisotropies.

For the purpose of this introduction we state our theorem in the case when all the p_i are greater than 2. We will also consider the case when some (but not all) of the p_i are less than or equal than 2, but the statement is less transparent, and we refer to the last section (namely, Theorem 5.3 therein) for further details.

Theorem 1.1. *Let \bar{p} be the harmonic mean of the p_i 's as per (1.3) and suppose that*

$$\bar{p} < N, \quad 2 < \min\{p_1, \dots, p_N\} \leq \max\{p_1, \dots, p_N\} < \bar{p} \left(1 + \frac{1}{N}\right).$$

Let u be a local weak solution of (1.1) in S_T under the growth conditions (1.2) with

$$u_0 \in L^2(\mathbb{R}^N), \quad \emptyset \neq \text{supp}(u_0) \subseteq [-R_0, R_0]^N.$$

Then there is a branch $\tilde{u} \neq 0$ of u such that

$$(1.6) \quad \text{supp}(\tilde{u}(\cdot, t)) \subseteq \prod_{j=1}^N [-R_j(t), R_j(t)],$$

for any $t < T$, where

$$(1.7) \quad R_j(t) = 2R_0 + Ct \frac{N(\bar{p}-p_j)+\bar{p}}{\lambda p_j} \|u_0\|_1^{\frac{\bar{p}}{p_j} \frac{p_j-2}{\lambda}}, \quad \lambda = N(\bar{p}-2) + \bar{p}.$$

Let us make some comments on the result.

(1) *Optimality of the estimated speed.*

First, the obtained velocities of propagation are optimal in each direction. Indeed, the Lebesgue measure of $\text{supp}(\tilde{u}(\cdot, t))$ is estimated for large times through (1.6) as

$$|\text{supp}(\tilde{u}(\cdot, t))| \leq Ct^{\sum_{i=1}^N \frac{N(\bar{p}-p_i)+\bar{p}}{\lambda p_i}} \|u_0\|_1^{\sum_{i=1}^N \frac{\bar{p}}{p_i} \frac{p_i-2}{\lambda}} = Ct^{\frac{N}{\lambda}} \|u_0\|_1^{\frac{N}{\lambda}(\bar{p}-2)}.$$

On the other hand, we will prove in Theorem 4.1 below the $L^\infty - L^1$ estimate

$$\|\tilde{u}(\cdot, t)\|_\infty \leq Ct^{-\frac{N}{\lambda}} \|u_0\|_1^{\frac{\bar{p}}{\lambda}}.$$

These two estimates imply

$$\|\tilde{u}(\cdot, t)\|_1 \leq \|\tilde{u}(\cdot, t)\|_\infty |\text{supp}(u(\cdot, t))| \leq C \|u_0\|_1.$$

Observe that if also $u_0 \geq 0$, the branch, being compactly supported, preserves the mass along the motion. Therefore, if any of the two estimates above were asymptotically (as $t \rightarrow +\infty$) less than optimal, this would contradict mass conservation.

(2) *Assumptions on the p_i 's.*

The main assumption on the parameters is the constraint

$$(1.8) \quad p_{\max} := \max\{p_1, \dots, p_N\} < \bar{p} \left(1 + \frac{1}{N}\right).$$

which is a purely anisotropic requirement, since for $p_i \equiv p$ the latter is always satisfied. The explicit power of t in (1.7) goes to zero as $p_{\max} \rightarrow \bar{p}(1 + 1/N)$, showing that condition (1.8) represents an actual threshold rather than being a technical assumption. It is possible that, for $p_{\max} > \bar{p}(1 + 1/N)$, zero-speed of propagation occurs in the corresponding direction (see [2, Ch. 5.3] for some instances of such phenomenon). In this regard, local boundedness of the solutions is ensured by the condition $\max\{2, p_{\max}\} < \bar{p}(1 + 2/N)$, hence the feasible range is $\bar{p}(1 + 1/N) < p_{\max} < \bar{p}(1 + 2/N)$.

(3) *Tools for the proof.*

A major tool in deriving $L^1 - L^\infty$ estimates for solutions of (1.1) is a well-known anisotropic Sobolev inequality, as proved for example in Troisi [29]. Unfortunately the latter alone seems to be insufficient to prove the optimal bound (1.7) and we will need

a variant of the anisotropic Sobolev inequality in the parabolic setting involving small powers of u . As the function $t \mapsto |t|^\alpha$ is not Lipschitz for $\alpha \in]0, 1[$, the Chain Rule turns out to be a quite delicate issue, however it can still be modified to prove the needed functional inequality in Proposition 2.3 below (see also Theorem 2.4 for its parabolic counterpart). We feel that these tools can have a wide range of applications to asymptotic and/or decay estimates for anisotropic equations.

(4) *Further investigation.*

Let us also mention that when $p_i \equiv p > 2$ (homogeneous principal part) and $u \geq 0$, the thesis of our main Theorem holds directly for u and there is no need to select the branch \tilde{u} . This is due to the fact that, (despite uniqueness is not guaranteed without additional monotonicity hypotheses) there are no local non-trivial and non-negative solutions with vanishing initial datum by [12, Proposition 18.1]. Indeed, even if the equation is nonlinear, subtracting a branch to a solution still gives a solution, possessing the aforementioned properties and thus being identically zero. We were not able to prove a similar statement in the anisotropic setting, since at present a weak Harnack inequality is still missing.

It is worth outlining that the optimality of disturbances' velocities, discussed in point (1) above, holds only when all the p_i 's lie in the degenerate range $p_i > 2$. While it is possible to obtain the same velocity bound when only *some* of the p_i are greater than 2 (again, see Theorem 5.3 in the last section), it seems less likely that for any remaining choices of the powers this bound is optimal. Indeed, it could happen that in some direction the diffusion is so fast that, in order to preserve the total mass, along the remaining ones the actual speed may be slower than what (1.7) dictates.

Finally, we did not tackle the question of extinction in finite time in the anisotropic setting, referring to [3] for a discussion of related results.

–*Outline of the paper*

In section 2 we gather some useful embedding of parabolic anisotropic Sobolev spaces. In section 3 we prove two types of energy inequalities and derive some contractivity estimates. Section 4 is devoted to *a-priori* L^∞ -estimates, both of global and local nature. In section 5 we first prove finite speed of propagation locally in time and space, then refine it to obtain a global result for a suitable branch. The final appendix is devoted to the Tikhonov example of non-uniqueness in the quasilinear setting.

2. PRELIMINARIES

Given $\mathbf{p} := (p_1, \dots, p_N) \in \mathbb{R}^N$, $\mathbf{p} > 1$, we let

$$\bar{p} := \left(\frac{1}{N} \sum_{i=1}^N \frac{1}{p_i} \right)^{-1},$$

and, assuming $\bar{p} < N$, we define

$$\bar{p}^* := \frac{N\bar{p}}{N - \bar{p}}.$$

Given $T > 0$ we set $S_T = \mathbb{R}^N \times]0, T[$ and for any rectangular domain $\Omega \subseteq \mathbb{R}^N$, $\Omega_T = \Omega \times]0, T[$.

We define the following spaces

$$\begin{aligned} W_0^{1,\mathbf{P}}(\Omega) &:= \{u \in W_0^{1,1}(\Omega) : D_i u \in L^{p_i}(\Omega)\} \\ W_{\text{loc}}^{1,\mathbf{P}}(\Omega) &:= \{u \in L_{\text{loc}}^1(\Omega) : D_i u \in L_{\text{loc}}^{p_i}(\Omega)\} \\ L^{\mathbf{P}}(0, T; W_0^{1,\mathbf{P}}(\Omega)) &:= \{u \in L^1(0, T; W_0^{1,1}(\Omega)) : D_i u \in L^{p_i}(\Omega_T)\} \\ L^{\mathbf{P}}(0, T; W_{\text{loc}}^{1,\mathbf{P}}(\Omega)) &:= \{u \in L^1(0, T; L_{\text{loc}}^1(\Omega)) : D_i u \in L^{p_i}(0, T; L_{\text{loc}}^{p_i}(\Omega))\} \\ L_{\text{loc}}^{\mathbf{P}}(0, T; W_{\text{loc}}^{1,\mathbf{P}}(\Omega)) &:= \{u \in L_{\text{loc}}^1(0, T; L_{\text{loc}}^1(\Omega)) : D_i u \in L_{\text{loc}}^{p_i}(0, T; L_{\text{loc}}^{p_i}(\Omega))\}. \end{aligned}$$

Let A be a measurable function obeying the growth conditions (1.2). By a local weak solution of

$$u_t = \text{div}(A(x, u, Du)) \quad \text{in } S_T$$

we mean a function $u \in C_{\text{loc}}^0(0, T; L_{\text{loc}}^2(\mathbb{R}^N)) \cap L_{\text{loc}}^{\mathbf{P}}(0, T; W_{\text{loc}}^{1,\mathbf{P}}(\mathbb{R}^N))$ such that for any $0 < t_1 < t_2 < T$ and any $\varphi \in C_{\text{loc}}^\infty(0, T; C_c^\infty(\mathbb{R}^N))$ it holds

$$(2.1) \quad \int u \varphi dx \Big|_{t_1}^{t_2} + \int_{t_1}^{t_2} \int -u \varphi_t + A(x, u, Du) \cdot D\varphi dx dt = 0,$$

where the integral is assumed henceforth to be on \mathbb{R}^N when no domain is specified. By density this actually holds for any $\varphi \in W_{\text{loc}}^{1,2}(0, T; L_{\text{loc}}^2(\mathbb{R}^N)) \cap L_{\text{loc}}^{\mathbf{P}}(0, T; W_0^{1,\mathbf{P}}(\Omega))$ for any rectangular $\Omega \Subset \mathbb{R}^N$.

For $T < +\infty$ and an initial data $u_0 \in L_{\text{loc}}^2(\mathbb{R}^N)$, we will consider the following Cauchy problem:

$$(2.2) \quad \begin{cases} u_t = \text{div}(A(x, u, Du)) & \text{in } S_T, \\ u(x, 0) = u_0(x), & x \in \mathbb{R}^N, \end{cases}$$

where u is a local solution of the equation such that, in addition, $u \in C^0(0, T; L_{\text{loc}}^2(\mathbb{R}^N)) \cap L^{\mathbf{P}}(0, T; W_{\text{loc}}^{1,\mathbf{P}}(\mathbb{R}^N))$ with $u(\cdot, 0) = u_0$. In other words, for u to solve (2.2) we require that the initial data is assumed strongly in $L_{\text{loc}}^2(\mathbb{R}^N)$ and that u has finite energy locally in space and globally in time. When $T = +\infty$, we say that u is a solution of (2.2) if it is so for any $T < +\infty$.

Definition 2.1. *Let u be a solution of (2.2). A branch \tilde{u} of u is a solution of (2.2) such that $\tilde{u} = u$ on $\text{supp}(\tilde{u})$.*

For any u , both 0 and u itself are obvious branches, which are the *trivial* ones (notice that 0 being a branch is due to the homogeneous structure assumed for the equation). Any other branch will be called *proper*, or *non trivial*.

Remark 2.2. *Examples of solutions having proper branches can be constructed simply by solving (in the finite energy space) the Cauchy problem for the p -Laplacian with initial datum φ being supported in two distant balls B_1 and B_2 , nontrivially in both. If $p > 2$, the finite speed of propagation provides us with a solution such that, up to some time $T > 0$, have two distinct branches, namely the ones starting from the initial datum $\varphi \chi_{B_i}$, $i = 1, 2$.*

More interesting examples are constructed in appendix A, where it is shown that for any $\beta > 0$ the slow-diffusion parabolic p -Laplacian equation has nontrivial solutions supported in $\{(x, t) : x \geq t^{-\beta} > 0\}$. The latter clearly have zero initial datum, are sign-changing and reproduce the Tikhonov phenomenon of non-uniqueness in the quasilinear setting.

Let now $\varphi \in C_c^\infty(\mathbb{R})$ be such that $\text{supp}(\varphi) \subseteq \{x < 0\}$ and let \tilde{v} be the unique energy solution of $u_t = \Delta_p u$ having φ as initial datum. Then, by finite speed of propagation, the function

$$v = \begin{cases} u & \text{on } \{x > 0\}, \\ \tilde{v} & \text{on } \{x \leq 0\}, \end{cases}$$

solves the equation in $\mathbb{R} \times]0, T]$ but, despite having compactly supported initial datum, has infinite support for any $t > 0$. In this case, selecting the branch \tilde{v} of v is necessary in order to recover finite speed of propagation of the support.

The following is a useful modification of the classical anisotropic Sobolev embedding by Troisi [29].

Proposition 2.3 (Sobolev embedding). *Let $\Omega \subseteq \mathbb{R}^N$ be a rectangular domain, $\bar{p} < N$ and $\alpha_i > 0$, $i = 1, \dots, N$. Then there exists a constant $C = C(N, \mathbf{p}, \alpha) > 0$ such that*

$$(2.3) \quad \|u\|_{L^{p_\alpha^*}(\Omega)} \leq C \prod_{i=1}^N \|D_i |u|^{\alpha_i}\|_{L^{p_i}(\Omega)}^{\frac{1}{\alpha_i}}$$

for all $u \in W_0^{1, \mathbf{p}}(\Omega)$, where

$$p_\alpha^* = \bar{p}^* \frac{\tilde{\alpha}}{N}, \quad \tilde{\alpha} = \sum_{i=1}^N \alpha_i.$$

Proof. We can suppose that the right-hand side is finite and, by a monotone convergence argument, that u and Ω are bounded.

First we claim that when $\alpha > 0$, $p \geq 1$ and $u \in W_0^{1,1}(\Omega) \cap L^\infty(\Omega)$ it holds

$$(2.4) \quad \int |D_i |u|^\alpha|^p dx = \alpha \int |u|^{\alpha-1} |D_i |u||^p dx,$$

whenever the left-hand side is finite, where we set

$$(2.5) \quad |u|^{\alpha-1} |D_i |u|| = 0 \quad \text{a.e. on } D_i u = 0.$$

This readily follows when $\alpha \geq 1$ from the boundedness of u and the classical Chain Rule for Sobolev functions. In the case $\alpha \in]0, 1[$ first observe that, since the left-hand side of (2.4) is finite, $D_i |u|^\alpha \equiv 0 \equiv D_i u$ a.e. on $\{u = 0\}$. Therefore it suffices to show that the weak derivatives of $|u|^\alpha$ and $|u|$ obey

$$(2.6) \quad D_i |u|^\alpha = \alpha |u|^{\alpha-1} D_i |u|, \quad \text{a.e. on } \{u \neq 0\}.$$

By the absolute continuity on lines characterization of Sobolev functions [25, Theorem 10.35], we can choose a representative such that for a.e. line l parallel to e_i , both¹ the restrictions of $|u|^\alpha$ and $|u|$ belong to $AC(l)$, with corresponding (classical) derivatives coinciding \mathcal{L}^1 a.e. with the i -th weak derivative. Moreover, $|u|^\alpha = f(|u|)$ with $f(t) := |t|^\alpha \in AC_{\text{loc}}(\mathbb{R})$ mapping

¹That this is true follows by suitably modifying the proof of [25, Theorem 10.35]: it suffices to start with $v_n \in C_c^\infty(\mathbb{R}^N)$ such that $D_i v_n \rightarrow D_i |u|^\alpha$ in $L^p(\mathbb{R}^N)$ and then observe that $D_i v_n^{1/\alpha} \rightarrow D_i |u|$ in $L^p(\mathbb{R}^N)$ as well, by the Lip_{loc} character of $t \mapsto |t|^{1/\alpha}$.

null-measure sets to null-measure sets. Therefore [25, Theorem 3.44] applies, giving, for \mathcal{L}^{N-1} a.e. x such that $x_i = 0$

$$\frac{d}{dt}|u|^\alpha(x + t e_i) = \alpha|u|^{\alpha-1}(x + t e_i) \frac{d}{dt}|u|(x + t e_i)$$

for \mathcal{L}^1 a.e. t such that $u(x + t e_i) \neq 0$. By Fubini's theorem and the absolute continuity on lines characterization of Sobolev functions, this shows (2.6) and thus (2.4).

Let us fix from now on a representative for which $|u|^\alpha \in AC(l)$ for almost every line parallel to e_i , and define $v_i = |u|^{\alpha_i}$. By the previous discussion, the classical partial derivative and the weak derivative coincide \mathcal{L}^1 a.e. on any line, with (2.6) holding as long as we assume (2.5). For $i = 1, \dots, N$ denote

$$\int_{\mathbb{R}_i} v(x) dt = \int_{\mathbb{R}} v(x + t e_i) dt.$$

Consider N parameters $s_i \geq 1$, $i = 1, \dots, N$ and observe that for any $i = 1, \dots, N$ and a.e. $x \in \mathbb{R}^N$

$$v_i^{s_i}(x) \leq s_i \int_{\mathbb{R}_i} v_i^{(s_i-1)}(x) |D_i v_i|(x) dt \leq s_i \left(\int_{\mathbb{R}_i} v_i^{(s_i-1)p'_i}(x) dt \right)^{\frac{1}{p'_i}} \left(\int_{\mathbb{R}_i} |D_i v_i|^{p_i}(x) dt \right)^{\frac{1}{p_i}}.$$

Therefore

$$\prod_{i=1}^N v_i^{\frac{s_i}{N-1}}(x) \leq C \prod_{i=1}^N \left(\int_{\mathbb{R}_i} v_i^{(s_i-1)p'_i}(x) dt \right)^{\frac{1}{(N-1)p'_i}} \left(\int_{\mathbb{R}_i} |D_i v_i|^{p_i}(x) dt \right)^{\frac{1}{(N-1)p_i}}$$

We successively integrate with respect to each variable x_j , observing that, at each step, the j -th factor of the product in the right-hand side does not depend on x_j . Regarding the other factors, it holds

$$\sum_{i \neq j} \frac{1}{(N-1)p'_i} + \frac{1}{(N-1)p_i} = 1,$$

so that at each step we can apply Hölder's inequality to obtain after N integrations

$$\int \prod_{i=1}^N v_i^{\frac{s_i}{N-1}} dx \leq C \prod_{i=1}^N \left(\int v_i^{(s_i-1)p'_i} dx \right)^{\frac{1}{(N-1)p'_i}} \left(\int |D_i v_i|^{p_i} dx \right)^{\frac{1}{(N-1)p_i}}.$$

i.e.

$$\int |u|^{\frac{1}{N-1} \sum_{i=1}^N \alpha_i s_i} dx \leq C \prod_{i=1}^N \left(\int |u|^{\alpha_i (s_i-1) p'_i} dx \right)^{\frac{1}{(N-1)p'_i}} \left(\int |D_i |u|^{\alpha_i}|^{p_i} dx \right)^{\frac{1}{(N-1)p_i}}.$$

Finally, the system in the $N+1$ variables s_j, q

$$\alpha_j (s_j - 1) p'_j = q, \quad j = 1, \dots, N, \quad \frac{1}{N-1} \sum_{i=1}^N \alpha_i s_i = q$$

is solvable with $q = p_\alpha^*$ (and $s_j \geq 1$), so that simple algebraic manipulations lead to (2.3). \square

When $\alpha_i \equiv 1$ we let $p_\alpha^* = \bar{p}^*$.

Theorem 2.4 (Parabolic Anisotropic Sobolev embedding). *Let $\Omega \subseteq \mathbb{R}^N$ be a rectangular domain, $\bar{p} < N$, $\alpha_i > 0$ for $i = 1, \dots, N$ and $\sigma \in [1, p_\alpha^*]$. For any $\theta \in [0, \frac{\bar{p}}{\bar{p}^*}]$ define*

$$q = q(\theta, \mathbf{p}, \alpha) = \theta p_\alpha^* + \sigma(1 - \theta).$$

Then there exists a constant $C = C(N, \mathbf{p}, \alpha, \theta, \sigma) > 0$ such that

$$(2.7) \quad \iint_{\Omega_T} |u|^q dx dt \leq C T^{1-\theta \frac{\bar{p}^*}{\bar{p}}} \left(\sup_{t \in [0, T]} \int_{\Omega} |u|^\sigma(x, t) dx \right)^{1-\theta} \prod_{i=1}^N \left(\iint_{\Omega_T} |D_i |u|^{\alpha_i}|^{p_i} dx dt \right)^{\frac{\theta \bar{p}^*}{N p_i}}$$

for any $u \in L^1(0, T; W_0^{1,1}(\Omega))$.

Proof. For a.e. $t \in [0, T]$, apply Hölder's inequality as

$$\int |u|^q(x, t) dx \leq \left(\int |u|^\sigma(x, t) dx \right)^{1-\theta} \left(\int u^{p_\alpha^*}(x, t) dx \right)^\theta$$

and through (2.3) deduce

$$\int |u|^q(x, t) dx \leq C \left(\int |u|^\sigma(x, t) dx \right)^{1-\theta} \prod_{i=1}^N \left(\int |D_i |u|^{\alpha_i}|^{p_i}(x, t) dx \right)^{\frac{\theta \bar{p}^*}{N p_i}}.$$

Finally we integrate in the t -variable and use Hölder's inequality in time as

$$\int_0^T \prod_{i=1}^N f_i^{\gamma_i} dt \leq T^{1-\frac{1}{r}} \sup_{t \in [0, T]} f_0^{\gamma_0} \left(\int_0^T \prod_{i=1}^N f_i^{\gamma_i r} dt \right)^{\frac{1}{r}} \leq T^{1-\frac{1}{r}} \sup_{t \in [0, T]} f_0^{\gamma_0} \prod_{i=1}^N \left(\int_0^T f_i dt \right)^{\gamma_i}$$

valid for

$$\sum_{i=1}^N r \gamma_i = 1, \quad r \geq 1, \quad \gamma_i > 0.$$

In the previous formula we set

$$\gamma_0 = 1 - \theta, \quad f_0(t) = \int |u|^\sigma(x, t) dx$$

and for $i = 1, \dots, N$

$$\gamma_i = \frac{\theta \bar{p}^*}{N p_i}, \quad r = \frac{\bar{p}}{\bar{p}^*} \frac{1}{\theta} \geq 1, \quad f_i(t) = \int |D_i |u|^{\alpha_i}|^{p_i}(x, t) dx$$

to get the claim. \square

Remark 2.5. *In the isotropic case, the previous theorem ensures an analogous local summability estimate without the assumption that u vanishes outside Ω , just by adding an L^1 term to the right-hand side. Unfortunately, this is no longer true in the anisotropic setting, and counter-examples to the corresponding embeddings are known (see [19, 23] in the non-parabolic case). More precisely, under the previous assumptions on the parameters, it may happen that $L^{\mathbf{p}}(0, T; W_{\text{loc}}^{1, \mathbf{p}}(\mathbb{R}^N)) \cap L^\infty(0, T; L_{\text{loc}}^\sigma(\mathbb{R}^N))$ fails to be contained in $L^q(0, T; L_{\text{loc}}^q(\mathbb{R}^N))$ for q given above. In order to remove the zero boundary condition, one is either forced to assume a-priori a suitable degree of summability, or to further constrain the location of the p_i 's.*

The following result deals with the problem mentioned in the previous remark when $\alpha_i \equiv 1$.

Theorem 2.6. *Let $1 \leq p_1 \leq \dots \leq p_N$, $\bar{p} < N$ and for any $\sigma \in [1, \bar{p}^*]$ define the critical parabolic exponent*

$$\bar{p}_\sigma = \bar{p} \left(1 + \frac{\sigma}{N}\right).$$

Then the embedding

$$(2.8) \quad L^{p_N}(0, T; L_{\text{loc}}^{p_N}(\mathbb{R}^N)) \cap L^{\mathbf{p}}(0, T; W_{\text{loc}}^{1, \mathbf{p}}(\mathbb{R}^N)) \cap L^\infty(0, T; L_{\text{loc}}^\sigma(\mathbb{R}^N)) \hookrightarrow L^{\bar{p}_\sigma}(0, T; L_{\text{loc}}^{\bar{p}_\sigma}(\mathbb{R}^N))$$

holds true. Moreover, under the assumption

$$(2.9) \quad \bar{p}_\sigma > p_N = \max\{p_1, \dots, p_N\}$$

we directly have

$$L^{\mathbf{p}}(0, T; W_{\text{loc}}^{1, \mathbf{p}}(\mathbb{R}^N)) \cap L^\infty(0, T; L_{\text{loc}}^\sigma(\mathbb{R}^N)) \hookrightarrow L^{\bar{p}_\sigma}(0, T; L_{\text{loc}}^{\bar{p}_\sigma}(\mathbb{R}^N)).$$

Proof. Apply the previous proposition with $\alpha_i \equiv 1$, $\theta = \bar{p}/\bar{p}^*$ to $u\eta$ for $\eta \in C_c^\infty(\mathbb{R}^N)$, $\eta \equiv 1$ on an arbitrary cube $K \subseteq \mathbb{R}^N$. With the notations of our statement, it holds $q = \bar{p}_\sigma$ and we get

$$\int_0^T \int_K |u|^{\bar{p}_\sigma} dx dt \leq \left(\sup_{t \in [0, T]} \int |u\eta|^\sigma(x, t) dx \right)^{1-\theta} \prod_{i=1}^N \left(\int_0^T \int |D_i(u\eta)|^{p_i} dx dt \right)^{\frac{\bar{p}}{N p_i}}.$$

The first factor on the right is finite by assumption, while for the others we have

$$\int_0^T \int |D_i(u\eta)|^{p_i} dx dt \leq C \int_0^T \int_{K'} |D_i u|^{p_i} + |u|^{p_i} dx dt,$$

where K' is another cube such that $\text{supp}(\eta) \subseteq K'$ and $C = C(\eta)$. To prove the first embedding (2.8), it suffices to observe that, by the ordering of the p_i and Hölder's inequality, $L^{p_i}(0, T; L_{\text{loc}}^{p_i}(\mathbb{R}^N)) \hookrightarrow L^{p_N}(0, T; L_{\text{loc}}^{p_N}(\mathbb{R}^N))$ for any $i = 1, \dots, N$.

Let us prove the second embedding under assumption (2.9). On vectors $\mathbf{q} \in \mathbb{R}^N$ we consider the component-wise partial ordering

$$(q_1, \dots, q_N) \geq (r_1, \dots, r_N) \iff q_i \geq r_i \quad \text{for all } i = 1, \dots, N.$$

In turn, this naturally defines a lattice structure on \mathbb{R}^N , with \wedge denoting the minimum operation. By abuse of notation we will say that, for $\lambda \in \mathbb{R}$ it holds $\mathbf{q} \geq \lambda$ to mean $\mathbf{q} \geq (\lambda, \dots, \lambda)$.

Define by recursion the following sequence $\mathbf{p}^n = (p_1^n, \dots, p_N^n) \in \mathbb{R}^N$:

$$\begin{cases} \mathbf{p}^1 = (p_1, \dots, p_N) \\ \mathbf{p}^{n+1} = \mathbf{p}^n \wedge (q^n, \dots, q^n), & q^n := \frac{N+\sigma}{\sum_1^N \frac{1}{p_i^n}}. \end{cases}$$

We claim that if (2.9) holds, then

$$(2.10) \quad q^n \geq p_N \quad \text{for sufficiently large } n.$$

Let us postpone the proof of (2.10) momentarily and show how this implies $u \in L^{p_N}(0, T; L_{\text{loc}}^{p_N}(\mathbb{R}^N))$.

For $n = 1$ the standard isotropic parabolic Sobolev embedding ensures that $u \in L^{q^1}(0, T; L_{\text{loc}}^{q^1}(\mathbb{R}^N))$.

Suppose that $u \in L^{q^n}(0, T; L_{\text{loc}}^{q^n}(\mathbb{R}^N))$ for some $q_n \leq p_N$. Then, since $\mathbf{p}^{n+1} \leq \mathbf{p}$, Hölder's inequality ensures $u \in L^{\mathbf{p}^{n+1}}(0, T; W_{\text{loc}}^{1, \mathbf{p}^{n+1}}(\mathbb{R}^N))$ and the embedding (2.8) with vector \mathbf{p}^{n+1} implies $u \in L^{q^{n+1}}(0, T; L_{\text{loc}}^{q^{n+1}}(\mathbb{R}^N))$. Therefore in a finite number of steps we get $u \in L^{\bar{p}_\sigma}(0, T; L_{\text{loc}}^{\bar{p}_\sigma}(\mathbb{R}^N))$ by the claim. Clearly, this process also proves the second stated embedding.

We next prove (2.10). Since q^n is a fixed multiple of the harmonic mean of \mathbf{p}^n and $q^1 \geq p_1 =: q^0$, it follows from

$$q^n \geq q^{n-1} \quad \Rightarrow \quad \mathbf{p}^{n+1} \geq \mathbf{p}^n \quad \Rightarrow \quad q^{n+1} \geq q^n$$

that $\{q^n\}$ is non-decreasing. Suppose there exists the smallest integer $1 \leq h < N$ such that $q^n \leq p_{h+1}$ for all $n \geq 0$, and let $q = \lim_n q^n \leq p_{h+1}$. Then we infer

$$\lim_n q^n = \lim_n q^{n+1} \quad \Leftrightarrow \quad q = \frac{N + \sigma}{\sum_1^h \frac{1}{p_i} + \frac{N-h}{q}} \quad \Leftrightarrow \quad q = r_h,$$

where we defined for all $k = 1, \dots, N$

$$r_k := \frac{k + \sigma}{\sum_1^k \frac{1}{p_i}}.$$

Notice that

$$r_k \leq p_{k+1} \quad \Leftrightarrow \quad \frac{k + \sigma}{p_{k+1}} \leq \sum_{i=1}^k \frac{1}{p_i}$$

so that, adding $1/p_{k+1}$ to both sides, rearranging and using the monotonicity of the p_i , we get

$$r_k \leq p_{k+1} \quad \Rightarrow \quad r_{k+1} \leq p_{k+2}.$$

Since $q = r_h \leq p_{h+1}$ by assumption, we eventually get by induction $r_{N-1} \leq p_N$, which is equivalent to say

$$p_N \geq \frac{N + \sigma}{\sum_1^N \frac{1}{p_i}} = \bar{p} \left(1 + \frac{\sigma}{N}\right) = \bar{p}_\sigma.$$

This contradicts (2.9), proving the claim (2.10) and the theorem. \square

The following is a straightforward generalization, needed to deal with the anisotropic growth, of [11, Ch. I, Lemma 4.1]. We omit its elementary proof.

Lemma 2.7. *Let $\beta_i > 0$ for $i = 1, \dots, N$ and suppose $Z_n \geq 0$ satisfies*

$$Z_{n+1} \leq C b^n \frac{1}{N} \sum_{i=1}^N Z_n^{1+\beta_i}$$

for some $C, b > 1$. Then letting $\beta = \min\{\beta_1, \dots, \beta_N\}$,

$$Z_0 \leq \frac{1}{C^{\frac{1}{\beta}} b^{\frac{1}{\beta^2}}} \quad \Rightarrow \quad Z_n \rightarrow 0.$$

3. BASIC PROPERTIES OF SOLUTIONS

The following energy estimate can be proved through a standard Steklov averaging procedure.

Lemma 3.1 (Energy Inequality). *Let u be a weak solution of (2.1). There exists a constant $C = C(N, \mathbf{p}, \Lambda) > 0$ such that for any $\eta \geq 0$ of the form*

$$(3.1) \quad \eta(x, t) = \prod_{i=1}^N \eta_i^{p_i}(x_i, t), \quad \eta_i \in C^\infty(0, T; C_c^\infty(\mathbb{R}))$$

and $0 < t_1 < t_2 < T$, $k \in \mathbb{R}$, it holds

$$(3.2) \quad \int (u - k)_+^2(x, t) \eta(x, t) dx \Big|_{t_1}^{t_2} + \frac{1}{C} \sum_{i=1}^N \int_{t_1}^{t_2} \int |D_i((u - k)_+ \eta)|^{p_i} dx dt \leq \\ \int_{t_1}^{t_2} \int (u - k)_+^2 |\eta_t| dx dt + C \sum_{i=1}^N \int_{t_1}^{t_2} \int (u - k)_+^{p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} dx dt.$$

Proof. Modulo a Steklov averaging process we can suppose that u_t exists and that (2.1) holds for the test function $\varphi := \eta(u - k)_+$. In this case (2.1) becomes

$$(3.3) \quad \int_{t_1}^{t_2} \int u_t \varphi dx dt + \int_{t_1}^{t_2} \int A(x, u, Du) \cdot D\varphi dx = 0.$$

The first integral is estimated, for the chosen test function, as

$$(3.4) \quad \int_{t_1}^{t_2} \int u_t \varphi dx dt = \int_{t_1}^{t_2} \int \left(\frac{(u - k)_+^2}{2} \right)_t \eta dx dt \\ = \int \frac{(u - k)_+^2}{2} \eta dx \Big|_{t_1}^{t_2} - \int_{t_1}^{t_2} \int \frac{(u - k)_+^2}{2} \eta_t dx dt.$$

For the second integral, let

$$\hat{\eta}_i = \frac{\eta}{\eta_i^{p_i}},$$

which does not depend on x_i , and use (1.2) to estimate

$$A(x, u, Du) \cdot D(\eta(u - k)_+) \geq \Lambda^{-1} \eta \sum_{i=1}^N |D_i(u - k)_+|^{p_i} - \Lambda(u - k)_+ \sum_{i=1}^N |D_i(u - k)_+|^{p_i-1} |D_i \eta|$$

The last term is estimated through Young inequality as

$$(3.5) \quad (u - k)_+ |D_i(u - k)_+|^{p_i-1} |D_i \eta| = p_i (u - k)_+ |D_i(u - k)_+|^{p_i-1} \eta_i^{p_i-1} |D_i \eta_i| \hat{\eta}_i \\ \leq \hat{\eta}_i \left(\frac{1}{2\Lambda} \eta_i^{p_i} |D_i(u - k)_+|^{p_i} + C(\mathbf{p}, \Lambda) (u - k)_+^{p_i} |D_i \eta_i|^{p_i} \right).$$

Using the definition of $\hat{\eta}_i$ we get

$$(3.6) \quad \hat{\eta}_i \eta_i^{p_i} = \eta, \quad \hat{\eta}_i |D_i \eta_i|^{p_i} = |D_i \eta^{\frac{1}{p_i}}|^{p_i},$$

therefore

$$(3.7) \quad \int_{t_1}^{t_2} \int A(x, u, Du) \cdot D\varphi dx \geq \frac{\Lambda^{-1}}{2} \sum_{i=1}^N \int_{t_1}^{t_2} \int |D_i(u - k)_+|^{p_i} \eta dx dt \\ - C \int_{t_1}^{t_2} \sum_{i=1}^N \int (u - k)_+^{p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} dx dt.$$

Finally, since $p_i > 1$ for all i and $\eta_i \in [0, 1]$, observe that for any $v \geq 0$,

$$|D_i(v \eta)|^{p_i} \leq c(\mathbf{p}) (|D_i v|^{p_i} \eta^{p_i} + v^{p_i} \hat{\eta}_i^{p_i} \eta_i^{p_i-1} |D_i \eta_i|^{p_i}) \leq c(\mathbf{p}) (|D_i v|^{p_i} \eta + v^{p_i} \hat{\eta}_i |D_i \eta_i|^{p_i}) \\ \leq c(\mathbf{p}) \left(|D_i v|^{p_i} \eta + v^{p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} \right)$$

so that when $v = (u - k)_+$

$$\begin{aligned} \int_{t_1}^{t_2} \int A(x, u, Du) \cdot D\varphi \, dx &\geq \frac{\Lambda^{-1}}{2c} \int_{t_1}^{t_2} \sum_{i=1}^N \int |D_i((u - k)_+ \eta)|^{p_i} \, dx \, dt \\ &\quad - \frac{C}{c} \int_{t_1}^{t_2} \sum_{i=1}^N \int (u - k)_+^{p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} \, dx \, dt \end{aligned}$$

and the claim follows inserting this last inequality and (3.4) into (3.3). \square

In general, we cannot always ensure that the right-hand side in (3.2) is finite. This can either be required as additional condition on the solution, or it can be deduced from Theorem 2.6 whenever $\max\{p_1, \dots, p_N\} < \bar{p}_2$, being a local solution *a priori* in $L^{\mathbf{p}}(0, T; W_{\text{loc}}^{1, \mathbf{p}}(\mathbb{R}^N)) \cap L^\infty(0, T; L_{\text{loc}}^2(\mathbb{R}^N))$.

Corollary 3.2. *Let $u \in \cap_{i=1}^N L^{p_i}(S_T)$ solve (2.2) in S_T for $u_0 \in L^2(\mathbb{R}^N)$. Then $u \in L^\infty(0, T; L^2(\mathbb{R}^N))$,*

$$(3.8) \quad \|u(\cdot, t)\|_2 \leq \|u_0\|_2$$

and for any $0 < t_1 < t_2 < T$, $\psi \in \text{Lip}([0, T]; \mathbb{R})$, $\psi \geq 0$ and $k \in \mathbb{R}$ it holds

(3.9)

$$\int (u - k)_+^2(x, t) \psi(t) \, dx \Big|_{t_1}^{t_2} + \frac{1}{C} \sum_{i=1}^N \int_{t_1}^{t_2} \int |D_i(u - k)_+|^{p_i} \psi \, dx \, dt \leq \int_{t_1}^{t_2} \int (u - k)_+^2 |\psi_t| \, dx \, dt.$$

Proof. To get (3.8) let $0 \leq \eta \leq 1$ be as in (3.1), independent on t and such that

$$(3.10) \quad \text{supp}(\eta_i) \subseteq [-2R, 2R], \quad \eta_i \equiv 1 \text{ on } [-R, R], \quad |D_i \eta^{\frac{1}{p_i}}| \leq \frac{C}{R}.$$

With such η , apply the (3.2) to both u and $-u$ for $k = 0$, adding the corresponding inequalities. Since $u \in C^0([0, T]; L_{\text{loc}}^2(\mathbb{R}^N))$ we can let $t_1 \rightarrow 0$ to get

$$\int u^2(x, t_2) \eta \, dx \leq \int u_0^2 \, dx + C \sum_{i=1}^N \int_0^{t_2} \int |u|^{p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} \, dx \, dt.$$

We let $R \rightarrow +\infty$, apply Fatou's lemma on the left-hand side, and observe that the right-hand side vanishes by the last property in (3.10) and $u \in \cap L^{p_i}(S_T)$. Therefore (3.8) is proved. To prove (3.9), repeat the proof of (3.2) with $\varphi = \eta \psi$ up to (3.7), obtaining

$$\begin{aligned} \int (u - k)_+^2(x, t) \eta(x, t) \psi(t) \, dx \Big|_{t_1}^{t_2} + \frac{1}{C} \sum_{i=1}^N \int_{t_1}^{t_2} \int |D_i((u - k)_+)|^{p_i} \eta \psi \, dx \, dt \leq \\ \int_{t_1}^{t_2} \int (u - k)_+^2 \eta |\psi_t| \, dx \, dt + C \sum_{i=1}^N \int_{t_1}^{t_2} \int (u - k)_+^{p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} \psi \, dx \, dt. \end{aligned}$$

Letting $R \rightarrow +\infty$ cancels the last term on the right as before, while we apply Fatou on the terms involving the spatial derivatives of u and dominated convergence to the others, to obtain (3.9). \square

A useful variant of the energy inequality (3.2) is the following.

Lemma 3.3 (Energy inequality 2). *Let $F \in C^{1,1}(\mathbb{R})$ with $|F'| \leq M$ and $M \geq F''(t) > 0$ for a.e. $t \in \mathbb{R}$ and for suitable $M \in \mathbb{R}$. If u is a local weak solution to (2.1) and η is of the form (3.1) and independent of t , then*

$$(3.11) \quad \begin{aligned} \int F(u(x, t)) \eta(x) dx \Big|_{t_1}^{t_2} + \frac{1}{2\Lambda} \sum_{i=1}^N \int_{t_1}^{t_2} \int F''(u) \eta |D_i u|^{p_i} dx dt \\ \leq C_\Lambda \sum_{i=1}^N \int_{t_1}^{t_2} \int |F'(u)|^{p_i} F''(u)^{1-p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} dx dt \end{aligned}$$

and any $0 < t_1 < t_2 < T$.

Proof. Test the equation with $\varphi = F'(u) \eta$, which is readily checked to be admissible since $F' \in \text{Lip}(\mathbb{R})$ is bounded and

$$|D_i \varphi| \leq M |D_i \eta| + M |D_i u| \eta.$$

Notice that $|F(s)| \leq M|s|$, hence $F(u(\cdot, \tau)) \in L^1_{\text{loc}}(\mathbb{R}^N)$ since $u(\cdot, \tau) \in L^2_{\text{loc}}(\mathbb{R}^N)$, so by Steklov averaging we can compute

$$\int_{t_1}^{t_2} \int u_t \varphi dx dt = \int_{t_1}^{t_2} \int (F(u) \eta)_t dx dt = \int F(u(x, t_2)) \eta(x) dx - \int F(u(x, t_1)) \eta(x) dx$$

for any $T > t_2 > t_1 > 0$. Therefore we have

$$\int F(u(x, t)) \eta(x) dx \Big|_{t_1}^{t_2} + \sum_{i=1}^N \int_{t_1}^{t_2} \int A_i(x, u, Du) (F''(u) \eta D_i u + F'(u) D_i \eta) dx dt = 0$$

which implies by (1.2)

$$(3.12) \quad \begin{aligned} \int F(u(x, t)) \eta(x) dx \Big|_{t_1}^{t_2} + \frac{1}{\Lambda} \sum_{i=1}^N \int_{t_1}^{t_2} \int F''(u) \eta |D_i u|^{p_i} dx dt \\ \leq \Lambda \sum_{i=1}^N \int_{t_1}^{t_2} \int |D_i u|^{p_i-1} |F'(u)| |D_i \eta| dx dt. \end{aligned}$$

Proceeding as in (3.5) and making use of (3.6) we can bound the right-hand side as

$$\begin{aligned} |D_i u|^{p_i-1} |F'(u)| |D_i \eta| &= \hat{\eta}_i \left(|D_i u|^{p_i-1} F''(u)^{1-\frac{1}{p_i}} \eta_i^{p_i-1} \frac{|F'(u)|}{F''(u)^{1-\frac{1}{p_i}}} |D_i \eta_i| \right) \\ &\leq \frac{1}{2\Lambda} F''(u) \eta |D_i u|^{p_i} + C_\Lambda |F'(u)|^{p_i} F''(u)^{1-p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i}, \end{aligned}$$

which, inserted into (3.12) gives the claim. \square

Corollary 3.4. *Suppose $u \in \cap_{i=1}^N L^{p_i}(S_T)$ solves (2.2) in S_T for $u_0 \in L^1(\mathbb{R}^N) \cap L^2(\mathbb{R}^N)$. Then it holds*

$$\int |u(x, t)| dx \leq \int |u_0| dx, \quad \forall t \in [0, T]$$

Proof. Let η be as in (3.10) and define, for $\alpha > 0$ to be determined later,

$$F_\varepsilon(s) = \int_0^s \frac{\tau}{(|\tau|^\alpha + \varepsilon)^{\frac{1}{\alpha}}} d\tau,$$

so that

$$F'_\varepsilon(s) = \frac{s}{(|s|^\alpha + \varepsilon)^{\frac{1}{\alpha}}}, \quad F''_\varepsilon(s) = \frac{\varepsilon}{(|s|^\alpha + \varepsilon)^{\frac{1}{\alpha}+1}} > 0.$$

All the assumptions of the previous Lemma are satisfied and (3.11) implies, with our choice of F_ε ,

$$\int F_\varepsilon(u(x, t)) \eta(x, t) dx \Big|_{t_1}^{t_2} \leq C \sum_{i=1}^N \varepsilon^{1-p_i} \int_{t_1}^{t_2} \int |u|^{p_i} (|u|^\alpha + \varepsilon)^{p_i-1-\frac{1}{\alpha}} |D_i \eta^{\frac{1}{p_i}}|^{p_i} dx dt$$

We then choose

$$\alpha = (\max\{p_1, \dots, p_N\} - 1)^{-1} > 0,$$

so that $p_i - 1 - \frac{1}{\alpha} \leq 0$ for all $i = 1, \dots, N$ and therefore

$$\int F_\varepsilon(u(x, t)) \eta(x, t) dx \Big|_{t_1}^{t_2} \leq C_\varepsilon \sum_{i=1}^N \int_{t_1}^{t_2} \int |u|^{p_i} |D_i \eta^{\frac{1}{p_i}}|^{p_i} dx dt$$

for any $\varepsilon \in]0, 1[$ and $C_\varepsilon = C(\Lambda, \mathbf{p}, \varepsilon) > 0$. Since F_ε is 1-Lipschitz and $u(\cdot, t) \rightarrow u_0$ in $L^1_{\text{loc}}(\mathbb{R}^N)$, we can let $t_1 \rightarrow 0$ in the previous estimate. By $F_\varepsilon(s) \leq |s|$, $0 \leq \eta \leq 1$ and $|D_i \eta^{\frac{1}{p_i}}| \leq \frac{C}{R}$, we get

$$\int F_\varepsilon(u(x, t_2)) \eta(x, t_2) dx \leq \int |u_0| dx + \sum_{i=1}^N \frac{C_\varepsilon}{R^{p_i}} \int_0^{t_2} \int |u|^{p_i} dx dt,$$

for all $\varepsilon \in]0, 1[$ and $R \geq 1$. Let first $R \rightarrow +\infty$ to cancel out the last term thanks to the hypothesis $u \in \cap L^{p_i}(S_T)$ obtaining through Fatou's Lemma

$$\int F_\varepsilon(u(x, t_2)) dx \leq \int |u_0| dx \quad \forall t_2 \in [0, T]$$

and since $0 \leq F_\varepsilon(s) \nearrow |s|$, we obtain the conclusion by monotone convergence. \square

4. L^∞ -ESTIMATES

Theorem 4.1. *Let $p_1 \leq \dots \leq p_N$, $\bar{p} < N$ and $u \in \cap_{i=1}^N L^{p_i}(S_T)$ solve (2.2) for $u_0 \in L^1(\mathbb{R}^N) \cap L^2(\mathbb{R}^N)$. Then:*

(1) *if $\bar{p}_2 > 2$, then $u \in L^\infty_{\text{loc}}(0, T; L^\infty(\mathbb{R}^N))$ and for any $q \in [2, \bar{p}_2]$ the following estimate holds true*

$$(4.1) \quad \sup_{t \in [0, T]} \|u(\cdot, t)\|_\infty \leq \frac{C}{\theta^{\frac{N+\bar{p}}{\lambda_q}}} \left(\int_{\theta/2}^T \int |u|^q dx dt \right)^{\frac{\bar{p}}{\lambda_q}}, \quad \lambda_q = N(\bar{p} - 2) + \bar{p}q, \quad \theta > 0.$$

(2) *If $\bar{p}_1 > 2$, the following $L^1 - L^\infty$ estimate holds true for any $\tau \in]0, T]$*

$$(4.2) \quad \|u(\cdot, \tau)\|_\infty \leq \frac{C}{\tau^{\frac{N}{\lambda}}} \|u_0\|_1^{\frac{\bar{p}}{\lambda}}, \quad \lambda := \lambda_1 = N(\bar{p} - 2) + \bar{p}.$$

Proof. First observe that $\bar{p}_1 \leq \bar{p}_2$, so that we can assume $\bar{p}_2 > 2$, and in that case $\lambda_q \geq \lambda_2 > 0$ for all $q \geq 2$. The global condition $u \in \cap_{i=1}^N L^{p_i}(S_T)$, together with Corollary 3.2 and Theorem 2.4, imply that $u \in L^{\bar{p}_2}(S_T) \cap L^2(S_T)$. Therefore, by interpolation,

$$u \in L^q(S_T), \quad \text{for all } q \in [\min\{2, p_1\}, \max\{\bar{p}_2, p_N\}].$$

Let $k > 0$ to be determined, $T > \theta > 0$ and define for any $n \geq 0$

$$k_n = k - \frac{k}{2^n}, \quad \theta_n = \theta - \frac{\theta}{2^{n+1}}, \quad S_n = \mathbb{R}^N \times [\theta_n, T], \quad \psi_n(t) = \min \left\{ 1, \frac{2^{n+2}}{\theta} (t - \theta_n)_+ \right\},$$

so that

$$\psi_n \equiv 1 \text{ on } [\theta_{n+1}, T], \quad |(\psi_n)_t| \leq \frac{2^{n+2}}{\theta}.$$

Since $\psi_n(0) = 0$, the energy estimate (3.9) reads

$$(4.3) \quad \sup_{t \in [\theta_n, T]} \int (u - k_n)_+^2 dx + \frac{1}{C} \sum_{i=1}^N \iint_{S_n} |D_i(u - k_n)_+|^{p_i} dx dt \leq \frac{2^{n+2}}{\theta} \iint_{S_n} (u - k_n)_+^2 dx dt.$$

Choose $q \in [2, \bar{p}_2]$. If $A_n := \{(x, t) \in S_n : u \geq k_n\}$, Tchebichev's inequality yields

$$(4.4) \quad \iint_{S_n} (u - k_{n-1})_+^q dx dt \geq (k_n - k_{n-1})^q |A_n| = \frac{k^q}{2^{nq}} |A_n|$$

so that, by Hölder's inequality and the monotonicity of $\{S_n\}$

$$\iint_{S_n} (u - k_n)_+^2 dx dt \leq \left(\iint_{S_n} (u - k_n)_+^q dx dt \right)^{\frac{2}{q}} |A_n|^{1 - \frac{2}{q}} \leq \frac{2^{n(q-2)}}{k^{q-2}} \iint_{S_{n-1}} (u - k_{n-1})_+^q dx dt.$$

Therefore (4.3) becomes

$$(4.5) \quad \sup_{t \in [\theta_n, T]} \int (u - k_n)_+^2 dx + \frac{1}{C} \sum_{i=1}^N \iint_{S_n} |D_i(u - k_n)_+|^{p_i} dx dt \leq C \frac{2^{n(q-1)}}{\theta k^{q-2}} \iint_{S_{n-1}} (u - k_{n-1})_+^q dx dt$$

By Hölder inequality and (4.4)

$$\begin{aligned} \iint_{S_n} (u - k_n)_+^q dx dt &\leq |A_n|^{1 - \frac{q}{\bar{p}_2}} \left(\iint_{S_n} (u - k_n)_{+}^{\bar{p}_2} dx dt \right)^{\frac{q}{\bar{p}_2}} \\ &\leq C \left(\frac{2^{nq}}{k^q} \iint_{S_{n-1}} (u - k_{n-1})_+^q dx dt \right)^{1 - \frac{q}{\bar{p}_2}} \left(\iint_{S_n} (u - k_n)_{+}^{\bar{p}_2} dx dt \right)^{\frac{q}{\bar{p}_2}}. \end{aligned}$$

Applying (2.7) for $\alpha_i \equiv 1$, $\sigma = 2$ and $\theta = \frac{\bar{p}}{\bar{p}^*} = \frac{N}{N+2}$ gives $q = \bar{p}_2$ and thus by (4.5)

$$\begin{aligned} \iint_{S_n} (u - k_n)_{+}^{\bar{p}_2} dx dt &\leq \left(\sup_{t \in [\theta_n, T]} \int (u - k_n)_+^2(x, t) dx \right)^{1 - \frac{\bar{p}}{\bar{p}^*}} \prod_{i=1}^N \left(\iint_{S_n} |D_i(u - k_n)_+|^{p_i} dx dt \right)^{\frac{\bar{p}}{N p_i}} \\ &\leq C \left(\frac{2^{n(q-1)}}{\theta k^{q-2}} \iint_{S_{n-1}} (u - k_{n-1})_+^q dx dt \right)^{1 + \frac{\bar{p}}{N}}. \end{aligned}$$

Gathering together the previous two estimates, we obtain, for suitable $b > 1$,

$$\iint_{S_n} (u - k_n)_+^q dx dt \leq \frac{C}{(\theta k^{q-2})^{\frac{q}{\bar{p}_2} (1 + \frac{\bar{p}}{N})}} \frac{b^n}{k^{q(1 - \frac{\bar{p}}{\bar{p}_2})}} \left(\iint_{S_{n-1}} (u - k_{n-1})_+^q dx dt \right)^{(1 + \frac{\bar{p}}{N}) \frac{q}{\bar{p}_2} + 1 - \frac{q}{\bar{p}_2}}.$$

Setting

$$X_n = \iint_{S_n} (u - k_n)_+^q dx dt, \quad \alpha = \frac{q}{N+2}, \quad \beta = \frac{q}{\bar{p}} \frac{N + \bar{p}}{N + 2}, \quad \gamma = \frac{q}{\bar{p}} \frac{N(\bar{p} - 2) + \bar{p}q}{N + 2}$$

we thus obtained the recursive inequality

$$X_n \leq \frac{C}{\theta^\beta k^\gamma} b^n X_{n-1}^{1+\alpha}, \quad n \geq 1.$$

By the classical version of Lemma 2.7 with $N = 1$, we have that $X_n \rightarrow 0$ provided $X_0 \leq C\theta^{\frac{\beta}{\alpha}} k^{\frac{\gamma}{\alpha}}$ for a suitable constant C depending only on the data. Choosing k such that equality holds we therefore get

$$\sup_{\mathbb{R}^N \times [\theta, T]} u \leq k = \frac{C}{\theta^{\frac{N+\bar{p}}{\lambda_q}}} \left(\int_{\theta/2}^T \int_{\mathbb{R}^N} u_+^q dx dt \right)^{\frac{\bar{p}}{\lambda_q}} \quad \forall \theta \in]0, T[$$

implying (4.1) (by considering $-u$ as well). To prove (4.2), assume $\bar{p}_1 \geq 2$ and choose $q \in [2, \bar{p}_1[$. The function $v(x, t) = u(x, t + \theta)$ still solves the equation on $S_{T-\theta}$ and is bounded there by the previous estimate. Moreover $v \in \cap L^{p_i}(S_{T-\theta})$ hence (4.1) holds, reading

$$\sup_{\mathbb{R}^N \times [\tilde{\theta}, T-\theta]} |v| \leq \frac{C}{\tilde{\theta}^{\frac{N+\bar{p}}{\lambda_q}}} \left(\int_{\frac{\tilde{\theta}}{2}}^{T-\theta} \int |v|^q dx dt \right)^{\frac{\bar{p}}{\lambda_q}} \quad \forall \tilde{\theta} \in]0, T-\theta[.$$

In the latter inequality we set

$$\tilde{\theta} = \theta_n = \frac{T-\theta}{2^n}, \quad M_n = \sup_{\mathbb{R}^N \times [\theta_n, T-\theta]} |v|$$

to obtain

$$M_n \leq \frac{C}{\theta_n^{\frac{N+\bar{p}}{\lambda_q}}} \left(\int_{\theta_{n+1}}^{T-\theta} \int |v|^q dx dt \right)^{\frac{\bar{p}}{\lambda_q}} \leq C \frac{2^n \frac{N+\bar{p}}{\lambda_q}}{(T-\theta)^{\frac{N+\bar{p}}{\lambda_q}}} \left(\int_0^{T-\theta} \int |v| dx dt \right)^{\frac{\bar{p}}{\lambda_q}} M_{n+1}^{(q-1)\frac{\bar{p}}{\lambda_q}}$$

By the boundedness of v we infer the boundedness of $\{M_n\}$, while

$$(q-1)\frac{\bar{p}}{\lambda_q} < 1 \quad \Leftrightarrow \quad \bar{p}_1 > 2.$$

Therefore the interpolation lemma [11, Ch. I, Lemma 4.3] provides

$$M_0 \leq C \left(\frac{1}{(T-\theta)^{\frac{N+\bar{p}}{\lambda_q}}} \left(\int_0^{T-\theta} \int |v| dx dt \right)^{\frac{\bar{p}}{\lambda_q}} \right)^{\frac{1}{1-(q-1)\frac{\bar{p}}{\lambda_q}}}.$$

Writing the latter in terms of u and noting that

$$\lambda_q \left(1 - (q-1)\frac{\bar{p}}{\lambda_q} \right) = \lambda_1,$$

we obtain

$$\sup_{\mathbb{R}^N} |u(\cdot, T-\theta)| \leq C \frac{1}{(T-\theta)^{\frac{N+\bar{p}}{\lambda_1}}} \left(\iint_{\mathbb{R}^N \times [\theta, T]} |u| dx dt \right)^{\frac{\bar{p}}{\lambda_1}}, \quad \forall \theta \in]0, T[.$$

Using Corollary 3.4 while setting $\tau := T - \theta \in]0, T[$ finally gives (4.2). \square

Another type of L^∞ estimate is the following, which instead is purely local.

Lemma 4.2. *Let*

$$(4.6) \quad p_1 \leq \cdots \leq p_N \quad \text{and} \quad \max\{2, p_N\} < \bar{p}_2.$$

Define for $k > 0$ the functions

$$(4.7) \quad g(k) = \sum_{i=1}^N k^{\bar{p}_2 - p_i}, \quad h(k) = \left(\sum_{i=1}^N k^{p_i - \bar{p}_2} \right)^{-1},$$

and for $T \in \mathbb{R}$, $M, \lambda > 0$ the intrinsic rectangle

$$Q_{\lambda, M} = \prod_{i=1}^N \left[-\lambda^{\frac{1}{p_i}}, \lambda^{\frac{1}{p_i}} \right] \times [T - M\lambda, T].$$

If u is a weak supersolution to (2.1) in $Q_{\lambda, M}$, then

$$(4.8) \quad \|u_+\|_{L^\infty(Q_{\lambda/2, M})} \leq g^{-1}(1/M) + h^{-1} \left(C \left(M \iint_{Q_{\lambda, M}} u_+^{\bar{p}_2} dx \right)^{\frac{\bar{p}}{N + \bar{p}}} \right).$$

Proof. Define for any $k > 0$ and λ, M, T

$$k_n = k - \frac{k}{2^n}, \quad \theta_n = T - \frac{M\lambda}{2} \left(1 + \frac{1}{2^n} \right), \quad r_{n,i} = \frac{\lambda^{\frac{1}{p_i}}}{2^{\frac{1}{p_i}}} \left(1 + \frac{1}{2^n} \right) \quad Q_n = \prod_{i=1}^N [-r_{n,i}, r_{n,i}] \times [\theta_n, T]$$

so that $Q_{n+1} \subseteq Q_n \subseteq Q_{\lambda, M}$ for all $n \geq 0$ and formally $Q_\infty = Q_{\lambda/2, M}$. Construct functions $\eta_n \in C^\infty(Q_n; [0, 1])$ of the form (3.1) such that

$$\eta_n|_{\partial_p Q_n} \equiv 0, \quad \eta_n|_{Q_{n+1}} \equiv 1, \quad |D_i \eta_n^{\frac{1}{p_i}}| \leq \frac{C 2^n}{\lambda^{\frac{1}{p_i}}}, \quad |(\eta_n)_t| \leq \frac{C 2^n}{M\lambda}$$

Apply (3.2) to obtain

$$\begin{aligned} & \sup_{t \in [\theta_n, T]} \int (u - k_n)_+^2 \eta_n(x, t) dx + \sum_{i=1}^N \iint_{Q_n} |D_i((u - k_n)_+ \eta_n)|^{p_i} dx dt \\ & \leq C 2^n \left(\frac{1}{M\lambda} \iint_{Q_n} (u - k_n)_+^2 dx dt + \frac{1}{\lambda} \sum_{i=1}^N \iint_{Q_n} (u - k_n)_+^{p_i} dx dt \right). \end{aligned}$$

Letting $p_0 = 2$, $A_n = \{(x, t) \in Q_n : u \geq k_n\}$ and recalling (4.4), it holds for any $i = 0, \dots, N$

$$\iint_{Q_n} (u - k_n)_+^{p_i} dx dt \leq \left(\iint_{Q_n} (u - k_n)_+^{\bar{p}_2} dx dt \right)^{\frac{p_i}{\bar{p}_2}} |A_n|^{1 - \frac{p_i}{\bar{p}_2}} \leq \frac{C b^n}{k^{\bar{p}_2 - p_i}} \iint_{Q_{n-1}} (u - k_{n-1})_+^{\bar{p}_2} dx dt,$$

for some numerical $b > 1$, so that by the definition (4.7) we obtain

$$\begin{aligned} & \sup_{t \in [\theta_n, T]} \int (u - k_n)_+^2 \eta_n(x, t) dx + \sum_{i=1}^N \iint_{Q_n} |D_i((u - k_n)_+ \eta_n)|^{p_i} dx dt \\ & \leq C b^n \left(\frac{1}{M\lambda k^{\bar{p}_2 - 2}} + \frac{1}{\lambda h(k)} \right) \iint_{Q_{n-1}} (u - k_{n-1})_+^{\bar{p}_2} dx dt. \end{aligned}$$

As $\{k_n\}$ and $\{\theta_n\}$ are increasing and $\eta_n \equiv 1$ on $\text{supp}(\eta_{n+1})$,

$$\begin{aligned} \sup_{t \in [\theta_{n+1}, T]} \int ((u - k_{n+1})_+ \eta_{n+1})^2(x, t) dx &\leq \sup_{t \in [\theta_n, T]} \int (u - k_n)_+^2 \eta_n(x, t) dx \\ &\leq C b^n \left(\frac{1}{M \lambda k^{\bar{p}_2 - 2}} + \frac{1}{\lambda h(k)} \right) \iint_{Q_{n-1}} (u - k_{n-1})_+^{\bar{p}_2} dx dt. \end{aligned}$$

Again by the monotonicity of $\{k_n\}$ and $\{Q_n\}$ it holds

$$\begin{aligned} \sum_{i=1}^N \iint_{Q_{n+1}} |D_i((u - k_{n+1})_+ \eta_{n+1})|^{p_i} dx dt &\leq C b^{n+1} \left(\frac{1}{M \lambda k^{\bar{p}_2 - 2}} + \frac{1}{\lambda h(k)} \right) \iint_{Q_n} (u - k_n)_+^{\bar{p}_2} dx dt \\ &\leq C b^n \left(\frac{1}{M \lambda k^{\bar{p}_2 - 2}} + \frac{1}{\lambda h(k)} \right) \iint_{Q_{n-1}} (u - k_{n-1})_+^{\bar{p}_2} dx dt \end{aligned}$$

Therefore, applying (2.7) with $\alpha_i \equiv 1$, $\sigma = 2$, $\theta = \bar{p}/\bar{p}^*$ and thus $q = \bar{p}_2$, we deduce, for some other constant $C, b \geq 1$, the recursive inequality

$$\iint_{Q_{n+1}} (u - k_{n+1})_+^{\bar{p}_2} dx dt \leq C b^n \left(\frac{1}{M \lambda k^{\bar{p}_2 - 2}} + \frac{1}{\lambda h(k)} \right)^{1 + \frac{\bar{p}}{N}} \left(\iint_{Q_{n-1}} (u - k_{n-1})_+^{\bar{p}_2} dx dt \right)^{1 + \frac{\bar{p}}{N}}$$

Now if k is so large that

$$(4.9) \quad \frac{1}{M k^{\bar{p}_2 - 2}} \leq \frac{1}{h(k)}, \quad \Leftrightarrow \quad k \geq g^{-1}(1/M),$$

then the previous iterative inequality reads

$$X_{n+1} \leq C b^n \left(\frac{1}{\lambda h(k)} \right)^{1 + \frac{\bar{p}}{N}} X_n^{1 + \frac{\bar{p}}{N}}, \quad n \geq 0$$

where

$$X_n = \iint_{Q_{2n}} (u - k_{2n})_+^{\bar{p}_2} dx dt.$$

By Lemma 2.7 for $N = 1$, $X_n \rightarrow 0$ whenever $X_0 \leq C(\lambda h(k))^{\frac{N+\bar{p}}{\bar{p}}}$ and, taking account of (4.9), this in turn implies that

$$\sup_{Q_{\lambda/2, M}} u_+ \leq \max \left\{ g^{-1}(1/M), h^{-1} \left(\frac{C}{\lambda} \left(\iint_{Q_0} u_+^{\bar{p}_2} dx dt \right)^{\frac{\bar{p}}{N+\bar{p}}} \right) \right\}.$$

Note that $Q_0 \subseteq Q_{\lambda, M}$ and

$$(4.10) \quad |Q_{\lambda, M}| = M \lambda \prod_{i=1}^N \lambda^{\frac{1}{p_i}} = M \lambda^{\frac{N+\bar{p}}{\bar{p}}},$$

so that being h monotone increasing we obtain

$$\sup_{Q_{\lambda/2, M}} u \leq \max \left\{ g^{-1}(1/M), h^{-1} \left(C \left(M \iint_{Q_{\lambda, M}} u_+^{\bar{p}_2} dx dt \right)^{\frac{\bar{p}}{N+\bar{p}}} \right) \right\}.$$

□

The next proof follows [24].

Corollary 4.3. *Let (4.6) hold and u be a weak supersolution of (2.1) in Ω_T . Then u has a lower semicontinuous representative.*

Proof. Recall that all the infima and suprema are taken in the essential sense. For any $M \in \mathbb{N}$ define a metric in Ω_T as

$$\text{dist}_M((x, t), (x', t')) = \max\{M|t - t'|, |x_1 - x'_1|^{\frac{1}{p_1}}, \dots, |x_N - x'_N|^{\frac{1}{p_N}}\},$$

with corresponding balls $B_{r,M}$. We will prove that there is a dist_1 -metric essential lower-semicontinuous representative of u and we start by fixing an arbitrary one, which we'll still denote by u . By (4.10), Ω_T with the induced metric and the Lebesgue measure is a locally doubling measure space. Therefore the set E_M of Lebesgue points for u has full measure, as well as $E = \bigcap_{M \in \mathbb{N}} E_M$. We can therefore suppose that for any $(x_0, t_0) \in E$ and for every $M \in \mathbb{N}$

$$\lim_{r \downarrow 0} \iint_{B_{r,M}} |u(x, t) - u(x_0, t_0)|^{\bar{p}_2} dx dt = 0.$$

We claim that for any $(x_0, t_0) \in E$

$$(4.11) \quad u(x_0, y_0) \leq \liminf_{r \rightarrow 0} \inf_{B_{r,1}(x_0, t_0)} u.$$

Suppose by contradiction that

$$(4.12) \quad u(x_0, y_0) - \inf_{B_{r,1}(x_0, t_0)} u \geq \varepsilon > 0 \quad \forall r < r_0$$

and consider the solution $v = u(x_0, t_0) - u$ to (2.1). Since $g(0) = h(0) = 0$, g and h are continuous and increasing, we can choose $M > 0$ such that

$$g^{-1}(1/M) + h^{-1}(C/M) < \varepsilon/2,$$

(C being the constant in (4.8)) and, being $(x_0, t_0) \in E_M$, choose $r(M) < r_0$ such that

$$B_{2r,M}(x_0, t_0) \subseteq \Omega_T, \quad \iint_{B_{2r,M}(x_0, t_0)} |u - u(x_0, t_0)|^{\bar{p}_2} dx dt \leq \frac{1}{M^{2+\frac{N}{\bar{p}}}}.$$

The previous Theorem (applied to $v(x - x_0, t)$) then assures that

$$\begin{aligned} \sup_{B_{r,1}(x_0, t_0)} u(x_0, t_0) - u &\leq \sup_{B_{r,M}(x_0, t_0)} u(x_0, t_0) - u \\ &\leq g^{-1}(1/M) + h^{-1} \left(C \left(M \iint_{B_{2r,M}(x_0, t_0)} (u - u(x_0, t_0))_+^{\bar{p}_2} dx dt \right)^{\frac{\bar{p}}{N+\bar{p}}} \right) \\ &\leq g^{-1}(1/M) + h^{-1}(C/M) < \varepsilon/2 \end{aligned}$$

contradicting (4.12). Finally, for $(x_0, t_0) \in \Omega_T \setminus E$ we modify the representative forcing the equality in (4.11). \square

5. PROOF OF THE MAIN THEOREM

Suppose that for a subset of M indexes, $1 \leq M \leq N$, it holds $p_{j_1}, \dots, p_{j_M} > 2$. We perform a permutation of the variables letting x_{j_1}, \dots, x_{j_N} be the last M ones and splitting \mathbb{R}^N as $\mathbb{R}^{N-M} \times \mathbb{R}^M$ with $\mathbb{R}^N \ni x = (x', x'') \in \mathbb{R}^{N-M} \times \mathbb{R}^M$. Therefore we can assume henceforth that

$$(5.1) \quad p_j > 2, \quad \text{for all } j = N - M + 1, \dots, N, \quad M \geq 1.$$

Furthermore, we will set, $K'_r = \{x' \in \mathbb{R}^{N-M} : |x'| \leq r\}$, $K''_r = \{x'' \in \mathbb{R}^M : |x''| \leq r\}$ and say that

$$u \in L^q_{\text{loc}}(\mathbb{R}^M; L^q(\mathbb{R}^{N-M} \times [0, T])) \Leftrightarrow \int_0^T \int_{\mathbb{R}^{N-M} \times K''_r} |u|^q dx' dx'' dt < +\infty$$

for any $r > 0$.

Lemma 5.1. *Assume (5.1), $\bar{p} < N$ and $\bar{p}_2 > \max\{p_1, \dots, p_N\}$. Let $u \in \cap_{i=1}^N L^q_{\text{loc}}(\mathbb{R}^M; L^{p_i}(\mathbb{R}^{N-M} \times [0, T]))$ be a weak solution of the Cauchy problem (2.2) in S_T , with $u_0 \in L^2(\mathbb{R}^N)$ and*

$$\text{supp}(u_0) \subseteq \mathbb{R}^{N-M} \times K''_{R_0}.$$

There exists $\alpha, \beta, c > 0$ such that, letting,

$$\tau(u, r, T) := cr^\alpha \left(\sum_{i=1}^N \int_0^T \int_{\mathbb{R}^{N-M} \times K''_{3r}} |u|^{p_i} dx \right)^{-\beta}, \quad r \geq 2R_0$$

then $u(\cdot, t) \equiv 0$ on $\mathbb{R}^{N-M} \times (K''_{2r} \setminus K''_r)$ for all $t \in [0, \tau(u, r, T)]$.

Proof. Let $\tau \in]0, \min\{1, T\}[$, $R \geq 1$, $r \geq 2R_0$ define

$$r_n = 2r + \frac{r}{2^n}, \quad s_n = r - \frac{r}{2^{n+1}}, \quad E_n = K''_{r_n} \setminus K''_{s_n}$$

so that $E''_{n+1} \subseteq E''_n$ and $\mathbb{R}^{N-M} \times E''_n \cap \text{supp } u_0 = \emptyset$. Choose $\eta' \in C_c^\infty(K'_{2R}; [0, 1])$ of the form (3.1) such that

$$\eta' \equiv 1 \text{ on } K'_R, \quad |D_i(\eta')^{\frac{1}{p_i}}| \leq \frac{C}{R},$$

for $i = 1, \dots, N-M$ and $\eta''_n \in C_c^\infty(E_n; [0, 1])$ of the form (3.1) and such that

$$\eta''_n \equiv 1 \text{ on } E_{n+1}, \quad |D_i(\eta''_n)^{\frac{1}{p_i}}| \leq C \frac{2^n}{r}$$

for $i = N-M+1, \dots, N$. We apply (3.2) for $k = 0$, $\eta_n(x) = \eta'(x')\eta''_n(x'')$ and $t_1 \rightarrow 0$ to both u and $-u$. Since $u \in C^0([0, T]; L^2_{\text{loc}}(\mathbb{R}^N))$, and $\text{supp}(u_0) \cap \text{supp}(\eta_n) = \emptyset$, we obtain

$$\int u^2(x, t) \eta_n(x) dx + \frac{1}{C} \sum_{i=1}^N \int_0^t \int |D_i(u \eta_n)|^{p_i} dx dt \leq C \sum_{i=1}^N \int_0^t \int |u|^{p_i} |D_i \eta_n^{\frac{1}{p_i}}|^{p_i} dx dt$$

for any $t \in [0, \tau]$. Letting $R \rightarrow +\infty$ and using the assumption $u \in \cap_{i=1}^N L^q_{\text{loc}}(\mathbb{R}^M; L^{p_i}(\mathbb{R}^{N-M} \times [0, T]))$ together with $|D_i(\eta')^{\frac{1}{p_i}}| \leq C/R$, all the terms for $i = 1, \dots, N-M$ on the right hand side vanish. Therefore, through Fatou's Lemma on the left and dominated convergence on the remaining terms on the right, we infer

$$\begin{aligned} \int u^2(x, t) \eta''_n(x'') dx + \frac{1}{C} \sum_{i=1}^N \int_0^t \int |D_i(u \eta''_n)|^{p_i} dx dt &\leq C \sum_{N-M+1}^N \int_0^t \int |u|^{p_i} |D_i(\eta''_n)^{\frac{1}{p_i}}|^{p_i} dx dt \\ &\leq C \sum_{N-M+1}^N \frac{2^{n p_i}}{r^{p_i}} \int_0^t \int_{\mathbb{R}^{N-M} \times E_n} |u|^{p_i} dx dt. \end{aligned}$$

By the usual monotonicity argument, the latter implies

$$\int (u \eta''_{n+1})^2(x, t) dx + \frac{1}{C} \sum_{i=1}^N \int_0^t \int |D_i(u \eta''_{n+1})|^{p_i} dx dt \leq C \sum_{N-M+1}^N \frac{2^{n p_i}}{r^{p_i}} \int_0^t \int_{\mathbb{R}^{N-M} \times E_n} |u|^{p_i} dx dt.$$

Let

$$X_n := \sum_{N-M+1}^N \frac{2^{n p_i}}{r^{p_i}} \int_0^t \int_{\mathbb{R}^M \times E_n} |u|^{p_i} dx dt.$$

For any $j = N - M + 1, \dots, N$, apply (2.7) with

$$\alpha_i \equiv 1, \quad \sigma = 2, \quad \theta = \theta_j = \frac{p_j - 2}{\bar{p}^* - 2}, \quad q = p_j$$

(notice that $2 < p_j$ and $\bar{p}_2 > p_N$ imply $0 < \theta_j < \frac{\bar{p}}{\bar{p}^*}$ for all $j = N - M + 1, \dots, N$), to obtain

$$\begin{aligned} \int_0^\tau \int_{\mathbb{R}^{N-M} \times E_{n+2}} |u|^{p_j} dx dt &\leq \int_0^\tau \int |u \eta''_{n+1}|^{p_j} dx dt \\ &\leq C \tau^{1-\theta_j \frac{\bar{p}^*}{\bar{p}}} \left(\sup_{t \in [0, \tau]} \int (u \eta''_{n+1})^2(x, t) dx \right)^{1-\theta_j} \prod_{i=1}^N \left(\int_0^\tau \int |D_i(u \eta''_{n+1})|^{p_i} dx dt \right)^{\frac{\theta_j \bar{p}^*}{N p_i}} \\ &\leq C \tau^{1-\theta_j \frac{\bar{p}^*}{\bar{p}}} X_n^{1-\theta_j} \prod_{i=1}^N X_n^{\frac{\theta_j \bar{p}^*}{N p_i}} = C \tau^{1-\theta_j \frac{\bar{p}^*}{\bar{p}}} X_n^{1+\theta_j \frac{\bar{p}^*}{N}}. \end{aligned}$$

Let

$$p_{\max} = \max\{p_{j_1}, \dots, p_{j_M}\}, \quad p_{\min} = \min\{p_{j_1}, \dots, p_{j_M}\}.$$

Since for any $n \geq 0$

$$X_n \leq C \frac{2^{n p_{\max}}}{r^{p_{\min}}} \sum_{N-M+1}^N \int_0^\tau \int_{\mathbb{R}^{N-M} \times E_n} |u|^{p_i} dx dt,$$

where defining

$$Y_n := \sum_{N-M+1}^N \int_0^\tau \int_{\mathbb{R}^{N-M} \times E_{3n}} |u|^{p_i} dx dt$$

we obtained the recursive inequality

$$Y_{n+1} \leq C \frac{2^{n p_{\max}} \tau^{1-\theta_{\max} \frac{\bar{p}^*}{\bar{p}}}}{r^{p_{\min}}} \frac{1}{M} \sum_{N-M+1}^N Y_n^{1+\theta_j \frac{\bar{p}^*}{N}}.$$

Setting

$$\gamma = \frac{N p_{\min}}{\theta_{\min} \bar{p}^*}, \quad \delta = \frac{N}{\theta_{\min}} \left(\frac{1}{\bar{p}^*} - \frac{\theta_{\max}}{\bar{p}} \right),$$

Lemma 2.7 ensures that for some other constant $C = C(N, \mathbf{p}, \Lambda)$

$$Y_0 \leq C \frac{r^\gamma}{\tau^\delta}, \quad \Rightarrow \quad Y_n \rightarrow 0,$$

which in turn implies $u(x, t) \equiv 0$ for $x \in K''_{2r} \setminus K''_r$ and $t \in [0, \tau]$ and proves the claim for $\alpha = \gamma/\delta$, $\beta = 1/\delta$. \square

It is useful to state the previous Lemma in the case when all the p_i 's are greater than 2.

Corollary 5.2. *Suppose $2 < p_1 \leq \dots \leq p_N < \bar{p}_2$ and let u solve (2.2) for $u_0 \in L^2(\mathbb{R}^N)$ with $\text{supp}(u_0) \subseteq K_{R_0}$. There exists $\alpha, \beta, c > 0$ such that, letting,*

$$\tau(u, r, T) := c r^\alpha \left(\sum_{i=1}^N \int_0^T \int_{K_{3r}} |u|^{p_i} dx \right)^{-\beta}, \quad r \geq 2R_0$$

then $u(\cdot, t) \equiv 0$ on $K_{2r} \setminus K_r$ for all $t \in [0, \tau(u, r, T)]$.

Proof. In this case the assumption $u \in \cap_{i=1}^N L_{\text{loc}}^{p_i}(\mathbb{R}^M; L^{p_i}(\mathbb{R}^{N-M} \times [0, T]))$ reduces to $u \in \cap_{i=1}^N L^{p_i}(0, T; L_{\text{loc}}^{p_i}(\mathbb{R}^N))$. But this follows from Theorem 2.6 and the condition $p_N < \bar{p}_2$. \square

We can finally prove the main result of the paper. In the case $p_i > 2$ for all $i = 1, \dots, N$ we can choose $M = N$, so that, as in the previous proof, the condition $u \in \cap_{i=1}^N L_{\text{loc}}^{p_i}(\mathbb{R}^M; L^{p_i}(\mathbb{R}^{N-M} \times [0, T]))$ reduces to $u \in \cap_{i=1}^N L^{p_i}(0, T; L_{\text{loc}}^{p_i}(\mathbb{R}^N))$. As long as $\max\{p_1, \dots, p_N\} < \bar{p}_2$, this directly follows from the condition of being a weak solution together with Theorem 2.6, as by Hölder's inequality

$$L^{p_i}(0, T; L_{\text{loc}}^{p_i}(\mathbb{R}^N)) \subseteq L^\infty(0, T; L_{\text{loc}}^2(\mathbb{R}^N)) \cap L^{\bar{p}_2}(0, T; L_{\text{loc}}^{\bar{p}_2}(\mathbb{R}^N))$$

for all $i = 1, \dots, N$.

Theorem 5.3. *Assume (5.1), $\bar{p} < N$ and*

$$(5.2) \quad \bar{p}_1 = \bar{p} \left(1 + \frac{1}{N} \right) > \max\{p_1, \dots, p_N\}.$$

Let $u \in \cap_{i=1}^N L_{\text{loc}}^{p_i}(\mathbb{R}^M; L^{p_i}(\mathbb{R}^{N-M} \times [0, T]))$ be a weak solution of the Cauchy problem (2.2) in S_T with $u_0 \in L^2(\mathbb{R}^N) \cap L^1(\mathbb{R}^N)$ such that

$$\emptyset \neq \text{supp}(u_0) \subseteq \mathbb{R}^{N-M} \times K_{R_0}''.$$

Then there is a branch $\tilde{u} \neq 0$ of u such that for any $j = N - M + 1, \dots, N$

$$(5.3) \quad \text{supp}(\tilde{u}(\cdot, t)) \subseteq \{|x_j| \leq R_j(t)\},$$

for any $t \in]0, T[$, where

$$R_j(t) = 2R_0 + Ct \frac{N(\bar{p}-p_j)+\bar{p}}{p_j^\lambda} \|u_0\|_1^{\frac{\bar{p}}{p_j} \frac{p_j-2}{\lambda}}, \quad \lambda = N(\bar{p}-2) + \bar{p}.$$

Proof. We start proving (5.3) for the whole u assuming that

$$(5.4) \quad u \in \cap_{i=1}^N L^{p_i}(S_T).$$

Since $2 < p_N \leq \max\{p_1, \dots, p_N\} < \bar{p}_1$, Theorem 4.1, point 2) applies, ensuring (4.2). Choose $\mu \in]0, 1[$ and for any $\varepsilon > 0$ apply (3.11) with

$$F_\varepsilon(s) = \int_0^s \tau(\tau^2 + \varepsilon^2)^{\frac{\mu-1}{2}} d\tau, \quad F_\varepsilon''(s) = \frac{\mu s^2 + \varepsilon^2}{(s^2 + \varepsilon^2)^{\frac{3-\mu}{2}}} > 0.$$

All the assumptions of Lemma 3.3 hold true, except the boundedness of F' , which however is not necessary being u bounded on $S_T \setminus S_t$, $t > 0$ by (4.2). Using

$$\mu(s^2 + \varepsilon^2)^{\frac{\mu-1}{2}} \leq F_\varepsilon''(s), \quad |F_\varepsilon'(u)|^{p_i} |F_\varepsilon''(u)|^{1-p_i} \leq |u|^{p_i} (u^2 + \varepsilon^2)^{\frac{\mu-1}{2}}$$

for $i = 1, \dots, N$, we get for all $0 < t_1 < t_2 < T$ and η of the form (3.1)

$$\begin{aligned} \int F_\varepsilon(u(x, t)) \eta(x) dx \Big|_{t_1}^{t_2} + \frac{\mu}{2\Lambda} \sum_{i=1}^N \int_{t_1}^{t_2} \int (u^2 + \varepsilon^2)^{\frac{\mu-1}{2}} \eta |D_i u|^{p_i} dx dt \\ \leq C_\Lambda \sum_{i=1}^N \int_{t_1}^{t_2} \int |u|^{p_i} (u^2 + \varepsilon^2)^{\frac{\mu-1}{2}} |D_i \eta^{\frac{1}{p_i}}|^{p_i} dx dt. \end{aligned}$$

Being $\mu \in]0, 1[$, by monotone convergence on all the terms we obtain

$$(5.5) \quad \begin{aligned} \int |u(x, t)|^{1+\mu} \eta(x) dx \Big|_{t_1}^{t_2} + \frac{1}{C} \sum_{i=1}^N \int_{t_1}^{t_2} \int \eta |u|^{\mu-1} |D_i u|^{p_i} dx dt \\ \leq C \sum_{i=1}^N \int_{t_1}^{t_2} \int |u|^{p_i+\mu-1} |D_i \eta^{\frac{1}{p_i}}|^{p_i} dx dt \end{aligned}$$

for some constant $C = C(\mu, \Lambda) > 0$. A further monotone convergence argument shows that the previous estimate holds true for any $0 \leq t_1 \leq t_2 \leq T$. Let $j \geq N - M + 1$ and choose $\tilde{\eta}, \psi \in C_c^\infty(\mathbb{R}; [0, 1])$

$$\eta(x) = \tilde{\eta}(x_j)^{p_j} \prod_{i \neq j} \psi(x_i)^{p_i}$$

with the properties

$$\tilde{\eta}|_{\{|s| \leq R_0\}} \equiv 0, \quad \psi|_{\{|s| \leq R\}} \equiv 1, \quad |\psi'| \leq C/R.$$

With this test function, we let $t_1 = 0$, $t_2 = T$ and $R \rightarrow +\infty$ in (5.5): all the terms on the right except the j -th one vanish, since (5.4) holds and $u \in L^1(S_T)$ by (3.4), therefore by interpolation $u \in L^{p_i+\mu-1}(S_T)$. On the left-hand side the term for $t_1 = 0$ vanishes since $\text{supp}(u_0) \subseteq \{|x_j| \leq R_0\}$ and on the other we apply Fatou's lemma, to obtain

$$(5.6) \quad \int |u(x, T)|^{1+\mu} \tilde{\eta} dx + \frac{1}{C} \sum_{i=1}^N \int_0^T \int \tilde{\eta}^{p_j} |u|^{\mu-1} |D_i u|^{p_i} dx dt \leq C \int_0^T \int |u|^{p_j+\mu-1} |D_j \tilde{\eta}|^{p_j} dx dt$$

where we set for brevity $\tilde{\eta}(x) = \tilde{\eta}(x_j)$. We specify further the function $\tilde{\eta}$ defining for $r > 2R_0$, $n \in \mathbb{N}$

$$r_n = 2r + \frac{r}{2^n}, \quad s_n = r - \frac{r}{2^{n+1}}, \quad E_n = \{x \in \mathbb{R}^N : s_n \leq |x_j| \leq r_n\},$$

and for suitable $\tilde{\eta}_n \in C^\infty(\mathbb{R}; [0, 1])$,

$$(5.7) \quad \tilde{\eta}_n \equiv 1 \text{ on } E_{n+1}, \quad |\tilde{\eta}'_n| \leq C \frac{2^n}{r}, \quad \text{supp}(\tilde{\eta}_n) \subseteq \{s_n \leq |s| \leq r_n\}.$$

Let finally

$$\beta_i = \frac{p_i + \mu - 1}{p_i} < 1, \quad \beta = \min\{\beta_i : j = 1, \dots, N\}, \quad \eta_n(x) := \tilde{\eta}_n^{1/\beta}(x_j).$$

Clearly η_n still satisfies (5.7), while being $0 \leq \tilde{\eta}_n \leq 1$

$$|D_i |\eta_n u|^{\beta_i}|^{p_i} = \eta_n^{\beta_i p_i} |D_i |u|^{\beta_i}|^{p_i} \leq \beta_i^{p_i} \tilde{\eta}_n^{p_i} |u|^{\mu-1} |D_i u|^{p_i},$$

for all $i \neq j$, where we used $\beta_i \geq \beta$ and (2.6) in the last inequality. Therefore (5.6) for $\tilde{\eta} = \tilde{\eta}_n$ provides

$$(5.8) \quad \int_0^T \int |D_i |\eta_n u|^{\beta_i}|^{p_i} dx dt \leq \frac{C 2^n}{r^{p_j}} \int_0^T \int_{E_{n-1}} |u|^{p_j+\mu-1} dx dt$$

for all $i \neq j$, where we used the properties in (5.7) and the monotonicity of E_n . When $i = j$ we similarly have

$$|D_j |\eta_n u|^{\beta_j}|^{p_j} \leq \frac{C 2^n}{r^{p_j}} |u|^{p_j+\mu-1} + \beta_j^{p_j} \eta_n^{\beta_j p_j} |u|^{\mu-1} |D_j u|^{p_j} \leq \frac{C 2^n}{r^{p_j}} |u|^{p_j+\mu-1} + \beta_j^{p_j} \tilde{\eta}_n^{p_j} |u|^{\mu-1} |D_j u|^{p_j},$$

giving (5.8) for $i = j$ as well. Since (5.6) also implies

$$\int |\tilde{\eta}_n u|^{1+\mu}(x, t) dx \leq \int_{E_n} \tilde{\eta}_{n-1} |u|^{1+\mu}(x, t) dx \leq \frac{C 2^n}{r^{p_j}} \int_0^T \int_{E_{n-1}} |u|^{p_j+\mu-1} dx dt$$

for any $t \in]0, T]$, we can apply Proposition 2.4 with parameters

$$\sigma = 1 + \mu, \quad \alpha_i = \beta_i = \frac{p_i + \mu - 1}{p_i}, \quad \theta = \frac{p_j - 2}{p_\alpha^* - \mu - 1}, \quad q = p_j + \mu - 1.$$

Substitution gives

$$\tilde{\alpha} = N \left(\frac{1}{\bar{p}'} + \frac{\mu}{\bar{p}} \right), \quad \theta = (p_j - 2) \frac{N - \bar{p}}{\lambda_{1+\mu}},$$

where we recall that

$$\lambda_{1+\mu} = N(\bar{p} - 2) + (1 + \mu)\bar{p}.$$

The necessary condition $\theta \in [0, \frac{\bar{p}}{\bar{p}^*}]$ reads, after some algebraic manipulations, as

$$p_j \leq \bar{p} \left(1 + \frac{\mu + 1}{N} \right) \Leftrightarrow \mu \geq N \left(\frac{p_j}{\bar{p}} - 1 \right) - 1,$$

and the latter quantity is always negative under assumption (5.2). Therefore for any $\mu \in]0, 1[$ (2.7) gives, through the previous estimates and some algebra

$$\int_0^T \int |\tilde{\eta}_n u|^{p_j+\mu-1} dx dt \leq C 2^n T^{1-(p_j-2)\frac{N}{\lambda_{1+\mu}}} \left(\frac{2}{r^{p_j}} \int_0^T \int_{E_{n-1}} |u|^{p_j+\mu-1} dx dt \right)^{1+(p_j-2)\frac{\bar{p}}{\lambda_{1+\mu}}}$$

which, being $\tilde{\eta}_n \equiv 1$ on E_{n+1} , implies

$$\int_0^T \int_{E_{n+1}} |u|^{p_j+\mu-1} dx dt \leq C 2^n T^{1-(p_j-2)\frac{N}{\lambda_{1+\mu}}} \left(\frac{2}{r^{p_j}} \int_0^T \int_{E_{n-1}} |u|^{p_j+\mu-1} dx dt \right)^{1+(p_j-2)\frac{\bar{p}}{\lambda_{1+\mu}}}.$$

Applying the classical form of Lemma 2.7 for $N = 1$ gives that the condition

$$(5.9) \quad \int_0^T \int_{E_0} |u|^{p_j+\mu-1} dx dt \leq C r^{p_j \left(1 + \frac{\lambda_{1+\mu}}{(p_j-2)\bar{p}} \right)} T^{\frac{N}{\bar{p}} - \frac{\lambda_{1+\mu}}{(p_j-2)\bar{p}}}$$

implies

$$\int_0^T \int_{E_{2n}} |u|^{p_j+\mu-1} dx dt \rightarrow 0$$

and hence

$$(5.10) \quad \text{supp}(u(\cdot, t)) \subseteq \mathbb{R}^N \setminus \{r \leq |x_j| \leq 2r\} \quad \forall t \in [0, T].$$

To obtain (5.9), we employ Corollary 3.4 and Theorem 4.1 as follows:

$$\begin{aligned} \int_0^T \int_{E_0} |u|^{p_j+\mu-1} dx dt &\leq \int_0^T \|u(\cdot, t)\|_1 \|u(\cdot, t)\|_{L^\infty}^{p_j+\mu-2} dt \\ &\leq C \|u_0\|_1 \int_0^T \frac{\|u_0\|_1^{\frac{\bar{p}}{\lambda}(p_j+\mu-2)}}{t^{\frac{N}{\lambda}(p_j+\mu-2)}} dt \\ &\leq C \|u_0\|_1^{1+\frac{\bar{p}}{\lambda}(p_j+\mu-2)} T^{1-\frac{N}{\lambda}(p_j+\mu-2)} \end{aligned}$$

where we recall that $\lambda = \lambda_1 = N(\bar{p}-2) + \bar{p}$ and, integrating in time at the last line, we assumed

$$\frac{N}{\lambda}(p_j + \mu - 2) < 1 \quad \Leftrightarrow \quad \mu < \bar{p}_1 - p_j,$$

the latter being positive due to (5.2). The previous discussion shows that if r and T obey

$$\|u_0\|_1^{1+\frac{\bar{p}}{\lambda}(p_j+\mu-2)} T^{1-\frac{N}{\lambda}(p_j+\mu-2)} \leq C r^{p_j \left(1 + \frac{\lambda_1 + \mu}{(p_j - 2)\bar{p}}\right)} T^{\frac{N}{\bar{p}} - \frac{\lambda_1 + \mu}{(p_j - 2)\bar{p}}}$$

for some constant C depending only on the data and on μ , then (5.10) holds. This inequality can be rewritten through some algebra as

$$r \geq CT^{\frac{N(\bar{p}-p_j)+\bar{p}}{\lambda p_j}} \|u_0\|_1^{\frac{\bar{p}}{p_j} \frac{p_j-2}{\lambda}}.$$

Thus (5.10) holds for any $r \geq 2R_0$ satisfying the previous one-sided inequality, implying (5.3).

Finally, we construct the claimed branch, removing assumption (5.4) in doing so. Notice that (5.2) implies (4.6), hence Lemma 5.1 ensures the existence of $\tau(u, 2R_0, T) > 0$ such that $u(\cdot, t) \equiv 0$ on $\mathbb{R}^{N-M} \times (K''_{4R_0} \setminus K''_{2R_0})$ for all $t \in [0, \tau]$. Then we define for $t \in [0, \tau]$

$$\tilde{u}(x, t) = \begin{cases} u(x, t) & \text{if } x \in \mathbb{R}^{N-M} \times K''_{3R_0}, \\ 0 & \text{otherwise,} \end{cases}$$

which is a non-zero branch of u on S_τ satisfying (5.4), due to

$$\int_{S_\tau} |\tilde{u}|^{p_i} dx dt = \int_0^\tau \int_{\mathbb{R}^{N-M} \times K''_{3R_0}} |u|^{p_i} dx' dx'' dt < +\infty$$

for all $i = 1, \dots, N$, by the assumption $u \in \cap_{i=1}^N L_{\text{loc}}^{p_i}(\mathbb{R}^M; L^{p_i}(\mathbb{R}^{N-M} \times [0, T]))$. Suppose that

$$T^* := \sup \{ \tau > 0 : u \text{ has a branch in } \cap_{i=1}^N L^{p_i}(S_\tau) \} < T,$$

and let

$$\bar{R} = \max\{R_j(T) : j = N - M + 1, \dots, N\}, \quad \varepsilon = \frac{1}{2}\tau(u, 2\bar{R}, T)$$

where τ is given in Lemma 5.1. Then, by (5.3), on the whole $S_{T^*-\varepsilon}$, u has a branch supported in $\mathbb{R}^{N-M} \times K''_{\bar{R}}$. Applying Lemma 5.1 as in the first step, \tilde{u} can be extended up to $S_{T^*+\varepsilon}$ staying in $\cap_{i=1}^N L^{p_i}(S_{T^*+\varepsilon})$, contradicting $T^* < T$. Thus $T^* = T$, there exists a branch $\tilde{u} \in \cap_{i=1}^N L^{p_i}(S_T)$, which therefore satisfies (5.3). \square

APPENDIX A. AN EXAMPLE Á LA TIKHONOV

In this appendix we will construct a nontrivial solution to the Cauchy problem

$$\begin{cases} u_t = \operatorname{div}(|\nabla u|^{p-2}\nabla u) & \text{in } \mathbb{R}^N \times]0, +\infty[, \\ u(\cdot, t) \rightarrow 0 & \text{as } t \rightarrow 0 \text{ in } L^2_{\text{loc}}(\mathbb{R}^N), \end{cases}$$

in the slow diffusion case $p > 2$. Suppose that $\alpha, \beta > 0$ satisfy

$$(A.1) \quad (p-2)\alpha = 1 + p\beta.$$

Then the equation $u_t = \Delta_p u$ in $\mathbb{R} \times]0, +\infty[$ with the ansatz $u(x, t) = t^{-\alpha}U(x t^\beta)$ reduces to

$$-\alpha U(s) + \beta U'(s) s = (|U'(s)|^{p-2}U'(s))', \quad s = x t^\beta$$

and letting $V = |U'|^{p-2}U'$, the latter can be rewritten as the non-autonomous ODE system

$$(A.2) \quad \begin{cases} U' = |V|^{\frac{2-p}{p-1}}V \\ V' = -\alpha U + \beta |V|^{\frac{2-p}{p-1}}V s. \end{cases}$$

Our aim is to construct a nontrivial solution for $s \in [1, +\infty[$ to this system with initial condition $U(1) = V(1) = 0$. Notice that $U \equiv V \equiv 0$ is certainly a solution, but the right-hand side of (A.2) is only Hölder continuous in V and the standard Picard uniqueness cannot be applied. We also observe that a major rôle in the following construction is some *anti-dissipative* feature of the system. Indeed, it is well known that, even if the ODE system $\mathbf{x}' = \mathbf{F}(s, \mathbf{x})$ has only continuous right hand side, the dissipativity condition

$$(\mathbf{F}(s, \mathbf{x}) - \mathbf{F}(s, \mathbf{y})) \cdot (\mathbf{x} - \mathbf{y}) \leq 0$$

is enough to ensure uniqueness of the Cauchy problem. Luckily, system (A.2) satisfies, for $\mathbf{x} = (0, V)$ and $\mathbf{y} = (0, 0)$ the opposite inequality $\mathbf{F}(s, 0, V) = \beta |V|^p s > 0$ near $s = 1$.

To construct the solution, first observe that it is enough to do it locally, since the right hand side of (A.2) has sublinear growth and by Gronwall's lemma any local solution can be extended to a global one. A local (forward) solution of (A.2) is a fixed point of the operator

$$T(\mathbf{x})(s) = \int_1^s \mathbf{F}(s, \mathbf{x}(s)) ds,$$

on the space

$$X_\delta = \{\mathbf{x} \in C([1, 1 + \delta], \mathbb{R}^2) : \mathbf{x}(1) = (0, 0)\}$$

with the standard uniform norm denoted by $\|\cdot\|_\delta$, where

$$\mathbf{x} = (x_1, x_2), \quad \mathbf{F}(s, \mathbf{x}) = (|x_2|^{\frac{2-p}{p-1}}x_2, -\alpha x_1 + \beta |x_2|^{\frac{2-p}{p-1}}x_2 s).$$

By the standard proof of the Peano existence theorem, there exists $\bar{\delta} \in]0, 1[$ and $M, L > 0$ such that, given any \bar{x} and $\delta \in]0, \bar{\delta}[$ with $\|\bar{x}\|_\delta \leq M$, the sequence

$$\mathbf{x}_0 = \bar{\mathbf{x}}, \quad \mathbf{x}_{n+1} = T(\mathbf{x}_n)$$

satisfies $\|\mathbf{x}_n\|_\delta \leq M$ for all $n \geq 0$, is L -equilipschitz for $n \geq 1$ and converges in X_δ to a fixed point of T .

We employ a sub-supersolution method with a nontrivial subsolution. The heuristic is that our Cauchy problem for (A.2) is, near $s = 1$, asymptotically equivalent to

$$\begin{cases} x_1' = |x_2|^{\frac{2-p}{p-1}} x_2, \\ x_2' = \beta |x_2|^{\frac{2-p}{p-1}} x_2, \end{cases} \quad x_1(1) = x_2(1) = 0,$$

whose maximal nontrivial solution is, up to multiplicative constants, $x_2(s) \simeq x_1(s) \simeq (s-1)^{\frac{p-1}{p-2}}$. For $0 < a \leq b$, $\delta \in]0, \bar{\delta}[$, consider the closed convex set

$$C_{\delta,a,b} := \left\{ \mathbf{x} \in X_\delta : 0 \leq x_1(s) \leq b(s-1)^{\frac{p-1}{p-2}}, a(s-1)^{\frac{p-1}{p-2}} \leq x_2(s) \leq b(s-1)^{\frac{p-1}{p-2}}, s \in [1, 1+\delta] \right\}.$$

We claim that $C_{\delta,a,b}$ is T -invariant for suitable choices of the parameters. If $T_i(x_1, x_2)$ is the i -th coordinate of $T(x_1, x_2)$, $i = 1, 2$, given $\mathbf{x} = (x_1, x_2) \in C_\delta$ we estimate for any $s \in]1, 1+\delta[$

$$\begin{aligned} T_2(x_1, x_2)(s) &= \int_1^s -\alpha x_1(\tau) + \beta x_2^{\frac{1}{p-1}}(\tau) \tau d\tau \\ &\geq -\alpha b \int_1^s (\tau-1)^{\frac{p-1}{p-2}} d\tau + \beta a^{\frac{1}{p-1}} \int_1^s (\tau-1)^{\frac{1}{p-2}} d\tau \\ (A.3) \quad &\geq -\alpha b \frac{p-2}{2p-3} (s-1)^{\frac{2p-3}{p-2}} + \beta a^{\frac{1}{p-1}} \frac{p-2}{p-1} (s-1)^{\frac{p-1}{p-2}}; \end{aligned}$$

$$(A.4) \quad T_2(x_1, x_2)(s) \leq \beta \int_1^s x_2^{\frac{1}{p-1}}(\tau) \tau \leq (1+\delta) \beta b^{\frac{1}{p-1}} \frac{p-2}{p-1} (s-1)^{\frac{p-1}{p-2}};$$

$$(A.5) \quad 0 \leq T_1(x_1, x_2)(s) = \int_1^s x_2^{\frac{1}{p-1}}(\tau) d\tau \leq b^{\frac{1}{p-1}} \int_1^s (\tau-1)^{\frac{1}{p-2}} d\tau \leq b^{\frac{1}{p-1}} \frac{p-2}{p-1} (s-1)^{\frac{p-1}{p-2}}.$$

Using $\delta \leq \bar{\delta} \leq 1$ in the upper bound (A.3) for T_2 , we first define b as per

$$\frac{p-1}{p-2} b^{\frac{p-2}{p-1}} = \max\{1, 2\beta\},$$

which, inserted into (A.4) and (A.5) gives through some algebra

$$(A.6) \quad T_2(x_1, x_2)(s) \leq b(s-1)^{\frac{p-1}{p-2}} \quad T_1(x_1, x_2)(s) \leq b(s-1)^{\frac{p-1}{p-2}}.$$

Then we choose $\delta \in]0, \bar{\delta}[$ such that

$$-\alpha b \frac{p-2}{2p-3} (s-1)^{\frac{2p-3}{p-2}} + \beta a^{\frac{1}{p-1}} \frac{p-2}{p-1} (s-1)^{\frac{p-1}{p-2}} \geq \frac{\beta}{2} a^{\frac{1}{p-1}} \frac{p-2}{p-1} (s-1)^{\frac{p-1}{p-2}}$$

(this is possible because from $p > 2$ we infer $(2p-3)/(p-2) > (p-1)/(p-2)$). Finally, we set

$$\frac{p-1}{p-2} a^{\frac{p-2}{p-1}} = \frac{\beta}{2},$$

(notice that $a \leq b$), to deduce from (A.4) the lower bound

$$T_2(x_1, x_2)(s) \geq a(s-1)^{\frac{p-1}{p-2}}.$$

The latter inequality, together with (A.6) and the trivial one $T_1(x_1, x_2) \geq 0$ concludes the proof of the claimed T -invariance of $C_{\delta,a,b}$. The starting point

$$\bar{\mathbf{x}}(s) = \left(0, a(s-1)^{\frac{p-1}{p-2}}\right)$$

satisfies $\|\bar{\mathbf{x}}\| \leq M$ by eventually further reducing δ , and thus produces by iteration a nontrivial solution (U, V) to (A.2) with initial conditions $U(1) = V(1) = 0$.

As remarked before, the solution can be extended to the whole $[1, +\infty[$, and it remains to observe that $\text{supp}(U) = [1, +\infty[$, due to the inequality

$$\left(\alpha U^2 + \frac{p-1}{p} |V|^{\frac{p}{p-1}}\right)' = \beta |V|^{\frac{2}{p-1}} s \geq 0.$$

For any $\alpha, \beta > 0$ obeying (A.1), such a trajectory defines a solution to $u_t = \Delta_p u$ in $\mathbb{R}^N \times]0, +\infty[$ through

$$u(x, t) = \begin{cases} t^{-\alpha} U(x_1 t^\beta) & \text{if } x_1 t^\beta \geq 1 \\ 0 & \text{otherwise,} \end{cases}$$

which has the claimed properties, since its support is $\{x_1 \geq t^{-\beta} > 0\}$.

Acknowledgement. We thank J. L. Vázquez (U. Autónoma de Madrid) for a discussion on non-uniqueness phenomena for the Cauchy problem. S. Mosconi and V. Vespri are members of GNAMPA (Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni) of INdAM (Istituto Nazionale di Alta Matematica). F. G. Düzgün is partially funded by Hacettepe University BAP through project FBI-2017-16260; S. Mosconi is partially funded by the grant PdR 2016-2018 - linea di intervento 2: “Metodi Variazionali ed Equazioni Differenziali” of the University of Catania.

REFERENCES

- [1] D. ANDREUCCI, S. F. TEDEEV, Finite speed of propagation and other higher-order parabolic equations with general nonlinearity. *Interf. Free Bound.* **3** (2001), 233–264. [3](#)
- [2] S. ANTONTSEV, S. SHMAREV, Evolution PDEs with Nonstandard Growth Conditions. Atlantis Studies in Differential Equations **4**, Atlantis Press, Paris, 2015. [2](#), [3](#), [4](#)
- [3] S. ANTONTSEV, S. SHMAREV, Localization of solutions of anisotropic parabolic equations. *Nonlinear Anal.* **71** (2009), 725–737. [5](#)
- [4] D. G. ARONSON, Widder’s inversion theorem and the initial distribution problem. *SIAM J. Math. Anal.* **12** (1981), 639–651. [2](#)
- [5] F. BERNIS, Qualitative properties for some nonlinear higher order degenerate parabolic equations. *Houston J. Math.* **14** (1988), 319–352. [3](#)
- [6] M. F. BIDAUT-VÉRON, Self-similar solutions of the p -Laplace heat equation: the case when $p > 2$. *Proc. Roy. Soc. Edinburgh Sect. A* **139** (2009), 1–43. [2](#), [3](#)
- [7] L. BOCCARDO, P. MARCELLINI, C. SBORDONE, L^∞ -regularity for variational problems with sharp non-standard growth conditions. *Boll. Unione Mat. Ital.* VII. Ser. A **4** (1990), 219–225. [2](#)
- [8] V. BÖGELEIN, F. RAGNEDDA, S. VERNIER PIRO, V. VESPRI, Moser–Nash kernel estimates for degenerate parabolic equations. *J. Func. Anal.* **272** (2017), 2956–2986. [3](#)
- [9] S. P. DEGTYAREV, A. F. TEDEEV, $L^1 - L^\infty$ estimates of the solution of the Cauchy problem for an anisotropic degenerate parabolic equation with double nonlinearity and growing initial data. *Mat. Sb.* **198** (2007), 46–66. [2](#)
- [10] S. P. DEGTYAREV, A. F. TEDEEV, On the solvability of the Cauchy problem with growing initial data for a class of anisotropic parabolic equations. *J. Math. Sci.* **181** (2012), 28–46. [2](#), [3](#)
- [11] E. DiBENEDETTO, Degenerate Parabolic Equations. *Universitext*, Springer-Verlag New York, 1993. [2](#), [11](#), [17](#)
- [12] E. DiBENEDETTO, U. GIANAZZA AND V. VESPRI, Harnack’s inequality for degenerate and singular parabolic equations. *Springer Monographs in Mathematics*, Springer-Verlag New York, 2012. [2](#), [5](#)
- [13] E. DiBENEDETTO, U. GIANAZZA AND V. VESPRI, Remarks on Local Boundedness and Local Holder Continuity of Local Weak Solutions to Anisotropic p -Laplacian Type Equations. *J. Elliptic Parabol. Equ.* **2** (2016) 157–169. [2](#)

- [14] E. DiBENEDETTO, M. A. HERRERO, On the Cauchy problem and initial traces for a degenerate parabolic equation. *Trans. AMS* **314** (1989), 187–224. [2](#)
- [15] E. DiBENEDETTO, M. A. HERRERO, Non negative solutions of the evolution p -Laplacian equation. Initial traces and Cauchy problem when $1 < p < 2$. *Arch. Rational Mech. Anal.* **111** (1990), 225–290. [2](#)
- [16] M. ELEUTERI, P. MARCELLINI, E. MASCOLO, Regularity for scalar integrals without structure conditions. To appear in *Adv. Calc. Var.*, DOI: 10.1515/acv-2017-0037. [2](#)
- [17] N. FUSCO, C. SBORDONE, Local boundedness of minimizers in a limit case. *Manuscr. Math.* **69** (1990), 19–25. [2](#)
- [18] M. GIAQUINTA, Growth conditions and regularity. A counterexample. *Manuscr. Math.* **59** (1987), 245–248. [2](#)
- [19] J. HAŠKOVEC, C. SCHMEISER, A note on the anisotropic generalizations of the Sobolev and Morrey embedding theorems. *Monatsh. Math.*, **158** (2009), 71–79. [9](#)
- [20] M.-C. HONG, Existence and partial regularity in the calculus of variations. *Ann. Mat. Pura Appl.*, **149** (1987), 311–328. [2](#)
- [21] J. HULSHOF, Similarity solutions for the porous medium equation with sign changes. *J. Math. Anal. Appl.* **157** (1991), 75111. [2](#), [3](#)
- [22] R. G. IAGAR, A. SÁNCHEZ, J. L. VÁZQUEZ, Radial equivalence for the two basic nonlinear degenerate diffusion equations. *J. Math. Pures Appl.*, **89** (2008), 1–24. [2](#)
- [23] S. N. KRUIZHKOVA, I. M. KOLODĪ, On the theory of embedding of anisotropic Sobolev spaces. *Uspekhi Mat. Nauk*, **38**, (1983), 207–208 (in Russian). English transl.: *Russian Math. Surveys* **38** (1983), 188–189. [2](#), [9](#)
- [24] T. KUUSI, Lower semicontinuity of weak supersolutions to nonlinear parabolic equations. *Differential Integral Equations* **22** (2009), 1211–1222. [19](#)
- [25] J. L. LIONS, Quelques méthodes de résolution des problèmes aux limites non linéaires. Dunod, Paris, 1969. [3](#), [7](#), [8](#)
- [26] P. MARCELLINI, Un esempio di soluzione discontinua di un problema variazionale nel caso scalare. *Ist. Mat. “U. Dini”* No. 11, Firenze, 1987. (In Italian). [2](#)
- [27] G. MINGIONE, Regularity of minima: an invitation to the Dark Side of the Calculus of Variations. *Appl. Math.* **51** (2006) 355–426. [2](#)
- [28] A. F. TEDEEV, V. VESPRI, Optimal behavior of the support of the solutions to a class of degenerate parabolic systems. *Interfaces Free Bound.* **17** (2015), 143–156. [3](#)
- [29] M. TROISI, Teoremi di inclusione per spazi di Sobolev non isotropi, *Ricerche Mat.* **18** (1969), 3–24. (In Italian). [4](#), [7](#)
- [30] A. N. TYCHONOV, Théorèmes d’unicité pour l’équation de la chaleur. *Mat. Sb.* **42** (1935), 199–216. [2](#)
- [31] M. YU, X. LIAN, Boundedness of solutions of parabolic equations with anisotropic growth conditions. *Can. J. Math.* **49** (1997), 798–809. [2](#)
- [32] J. L. VÁZQUEZ, New selfsimilar solutions of the porous medium equation and the theory of solutions with changing sign. *Nonlinear Analysis* **15** (1990), 931–942. [2](#)
- [33] D. V. WIDDER, Positive temperatures in an infinite rod. *Trans. Amer. Math. Soc.* **55** (1944), 85–95.

[2](#)

(F. G. Düzgün) DEPARTMENT OF MATHEMATICS,
HACETTEPE UNIVERSITY, 06800, BEYTEPE, ANKARA, TURKEY
E-mail address: gamzeduz@hacettepe.edu.tr

(S. Mosconi) DIPARTIMENTO DI MATEMATICA E INFORMATICA,
UNIVERSITÀ DEGLI STUDI DI CATANIA, VIALE A. DORIA 6, 95125 CATANIA, ITALY
E-mail address: mosconi@dmi.unict.it

(V. Vespri) DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”,
UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY
E-mail address: vespri@math.unifi.it