The isoperimetric spectrum of finitely presented groups

M.V. Sapir^{*}

Abstract

The isoperimeric spectrum consists of all real positive numbers α such that $O(n^{\alpha})$ is the Dehn function of a finitely presented group. In this note we show how a recent result of Olshanskii completes the description of the isoperimetric spectrum modulo the celebrated Computer Science conjecture (and one of the seven Millennium Problems) $\mathbf{P} = \mathbf{NP}$ and even a formally weaker conjecture.

The goal of this note is to show that the recent paper by Olshanskii [7] completes a description of the isoperimetric spectrum of finitely presented groups modulo the $\mathbf{P} = \mathbf{NP}$ conjecture.

Since in this note we consider only polynomially bounded functions $\mathbb{N} \to \mathbb{R}$, we call two functions f, g equivalent if $af(n) \leq g(n) \leq bf(b)$ for some positive constants a, b.

Recall that Brady and Bridson [5] called the set of all real numbers $\alpha \geq 1$ such that n^{α} is equivalent to the Dehn function of a finitely presented group the *isoperimetric spectrum*. When it was introduced, it was known only that all natural numbers belong to the isoperimetric spectrum (the free nilpotent group of class c with at least 2 generators has Dehn function n^{c+1} [1]), and that by Gromov's theorem the intersection of the isoperimetric spectrum with the open interval (1, 2) is empty. It is obvious also that the isoperimetric spectrum is a countable set since the set of all finite group presentation is countable. Bridson [5] found the first examples of non-integral numbers in the spectrum.

Soon after, we proved in [10] that for $\alpha \geq 4$ to be in the isoperimetric spectrum, it is enough that α is computed in time $\leq 2^{2^m}$. Recall [10] that a real number α is called computable in time T(m) where T(m) is a function $\mathbb{N} \to \mathbb{N}$, if there exists a deterministic Turing machine which, given a natural number n, computes a binary rational approximation of α with an error at most $1/2^{n+1}$ in at most T(n) steps. Thus all algebraic numbers ≥ 4 and many transcendental numbers such as $\pi + 1$ are in the isoperimetric spectrum. On the other hand, we proved in [10] that every number in the isoperimetric spectrum can be computed in time $\leq 2^{2^{2^{cm}}}$ for some constant c. It can be seen from the proof of this result that the number of 2's in this estimate can be reduced to two as in the lower bound if we had $\mathbf{P} = \mathbf{NP}$. I mentioned it (without a proof) in my ICM talk [8] and my Bulletin of Mathematical Sciences survey [9].

For $\alpha \leq 4$, the situation was more complicated. On the one hand the tools used in [10] were too weak to handle $\alpha \leq 4$. On the other hand, Brady, Bridson, Forester and Shankar found more numbers from the interval (2, 4) in the isoperimetric spectrum, showing that the set of these numbers is dense in the interval (2, 4) [2] and even contains all rational numbers [3]. Their numbers from the isoperimetric spectrum were constructed using

^{*}Supported in part by the NSF grant DMS-1500180.

[†]Key words: Dehn function, finitely presented group, isoperimetric spectrum, P=NP

[‡]AMS Mathematical Subject Classification: 20F05, 20F06, 20F65, 20F69, 03D10.

algebraic rather than computational properties. (Note also that the groups constructed in [5, 2, 4, 3], are given by very small presentations comparing to the groups in [10] and are subgroups of CAT(0) groups which is quite remarkable.) But the paper by Olshanskii [7] showed that the intersection of the isoperimetric spectrum with (2, 4) can be described in the same terms as in [10].

Combining results of [10] and [7] we get:

Theorem 1 (The first part of Corollary 1.4 from [10], and Corollary 1.4 from [7]). If a number $\alpha \geq 2$ can be computed in time $\leq 2^{c2^m}$ for some c, then α belongs to the isoperimetric spectrum.

Notice that one needs to modify a little the proof of the first part of Corollary 1.4 from [10] to obtain the estimate $\leq 2^{c2^m}$ instead of $\leq 2^{2^m}$. For this, one should take natural number $d > \log_2 c$ which is a power 2^k for some k and consider d-ary representations of numbers instead of binary representations as in [10]. Each d-ary digit of α is k binary digits. So if the first m binary digits of α are computed in time $\leq 2^{c2^m}$, then the first m d-ary digits of α are computed in time at most d^{d^m} and the rest of the proof of [10, Corollary 1.4] carries by replacing 2 by d everywhere.

Now we will prove the main result of the note.

Theorem 2. Provided $\mathbf{P} = \mathbf{NP}$, a number α is in the isoperimetric spectrum if and only if it can be computed in time $\leq 2^{c2^m}$ for some $c \geq 1$.

Proof. Theorem 1 gives one part of Theorem 2.

To prove the other part, suppose that n^{α} is the Dehn function of a finitely presented group. Then by [10, Theorem 1.1], n^{α} is equivalent to the time function T(n) of some (non-deterministic) Turing machine M which recognizes the word problem in a finitely presented group.

As explained by Emil Jeřábek [6], the following property follows from $\mathbf{P} = \mathbf{NP}$:

(*) There is a deterministic Turing machine M' computing a function T'(n) which is equivalent to T(n) and having time function at most $T(n)^d$ for some constant d.

Since $\alpha \geq 2$, for any n > 0

 $\epsilon_1 n^{\alpha} \le T'(n) \le \epsilon_2 n^{\alpha}$

for some positive constants $\epsilon_1 \leq 1$ and $\epsilon_2 \geq 1$. Let number n_0 be such that $2^n > \log_2(\epsilon_2/\epsilon_1)$ for every $n \geq n_0$. Let $q = [\alpha] + 1$.

Given this machine M' with, say, k tapes, consider the following Turing machine M''which will calculate the first m digits of α (for every m). This machine has k + 3 tapes with tape k + 3 being the input tape. It starts with number m in binary written on tape k + 3 and all other tapes empty. Then it calculates the number $n = 2^{2^{m+n_0}}$ and writes it on tape k + 1 (using tape k + 2 as an auxiliary tape and cleaning it after nis computed). Then M'' turns on the machine M' and produces T'(n) on tape k + 2. Then it calculates $p = [(\log_2 T'(n) - \log_2 \epsilon_1)/2^{n_0}]$ and writes it on tape k + 2. Notice that $\alpha \log_2 n + \log_2 \epsilon_1 \leq \log_2 T'(n) \leq \alpha \log_2 n + \log_2 \epsilon_2$. Therefore

$$[\alpha 2^m] \le p \le \alpha 2^m + \log_2(\epsilon_2/\epsilon_1)/2^{n_0}.$$

Hence $p = [\alpha 2^m]$, so $p/2^m$ a rational approximation of α which is within $1/2^m$ from α . From the construction of M'', it is clear that the time complexity of M'' does not exceed 2^{c2^m} for some constant c.

As explained by Emil Jeřábek [6], Property (*) follows also from the property $\mathbf{E} = \Sigma_2^{\mathbf{E}}$, which is not known to imply $\mathbf{P} = \mathbf{NP}$. Here \mathbf{E} denotes the class of languages recognized by deterministic Turing machines in time $2^{O(n)}$ and $\Sigma_2^{\mathbf{E}}$ is the second level of the exponential hierarchy (with linear exponent). Thus in Theorem 2, one can replace $\mathbf{P} = \mathbf{NP}$ by $\mathbf{E} = \Sigma_2^{\mathbf{E}}$.

Acknowledgement. I am grateful to Emil Jeřábek for answering my question [6] and several useful remarks. I am also grateful to Jean-Camille Birget and A. Yu. Olshanskii for their comments.

References

- G. Baumslag, C. F. Miller III, and H. Short, Isoperimetric inequalities and the homology of groups, Invent. Math. 113 (1993), 531–560.
- [2] N. Brady and M. Bridson, There is only one gap in the isoperimetric spectrum, Geometric and Functional Analysis, 10 (2000), 1053–1070.
- [3] Noel Brady, Martin R. Bridson, Max Forester, Krishnan Shankar, Snowflake groups, Perron-Frobenius eigenvalues and isoperimetric spectra. Geom. Topol. 13 (2009), no. 1, 141–187.
- [4] Noel Brady, Max Forester, Snowflake geometry in CAT(0) groups. J. Topol. 10 (2017), no. 4, 883–920.
- [5] M. Bridson, Fractional isoperimetric inequalities and subgroup distortion. J. Amer. Math. Soc. 12 (1999), no. 4, 1103–1118.
- [6] Mathoverflow, question 307629.
- [7] A. Yu. Olshanskii, Polynomially-bounded Dehn functions of groups, arXiv:1710.00550, accepted in Journal of Combinatorial Algebra, 2018.
- [8] M. V. Sapir, Algorithmic and asymptotic properties of groups. International Congress of Mathematicians. Vol. II, 223–244, Eur. Math. Soc., Zürich, 2006.
- [9] M. V. Sapir, Asymptotic invariants, complexity of groups and related problems. Bull. Math. Sci. 1 (2011), no. 2, 277–364.
- [10] M. V. Sapir, J. C. Birget, E. Rips, Isoperimetric and isodiametric functions of groups, Annals of Mathematics, 157, 2(2002), 345-466.