

# RATE OF CONVERGENCE FOR PERIODIC HOMOGENIZATION OF CONVEX HAMILTON-JACOBI EQUATIONS IN ONE DIMENSION

SON N.T. TU

ABSTRACT. In this paper we study the rate of convergence  $u^\varepsilon \rightarrow u$  as  $\varepsilon \rightarrow 0^+$  in periodic homogenization of Hamilton-Jacobi equations. Here  $u^\varepsilon$  and  $u$  are viscosity solutions to the oscillatory Hamilton-Jacobi equation

$$\begin{cases} u_t^\varepsilon + H\left(x, \frac{x}{\varepsilon}, Du^\varepsilon\right) = 0 & \text{in } \mathbb{R} \times [0, \infty) \\ u^\varepsilon(x, 0) = u_0(x) & \text{on } \mathbb{R}, \end{cases}$$

and its effective equation

$$\begin{cases} u_t + \bar{H}(x, Du) = 0 & \text{in } \mathbb{R} \times [0, \infty) \\ u(x, 0) = u_0(x) & \text{on } \mathbb{R}, \end{cases}$$

respectively. Assuming that the initial data  $u_0$  is bounded and Lipschitz continuous, we provide a simple proof to get the optimal rate  $\mathcal{O}(\varepsilon)$  for a class of Hamiltonians including the classical mechanics one with separable potential

$$H(x, y, p) = \frac{|p|^2}{2} + (a(x)b(y) + C) \quad \text{for } (x, y, p) \in \mathbb{R}^3,$$

where  $a(\cdot) : \mathbb{R} \rightarrow (0, +\infty)$  is continuously differentiable,  $b(\cdot) : \mathbb{R} \rightarrow (-\infty, 0]$  is continuous and 1-periodic.

## 1. INTRODUCTION

We first give a brief description of the periodic homogenization theory for Hamilton-Jacobi equations in the framework of viscosity solutions (see [7, 11, 2] or appendix of [10]). For each  $\varepsilon > 0$ , let  $u^\varepsilon \in C(\mathbb{R}^n \times [0, \infty))$  be the viscosity solution to the following Hamilton-Jacobi equation:

$$\begin{cases} u_t^\varepsilon(x, t) + H\left(x, \frac{x}{\varepsilon}, Du^\varepsilon(x, t)\right) = 0 & \text{in } \mathbb{R}^n \times (0, \infty), \\ u^\varepsilon(x, 0) = u_0(x) & \text{on } \mathbb{R}^n. \end{cases} \quad (\text{C}_\varepsilon)$$

The initial data  $u_0 \in \text{BUC}(\mathbb{R}^n)$ , the set of bounded, uniformly continuous functions on  $\mathbb{R}^n$  and the Hamiltonian  $H = H(x, y, p) \in C(\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n)$  satisfies the following:

- (H1) For each  $(x, p) \in \mathbb{R}^n \times \mathbb{R}^n$ ,  $y \mapsto H(x, y, p)$  is  $\mathbb{Z}^n$ -periodic.
- (H2)  $p \mapsto H(x, y, p)$  is coercive uniformly in  $(x, y) \in \mathbb{R}^n \times \mathbb{T}^n$ , that is,

$$\lim_{|p| \rightarrow +\infty} \left( \inf_{(x, y) \in \mathbb{R}^n \times \mathbb{T}^n} H(x, y, p) \right) = +\infty,$$

---

2010 *Mathematics Subject Classification.* 35B40, 37J50, 49L25 .

*Key words and phrases.* Cell problems; Periodic homogenization; first order Hamilton-Jacobi equations; Rate of convergence; viscosity solutions.

where  $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$ .

(H3)  $\sup \{|H(x, y, p)| : (x, y) \in \mathbb{R}^{2n}, |p| \leq R\} < \infty$  for all  $R > 0$ .

(H4) For each  $R > 0$ , there exists  $\omega_R(\cdot) \in C([0, \infty))$ , with  $\omega_R(0) = 0$ , such that for all  $x, y \in \mathbb{R}^n, p, q \in B(0, R)$  then

$$|H(x, y, p) - H(x, y, q)| \leq \omega_R(|p - q|)$$

where  $B(0, R)$  denotes the open ball centered at 0 with radius  $R$  in  $\mathbb{R}^n$ .

Some first general results on homogenization of Hamilton-Jacobi equations are due to P.-L. Lions, G. Papanicolau and S.R.S. Varadhan [12] for the case  $H$  is independent of  $x$ , namely  $H(x, y, p) = H(y, p)$ . The more general case  $H = H(x, y, p)$  was considered later by L. C. Evans [8, 9], who developed the perturbed test functions methods for studying the homogenization problem in the framework of the theory of viscosity solutions. The main result is, under assumptions (H1)–(H4) on  $H$ ,  $u^\varepsilon \rightarrow u$  locally uniformly on  $\mathbb{R}^n \times [0, \infty)$  as  $\varepsilon \rightarrow 0^+$  and  $u$  solves the following effective equation

$$\begin{cases} u_t(x, t) + \overline{H}(x, Du(x, t)) = 0 & \text{in } \mathbb{R}^n \times (0, \infty), \\ u(x, 0) = u_0(x) & \text{on } \mathbb{R}^n. \end{cases} \quad (\text{C})$$

The effective Hamiltonian  $\overline{H}(x, p) : \mathbb{R}^{2n} \rightarrow \mathbb{R}$  is determined by  $H$  in a very nonlinear way through the cell problem as follows: For each  $(x, p) \in \mathbb{R}^n \times \mathbb{R}^n$ , it can be shown (see [12] and [8, 9]) that there is a unique constant  $\lambda = \lambda(x, p) \in \mathbb{R}$  for which the following cell problem

$$H(x, y, p + D_y v(y)) = \lambda \quad \text{in } \mathbb{T}^n \quad (\text{CP})$$

has a continuous solution  $v(y) = v(y; x, p)$ . The effective Hamiltonian  $\overline{H}(x, p)$  is then defined by setting

$$\overline{H}(x, p) = \lambda.$$

It is worth mentioning that in general  $v(y; x, p)$  is not unique even up to adding a constant. More research in understanding the effective Hamiltonians  $\overline{H}$  is reported in [5, 6, 13, 18] and the references therein.

The main goal of this paper is to obtain optimal rate of convergence of  $u^\varepsilon$  to  $u$  in one dimension, that is, an optimal bound for  $\|u^\varepsilon - u\|_{L^\infty([-R, R] \times [0, T])}$  for any given  $R, T > 0$  as  $\varepsilon \rightarrow 0^+$ . Heuristically, owing to the two-scale asymptotic expansion (see [16])

$$u^\varepsilon(x, t) \approx u(x, t) + \varepsilon v\left(\frac{x}{\varepsilon}; x, Du(x, t)\right) + \mathcal{O}(\varepsilon^2),$$

the optimal rate looks like  $\mathcal{O}(\varepsilon)$ . However, as pointed out in [16] by H. Mitake, H. V. Tran and Y. Yu, it is hard to justify this expansion rigorously due to two reasons:

- In general, there does not even exist a continuous selection of  $v(\cdot; x, p)$  with respect to  $p$ , let alone Lipschitz continuous selection.
- The solution  $u(x, t)$  to (C) is only Lipschitz in  $(x, t)$ , and is usually not  $C^1$ .

The rate of convergence  $u^\varepsilon \rightarrow u$  was first studied by I. Capuzzo-Dolcetta and H. Ishii in [4] using a PDE approach. They established the uniform rate  $\mathcal{O}(\varepsilon^{1/3})$  under general assumptions on  $H$  (e.g., (H1)–(H4)), including non-convex Hamiltonians.

To be precise, [4] deals with stationary problems, but the approach can be easily adjusted to handle the Cauchy problem.

Recently, in [16], H. Mitake, H. V. Tran and Y. Yu considered the case that  $H(x, y, p)$  does not depend on  $x$ , namely  $H(x, y, p) = H(y, p)$ . They established an optimal rate  $\mathcal{O}(\varepsilon)$  in the one dimensional case with convex Hamiltonians along with other important results in higher dimensional spaces using tools from dynamical systems and weak KAM theory. They also presented an essential obstacle to improve the convergence rate  $\mathcal{O}(\varepsilon^{1/3})$  by the method in [4]. See also [1, 3, 14, 15] and the references therein for related results on rate of convergence of Hamilton-Jacobi equations in various settings.

Within the convex setting, in the case  $H$  depends on  $x$ , namely  $H = H(x, y, p)$ , the situation is more complicated and requires harder analysis of the dynamics of optimal paths in the optimal control formula. Up to now, the best known convergence rate in this setting is  $\mathcal{O}(\varepsilon^{1/3})$  obtained in [4].

In this paper, we consider the one dimensional case  $n = 1$  and the convex Hamiltonian is of the form:

$$H(x, y, p) = H(p) + V(x, y) \quad \text{for all } (x, y, p) \in \mathbb{R} \times \mathbb{T} \times \mathbb{R}.$$

The main results are the following theorem and its generalization.

**Theorem 1.1.** *Assume  $n = 1$  and  $H(x, y, p) = \frac{1}{2}|p|^2 + V(x, y)$  where  $V$  is of the separable form  $V(x, y) = a(x)b(y) + C_0$  where  $C_0$  is a constant and*

- (i)  $a(x) \in C^1(\mathbb{R})$  is bounded with  $a(x) > 0$  for all  $x \in \mathbb{R}$ ,
- (ii)  $b(y) \in C(\mathbb{T})$  and  $\max_{y \in \mathbb{T}} b(y) = 0$ .

Assume  $u_0 \in \text{Lip}(\mathbb{R}) \cap \text{BUC}(\mathbb{R})$ , then for each  $R, T > 0$  we have

$$\|u^\varepsilon - u\|_{L^\infty([-R, R] \times [0, T])} \leq C\varepsilon \tag{1.1}$$

where  $C$  is a constant depends on  $R, T, \text{Lip}(u_0), a(x)$  and  $\max |b(y)|$ .

**Remark 1.** In Theorem 1.1 if  $V(x, y) = V(y)$  does not depend on  $x$ , then as it is proved in Section 2 (Proposition 2.2 is no longer needed in this case),  $u^\varepsilon \rightarrow u$  uniformly in  $\mathbb{R} \times (0, \infty)$  and the constant  $C$  in (1.1) can be chosen explicitly as

$$C = 2 \left( \|u'_0\|_{L^\infty(\mathbb{R})} + 2 (\|V\|_{L^\infty})^{1/2} \right).$$

**Theorem 1.2.** *Let  $n = 1$  and  $H(x, y, p) = H(p) + V(x, y) + C_0$  where  $C_0$  is a constant,  $H \in C^2(\mathbb{R}, [0, \infty))$  is strictly convex with  $\min_{p \in \mathbb{R}} H(p) = H(0) = 0$ . Define*

$$G_1 = (H'|_{[0, \infty)}) \circ (H|_{[0, \infty)})^{-1} \quad \text{and} \quad G_2 = (H'|_{(-\infty, 0]}) \circ (H|_{(-\infty, 0]})^{-1}.$$

Assume (H1)–(H4),  $V(x, y)$  is continuously differentiable in  $x$  variable for each  $y \in \mathbb{T}$ , and:

(A0)

$$\limsup_{p \rightarrow 0} \left| \frac{H''(p)}{H'(p)} \sqrt{H(p)} \right| < \infty. \tag{1.2}$$

(A1)  $\max_{\mathbb{R} \times \mathbb{T}} V(x, y) = 0$ , there exists  $y_0 \in \mathbb{T}$  such that  $V(x, y_0) = 0$  for all  $x \in \mathbb{R}$ .

For each compact interval  $I \subset \mathbb{R}$  and  $i = 1, 2$  we have:

(A2)

$$\limsup_{r \rightarrow 0^+} \left\{ |V_x(x, y)| \cdot \frac{|G'_i(r - V(x, y))|}{|G_i(r - V(x, y))|} : (x, y) \in I \times \mathbb{T} \right\} \leq A_I.$$

(A3)

$$\sup_{(x, y) \in I \times \mathbb{T}} \left| \frac{V_x(x, y)}{G_i(|V(x, y)|)} \right| \leq B_I.$$

(A4)

$$\limsup_{r \rightarrow 0^+} \left( \frac{\max_{x \in I} \int_0^1 \frac{dy}{|G_i(r - V(x, y))|}}{\min_{x \in I} \int_0^1 \frac{dy}{|G_i(r - V(x, y))|}} \right) \leq C_I.$$

If  $u_0 \in \text{Lip}(\mathbb{R}) \cap \text{BUC}(\mathbb{R})$  then for any  $R, T > 0$  we have

$$\|u^\varepsilon - u\|_{L^\infty([-R, R] \times [0, T])} \leq C\varepsilon \quad (1.3)$$

where  $C$  is a constant depends only on  $R, T, \text{Lip}(u_0), H(p)$  and  $V(x, y)$ .

Our paper is the first work improving the rate of convergence  $u^\varepsilon \rightarrow u$  for  $(C_\varepsilon)$  in one dimensional case as far as we know. Our method develops further that in [16] and uses deep dynamical properties of optimal paths in the optimal control formula. Higher dimensional cases will be investigated in future works.

### Remark 2.

- (i) Condition (A0) is satisfied for a vast class of strictly convex  $C^2$  Hamiltonians, including ones with  $H''(0) > 0$ ,  $H \in C^3$ , or  $|p|^\gamma$  with  $\gamma \geq 2$  (Lemma 2.7).
- (ii) If  $H''(0) > 0$  and  $\sup_{I \times \mathbb{T}} \left| \frac{V_x(x, y)}{V(x, y)} \right|$  is bounded for any compact interval  $I \subset \mathbb{R}$ , then (A2), (A3) hold (Corrolary 1.3).
- (iii) If  $V(x, y) = V(y)$  does not depend on  $x$ , then assumptions (A2)–(A4) automatically hold, while (A1) is satisfied after approximating  $H$  by uniformly convex Hamiltonians. Indeed, the method can be used to get the result for general convex Hamiltonians. We thus recover Theorem 1.3 in [16] and the convergence is uniform in this case. By Proposition 4.3 in [16], the rate  $\mathcal{O}(\varepsilon)$  is optimal.

The following is an important consequence of Theorem 1.2.

**Corollary 1.3.** *If  $H(x, y, p) = H(p) + V(x, y)$  where  $H(p) \geq H(0) = 0$  such that:*

- $H(p) \in C^2(\mathbb{R})$  is strictly convex with  $H''(0) > 0$ , or  $H(p) = |p|^\gamma$  where  $\gamma \geq 2$ .
- $\max_{\mathbb{R} \times \mathbb{T}} V(x, y) = 0$ , there exists  $y_0 \in \mathbb{T}$  such that  $V(x, y_0) = 0$  for all  $x \in \mathbb{R}$ .
- For every compact interval  $I \subset \mathbb{R}$  then  $\alpha_I f_I(y) \leq |V(x, y)| \leq \beta_I f_I(y)$  for  $\alpha_I, \beta_I > 0$  and  $f_I \in C(\mathbb{R}, [0, \infty))$ , and

$$\sup_{(x, y) \in I \times \mathbb{T}} \left| \frac{V_x(x, y)}{V(x, y)} \right| \leq C_I < \infty. \quad (1.4)$$

If  $u_0 \in \text{Lip}(\mathbb{R}) \cap \text{BUC}(\mathbb{R})$  then for any  $R, T > 0$  we have

$$\|u^\varepsilon - u\|_{L^\infty([-R, R] \times [0, T])} \leq C\varepsilon$$

where  $C$  is a constant depends only on  $R, T, \text{Lip}(u_0), H(p)$  and  $V(x, y)$ .

## 2. CLASSICAL MECHANICS HAMILTONIAN SETTING

*Proof of Theorem 1.1.* We observe that the estimate (1.1) does not depend on the smoothness of  $b(\cdot)$ , by approximation, without loss of generality, we can assume that  $V \in C^2(\mathbb{R} \times \mathbb{T})$ . Also, by replacing  $u$  by  $u + C$  we can normalize that  $C_0 = 0$ . Let us fix  $R, T > 0$ ,  $\varepsilon \in (0, 1)$  and  $(x_0, t_0) \in [-R, R] \times [0, T]$ , thanks to the optimal control formula (see [2, 10]) we have

$$u^\varepsilon(x_0, t_0) = \inf_{\eta \in \mathcal{T}} \left\{ \varepsilon \int_0^{\varepsilon^{-1}t_0} \left( \frac{|\dot{\eta}(s)|^2}{2} - V(\varepsilon\eta(s), \eta(s)) \right) ds + u_0(\varepsilon\eta(\varepsilon^{-1}t_0)) \right\}, \quad (2.1)$$

where  $\mathcal{T} = \{\eta(\cdot) \in \text{AC}([0, \varepsilon^{-1}t_0]), \varepsilon\eta(0) = x_0\}$ . Here  $\text{AC}([a, b])$  denotes the set of absolutely continuous functions from  $[a, b]$  to  $\mathbb{R}$ . Let  $\eta_\varepsilon(\cdot) \in \mathcal{T}$  be a minimizer to the optimization problem (2.1), it is clear that  $\eta_\varepsilon(\cdot)$  must satisfy the following Euler-Lagrange equation

$$\begin{cases} \ddot{\eta}_\varepsilon(s) &= -\nabla V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \cdot (\varepsilon, 1) & \text{on } (0, \varepsilon^{-1}t_0), \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0. \end{cases} \quad (2.2)$$

Here  $\nabla V$  means the full gradient of  $V$ . In particular, this implies the following conservation of energy:

$$\frac{d}{ds} \left( \frac{|\dot{\eta}_\varepsilon(s)|^2}{2} + V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \right) = \dot{\eta}_\varepsilon(s) \left( \ddot{\eta}_\varepsilon(s) + \nabla V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \cdot (\varepsilon, 1) \right) = 0$$

for all  $s \in (0, \varepsilon^{-1}t_0)$ . Hence there exists a constant  $r = r(\eta_\varepsilon) \in [V(0, 0), +\infty)$  such that

$$\frac{|\dot{\eta}_\varepsilon(s)|^2}{2} + V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) = r \quad \text{for all } s \in (0, \varepsilon^{-1}t_0). \quad (2.3)$$

For each  $r \in [V(0, 0), \infty)$  the Euler-Lagrange equation (2.2) can be written as:

$$\begin{cases} \ddot{\eta}_\varepsilon(s) &= -\nabla V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \cdot (\varepsilon, 1) & \text{on } (0, \varepsilon^{-1}t_0), \\ |\dot{\eta}_\varepsilon(0)| &= \sqrt{2(r - V(0, 0))}, \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0. \end{cases} \quad (2.4)$$

For simplicity, let us define the action functional

$$A^\varepsilon[\eta] = \varepsilon \int_0^{\varepsilon^{-1}t_0} \left( \frac{|\dot{\eta}(s)|^2}{2} - V(\varepsilon\eta(s), \eta(s)) \right) ds + u_0(\varepsilon\eta(\varepsilon^{-1}t_0))$$

for  $\eta(\cdot) \in \mathcal{T}$ . Thanks to the conservation of energy (2.3), the optimization problem (2.1) is equivalent to

$$u^\varepsilon(x_0, t_0) = \inf_r \left\{ A^\varepsilon[\eta_\varepsilon] : \text{among all } \eta_\varepsilon(\cdot) \text{ solve (2.2) with energy } r \right\}. \quad (2.5)$$

We proceed to get different estimates for  $r \leq 0$  and  $r > 0$ . For simplicity, let us introduce the following notation. For  $I$  be an interval of  $\mathbb{R}$ , we define

$$\inf_{r \in I} A^\varepsilon[\eta_\varepsilon]$$

which means the infimum over all solutions  $\eta_\varepsilon(\cdot)$  that solve (2.2) and with all energies  $r \in I$ .

**Proposition 2.1.** *When  $r \leq 0$ , we have the following estimate:*

$$\left| \inf_{r \leq 0} A^\varepsilon[\eta_\varepsilon] - u_0(x_0) \right| \leq \left( \sqrt{2\|V\|_{L^\infty}} + \|u'_0\|_{L^\infty} \right) \varepsilon. \quad (2.6)$$

*Proof.* Let  $\eta_\varepsilon(\cdot)$  be a solution to (2.4) with  $r \in [V(0, 0), 0]$  we claim that

$$\underline{y}_0 \leq \eta_\varepsilon(s) \leq \overline{y}_0 \quad \text{for all } s \in [0, \varepsilon^{-1}t_0], \quad (2.7)$$

where

$$\begin{aligned} \overline{y}_0 &= \min \{ y \in [\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1) : b(y) = 0 \}, \\ \underline{y}_0 &= \max \{ y \in (\varepsilon^{-1}x_0 - 1, \varepsilon^{-1}x_0] : b(y) = 0 \}. \end{aligned}$$

The existence of  $\underline{y}_0$  and  $\overline{y}_0$  is due to the periodicity of  $b(\cdot)$  and  $b(y_0) = 0$ . Recall that  $\eta_\varepsilon(\cdot)$  satisfies the following equation thanks to the conservation of energy (2.3):

$$\begin{cases} |\dot{\eta}_\varepsilon(s)| &= \sqrt{2(r - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)))}, & s \in (0, \varepsilon^{-1}t_0), \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0. \end{cases}$$

Let us define  $\gamma_+ : [0, \infty) \rightarrow \mathbb{R}$  and  $\gamma_- : [0, \infty) \rightarrow \mathbb{R}$  such that

$$\begin{cases} \dot{\gamma}_+(s) &= \sqrt{-2V(\varepsilon\gamma_+(s), \gamma_+(s))} & \text{on } (0, +\infty), \\ \gamma_+(0) &= \varepsilon^{-1}x_0, \end{cases} \quad (2.8)$$

and

$$\begin{cases} \dot{\gamma}_-(s) &= -\sqrt{-2V(\varepsilon\gamma_-(s), \gamma_-(s))} & \text{on } (0, +\infty), \\ \gamma_-(0) &= \varepsilon^{-1}x_0, \end{cases} \quad (2.9)$$

respectively. To be precise, there are two cases:

- $V(x_0, \varepsilon x_0) = 0$ , by Lemma 2.7 we have  $x \mapsto \sqrt{-V(\varepsilon x, x)}$  is Lipschitz on  $[\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1]$ . By uniqueness of solutions to (2.8) and (2.9) we have  $\gamma_-(s) \equiv \gamma_+(s) \equiv \varepsilon^{-1}x_0$  for all  $s \in [0, +\infty)$ .
- $V(x_0, \varepsilon x_0) \neq 0$ , the solution  $\gamma_+(\cdot)$  exists at least until  $\gamma_+(\cdot)$  goes passing  $\varepsilon^{-1}x_0 + 1$ . Indeed,  $\gamma_+(\cdot)$  remains stay inside  $[\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1]$  and hence solution exists on  $(0, +\infty)$ . To see this, we first observe that  $\gamma_+(\cdot)$  is increasing and for each time  $t > 0$ , from (2.8) we have

$$t = \int_{\gamma_+(0)}^{\gamma_+(t)} \frac{dx}{\sqrt{-V(\varepsilon x, x)}},$$

hence the amount of time  $\gamma_+(\cdot)$  needs to reach  $\overline{y}_0$  is

$$\int_{\gamma_+(0)}^{\overline{y}_0} \frac{dx}{\sqrt{-V(\varepsilon x, x)}} = +\infty$$

since  $x \mapsto \sqrt{-V(\varepsilon x, x)}$  is Lipschitz on  $[\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1]$  by Lemma 2.7. We conclude that  $\gamma_+(s) \rightarrow \overline{y_0}$  and similarly  $\gamma_-(s) \rightarrow \underline{y_0}$  as  $s \rightarrow \infty$ .

As a consequence, we have

$$\underline{y_0} \leq \gamma_-(s) \leq \eta_\varepsilon(s) \leq \gamma_+(s) \leq \overline{y_0} \quad \text{for all } s \in [0, \varepsilon^{-1}t_0] \quad (2.10)$$

and thus (2.7) follows. Now we utilize (2.7) to estimate  $A^\varepsilon[\eta_\varepsilon]$ . For any  $\eta_\varepsilon$  solves (2.4) we have

$$A^\varepsilon[\eta_\varepsilon] \geq u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \geq u_0(\varepsilon\eta_\varepsilon(0)) - \|u'_0\|_{L^\infty}\varepsilon. \quad (2.11)$$

On the other hand,

$$\begin{aligned} \inf_{r \leq 0} A^\varepsilon[\eta_\varepsilon] &\leq A^\varepsilon[\gamma_+] = \varepsilon \int_{\gamma_+(0)}^{\gamma_+(\varepsilon^{-1}t_0)} \sqrt{-2V(\varepsilon x, x)} dx + u_0(\varepsilon\gamma_+(\varepsilon^{-1}t_0)) \\ &\leq u_0(\varepsilon\eta_\varepsilon(0)) + \left( \sqrt{2\|V\|_{L^\infty}} + \|u'_0\|_{L^\infty} \right) \varepsilon. \end{aligned} \quad (2.12)$$

thanks to (2.21). From (2.11) and (2.12) we obtain our claim (2.6).  $\square$

For each  $r \in (0, \infty)$ , equation (2.4) has exactly two distinct solutions  $\eta_{1,r,\varepsilon}(\cdot)$  and  $\eta_{2,r,\varepsilon}(\cdot)$  thanks to the conservation of energy (2.3). They are

$$\begin{cases} \dot{\eta}_\varepsilon(s) &= \sqrt{2(r - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)))} & \text{on } (0, \varepsilon^{-1}t_0), \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0, \end{cases} \quad (2.13)$$

and

$$\begin{cases} \dot{\eta}_\varepsilon(s) &= -\sqrt{2(r - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)))} & \text{on } (0, \varepsilon^{-1}t_0), \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0, \end{cases} \quad (2.14)$$

respectively. Let us consider the first case  $\eta_\varepsilon(\cdot)$  solves (2.13) since the other case is similar. Since  $\dot{\eta}_\varepsilon(s) > 0$  we have

$$t_0 = \varepsilon \int_0^{\varepsilon^{-1}t_0} \frac{\dot{\eta}_\varepsilon(s)}{\dot{\eta}_\varepsilon(s)} ds = \varepsilon \int_{\eta_\varepsilon(0)}^{\eta_\varepsilon(\varepsilon^{-1}t_0)} \frac{dx}{\sqrt{2(r - V(\varepsilon x, x))}}. \quad (2.15)$$

This holds true for every  $\varepsilon > 0$ , thus we deduce that  $\eta_\varepsilon(\varepsilon^{-1}t_0) \rightarrow +\infty$  as  $\varepsilon \rightarrow 0^+$ . It is also clear that for all  $\varepsilon > 0$  then

$$t_0\sqrt{2r} \leq \varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - x_0 \leq t_0\sqrt{2(r + \|V\|_{L^\infty})}. \quad (2.16)$$

By the conservation of energy (2.3) we can write the action functional as

$$A^\varepsilon[\eta_\varepsilon] = rt_0 + 2\varepsilon \int_{\eta_\varepsilon(0)}^{\eta_\varepsilon(\varepsilon^{-1}t_0)} \frac{-V(\varepsilon x, x)}{\sqrt{2(r - V(\varepsilon x, x))}} dx + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)). \quad (2.17)$$

We observe that the infimum of the optimization problem (2.5) should be taken over  $r$  not too big.

**Proposition 2.2.** *There exists  $r_0 > 0$  depends only on  $\text{Lip}(u_0)$  and  $\|V\|_{L^\infty}$  such that*

$$\inf_{r \geq r_0} A^\varepsilon[\eta_\varepsilon] \geq u^\varepsilon(x_0, t_0) + t_0. \quad (2.18)$$

*Proof.* If  $\eta_\varepsilon$  is a solution to (2.8) with  $r > 0$ , then from (2.17) we have

$$\begin{aligned} A^\varepsilon[\eta_\varepsilon] &\geq rt_0 + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \\ &\geq rt_0 + u_0(x_0) - \|u'_0\|_{L^\infty} |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - x_0| \\ &\geq rt_0 + u_0(x_0) - \|u'_0\|_{L^\infty} t_0 \sqrt{2(r + \|V\|_{L^\infty})} \end{aligned} \quad (2.19)$$

thanks to (2.16). On the other hand, by assumption (H3) we can define

$$\overline{C} = \sup_{(x,y)} \left\{ |H(x,y,p)| : |p| \leq \|u'_0\|_{L^\infty} \right\} < \infty$$

then  $\bar{u}(x,t) = u_0(x) + \overline{C}t$  is a super-solution to  $(C_\varepsilon)$ . The well-known comparison principle for  $(C_\varepsilon)$  gives us

$$u^\varepsilon(x_0, t_0) \leq \bar{u}(x_0, t_0) = u_0(x_0) + \overline{C}t_0. \quad (2.20)$$

There exists  $r_0 > 0$  such that for  $r \geq r_0$  we have

$$r \geq \overline{C} + 1 + \|u'_0\|_{L^\infty} \sqrt{2(r + \|V\|_{L^\infty})},$$

which is equivalent to

$$rt_0 + u_0(x_0) - \|u'_0\|_{L^\infty} t_0 \sqrt{2(r + \|V\|_{L^\infty})} \geq u_0(x_0) + (\overline{C} + 1)t_0.$$

This estimate together with (2.19) and (2.20) gives us

$$A^\varepsilon[\eta_\varepsilon] \geq u^\varepsilon(x_0, t_0) + t_0 \quad \text{for all } r \geq r_0$$

which proves our claim (2.18), since the case  $\eta_\varepsilon$  solves (2.9) can be done similarly.  $\square$

With (2.18), the optimization problem (2.5) can be reduced to

$$u^\varepsilon(x_0, t_0) = \min \left\{ \inf_{r \leq 0} A^\varepsilon[\eta_\varepsilon], \inf_{0 < r < r_0} A^\varepsilon[\eta_\varepsilon] \right\}. \quad (2.21)$$

Thanks to (2.6), we only need to focus on the case  $0 < r < r_0$ . For simplicity, let us define the following interval  $I_0 \subset \mathbb{R}$  to be

$$I_0 = I_0(T, R) = [-R, c_0 + R] \quad \text{where} \quad c_0 = T \sqrt{2(r_0 + \|V\|_{L^\infty})}.$$

Since (2.16) is true for all  $0 < r < r_0$ , for all  $(x_0, t_0) \in [-R, R] \times [0, T]$  we have

$$\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) \in I_0.$$

Let us define  $c_{1,r} > 0$  and  $c_{2,r} < 0$  be unique numbers such that

$$\int_{x_0}^{c_{1,r}} \int_0^1 \frac{dydx}{\sqrt{2(r - V(x,y))}} = t_0, \quad (2.22)$$

and

$$\int_{c_{2,r}}^{x_0} \int_0^1 \frac{dydx}{\sqrt{2(r - V(x,y))}} = t_0,$$

respectively.

**Proposition 2.3.** *Let  $\alpha_T = \min_{x \in I_0} a(x)$  and  $\beta_T = \max_{x \in I_0} a(x)$ , then*

$$|\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - c_{1,r}| \leq C_K \varepsilon \quad (2.23)$$

for  $0 < r < r_0$  where  $C_K$  is a constant only depends on  $R, T$  and  $V$ .



*Proof.* Let us define

$$\mathcal{K}_r(x, y) = \frac{1}{\sqrt{2(r - V(x, y))}}, \quad (x, y) \in \mathbb{R} \times \mathbb{T}.$$

From (2.15) and (2.22) we have

$$t_0 = \int_{x_0}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \mathcal{K}_r\left(x, \frac{x}{\varepsilon}\right) dx = \int_{x_0}^{c_{1,r}} \int_0^1 \mathcal{K}_r(x, y) dy dx. \quad (2.24)$$

Using Lemma 2.6 we obtain

$$\left| \int_{x_0}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \mathcal{K}_r\left(x, \frac{x}{\varepsilon}\right) dx - \int_{x_0}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \int_0^1 \mathcal{K}_r(x, y) dy dx \right| \leq K\varepsilon \quad (2.25)$$

where

$$K = 2 \max_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy + c_0 \max_{x \in I_0} \int_0^1 \frac{\partial \mathcal{K}_r}{\partial x}(x, y) dy. \quad (2.26)$$

Using (2.24) in (2.25) we have

$$\left| \int_{c_{1,r}}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \int_0^1 \mathcal{K}_r(x, y) dy dx \right| \leq K\varepsilon$$

which implies that

$$\left( \min_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \right) |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - c_{1,r}| \leq K\varepsilon. \quad (2.27)$$

On  $I_0$  we have  $0 < \alpha_T \leq a(x) \leq \beta_T$ , which implies that

$$\begin{aligned} \int_0^1 \frac{dy}{\sqrt{2(r - \beta_T b(y))}} &\leq \min_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \\ &\leq \max_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \leq \int_0^1 \frac{dy}{\sqrt{2(r - \alpha_T b(y))}}. \end{aligned} \quad (2.28)$$

Since  $\alpha_T \leq \beta_T$ , it is clear that

$$\begin{aligned} \int_0^1 \frac{dy}{\sqrt{2(r - \alpha_T b(y))}} &= \frac{1}{\sqrt{\alpha_T}} \int_0^1 \frac{dy}{\sqrt{2((r/\alpha_T) - b(y))}} \\ &\leq \frac{1}{\sqrt{\alpha_T}} \int_0^1 \frac{dy}{\sqrt{2((r/\beta_T) - b(y))}} = \sqrt{\frac{\beta_T}{\alpha_T}} \int_0^1 \frac{dy}{\sqrt{2(r - \beta_T b(y))}}. \end{aligned} \quad (2.29)$$

From direct calculation

$$\frac{\partial \mathcal{K}_r}{\partial x}(x, y) = \frac{a'(x)b(y)}{2(r - a(x)b(y))} \frac{1}{\sqrt{2(r - a(x)b(y))}}$$

we have

$$\max_{x \in I_0} \int_0^1 \left| \frac{\partial \mathcal{K}_r}{\partial x}(x, y) \right| dy \leq \frac{1}{2} \max_{x \in I_0} \left| \frac{a'(x)}{a(x)} \right| \int_0^1 \frac{dy}{\sqrt{2(r - \alpha_T b(y))}}. \quad (2.30)$$

Use (2.28) and (2.30) in (2.26) to deduce that

$$K \leq \left( 2 + \frac{c_0}{2} \max_{x \in I_0} \left| \frac{a'(x)}{a(x)} \right| \right) \left( \int_0^1 \frac{dy}{\sqrt{2(r - \alpha_T b(y))}} \right). \quad (2.31)$$

Next, we use (2.28), (2.31) in (2.27) to deduce that

$$\int_0^1 \frac{dy}{\sqrt{2(r - \beta_T b(y))}} |\varepsilon \eta_\varepsilon(\varepsilon^{-1}t) - c_{1,r}| \leq \left( 2 + \frac{c_0}{2} \max_{x \in I_0} \frac{|a'(x)|}{a(x)} \right) \left( \int_0^1 \frac{dy}{\sqrt{2(r - \alpha_T b(y))}} \right) \varepsilon. \quad (2.32)$$

From (2.29) and (2.32) we obtain our claim (2.23) with

$$C_K = \sqrt{\frac{\beta_T}{\alpha_T}} \left( 2 + \frac{c_0}{2} \max_{x \in I_0} \left| \frac{a'(x)}{a(x)} \right| \right). \quad (2.33)$$

It is clear that  $C_K$  depends only on  $R, T$  and  $a(x)$ .  $\square$

In view of (2.17), for  $0 < r < r_0$  we aim to show that

$$\left| \varepsilon \int_{x_0}^{\eta_\varepsilon(\varepsilon^{-1}t_0)} \frac{-V(\varepsilon x, x)}{\sqrt{2(r - V(\varepsilon x, x))}} dx - \int_{x_0}^{c_{1,r}} \int_0^1 \frac{-V(x, y)}{\sqrt{2(r - V(x, y))}} dy dx \right| \leq C_F \varepsilon \quad (2.34)$$

where  $C_F$  is some constant only depends on  $R, T$  and  $V$ . To see it, let

$$\mathcal{F}_r(x, y) = \frac{-V(x, y)}{\sqrt{2(r - V(x, y))}}, \quad (x, y) \in \mathbb{R} \times \mathbb{T}.$$

Using Lemma 2.6 we obtain

$$\left| \int_{x_0}^{c_{1,r}} \frac{-V(x, \varepsilon^{-1}x)}{\sqrt{2(r - V(x, \varepsilon^{-1}x))}} dx - \int_{x_0}^{c_{1,r}} \int_0^1 \frac{-V(x, y)}{\sqrt{2(r - V(x, y))}} dy dx \right| \leq (2F_1 + c_0 F_2) \varepsilon \quad (2.35)$$

where

$$F_1 := (\|V\|_{L^\infty})^{1/2} \geq \max_{\mathbb{R} \times \mathbb{T}} |\mathcal{F}_r(x, y)|$$

and

$$F_2 := \frac{3}{\sqrt{2}} (\|V\|_{L^\infty})^{1/2} \max_{x \in I_0} \left| \frac{a'(x)}{a(x)} \right| \geq \max_{I_0 \times \mathbb{T}} \left| \frac{\partial \mathcal{F}_r}{\partial x}(x, y) \right|.$$

On the other hand, we have

$$\left| \int_{x_0}^{\varepsilon \eta_\varepsilon(\varepsilon^{-1}t_0)} \frac{-V(x, \varepsilon^{-1}x)}{\sqrt{2(r - V(x, \varepsilon^{-1}x))}} dx - \int_{x_0}^{c_{1,r}} \frac{-V(x, \varepsilon^{-1}x)}{\sqrt{2(r - V(x, \varepsilon^{-1}x))}} dx \right| \leq F_1 |\varepsilon \eta_\varepsilon(\varepsilon^{-1}t_0) - c_{1,r}| \leq F_1 C_K \varepsilon \quad (2.36)$$

thanks to (2.23). From (2.35) and (2.36) we deduce that

$$\left| \int_{x_0}^{\varepsilon \eta_\varepsilon(\varepsilon^{-1}t_0)} \frac{-V(x, \varepsilon^{-1}x)}{\sqrt{2(r - V(x, \varepsilon^{-1}x))}} dx - \int_{x_0}^{c_{1,r}} \int_0^1 \frac{-V(x, y)}{\sqrt{2(r - V(x, y))}} dy dx \right| \leq (2F_1 + F_2 c_0 + F_1 C_K) \varepsilon. \quad (2.37)$$

From (2.37) we obtain our claim (2.34) with

$$C_F = (\|V\|_{L^\infty})^{1/2} \left( 2 + 2\sqrt{\frac{\beta_T}{\alpha_T}} + c_0 \left( \frac{3}{\sqrt{2}} + \frac{1}{2}\sqrt{\frac{\beta_T}{\alpha_T}} \right) \max_{x \in I_0} \left| \frac{a'(x)}{a(x)} \right| \right). \quad (2.38)$$

**Proposition 2.4.** *We have the following estimate:*

$$\left| \inf_{\substack{0 < r < r_0 \\ i=1,2}} A^\varepsilon[\eta_{i,r,\varepsilon}] - \inf_{0 < r < r_0} I(r) \right| \leq C\varepsilon \quad (2.39)$$

where  $C$  is a constant depends only on  $R, T, a(x)$  and  $\|V\|_{L^\infty}$ ,  $I(r) = \min\{I_1(r), I_2(r)\}$  where

$$I_1(r) = rt_0 + 2 \int_{x_0}^{c_{1,r}} \int_0^1 \frac{-V(x, y)}{\sqrt{2(r - V(x, y))}} dy dx + u_0(c_{1,r}), \quad (2.40)$$

$$I_2(r) = rt_0 + 2 \int_{c_{2,r}}^{x_0} \int_0^1 \frac{-V(x, y)}{\sqrt{2(r - V(x, y))}} dy dx + u_0(c_{2,r}). \quad (2.41)$$

*Proof.* Within our notation  $\eta_\varepsilon \equiv \eta_{1,r,\varepsilon}$ , we have

$$|u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) - u_0(c_{1,r})| \leq \|u'_0\|_{L^\infty} |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - c_{1,r}|. \quad (2.42)$$

since  $u_0 \in \text{Lip}(\mathbb{R})$ . In view of (2.17) and (2.23), (2.34), (2.42) we conclude that

$$\begin{aligned} |A^\varepsilon[\eta_\varepsilon] - I_1(r)| &\leq 2C_F\varepsilon + \|u'_0\|_{L^\infty} |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t) - c_{1,r}| \\ &\leq (2C_F + C_K\|u'_0\|_{L^\infty})\varepsilon. \end{aligned} \quad (2.43)$$

Taking the infimum over  $0 < r < r_0$  we obtain

$$\left| \inf_{0 < r < r_0} A^\varepsilon[\eta_{1,r,\varepsilon}] - \inf_{0 < r < r_0} I_1(r) \right| \leq C_1\varepsilon \quad (2.44)$$

where

$$C_1 = 2C_F + C_K\|u'_0\|_{L^\infty}. \quad (2.45)$$

Doing similarly for the case  $\eta_{2,r,\varepsilon}$  solves (2.14), we obtain

$$\left| \inf_{0 < r < r_0} A^\varepsilon[\eta_{2,r,\varepsilon}] - \inf_{0 < r < r_0} I_2(r) \right| \leq C_2\varepsilon \quad (2.46)$$

where  $C_2$  is some constant depends on  $R, T, a(x)$  and  $\|V\|_{L^\infty}$  in the same manner as (2.45). Thus our claim (2.39) is correct with  $C = \max\{C_1, C_2\}$ .  $\square$

From (2.6), (2.21) and (2.39) we conclude that

$$|u^\varepsilon(x_0, t_0) - u(x_0, t_0)| \leq \left( \max \left\{ \sqrt{2\|V\|_{L^\infty}} + \|u'_0\|_{L^\infty}, C \right\} \right) \varepsilon$$

and the proof is complete.  $\square$

**Corollary 2.5.** *We have the following representation formula*

$$u(x_0, t_0) = \min \left\{ u_0(x_0), \min \left\{ \inf_{0 < r < r_0} I_1(r), \inf_{0 < r < r_0} I_2(r) \right\} \right\}$$

where  $I_1(r)$  and  $I_2(r)$  are defined in (2.40) and (2.41) respectively.

**Remark 3.** If  $V(x, y) = V(y)$  is independent of  $x$ , then the constants  $C_K$  in (2.38) and  $C_F$  in (2.38) are independent of  $R$  and  $T$ . Therefore the convergence is uniform in  $\mathbb{R} \times [0, \infty)$  and by carefully keeping track of all constants, we get

$$C = 2 \left( \|u'_0\|_{L^\infty(\mathbb{R})} + 2\sqrt{\max_{y \in \mathbb{T}} |V(y)|} \right).$$

For the sake of completeness, we provide here a proof for Lemma 2.6 (see [17]), which is a quantitative version of the ergodic Theorem for periodic functions in one dimension. This is a generalized version of Lemma 4.2 in [16].

**Lemma 2.6.** *If  $F(x, y) \in C^1(\mathbb{R} \times \mathbb{T})$  then for any real numbers  $a < b$  we have*

$$\left| \int_a^b F\left(x, \frac{x}{\varepsilon}\right) dx - \int_a^b \left( \int_0^1 F(x, y) dy \right) dx \right| \leq C\varepsilon$$

where

$$C = 2 \max_{x \in [a, b]} \int_0^1 |F(x, y)| dy + (b - a) \max_{x \in [a, b]} \int_0^1 \left| \frac{\partial F}{\partial x}(x, y) \right| dy.$$

*Proof.* Since  $y \mapsto F(x, y)$  is periodic, we have  $y \mapsto \frac{\partial F}{\partial x} F(x, y)$  is also periodic. Let us define

$$G(x, y) = \int_0^y \left( F(x, z) - \int_0^1 F(x, \zeta) d\zeta \right) dz$$

then

$$\frac{\partial G}{\partial y}(x, y) = F(x, y) - \int_0^1 F(x, \zeta) d\zeta.$$

Since  $G$  is periodic in  $y$ ,  $\frac{\partial G}{\partial x}$  is also periodic in  $y$  because

$$\begin{aligned} \frac{\partial G}{\partial x}(x, y+1) &= \int_0^{y+1} \left( \frac{\partial F}{\partial x}(x, z) - \int_0^1 \frac{\partial F}{\partial x}(x, \zeta) d\zeta \right) dz \\ &= \int_0^y \left( \frac{\partial F}{\partial x}(x, z+1) - \int_0^1 \frac{\partial F}{\partial x}(x, \zeta) d\zeta \right) dz = \frac{\partial G}{\partial x}(x, y). \end{aligned}$$

Thus  $G$  and  $\frac{\partial G}{\partial x}$  are bounded in  $y$ . The fact that  $\frac{\partial F}{\partial x}$  is bounded in  $x$  implies  $\frac{\partial G}{\partial x}$  is bounded in  $x$  as well. Let  $g_\varepsilon(x) = \varepsilon G\left(x, \frac{x}{\varepsilon}\right)$  we obtain

$$\frac{d}{dx}\left(g_\varepsilon(x)\right) = \varepsilon \frac{\partial G}{\partial x}\left(x, \frac{x}{\varepsilon}\right) + \frac{\partial G}{\partial y}\left(x, \frac{x}{\varepsilon}\right) = \varepsilon \frac{\partial G}{\partial x}\left(x, \frac{x}{\varepsilon}\right) + F\left(x, \frac{x}{\varepsilon}\right) - \int_0^1 F(x, \zeta) d\zeta.$$

Thus

$$\begin{aligned} \int_a^b F\left(x, \frac{x}{\varepsilon}\right) dx - \int_a^b \int_0^1 F(x, \zeta) d\zeta dx &= \int_a^b \left( \frac{d}{dx}\left(g_\varepsilon(x)\right) - \varepsilon \frac{\partial G}{\partial x}\left(x, \frac{x}{\varepsilon}\right) \right) dx \\ &= \varepsilon \left[ G\left(b, \frac{b}{\varepsilon}\right) - G\left(a, \frac{a}{\varepsilon}\right) - \int_a^b \frac{\partial G}{\partial x}\left(x, \frac{x}{\varepsilon}\right) dx \right]. \end{aligned}$$

Note that by the way we defined  $G$ , we also have

$$\max_{(x, y)} |G(x, y)| \leq \max_{x \in [a, b]} \int_0^1 |F(x, y)| dy$$

and

$$\frac{\partial G}{\partial x}(x, y) = \int_0^y \frac{\partial F}{\partial x}(x, \zeta) d\zeta - y \int_0^1 \frac{\partial F}{\partial x}(x, \zeta) d\zeta$$

which implies

$$\left| \int_a^b \frac{\partial G}{\partial x} \left( x, \frac{x}{\varepsilon} \right) dx \right| \leq \int_a^b \max_{(x,y)} \left| \frac{\partial G}{\partial x}(x, y) \right| dx \leq (b-a) \max_{x \in [a,b]} \int_0^1 \left| \frac{\partial F}{\partial x}(x, y) \right| dy$$

and hence the proof is complete.  $\square$

**Lemma 2.7.** *Let  $\mathcal{V} \in C^2([0, 1], [0, \infty))$  with  $\min_{x \in [0,1]} \mathcal{V}(x) = 0$  and  $\mathcal{V}(0) = \mathcal{V}(1)$ . There exists a constant  $L > 0$  such that*

$$|\mathcal{V}'(x)| \leq L\sqrt{\mathcal{V}(x)} \quad \text{for all } x \in [0, 1]. \quad (2.47)$$

As a consequence,  $x \mapsto \sqrt{\mathcal{V}(x)}$  is Lipschitz in  $[0, 1]$ .

*Proof.* For each  $a \in [0, 1]$ , an  $\delta$ -neighborhood  $\mathcal{N}_{a,\delta}$  of  $a$  is defined as  $(a - \delta, a + \delta)$  if  $a \in (0, 1)$  and  $[0, \delta) \cup (1 - \delta, 1]$  if  $a \in \{0, 1\}$ . It is clear that  $\mathcal{N}_{a,\delta}$  is open in  $[0, 1]$ . We claim that there exists  $\delta = \delta(a) > 0$  such that

$$\sup_{x \in \mathcal{N}_{a,\delta}^*} \frac{|\mathcal{V}'(x)|}{\sqrt{\mathcal{V}(x)}} \leq C_a < \infty \quad (2.48)$$

for some constant  $C_a$ , where  $\mathcal{N}_{a,\delta}^* = \{x \in \mathcal{N}_{a,\delta} : \mathcal{V}(x) \neq 0\}$ . Assume that (2.48) is false, then without loss of generality there exists a sequence  $x_k \rightarrow a^+$  such that  $\mathcal{V}(x_k) \neq 0$  for all  $k \in \mathbb{N}$  and

$$\lim_{k \rightarrow \infty} \frac{|\mathcal{V}'(x_k)|}{\sqrt{\mathcal{V}(x_k)}} = +\infty. \quad (2.49)$$

It is clear that  $\mathcal{V}'(x_k) \neq 0$  for all  $k \in \mathbb{N}$ . Without loss of generality again we can assume that  $\mathcal{V}'(x_k) > 0$  for all  $k$ . Let  $g_k = \sqrt{\mathcal{V}(x_k)}$  and  $h_k = \mathcal{V}'(x_k)$ , and

$$a_k = \sup \left\{ r > 0 : \mathcal{V}'(x) \geq \frac{h_k}{2} \text{ for all } x \in (x_k - r, x_k) \right\}.$$

Clearly  $\mathcal{V}'(x_k - a_k) = \frac{h_k}{2}$ . By the mean value theorem we have

$$g_k^2 = \mathcal{V}(x_k) \geq \mathcal{V}(x_k) - \mathcal{V}(x_k - a_k) \geq \frac{h_k a_k}{2}.$$

By the mean value theorem again, there exists  $\xi_k \in (x_k - a_k, x_k)$  such that

$$\mathcal{V}''(\xi_k) = \frac{\mathcal{V}'(x_k) - \mathcal{V}'(x_k - a_k)}{a_k} = \frac{1}{2} \frac{h_k}{a_k} \geq \frac{1}{4} \left( \frac{h_k}{g_k} \right)^2 \rightarrow \infty$$

as  $k \rightarrow \infty$  due to (2.49). It is a contradiction since  $\mathcal{V} \in C^2([0, 1])$ , thus (2.48) must be correct. By compactness of  $[0, 1]$ , we can pick a finite subcover of  $[0, 1]$  from the open cover  $\{\mathcal{N}_{a,\delta} : a \in [0, 1]\}$ . From (2.48) there exists a constant  $L > 0$  such that

$$\frac{|\mathcal{V}'(x)|}{\sqrt{\mathcal{V}(x)}} \leq L \quad \text{whenever } \mathcal{V}(x) \neq 0. \quad (2.50)$$

From that we obtain our claim (2.47). For  $0 < \varepsilon < 1$  let  $f_\varepsilon(x) = \sqrt{\mathcal{V}(x) + \varepsilon} \in C^2([0, 1])$ . It  $\mathcal{V}(x) = 0$  then  $\mathcal{V}'(x) = 0$ , hence  $f'_\varepsilon(x) = 0$  as well, while if  $\mathcal{V}(x) \neq 0$  then from (2.50) we have

$$|f'_\varepsilon(x)| = \left| \frac{\mathcal{V}'(x)}{2\sqrt{\mathcal{V}(x) + \varepsilon}} \right| \leq \frac{1}{2} \frac{|\mathcal{V}'(x)|}{\sqrt{\mathcal{V}(x)}} \leq \frac{L}{2}$$

Thus  $f'_\varepsilon(x)$  is Lipschitz on  $[0, 1]$  with the Lipschitz constant independent of  $\varepsilon$ . Let  $\varepsilon \rightarrow 0$  we deduce that  $x \mapsto \sqrt{\mathcal{V}(x)}$  is Lipschitz on  $[0, 1]$ .  $\square$

### 3. GENERAL STRICTLY CONVEX HAMILTONIANS SETTING

**3.1. Setting and simplifications.** First of all, by replacing  $u$  by  $u + C_0$ , it suffices to prove the theorem for  $C_0 = 0$ . By approximation we can assume without loss of generality that  $V \in C^2(\mathbb{R} \times \mathbb{T})$ . It is well-known that (see appendix [10]) for  $(C_\varepsilon)$  we have the following estimate:

$$\sup_{(x,t) \in \mathbb{R} \times [0, \infty)} \left( |u_t^\varepsilon(x, t)| + |u_x^\varepsilon(x, t)| \right) \leq M \quad (3.1)$$

in the viscosity sense for all  $\varepsilon > 0$ . Accordingly, values of  $H(p)$  for  $|p| > M$  are irrelevant, hence without loss of generality we can assume that  $H$  grows quadratically, i.e.,

$$\frac{|p|^2}{2} - K_0 \leq H(p) \leq \frac{|p|^2}{2} + K_0 \quad (3.2)$$

for all  $p \in \mathbb{R}$  and for some  $K_0 > 0$ . The Legendre transform of  $H$  is defined by

$$L(v) = \sup_{p \in \mathbb{R}} \left( p \cdot v - H(p) \right) \quad \text{for } v \in \mathbb{R}.$$

It is clear that  $L$  is also smooth and strictly convex, with the same properties as  $H$ :

$$\frac{|v|^2}{2} - K_0 \leq L(v) \leq \frac{|v|^2}{2} + K_0 \quad \text{and} \quad L(v) > L(0) = 0 \quad \text{for } v \neq 0. \quad (3.3)$$

Using  $H(0) = H'(0) = 0$  and  $L(0) = L'(0) = 0$ , we can modify  $K_0$  such that beside (3.2) and (3.3) we also have

$$\frac{|v|^2}{2} - K_0|v| \leq L(v) \leq \frac{|v|^2}{2} + K_0|v| \quad \text{for all } v \in \mathbb{R} \quad (3.4)$$

and

$$\frac{|p|^2}{2} - K_0|p| \leq H(p) \leq \frac{|p|^2}{2} + K_0|p| \quad \text{for all } p \in \mathbb{R} \quad (3.5)$$

For each  $p \in \mathbb{R}$ , we have

$$v = H'(p) \iff p = L'(v) \iff p \cdot v = L(v) + H(p). \quad (3.6)$$

From that we obtain  $p \cdot v \geq \max\{H(p), L(v)\}$ , which implies that

$$\begin{cases} p = L'(v) \geq \frac{L(v)}{v} \geq \frac{v}{2} - K_0 & \text{if } v \geq 0, \\ p = L'(v) \leq \frac{L(v)}{v} \leq \frac{v}{2} - K_0 & \text{if } v \leq 0. \end{cases} \quad (3.7)$$

and similarly

$$\begin{cases} v = H'(p) \geq \frac{H(p)}{p} \geq \frac{p}{2} - K_0 & \text{if } p \geq 0, \\ v = H'(p) \leq \frac{H(p)}{p} \leq \frac{p}{2} - K_0 & \text{if } p \leq 0. \end{cases} \quad (3.8)$$

We also denote:

$$\begin{cases} H_1^{-1} := (H|_{[0,+\infty)})^{-1} : [0, +\infty) \longrightarrow [0, +\infty), \\ (L'_1)^{-1} := (L'|_{[0,+\infty)})^{-1} : [0, +\infty) \longrightarrow [0, +\infty), \\ \tilde{G}_1 := (L'_1)^{-1} \circ H_1^{-1} : [0, +\infty) \longrightarrow [0, +\infty). \end{cases}$$

and

$$\begin{cases} H_2^{-1} := (H|_{(-\infty,0]})^{-1} : (-\infty, 0] \longrightarrow (-\infty, 0], \\ (L'_2)^{-1} := (L'|_{(-\infty,0]})^{-1} : (-\infty, 0] \longrightarrow (-\infty, 0], \\ \tilde{G}_2 := (L'_2)^{-1} \circ H_2^{-1} : (-\infty, 0] \longrightarrow (-\infty, 0]. \end{cases}$$

From (3.6) we have that  $H'_i = (L'_i)^{-1}$  and thus  $\tilde{G}_i \equiv G_i$  for  $i = 1, 2$  where  $G_i$  are defined in the statement of Theorem 1.2. It is clear that for  $i = 1, 2$  we have  $x \mapsto |G_i(x)|$  is increasing on  $[0, \infty)$ . Furthermore since  $L$  is strictly convex we deduce that  $q \mapsto L'(q)$  is strictly increasing. Thus  $x \mapsto (L'_i)^{-1}(x)$  is increasing as well and from (3.6), (3.8) and (3.7) we obtain

$$\begin{cases} \frac{p}{2} - K_0 \leq (L'_1)^{-1}(p) \leq 2(K_0 + p) & \text{if } p \geq 0, \\ \frac{p}{2} - K_0 \geq (L'_2)^{-1}(p) \geq 2(K_0 + p) & \text{if } p \leq 0. \end{cases}$$

Here we used the fact that  $L_i^{-1}(p) = v = H'_i(p)$  for  $i = 1, 2$ . From (3.2) we obtain

$$\begin{cases} \sqrt{2(x - K_0)} \leq H_1^{-1}(x) \leq \sqrt{2(x + K_0)}, \\ -\sqrt{2(x + K_0)} \leq H_2^{-1}(x) \leq -\sqrt{2(x - K_0)} \end{cases} \quad \text{for } x \geq K_0.$$

From these observations we deduce that for all  $x \geq K_0$  then

$$\begin{cases} \sqrt{\frac{x - K_0}{2}} \leq G_1(x) \leq 2K_0 + 2\sqrt{2(x + K_0)}, \\ -\sqrt{\frac{x - K_0}{2}} \geq G_2(x) \geq 2K_0 - 2\sqrt{2(x + K_0)}. \end{cases} \quad (3.9)$$

As a consequence, we have  $|G_i(x)| \rightarrow +\infty$  as  $x \rightarrow \infty$  for  $i = 1, 2$ .

**3.2. Proof of Theorem 1.2.** Let us fix  $\varepsilon > 0$  and  $R, T > 0$  and consider a point  $(x_0, t_0) \in [-R, R] \times [0, T]$ . Thanks to the optimal control formula we have

$$u^\varepsilon(x_0, t_0) = \inf_{\eta(\cdot) \in \mathcal{T}} \left\{ \varepsilon \int_0^{\varepsilon^{-1}t_0} \left( L(\dot{\eta}(s)) - V(\varepsilon\eta(s), \eta(s)) \right) ds + u_0(\varepsilon\eta(\varepsilon^{-1}t_0)) \right\}, \quad (3.10)$$

where  $\mathcal{T} = \{\eta(\cdot) \in AC([0, \varepsilon^{-1}t_0]), \varepsilon\eta(0) = x_0\}$ . Let  $\eta_\varepsilon(\cdot) \in \mathcal{T}$  be a minimizer to the optimization problem (3.10), it is clear that  $\eta_\varepsilon(\cdot)$  satisfies the following Euler-Lagrange equation

$$\begin{cases} L''(\dot{\eta}_\varepsilon(s))\ddot{\eta}_\varepsilon(s) = -\nabla V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \cdot (\varepsilon, 1) & \text{on } (0, \varepsilon^{-1}t_0), \\ \eta_\varepsilon(0) = \varepsilon^{-1}x_0. \end{cases} \quad (3.11)$$

From (3.6) we deduce that

$$H'(L'(\dot{\eta}_\varepsilon(s))) = \dot{\eta}_\varepsilon(s) \quad \text{for all } s \in (0, \varepsilon^{-1}t_0). \quad (3.12)$$

Using the Euler-Lagrange equation (3.11) we have the conservation of energy:

$$\begin{aligned} \frac{d}{ds} \left( H(L'(\dot{\eta}_\varepsilon(s))) + V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \right) &= \dot{\eta}_\varepsilon(s) \left( \frac{d}{ds} L'(\dot{\eta}_\varepsilon(s)) + \nabla V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \cdot (\varepsilon, 1) \right) \\ &= 0 \end{aligned}$$

for all  $s \in (0, \varepsilon^{-1}t_0)$ . Hence there exists  $r = r(\eta_\varepsilon) \in [V(0, 0), +\infty)$  such that

$$H(L'(\dot{\eta}_\varepsilon(s))) + V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) = r \quad \text{for all } s \in (0, \varepsilon^{-1}t_0). \quad (3.13)$$

For each  $r \in [V(0, 0), \infty)$  the Euler-Lagrange equation (3.11) can be written as:

$$\begin{cases} L''(\dot{\eta}_\varepsilon(s))\ddot{\eta}_\varepsilon(s) = -\nabla V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \cdot (\varepsilon, 1) & \text{on } (0, \varepsilon^{-1}t_0), \\ \dot{\eta}_\varepsilon(0) = G_i(r - V(0, 0)), \\ \eta_\varepsilon(0) = \varepsilon^{-1}x_0. \end{cases} \quad (3.14)$$

where  $i = 1, 2$ . For simplicity, let us define the following action functional

$$A^\varepsilon[\eta] = \varepsilon \int_0^{\varepsilon^{-1}t_0} \left( L(\dot{\eta}(s)) - V(\varepsilon\eta(s), \eta(s)) \right) ds + u_0(\varepsilon\eta(\varepsilon^{-1}t_0)) \quad (3.15)$$

for  $\eta(\cdot) \in \mathcal{T}$ . Thanks to the conservation of energy (3.13), the optimization problem (3.10) is equivalent to

$$u^\varepsilon(x_0, t_0) = \inf_r \left\{ A^\varepsilon[\eta_\varepsilon] : \text{among all } \eta_\varepsilon(\cdot) \text{ solve (3.11) with energy } r \right\}. \quad (3.16)$$

We proceed to get different estimates for  $r \leq 0$  and  $r > 0$ . Similarly to the proof of Theorem 1.1, for an interval  $I \subset \mathbb{R}$  we denote

$$\inf_{r \in I} A^\varepsilon[\eta_\varepsilon] \quad (3.17)$$

which means the infimum over all solutions  $\eta_\varepsilon(\cdot)$  that solve (3.11) and with all energies  $r \in I$ .

**Proposition 3.1.** *When the energy  $r$  is negative, we have the following estimate:*

$$\left| \inf_{r \leq 0} A^\varepsilon[\eta_\varepsilon] - u_0(x_0) \right| \leq \left( H_1^{-1}(\|V\|_{L^\infty}) + \|u'_0\|_{L^\infty} \right) \varepsilon. \quad (3.18)$$

*Proof.* Let  $\eta_\varepsilon(\cdot)$  be a solution to (3.14) with  $r \in [V(0, 0), 0]$  we claim that

$$\underline{y_0} \leq \eta_\varepsilon(s) \leq \overline{y_0} \quad \text{for all } s \in [0, \varepsilon^{-1}t_0] \quad (3.19)$$



where

$$\begin{aligned}\overline{y_0} &= \min \{y \in [\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1] : V(\varepsilon y, y) = 0\}, \\ \underline{y_0} &= \max \{y \in (\varepsilon^{-1}x_0 - 1, \varepsilon^{-1}x_0] : V(\varepsilon y, y) = 0\}.\end{aligned}$$

The existence of  $\overline{y_0}$  and  $\underline{y_0}$  is due to assumption (A1). Recall that  $\eta_\varepsilon(\cdot)$  satisfies the following equation thanks to the conservation of energy (3.13)

$$\begin{cases} \dot{\eta}_\varepsilon(s) &= G_i(r - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s))), & s > 0, \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0 \end{cases}$$

where  $i = 1, 2$ . Let us define  $\gamma_+ : [0, \infty) \rightarrow \mathbb{R}$  and  $\gamma_- : [0, \infty) \rightarrow \mathbb{R}$  such that

$$\begin{cases} \dot{\gamma}_+(s) &= G_1(-V(\varepsilon\gamma_+(s), \gamma_+(s))) & \text{on } (0, +\infty), \\ \gamma_+(0) &= \varepsilon^{-1}x_0, \end{cases} \quad (3.20)$$

and

$$\begin{cases} \dot{\gamma}_-(s) &= G_2(-V(\varepsilon\gamma_-(s), \gamma_-(s))) & \text{on } (0, +\infty), \\ \gamma_-(0) &= \varepsilon^{-1}x_0, \end{cases} \quad (3.21)$$

respectively. To be precise, there are two cases:

- $V(x_0, \varepsilon x_0) = 0$ , by Lemma 3.5 we have  $x \mapsto G_i(-V(\varepsilon x, x))$  is Lipschitz on  $[\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1]$  for  $i = 1, 2$ . By uniqueness of solutions to (3.20) and (3.21) we have  $\gamma_-(s) \equiv \gamma_+(s) \equiv \varepsilon^{-1}x_0$  for all  $s \in [0, +\infty)$ .
- $V(x_0, \varepsilon x_0) \neq 0$ , the solution  $\gamma_+(\cdot)$  exists at least until  $\gamma_+(\cdot)$  goes passing  $\varepsilon^{-1}x_0 + 1$ . Indeed,  $\gamma_+(\cdot)$  remains staying inside  $[\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1]$  and hence solution exists on  $(0, +\infty)$ . To see this, we observe that  $\gamma_+(\cdot)$  is increasing and for each time  $t > 0$ , from (3.20) we have

$$t = \int_{\gamma_+(0)}^{\gamma_+(t)} \frac{dx}{G_1(-V(\varepsilon x, x))},$$

hence the amount of time  $\gamma_+(\cdot)$  needs to reach  $\overline{y_0}$  is

$$\int_{\gamma_+(0)}^{\overline{y_0}} \frac{dx}{G_1(-V(\varepsilon x, x))} = +\infty$$

since  $x \mapsto G_1(-V(\varepsilon x, x))$  is Lipschitz on  $[\varepsilon^{-1}x_0, \varepsilon^{-1}x_0 + 1]$  by Lemma 3.5. We conclude that  $\gamma_+(s) \rightarrow \overline{y_0}$  and similarly  $\gamma_-(s) \rightarrow \underline{y_0}$  as  $s \rightarrow \infty$ .

As a consequence, we have

$$\underline{y_0} \leq \gamma_-(s) \leq \eta_\varepsilon(s) \leq \gamma_+(s) \leq \overline{y_0} \quad \text{for all } s \in [0, \varepsilon^{-1}t_0] \quad (3.22)$$

and thus (3.19) follows. Now we utilize (3.19) to estimate  $A^\varepsilon[\eta_\varepsilon]$ . For any  $\eta_\varepsilon$  solves (3.20) we have

$$A^\varepsilon[\eta_\varepsilon] \geq u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \geq u_0(\varepsilon\eta_\varepsilon(0)) - \|u'_0\|_{L^\infty}\varepsilon. \quad (3.23)$$

On the other hand, by the conservation of energy (3.13) and (3.6) we have

$$L(\dot{\gamma}_+(s)) - V(\varepsilon\gamma_+(s), \gamma_+(s)) = L(\dot{\gamma}_+(s)) + H(L'(\dot{\gamma}_+(s))) = \dot{\gamma}_+(s)L'(\dot{\gamma}_+(s)).$$

Together with the fact that  $|\bar{y}_0 - \varepsilon^{-1}x_0| \leq 1$  from (3.22) we deduce that

$$\begin{aligned}
\inf_{r \leq 0} A^\varepsilon[\eta_\varepsilon] &\leq A^\varepsilon[\gamma_+] = \varepsilon \int_0^{\varepsilon^{-1}t_0} \dot{\gamma}_+(s) L'(\dot{\gamma}_+(s)) ds + u_0(\varepsilon\gamma_+(\varepsilon^{-1}t_0)) \\
&= \varepsilon \int_{\gamma_+(0)}^{\gamma_+(\varepsilon^{-1}t_0)} L'(G_1(-V(\varepsilon x, x))) dx + u_0(\varepsilon\gamma_+(\varepsilon^{-1}t_0)) \\
&= \varepsilon \int_{\gamma_+(0)}^{\gamma_+(\varepsilon^{-1}t_0)} H_1^{-1}(-V(\varepsilon x, x)) dx + u_0(\varepsilon\gamma_+(\varepsilon^{-1}t_0)) \\
&\leq \left( H_1^{-1}(\|V\|_{L^\infty}) + \|u'_0\|_{L^\infty} \right) \varepsilon + u_0(x_0). \tag{3.24}
\end{aligned}$$

From (3.23) and (3.24) we obtain our claim (3.18).  $\square$

For each  $r \in (0, \infty)$ , (3.14) has exactly two distinct solutions  $\eta_{1,r,\varepsilon}(\cdot)$  and  $\eta_{2,r,\varepsilon}(\cdot)$  thanks to the conservation of energy (3.13):

$$\begin{cases} \dot{\eta}_\varepsilon(s) &= G_1(r - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s))) & \text{on } (0, \varepsilon^{-1}t_0), \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0, \end{cases} \tag{3.25}$$

and

$$\begin{cases} \dot{\eta}_\varepsilon(s) &= G_2(r - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s))) & \text{on } (0, \varepsilon^{-1}t_0), \\ \eta_\varepsilon(0) &= \varepsilon^{-1}x_0, \end{cases} \tag{3.26}$$

respectively. Let us consider the first case  $\eta_\varepsilon(\cdot)$  solves (3.25) since the other case is similar. Since  $\dot{\eta}_\varepsilon(s) > 0$  for all  $s \geq 0$ , we have

$$t_0 = \varepsilon \int_0^{\varepsilon^{-1}t_0} \frac{\dot{\eta}_\varepsilon(s)}{\dot{\eta}_\varepsilon(s)} ds = \varepsilon \int_{\eta_\varepsilon(0)}^{\eta_\varepsilon(\varepsilon^{-1}t_0)} \frac{dx}{G_1(r - V(\varepsilon x, x))}. \tag{3.27}$$

Let  $\varepsilon \rightarrow 0$  we deduce that  $\eta_\varepsilon(\varepsilon^{-1}t_0) \rightarrow +\infty$ . It is also clear from (3.25) that

$$t_0 G_1(r) \leq \varepsilon \eta_\varepsilon(\varepsilon^{-1}t_0) - \varepsilon \eta_\varepsilon(0) \leq t_0 G_1(r + \max|V|). \tag{3.28}$$

For a fixed  $\varepsilon > 0$ , we observe that the infimum of the optimization problem (3.16) should be taken over  $r$  not too big.

**Proposition 3.2.** *There exists  $r_0 > 0$  depends on  $\text{Lip}(u_0)$  and  $H(p)$  such that*

$$\inf_{r \geq r_0} A^\varepsilon[\eta_\varepsilon] = \inf_{r \geq r_0} \left\{ A^\varepsilon[\eta_{1,r,\varepsilon}], A^\varepsilon[\eta_{2,r,\varepsilon}] \right\} \geq u^\varepsilon(x_0, t_0) + t_0. \tag{3.29}$$

*Proof.* From assumption (H3) we can define

$$\bar{C} = \sup_{(x,y)} \left\{ |H(x, y, p)| : |p| \leq \|u'_0\|_{L^\infty} \right\} < \infty$$

then  $\bar{u}(x, t) = u_0(x) + \bar{C}t$  is a super-solution to  $(C_\varepsilon)$ . By comparison principle for  $(C_\varepsilon)$  we have

$$u^\varepsilon(x_0, t_0) \leq \bar{u}(x_0, t_0) = u_0(x_0) + \bar{C}t_0. \tag{3.30}$$

On the other hand, using (3.25) in the formula of  $A^\varepsilon[\eta]$  for  $\eta_\varepsilon(\cdot) = \eta_{1,r,\varepsilon}(\cdot)$  we have

$$\begin{aligned} A^\varepsilon[\eta_\varepsilon] &\geq \varepsilon \int_0^{\varepsilon^{-1}t_0} L(\dot{\eta}_\varepsilon(s)) ds + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \\ &= \varepsilon \int_{\eta_\varepsilon(0)}^{\eta_\varepsilon(\varepsilon^{-1}t_0)} \frac{L(G_1(r - V(\varepsilon x, x)))}{G_1(r - V(\varepsilon x, x))} dx + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \\ &\geq \left( \varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - \varepsilon\eta_\varepsilon(0) \right) \left( \frac{G_1(r)}{2} - K_0 \right) + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)). \end{aligned}$$

The deduction from the second line to the third line in the above is due to (3.4). Now utilizing (3.28) and the fact that  $G_1$  is increasing we have further that

$$\begin{aligned} A^\varepsilon[\eta_\varepsilon] &\geq \left( \varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - \varepsilon\eta_\varepsilon(0) \right) \left( \frac{G_1(r)}{2} - K_0 \right) + u_0(x_0) - \|u'_0\|_{L^\infty} |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - \varepsilon\eta_\varepsilon(0)| \\ &= \left( \varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - \varepsilon\eta_\varepsilon(0) \right) \left( \frac{G_1(r)}{2} - K_0 - \|u'_0\|_{L^\infty} \right) + u_0(x_0) \\ &\geq t_0 G_1(r) \left( \frac{G_1(r)}{2} - K_0 - \|u'_0\|_{L^\infty} \right) + u_0(x_0) \end{aligned}$$

for  $r$  large enough such that  $G_1(r) \geq 2(K_0 + \|u'_0\|_{L^\infty})$ . From (3.9) there exists  $r_0 > 0$  such that

$$G_1(r) \left( \frac{G_1(r)}{2} - K_0 - \|u'_0\|_{L^\infty} \right) \geq \bar{C} + 1 \quad \text{for } r \geq r_0.$$

We observe that  $r_0$  only depends on  $H(p)$  and  $u_0$ . This estimate together with (3.30) conclude that  $A^\varepsilon[\eta_{1,r,\varepsilon}] \geq u^\varepsilon(x_0, t_0) + t_0$  for all  $r \geq r_0$ . The case  $\eta_{2,r,\varepsilon}$  solves (3.26) can be done in the same manner, hence we have proved our claim (3.29).  $\square$

With (3.29), the optimization problem (3.16) can be reduced to

$$u^\varepsilon(x_0, t_0) = \min \left\{ \inf_{r \leq 0} A^\varepsilon[\eta_\varepsilon], \inf_{0 < r < r_0} A^\varepsilon[\eta_{1,r,\varepsilon}], \inf_{0 < r < r_0} A^\varepsilon[\eta_{2,r,\varepsilon}] \right\}. \quad (3.31)$$

By (3.6) and the conservation of energy (3.13) and (3.6) with  $\eta_\varepsilon = \eta_{1,r,\varepsilon}$  solves (3.25) we have

$$L(\dot{\eta}_\varepsilon(s)) - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) = -r + \dot{\eta}_\varepsilon(s)L'(\dot{\eta}_\varepsilon(s)). \quad (3.32)$$

Using (3.32) together with (3.25) we can rewrite the action functional as

$$\begin{aligned} A^\varepsilon[\eta_\varepsilon] &= \varepsilon \int_0^{\varepsilon^{-1}t_0} \left( L(\dot{\eta}_\varepsilon(s)) - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s)) \right) ds + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \\ &= -rt_0 + \varepsilon \int_0^{\varepsilon^{-1}t_0} L'_1(\dot{\eta}_\varepsilon(s)) \dot{\eta}_\varepsilon(s) ds + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \\ &= -rt_0 + \varepsilon \int_0^{\varepsilon^{-1}t_0} L'_1 \left( G_1(r - V(\varepsilon\eta_\varepsilon(s), \eta_\varepsilon(s))) \right) \dot{\eta}_\varepsilon(s) ds + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) \\ &= -rt_0 + \varepsilon \int_{\eta_\varepsilon(0)}^{\eta_\varepsilon(\varepsilon^{-1}t_0)} H_1^{-1}(r - V(\varepsilon x, x)) dx + u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)). \end{aligned} \quad (3.33)$$

We now focus on  $0 < r < r_0$ . For simplicity, let us define the following interval  $I_0 \subset \mathbb{R}$  to be

$$I_0 = I_0(T, R) = [-R, c_0 + R] \quad \text{where} \quad c_0 = TG_1(r_0 + \|V\|_{L^\infty}).$$

Since (3.28) is true for all  $0 < r < r_0$ , for all  $(x_0, t_0) \in [-R, R] \times [0, T]$  we have

$$\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) \in I_0.$$

Let  $c_{1,r}$  and  $c_{2,r}$  be unique numbers such that

$$\int_{x_0}^{c_{1,r}} \int_0^1 \frac{dydx}{G_1(r - V(x, y))} = t_0, \quad (3.34)$$

and

$$\int_{c_{2,r}}^{x_0} \int_0^1 \frac{dydx}{G_2(r - V(x, y))} = t_0,$$

respectively.

**Proposition 3.3.** *For  $0 < r < r_0$  we have*

$$|\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - c_{1,r}| \leq C_K\varepsilon. \quad (3.35)$$

where  $C_K = C_K(R, T, H, V)$  is a constant independent of  $r$ .

*Proof.* Let us define

$$\mathcal{K}_r(x, y) = \frac{1}{G_1(r - V(x, y))}, \quad (x, y) \in \mathbb{R} \times \mathbb{T}.$$

From (3.27) and (3.34) we have

$$t_0 = \int_{x_0}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \mathcal{K}_r\left(x, \frac{x}{\varepsilon}\right) dx = \int_{x_0}^{c_{1,r}} \int_0^1 \mathcal{K}_r(x, y) dy dx. \quad (3.36)$$

Using Lemma 2.6 we obtain

$$\left| \int_{x_0}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \mathcal{K}_r\left(x, \frac{x}{\varepsilon}\right) dx - \int_{x_0}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \int_0^1 \mathcal{K}_r(x, y) dy dx \right| \leq K\varepsilon \quad (3.37)$$

where

$$K = 2 \max_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy + 4c_0 \max_{x \in I_0} \int_0^1 \frac{\partial \mathcal{K}_r}{\partial x}(x, y) dy. \quad (3.38)$$

Using (3.36) in (3.37) we have

$$\left| \int_{c_{1,r}}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} \int_0^1 \mathcal{K}_r(x, y) dy dx \right| \leq K\varepsilon.$$

which implies that

$$\left( \min_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \right) |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - c_{1,r}| \leq K\varepsilon. \quad (3.39)$$

On the other hand, we have

$$\frac{\partial \mathcal{K}_r}{\partial x}(x, y) = \left( -V_x(x, y) \frac{G_1'(r - V(x, y))}{G_1(r - V(x, y))} \right) \mathcal{K}_r(x, y)$$

which implies that

$$\max_{x \in I_0} \int_0^1 \left| \frac{\partial \mathcal{K}_r}{\partial x}(x, y) \right| dy \leq \tilde{K} \left( \max_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \right) \quad (3.40)$$

where

$$\tilde{K} = \sup_{0 < r < r_0} \left\{ |V_x(x, y)| \cdot \left| \frac{G_1'(r - V(x, y))}{G_1(r - V(x, y))} \right| : (x, y) \in I_0 \times \mathbb{T} \right\} < \infty$$

by assumption (A2). Now using (3.40) in (3.38) we deduce that

$$K \leq 2 \left( 1 + 2c_0 \tilde{K} \right) \left( \max_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \right). \quad (3.41)$$

Use (3.41) in (3.39) we deduce that

$$\left( \min_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \right) |\varepsilon \eta_\varepsilon(\varepsilon^{-1}t) - c_{1,r}| \leq 2 \left( 1 + 2c_0 \tilde{K} \right) \left( \max_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy \right) \varepsilon. \quad (3.42)$$

From (3.42) we obtain our claim (3.35), where

$$C_K = 2 \left( 1 + 2c_0 \tilde{K} \right) \sup_{0 < r < r_0} \left( \frac{\max_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy}{\min_{x \in I_0} \int_0^1 \mathcal{K}_r(x, y) dy} \right) < \infty$$

by assumption (A4).  $\square$

In view of (3.33), we aim to show that for  $0 < r < r_0$  then

$$\left| \int_{x_0}^{\varepsilon \eta_\varepsilon(\varepsilon^{-1}t_0)} H_1^{-1}(r - V(x, \varepsilon^{-1}x)) dx - \int_{x_0}^{c_{1,r}} \int_0^1 H_1^{-1}(r - V(x, y)) dy dx \right| \leq C_F \varepsilon \quad (3.43)$$

where  $C_F$  is a constant independent of  $r$ . To see that, let

$$\mathcal{F}_r(x, y) = H_1^{-1}(r - V(x, y)), \quad (x, y) \in \mathbb{R} \times \mathbb{T}.$$

By Lemma 2.6 we have

$$\left| \int_{x_0}^{c_{1,r}} H_1^{-1}(r - V(x, \varepsilon^{-1}x)) dx - \int_{x_0}^{c_{1,r}} \int_0^1 H_1^{-1}(r - V(x, y)) dy dx \right| \leq (2F_1 + c_0 F_2) \varepsilon. \quad (3.44)$$

where

$$F_1 := H_1^{-1}(r_0 + \|V\|_{L^\infty}) \geq \max_{x \in I_0} \int_0^1 \mathcal{F}_r(x, y) dy \quad (3.45)$$

and

$$F_2 := \sup \left\{ \frac{|V_x(x, y)|}{|G_1(-V(x, y))|} \mid (x, y) \in I_0 \times \mathbb{T} \right\} \geq \max_{x \in I_0} \int_0^1 \left| \frac{\partial \mathcal{F}_r}{\partial x}(x, y) \right| dy \quad (3.46)$$

which is finite by assumption (A3). Here we used the fact that  $\frac{d}{dx} (H_1^{-1}(x)) = \frac{1}{G_1(x)}$  for all  $x > 0$ . On the other hand, it is obvious that

$$\left| \int_{x_0}^{\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)} H_1^{-1}(r - V(x, \varepsilon^{-1}x)) dx - \int_{x_0}^{c_{1,r}} H_1^{-1}(r - V(x, \varepsilon^{-1}x)) dx \right| \leq F_1 |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - c_{1,r}| \leq C_K F_1 \varepsilon \quad (3.47)$$

thanks to (3.35). From (3.44) and (3.47) we obtain our claim (3.43) with

$$C_F = 2F_1 + c_0 F_2 + C_K F_1.$$

**Proposition 3.4.** *We have the following estimate:*

$$\left| \inf_{\substack{0 < r < r_0 \\ i=1,2}} A^\varepsilon[\eta_{i,r,\varepsilon}] - \inf_{0 < r < r_0} I(r) \right| \leq C\varepsilon \quad (3.48)$$

where  $C$  is a constant independent of  $r$  and  $I(r) = \min \{I_1(r), I_2(r)\}$  where

$$I_1(r) = -rt_0 + \int_{x_0}^{c_{1,r}} \int_0^1 H_1^{-1}(r - V(x, y)) dy dx + u_0(c_{1,r}), \quad (3.49)$$

$$I_2(r) = -rt_0 + \int_{c_{2,r}}^{x_0} \int_0^1 H_2^{-1}(r - V(x, y)) dy dx + u_0(c_{2,r}). \quad (3.50)$$

*Proof.* Within our notation  $\eta_{1,r,\varepsilon} = \eta_\varepsilon$ , we have

$$|u_0(\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0)) - u_0(c_{1,r})| \leq \|u'_0\|_{L^\infty} |\varepsilon\eta_\varepsilon(\varepsilon^{-1}t_0) - c_r| \leq (C_K \|u'_0\|_{L^\infty}) \varepsilon \quad (3.51)$$

since  $u_0 \in \text{Lip}(\mathbb{R})$ . In view of (3.33), (3.43) and (3.51) we conclude that

$$|A^\varepsilon[\eta_{1,r,\varepsilon}] - I_1(r)| \leq (C_F + C_K \|u'_0\|_{L^\infty}) \varepsilon. \quad (3.52)$$

Taking the infimum over  $0 < r < r_0$  in (3.52) we obtain our claim (3.48) with  $i = 1$  where

$$C_1 = C_F + C_K \|u'_0\|_{L^\infty}. \quad (3.53)$$

Doing similarly with the other case  $\eta_{2,r,\varepsilon}$  solves (3.26) we obtain

$$\left| \inf_{0 < r < r_0} A^\varepsilon[\eta_{2,r,\varepsilon}] - \inf_{0 < r < r_0} I_2(r) \right| \leq C_2 \varepsilon \quad (3.54)$$

where  $C_2$  is a constant independent of  $r$  in the same manner as  $C_1$  in (3.53). Thus our claim(3.48) is correct with  $C = \max\{C_1, C_2\}$ .  $\square$

Finally, using (3.18) and (3.48) in (3.31) we conclude that

$$|u^\varepsilon(x_0, t_0) - u(x_0, t_0)| \leq \left( \max \left\{ H_1^{-1}(\|V\|_{L^\infty}) + \|u'_0\|_{L^\infty}, C \right\} \right) \varepsilon.$$

Thus the proof is complete.

**Lemma 3.5.** *Let  $\mathcal{V} \in C^2([0, 1], [0, \infty))$  with  $\min_{x \in [0,1]} \mathcal{V}(x) = 0$  and  $\mathcal{V}(0) = \mathcal{V}(1)$ .*

(i) Let  $H, G_1, G_2$  be defined as in Theorem 1.2. If

$$\limsup_{p \rightarrow 0} \left| \frac{H''(p)}{H'(p)} \sqrt{H(p)} \right| < \infty, \quad (3.55)$$

then  $x \mapsto G_i(\mathcal{V}(x))$  is Lipschitz on  $[0, 1]$  for  $i = 1, 2$ .

(ii) If  $H$ , defined in Theorem 1.2, satisfies  $H''(0) > 0$  then we have something stronger than (3.55):

$$\limsup_{p \rightarrow 0} \left| \frac{\sqrt{H(p)}}{H'(p)} \right| < \infty. \quad (3.56)$$

In this case we have further that

$$C_{1,I}\sqrt{x} \leq |G_i(x)| \leq C_{2,I}\sqrt{x}$$

on any bounded subset  $I \subset \mathbb{R}$ , where  $i = 1, 2$  and  $C_{I,1}, C_{I,2} > 0$ .

(iii) If  $H$ , defined in Theorem 1.2, satisfies  $H \in C^3(\mathbb{R})$  then

$$\limsup_{p \rightarrow 0} \frac{|H''(p)|}{\sqrt{|H'(p)|}} < \infty. \quad (3.57)$$

As a consequence, we have something stronger than (3.55):

$$\limsup_{p \rightarrow 0} \left| \frac{H''(p)}{H'(p)} \sqrt{\frac{H(p)}{|p|}} \right| < \infty. \quad (3.58)$$

(iv) If  $H(p) = |p|^\gamma$  where  $\gamma \geq 2$  then (3.55) holds true.

*Proof.*

(i) It suffices to show for  $G_1$  since the argument is similar for  $G_2$ . For simplicity, let us denote  $G_1, H_1^{-1}$  by  $G, H^{-1}$ . For  $0 < \varepsilon < 1$  let  $f_\varepsilon(x) = G(\mathcal{V}(x) + \varepsilon)$  then  $f_\varepsilon \in C^2([0, 1])$  and

$$f'_\varepsilon(x) = \frac{\mathcal{V}'(x)}{\sqrt{\mathcal{V}(x) + \varepsilon}} \left( \frac{H''(H^{-1}(\mathcal{V}(x) + \varepsilon))}{H'(H^{-1}(\mathcal{V}(x) + \varepsilon))} \sqrt{\mathcal{V}(x) + \varepsilon} \right).$$

For  $x \in [0, 1]$  such that  $\mathcal{V}(x) = \mathcal{V}'(x) = 0$  then obviously  $f'_\varepsilon(x) = 0$ , while if  $x \in [0, 1]$  such that  $\mathcal{V}(x) \neq 0$  then from (3.55) and Lemma 2.7 we have

$$|f'_\varepsilon(x)| \leq \left| \frac{\mathcal{V}'(x)}{\sqrt{\mathcal{V}(x)}} \right| \cdot \left| \frac{H''(\xi)}{H'(\xi)} \sqrt{H(\xi)} \right| \leq L \left( \sup_{[0, p^*]} \left| \frac{H''(p)}{H'(p)} \sqrt{H(p)} \right| \right) < \infty$$

where  $\xi = H^{-1}(\mathcal{V}(x) + \varepsilon)$  and  $p^* = H^{-1}(\|V\|_{L^\infty} + 1)$ . Therefore  $f_\varepsilon$  is Lipschitz on  $[0, 1]$  with a Lipschitz constant independent of  $\varepsilon > 0$ . Let  $\varepsilon \rightarrow 0$  we deduce that  $x \mapsto G(\mathcal{V}(x))$  is Lipschitz on  $[0, 1]$ .

(ii) If  $H''(0) > 0$  then there exists  $\delta > 0$  so that  $H''(p) \geq c > 0$  for  $p \in (-\delta, \delta)$ , hence by Taylor's expansion we deduce that there are some  $m, M > 0$  such that

$$m|p|^2 \leq H(p) \leq M|p|^2 \quad \text{and} \quad m|p| \leq |H'(p)| \leq M|p|. \quad (3.59)$$

From that (3.56) follows easily. On the other hand, by definition  $G_i(x) = H'(H_i^{-1}(x))$  and (3.59) we deduce that for all  $x$  small then

$$\sqrt{\frac{m}{M}}\sqrt{x} \leq |H'(H_i^{-1}(x))| \leq \sqrt{\frac{M}{m}}\sqrt{x}. \quad (3.60)$$

Since  $G_i(x) = 0$  if and only if  $x = 0$ , we have (3.60) is true for any bounded set of  $\mathbb{R}$  after modifying the two upper bound and lower bound constants.

(iii) Using the convexity we have  $H(p) \leq pH'(p)$  for all  $p$ , hence

$$\left| \frac{H''(p)}{H'(p)} \sqrt{H(p)} \right| \leq \frac{H''(p)}{\sqrt{|H'(p)|}} \sqrt{|p|}. \quad (3.61)$$

Let  $g(p) = H'(p) \in C^2(\mathbb{R})$  is strictly increasing on  $(0, \infty)$  and is strictly decreasing on  $(-\infty, 0)$  with  $g(0) = 0$ , we claim that indeed

$$\limsup_{p \rightarrow 0} \frac{g'(p)}{\sqrt{|g(p)|}} < \infty. \quad (3.62)$$

This can be done by a similar argument to Lemma 2.7, hence (3.58) follows.

(iv) It is clear from direct computation. □

**Corollary 3.6.** *We have the following representation formula*

$$u(x_0, t_0) = \min \left\{ u_0(x_0), \min \left\{ \inf_{0 < r < r_0} I_1(r), \inf_{0 < r < r_0} I_2(r) \right\} \right\}.$$

where  $I_1(r)$  and  $I_2(r)$  are defined in (3.49) and (3.50), respectively.

*Proof of Corollary 1.3.* We need to check conditions (A0),(A2),(A3),(A4) of Theorem 1.2 to apply that theorem. Let us fix a compact interval  $I \subset \mathbb{R}$ , for simplicity in the assumption of  $V$  let us denote  $\alpha, \beta, f$  by  $\alpha_I, \beta_I, f_I$ .

If  $H(p) = |p|^\gamma$  where  $\gamma \geq 2$  then  $|G_i(p)| = \gamma|p|^{1-\frac{1}{\gamma}}$  and  $|G'_i(p)| = (\gamma - 1)|p|^{-\frac{1}{\gamma}}$ . Therefore conditions (A0),(A2),(A3) follow from direct computation. (A4) follows from the fact that  $p \mapsto |G_i(p)|$  is increasing and for any compact interval  $I \subset \mathbb{R}$  then

$$\begin{aligned} \max_{x \in I} \int_0^1 \frac{dy}{|G_i(r - V(x, y))|} &\leq \int_0^1 \frac{dy}{|G_i(r + \alpha f(y))|} \\ &\leq \int_0^1 \frac{dy}{|G_i(\alpha/\beta (r + \beta f(y)))|} \\ &= \frac{1}{\gamma} \left( \frac{\beta}{\alpha} \right)^{1-\frac{1}{\gamma}} \int_0^1 \frac{dy}{|G_i(r + \beta f(y))|} \\ &\leq \frac{1}{\gamma} \left( \frac{\beta}{\alpha} \right)^{1-\frac{1}{\gamma}} \min_{x \in I} \int_0^1 \frac{dy}{|G_i(r - V(x, y))|}. \end{aligned}$$

In case  $H''(0) > 0$ , condition (A0) follows from Lemma 3.5. Now on the bounded set  $[0, \|V\|_{L^\infty} + 1]$  by Lemma 3.5 we have

$$C_1\sqrt{x} \leq |G_i(x)| \leq C_2\sqrt{x} \quad (3.63)$$



for  $i = 1, 2$  and for some  $C_1, C_2 > 0$ . For  $i = 1, 2$ ,  $0 < r < 1$  and  $x \in I$  we have

$$|V_x(x, y)| \cdot \frac{|G'_i(r - V(x, y))|}{|G_i(r - V(x, y))|} \leq \frac{|V_x(x, y)|}{|V(x, y)|} \left( \frac{\sqrt{H(\xi)}}{|G_i(H(\xi))|} \right) \left( \frac{|H''(\xi)|}{|H'(\xi)|} \sqrt{H(\xi)} \right)$$

where  $\xi = H_i^{-1}(r - V(x, y))$ . The right hand side is bounded as  $r \rightarrow 0^+$  due to (A0), (3.63) and (1.4), thus (A2) follows. Condition (A3) is true since for  $x \in I$  we have

$$\left| \frac{V_x(x, y)}{G_i(V(x, y))} \right| \leq \left| \frac{V_x(x, y)}{V(x, y)} \right| \cdot \frac{\sqrt{|V(x, y)|}}{|G_i(|V(x, y)|)|} \cdot \sqrt{|V(x, y)|}.$$

The right hand side is bounded as well, due to (1.4), (3.63) and the fact that  $V$  bounded. Finally, for  $i = 1, 2$  we have  $x \mapsto |G_i(x)|$  is increasing, from (3.63) we deduce that for  $0 < r < 1$  then

$$\max_{x \in I} \int_0^1 \frac{dy}{|G_i(r - V(x, y))|} \leq \int_0^1 \frac{dy}{|G_i(r + \alpha f(y))|} \leq \int_0^1 \frac{dy}{C_1 \sqrt{r + \alpha f(y)}}$$

while

$$\min_{x \in I} \int_0^1 \frac{dy}{|G_i(r - V(x, y))|} \geq \int_0^1 \frac{dy}{|G_i(r + \beta f(y))|} \geq \int_0^1 \frac{dy}{C_2 \sqrt{r + \beta f(y)}}.$$

Since  $\alpha \leq \beta$ , we have  $\sqrt{r + \alpha f(y)} \geq \sqrt{\frac{\alpha}{\beta}(r + \beta f(y))}$ , therefore

$$\int_0^1 \frac{dy}{C_1 \sqrt{r + \alpha f(y)}} \leq \left( \frac{C_2}{C_1} \sqrt{\frac{\beta}{\alpha}} \right) \int_0^1 \frac{dy}{C_2 \sqrt{r + \beta f(y)}}$$

and thus (A4) follows.  $\square$

#### ACKNOWLEDGMENTS

The author would like to express his appreciation to his advisor, Hung V. Tran for giving him this interesting problem and for his great guidance. The author also would like to thank Jingrui Cheng for helpful comments, discussions and being a great friend with a lot of supports.

#### REFERENCES

- [1] ARMSTRONG, S., CARDALIAGUET, P., AND SOUGANIDIS, P. Error estimates and convergence rates for the stochastic homogenization of Hamilton-Jacobi equations. *Journal of the American Mathematical Society* 27, 2 (2014), 479–540.
- [2] BARDI, M., AND CAPUZZO-DOLCETTA, I. *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations*. Modern Birkhäuser Classics. Birkhäuser Basel, 1997.
- [3] CAMILLI, F., ANNALISA, C., AND CLAUDIO, M. Homogenization and vanishing viscosity in fully nonlinear elliptic equations: Rate of convergence estimates. *Advanced Nonlinear Studies* 11 (03 2016), 405–428.
- [4] CAPUZZO-DOLCETTA, I., AND ISHII, H. On the rate of convergence in homogenization of Hamilton-Jacobi equations. *Indiana University Mathematics Journal* 50, 3 (2001), 1113–1129.
- [5] CONCORDEL, M. C. Periodic homogenization of Hamilton-Jacobi equations: Additive eigenvalues and variational formula. *Indiana University Mathematics Journal* 45, 4 (1996), 1095–1117.

- [6] CONCORDEL, M. C. Periodic homogenisation of HamiltonJacobi equations: 2. eikonal equations. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics* 127, 4 (1997), 665–689.
- [7] CRANDALL, M. G., ISHII, H., AND LIONS, P.-L. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)* 27 (1992) 1–67 (July 1992).
- [8] EVANS, L. C. The perturbed test function method for viscosity solutions of nonlinear pde. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics* 111, 3–4 (1989), 359–375.
- [9] EVANS, L. C. Periodic homogenization of certain fully nonlinear partial differential equations. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics* 120 (1992), 245 – 265.
- [10] LE, Q. N., MITAKE, H., AND TRAN, H. V. *Dynamical and Geometric Aspects of Hamilton-Jacobi and Linearized Monge-Ampre Equations*, vol. 2183 of *Lecture Notes in Mathematics*. Springer International Publishing, 2017.
- [11] LIONS, P.-L. *Generalized Solutions of Hamilton-Jacobi Equations*. Chapman & Hall/CRC research notes in mathematics series. Pitman, 1982.
- [12] LIONS, P.-L., PAPANICOLAOU, G., AND VARADHAN, S. R. Homogenization of Hamilton-Jacobi equations. *Unpublished preprint* (1986).
- [13] LUO, S., TRAN, H. V., AND YU, Y. Some inverse problems in periodic homogenization of Hamilton-Jacobi equations. *Archive for Rational Mechanics and Analysis* 221, 3 (Sep 2016), 1585–1617.
- [14] LUO, S., YU, Y., AND ZHAO, H. A new approximation for effective hamiltonians for homogenization of a class of HamiltonJacobi equations. *Multiscale Modeling & Simulation* 9, 2 (2011), 711–734.
- [15] MITAKE, H., AND TRAN, H. V. Homogenization of weakly coupled systems of Hamilton-Jacobi equations with fast switching rates. *Archive for Rational Mechanics and Analysis* 211, 3 (Mar 2014), 733–769.
- [16] MITAKE, H., TRAN, H. V., AND YU, Y. Rate of convergence in periodic homogenization of Hamilton-Jacobi equations: the convex setting. *ArXiv e-prints* (Dec 2018).
- [17] NEUKAMM, S. An introduction to the qualitative and quantitative theory of homogenization. *ArXiv e-prints* (July 2017).
- [18] QIAN, J., TRAN, H. V., AND YU, Y. Min–max formulas and other properties of certain classes of nonconvex effective Hamiltonians. *Mathematische Annalen* (Oct 2017).

(S. N.T. Tu) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN MADISON, 480 LINCOLN DRIVE, MADISON, WI 53706, USA

*E-mail address:* thaison@math.wisc.edu