

# $p$ -Branes with $AdS_{p+1}$ vacuum as models of $R^2$ gravity

A. A. Zheltukhin <sup>a,b\*</sup>

<sup>a</sup> Kharkov Institute of Physics and Technology, Kharkov, 61108, Ukraine

<sup>b</sup> Nordita, KTH Royal Institute of Technology and Stockholm University  
SE 106 91 Stockholm, Sweden

October 16, 2019

## Abstract

Branes with *constant mean curvature* of their hyper-worldsheets of codim 1 are treated as the Nambu-Goldstone fields of the broken Poincare symmetry. Mapping of their action into quadratic curvature gravity action in hyper-ws is shown. Equation for the brane *potential* extremals and its solution describing hyper-ws of *constant curvature* are found. For membranes in  $\mathbf{R}^{1,3}$  this extremum is shown to be a *saddle* 3-dim. hypersurface which defines classically unstable vacuum.

## 1 Introduction

Differential geometric description of hyper-worldsheets (h-ws) [1 - 4] of the Dirac (or fundamental) branes [5] as hypersurfaces embedded into Riemannian spaces revealed new connections between the theories of strings, gauge and gravitational fields [6 - 9]. Embedding of a h-ws  $\Sigma_{p+1}^{min}$  of fundamental  $p$ -brane into  $D$ -dim. Minkowski space results in the spontaneous breaking of the Poincare group  $ISO(1, D - 1)$  to its subgroup  $ISO(1, p) \times SO(D - p - 1)$  [10 - 13]. The corresponding Nambu-Goldstone (N-G) fields (see [14] and

---

\*e-mail: aaz@nordita.org

refs. there) were used for construction of the invariant action of fundamental  $p$ -branes in the gauge theory formalism on the curved h-ws [15]. The latter introduced the  $SO(D - p - 1)$  Y-M multiplet  $B_\mu^{ab}$ , the N-G tensor multiplet  $l_{\mu\nu}^a$  identified with the second fundamental form of  $\Sigma_{p+1}^{min}$  equipped with the metric tensor  $g_{\mu\nu}$ . As shown in [16] for codim 1, the action [15] realizes a  $(p + 1)$ -dim. h-ws version of quadratic curvature gravity investigated in [17 - 21]. The pure  $R^2$  term and cosmological constant, represented in the action  $S_{Dir}$  [16] by an integration constant, form the quartic scalar potential of interaction  $V_{Dir}(l, g)$  for the fields  $l_{\mu\nu}$  and  $g_{\mu\nu}$ . For zero cosmological constant the action  $S_{Dir}$  contains only one coupling constant  $\kappa_p \sim T_p^{\frac{3-p}{2(p+1)}}$  depending on the  $p$ -brane tension  $T_p$ . The power law provides three different regimes of the behavior of  $k_p$  as a function of the tension and, consequently, energy corresponding to the cases  $p < 3$ ,  $p = 3$  and  $p > 3$ . These regimes show the presence of the asymptotic freedom and confinement phases and their connection with inflation and collapse of the branes. Exact solutions for  $p$ -brane collapse were earlier obtained in [22], [23]. The dimensionless coupling  $k_3$  associated with three-branes characterizes a scale invariant model belonging to those considered in [24 - 29]. The model [16] does not contain invariants linear in components of the h-ws Riemannian tensor. This necessitates the model extension which could encode the Hilbert-Einstein(H-E) term [30]. To this objective we note that h-ws of *fundamental* branes are *minimal* hypersurfaces with *zero mean curvature*, i.e.  $Spl \equiv g^{\nu\mu} l_{\nu\mu} = 0$ . A natural generalization of the latter condition is  $Spl = \mu$ , where  $\mu$  is a constant with the dimension of mass  $[\mu] = [L^{-1}]$ . This condition selects hypersurfaces with the *constant mean curvature* (cmc) equal to  $\mu$ . The string model with cmc ws embedded into  $\mathbf{R}^{1,2}$  was studied in [31] by the extension of the Nambu-Goto action. In the gauge formalism  $\mu$  is treated as a physical constant fixing a characteristic scale of the symmetry breaking. A desired extension is supposed to have two fundamental couplings  $T_p$  and  $\mu$  defining the brane dynamics in the presence of gravitational and elastic forces. A similar dynamics is observed for string in curved space [32]. Here we solve the problem in question and build the potential  $V_{cmc}(l, g, \mu)$  which encodes the H-E and cosmological terms (50) in addition to the pure  $R^2$  term. The proposed potential extends the Starobinsky model [18] and its generalization [33] including a massless scalar field  $\phi$ . In particular, the potential contains the cosmological and interaction terms. The latter term is proportional to  $\mu R_{\nu\lambda} l^{\lambda\nu}$ . It should be also noted that the field  $\phi$  from [33] can be identified with  $Spl$  treated as the *dynamical mean*

*curvature field* constrained to be constant in our approach.

Then we derive the equation for extremals of the functional  $V_{cmc}(l, g, \mu)$  and find its general solution for the most interesting case  $R \neq 0$ . Consistency of the solution with the Gauss and Peterson-Codazzi (P-C) embedding conditions results in the matrix equation for null vectors of the  $(p+1)$ -dim. Einstein tensor (matrix)  $G_{\mu\nu}$ . Its solution reveals the presence of the extremal  $l_{o\mu\nu}(\xi) = \frac{\mu}{p+1}g_{\mu\nu}(\xi)$ . This critical point corresponds to an  $AdS_{p+1}$ -like hyper-ws  $\Sigma_{p+1}^o$  of *constant negative curvature*. Next we analyse the behavior of  $V_{cmc}$  in the vicinity of  $l_o(\xi)$ . The second partial derivatives of  $V_{cmc}$  with respect to  $l$  at  $l_o(\xi)$  have different signs depending, in particular, on the brane dimension  $p$ . So, for string the extremum is the minimum that shows stability of the string potential in  $\mathbf{R}^{1,2}$ . But for membrane ( $p=2$ ) the critical point  $l_o(\xi)$  is proved to be the *saddle* 3-dim. hypersurface. It points to the classical unstability of the *cmc membrane* potential in 4-dim. Minkowski space-time. The classical unstability of the fundamental (super)membrane [36], described by the potential  $V(x(\xi))$  [37], is well known. Since *fundamental* membrane sweeps *minimal* h-ws, its change by *cmc* h-ws embedded into  $\mathbf{R}^{1,3}$  does not remove the unstability. The obtained results can be used while studying the braneworld cosmological models [34], [35], as well as for investigation of inflation and reheating mechanisms in  $R^2$  gravity models.

## 2 Branes with minimal hyper-worldsheets

In the geometrical approach Dirac  $p$ -branes embedded into  $\mathbf{R}^{1,D-1}$  with the signature  $(+ - \dots -)$  are described as dynamical systems with the Poincare symmetry  $ISO(1, D-1)$  spontaneously broken to  $ISO(1, p) \times SO(D-p-1)$ . The  $p$ -branes sweep minimal h-ws  $\Sigma_{p+1}^{min}$  with zero mean curvature

$$Sp l^a := g^{\mu\nu} l_{\nu\mu}^a = 0, \quad (1)$$

where  $g_{\mu\nu}$  is the induced metric in  $\Sigma_{p+1}^{min}$ . The symmetric tensor  $l_{\mu\nu}^a(\xi)$  is the second fundamental form of the h-ws treated as a constrained multiplet of the local group  $SO(D-p-1)$ . The indexes  $a, b = p+1, p+2, \dots, D-1$  enumerate the orts  $\mathbf{n}_a(\xi^\rho)$  of an orthonormal moving frame attached to  $\Sigma_{p+1}^{min}$  and orthogonal to it. The internal coordinates  $\xi^\mu = (\tau, \sigma^r)$ ,  $r = 1, 2, \dots, p$  parametrize the h-ws world vector  $\mathbf{x}(\xi^\rho)$ . The map of the Dirac  $p$ -brane action to the diff and  $SO(D-p-1)$  invariant h-ws action is realized by the

field action [15]

$$S_{Dir} = \frac{1}{k_p^2} \int d^{p+1} \xi \sqrt{|g|} \left\{ -\frac{1}{4} Sp(H_{\mu\nu} H^{\nu\mu}) + \frac{1}{2} \nabla_{\mu}^{\perp} l_{\nu\rho a} \nabla^{\perp \{\mu} l^{\nu\} \rho a} - \nabla_{\mu}^{\perp} l_{\rho a}^{\mu} \nabla_{\nu}^{\perp} l^{\nu\rho a} + V(l, g) \right\}, \quad (2)$$

where the brackets  $\{\mu\nu\}$ ,  $[\mu\nu]$  imply the  $\mu, \nu$  symmetrization and antisymmetrization, respectively. The potential energy term  $V(l, g)$  is equal to

$$V(l, g) = V_{Dir} := -\frac{1}{2} Sp(l_a l_b) Sp(l^a l^b) + Sp(l_a l_b l^a l^b) - Sp(l_a l^a l_b l^b) + c_p. \quad (3)$$

The kinetic term for the  $SO(D-p-1)$  gauge field  $B_{\mu}^{ab}(\xi) = -B_{\mu}^{ba}(\xi)$  in the fundamental representation is given by the trace  $Sp(H_{\mu\nu} H^{\nu\mu}) := H_{\mu\nu}^{ab} H_{ba}^{\nu\mu}$

$$H_{\mu\nu a}{}^b := (\partial_{[\mu} B_{\nu]} + [B_{\mu}, B_{\nu}])_a{}^b. \quad (4)$$

The metric and Y-M covariant derivative  $\nabla_{\mu}^{\perp} l_{\nu\rho}{}^a$  is defined by the expression

$$\nabla_{\mu}^{\perp} l_{\nu\rho}{}^a := \partial_{\mu} l_{\nu\rho}{}^a - \Gamma_{\mu\nu}^{\lambda} l_{\lambda\rho}{}^a - \Gamma_{\mu\rho}^{\lambda} l_{\nu\lambda}{}^a + B_{\mu}^{ab} l_{\nu\rho b}. \quad (5)$$

The h-ws metric  $g_{\mu\nu}(\xi)$  in (2) is treated as a background field since its equations are encoded by the Gauss's Theorema Egregium

$$R_{\mu\nu}{}^{\gamma}{}_{\lambda} = l_{[\mu}{}^{\gamma a} l_{\nu]\lambda a} \quad (6)$$

used under construction of action (2). The discussed gauge approach can be generalized for the construction of h-ws with non-zero mean curvature. Solution of this problem is reduced to such a deformation of  $V_{Dir}$  (3) which makes it consistent with the cmc condition. Below we solve this problem.

### 3 Brane h-ws with constant mean curvature

For h-ws of codimension 1 the gauge field  $B_{\mu}^{ab} \equiv 0$  since  $a = b = p + 1$  and (2) encodes description of quadratic curvature gravity on the string/brane-ws with zero mean curvature by means of the reduced potential (3)

$$V_{Dir} = -\frac{1}{2} Sp(l^2) Sp(l^2) + c_p = -\frac{1}{2} R^2 + c_p, \quad R := g^{\mu\nu} R_{\mu\nu} \quad (7)$$

as shown in [16]. To find a deformed potential  $V_{cmc}$  corresponding to a cmc hypersurface  $\Sigma_{p+1}^{cmc}$  of codim 1 we consider the h-ws action

$$S = \frac{1}{k_p^2} \int d^{p+1} \xi \sqrt{|g|} \left( \frac{1}{2} \nabla_\mu l_{\nu\rho} \nabla^{\{\mu} l^{\nu\}\rho} - \nabla_\mu l_\rho^\mu \nabla_\nu l^{\nu\rho} - V(l, g) \right) \quad (8)$$

and obtain the following EOM for the reduced N-G field  $l_{\mu\nu} := -l_{\mu\nu}^{(p+1)}$

$$\frac{1}{2} \nabla_\mu \nabla^{[\mu} l^{\nu]\rho\} = -[\nabla^\mu, \nabla^{\{\nu} l_\mu^{\rho\}}] + \frac{\partial V}{\partial l_{\nu\rho}}. \quad (9)$$

The (P-C) conditions for the covariant derivatives (5) with  $B_\mu = 0$

$$\nabla_{[\mu} l_{\nu]\rho} = 0, \quad (10)$$

used in (9) nullify their l.h.s. and yield the consistency condition for  $V(l, g)$

$$\frac{\partial V}{\partial l_{\nu\rho}} = [\nabla^\mu, \nabla^{\{\nu} l_\mu^{\rho\}}]. \quad (11)$$

Using the Bianchi identities (BI) for the commutator in (11)

$$[\nabla_\mu, \nabla_\nu] l^{\gamma\rho} = R_{\mu\nu}{}^{\gamma\lambda} l^{\lambda\rho} + R_{\mu\nu}{}^{\rho\lambda} l^{\gamma\lambda} \quad (12)$$

and the Gauss eqs. (6) we present the r.h.s. of Eqs. (11) in the form

$$\frac{1}{2} [\nabla^\mu, \nabla^{\{\nu} l_\mu^{\rho\}}] = (l^2)^{\nu\rho} Sp l - l^{\nu\rho} Sp(l^2), \quad (13)$$

where  $Sp(l^2) = l_{\mu\rho} l_\nu^\rho g^{\mu\nu}$ . Substitution of (13) into (11) yields the PDE's

$$\frac{1}{2} \frac{\partial V}{\partial l_{\nu\rho}} = (l^2)^{\nu\rho} Sp l - l^{\nu\rho} Sp(l^2). \quad (14)$$

for  $V(l, g)$ . Solution of Eqs. (14) corresponding to the cmc  $\mu(p)$  is

$$V = V_{cmc} := \frac{2}{3} \mu Sp(l^3) - \frac{1}{2} Sp(l^2) Sp(l^2) + c_p, \quad Sp l = \mu(p), \quad (15)$$

where  $c_p$  is an integration constant and the cmc condition is treated as the inverse Higgs effect. So, the action (8) with  $V_{cmc}$  (15)

$$S_{cmc} = \frac{1}{k_p^2} \int d^{p+1} \xi \sqrt{|g|} \left( \frac{1}{2} \nabla_\mu l_{\nu\rho} \nabla^{\{\mu} l^{\nu\}\rho} - \nabla_\mu l_\rho^\mu \nabla_\nu l^{\nu\rho} - \frac{2}{3} \mu Sp(l^3) + \frac{1}{2} Sp(l^2) Sp(l^2) - c_p \right) \quad (16)$$

is consistent with the Gauss (6) and P-C (10) conditions. This action describes diff invariant self-interaction of the N-G field  $l_{\mu\nu}$ . From the viewpoint of the variational principle the h-ws metric  $g_{\mu\nu}$  works as a background metric since its evolution is defined by Eqs.(6). The latter permits to present the Euler-Lagrange eqs. produced by (16) as

$$\frac{1}{2}\nabla_{\mu}\nabla^{[\mu}l^{\nu]\rho\}} = 0, \quad Spl = \mu. \quad (17)$$

Using the identity  $\nabla^{\perp[\mu}l^{\nu]\rho]} = -\nabla^{\perp[\nu}l^{\rho]\mu]}$  transforms (17) into the equation

$$\frac{1}{2}\nabla_{\mu}\nabla^{[\mu}l^{\nu]\rho\}} \equiv \nabla_{\mu}(\nabla^{[\mu}l^{\nu]\rho} + \frac{1}{2}\nabla^{[\nu}l^{\rho]\mu}) = 0. \quad (18)$$

Contracting (18) with  $g_{\rho\nu}$  and using the cmc  $Spl = \mu$  we obtain the covariant conservation law for the divergence  $\nabla^{\rho}l_{\rho}^{\mu}$

$$\nabla_{\mu}(\nabla^{\rho}l_{\rho}^{\mu}) = 0. \quad (19)$$

Now it is suitable to rewrite EOM (18) in the form

$$\nabla_{\mu}\nabla^{[\mu}l^{\nu]\rho} = -\frac{1}{2}\{[\nabla^{\mu}, \nabla^{[\nu]}l^{\rho]}_{\mu} + \nabla^{[\nu}\nabla_{\mu}l^{\rho]\mu}\}.$$

Using the relation for the  $(\nu, \rho)$ -antisymmetric part of the commutator

$$\frac{1}{2}[\nabla^{\perp\mu}, \nabla^{\perp[\nu]}l^{\rho]a}] = \frac{1}{2}([l^a, l^b])^{\nu\rho}Spl_b, \quad (20)$$

given by the Bianchi identities and Gauss conditions, shows vanishing of (20) for codim 1. As a result, EOM (17) are represented in the form

$$\nabla_{\mu}\nabla^{[\mu}l^{\nu]\rho} + \frac{1}{2}\nabla^{[\nu}\nabla_{\mu}l^{\rho]\mu} = 0, \quad Spl = \mu. \quad (21)$$

Eqs. (21) are consistent with the P-C eqs. (10) in view of the identity

$$\nabla^{\mu}l_{\mu\lambda} = \nabla_{\lambda}Spl \equiv \partial_{\lambda}Spl = 0 \quad (22)$$

following from (10) after contraction of  $\mu$  and  $\rho$  and using the cmc condition. Consideration of the local conservation law (19) together with (22) hints that the latter can be treated as initial data constraints (IDC) preserved by EOM (21). Moreover, we intend to check a possibility to interpret all the P-C eqs. (10) as the IDC preserved by PDEs (21).

## 4 Cauchy constraints for cmc brane EOM

The Cauchy problem for PDEs (21) is studied here treating all eqs. (10) as the IDC fixed at the moment  $\tau = 0$

$$\nabla^{[\mu} l^{\nu]\rho}(0, \sigma^r) = 0, \quad \nabla_{\mu} l^{\mu\rho}(0, \sigma^r) = 0. \quad (23)$$

Using the results [15] we prove that the constraints (23) are always fulfilled during the evolution prescribed by EOM (21). Thereat we take into account vanishing of the space covariant derivatives of the IDC (23)

$$\nabla_r \nabla_{\mu} l^{\mu\rho}|_{\tau=0} = 0, \quad \nabla_r \nabla^{[\mu} l^{\nu]\rho}|_{\tau=0} = 0, \quad (r = 1, 2, \dots, p). \quad (24)$$

Consider the power series expansion of  $\nabla^{[\mu} l^{\nu]\rho}$  and  $\chi^{\rho} \equiv \nabla_{\mu} l^{\mu\rho}$

$$\chi^{\rho}(\delta\tau, \sigma^r) = \chi^{\rho}|_{\tau=0} + \partial_{\tau} \chi^{\rho}|_{\tau=0} \delta\tau + \dots = \nabla_{\tau} \chi^{\rho}|_{\tau=0} \delta\tau + \dots, \quad (25)$$

$$\nabla^{[\tau} l^{\nu]\rho}(\delta\tau, \sigma^r) = \partial_{\tau} \nabla^{[\tau} l^{\nu]\rho}|_{\tau=0} \delta\tau + \dots = \nabla_{\tau} \nabla^{[\tau} l^{\nu]\rho}|_{\tau=0} \delta\tau + \dots \quad (26)$$

The use of Eqs. (19)  $\nabla_{\tau} \chi^{\tau}(\xi) = -\nabla_r \chi^r(\xi)$  and (24)  $\nabla_r \chi^r|_{\tau=0} = 0$  gives

$$\chi^{\tau}(\delta\tau, \sigma^r) = \chi^{\tau}|_{\tau=0} = 0. \quad (27)$$

To prove that  $\chi^r(\delta\tau, \sigma^r) = 0$  we consider EOM (21) for  $\nu = \tau$  and  $\rho = r$

$$\nabla_s \nabla^{[s} l^{\tau]r} + \frac{1}{2} \nabla^{\tau} \chi^r = 0, \quad (28)$$

and taking into account (24), (27) find

$$\nabla_{\tau} \chi^r|_{\tau=0} = \partial_{\tau} \chi^r|_{\tau=0} = 0 \longrightarrow \chi^r(\delta\tau, \sigma^r) = \chi^r|_{\tau=0} = 0. \quad (29)$$

Eqs.  $\partial_{\tau} \chi^{\rho}|_{\tau=0} = 0$  and (27) complete the proof of the desired statement

$$\nabla_{\mu} l^{\mu\rho}(\xi) = 0. \quad (30)$$

Eqs. (30) prove equivalence of EOM (21) to the system

$$\nabla_{\mu} \nabla^{[\mu} l^{\nu]\rho} = 0, \quad \nabla_{\mu} l^{\mu\rho}(\xi) = 0 \quad (31)$$

showing the local conservation of the P-C conditions (10).

Next we prove that the IDC (23) together with the local conservation laws (31) provide fulfillment of the P-C conditions (10) at any  $\tau$

$$\nabla^{[\mu\nu]\rho}(\xi) = 0. \quad (32)$$

To this objective consider  $\nu = r$  and  $\rho = \tau$  in (31) and obtain the relations

$$\nabla_\tau \nabla^{[\tau l^r]\tau} = -\nabla_s \nabla^{[s l^r]\tau} \quad (33)$$

which in combination with the IDC (24) show that

$$\nabla_\tau \nabla^{[\tau l^r]\tau}|_{\tau=0} = \partial_\tau \nabla^{[\tau l^r]\tau}|_{\tau=0} = 0. \quad (34)$$

The substitution of (34) into the expansion (26) proves that

$$\nabla^{[\tau l^r]\tau}(\xi) = 0. \quad (35)$$

The choice of both indices  $\nu, \rho$  in (31) as space ones  $\nu = r, \rho = s$  gives

$$\nabla_\tau \nabla^{[\tau l^r]s} = -\nabla_q \nabla^{[q l^r]s}. \quad (36)$$

Eqs. (36) and (24) show that  $\nabla_\tau \nabla^{[\tau l^r]s}|_{\tau=0} = \partial_\tau \nabla^{[\tau l^r]s}|_{\tau=0} = 0$  that provides

$$\nabla^{[\tau l^r]s}(\xi) = 0. \quad (37)$$

Eqs. (37) and (35) testify to fulfillment of the subset

$$\nabla^{[\tau l^r]\rho}(\xi) = 0 \quad (38)$$

of the P-C equations. Antisymmetrization in the  $\rho, r$  indices in (38) shows fulfillment of the next subset of Eqs. (10)

$$\nabla^{[r l^s]\tau}(\xi) = 0. \quad (39)$$

The remaining subset of the P-C eqs. to be proved

$$\nabla^{[r l^s]q}(\xi) = 0 \quad (40)$$

contains only space-like indices  $r, s, q$ . Therefore, we consider the expansion

$$\nabla^{[r l^s]q}(\delta\tau, \sigma^r) = \partial_\tau \nabla^{[r l^s]q}|_{\tau=0} \delta\tau + \dots = g_{\tau\tau} \nabla^\tau \nabla^{[r l^s]q}|_{\tau=0} \delta\tau + \dots, \quad (41)$$

where the relation  $\nabla_\tau \nabla^{[r]l^s]q}|_{\tau=0} = g_{\tau\tau} \nabla^\tau \nabla^{[r]l^s]q}|_{\tau=0}$  together with (23), (24) are used. The proof of (40) reduces to the that of the relation

$$\nabla^\tau \nabla^{[r]l^s]q}|_{\tau=0} \equiv ([\nabla^\tau, \nabla^{[r]l^s]q} + \nabla^{[r]l^s]q} \nabla^\tau]|_{\tau=0} = 0. \quad (42)$$

Using the BI (12) we present the first term in the r.h.s. of (42) as

$$[\nabla^\tau, \nabla^{[r]l^s]q} = R^{\tau[r]s]_\mu} l^{\mu q} + R^{\tau[r]q]_\mu} l^{s] \mu}. \quad (43)$$

The conditions (37) permit to rewrite the second term in the form

$$\begin{aligned} \nabla^{[r]l^s]q} \nabla^\tau = \nabla^{[r]l^s]q} \nabla^\tau &= [\nabla^{[r]l^s]q}, \nabla^\tau] l^s]^\tau + \nabla^q \nabla^{[r]l^s]q} \tau = \\ &= R^{[r]q\tau]_\mu} l^{\mu s]^\tau} + R^{[r]q]^\tau_\mu} l^{s] \tau \mu} + \nabla^q \nabla^{[r]l^s]q} \tau. \end{aligned} \quad (44)$$

The sum of (43) and (44) transformss (42) into

$$\nabla^\tau \nabla^{[r]l^s]q}|_{\tau=0} = (R^{\tau[r]s]_\mu} l^{\mu q} + R^{\tau[r]q]_\mu} l^{s] \mu} + R^{[r]q\tau]_\mu} l^{\mu s]^\tau} + R^{[r]q]^\tau_\mu} l^{s] \tau \mu})|_{\tau=0} \quad (45)$$

after using (38) and (24). Further, we apply the Gauss theorem (6)

$$R^{\gamma\lambda\nu}{}_\mu l^{\mu r} = (l^2)^{r[\gamma} l^{\lambda]\nu} \quad (46)$$

and obtain the following expressions for the terms in (45)

$$R^{\tau[r]s]_\mu} l^{\mu q} = -(l^2)^{q[r} l^{s]\tau}, \quad R^{[r]q]^\tau_\mu} l^{s] \mu} = (l^2)^{\tau[r} l^{s]q}, \quad (47)$$

$$R^{[r]q\tau]_\mu} l^{\mu s]^\tau} + R^{[r]q]^\tau_\mu} l^{s] \tau \mu} = (l^2)^{\tau[r} l^{s]q} - (l^2)^{q[s} l^{r]\tau}. \quad (48)$$

Here we observe mutual cancellation of the corresponding terms in the sum of (47) and (48). This proves Eqs. (40) and shows validity of Eqs.(32) . Then the cmc condition  $Sp l = \mu$  emerges as a consequence of Eqs. (30) and (32). So, we see that EOM (17) together with the Gauss eqs. (6) provide conservation of the IDC (30) and (32). The latter select a *closed* sector of the solutions descrbing cmc h-ws of codim 1.

## 5 Extremals of the brane potential

The Gauss's theorem permits to express  $V_{cmc}$  (15) associated with the h-ws  $\Sigma_{p+1}^{cmc}$  in terms of the Riemann tensor. Actually, using Eqs. (6) we obtain

$$R_{\nu\lambda} = (l^2)_{\nu\lambda} - \mu l_{\nu\lambda}, \quad R_{\nu\lambda} l^{\lambda\nu} = Sp(l^3) - \mu Sp(l^2), \quad R = Sp(l^2) - \mu^2 \quad (49)$$

where  $R_{\nu\rho}$  is the Ricci tensor. Using (49) we obtain the  $R$ -reps. for  $V_{cmc}$

$$V_{cmc} = -\frac{1}{2}R^2 - \frac{\mu^2}{3}R + \frac{2\mu}{3}R_{\nu\lambda}l^{\lambda\nu} + \frac{\mu^4}{6} - c_p \quad (50)$$

which reduces to  $V_{Dir}$  when  $\mu = 0$ . Reprs. (50) can be rewritten as

$$V_{cmc} = -\frac{\mu^2}{3}\left\{(R - 2\Lambda) + \frac{3}{2\mu^2}R^2 - \frac{2}{\mu}R_{\nu\lambda}l^{\nu\lambda}\right\}, \quad (51)$$

where  $\Lambda(p) := \left(\frac{\mu^2(p)}{4} - \frac{3c_p}{\mu^2(p)}\right)$  is the h-ws cosmological constant. We see emergence of the H-E term together with the interaction term

$$V_{cmc} = V_{Dir} - \frac{\mu^2}{3}\left(R - \frac{\mu^2}{2} - \frac{2}{\mu}R_{\nu\lambda}l^{\lambda\nu}\right), \quad \Lambda|_{c_p=0} = \frac{1}{4}\mu^2(p) \quad (52)$$

which extend the pure  $R^2$  gravity encoded by  $V_{Dir}$ . The potential  $V_{cmc}$  reduces to the constant  $\frac{2}{3}\mu^2\Lambda(p)$  when  $R_{\nu\rho} = 0$ . Then (49) yields the relations

$$(l^n)_{\nu\rho} = \mu^{n-1}l_{\nu\rho}, \quad (n = 2, 3, \dots) \implies l_{\nu\rho} = \mu g_{\nu\rho}, \quad \det l_\nu^\alpha \neq 0 \quad (53)$$

and  $Spl = (p+1)\mu$ , respectively. The latter is compatible with the cmc condition only if  $p = 0$  that describes a worldline. So, the case when  $R_{\nu\rho} = 0$ , but  $\mu \neq 0$ ,  $\det l_{\nu\rho} \neq 0$  describes point-like 0-branes.

Further we study extremals of  $V_{cmc}$  defined by solutions of the equation

$$\frac{1}{2} \frac{\partial V_{cmc}}{\partial l_{\nu\rho}} = \mu(l^2)^{\nu\rho} - l^{\nu\rho} Sp(l^2) = 0. \quad (54)$$

In the  $R$ -reps. this equation takes the form

$$Rl_{\nu\rho}(\xi) = \mu R_{\nu\rho} \quad (55)$$

after using relations (49). The general solution of (55) is

$$l_{\nu\rho}(\xi) = \mu \frac{R_{\nu\rho}}{R}, \quad R \neq 0. \quad (56)$$

The substitution of this extremal in the Gauss equation yields the relation

$$R_{\mu\nu\gamma\lambda} = -\left(\frac{\mu}{R}\right)^2 (R_{\mu\gamma}R_{\nu\lambda} - R_{\nu\gamma}R_{\mu\lambda}). \quad (57)$$

The extremal (56) should obey the P-C eqs. (10)

$$\nabla_{[\mu} R_{\nu]\rho} - \frac{1}{R} R_{\rho[\nu} \nabla_{\mu]} R = 0. \quad (58)$$

and Eq. (22) which follows from (10) and has the form

$$\nabla_{\nu} R^{\nu\rho} - \frac{1}{R} R^{\rho\nu} \partial_{\nu} R = 0 \quad (59)$$

after substitution of (56) in (22). The use of the relation

$$\nabla_{\nu} R^{\nu}_{\rho} = \frac{1}{2} \partial_{\rho} R, \quad (60)$$

which follows from the Bianchi identities

$$\nabla_{\alpha} R^{\mu}_{\nu\gamma\lambda} + \nabla_{\gamma} R^{\mu}_{\nu\lambda\alpha} + \nabla_{\lambda} R^{\mu}_{\nu\alpha\gamma} = 0, \quad (61)$$

transforms Eq. (59) into the equation

$$(R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R) \partial_{\nu} R = 0. \quad (62)$$

The latter means that  $\partial_{\mu} R$  should be a null vector for the Einstein matrix  $G^{\mu\nu} := R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R$  associated with  $\Sigma_{p+1}^{extr}$ . In the  $R$ -reps. this h-ws is fixed by Eqs. (57), (62). Their solution for the case when  $\det G^{\mu\nu} \neq 0$  is

$$\partial_{\nu} R = 0 \implies R = R_o = \text{constant}. \quad (63)$$

This solution defines a h-ws  $\Sigma_{p+1}^o$  of *constant curvature* (cc) characterized by the Riemann tensor and the Gauss curvature  $K_o$

$$R_{o\mu\nu\gamma\lambda} = K_o (g_{\mu\gamma} g_{\nu\lambda} - g_{\nu\gamma} g_{\mu\lambda}). \quad (64)$$

The substitution of (64) in the Gauss eqs. (57) results in the relation

$$K_o (g_{\mu\gamma} g_{\nu\lambda} - g_{\nu\gamma} g_{\mu\lambda}) = -\left(\frac{\mu}{R}\right)^2 (R_{\mu\gamma} R_{\nu\lambda} - R_{\nu\gamma} R_{\mu\lambda}) \quad (65)$$

which shows proportionality of  $R_{o\mu\nu}$  to  $g_{\mu\nu}$  and defines  $K_o$

$$R_{o\mu\nu} = \Lambda_o g_{\mu\nu}, \quad R_o = (p+1)\Lambda, \quad K_o = -\left(\frac{\mu}{p+1}\right)^2. \quad (66)$$

The latter completely fixes the Riemann tensor of  $\Sigma_{p+1}^o$

$$R_{o\mu\nu\gamma\lambda} = -\left(\frac{\mu}{p+1}\right)^2 (g_{\mu\gamma}g_{\nu\lambda} - g_{\nu\gamma}g_{\mu\lambda}) \quad (67)$$

and the function  $\Lambda_o = pK_o$ , respectively. Then we obtain

$$R_{o\mu\nu} = -\frac{p}{(p+1)^2}\mu^2 g_{\mu\nu}, \quad R_o := g^{\mu\nu}R_{o\mu\nu} = -\frac{p}{p+1}\mu^2. \quad (68)$$

As shows the substitution of (68) in (56), this extremal acquires the form

$$l_{o\mu\nu} = \frac{\mu}{p+1}g_{\mu\nu} \implies R_{o\mu\nu} = -\frac{p}{p+1}\mu l_{o\mu\nu}. \quad (69)$$

Solutions (69) are proportional to  $g_{\mu\nu}(\xi)$ , and therefore they are solutions of the P-C eqs. (10) in the  $l$ -reps. and eqs. (58) in the  $R$ -reps., respectively.

Coming back to Eq. (54) in the  $l$ -reps. we rewrite it in the form

$$l_\alpha^\nu (\mu l^{\alpha\rho} - g^{\alpha\rho} Sp(l^2)) = 0$$

which shows that there exist two sets of the solutions. For the first of them  $detl_{\mu\nu} = 0$ , for the other it is nonzero. In the latter case (54) reduces to

$$l^{\alpha\rho} = \frac{Sp(l^2)}{\mu} g^{\alpha\rho}, \quad detl_{\mu\nu} \neq 0 \quad (70)$$

and corresponds to the above-considered solution (69). Indeed, in this case contraction of (70) with  $l^{\rho\alpha}$  and  $g^{\rho\alpha}$ , respectively, yields the conditions

$$Sp l_o = \mu, \quad Sp(l_o^2) = \frac{\mu}{p+1} Sp l_o = \frac{\mu^2}{p+1}. \quad (71)$$

Then the substitution of (71) in (70) reproduces the solution (69) for  $l_{o\mu\nu}$ . The substitution of  $l_{o\mu\nu}$  in the Gauss eqs. (6) yields the solution (67) for the Riemann tensor of the extremal h-ws  $\Sigma_{p+1}^o$ . The cc h-ws  $\Sigma_{p+1}^o$  is a hyperbolic one with the negative *constant curvature*  $K_o$ . The use of the relation

$$R_{o\mu\nu} = \frac{R_o}{p+1} g_{\mu\nu}, \quad \Lambda_o = -\frac{p}{(p+1)^2} \mu^2$$

shows that  $\Sigma_{p+1}^o$  is a  $(p+1)$ -dim. Einstein space with the negative cosmological constant  $\Lambda_o$  which can be treated as the anti-de Sitter space  $AdS_{p+1}$ .

The vacuum value of  $V_{cmc}$  (15) is the constant given by

$$V_o := V_{cmc}(l = l_o, g, \mu) = \frac{\mu^4(p)}{6(p+1)^2} - c_p. \quad (72)$$

The action (16) at the extremum  $l_o(\xi)$  is also constant equal to

$$S_o := S_{cmc}|_{vac} = \frac{1}{k_p^2} \int d^{p+1}\xi \sqrt{|g|} \left( \frac{\mu^4(p)}{6(p+1)^2} - c_p \right), \quad (73)$$

due to vanishing of the kinetic term of  $l_o(\xi)$  because  $l_{o\mu\nu} \sim g_{\mu\nu}$ .

The differential of Eq. (14) yields the second partial derivative of  $V_{cmc}$

$$\begin{aligned} \frac{\partial^2 V_{cmc}}{\partial l_{\alpha\beta} \partial l_{\nu\rho}} &= [g^{\beta\{\nu} l^{\rho\}\alpha} + g^{\alpha\{\nu} l^{\rho\}\beta}] S_{pl} - g^{\alpha\{\nu} g^{\rho\}\beta} S_p(l^2) \\ &+ [(l^2)^{\nu\rho} g^{\alpha\beta} + (l^2)^{\alpha\beta} g^{\nu\rho}] - 4l^{\nu\rho} l^{\alpha\beta}. \end{aligned} \quad (74)$$

The substitution of  $l_{\mu\nu} = l_{o\mu\nu}$  (69) in Eq. (74) gives its value

$$\frac{\partial^2 V_{cmc}}{\partial l_{\alpha\beta} \partial l_{\nu\rho}}|_{vac} = \left( \frac{\mu}{p+1} \right)^2 [(p+1)g^{\beta\{\nu} g^{\rho\}\alpha} - 2g^{\nu\rho} g^{\alpha\beta}] \quad (75)$$

at the critical point  $l_{o\mu\nu}(\xi)$ . This value is not positively defined that hints on possible classical unstability of the vacuum. The latter is easily checked for strings and mebranes where the diagonal gauge for  $g_{\mu\nu}$  can be chosen. For string ws we obtain two non-zero components  $l_{\tau\tau} \equiv l_\tau, l_{\sigma\sigma} \equiv l_\sigma$  that gives

$$\frac{\partial^2 V_{cmc}}{\partial l_\tau \partial l_\tau}|_{vac} = \frac{(\mu g^\tau)^2}{2}, \quad \frac{\partial^2 V_{cmc}}{\partial l_\sigma \partial l_\sigma}|_{vac} = \frac{(\mu g^\sigma)^2}{2}, \quad \frac{\partial^2 V_{cmc}}{\partial l_\tau \partial l_\sigma}|_{vac} = -\left(\frac{\mu}{2}\right)^2 g^\tau g^\sigma. \quad (76)$$

Since  $g^\tau g^\sigma < 0$ , we find the extremum to be the *minimum* defining classically stable *string* vacuum. For membrane  $g_{\tau r} = g_{\sigma \eta} = 0$ , where  $\sigma^1 \equiv \sigma$  and  $\sigma^2 \equiv \eta$ . Then we obtain three non-zero diagonal components:  $l_{\tau\tau} \equiv l_\tau, l_{\sigma\sigma} \equiv l_\sigma, l_{\eta\eta} \equiv l_\eta$ , and Eqs. (76) take the form

$$\frac{\partial^2 V_{cmc}}{\partial l_\tau \partial l_\tau}|_{vac} = \frac{(2\mu g^\tau)^2}{2}, \quad \frac{\partial^2 V_{cmc}}{\partial l_\sigma \partial l_\sigma}|_{vac} = \frac{(2\mu g^\sigma)^2}{2}, \quad \frac{\partial^2 V_{cmc}}{\partial l_\eta \partial l_\eta}|_{vac} = \frac{(2\mu g^\eta)^2}{2} \quad (77)$$

with all *positive* derivatives. The same we see for the mixed derivatives

$$\frac{\partial^2 V_{cmc}}{\partial l_\tau \partial l_\sigma}|_{vac} = -2\left(\frac{\mu}{3}\right)^2 g^\tau g^\sigma, \quad \frac{\partial^2 V_{cmc}}{\partial l_\tau \partial l_\eta}|_{vac} = -2\left(\frac{\mu}{3}\right)^2 g^\tau g^\eta, \quad (78)$$

since  $g^\tau g^\sigma < 0$  and  $g^\tau g^\eta < 0$  due to the time-like character of  $\Sigma_3^g$ .

However, the mixed derivative with respect to  $l_\sigma$  and  $l_\eta$  is *negative*

$$\frac{\partial^2 V_{cmc}}{\partial l_\sigma \partial l_\eta} \Big|_{vac} = -2 \left(\frac{\mu}{3}\right)^2 g^\sigma g^\eta, \quad (79)$$

since  $g^\sigma g^\eta > 0$ . It means that the extremal of the potential  $V_{cmc}(l, g, \mu)$  is a 3-dim. *saddle* hypersurface that shows classical *unstability* of the membrane vacuum. So, the potential of cmc 3-brane with  $R_{\mu\nu} \neq 0$  in 4-dim. Minkowski space turns out to be unstable. This result based on the sign indefiniteness of the second derivatives in (75) hints that extremals of  $V_{cmc}(l, g, \mu)$  for branes with codim 1 and higher  $p$  might also be *saddle* hypersurfaces.

## 6 Conclusion

We considered the extension of the gauge theory approach [15] to branes sweeping hyper-worldsheets of codimension 1 with *constant mean curvature*. Constructed were invariant action and interaction potential for the h-ws Nambu-Goldstone bosons  $l_{\mu\nu}$  of the broken Poincare symmetry. The potential was shown to encode the generalized h-ws action of quadratic curvature gravity. The equation for the brane *potential* extremals and its solution describing h-ws of negative *constant curvature*, were revealed. This set of maximally symmetric spaces also includes anti-de Sitter space  $AdS_{p+1}$  which is the exact solution of the H-E equations for an empty universe with negative cosmological constant. The found extremum for  $p = 2$  was proved to be realized by a *saddle* 3-dim. hypersurface. This points to classical unstability of the constant mean curvature *membrane* potential in  $\mathbf{R}^{1,3}$ . It is interesting to connect the saddle-like unstability with the inflation and reheating mechanisms studied in quadratic curvature models of gravity.

### Acknowledgments

The author would like to thank NORDITA and Physics Department of Stockholm University for kind hospitality and support, I. Bengtsson, S. Lukyanov for stimulating discussions and G. Huicken for his interest to the work and valuable remarks.

## References

- [1] E. Cartan, Riemannian Geometry in an Orthogonal Frame (World Scientific, Singapore, 2001).
- [2] L.F. Eisenhart, Riemannian Geometry (Princeton University Press, Princeton, 1968).
- [3] James Simons, Minimal varieties in riemannian manifolds, *Annals of Mathematics*, 88 (1968) 62-105.
- [4] R. Schoen, L. Simon and S.T. Yau, Curvature estimates for minimal surfaces, *Acta Mathematica*, 134 (1975) 255-288.
- [5] J. Hoppe and H. Nicolai, Relativistic minimal surfaces, *Phys. Lett. B* 196 (1987) 451-455.
- [6] F. Lund and T. Regge, Unified approach to strings and vortices with soliton solutions, *Phys. Rev. D* 14 (1976) 1524-1535.
- [7] R. Omnes, A new geometric approach to the relativistic string, *Nucl. Phys. B* 149 (1979) 269-284.
- [8] B.M. Barbashov and V.V. Nesterenko, Introduction to the Relativistic String Theory (World Scientific Pub Co Inc, Singapore, 1990).
- [9] A.A. Zheltukhin, Classical relativistic string as an exactly solvable sector of  $SO(1,1) \times SO(2)$  gauge model, *Phys. Lett. B* 116 (1982) 147-150; Gauge description and nonlinear string equations in d-dimensional space-time, *Theor. Math. Phys.* 56 (1983) 785-795.
- [10] J. Gomis, K. Kamimura and P. West, The construction of brane and superbrane actions using non-linear realizations, *Class. Quant. Grav.* 23 (2006) 7369-7381.
- [11] F. Gliozzi, M. Meineri, Lorentz completion of effective string (and p-brane) action, *JHEP* 1208 (2012) 056.
- [12] J. Gomis, K. Kamimura and J. M. Pons, Non-linear realizations, Goldstone bosons of broken Lorentz rotations and effective actions for p-branes, *Nucl. Phys. B* 871 (2013) 420-451.

- [13] I.A. Bandos and A.A. Zheltukhin, Null super p-brane quantum theory in 4-dimensional space-time, *Fortschr. Phys.* 4 (1993) 619-676.
- [14] M. P. Bogers and T. Brauner, Lie-algebraic classification of effective theories with enhanced soft limits, *JHEP* 1805 (2018) 076.
- [15] A.A. Zheltukhin, Gauge theory approach to branes and spontaneous symmetry breaking, *Rev. Math. Phys.* 29, No. 3 (2017) 1750009.
- [16] A.A. Zheltukhin, Inflation versus collapse in brane matter, *Modern Phys. Lett. A* 32, No.36 (2017) 1750199(1-9).
- [17] K. S. Stelle, Abdus Salam and quadratic curvature gravity: Classical solutions, *Int.J.Mod.Phys. A*32 (2017) no.09, 1741012.
- [18] Starobinsky, A. A. , A new type of isotropic cosmological models without singularity, *Phys. Lett. B* 91 (1980) 99102.
- [19] E. S. Fradkin and A. A. Tseytlin, Renormalizable asymptotically free quantum theory of gravity, *Phys. Lett. B* 104 (1981) 377.
- [20] B. Zwiebach, Curvature squared terms and string theories, *Phys. Lett. B* 156 (1985) 315.
- [21] H. Lü, A. Perkins, C.N. Pope and K.S. Stelle, Spherically Symmetric Solutions in Higher-Derivative Gravity, *Phys. Rev. D* 92 (2015) 124019.
- [22] M. Trzetrzelewski and A.A. Zheltukhin, Exact solutions for U(1) globally invariant membranes, *Phys. Lett. B* 679 (2009) 523-528.
- [23] A.A. Zheltukhin, Toroidal p-branes, anharmonic oscillators and (hyper)elliptic solutions, *Nucl. Phys. B* 858 (2012) 142-154; Laplace-Beltrami operator and exact solutions for branes, *Nucl. Phys. B* 867 (2013) 763-778.
- [24] P.A.M. Dirac, Long range forces and broken symmetries, *Proc. R. Soc. Lond., A* 333 (1973) 403.
- [25] V. De Alfaro, S. Fubini and G. Furlan, *Nuovo Cimento* 50 A (1979) 523.
- [26] L. Smolin, Towards a Theory of Space-Time Structure at Very Short Distances, *Nucl. Phys. B* 160 (1979) 253-258.

- [27] A. Zee, Spontaneously generated gravity, *Phys. Rev. D* 23 (1980) 858-866.
- [28] H. Johansson and J. Nohle, Conformal Gravity from Gauge Theory, arXiv:1707.02965[hep-th].
- [29] V. Pervushin and A. Pavlov, Principles of Quantum Universe, LAP LAMBERT Academic Publishing, Saarbrucken, 2013.
- [30] C.G. Callan, C. Lovelace, C.R. Nappi and S.A. Yost, String loop corrections to beta functions, *Nucl. Phys. B* 288 (1987) 525-550.
- [31] B.M. Barbashov, V.V Nesterenko and A.M. Chervjakov, Generalization of the relativistic string model in the geometrical approach, *Lett. Math. Phys.* 3 (1979) 359-365.
- [32] A.A. Zheltukhin, Tension as a perturbative parameter in nonlinear string equations in curved space-time, *Class.Quant.Grav.* 13 (1996) 2357-23604; S.N. Roshchupkin and A.A. Zheltukhin, Variational principle and a perturbative solution of nonlinear string equations in curved space, *Nucl. Phys. B* 543 (1999) 365-386.
- [33] M. Rinaldi and L. Vanzo, Inflation and reheating in theories with spontaneous scale invariance symmetry breaking, *Phys. Rev. D* 94 (2016) no.2, 024009; G. Tambalo and M. Rinaldy, Inflation and reheating in scale-invariant scalar-tensor gravity, *Gen. Rel. Grav.* 49 (2017) no.4, 52.
- [34] V. A. Rubakov and M. E. Shaposhnikov, Do we live inside a domain wall?, *Phys. Lett. B* 125 (1983) 136-138.
- [35] S.N. Roshchupkin and A.A. Zheltukhin, On a possibility of membrane cosmology, e-Print: hep-th/9607119; On an approximate description of string and p-brane dynamics in curved space-time, *Theor. Math. Phys.* 111 (1997) 714-722.
- [36] B. de Witt, M. Lusher and G. Nicolai, The supermembrane is unstable, *Nucl. Phys. B* 320 (1989) 135-159.
- [37] J. Goldstone, unpublished: J. Hoppe, MIT-PhD Thesis, 1982: in *Proc. Int. Workshop on Constraints Theory and Relativistic Dynamics*, eds. G. Longhi and L. Lusanna (World Scientific, Singapore, 1987).