

The multistationarity structure of networks with intermediates and a binomial core network

AmirHosein Sadeghimanesh¹, Elisenda Feliu^{1,2}

December 8, 2021


Abstract

This work addresses whether a reaction network, taken with mass-action kinetics, is multistationary, that is, admits more than one positive steady state in some stoichiometric compatibility class. We build on previous work on the effect that removing or adding intermediates has on multistationarity, and also on methods to detect multistationarity for networks with a binomial steady state ideal. In particular, we provide a new determinant criterion to decide whether a network is multistationary, which applies when the network obtained by removing intermediates has a binomial steady state ideal. We apply this method to easily characterize which subsets of complexes are responsible for multistationarity; this is what we call the *multistationarity structure* of the network. We use our approach to compute the multistationarity structure of the n -site sequential distributive phosphorylation cycle for arbitrary n .

Keywords: binomial ideal, phosphorylation cycle, multistationarity, model reduction, determinant criterion, toric

Introduction

Given a reaction network, an *intermediate* is a species that does not interact with any other species, is produced by at least one reaction, and consumed by at least one reaction. Typical intermediates Y arise in Michaelis-Menten type mechanisms as



where c, c' are arbitrary complexes. Removal of the intermediates of a network yields a new network, called the *core network*, as introduced in [8] (and further generalized in [15]). For example, removal of Y from the mechanism above gives the reaction $c \rightarrow c'$.

We consider mass-action kinetics, such that the system of ordinary differential equations (ODEs) modeling the evolution of the concentrations of the species in time is polynomial. As shown in [8, Theorem 5.1], multistationarity of the core network implies multistationarity of the original (extended) network, provided a technical realization condition is satisfied. Further, whether an extended network is multistationary depends only on the set of complexes of the core network that react to an intermediate. These complexes are called *inputs*. The subsets of complexes that give rise to multistationarity define the *multistationarity structure* of the core network.

In this work we present an approach to find the multistationarity structure of core networks, and thereby provide a fast way to decide whether a given extension network

¹Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark

²Corresponding author: efeliu@math.ku.dk

is multistationary or not; indeed, it suffices to find the set of inputs of our network and check whether it belongs to the multistationarity structure.

The method applies to core networks that are *binomial* (the ideal generated by the steady state polynomials is binomial). For these networks, a method to decide on multistationarity was introduced in [13], based on the computation of sign vectors. Under some extra assumptions, another method relying on the computation of a symbolic determinant and inspection of the sign of its coefficients is presented in [12] (see Theorem 2.7).

The first main result of this paper is Theorem 3.11, where we combine the results in [8, 12] into a new determinant criterion for multistationarity that applies to extended networks for which the core network is binomial, even though the original network might not be binomial.

The second main result is Theorem 4.2, which removes the technical realization condition in [8] for concluding that an extended network is multistationary provided the core network is. Instead, we require that both the core and the extended networks are binomial (in a compatible way).

The third main contribution is Algorithm 4.8, which returns the multistationarity structure of a binomial core network based on the determinant criterion. The algorithm relies on the study of the signs of a polynomial obtained after the computation of the determinant of a symbolic matrix. Our approach is more direct than testing if the extended network is multistationary for all subsets of input complexes. We apply our method to find the multistationarity structure of the n -site distributive sequential phosphorylation cycle for arbitrary n in §4.3. This illustrates how our results allow us to study a family of networks at once.

We conclude with an investigation of when the technical realization conditions in [8] are satisfied. In particular, we show that the conditions hold for typical types of intermediates like the Michaelis-Menten mechanism above.

The structure of the paper is as follows. In §1 we introduce basic concepts on reaction networks and multistationarity. In §2 we introduce (complete) binomial networks and the determinant criterion for determining multistationarity. In §3 we focus on intermediates and give the determinant criterion for multistationarity applicable to extended networks with a binomial core network. In §4 we link multistationarity of the core and extended networks, and in particular study the multistationary structure. Finally, in §5 we expand on how to check the realization conditions.

Notation. Subscripts $\geq 0, > 0$ for \mathbb{R} refer to the non-negative and positive real numbers. The sets $\{1, \dots, m\}$ and $\{m_1, \dots, m_2\}$ are respectively denoted by $[m]$ and $[m_1, m_2]$. In particular $[m] = [1, m]$. The cardinality of a set A is denoted by $|A|$.

Consider two vectors $u, v \in \mathbb{R}^n$. The scalar product of u and v is denoted by $u \cdot v$. The vector v^u is defined as $\prod_{j=1}^n v_j^{u_j}$, and for a matrix $M \in \mathbb{R}^{n \times m}$ with column vectors $u^{(i)}$, $i \in [m]$, the vector v^M is the vector whose i -th entry is $v^{u^{(i)}}$. We let $\text{diag}(v)$ be the diagonal matrix with diagonal v and

$$M_v = M \text{diag}(v).$$

The sign vector of v , $\sigma(v) \in \{-1, 0, 1\}^n$, is defined for $i \in [n]$ as

$$\sigma(v)_i = \begin{cases} 1 & \text{if } v_i > 0 \\ 0 & \text{if } v_i = 0 \\ -1 & \text{if } v_i < 0. \end{cases}$$

1 Reaction networks

In this section we briefly introduce the ingredients from chemical reaction network theory needed in the sequel. See for example [5, 10]. A reaction network, also called a *network*, is a triplet of finite sets $\mathcal{N} = (\mathcal{S}, \mathcal{C}, \mathcal{R})$. The three sets are called respectively the set of *species*, *complexes* and *reactions*. The elements of \mathcal{C} are finite linear combinations of the species with non-negative integer coefficients. The set of reactions consists of ordered pairs (c, c') of complexes, denoted $c \rightarrow c'$.

After fixing an order on \mathcal{S} , write $\mathcal{S} = \{X_1, \dots, X_n\}$. We identify a complex $c \in \mathcal{C}$ with the vector in \mathbb{R}^n whose i -th entry is the coefficient of X_i in c . Therefore a complex c is either given as $\sum_{X \in \mathcal{S}} c_X X$ or by the corresponding vector (again denoted c).

There is a natural digraph associated with a network, with vertex set \mathcal{C} and edge set \mathcal{R} . We often identify the reaction network with the digraph for simplicity. The stoichiometric matrix N is an $|\mathcal{S}| \times |\mathcal{R}|$ matrix whose column vectors are $c' - c$ for each reaction $c \rightarrow c' \in \mathcal{R}$. This matrix depends on a fixed order of the set of reactions. The (real) column space of N is called the *stoichiometric subspace* and is denoted by S . Its dimension, that is, the rank of N , is the *rank* of the network.

In this work we consider so-called *mass-action kinetics*. Under this assumption, the evolution of the concentration of the species in time is modeled by means of a polynomial ODE system as follows. First, a positive real number $k_{c \rightarrow c'}$ is assigned to each reaction $c \rightarrow c'$. This number is called the *reaction rate constant* and often written as a label of the reaction in the associated digraph. We interchangeably write $k_i = k_{c \rightarrow c'}$ if $c \rightarrow c'$ is the i -th reaction and write the vector of reaction rate constants $k \in \mathbb{R}_{>0}^{|\mathcal{R}|}$, if the order of \mathcal{R} is relevant, and $k \in \mathbb{R}_{>0}^{\mathcal{R}}$ otherwise.

Next, we let $x = (x_1, \dots, x_n)$ denote the vector of the concentrations of X_1, \dots, X_n ; note that in examples we simply use corresponding lower-case letters to denote concentrations. Given $x \in \mathbb{R}^n$, we define the vector $\psi(x) \in \mathbb{R}^{|\mathcal{R}|}$ as

$$\psi(x)_i = x^c, \quad \text{if } c \rightarrow c' \text{ is the } i\text{-th reaction.}$$

Now the ODE system associated with the network and $k \in \mathbb{R}_{>0}^{|\mathcal{R}|}$ is

$$\frac{dx}{dt} = F_k(x), \quad \text{where } F_k(x) = N_k \psi(x), \quad x \in \mathbb{R}_{>0}^n. \quad (1)$$

Recall that $N_k = N \text{diag}(k)$.

The solution to (1) with an initial condition $x_0 \in \mathbb{R}_{>0}^n$ is confined to the *stoichiometric compatibility class* of x_0 : $(x_0 + S) \cap \mathbb{R}_{>0}^n$ [6, Remark 3.4]. Equations for these classes are found as follows. Let d be the corank of the network, that is, $d = n - \text{rank}(N)$. A matrix $Z \in \mathbb{R}^{d \times n}$ whose rows form a basis of the orthogonal complement S^\perp of S is called a *matrix of conservation laws*. Then the set $(x_0 + S) \cap \mathbb{R}_{>0}^n$ agrees with the set

$$\{u \in \mathbb{R}_{>0}^n \mid Zu = Zx_0\}.$$

A positive *steady state* is a solution to the system $F_k(x) = 0$ in $\mathbb{R}_{>0}^n$. Since we would like to treat the values of k as unknown, we view the polynomials $F_{k,i}(x)$ as polynomials in the ring $\mathbb{R}(k)[x]$ by regarding k as parameters instead of positive real numbers. When we do this, we write $F_i(x)$. Then $F_1(x), \dots, F_n(x)$ are called the *steady state polynomials* of the network, and the ideal I they generate is the *steady state ideal*:

$$I = \langle F_1(x), \dots, F_n(x) \rangle \subseteq \mathbb{R}(k)[x].$$

Throughout this work, given an element or subset B of $\mathbb{R}(k)[x]$, we denote by B_k the specialization of B to a given value of k .

It is always possible to find a basis (a set of generators) of the steady state ideal of a network with cardinality equal to the rank of the network. Indeed, d of the steady state polynomials are redundant since they can be expressed as a linear combination of the $n - d$ remaining polynomials.

Definition 1.1. We say that a reaction network is *multistationary* if there exists a strictly positive vector $k \in \mathbb{R}_{>0}^{|\mathcal{R}|}$ such that the system $F_{k,i}(x) = 0$, $i \in [n]$, has more than one positive solution in a stoichiometric compatibility class. Alternatively, given a matrix of conservation laws Z , the system

$$N_k \psi(x) = 0 \quad \text{and} \quad Zx = T$$

has at least two positive solutions for some positive k and $T \in \mathbb{R}^d$.

2 Binomial networks and multistationarity

In this section we discuss and expand known results on determining whether a network is multistationary when the steady state ideal is binomial. The main references are [12, 13]. An ideal is binomial if it admits a binomial basis, that is, a basis with all polynomials having at most two terms. It is well known that an ideal is binomial if and only if the reduced Gröbner basis in an arbitrary monomial order consists only of binomials [3, Corollary 1.2].

We start with an observation on changing bases in $\mathbb{R}(k)[x]$ for $k = (k_1, \dots, k_r)$ and $x = (x_1, \dots, x_n)$. For a set of polynomials A in a polynomial ring $K[x]$, we let $V(A) \subseteq K^n$ denote its solution set, which agrees with $V(\langle A \rangle)$. If B and B' are bases of the same ideal I in $\mathbb{R}(k)[x]$, then $V(B) = V(B') \subseteq (\mathbb{R}(k))^n$. However, this does not imply that the specializations to real values $k \in \mathbb{R}^r$ agree, that is, it can happen that $V(B_k) \neq V(B'_k) \subseteq \mathbb{R}^n$. Since we want to study the steady state ideal in $\mathbb{R}(k)[x]$ but obtain results for specific values of k , we introduce the following definition.

Definition 2.1. Let \mathcal{N} be a reaction network and $B \subseteq \mathbb{R}(k)[x]$ the set of steady state polynomials. A basis B' of the steady state ideal of \mathcal{N} is called *admissible* if for every $k \in \mathbb{R}_{>0}^r$ it holds that $V(B_k) = V(B'_k)$. The network \mathcal{N} is a *binomial network* if the steady state ideal has an admissible binomial basis.

We consider a sufficient condition to decide whether the solution sets of two parametric systems agree for any specialization of the parameters, and in particular, for when a basis of the steady state ideal is admissible. Let $B = \{f_1, \dots, f_\ell\}$ and $B' = \{f'_1, \dots, f'_{\ell'}\}$. We consider representations of B in terms of B' and *vice versa*, that is, we write

$$f_i = \sum_{j \in [\ell']} \frac{h_{ij}}{h_i} f'_j, \quad \text{for } i \in [\ell], \quad \text{and} \quad f'_i = \sum_{j \in [\ell]} \frac{h'_{ij}}{h'_i} f_j, \quad \text{for } i \in [\ell'], \quad (2)$$

with $h_i, h'_i \in \mathbb{R}[k]$ and $h_{i1}, \dots, h_{i\ell'}, h'_{i1}, \dots, h'_{i\ell} \in \mathbb{R}[k][x]$. Note that these representations might not be unique.

Lemma 2.2. *With the notation above, given two bases B and B' of an ideal in $\mathbb{R}(k)[x]$, if k^* is not in the zero set of $(\prod_{i=1}^{\ell} h_i)(\prod_{i=1}^{\ell'} h'_i)$, then $\langle B_{k^*} \rangle = \langle B'_{k^*} \rangle$.*

Proof. For all $i \in [\ell]$ and $j \in [\ell']$ we have that $\frac{h_{k^*,ij}}{h_{k^*,i}}, \frac{h'_{k^*,ji}}{h'_{k^*,j}} \in \mathbb{R}[x]$ and the equalities in (2) specialize to k^* . Hence $B_{k^*} \subseteq \langle B'_{k^*} \rangle$ and $B'_{k^*} \subseteq \langle B_{k^*} \rangle$ and so $\langle B_{k^*} \rangle = \langle B'_{k^*} \rangle$. \square

In particular, if $(\prod_{i=1}^{\ell} h_i)(\prod_{i=1}^{\ell'} h'_i)$ has no positive solution, then $V(B_k) = V(B'_k)$ for all positive k .

Example 2.3. Consider the following reaction network



The set of steady state polynomials is

$$B = \{(k_1 - k_2)x_1 + (2k_3)x_2, (2k_2)x_1 + (-k_3 + k_4)x_2\},$$

and the set $B' = \{x_1 - x_2, x_1 - 2x_2\}$ is another basis of the steady state ideal in $\mathbb{R}(k)[x]$. We have that $(a, b) \in \mathbb{R}_{>0}^2$ belongs to $V(B_k)$ with $k = (2\frac{b}{a} + 1, 1, 1, 2\frac{a}{b} + 1)$. But $V(B'_k) \cap \mathbb{R}_{>0}^2 = \emptyset$ for every choice of k . Hence B' is not an admissible basis.

The connection between binomial ideals and multistationarity is as follows. Consider a system of binomial equations in $\mathbb{R}(k)[x]$, say

$$p_1(k)x^{c_1} - p'_1(k)x^{c'_1} = 0, \quad \dots \quad p_s(k)x^{c_s} - p'_s(k)x^{c'_s} = 0. \quad (3)$$

If one of the equations has only one term, or the two terms of a binomial have the same sign, then the system does not admit positive solutions. If $p_i \neq 0$ and $p'_i \neq 0$ in $\mathbb{R}(k)$, the positive solutions to (3) are the positive solutions of the following system

$$x^{c_1 - c'_1} = \frac{p'_1(k)}{p_1(k)}, \quad \dots \quad x^{c_s - c'_s} = \frac{p'_s(k)}{p_s(k)}.$$

Letting

$$\gamma(k) := \begin{bmatrix} \frac{p'_1(k)}{p_1(k)} \\ \vdots \\ \frac{p'_s(k)}{p_s(k)} \end{bmatrix} \quad \text{and} \quad M := \begin{bmatrix} c_1 - c'_1 \\ \vdots \\ c_s - c'_s \end{bmatrix}^t \in \mathbb{R}^{n \times s}, \quad (4)$$

the set of positive solutions of (3) for a positive vector k such that $p_i(k) \neq 0$ for all $i \in [s]$ is

$$\{x \in \mathbb{R}_{>0}^n \mid x^M = \gamma(k)\}. \quad (5)$$

Let M' be a matrix whose rows form a basis of the orthogonal complement of the row space of M . The solution set of $x^M = \gamma(k)$ is non-empty if and only if $M' \ln(\gamma(k)) = 0$, where $\ln(\gamma(k))$ is defined component-wise. To see why, take the logarithm of both sides, $M^t \ln(x) = \ln(\gamma(k))$, and impose that $\ln(\gamma(k))$ belongs to the image of M^t .

The parametrization in (5) of positive solutions of a binomial system (3) makes it possible to use results of [12, 13] for detecting multistationarity of binomial networks.

Theorem 2.4 ([12], Proposition 3.9 and Corollary 3.11). *Let \mathcal{N} be a binomial network. Let M and $\gamma(k)$ be as in (4), obtained from an admissible binomial basis B of I , and Z be a matrix of conservation laws. Consider the following conditions:*

(surj) *Surjectivity condition: for all $x \in \mathbb{R}_{>0}^n$ there exists $k \in \mathbb{R}_{>0}^{\mathcal{R}}$ such that $x^M = \gamma(k)$ (equivalently, $x \in V(B_k)$).*

(sign) *Sign Condition: there exist $u, v \in \mathbb{R}^n \setminus \{0\}$ such that $M^t u = Zv = 0$ and $\sigma(u) = \sigma(v)$.*

Then we have:

- (i) Assume (surj) is satisfied. Then \mathcal{N} is multistationary if and only if (sign) holds.
- (ii) If (sign) does not hold, then \mathcal{N} is not multistationary.

Lemma 2.5 ([13]). *A binomial network \mathcal{N} with stoichiometric matrix N satisfies (surj) if and only if $\ker N \cap \mathbb{R}_{>0}^{|\mathcal{R}|} \neq \emptyset$.*

Proof. The solution set of an admissible binomial basis agrees with the solution set of the set of steady state polynomials for every $k \in \mathbb{R}_{>0}^{|\mathcal{R}|}$. Therefore (surj) is equivalent to the statement

$$\text{for all } x \in \mathbb{R}_{>0}^n, \text{ there exists } k \in \mathbb{R}_{>0}^{|\mathcal{R}|} \text{ such that } N_k \psi(x) = 0.$$

Now $N_k \psi(x) = 0$ is equivalent to $\text{diag}(k) \psi(x) \in \ker N \cap \mathbb{R}_{>0}^{|\mathcal{R}|}$. Thus if $\ker N \cap \mathbb{R}_{>0}^{|\mathcal{R}|} = \emptyset$, then (surj) fails. Conversely given $v \in \ker N \cap \mathbb{R}_{>0}^{|\mathcal{R}|}$ and any $x \in \mathbb{R}_{>0}^n$, by taking $k = \frac{v}{\psi(x)}$ (defined component-wise), we see that (surj) holds. \square

Networks fulfilling the condition of the lemma above are often called *consistent*. Using Lemma 2.5 we can check (surj) algorithmically. For a matrix N , $U = \ker(N) \cap \mathbb{R}_{\geq 0}^{|\mathcal{R}|}$ is a convex set. A set of vectors in $\mathbb{R}_{\geq 0}^{|\mathcal{R}|}$ is an *extremal generating set* for U if their non-negative linear combinations generate U and none of them is a non-negative combination of the rest. Then U contains a strictly positive vector if and only if the sum of the vectors in an extremal generating set is positive. To find an extremal generating set for a convex set one can use existing algorithms, e.g. [16, Appendix B].

Let \mathcal{N} be a binomial reaction network with an admissible binomial basis B . Let M be the exponent matrix in the parametrization of its positive steady states as in (5), and Z be a matrix of conservation laws. For $\lambda = (\lambda_1, \dots, \lambda_n)$ a vector of indeterminates, define

$$\Gamma = \begin{bmatrix} M^\lambda \\ Z \end{bmatrix} \in (\mathbb{R}[\lambda])^{n \times n} \quad (6)$$

and consider the following conditions:

(rank) *Rank Condition*: The number of elements of B , equivalently the number of columns of M , is equal to the rank of the network.


(det) *Determinant Condition*: Viewed as a polynomial in λ , $\det(\Gamma)$ is either zero or has at least one positive and at least one negative coefficient.

Note that by [12, Lemma 2.11], $\det(\Gamma)$ is a polynomial in λ that is linear or constant in each λ_i . Theorem 2.13 of [12] states that provided (rank) is fulfilled, then (sign) holds if and only if (det) holds. Combining this with Theorem 2.4, if (surj) and (rank) are satisfied, then \mathcal{N} is multistationary if and only if (det) holds. This yields to the following definition and theorem.

Definition 2.6. A binomial network \mathcal{N} is *complete* if (surj) and (rank) hold for an admissible binomial basis.

Theorem 2.7 (Determinant criterion for complete binomial networks). *Consider a complete binomial network, and let M be as in (5) obtained from an admissible binomial basis of the steady state ideal that satisfies both (surj) and (rank). Then the network is multistationary if and only if (det) holds.*

Example 2.8. Consider the following network modeling a simple biological circuit:



An admissible binomial basis of the steady state ideal is

$$B = \left\{ \kappa_1 - \frac{\kappa_2 \kappa_4}{\kappa_3 + \kappa_4} x_1 e, \frac{\kappa_2 \kappa_4}{\kappa_3 + \kappa_4} x_1 e - \kappa_7 x_2, \kappa_2 x_1 e - (\kappa_3 + \kappa_4) y_1, \kappa_5 x_1^2 e - \kappa_6 y_2 \right\}.$$

The matrices M , Z and N are

$$M = \begin{bmatrix} 1 & 1 & -1 & -2 \\ 0 & -1 & 0 & 0 \\ 1 & 1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, Z = [0 \ 0 \ 1 \ 1 \ 1], N = \begin{bmatrix} 1 & -1 & 1 & 0 & -2 & 2 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & -1 & 1 & 1 & -1 & 1 & 0 \\ 0 & 1 & -1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 \end{bmatrix}.$$

The set $P = \{(1, 1, 0, 1, 0, 0, 1), (0, 1, 1, 0, 0, 0, 0), (0, 0, 0, 0, 1, 1, 0)\}$ is an extremal generating set for $\ker(N) \cap \mathbb{R}_{\geq 0}^7$. Since the sum of the vectors in P has all entries positive, (surj) holds by Lemma 2.5. Now taking $u = v = (-1, 0, 1, 0, -1)$, the condition (sign) holds. Therefore by Theorem 2.4 this network is multistationary.

Alternatively, we have $\text{rank}(N) = 4$ and B has 4 elements; hence (rank) holds and this binomial network is complete. We have

$$\det(\Gamma) = \begin{vmatrix} \lambda_1 & 0 & \lambda_3 & 0 & 0 \\ \lambda_1 & -\lambda_2 & \lambda_3 & 0 & 0 \\ -\lambda_1 & 0 & -\lambda_3 & \lambda_4 & 0 \\ -2\lambda_1 & 0 & -\lambda_3 & 0 & \lambda_5 \\ \hline 0 & 0 & 1 & 1 & 1 \end{vmatrix} = \lambda_1 \lambda_2 \lambda_3 \lambda_4 - \lambda_1 \lambda_2 \lambda_4 \lambda_5.$$

The network is multistationary by Theorem 2.7 since (det) holds.

3 Intermediates and multistationarity

In this section we introduce a particular type of species, intermediates, and extended and core networks obtained by adding or removing intermediates. We proceed to present results on multistationarity of extended and core networks from [8] and specifically for binomial networks from [14].

3.1 Intermediates

A species Y is an *intermediate* if it is also a complex, that is belongs to \mathcal{C} , only appears in the complex Y , and further both the outdegree and indegree of Y are at least one in the digraph of the network [8]. Given a set of intermediates $\mathcal{Y} = \{Y_1, \dots, Y_m\}$, let $\mathcal{X} = \mathcal{S} \setminus \mathcal{Y} = \{X_1, \dots, X_n\}$ be the set of non-intermediates. Then \mathcal{S} is the disjoint union of \mathcal{X} and \mathcal{Y} . From now on, the species are ordered such that intermediates are after non-intermediates. Then by (x, y) we mean the vector $(x_1, \dots, x_n, y_1, \dots, y_m)$. A complex that is not an intermediate is called a non-intermediate complex.

Given an intermediate Y , an *input* for Y is a non-intermediate complex c such that there exists a directed reaction path from c to Y with all vertices other than c being intermediates. The intermediate Y is an ℓ -input intermediate if it has ℓ inputs [14].

Definition 3.1 ([8]). Let $\mathcal{N} = (\mathcal{S}, \mathcal{C}, \mathcal{R})$ and $\tilde{\mathcal{N}} = (\tilde{\mathcal{S}}, \tilde{\mathcal{C}}, \tilde{\mathcal{R}})$ be two reaction networks. We say that $\tilde{\mathcal{N}}$ is an *extension* of \mathcal{N} via the addition of intermediates Y_1, \dots, Y_m if

- (i) $\mathcal{Y} = \{Y_1, \dots, Y_m\}$ is a set of intermediates of $\tilde{\mathcal{N}}$.
- (ii) $\mathcal{S} \cup \mathcal{Y} = \tilde{\mathcal{S}}$ and $\mathcal{C} \cup \mathcal{Y} = \tilde{\mathcal{C}}$.

- (iii) $c \rightarrow c' \in \mathcal{R}$ if and only if there is a directed path from c to c' in the digraph associated with $\tilde{\mathcal{N}}$, such that all vertices other than c and c' belong to \mathcal{Y} (there might be none).

In this case \mathcal{N} is called the *core network* of $\tilde{\mathcal{N}}$.

Notations κ , and symbols with tilde are used to refer to reaction rate constants and the corresponding objects of the extended network respectively. Let \mathcal{N} be a reaction network and $\tilde{\mathcal{N}}$ an extension of it via the addition of m intermediates, Y_1, \dots, Y_m . Choose an input complex c_i for each intermediate Y_i and let $[c_1 \ \dots \ c_m] \in \mathbb{R}^{n \times m}$ be the matrix whose columns are c_1, \dots, c_m . It follows from Theorem 2.1 of [8] that if Z is a matrix of conservation laws for \mathcal{N} , then a matrix of conservation laws for $\tilde{\mathcal{N}}$ is

$$\tilde{Z} = [Z \ ; \ Z [c_1 \ \dots \ c_m]] \in \mathbb{R}^{d \times (n+m)}. \quad (7)$$

In particular the corank of \mathcal{N} and $\tilde{\mathcal{N}}$ agree and the rank of $\tilde{\mathcal{N}}$ is the rank of \mathcal{N} plus m .

We next introduce a simple type of extended networks via the addition of intermediates that are useful in the study of multistationarity, see Theorem 3.11.

Definition 3.2. Let \mathcal{N} be a network and $C = \{c_1, \dots, c_m\} \subseteq \mathcal{C}$. The *canonical extension* of \mathcal{N} associated with C , denoted by $\tilde{\mathcal{N}}_C = (\tilde{\mathcal{S}}_C, \tilde{\mathcal{C}}_C, \tilde{\mathcal{R}}_C)$, is the extension of \mathcal{N} via the addition of 1-input intermediates Y_1, \dots, Y_m such that

$$\tilde{\mathcal{R}}_C = \mathcal{R} \cup \{c_i \Rightarrow Y_i \mid i \in [m]\}.$$

The canonical extension associated with $C = \mathcal{C}$ is called the *largest canonical extension*.

We now review the key results in [8] that relate the steady states of extended and core networks. We start by studying the steady state polynomials of the two networks. Let $\tilde{\mathcal{N}}$ be an extension of \mathcal{N} via the addition of intermediates Y_1, \dots, Y_m and $C \subseteq \mathcal{C}$ be the set of input complexes. The steady state polynomials associated with the intermediates yield a system $F_{n+1}(x, y) = \dots = F_{n+m}(x, y) = 0$ that is linear in y and square. As shown in [8], the solution to this linear system is of the form

$$y_i = \sum_{c \in C} \mu_{i,c} x^c, \quad i \in [m] \quad (8)$$

where $\mu_{i,c}$ is a rational function in $\mathbb{R}(\kappa)$ with all non-zero coefficients positive (see also (22)). Consider the following map

$$\begin{aligned} \phi: \quad \mathbb{R}[k] &\longrightarrow \mathbb{R}(\kappa) \\ k_{c \rightarrow c'} &\longmapsto \phi_{c \rightarrow c'}(\kappa) = \kappa_{c \rightarrow c'} + \sum_{i=1}^m \kappa_{Y_i \rightarrow c'} \mu_{i,c}, \end{aligned} \quad (9)$$

where it is understood that $\kappa_{c \rightarrow c'} = 0$ and $\kappa_{Y_i \rightarrow c'} = 0$ if $c \rightarrow c'$ and $Y_i \rightarrow c'$ do not belong to $\tilde{\mathcal{R}}$ respectively. Then the steady state polynomials F, \tilde{F} of \mathcal{N} and $\tilde{\mathcal{N}}$ for non-intermediate species relate in the following way

$$\tilde{F}_{\kappa,i}(x, \sum_{c \in C} \mu_{1,c} x^c, \dots, \sum_{c \in C} \mu_{m,c} x^c) = F_{\phi(k),i}(x), \quad i \in [n].$$

Given $f/g \in \mathbb{R}(k)$ such that $\phi(g) \neq 0$, then $\phi(f/g)$ is well defined in $\mathbb{R}(\kappa)$. If $G \in \mathbb{R}(\kappa)[x]$ is a polynomial in x such that all coefficients are rational functions with non-vanishing denominator upon applying ϕ , then we consider the polynomial $\Phi(G)$ in $\mathbb{R}(\kappa)[x, y]$ obtained by applying ϕ on the coefficients of G . In particular, if the rational functions $\phi_{c \rightarrow c'}$ are algebraically independent over \mathbb{R} , then there is no polynomial with coefficients in \mathbb{R} that

identically vanishes when evaluated on the image of ϕ . Then the map ϕ extends to a map of polynomial rings $\Phi: \mathbb{R}(k)[x] \rightarrow \mathbb{R}(\kappa)[x, y]$. Strategies to check this algebraic independence condition as well as classes of intermediates that satisfy it are described in [14, §4]. In particular the rational functions $\phi_{c \rightarrow c'}$ are algebraically independent over \mathbb{R} for all canonical extensions by [14, Corollary 4.6].

In order to introduce Theorem 3.3 below, we need to consider the following conditions. Let $\omega_1, \dots, \omega_d$ be a basis of S^\perp and $C' \subseteq C$ consist of the complexes c such that $\omega_j \cdot c \neq 0$ for some $j \in [d]$ (i.e. $c \notin S$). We define two realization conditions on the reaction rate constants of \mathcal{N} and $\tilde{\mathcal{N}}$:

(i) *Realization condition:*

For all $k \in \mathbb{R}_{>0}^{\mathcal{R}}$, there exists $\kappa \in \mathbb{R}_{>0}^{\tilde{\mathcal{R}}}$ such that $k = \phi(\kappa)$.

(ii) *Generalized realization condition:*

For all $k \in \mathbb{R}_{>0}^{\mathcal{R}}$ and $r \in \mathbb{R}_{>0}^{C'}$, there exists $\kappa \in \mathbb{R}_{>0}^{\tilde{\mathcal{R}}}$ such that

$$k = \phi(\kappa) \quad \text{and} \quad r_c = \sum_{i \in [m]} \mu_{i,c} \quad \text{for all } c \in C'.$$

Note that $\mu_{i,c}$ depends as well on κ in the last statement. In §5 we focus on how to check whether these realization conditions are satisfied. The proof of the next theorem is found in [8]. The first part is Theorem 5.1, and the second part is discussed in the text.

Theorem 3.3 ([8]). *Let $\tilde{\mathcal{N}}$ be an extension of \mathcal{N} via the addition of intermediates Y_1, \dots, Y_m .*

(i) *If the realization condition holds, then multistationarity of \mathcal{N} implies multistationarity of $\tilde{\mathcal{N}}$.*

(ii) *Let $C \subseteq \mathcal{C}$ be the set of inputs of Y_1, \dots, Y_m . If the generalized realization condition holds for $\tilde{\mathcal{N}}$, then $\tilde{\mathcal{N}}$ is multistationary if and only if the canonical extension $\tilde{\mathcal{N}}_C$ is multistationary.*

Definition 3.4. Let \mathcal{N} be a reaction network and $C \subseteq \mathcal{C}$. The *canonical class* associated with C is the set of all extensions of \mathcal{N} via the addition of intermediates with input set C that satisfy the generalized realization condition.

Proposition 3.5. *The generalized realization condition holds for canonical extensions. Therefore, a canonical class is not empty.*


Proof. Let $\tilde{\mathcal{N}}$ be the canonical extension of a network \mathcal{N} associated with $C = \{c_1, \dots, c_m\} \subseteq \mathcal{C}$. For every $i \in [m]$, we have $\mu_{i,c} = \frac{\kappa_{c_i \rightarrow Y_i}}{\kappa_{Y_i \rightarrow c_i}}$ if $c = c_i$ and zero otherwise. Since no two intermediates have a common input, the generalized realization condition holds if for every $k \in \mathbb{R}_{>0}^{\mathcal{R}}$ and $r \in \mathbb{R}_{>0}^m$, there exists $\kappa \in \mathbb{R}_{>0}^{\tilde{\mathcal{R}}}$ such that

$$k_{c \rightarrow c'} = \kappa_{c \rightarrow c'} \quad \text{for all } c \rightarrow c' \in \mathcal{R} \quad \text{and} \quad r_i = \frac{\kappa_{c_i \rightarrow Y_i}}{\kappa_{Y_i \rightarrow c_i}} \quad \text{for all } i \in [m].$$

This condition clearly holds. □

Theorem 3.3 implies that multistationarity of an extended network $\tilde{\mathcal{N}}$ satisfying the generalized realization condition is equivalent to multistationarity of any network in the same canonical class of $\tilde{\mathcal{N}}$, in particular of the canonical extensions in the class. Canonical extensions have a simple structure and preserve some important properties of \mathcal{N} as we will see below. Hence they are chosen as representatives of the class.


Example 3.6. The following digraph defines a reaction network corresponding to the Mitogen-Activated Protein Kinase cascade [2]:



If we consider Y_1, \dots, Y_6 as intermediates, then the associated core network is



The generalized realization condition holds by Example 5.2 in §5. The canonical extension in the canonical class of (10) is the following network:



3.2 Binomial networks and intermediates

In this subsection, building on the results in [14], we relate the condition (det) for the core and extended network. This leads to a determinant criterion for multistationarity of extended networks with a complete binomial core network.

Let $\tilde{\mathcal{N}}$ be an extension of a binomial reaction network \mathcal{N} via the addition of intermediates Y_1, \dots, Y_m . For a binomial basis B of the steady state ideal $I \subseteq \mathbb{R}(k)[x]$ of \mathcal{N} , let $\text{rem}(f, B)$ denote the remainder of the division of a polynomial f by B . Assume $\Phi(B)$ is well defined and consider

$$\tilde{B} = \Phi(B) \cup \left\{ y_i - \sum_{c \in \mathcal{C}} \mu_{i,c} x^c, i \in [m] \right\}, \tag{11}$$

$$\tilde{B}' = \Phi(B) \cup \left\{ y_i - \text{rem} \left(\sum_{c \in \mathcal{C}} \mu_{i,c} x^c, \Phi(B) \right), i \in [m] \right\}. \tag{12}$$

If the set on the right-hand side of the union in either \tilde{B} or \tilde{B}' consists of binomials, then they have the form

$$\Phi(B) \cup \{ y_i - p_i(\kappa) x^{\alpha_i}, i \in [m] \} \tag{13}$$

where $p_i(\kappa) \in \mathbb{R}(\kappa)$ and α_i is a vector of non-negative integers. If all intermediates are 1-input, then a binomial basis of \tilde{I} is

$$\tilde{B} = \Phi(B) \cup \{ y_i - \mu_{i,c} x^{c_i}, i \in [m] \}, \tag{14}$$

where c_i is the only input of Y_i , and this basis is admissible provided B is an admissible binomial basis of I and $\Phi(B)$ is defined. This applies in particular to canonical extensions.

Definition 3.7. Let \mathcal{N} be a binomial reaction network and $\tilde{\mathcal{N}}$ an extension of it via the addition of intermediates. $\tilde{\mathcal{N}}$ is a *binomial extension* of \mathcal{N} if there exists an admissible binomial basis B of \mathcal{N} such that $\Phi(B)$ is well defined, no coefficient of B becomes zero under Φ , and further, either \tilde{B} or \tilde{B}' is an admissible binomial basis of $\tilde{\mathcal{N}}$.

Remark 3.8. If the functions $\phi_{c \rightarrow c'}$ are algebraically independent over \mathbb{R} , then $\Phi(B)$ is well defined and no coefficient of B vanishes. By [14, Lemma 3.3], (11) and (12) are bases of the steady state ideal of the extended network $\tilde{I} \subseteq \mathbb{R}(\kappa)[x, y]$, and if B is admissible, then

so is \tilde{B} . To decide whether \tilde{B}' is also admissible, it suffices to check that the representations of $y_i - \text{rem}(\sum_{c \in \mathcal{C}} \mu_{i,c} x^c, \Phi(G))$ in terms of \tilde{B} and that of $y_i - \sum_{c \in \mathcal{C}} \mu_{i,c} x^c$ in terms of \tilde{B}' are well defined for all κ (c.f. Lemma 2.2).

Further, by [14, Theorem 3.10], the steady state ideal \tilde{I} of $\tilde{\mathcal{N}}$ is binomial if and only if the steady state ideal I of \mathcal{N} is binomial and for any reduced Gröbner basis G of I , $\text{rem}(\sum_{c \in \mathcal{C}} \mu_{i,c} x^c, \Phi(G))$ has at most one term for all $i \in [m]$.

Lemma 3.9. *Let $\tilde{\mathcal{N}}$ be a binomial extension of a binomial network \mathcal{N} . Condition (rank) holds for $\tilde{\mathcal{N}}$ if and only if it holds for \mathcal{N} .*

Proof. Let B and \bar{B} be admissible binomial bases of the steady state ideals of \mathcal{N} and $\tilde{\mathcal{N}}$ respectively, such that \bar{B} is either \tilde{B} in (11) or \tilde{B}' in (12). Then $|\bar{B}| = |B| + m$. If $n - d$ is the rank of \mathcal{N} , then by (7) and the text below it, the condition (rank) for $\tilde{\mathcal{N}}$ is $n + m - d = |\bar{B}| = |B| + m$, which is the rank condition for \mathcal{N} , $|B| = n - d$. \square

It follows from the lemma above that a binomial extension of a complete binomial network satisfying (surj) is a complete binomial network. In general, for an arbitrary extended network $\tilde{\mathcal{N}}$, we cannot guarantee that (surj) holds provided it holds for \mathcal{N} . However, it does for canonical extensions.

Proposition 3.10. *Let \mathcal{N} be a binomial network. Any canonical extension $\tilde{\mathcal{N}}_C$ of \mathcal{N} is a binomial extension. Further, (surj) holds for \mathcal{N} with an admissible binomial basis B if and only if (surj) holds for $\tilde{\mathcal{N}}_C$ with \tilde{B} as in (11). Therefore, a canonical extension of a complete binomial network is also complete.*

Proof. By [14, Corollary 4.6], the functions $\phi_{c \rightarrow c'}$ are algebraically independent for canonical extensions. By (14) and Remark 3.8, $\tilde{\mathcal{N}}_C$ is a binomial extension.

For the second part of the proposition, write $C = \{c_1, \dots, c_m\}$. Denote the reaction rate constants of \mathcal{N} by k_1, k_2, \dots, k_r following the order of the reaction set. The network $\tilde{\mathcal{N}}_C$ has $r + 2m$ reactions. We denote the reaction rate constants of the reactions of $\tilde{\mathcal{N}}_C$ that are also in \mathcal{N} with $\kappa_1, \dots, \kappa_r$ and of the other reactions by $c_i \xrightleftharpoons[\kappa_{r+2i}]{\kappa_{r+2i-1}} Y_i$ for $i \in [m]$. Then $\mu_{i,c_i} = \frac{\kappa_{r+2i-1}}{\kappa_{r+2i}}$. Let M be the matrix constructed in (4) for the basis B . Now (surj) holds for $\tilde{\mathcal{N}}_C$ if for every $(x, y) \in \mathbb{R}_{>0}^{n+m}$ there exists $(\kappa_1, \dots, \kappa_{r+2m}) \in \mathbb{R}_{>0}^{r+2m}$ such that $x^M = \gamma(\kappa)$ and $\frac{y_i}{x^{c_i}} = \frac{\kappa_{r+2i-1}}{\kappa_{r+2i}}$ for all $i \in [m]$. Since the first part of the system does not depend on $\kappa_{r+1}, \dots, \kappa_{r+2m}$, the second part is always satisfied independently from the first part by letting $\kappa_{r+2i} = x^{c_i}$ and $\kappa_{r+2i-1} = y_i$. The first part of the system is exactly the same as (surj) for \mathcal{N} after replacing k_i by κ_i . Hence (surj) holds for $\tilde{\mathcal{N}}_C$ if and only if it holds for \mathcal{N} . The last statement follows from Lemma 3.9. \square

Recall that criterion (det) can be used to determine multistationarity for complete networks, c.f. Theorem 2.7. Consider a binomial extension $\tilde{\mathcal{N}}$ of a complete binomial network \mathcal{N} . The steady state ideal of $\tilde{\mathcal{N}}$ has an admissible binomial basis \tilde{B} of the form (13), with B an admissible binomial basis of the steady state ideal of \mathcal{N} . Let $M \in \mathbb{R}^{n \times s}$ be the exponent matrix associated with the binomials in B , c.f. (4). Since no coefficient of B becomes zero under ϕ , the exponents of the monomials in B and $\Phi(B)$ agree. Hence the exponent matrix \tilde{M} associated with the binomials in \tilde{B} has the following form:

$$(\tilde{M})^t = \left[\begin{array}{c|c} M^t & 0 \\ \hline -\alpha_1 & \\ \vdots & I_m \\ -\alpha_m & \end{array} \right] \in \mathbb{R}^{(s+m) \times (n+m)}.$$

Using (7), the corresponding matrices Γ and $\tilde{\Gamma}$ in (6), for \mathcal{N} and $\tilde{\mathcal{N}}$ respectively, are:

$$\Gamma = \begin{bmatrix} M_\lambda^t \\ Z \end{bmatrix} \in \mathbb{R}^{n \times n}, \quad \tilde{\Gamma} = \begin{bmatrix} M_\lambda^t & & & & 0 \\ \hline -\alpha_1 & \lambda_{n+1} & & & 0 \\ \vdots & & \ddots & & \\ -\alpha_m & 0 & & & \lambda_{n+m} \\ \hline Z & Zc_1^t & \dots & Zc_m^t & \end{bmatrix} \in \mathbb{R}^{(n+m) \times (n+m)}, \quad (15)$$

where Z is a matrix of conservation laws for \mathcal{N} , c_1, \dots, c_m are chosen inputs for Y_1, \dots, Y_m and $\lambda = (\lambda_1, \dots, \lambda_n)$. If $\tilde{\mathcal{N}}_C$ is a canonical extension, then $\alpha_i = c_i$ in $\tilde{\Gamma}$, c.f. (14), in which case we denote the matrix by $\tilde{\Gamma}_C$.

The results in this subsection combined with Theorem 3.3(ii) imply that we can use (det) to detect multistationarity of networks that are not necessarily binomial, but such that the core network is binomial and complete.

Theorem 3.11 (Determinant criterion for extensions of complete binomial networks). *Let \mathcal{N} be a complete binomial network and $\tilde{\mathcal{N}}$ an extended network in the canonical class associated with $C \subseteq \mathcal{C}$. Then $\tilde{\mathcal{N}}$ is multistationary if and only if $\det(\tilde{\Gamma}_C)$ is either identically zero or has both positive and negative coefficients.*

Proof. By Theorem 3.3(ii), $\tilde{\mathcal{N}}$ is multistationary if and only if $\tilde{\mathcal{N}}_C$ is. Now $\tilde{\mathcal{N}}_C$ is a complete binomial network and the matrix $\tilde{\Gamma}_C$ is defined as in (6) for an admissible binomial basis of the steady state ideal that satisfies both (surj) and (rank). Thus $\tilde{\mathcal{N}}_C$ is multistationary if and only if (det) holds with the matrix $\tilde{\Gamma}_C$ by Theorem 2.7. \square

Example 3.12. (Continued from Example 3.6) The network in Example 3.6 is not binomial by [14, Example 3.13], but the core network is a complete binomial network. The extended network belongs to the canonical class associated with $C = \{X_0 + E, X_1 + E, X_2 + F, X_1 + F, X_0 + F\}$. With a suitable choice of basis B , the matrix $\tilde{\Gamma}_C$ is as follows:

$$\begin{bmatrix} -\lambda_1 & \lambda_2 & 0 & -\lambda_4 & \lambda_5 & & & & & 0 \\ 0 & -\lambda_2 & \lambda_3 & -\lambda_4 & \lambda_5 & & & & & \\ \hline -\lambda_1 & 0 & 0 & -\lambda_4 & 0 & \lambda_6 & & & & 0 \\ 0 & -\lambda_2 & 0 & -\lambda_4 & 0 & & \lambda_7 & & & \\ 0 & 0 & -\lambda_3 & 0 & -\lambda_5 & & & \lambda_8 & & \\ 0 & -\lambda_2 & 0 & 0 & -\lambda_5 & & & & \lambda_9 & \\ -\lambda_1 & 0 & 0 & 0 & -\lambda_5 & 0 & & & & \lambda_{10} \\ \hline 1 & 1 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 & 1 \end{bmatrix}.$$

The polynomial $\det(\tilde{\Gamma}_C)$ has terms with different signs. Therefore by Theorem 3.11, the network in (10) is multistationary.

4 Lifting multistationarity and the multistationarity structure

In this section we use the following notation: for $J \subseteq [n]$ and λ an n -tuple of indeterminates/numbers, we define $\lambda_J = \prod_{i \in J} \lambda_i$.

4.1 Lifting multistationarity

Theorem 3.3 tells us that multistationarity of the core network implies multistationarity of the extended network if the realization condition is satisfied. In this scenario we informally say that multistationarity is lifted. In this subsection we show that multistationarity is lifted for binomial extensions of complete binomial networks, even if the realization condition is not satisfied. Before that, we start with a lemma on the structure of $\tilde{\Gamma}$.

Lemma 4.1. *Let $\tilde{\mathcal{N}}$ be a binomial extension of a complete binomial network \mathcal{N} via the addition of intermediates Y_1, \dots, Y_m , and let $\Gamma, \tilde{\Gamma}$ be as in (15). Then*

$$\det(\tilde{\Gamma}) = \lambda_{[n+1, n+m]} \det(\Gamma) + p'(\lambda),$$

where $p'(\lambda)$ is a polynomial in λ such that none of its terms is divisible by $\lambda_{[n+1, n+m]}$.

Proof. Let $s = \text{rank}(\mathcal{N})$ and $\tilde{\Gamma}_{[s+1, s+m], [n+1, n+m]}$ be the submatrix of $\tilde{\Gamma}$ obtained by removing the rows with index in $[s+1, s+m]$ and the columns with index in $[n+1, n+m]$. By the generalized Laplacian expansion of $\det(\tilde{\Gamma})$ along rows $s+1, \dots, s+m$ we have that

$$\det(\tilde{\Gamma}) = \lambda_{[n+1, n+m]} \det(\tilde{\Gamma}_{[s+1, s+m], [n+1, n+m]}) + p'(\lambda) = \lambda_{[n+1, n+m]} \det(\Gamma) + p'(\lambda),$$


where $p'(\lambda)$ is a polynomial in λ . By construction, $p'(\lambda)$ does not have any monomial multiple of $\lambda_{[n+1, n+m]}$. \square

Theorem 4.2 (Lifting multistationarity). *Let $\tilde{\mathcal{N}}$ be a binomial extension of a complete binomial network \mathcal{N} via the addition of m intermediates Y_1, \dots, Y_m and let $\Gamma, \tilde{\Gamma}$ be as in (15). Assume in addition that (surj) holds for $\tilde{\mathcal{N}}$. If \mathcal{N} is multistationary and $\det(\Gamma) \neq 0$, then $\tilde{\mathcal{N}}$ is multistationary.*


Proof. Let $\lambda = (\lambda_1, \dots, \lambda_{n+m})$ and $\bar{\lambda} = (\lambda_1, \dots, \lambda_n)$. Since \mathcal{N} is multistationary and complete, by Theorem 2.7 we have that $\det(\Gamma)$ is a polynomial in $\lambda_1, \dots, \lambda_n$ with two terms of different non-zero sign, namely $\alpha \bar{\lambda}^u$ and $-\beta \bar{\lambda}^v$ with $\alpha, \beta > 0$ and $u, v \in \mathbb{Z}_{\geq 0}^n$. By Lemma 4.1, $\det(\tilde{\Gamma})$ has two terms with different non-zero sign $\alpha \bar{\lambda}^u \lambda_{[n+1, n+m]}$ and $-\beta \bar{\lambda}^v \lambda_{[n+1, n+m]}$. Since $\tilde{\mathcal{N}}$ is a complete binomial network, $\tilde{\mathcal{N}}$ is multistationary by Theorem 2.7. \square

We finish this subsection with an example where Theorem 4.2 allows us to conclude that an extended network is multistationary, while the realization condition is not satisfied (and hence Theorem 3.3 cannot be applied).

Example 4.3. Consider the following network \mathcal{N} :



\mathcal{N} has rank 4 and the steady state ideal has the following admissible binomial basis with 4 elements: $B = \{-k_1 x_0 e + k_5 x_1 f, -k_2 x_1 e + k_3 k_4 x_2 f, k_4 y_1 - k_3 x_2 f, (k_9 + 2k_{10} + 3k_{11})e - (3k_6 + 2k_7 + k_8)e^5\}$. Since $(1, 1, 1, 1, 1, 4, 5, 6, 6, 5, 4) \in \ker(N) \cap \mathbb{R}_{>0}^{11}$, (surj) holds. Therefore \mathcal{N} is a complete binomial network. Using (det), we see that \mathcal{N} is multistationary and $\det(\Gamma) \neq 0$. Now consider the following extension $\tilde{\mathcal{N}}$ of \mathcal{N} via the addition of one intermediate Y_2 :




This network does not satisfy the realization condition (see §2.2 in the electronic supplementary material of [8]). The well-defined set $\tilde{B}' = \Phi(B) \cup \{y_2 - \frac{4\kappa_7 e}{3\kappa_8 + 2\kappa_9 + \kappa_{10}}\}$ in (12) makes $\tilde{\mathcal{N}}$ a binomial extension. Further $(1, 1, 1, 1, 1, 5, 5, 4, 2, 4) \in \ker(\tilde{N}) \cap \mathbb{R}_{>0}^{10}$, and hence (surj) holds for \tilde{N} . By Theorem 4.2, we conclude that $\tilde{\mathcal{N}}$ is also multistationary.

4.2 Multistationarity structure

In this subsection we introduce the *multistationarity structure* of a core network, consisting of the subsets of complexes that give rise to multistationarity when being the input of some intermediate. Note that if $C_1 \subseteq C_2 \subseteq \mathcal{C}$, then $\tilde{\mathcal{N}}_{C_2}$ is a binomial extension of $\tilde{\mathcal{N}}_{C_1}$. Using this and Lemma 4.1, we devise a strategy to determine the multistationarity structure of complete binomial networks by computing the determinant of $\tilde{\Gamma}_{\mathcal{C}}$ corresponding to the largest canonical extension. We start with an example that illustrates the approach.

Example 4.4. Consider the network in Example 2.8, where Y_1 and Y_2 are intermediates. The associated core network is



which gives $\mathcal{C} = \{0, X_1, X_1 + E, X_2 + E, 2X_1 + E, X_2\}$. An admissible binomial basis of the steady state ideal of (16) is $\{k_1 - k_2 x_1 e, k_2 x_1 e - k_3 x_2\}$. Since the rank of (16) is two and $(1, 1, 1) \in \ker(N) \cap \mathbb{R}_{>0}^3$, (16) is a complete binomial network. The polynomial $\det(\tilde{\Gamma}_{\mathcal{C}})$ for the largest canonical extension $\tilde{\mathcal{N}}_{\mathcal{C}}$ is as follows:

$$\det(\tilde{\Gamma}_{\mathcal{C}}) = \begin{vmatrix} -\lambda_1 & 0 & -\lambda_3 & & & & & & & 0 \\ \lambda_1 & -\lambda_2 & \lambda_3 & & & & & & & \\ \hline 0 & 0 & 0 & \lambda_4 & & & & & & 0 \\ -\lambda_1 & 0 & 0 & & \lambda_5 & & & & & \\ -\lambda_1 & 0 & -\lambda_3 & & & \lambda_6 & & & & \\ 0 & -\lambda_2 & -\lambda_3 & & & & \lambda_7 & & & \\ -2\lambda_1 & 0 & -\lambda_3 & & & & & \lambda_8 & & \\ 0 & -\lambda_2 & 0 & 0 & & & & & \lambda_9 & \\ \hline 0 & 0 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \end{vmatrix} = \begin{aligned} & -\lambda_1 \lambda_2 \lambda_3 \lambda_4 \lambda_5 \lambda_6 \lambda_7 \lambda_9 \\ & + \lambda_1 \lambda_2 \lambda_3 \lambda_4 \lambda_5 \lambda_6 \lambda_8 \lambda_9 \\ & + \lambda_1 \lambda_2 \lambda_4 \lambda_5 \lambda_6 \lambda_7 \lambda_8 \lambda_9. \end{aligned}$$

Since $\mathcal{N}_{\mathcal{C}}$ is a binomial extension of all canonical extensions, and all canonical extensions are complete, we can use Lemma 4.1 to find $\det(\tilde{\Gamma}_C)$ for all $C \in \mathcal{C}$. For example, the 3rd and 5th complexes form the set $C = \{X_1 + E, 2X_1 + E\}$ and $\det(\tilde{\Gamma}_C)$ is the coefficient of $\lambda_4 \lambda_5 \lambda_7 \lambda_9$ in $\det(\tilde{\Gamma}_{\mathcal{C}})$:

$$\det(\tilde{\Gamma}_{\{X_1 + E, 2X_1 + E\}}) = -\lambda_1 \lambda_2 \lambda_3 \lambda_6 + \lambda_1 \lambda_2 \lambda_6 \lambda_8.$$

For any subset $J \subseteq \{4, \dots, 9\}$, the coefficient of λ_J in $\det(\tilde{\Gamma}_{\mathcal{C}})$ is $\det(\tilde{\Gamma}_C)$ for the set of complexes c_i such that $i + 3 \notin J$.

By Theorem 3.11, all networks in the canonical class associated with $\{X_1 + E, 2X_1 + E\}$ are multistationary. In particular, the network in Example 2.8 belongs to this canonical class (and hence is multistationary), since it satisfies the generalized realization condition:

for every $(k, r) \in \mathbb{R}_{>0}^{3+2}$ there exists $\kappa \in \mathbb{R}_{>0}^7$ such that

$$k_1 = \kappa_1, \quad k_2 = \frac{\kappa_2 \kappa_4}{\kappa_3 + \kappa_4}, \quad k_3 = \kappa_7, \quad r_1 = \frac{\kappa_2}{\kappa_3 + \kappa_4}, \quad r_2 = \frac{\kappa_5}{\kappa_6}.$$

Motivated by this example, we proceed as follows.

Definition 4.5. Let \mathcal{N} be a reaction network, with set of complexes \mathcal{C} . Let Mult be the set of all subsets of complexes $C \subseteq \mathcal{C}$ for which the canonical extension $\tilde{\mathcal{N}}_C$ of \mathcal{N} associated with C is multistationary. Denote the set of minimal elements of Mult with respect to inclusion by Circuits. The set Mult is called the *multistationarity structure* of \mathcal{N} and the elements of Circuits are called the *circuits of multistationarity* of \mathcal{N} .

By Theorem 3.3(i), the sets Mult and Circuits associated with a complete binomial network determine each other. Let $D_{\mathcal{N}} = \det(\tilde{\Gamma}_C)$, that is the determinant of $\tilde{\Gamma}$ associated with the largest canonical extension of \mathcal{N} . We assume that the set of complexes is ordered $\mathcal{C} = \{c_1, \dots, c_m\}$.

Lemma 4.6. *Assume \mathcal{N} is a complete binomial network. Then $C \in \text{Mult}$ if and only if the coefficient of $\prod_{c_i \in \mathcal{C}-C} \lambda_{n+i}$ in $D_{\mathcal{N}}$ is zero or has two terms with different non-zero sign.*

Proof. We assume without loss of generality that $C = \{c_1, \dots, c_t\}$ and $\mathcal{C}-C = \{c_{t+1}, \dots, c_m\}$. By Lemma 4.1 applied to the complete binomial network $\tilde{\mathcal{N}}_C$ and the binomial extension $\tilde{\mathcal{N}}_{\mathcal{C}}$, we have

$$D_{\mathcal{N}} = \det(\tilde{\Gamma}_C) \lambda_{[n+t+1, n+m]} + p'(\lambda),$$

where $\lambda_{[n+t+1, n+m]}$ does not divide any term of p' . Hence $\det(\tilde{\Gamma}_C)$ is the coefficient of $\lambda_{[n+t+1, n+m]}$ in $D_{\mathcal{N}}$. The statement now follows by Theorem 2.7. \square

Lemma 4.7. *Let \mathcal{N} be a complete binomial network that is not multistationary. Then $C \in \text{Mult}$ if and only if a term of $D_{\mathcal{N}}$ is a multiple of $\prod_{c_i \in \mathcal{C}-C} \lambda_{n+i}$ with sign different than the sign of multiples of $\lambda_{[n+1, n+t]}$ in $D_{\mathcal{N}}$.*

Proof. Assume $C = \{c_1, \dots, c_t\}$ and $\mathcal{C}-C = \{c_{t+1}, \dots, c_m\}$. We consider \mathcal{N} , $\tilde{\mathcal{N}}_C$ and $\tilde{\mathcal{N}}_{\mathcal{C}}$, and note that each network is a binomial extension of the previous. We apply Lemma 4.1 twice and obtain

$$D_{\mathcal{N}} = \det(\tilde{\Gamma}_C) \lambda_{[n+t+1, n+m]} + p' = (\det(\Gamma) \lambda_{[n+1, n+t]} + p'') \lambda_{[n+t+1, n+m]} + p'.$$

Since \mathcal{N} is not multistationary, then by Theorem 2.7, $\det(\Gamma)$ is non-zero and all terms have the same sign. Hence the coefficient of $\lambda_{[n+1, n+m]}$ is a polynomial where all coefficients have the same sign, σ . Now by Lemma 4.6, $C \in \text{Mult}$ if and only if the coefficient of $\lambda_{[n+t+1, n+m]}$ has terms with different sign. Therefore $C \in \text{Mult}$ if and only if there is a term in p'' with sign $-\sigma$. This proves the lemma. \square

We are now ready to introduce an algorithm to determine the multistationarity structure of a complete binomial network. If the network is multistationary, then all canonical extensions are multistationary and hence Circuits = $\{\emptyset\}$, and Mult is the set of all subsets of \mathcal{C} . If \mathcal{N} is not multistationary, then we use the following algorithm.

Algorithm 4.8.

Input: A complete binomial network \mathcal{N} that is not multistationary.

Output: Circuits for \mathcal{N} .

Procedure:

Initializing step: Circuits = \emptyset .

1. Compute $D_{\mathcal{N}}$ and let σ be the sign of any term divisible by $\lambda_{[n+1, n+m]}$.


2. For every term $T = \alpha\lambda^u$ of $D_{\mathcal{N}}$ with sign $-\sigma$:

2a. Define $C = \{c_i \mid u_{n+i} = 0, i = 1, \dots, m\}$.

2b. Add C to Circuits if no subset of C is already in Circuits, and subsequently remove the supersets of C from Circuits, if any.

Step 2 can be analyzed directly on the exponents u of the terms with sign $-\sigma$: restrict u to the components $n + 1, \dots, n + m$ and choose the corresponding sets C yielding to vectors with maximal support.

Example 4.9. Consider the following complete binomial networks:



We order the species of \mathcal{N}_1 as X_0, X_1, X_2, E, F and for $\mathcal{N}_2, \mathcal{N}_3$ we consider P_0, P_1, S_0, S_1, E, F . Complexes are ordered as they appear in the reaction network from left to right and from up to down. For suitable choices of admissible binomial bases, we obtain the following exponent matrices M_1, M_2, M_3 for the three networks respectively, c.f. (4):

$$M_1^t = \begin{bmatrix} -1 & 1 & 0 & -1 & 1 \\ 0 & -1 & 1 & -1 & 1 \end{bmatrix}, \quad M_2^t = \begin{bmatrix} 0 & 0 & -1 & 1 & -1 & 1 \\ -1 & 1 & 0 & 0 & -1 & 1 \end{bmatrix}, \quad M_3^t = \begin{bmatrix} 0 & 0 & -1 & 1 & -1 & 1 \\ -1 & 1 & 0 & -1 & 0 & 1 \end{bmatrix},$$

and we choose the following matrices of conservation laws:

$$Z_1 = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad Z_2 = Z_3 = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Using this data, we construct the matrices $\tilde{\Gamma}_1, \tilde{\Gamma}_2, \tilde{\Gamma}_3$ defined as in (15) for the largest canonical extensions associated with $\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_3$ and find their determinants $D_{\mathcal{N}_1}, D_{\mathcal{N}_2}$ and $D_{\mathcal{N}_3}$. We have $\sigma = 1$ for all three cases. The sets of monomials with negative coefficients in the determinant of $D_{\mathcal{N}_1}, D_{\mathcal{N}_2}$ and $D_{\mathcal{N}_3}$ are respectively

$$\begin{aligned} A_1 &= \{ \lambda_{\{1,2,3,4,7,8,10,11\}}, \lambda_{\{1,2,3,5,7,8,10,11\}}, \lambda_{\{1,2,4,5,7,8,10,11\}}, \lambda_{\{1,2,4,7,8,9,10,11\}}, \lambda_{\{1,3,4,5,7,8,10,11\}}, \\ & \quad \lambda_{\{2,3,4,5,7,8,10,11\}}, \lambda_{\{2,3,5,6,7,8,10,11\}} \}, \\ A_2 &= \{ \lambda_{\{1,4,5,6,7,8,10,12,13,14\}}, \lambda_{\{2,3,5,6,8,9,10,11,12,14\}} \}, \\ A_3 &= \{ \lambda_{\{1,3,4,5,6,7,9,12,13,14\}}, \lambda_{\{1,3,4,5,6,7,10,12,13,14\}}, \lambda_{\{1,3,4,5,6,8,9,12,13,14\}}, \lambda_{\{1,3,4,5,6,7,9,11,12,14\}}, \\ & \quad \lambda_{\{1,3,4,5,6,7,10,11,12,14\}}, \lambda_{\{1,3,4,5,6,8,9,11,12,14\}}, \lambda_{\{1,3,4,5,7,9,10,11,12,14\}}, \lambda_{\{1,3,4,5,8,9,10,11,12,14\}}, \\ & \quad \lambda_{\{1,3,4,6,7,8,9,12,13,14\}}, \lambda_{\{1,3,4,6,7,8,10,12,13,14\}}, \lambda_{\{1,3,4,6,7,8,9,11,12,14\}}, \lambda_{\{1,3,4,6,7,8,10,11,12,14\}}, \\ & \quad \lambda_{\{1,3,4,6,8,9,10,11,12,14\}}, \lambda_{\{2,4,6,7,8,9,10,11,12,14\}} \}. \end{aligned}$$

According to the algorithm, the monomial $\lambda_{\{1,2,3,4,7,8,10,11\}}$ in A_1 gives rise to the set $C_1 = \{c_6, c_9\}$, while the monomial $\lambda_{\{1,2,4,7,8,9,10,11\}}$ yields $C_2 = \{c_6\}$. Thus C_2 belongs to Circuits₁ while C_1 does not. Proceeding in this way for all monomials, we obtain

$$\begin{aligned} \text{Circuits}_1 &= \{ \{c_6\}, \{c_9\} \} = \{ \{X_0 + E\}, \{X_2 + F\} \}, \\ \text{Circuits}_2 &= \{ \{c_1, c_7\}, \{c_3, c_5\} \} = \{ \{S_0 + E, P_1 + F\}, \{S_1 + F, P_0 + E\} \}, \\ \text{Circuits}_3 &= \{ \{c_7\}, \{c_3, c_5\}, \{c_4, c_5\} \} = \{ \{P_1 + F\}, \{S_1 + F, P_0 + S_1\}, \{S_0 + F, P_0 + S_1\} \}. \end{aligned}$$

We conclude that motifs (g), (i) and (k) in [7] are multistationary, since they are extensions of $\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_3$ respectively, which satisfy the generalized realization condition and the set of inputs of their intermediates belong to the respective multistationarity structures.

As illustrated by these three examples, the elements of the set of circuits might not have the same cardinality.

Remark 4.10. Algorithm 4.8 provides a direct way to detect the sets of complexes that contribute to multistationarity. The method is appealing because, for small networks, the multistationarity structure can be found by simple visual inspection of one multivariate polynomial. The alternative strategy for finding the multistationarity structure would consist in searching for the circuits by computing $\det(\tilde{\Gamma}_C)$ for the canonical extensions. One could start from one of the smallest subsets C of \mathcal{C} and compute $\det(\tilde{\Gamma}_C)$. If this polynomial had terms with different sign or were zero, we would add C to Circuits, and remove C and all its supersets from \mathcal{C} before proceeding in the same way with the next set. The search could alternatively start from the largest subsets C of \mathcal{C} and compute $\det(\tilde{\Gamma}_C)$. If the determinant did not have terms with different sign or is zero, then we would remove all subsets of C from \mathcal{C} . If it had terms with different signs, then we would check the subsets of C with one less element. If none of them were multistationary, then we would add C to Circuits and remove all its subsets from the search.


Going from small to large sets has the advantage of involving the computation of smaller determinants. Our algorithm requires the computation of only one determinant, but it can be large. So, for large networks, it might be advantageous to adopt the search approach starting with small sets described here.

4.3 n -site phosphorylation network

In this section we find the multistationarity structure of the n -site distributive sequential phosphorylation network given as follows (see e.g. [13, 17]):



By removing the intermediates Y_1, \dots, Y_{2n} , the core network associated with the n -site phosphorylation network is



Since $(1, \dots, 1)$ is in the kernel of the stoichiometric matrix of \mathcal{N} , (surj) holds. Further, the rank of \mathcal{N} is n and an admissible binomial basis of the steady state ideal is

$$B := \{ -k_1 x_0 e + k_{2n} x_1 f, \dots, -k_n x_{n-1} e + k_{n+1} x_n f \}.$$

Therefore (rank) also holds and \mathcal{N} is a complete binomial network. By Proposition 5.3 (ii), the generalized realization condition holds for the n -site distributive sequential phosphorylation networks given in (17).

We order the set of species as $X_0, X_1, \dots, X_n, E, F$, and denote the complexes of the core network as

$$c_1 = X_0 + E, \quad \dots \quad c_{n+1} = X_n + E, \quad c_{n+2} = X_n + F, \quad \dots \quad c_{2n+2} = X_0 + F.$$

The largest canonical network consists of the reactions of \mathcal{N} together with the reactions $c_i \rightleftharpoons Y_i$. The matrix $M_\lambda^t \in \mathbb{R}^{n \times (n+3)}$ associated with B and a choice of $Z \in \mathbb{R}^{3 \times (n+3)}$ are

$$M_\lambda^t = \begin{bmatrix} -\lambda_1 & \lambda_2 & & & 0 & -\lambda_{n+2} & \lambda_{n+3} \\ & -\lambda_2 & \lambda_3 & & & -\lambda_{n+2} & \lambda_{n+3} \\ & & & \ddots & & \vdots & \vdots \\ & & & & -\lambda_n & \lambda_{n+1} & -\lambda_{n+2} & \lambda_{n+3} \\ 0 & & & & & & & \end{bmatrix}, \quad Z = \begin{bmatrix} 1 & \cdots & 1 & 0 & 0 \\ 0 & \cdots & 0 & 1 & 0 \\ 0 & \cdots & 0 & 0 & 1 \end{bmatrix}.$$

Proposition 4.11. *For the n -site phosphorylation network with $n \geq 2$, we have*

$$\text{Circuits} = \{\{c_i\} \mid i \neq n, n+1, 2n+1, 2n+2\} = \{X_0+E, \dots, X_{n-2}+E, X_n+F, \dots, X_2+F\}.$$

If $n = 1$, then $\text{Circuits} = \emptyset$, since the largest canonical extension is not multistationary.

Proof. The case $n = 1$ follows by computing the determinant of the largest canonical extension and checking that it is non-zero and that all coefficients have the same sign.

Hence assume that $n \geq 2$. It is enough to first show that $\{c_i\} \in \text{Circuits}$ if $i \neq n, n+1, 2n+1, 2n+2$ and then that $\{c_n, c_{n+1}, c_{2n+1}, c_{2n+2}\} \notin \text{Mult}$.

So let $i \in [n+1]$ and define

$$\Omega(i) = \begin{bmatrix} -1 & 1 & & & & 0 & -1 & 1 & 0 \\ & & \ddots & \ddots & & & \vdots & \vdots & \vdots \\ & & & -1 & 1 & & -1 & 1 & 0 \\ & & & & -1 & 1 & & -1 & 1 & 0 \\ & & & & & \ddots & \ddots & \vdots & \vdots & \vdots \\ 0 & & & & & & -1 & 1 & -1 & 1 & 0 \\ \hline 0 & \cdots & 0 & -1 & 0 & \cdots & 0 & -1 & 0 & 1 \end{bmatrix} \in \mathbb{R}^{(n+1) \times (n+4)},$$

where the -1 in the last row is in the i -th column. Then we have that

$$\tilde{\Gamma}_{\{c_i\}} = \begin{bmatrix} \Omega(i) \text{diag}(\lambda_1, \dots, \lambda_{n+4}) \\ \hline Z \\ \vdots \\ 1 \\ 0 \\ 0 \end{bmatrix} \in \mathbb{R}^{(n+4) \times (n+4)}.$$

For $J \subseteq [n+4]$ of cardinality 3, we denote by $\Omega(i)_J$ the $(n+1) \times (n+1)$ submatrix of $\Omega(i)$ obtained by deleting the columns with index in J . We expand the determinant of

$\tilde{\Gamma}_{\{c_i\}}$ along the last three rows and obtain

$$\begin{aligned}
\det(\tilde{\Gamma}_{\{c_i\}}) &= \sum_{j=1}^{n+1} (-1)^{5n+14+j} \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} \det(\Omega(i)_{\{j,n+2,n+3\}}) \lambda_{[n+4] \setminus \{j,n+2,n+3\}} + \\
&\quad \sum_{j=1}^{n+1} (-1)^{5n+16+j} \begin{vmatrix} 1 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{vmatrix} \det(\Omega(i)_{\{j,n+3,n+4\}}) \lambda_{[n+4] \setminus \{j,n+3,n+4\}} + \\
&\quad (-1)^{6n+18} \begin{vmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{vmatrix} \det(\Omega(i)_{\{n+2,n+3,n+4\}}) \lambda_{[n+4] \setminus \{n+2,n+3,n+4\}} \\
&= \sum_{j=1}^{n+1} (-1)^{n+j} \det(\Omega(i)_{\{j,n+2,n+3\}}) \lambda_{[n+4] \setminus \{j,n+2,n+3\}} + \\
&\quad \sum_{j=1}^{n+1} (-1)^{n+j+1} \det(\Omega(i)_{\{j,n+3,n+4\}}) \lambda_{[n+4] \setminus \{j,n+3,n+4\}} + \\
&\quad \det(\Omega(i)_{\{n+2,n+3,n+4\}}) \lambda_{[n+4] \setminus \{n+2,n+3,n+4\}}.
\end{aligned}$$

We see from this expansion that the coefficient of $\lambda_{[n+4] \setminus \{1,n+2,n+3\}}$ is $(-1)^{n+1} \det(\Omega(i)_{\{1,n+2,n+3\}})$. We have that

$$\Omega(i)_{\{1,n+2,n+3\}} = \left[\begin{array}{cccc|cc} 1 & & & & 0 & 0 \\ -1 & \ddots & & & & \\ & \ddots & \ddots & & & \vdots \\ & & \ddots & \ddots & & \\ & & & \ddots & & \\ 0 & & & -1 & 1 & 0 \\ \hline 0 & \dots & -1 & \dots & 0 & 1 \end{array} \right] \in \mathbb{R}^{(n+1) \times (n+1)}, \quad (19)$$

where the -1 in the last row is in position $i-1$ if $i > 1$ and is not there if $i = 1$. Clearly, $(-1)^{n+1} \det(\Omega(i)_{\{1,n+2,n+3\}}) = (-1)^{n+1}$.

Consider now the coefficient of $\lambda_{[n+4] \setminus \{n+1,n+3,n+4\}}$, which is $(-1)^{n+n+1+1} \det(\Omega(i)_{\{n+1,n+3,n+4\}})$. We have that

$$\Omega(i)_{\{n+1,n+3,n+4\}} = \left[\begin{array}{cccc|cc} -1 & 1 & & & 0 & -1 \\ & -1 & \ddots & & & \vdots \\ & & \ddots & \ddots & & \\ & & & \ddots & & \\ & & & & 1 & \\ 0 & & & & -1 & -1 \\ \hline 0 & \dots & -1 & \dots & 0 & 1 \end{array} \right] \in \mathbb{R}^{(n+1) \times (n+1)},$$

where the -1 in the last row is in position i if $i \leq n$, and there is no -1 if $i = n+1$. Replacing the last row of $\Omega(i)_{\{n+1,n+3,n+4\}}$ with minus the sum of the rows from i to n , we obtain the matrix

$$\left[\begin{array}{cccc|cc} -1 & 1 & & 0 & -1 & \\ & -1 & \ddots & & \vdots & \\ & & \ddots & 1 & -1 & \\ 0 & & & -1 & -1 & \\ \hline & & 0 & & n-i & \end{array} \right] \in \mathbb{R}^{(n+1) \times (n+1)}.$$

The determinant of $\tilde{\Gamma}_C$ is therefore equal to $(-1)^n \lambda_{[n]}$ times the determinant of the inferior diagonal block of size 7×7 of the matrix above. We compute this determinant and obtain the following expression:

$$\begin{aligned} & \lambda_{n+2} \lambda_{n+4} \lambda_{n+6} \lambda_{n+7} (1 + (z_1 + z_2) \lambda_{n+1}) + \lambda_{n+1} \lambda_{n+4} (\lambda_{n+5} \lambda_{n+7} + \lambda_{n+6} \lambda_{n+7} + \lambda_{n+5} \lambda_{n+6}) \\ & + \lambda_{n+1} \lambda_{n+5} \lambda_{n+6} \lambda_{n+7} (1 + z_1 \lambda_{n+4} + z_2 \lambda_{n+2}) + ((n z_1 - z_2) \lambda_{n+1} + n) \lambda_{n+3} \lambda_{n+4} \lambda_{n+5} \lambda_{n+7} \\ & + ((n z_1 + z_1 - z_2) \lambda_{n+1} + n + 1) \lambda_{n+3} \lambda_{n+4} \lambda_{n+5} \lambda_{n+6} + \lambda_{n+4} \lambda_{n+5} \lambda_{n+6} \lambda_{n+7} \\ & + (z_1 \lambda_{n+1} \lambda_{n+2} \lambda_{n+3} + \lambda_{n+1} \lambda_{n+2} + \lambda_{n+1} \lambda_{n+3} + \lambda_{n+2} \lambda_{n+3}) ((n + 1) \lambda_{n+4} \lambda_{n+6} + n \lambda_{n+4} \lambda_{n+7} \\ & + n \lambda_{n+5} \lambda_{n+6} + (n - 1) \lambda_{n+5} \lambda_{n+7}). \end{aligned}$$

Since $n z_1 \geq z_2$ and $n \geq 2$, this determinant is strictly positive. Hence, the determinant of $\tilde{\Gamma}_C$ has sign $(-1)^n$. By Theorem 2.7, we conclude that $\{c_n, c_{n+1}, c_{2n+1}, c_{2n+2}\} \notin \text{Mult}$. \square

In view of Proposition 4.11 and Theorem 3.3(ii) we obtain the following theorem.

Theorem 4.12. *Let $\tilde{\mathcal{N}}$ be an extension of the core n -site phosphorylation network in (18) via the addition of intermediates that satisfies the generalized realization condition. Then $\tilde{\mathcal{N}}$ is multistationary if and only if at least one of $X_0 + E, \dots, X_{n-2} + E, X_n + F, \dots, X_2 + F$ is an input of an intermediate.*

Note that the network in Example 3.6 is an extension of the 2-site phosphorylation network, with set of inputs $C = \{X_0 + E, X_1 + E, X_2 + F, X_1 + F, X_0 + F\}$. This network satisfies the generalized realization condition by Example 5.2. By Theorem 4.12, we conclude that the network is multistationary.

For the n -site phosphorylation network for a fixed n , Algorithm 4.8 requires the computation of one large determinant. The search approach described in Remark 4.10, stops after computing $2n + 14$ determinants, if we start with the small subsets, while it stops after computing

$$\sum_{i=1}^3 \binom{2n-2}{i} + \sum_{i=4}^{2n+2} \binom{2n+2}{i}$$

determinants if we start with large subsets. For example, if $n = 2, 3$, the first approach requires the computation of 18 and 20 determinants, and the second approach requires the computation of 25 and 177 determinants respectively. In these cases, the computation of the determinants takes negligible time, and therefore our algorithm is the fastest strategy.

5 Realization conditions

In this section we discuss methods to decide whether the realization conditions are satisfied. The two realization conditions concern the surjectivity of a rational map on the positive orthant. Specifically, let $\tilde{\mathcal{N}}$ be an extension of \mathcal{N} via the addition of the intermediates Y_1, \dots, Y_m and C' be the set of input complexes that do not belong to the stoichiometric subspace. Consider the following maps from $\mathbb{R}_{>0}^{\tilde{\mathcal{R}}}$:

$$\phi(\kappa) = (\phi_{c \rightarrow c'}(\kappa) \mid c \rightarrow c' \in \mathcal{R}) \in \mathbb{R}_{>0}^{\mathcal{R}}, \quad (20)$$

$$\phi'(\kappa) = \left(\phi(\kappa), \left(\sum_{i \in [m]} \mu_{i,c}(\kappa) \mid c \in C' \right) \right) \in \mathbb{R}_{>0}^{\mathcal{R}} \times \mathbb{R}_{>0}^{C'}. \quad (21)$$

The generalized realization condition is equivalent to the surjectivity of ϕ' and the realization condition to the surjectivity of ϕ . So let $f = (\frac{f_1}{g_1}, \dots, \frac{f_m}{g_m})$ be an arbitrary map from $\mathbb{R}_{>0}^n$ to $\mathbb{R}_{>0}^m$, defined by rational functions $\frac{f_i}{g_i} \in \mathbb{R}(x_1, \dots, x_n)$. Consider the ideal

$I = \langle g_1 y_1 - f_1, \dots, g_m y_m - f_m, 1 - z \prod_{i=1}^m g_i \rangle \subseteq \mathbb{R}[y_1, \dots, y_m, x_1, \dots, x_n, z]$. As discussed in §2.3 of the electronic supplementary material of [8], if $I \cap \mathbb{R}[y_1, \dots, y_m] \neq \{0\}$, then f is not surjective, but the reverse does not necessarily hold.

Another approach is to use Cylindrical Algebraic Decomposition (CAD) [1, 9, 11]. Consider the parametric multivariate system of equations $(x_1, \dots, x_n, z) \in V(I)$ with y_1, \dots, y_m treated as parameters and all variables and parameters constrained to be real and positive. The map f is surjective if and only if this system has at least one positive real solution when evaluated at the sample parameter point of all cells obtained after performing CAD. This approach fully characterizes whether f is surjective, but CAD is computationally expensive. In particular, the number of cells is doubly exponential in the number of variables and parameters, and depends also on the degree and number of polynomials in the system [4, Theorem 5]. Therefore the use of CAD is impractical already in relatively small examples.

Example 5.1. We consider the following core network and its extension via the addition of one intermediate Y :



Using CAD on the system of equations describing the realization condition, we obtain three cells. The sample point of each cell yields a system with infinitely many positive solutions. Therefore the realization condition holds.

In view of the difficulties of checking the realization conditions in practice, we start by understanding how the coefficients $\mu_{i,c}$ are found. Let $\tilde{\mathcal{N}}$ be an extension of \mathcal{N} via the addition of intermediates Y_1, \dots, Y_m . Consider the digraph associated with $\tilde{\mathcal{N}}$ and let $\mathcal{Y}_1, \dots, \mathcal{Y}_t$ denote the vertex sets of the connected components of the subgraph induced by the subset of vertices $\{Y_1, \dots, Y_m\}$. For each non-intermediate complex c and intermediate Y_i , consider the labeled digraph $G_{i,c}$ with vertex set $\mathcal{Y}_\ell \cup \{\star\}$ if $Y_i \in \mathcal{Y}_\ell$. Labeled edges are $Y_i \xrightarrow{\kappa_{Y_i \rightarrow Y_j}} Y_j$ if $Y_i \rightarrow Y_j \in \tilde{\mathcal{R}}$, $\star \xrightarrow{\kappa_{c \rightarrow Y_i}} Y_i$ if $c \rightarrow Y_i \in \tilde{\mathcal{R}}$ and $Y_i \xrightarrow{\beta_i} \star$ with $\beta_i = \sum_{Y_i \rightarrow c'} \kappa_{Y_i \rightarrow c'}$ if $\beta_i \neq 0$. For each vertex v of $G_{i,c}$, define $\Theta_{i,c}(v)$ to be the set of all spanning trees rooted at v , that is, v is the only vertex with zero outdegree. Given a tree τ , let $\pi(\tau)$ be the product of all labels of the edges of τ . Then


$$\mu_{i,c} = \frac{\sum_{\tau \in \Theta_{i,c}(Y_i)} \pi(\tau)}{\sum_{\tau \in \Theta_{i,c}(\star)} \pi(\tau)}. \quad (22)$$

The numerator of $\mu_{i,c}$ is linear in the reaction rate constants of the form $\kappa_{c \rightarrow Y_j}$, and these reaction rate constants do not appear in the denominator. To read more about properties of the $\mu_{i,c}$'s and how to compute them using the Matrix-Tree theorem, see [8].

The components of ϕ and ϕ' might not depend on the reaction rate constants of all reactions in the network. Specifically, from (22) and (9) it follows that $\phi_{c \rightarrow c'}$ depends on $c \rightarrow c'$, if this reaction belongs to $\tilde{\mathcal{R}}$, and possibly on the reactions involving intermediates in the sets \mathcal{Y}_j such that there exists a path from c to c' with all intermediates in \mathcal{Y}_j . So for each reaction, we consider the union of these relevant sets of intermediates \mathcal{Y}_j . Then $\phi_{c_1 \rightarrow c'_1}$ and $\phi_{c_2 \rightarrow c'_2}$ do not depend on a common reaction rate constant if the sets of intermediates corresponding to $c_1 \rightarrow c'_1$ and $c_2 \rightarrow c'_2$ are disjoint. In this way we obtain a partition of $\tilde{\mathcal{R}}$ into subsets of reactions, that is, subnetworks, for which surjectivity of the map ϕ can be checked independently on each smaller network.

We proceed similarly for ϕ' , but in this case the relevant sets of intermediates \mathcal{Y}_j are those for which there exists a path from c to at least one $Y_i \in \mathcal{Y}_j$ (or equivalently, $\mu_{i,c} \neq 0$).

Example 5.2. We consider the generalized realization condition for Example 3.6. By the discussion above, this condition needs to be checked independently on the following three subnetworks:




For the three subnetworks the generalized realization condition holds due to Proposition 5.3(ii) below.

We next show that the realization condition holds for specific classes of intermediates without the need to do any extra computations.

Proposition 5.3. *The realization condition holds for the following types of extension networks via the addition of intermediates Y_1, \dots, Y_m .*


(i)



with an arbitrary digraph structure among the complexes c, Y_1, \dots, Y_m such that there is a path from c to all Y_i , and where some reactions with label ℓ_i might not exist.

(ii) $c \longleftrightarrow Y_1 \longleftrightarrow Y_2 \longleftrightarrow \dots \longleftrightarrow Y_m \longleftrightarrow c'$, provided $\{Y_1, \dots, Y_m\}$ is a set of intermediates, and where \longleftrightarrow means the reaction can be either irreversible or reversible. These networks satisfy also the generalized realization condition. Further, a union of subnetworks of this form such that the sets of intermediates of each subnetwork do not intersect, satisfies also the generalized realization condition.

(iii)



with an arbitrary digraph structure among the complexes c_0, Y_1, \dots, Y_m such that there exists a directed path from c_0 to Y_m .

Proof. (i) The realization condition is equivalent to the scalar-valued map $\ell_0 + \sum_{i=1}^m \ell_i \mu_{i,c}$ being surjective. This map is linear in $\ell_0, \kappa_{c \rightarrow Y_1}, \dots, \kappa_{c \rightarrow Y_m}$ (some might be zero, but at least one is non-zero). Hence the statement is clear.

(ii) We start with the case with only one such block. We write

$$\tilde{\mathcal{N}}: c \xrightleftharpoons[\kappa_2]{\kappa_1} Y_1 \xrightleftharpoons[\kappa_4]{\kappa_3} Y_2 \xrightleftharpoons[\kappa_6]{\kappa_5} \dots \xrightleftharpoons[\kappa_{2m}]{\kappa_{2m-1}} Y_m \xrightleftharpoons[\kappa_{2m+2}]{\kappa_{2m+1}} c'. \quad (23)$$

If not all reactions are reversible, then we assume that the reaction of the core network is $c \rightarrow c'$. This means that all reactions with label with odd subindex are present, and the reverse reactions might or might not be present.

We can assume without loss of generality that neither $c \rightarrow c'$ nor $c' \rightarrow c$ belong to $\tilde{\mathcal{N}}$ (if a map is surjective between two positive orthants, adding an extra variable that sums to one component preserves surjectivity).


We have that $\phi_{c \rightarrow c'}(\kappa) = \kappa_{2m+1}\mu_{m,c}(\kappa)$ and $\phi_{c' \rightarrow c}(\kappa) = \kappa_2\mu_{1,c'}(\kappa)$ (the latter being zero in the irreversible case). Throughout we assume that the set C' used to define ϕ' equals $\{c, c'\}$. This is the worst case scenario.

We show by induction on m that this network satisfies the generalized realization condition. For $m = 1$, if all reverse reactions are present we have that

$$\phi'(\kappa_1, \kappa_2, \kappa_3, \kappa_4) = (\phi_{c \rightarrow c'}, \phi_{c' \rightarrow c}, \mu_{1,c}, \mu_{1,c'}) = \left(\frac{\kappa_1\kappa_3}{\kappa_2+\kappa_3}, \frac{\kappa_2\kappa_4}{\kappa_2+\kappa_3}, \frac{\kappa_1}{\kappa_2+\kappa_3}, \frac{\kappa_4}{\kappa_2+\kappa_3} \right).$$

A missing reverse reaction corresponds to setting the reaction rate constant equal to zero, and projecting ϕ' away from the components that become zero. We confirm using CAD that this map is surjective when restricted to the positive orthants, in the four scenarios obtained by considering none, one or both reverse reactions.

Assume now that (23) satisfies the generalized realization condition for $m - 1$. We view $\tilde{\mathcal{N}}$ as an extended network of



via the addition of one intermediate Y_1 . If we let $\tilde{\kappa} = (\kappa_1, \kappa_2, \kappa_3, \kappa_4)$, this gives rise to the following relevant functions

$$\tilde{\mu}_{1,c}(\tilde{\kappa}) = \frac{\kappa_1}{\kappa_2+\kappa_3}, \quad \tilde{\mu}_{1,Y_2}(\tilde{\kappa}) = \frac{\kappa_4}{\kappa_2+\kappa_3}, \quad \bar{\kappa}_1 = \kappa_3\tilde{\mu}_{1,c}(\tilde{\kappa}), \quad \bar{\kappa}_2 = \kappa_2\tilde{\mu}_{1,Y_2}(\tilde{\kappa}). \quad (24)$$

By the case $m = 1$, the right-hand sides of these equalities define a surjective map when restricted to the positive orthant (by omitting the zero components if some reaction rate constants are set to zero). In turn, $\overline{\mathcal{N}}$ is an extended network of $c \rightleftharpoons c'$ via the addition of the intermediates Y_2, \dots, Y_m . By the induction hypothesis, $\overline{\mathcal{N}}$ satisfies the generalized realization condition. Let $\bar{\mu}_{i,c}, \bar{\mu}_{i,c'}$ for $i = 2, \dots, m$ correspond to this extension.

Recall that $\mu_{i,c}$ and $\mu_{i,c'}$ are the coefficients of x^c and $x^{c'}$ respectively after writing y_1, \dots, y_m in terms of x by solving the steady state equations corresponding to the intermediates. This system can be solved iteratively, by first finding y_1 and then y_2, \dots, y_m . If we let $\varphi(\kappa) = (\kappa_3\tilde{\mu}_{1,c}(\tilde{\kappa}), \kappa_2\tilde{\mu}_{1,Y_2}(\tilde{\kappa}), \kappa_5, \kappa_6, \dots, \kappa_{2m+2})$, it follows that

$$\mu_{i,c} = \bar{\mu}_{i,c}(\varphi(\kappa)), \quad \mu_{i,c'} = \bar{\mu}_{i,c'}(\varphi(\kappa)), \quad \text{for } i = 2, \dots, m.$$

For $i = 1$, iterative elimination of y_1 and $y_2 = \bar{\mu}_{2,c}(\varphi(\kappa))x^c + \bar{\mu}_{2,c'}(\varphi(\kappa))x^{c'}$ gives that

$$y_1 = \tilde{\mu}_{1,Y_2}(\tilde{\kappa})y_2 + \tilde{\mu}_{1,c}(\tilde{\kappa})x^c = \tilde{\mu}_{1,Y_2}(\tilde{\kappa})\bar{\mu}_{2,c'}(\varphi(\kappa))x^{c'} + (\tilde{\mu}_{1,Y_2}(\tilde{\kappa})\bar{\mu}_{2,c}(\varphi(\kappa)) + \tilde{\mu}_{1,c}(\tilde{\kappa}))x^c.$$

Hence

$$\mu_{1,c}(\kappa) = \tilde{\mu}_{1,Y_2}(\tilde{\kappa})\bar{\mu}_{2,c}(\varphi(\kappa)) + \tilde{\mu}_{1,c}(\tilde{\kappa}), \quad \mu_{1,c'}(\kappa) = \tilde{\mu}_{1,Y_2}(\tilde{\kappa})\bar{\mu}_{2,c'}(\varphi(\kappa)).$$

Therefore $\phi'(\kappa) = (\kappa_{2m+1}\mu_{m,c}(\kappa), \kappa_2\mu_{1,c'}(\kappa), \sum_{i=1}^m \mu_{i,c}(\kappa), \sum_{i=1}^m \mu_{i,c'}(\kappa))$ can be written as

$$\begin{aligned} \phi'(\kappa) &= (\kappa_{2m+1}\bar{\mu}_{m,c}(\varphi(\kappa)), \kappa_2\tilde{\mu}_{1,Y_2}(\tilde{\kappa})\bar{\mu}_{2,c'}(\varphi(\kappa)), \sum_{i=2}^m \bar{\mu}_{i,c}(\varphi(\kappa)), \sum_{i=2}^m \bar{\mu}_{i,c'}(\varphi(\kappa))) \\ &\quad + (0, 0, \tilde{\mu}_{1,Y_2}(\tilde{\kappa})\bar{\mu}_{2,c}(\varphi(\kappa)) + \tilde{\mu}_{1,c}(\tilde{\kappa}), \tilde{\mu}_{1,Y_2}(\tilde{\kappa})\bar{\mu}_{2,c'}(\varphi(\kappa))). \end{aligned}$$

Let $(k_1, k_2, r_c, r_{c'}) \in \mathbb{R}_{>0}^4$, with $k_2 = 0$ in the irreversible case and $r_{c'} = 0$ if c' is not an input of any intermediate. Write $r_c = r_{c,1} + r_{c,2}$, $r_{c'} = r_{c',1} + r_{c',2}$ such that

$r_{c',2} < r_{c,2}$ and $r_{c,1}, r_{c,2}, r_{c',1}, r_{c',2} > 0$ ($= 0$ as appropriate). We want to show that $(k_1, k_2, r_c, r_{c'}) = \phi'(\kappa)$ for some κ . First note that by the induction hypothesis, we can find $\bar{\kappa} = (\bar{\kappa}_1, \bar{\kappa}_2, \bar{\kappa}_3, \dots, \bar{\kappa}_{2m+2})$ such that

$$(k_1, k_2, r_{c,1}, r_{c',1}) = \left(\kappa_{2m+1} \bar{\mu}_{m,c}(\bar{\kappa}), \bar{\kappa}_2 \bar{\mu}_{2,c'}(\bar{\kappa}), \sum_{i=2}^m \bar{\mu}_{i,c}(\bar{\kappa}), \sum_{i=2}^m \bar{\mu}_{i,c'}(\bar{\kappa}) \right).$$

By the last two equalities in (24), the decomposition of $\phi'(\kappa)$ above and the definition of φ , all we need is to show that there exists $\tilde{\kappa} = (\tilde{\kappa}_1, \tilde{\kappa}_2, \tilde{\kappa}_3, \tilde{\kappa}_4)$ such that

$$\begin{aligned} \bar{\kappa}_1 &= \kappa_3 \tilde{\mu}_{1,c}(\tilde{\kappa}), & \bar{\kappa}_2 &= \kappa_2 \tilde{\mu}_{1,Y_2}(\tilde{\kappa}), \\ r_{c,2} &= \tilde{\mu}_{1,Y_2}(\tilde{\kappa}) \bar{\mu}_{2,c}(\bar{\kappa}) + \tilde{\mu}_{1,c}(\tilde{\kappa}), & r_{c',2} &= \tilde{\mu}_{1,Y_2}(\tilde{\kappa}) \bar{\mu}_{2,c'}(\bar{\kappa}). \end{aligned}$$

This gives in particular that $\bar{\kappa} = \varphi(\kappa)$. Since $\bar{\kappa}$ has now been fixed, we want

$$\bar{\kappa}_1 = \kappa_3 \tilde{\mu}_{1,c}(\tilde{\kappa}), \quad \bar{\kappa}_2 = \kappa_2 \tilde{\mu}_{1,Y_2}(\tilde{\kappa}), \quad \tilde{\mu}_{1,Y_2}(\tilde{\kappa}) = \frac{r_{c',2}}{\bar{\mu}_{2,c'}(\bar{\kappa})} > 0, \quad \tilde{\mu}_{1,c}(\tilde{\kappa}) = r_{c,2} - r_{c',2} > 0.$$

Since the generalized realization condition holds for $m = 1$, there exist $\kappa_1, \dots, \kappa_4$ such that this system holds (or the equivalent system if some reactions are irreversible). This finishes the proof for the case where there is only one block.

If there are several blocks with the same structure as (23), then we simply need to notice that ϕ' can be written as the Cartesian product of the corresponding map for each block, and $\sum_{i=1}^m \mu_{i,c}$ can be split as a sum of the $\mu_{i,c}$'s of each block. Since the generalized realization condition holds for each block, it also holds for the whole network by splitting r_c accordingly for each complex c .

(iii) The core network has p reactions $c_0 \rightarrow c_1, \dots, c_0 \rightarrow c_p$. We have $\phi_{c_0 \rightarrow c_i}(\kappa) = \ell_i \mu_{m,c_0}$. The denominator of μ_{m,c_0} is a multiple of $\sum_{i=1}^p \ell_i$ and $\mu_0 = (\sum_{i=1}^p \ell_i) \mu_{m,c_0}$ does not depend on any ℓ_i . Note that the scalar-valued function μ_0 is positive and linear in $(\kappa_{c_0 \rightarrow Y_1}, \dots, \kappa_{c_0 \rightarrow Y_m})$. Hence by varying the reaction rate constants different from ℓ_i , μ_0 covers $\mathbb{R}_{>0}$. With this we have that given $k_1, \dots, k_p > 0$, we define $\ell_i = k_i$ and choose the rest of reaction rate constants such that $\mu_0 = \sum_{i=1}^p k_i$. Then $\phi_{c_0 \rightarrow c_i}(\kappa) = k_i$, showing that ϕ is surjective. \square

Acknowledgements. This work has been supported by the Independent Research Fund of Denmark. We thank Martin Helmer and Angélica Torres for comments on a preliminary version of this manuscript.

References

- [1] S. Basu, R. Pollack, and M. F. Coste-Roy. *Algorithms in real algebraic geometry*, volume 10. Springer Science & Business Media, 2007.
- [2] R. Bradford, J. H. Davenport, M. England, H. Errami, V. Gerdt, D. Grigoriev, C. Hoyt, M. Kořta, O. Radulescu, T. Sturm, and A. Weber. A case study on the parametric occurrence of multiple steady states. In *Proceedings of the International Symposium on Symbolic and Algebraic Computation, ISSAC*, pages 45–52. Association for Computing Machinery, 2017.
- [3] D. Eisenbud and B. Sturmfels. Binomial ideals. *Duke Math. J.*, 84(1):1–45, 1996.
- [4] M. England, R. Bradford, and J. H. Davenport. Improving the use of equational constraints in cylindrical algebraic decomposition. In *Proceedings of the International Symposium on Symbolic and Algebraic Computation, ISSAC*, pages 165–172. Association for Computing Machinery, 2015.

- [5] M. Feinberg. Lectures on chemical reaction networks. Available online at <http://www.crnt.osu.edu/LecturesOnReactionNetworks>, 1980.
- [6] M. Feinberg. The existence and uniqueness of steady states for a class of chemical reaction networks. *Arch. Ration. Mech. Anal.*, 132(4):311–370, 1995.
- [7] E. Feliu and C. Wiuf. Enzyme-sharing as a cause of multi-stationarity in signalling systems. *J. R. Soc. Interface*, 9(71):1224–1232, 2012.
- [8] E. Feliu and C. Wiuf. Simplifying biochemical models with intermediate species. *J. R. Soc. Interface*, 10(87):20130484, 2013.
- [9] J. Gerhard, D. Jeffrey, and G. Moroz. A package for solving parametric polynomial systems. *ACM Commun. Comput. Algebra*, 43(3/4):61–72, 2010.
- [10] J. Gunawardena. Chemical reaction network theory for in-silico biologists. Available online at <http://vcp.med.harvard.edu/papers/crnt.pdf>, 2003.
- [11] D. Lazard and F. Rouillier. Solving parametric polynomial systems. *J. Symb. Comput.*, 42(6):636–667, 2007.
- [12] S. Müller, E. Feliu, G. Regensburger, C. Conradi, A. Shiu, and A. Dickenstein. Sign conditions for injectivity of generalized polynomial maps with applications to chemical reaction networks and real algebraic geometry. *Found. Comput. Math.*, 16(1):69–97, 2016.
- [13] M. Pérez Millán, A. Dickenstein, A. Shiu, and C. Conradi. Chemical reaction systems with toric steady states. *Bull. Math. Biol.*, 74(5):1027–1065, 2012.
- [14] A. H. Sadeghimanesh and E. Feliu. Gröbner bases of reaction networks with intermediate species. *arXiv: 1804.01381*, 2018.
- [15] M. Sáez, C. Wiuf, and E. Feliu. Graphical reduction of reaction networks by linear elimination of species. *J. Math. Biol.*, 74(1):195–237, 2017.
- [16] C. H. Schilling, D. Letscher, and Palsson B. Ø. Theory for the systemic definition of metabolic pathways and their use in interpreting metabolic function from a pathway-oriented perspective. *J. Theor. Biol.*, 203(3):229–248, 2000.
- [17] L. Wang and E. D. Sontag. On the number of steady states in a multiple futile cycle. *J. Math. Biol.*, 57(1):29–52, 2008.