

Parametric heating in a 2D periodically-driven bosonic system: Beyond the weakly-interacting regime

T. Boulier,^{1,2,*} J. Maslek,^{1,†} M. Bukov,^{3,†} C. Bracamontes,¹ E. Magnan,^{1,2} S. Lellouch,⁴ E. Demler,⁵ N. Goldman,⁶ and J. V. Porto^{1,‡}

¹*Joint Quantum Institute, National Institute of Standards and Technology and the University of Maryland, College Park, Maryland 20742 USA*

²*Laboratoire Charles Fabry, Institut d'Optique Graduate School, CNRS, Université Paris-Saclay, 91127 Palaiseau cedex, France*

³*Department of Physics, University of California Berkeley, CA 94720, USA*

⁴*Laboratoire de Physique des Lasers, Atomes et Molécules, Université Lille 1 Sciences et Technologies, CNRS; F-59655 Villeneuve d'Ascq Cedex, France*

⁵*Department of Physics, Harvard University, Cambridge, MA 02138, USA*

⁶*Center for Nonlinear Phenomena and Complex Systems, Université Libre de Bruxelles, CP 231, Campus Plaine, B-1050 Brussels, Belgium*

(Dated: August 22, 2022)

We experimentally investigate the effects of parametric instabilities on the short-time heating process of periodically-driven bosons in 2D optical lattices with a continuous transverse (tube) degree of freedom. We analyze three types of periodic drives: (i) linear along the x -lattice direction only, (ii) linear along the lattice diagonal, and (iii) circular in the lattice plane. In all cases, we demonstrate that the BEC decay is dominated by the emergence of unstable Bogoliubov modes, rather than scattering in higher Floquet bands, in agreement with recent theoretical predictions. The observed BEC depletion rates are much higher when shaking both along x and y directions, as opposed to only x or only y . We also report an explosion of the decay rates at large drive amplitudes, and suggest a phenomenological description beyond Bogoliubov theory. In this strongly-coupled regime, circular drives heat faster than diagonal drives, which illustrates the non-trivial dependence of the heating on the choice of drive.

An area of increasing interest in ultracold atoms concerns the engineering of novel states of matter using highly-controllable optical lattices [1]. In this context, a promising approach relies on applying time-periodic modulation to the system, in view of designing an effective time-independent Hamiltonian featuring the desired properties [2–4]. This *Floquet engineering* has emerged as a promising and conceptually straightforward way to expand the quantum simulation toolbox, enabling appealing features such as suppressed [5, 6] or laser-assisted [7] tunneling in optical lattices, enhanced magnetic correlations [8], state-dependent lattices [9], sub-wavelength optical lattices [10], as well as synthetic dimensions [11, 12], synthetic gauge fields [13, 14] and topological band structures [15].

Despite these promising applications, progress in Floquet engineering for interacting systems has been hindered by heating due to uncontrolled energy absorption. This heating, which stems from a rich interplay between the periodic drive and inter-particle interactions, is a particularly challenging problem in interacting systems, where it is known to occur due to proliferation of resonances between many-body Floquet states, not captured by the inverse-frequency expansion [4, 16]. This constrains the applicability of Floquet engineering to regimes where heating is slower than the engineered dynamics [9, 17–21]. A deeper understanding of the underlying processes is essential to determine stable regions of the (large) parameter space, where the system is amenable to

Floquet engineering. Additionally, interaction-mediated heating is itself an interesting nontrivial quantum many-body process. Energy absorption and entanglement production in periodically driven systems have recently been the focus of theoretical studies [16, 22–29] and experimental investigations [5, 9, 20, 21]. It was predicted that, whenever the drive frequency is larger than all single-particle energy scales of the problem, heating succumbs to a stable long-lived prethermal steady state, before it can occur at exponentially-long times [25, 30–35]. However, it is unclear whether this physics is accessible in interacting bosonic experiments

A perturbative approach to understanding drive-induced heating is to analyze the underlying two-body scattering processes using Fermi's Golden rule (FGR) [21, 23, 36–38]. In the weakly-interacting limit, interactions provide a small coupling between noninteracting Floquet states. However, Floquet states cannot be treated as noninteracting when the Floquet-modified *excitation spectrum* is itself unstable [22, 24–26]. These instabilities indicate that heating can occur on a shorter timescale than expected from scattering theory alone.

For Bose-Einstein condensates (BEC) in optical lattices, increased decay rates arise due to the emergence of unstable collective modes. The resulting parametric heating can be described using a Floquet-Bogoliubov-deGennes (FBdG) approach [24], and the short-time dynamics is dominated by an exponential growth of the unstable excited modes in the BEC. The depletion time of

the condensate fraction provides an experimental window to observe this and related effects. Qualitatively different behavior is expected between scattering and parametric instability rates; most notably, different power laws as a function of the interaction strength, tunneling rate and drive amplitude.

We experimentally explore these predictions in a 2D lattice subject to 1D and 2D periodic drives, by measuring decay of the BEC condensed fraction. We provide strong experimental evidence that parametric instabilities dominate the short-time dynamics over FGR-type scattering processes, which are responsible for long-time thermalization [21]. Our experiment reveals effects beyond Floquet-Bogoliubov predictions, and points out limitations in the applicability of FBdG theory.

The experiments are performed on a BEC of ^{87}Rb atoms loaded into a square 2D optical lattice [39, 40] with principal axes along x and y , formed by two pairs of counter-propagating laser beams with wavelength $\lambda = 814$ nm. The total atom number is $N \simeq 10^5$ ($\pm 20\%$ systematic uncertainty). Two piezo-actuated mirrors [41] sinusoidally translate the lattice along x and y with any desired amplitude, relative phase and angular frequency: $\mathbf{r}(t) = \{\Delta x \sin(\omega t), \Delta y \sin(\omega t + \phi)\}$. We consider the effect of three drive trajectories on the decay rate: translation along x only ($\Delta y = 0$), diagonal translation along x and y ($\Delta y = \Delta x$ and $\phi = 0$), and circular translation ($\Delta y = \Delta x$ and $\phi = \pi/2$). Therefore, the driving is 1D (x only) or 2D (diagonal or circular), in a 2D system (2D array of tubes), as shown on Fig. 1. We express the amplitude Δx in terms of the drive-induced maximum effective energy offset between neighboring lattice sites in the co-moving frame, $K_0 = \Delta E/\hbar\omega$, where $\Delta E = m\omega^2 a \Delta x$, a is the lattice spacing and m is the ^{87}Rb mass. The physical displacement is $\Delta x = \hbar K_0 / a\omega m$. The lattice depth V_0 is held constant during shaking, and is measured in units of lattice recoil energy $E_R = \hbar^2/m\lambda^2$. The lattice tunneling energy $\hbar J$ and the interaction strength g are controlled via V_0 . The value of $J(V_0)$ and $g(V_0)$ are calculated from the band structure, peak atom density and scattering length (see appendix). This results in the following periodically-driven Bose-Hubbard Hamiltonian:

$$\hat{H}(t) = \int_z \sum_{i,j} \left\{ -\hbar J \left(\hat{a}_{i+1,j}^\dagger(z) \hat{a}_{i,j}(z) + \hat{a}_{i,j+1}^\dagger(z) \hat{a}_{i,j}(z) + \text{h.c.} \right) - \hat{a}_{i,j}^\dagger(z) \frac{\hbar^2 \partial_z^2}{2m} \hat{a}_{i,j}(z) + \frac{\hbar U}{2} \hat{a}_{i,j}^\dagger(z) \hat{a}_{i,j}^\dagger(z) \hat{a}_{i,j}(z) \hat{a}_{i,j}(z) + \hbar\omega K_0 (i \sin(\omega t) + \kappa j \sin(\omega t + \phi)) \hat{a}_{i,j}^\dagger(z) \hat{a}_{i,j}(z) \right\} \quad (1)$$

where $\hat{a}_{i,j}^{(\dagger)}(z)$ is the annihilation (creation) operator at lattice site (i, j) and transverse position z , and $\kappa = 0$ for x -only and $\kappa = 1$ for 2D drives. The interaction U is defined such as $U/V \sum_{i,j} \int_z \langle \hat{a}_{i,j}^\dagger(z) \hat{a}_{i,j}(z) \rangle = g$ with V the volume of the system.




FIG. 1. **Schematic of the lattice driving.** (up) Lattice translation is performed along the x -direction (green), the diagonal (blue), and in circles (red). The normalized drive amplitude projected along the x -axis, K_0 , is used to characterize the drive strength for all trajectories. (down) Example of periodic drive for a circular driving.

In order to avoid micro-motion effects during a drive period [10], the experiments are performed at integer multiples of the period $T = 2\pi/\omega$ (see appendix). The drive amplitude is ramped up smoothly [10, 42, 43] in a fixed time (minimum 2 ms) corresponding to an integer number of periods [Fig. 1]. The shaking is then held at a constant amplitude for a time τ . Finally, the amplitude is ramped down to zero in a few periods. We checked the effect of adiabaticity of this procedure (or lack thereof) for the three drive trajectories, and describe it in the Appendix (see appendix). Once the lattice is at rest, we turn it off in 300 μs to determine the atomic distribution. We use absorption imaging after time of flight to measure the condensate fraction as a function of τ .

For most conditions, the condensate decay agrees with an exponential decay ($N(t) = N(0)e^{-\Gamma_{\text{cf}}t}$), whose rate Γ_{cf} we extract from a least-square fit (see appendix). We

measure Γ_{cf} for the three drives at different values of ω , K_0 and V_0 . FBdG predicts an undamped parametric instability, characterized by exponential growth of unstable modes, i.e. *accelerated* condensate loss. This behavior is inconsistent with the measured exponential decay of the BEC. Hence, the undamped FBdG regime does not last long compared to the typical BEC lifetime for our parameters, and interactions between the excited unstable modes and the BEC play a significant role in the observed heating process. Nonetheless, as we discuss below, the magnitude and scaling of Γ_{cf} are well captured by a FBdG description.

Since the Floquet-renormalized hopping is $J_{\text{eff}} = J\mathcal{J}_0(K_0)$ ($\mathcal{J}_\nu(K_0)$ is the ν -th order Bessel function), $J_{\text{eff}} < 0$ for $K_0 > 2.4$ and the lowest Floquet band is inverted [5]; the BEC then becomes dynamically unstable at $\mathbf{q} = (0, 0)$ [44], but a stable equilibrium occurs at the band edge and a sudden change in the stability point can trigger a dynamical transition [45] between the two equilibrium configurations. In the band inverted regime, the stable quasi-momenta are $\mathbf{q} = (\pm\pi, 0)$ for a 1D drive along x , and at $\mathbf{q} = (\pm\pi, \pm\pi)$ for a 2D drive along x and y , where the components of the crystal momentum \mathbf{q} are measured in units of the inverse lattice spacing a^{-1} . FBdG assumes an initial macroscopic occupation of these modes [24]. Unless stated otherwise, for data taken at $K_0 > 2.4$ we first accelerate the BEC to the appropriate stable point while simultaneously turning on the Floquet drive (see appendix).

Lattice depth scans: $\Gamma_{\text{cf}}(V_0)$ – A major difference between FGR and FBdG theory is the scaling of the decay rate with the hopping J and interaction strength g . Whereas FGR predicts $\Gamma_{\text{cf}} \propto (gJ)^2$, the parametric instability rate is expected to be linear ($\Gamma_{\text{cf}} \propto gJ$) [24]. Figure 2 shows the condensed fraction decay rate Γ_{cf} measured at different V_0 (between $7.3 E_R$ and $17.0 E_R$) and ω , plotted as a function of gJ/ω , for the 2D-diagonal drive and 1D x -only drive. The solid lines show the FBdG theory, and the dashed lines are linear fits to the data. For the x -only drive, the magnitude and slope extracted from the data are well described by the FBdG theory, and while some deviations appear for the 2D drive, both results are clearly inconsistent with a quadratic dependence.

Amplitude scans: $\Gamma_{\text{cf}}(K_0)$ – Figure 3 shows the decay rate as a function of the drive amplitude K_0 , at lattice depth $V_0 = 11 E_R$ (which gives $J = 2\pi \times 50$ Hz, $g = 2\pi \times 700$ Hz and a gap $\Delta E_v/h = 21$ kHz to the next vibrational level) and a drive frequency $\omega = 2\pi \times 2.5$ kHz. The same data is shown on a full range (top) and zoomed in (bottom).

Consider first the x -only drive. The instability growth rate predicted from FBdG theory [16, 46] $\Gamma_{\text{mum}} = 8J\mathcal{J}_2(K_0)g/\omega$, which we detail in the Appendix (see appendix), agrees with the measured decay rates [Fig. 3]. The appearance of the Bessel function $\mathcal{J}_2(K_0)$ in the ex-




FIG. 2. $\Gamma_{\text{cf}}(Jg/\omega)$: Measured decay rates for $K_0 = 2.1$, $\omega = 2\pi \times 4$ kHz (filled circles) and $\omega = 2\pi \times 2.5$ kHz (empty circles) and various lattice depths, compared to FBdG theory. The horizontal axis is $Jg/(2 \times \omega)$, since FBdG theory predicts the modes growth rate to be linear in J , g and $1/\omega$, as derived in the Appendix (see appendix). The dashed lines are linear fits to the data.

pression for the rate can be traced back to the parametric resonance condition, requiring the excitation spectrum to match the drive frequency, cf. the Appendix. In contrast, the FGR scattering approach prediction is too low by a factor of about 30, and the predicted scaling, $\propto |\mathcal{J}_2(K_0)|^2$ to leading order, does not describe the data as well. The agreement with the FBdG theory, despite evidence for effects beyond simple undamped parametric instability, is consistent over a range of parameter space in K_0 , ω and V_0 . As expected, the decay dramatically increases when $J_{\text{eff}} < 0$ ($K_0 > 2.4$) for $\mathbf{q} = (0, 0)$, while it is partially stabilized when accelerating the BEC to $\mathbf{q} = (\pi, 0)$. We note that significant heating occurs during the drive turn-on and acceleration phase for the $\mathbf{q} = (\pi, 0)$ data, resulting in partial BEC losses.

For the two 2D drives (circular and diagonal) we observe decay rates that follow roughly the same functional form as the 1D drive, but about $3 \times$ larger. Additionally, the sudden increase in the decay rate Γ_{cf} for the 2D drives consistently occurs at a critical amplitude K_0^c below 2.4. At $11 E_R$, $K_0^c \simeq 2.15$ [Fig. 3]. Above K_0^c , both 2D rates increase dramatically beyond any prediction and the circular rate increases faster than the diagonal rate. This drive dependence and drastic rate increase for $K_0^c < K_0 < 2.4$ suggest effects beyond FBdG theory, distinct from the simple parametric instability, and are discussed at the end of this Letter. For $K_0 \gtrsim 2.4$, the rates are essentially unmeasurable since $\Gamma_{\text{cf}} > \omega$. As with the 1D drive, accelerating the BEC to $\mathbf{q} = (\pi, \pi)$ partially stabilizes the decay, just enough to be measurable.

Frequency scans – $\Gamma_{\text{cf}}(\omega)$: FBdG theory predicts distinct behavior at low and high drive frequencies. In the low-frequency regime, the momenta of the maximally-unstable mode \mathbf{q}_{mum} evolve as one increases the drive fre-




FIG. 3. $\Gamma_{\text{cf}}(K_0)$: Comparison between circular (red), diagonal (blue) and x -only (green) drives for $V_0 = 11 E_R$ and $\omega = 2\pi \times 2.5$ kHz. The rates are in units of $J = 2\pi \times 50$ Hz, shown full scale (top) and zoomed in (bottom). The Floquet-Bogoliubov-deGennes theory [24, 46] Γ_{mum} is shown for each drive as solid lines, and the 1D FGR-based scattering theory is shown as the green dashed line (zoom only). Filled circles indicate data taken at $\mathbf{q} = (0, 0)$, while open circles indicate data taken at $\mathbf{q} = (\pi, \pi)$ [see text] to keep the BEC in the stable region of the band (illustrated with the bottom plot cartoon). A dramatic increase in the rate occurs at $K_0^c \gtrsim 2.4$ for the circular and diagonal drives, highlighted by the light red region.

quency, until it saturates at the Bogoliubov band edge at $\mathbf{q}_{\text{mum}} = \{(\pi, 0)\}, \{(\pi, 0), (0, \pi)\}$, and $\{(\pi, \pi)\}$ for x -only, circular and diagonal drive, respectively. The periodic drive results in interference which depends on the relative phase of the two components of the drive, leading to different most unstable modes for the three shaking geometries. The saturation frequency $\omega_c = E_{\text{eff}}^{\text{Bog}}(\mathbf{q}_{\text{mum}})$ marks the onset of the high-frequency regime [24]. Note that unlike in 1D lattices [24] the energy of the maximally unstable mode can be lower than the full effective bandwidth (see appendix). The rate is predicted to increase quasi-linearly for $\omega \leq \omega_c$ while $\Gamma_{\text{mum}} \propto \omega^{-1}$ for

$\omega \geq \omega_c$ [21, 24], resulting in a cusp in the rate at ω_c . A detailed derivation of the expressions for the most unstable modes and the instability rates are given in the Appendix.

Figure 4 shows the experimental values for $\Gamma_{\text{cf}}(\omega)$ compared with FBdG theory. Since there is no observed rate explosion for the x -only drive, we use the cusp in $\Gamma_{\text{cf}}^x(\omega)$ to calibrate our experimental value for g , which agrees to within 20% with an estimate calculated from the lattice parameters (see appendix). Using this value of g , the prediction for the diagonal drive ω_c^{Diag} matches the experiment. While FBdG theory predicts the same ω_c for circular and x -only drives, the measured cusp for the circular drive lies between the cusps of the two linear drives.

The observed behavior qualitatively fits FBdG theory, with rates generally higher than predicted for $\omega \sim \omega_c$. For the x -only drive, the measured rates are slightly above the prediction below $2\pi \times 1.5$ kHz, and the agreement is excellent above $2\pi \times 1.5$ kHz, as observed with $\Gamma_{\text{cf}}(K_0)$ at $2\pi \times 2.5$ kHz [Fig. 3]. The 2D rates show a larger discrepancy at low frequencies, and a decent quantitative agreement for $\omega \gtrsim 2\pi \times 2$ kHz. This is related to the rates explosion appearing for $K_0 > K_0^c$. As we discuss below, the observed value of K_0^c increases with the frequency. This implies a similar rate explosion should happen when decreasing ω at fixed K_0 . This is especially visible with the diagonal drive [Fig. 4]: for $\omega < 2\pi \times 1$ kHz the data abruptly departs from the prediction. This increased rate at low frequencies for 2D drives is likely responsible for the discrepancy between experiment and theory. The presence of the cusps in the rate explosion region is still expected since, for ω low enough, some modes are energetically inaccessible, and the limit $\Gamma_{\text{cf}}(\omega \rightarrow 0) \rightarrow 0$ must be fulfilled.

Rates explosion – Beyond a critical amplitude K_0^c , we observe a sudden increase of the 2D-driven decay rate [Fig. 3]. The dependence of K_0^c on the frequency for circular driving at $V_0 = 11 E_R$ is shown in Fig. 5. We observe that $K_0^c \rightarrow 2.4$ as $\omega \rightarrow \infty$, suggesting that the giant heating arises from a finite-frequency effect. Because the effect appears for both diagonal and circular drives, whereas a perturbative argument using the inverse-frequency expansion shows that there are finite $1/\omega$ corrections only for the circular drive, we surmise that the effect is likely non-perturbative. Assuming the anomalously strong heating results from an interplay of correlated physics beyond the Bogoliubov regime, and recalling that resonance effects beyond the infinite-frequency approximation lead to a $1/\omega$ dependence in the instability rates [2–4], we make the following scaling argument. Viewing the system as an effective Bose-Hubbard model, the strongly-correlated regime is reached for $g/J_{\text{eff}} \gtrsim 1$. On the other hand, we note that corrections to the infinite-frequency Floquet Hamiltonian scale as J/ω . We make the phenomenological




FIG. 4. **Decay rates vs drive frequency.** $\Gamma_{\text{cf}}(\omega)$ for the three drive trajectories at $K_0 = 1.25$ and $V_0 = 11 E_R$. The Floquet-Bogoliubov-deGennes theory $\Gamma_{\text{mum}}(\omega)$ is shown for each trajectory as dashed lines. The theoretical cusps positions are marked as vertical black dashed lines. The value we use for g is fitted to the x -only drive cusp, and is close to the expected value (see appendix). Rate explosion occurs at low frequencies when $K_0 > K_0^c(\omega)$ [cf. Fig. 3], represented as the light red zone.

observation that the dimensionless ratio $(g/J_{\text{eff}})(J/\omega)$ should be relevant to a combination of beyond-mean-field and finite-frequency effects. When J_{eff} is low in all lattice directions, $(g/J_{\text{eff}})(J/\omega)$ is large and, therefore, these effects should be large. The simple scaling relation $g/\omega \mathcal{J}_0(K_0) = 1$ gives $K_0^c(\omega) = \mathcal{J}_0^{-1}(g/\omega)$, and is shown as the red line in Fig. 5 with the experimental data. The agreement is surprisingly good for such a simple argument. The quantum many-body nature of the rates explosion calls for more extensive study, that promises new insights into periodically-driven strongly-correlated quantum lattice systems.

We present a detailed investigation of heating for interacting bosons in a periodically driven 2D lattice. The observed decay rates are substantially larger than expected from a scattering theory based on Fermi's golden rule [21], and scale as expected for interaction-driven parametric instabilities [24]. The observed exponential decay of the condensed fraction suggests that interactions between these excited modes and the BEC, not captured by FBdG theory, play an important role in the dynamics. Nonetheless, the linear scaling of the condensate loss rate with gJ/ω is indicative of direct, interaction-induced *instabilities*. Importantly, these instabilities arise from collective modes and involve coherent processes, unlike scattering in a purely FGR approach. In addition, for 2D driving, there exist regions where the heating is even larger than predicted by FBdG, which is not explained




FIG. 5. $K_0^c(\omega)$: Critical drive amplitude measured for various frequencies ω at $V_0 = 11 E_R$. For $K_0 > K_0^c$ (light red region), the decay rate for both 2D drive trajectories increases dramatically, and we observe $K_0^c \rightarrow 2.4$ at large ω . A simple relation captures this feature [red line].

by current theories. Altogether, our observations provide important insight into a major heating mechanism in bosonic systems subject to a position-modulation drive, a valuable knowledge for future many-body Floquet engineering schemes.

We note that complementary signatures of parametric instabilities have been recently investigated with bosonic atoms in periodically-driven 1D optical lattices [47].

We would like to thank M. Aidelsburger, I. Bloch, J. Näger, A. Polkovnikov, D. Sels, and K. Wintersperger for valuable discussions. This work was partially supported by the NSF Physics Frontier Center, PFC@JQI (PHY1430094) and the ARO MURI program. Work in Brussels was supported by the FRS-FNRS (Belgium) and the ERC Starting Grant TopoCold. T. B. acknowledges the support of the European Marie Skłodowska-Curie Actions (H2020-MSCA-IF-2015 Grant 701034). M.B. acknowledges support from the Emergent Phenomena in Quantum Systems initiative of the Gordon and Betty Moore Foundation, the ERC synergy grant UQUAM, and the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Quantum Algorithm Teams Program. E. D. acknowledges fundings from Harvard-MIT CUA (NSF Grant No. DMR-1308435), AFOSR-MURI Quantum Phases of Matter (grant FA9550-14-1-0035) and AFOSR-MURI: Photonic Quantum Matter, (award FA95501610323). E. M. acknowledges the support of the Fulbright Program. We used QuSpin [48, 49] to perform the numerical simulations. The authors are pleased to acknowledge that the computational work reported on in this paper was performed on the Shared Computing Cluster which is administered by Boston University's Research Computing Services.

* thomas.boulier@institutoptique.fr

† These authors contributed equally.

‡ porto@jqj.umd.edu

- [1] Immanuel Bloch, Jean Dalibard, and Sylvain Nascimbene, “Quantum simulations with ultracold quantum gases,” *Nature Physics* **8**, 267 (2012).
- [2] N. Goldman and J. Dalibard, “Periodically driven quantum systems: Effective hamiltonians and engineered gauge fields,” *Phys. Rev. X* **4**, 031027 (2014).
- [3] Marin Bukov, Luca D’Alessio, and Anatoli Polkovnikov, “Universal high-frequency behavior of periodically driven systems: from dynamical stabilization to floquet engineering,” *Advances in Physics* **64**, 139–226 (2015).
- [4] André Eckardt, “Colloquium: Atomic quantum gases in periodically driven optical lattices,” *Rev. Mod. Phys.* **89**, 011004 (2017).
- [5] H Lignier, Carlo Sias, D Ciampini, Y Singh, A Zenesini, O Morsch, and E Arimondo, “Dynamical control of matter-wave tunneling in periodic potentials,” *Phys. Rev. Lett.* **99**, 220403 (2007).
- [6] Alessandro Zenesini, Hans Lignier, Donatella Ciampini, Oliver Morsch, and Ennio Arimondo, “Coherent control of dressed matter waves,” *Phys. Rev. Lett.* **102**, 100403 (2009).
- [7] Carlo Sias, H Lignier, YP Singh, A Zenesini, D Ciampini, O Morsch, and E Arimondo, “Observation of photon-assisted tunneling in optical lattices,” *Phys. Rev. Lett.* **100**, 040404 (2008).
- [8] Frederik Görg, Michael Messer, Kilian Sandholzer, Gregor Jotzu, Rémi Desbuquois, and Tilman Esslinger, “Enhancement and sign change of magnetic correlations in a driven quantum many-body system,” *Nature* **553**, 481 (2018).
- [9] Gregor Jotzu, Michael Messer, Frederik Görg, Daniel Greif, Rémi Desbuquois, and Tilman Esslinger, “Creating state-dependent lattices for ultracold fermions by magnetic gradient modulation,” *Phys. Rev. Lett.* **115**, 073002 (2015).
- [10] Sylvain Nascimbene, Nathan Goldman, Nigel R Cooper, and Jean Dalibard, “Dynamic optical lattices of subwavelength spacing for ultracold atoms,” *Phys. Rev. Lett.* **115**, 140401 (2015).
- [11] Hannah M Price, Tomoki Ozawa, and Nathan Goldman, “Synthetic dimensions for cold atoms from shaking a harmonic trap,” *Phys. Rev. A* **95**, 023607 (2017).
- [12] Ivar Martin, Gil Refael, and Bertrand Halperin, “Topological frequency conversion in strongly driven quantum systems,” *Phys. Rev. X* **7**, 041008 (2017).
- [13] Jean Dalibard, Fabrice Gerbier, Gediminas Juzeliūnas, and Patrik Öhberg, “Colloquium: Artificial gauge potentials for neutral atoms,” *Reviews of Modern Physics* **83**, 1523 (2011).
- [14] Nathan Goldman, G Juzeliūnas, P Öhberg, and Ian B Spielman, “Light-induced gauge fields for ultracold atoms,” *Reports on Progress in Physics* **77**, 126401 (2014).
- [15] NR Cooper, J Dalibard, and IB Spielman, “Topological bands for ultracold atoms,” arXiv preprint arXiv:1803.00249 (2018).
- [16] Marin Bukov, Markus Heyl, David A Huse, and Anatoli Polkovnikov, “Heating and many-body resonances in a periodically driven two-band system,” *Phys. Rev. B* **93**, 155132 (2016).
- [17] Monika Aidelsburger, Michael Lohse, C Schweizer, Marcos Atala, Julio T Barreiro, S Nascimbene, NR Cooper, Immanuel Bloch, and Nathan Goldman, “Measuring the chern number of hofstadter bands with ultracold bosonic atoms,” *Nature Physics* **11**, 162 (2015).
- [18] Colin J Kennedy, William Cody Burton, Woo Chang Chung, and Wolfgang Ketterle, “Observation of bose-einstein condensation in a strong synthetic magnetic field,” *Nature Physics* **11**, 859–864 (2015).
- [19] Hirokazu Miyake, Georgios A. Siviloglou, Colin J. Kennedy, William Cody Burton, and Wolfgang Ketterle, “Realizing the harper hamiltonian with laser-assisted tunneling in optical lattices,” *Physical Review Letters* **111**, 1–5 (2013), arXiv:1308.1431.
- [20] Michael Messer, Kilian Sandholzer, Frederik Görg, Joaquín Minguzzi, Rémi Desbuquois, and Tilman Esslinger, “Floquet dynamics in driven fermi-hubbard systems,” *Physical Review Letters* **121**, 233603 (2018).
- [21] Martin Reitter, Jakob Näger, Karen Wintersperger, Christoph Sträter, Immanuel Bloch, André Eckardt, and Ulrich Schneider, “Interaction dependent heating and atom loss in a periodically driven optical lattice,” *Phys. Rev. Lett.* **119**, 200402 (2017).
- [22] CE Creffield, “Instability and control of a periodically driven bose-einstein condensate,” *Phys. Rev. A* **79**, 063612 (2009).
- [23] Sayan Choudhury and Erich J Mueller, “Stability of a floquet bose-einstein condensate in a one-dimensional optical lattice,” *Phys. Rev. A* **90**, 013621 (2014).
- [24] S Lellouch, M Bukov, E Demler, and N Goldman, “Parametric instability rates in periodically driven band systems,” *Phys. Rev. X* **7**, 021015 (2017).
- [25] Marin Bukov, Sarang Gopalakrishnan, Michael Knap, and Eugene Demler, “Prethermal floquet steady states and instabilities in the periodically driven, weakly interacting bose-hubbard model,” *Phys. Rev. Lett.* **115**, 205301 (2015).
- [26] S Lellouch and N Goldman, “Parametric instabilities in resonantly-driven bose-einstein condensates,” *Quantum Science and Technology* **3**, 024011 (2018).
- [27] Luca D’Alessio and Marcos Rigol, “Long-time behavior of isolated periodically driven interacting lattice systems,” *Phys. Rev. X* **4**, 041048 (2014).
- [28] Achilleas Lazarides, Arnab Das, and Roderich Moessner, “Equilibrium states of generic quantum systems subject to periodic driving,” *Phys. Rev. E* **90**, 012110 (2014).
- [29] Adam Nahum, Jonathan Ruhman, and David A Huse, “Dynamics of entanglement and transport in one-dimensional systems with quenched randomness,” *Physical Review B* **98**, 035118 (2018).
- [30] Dmitry A. Abanin, Wojciech De Roeck, and Fran çois Huvneers, “Exponentially slow heating in periodically driven many-body systems,” *Phys. Rev. Lett.* **115**, 256803 (2015).
- [31] Takashi Mori, Tomotaka Kuwahara, and Keiji Saito, “Rigorous bound on energy absorption and generic relaxation in periodically driven quantum systems,” *Phys. Rev. Lett.* **116**, 120401 (2016).
- [32] Tomotaka Kuwahara, Takashi Mori, and Keiji Saito, “Floquet–magnus theory and generic transient dynamics in periodically driven many-body quantum systems,” *Annals of Physics* **367**, 96–124 (2016).

- [33] Simon A Weidinger and Michael Knap, “Floquet prethermalization and regimes of heating in a periodically driven, interacting quantum system,” *Scientific reports* **7**, 45382 (2017).
- [34] Roderich Moessner and SL Sondhi, “Equilibration and order in quantum floquet matter,” *Nature Physics* **13**, 424 (2017).
- [35] Owen Howell, Phillip Weinberg, Dries Sels, Anatoli Polkovnikov, and Marin Bukov, “Frequency-controlled thermalization phase transition in a chaotic periodically-driven classical spin chain,” arXiv preprint arXiv:1802.04910 (2018).
- [36] Thomas Bilitewski and Nigel R Cooper, “Scattering theory for floquet-bloch states,” *Phys. Rev. A* **91**, 033601 (2015).
- [37] Sayan Choudhury and Erich J Mueller, “Transverse collisional instabilities of a bose-einstein condensate in a driven one-dimensional lattice,” *Phys. Rev. A* **91**, 023624 (2015).
- [38] C Cabrera-Gutiérrez, E Michon, M Arnal, V Brunaud, T Kawalec, J Billy, and D Guéry-Odelin, “Resonant excitations of a bose einstein condensate in an optical lattice,” arXiv preprint arXiv:1808.00345 (2018).
- [39] Y-J Lin, Abigail R Perry, Robert L Compton, Ian B Spielman, and James V Porto, “Rapid production of r 87 b bose-einstein condensates in a combined magnetic and optical potential,” *Phys. Rev. A* **79**, 063631 (2009).
- [40] Jennifer Sebby-Strabley, Marco Anderlini, PS Jessen, and James V Porto, “Lattice of double wells for manipulating pairs of cold atoms,” *Phys. Rev. A* **73**, 033605 (2006).
- [41] E. Magnan, J. Maslek, C. Bracamontes, A. Restelli, T. Boulter, and J. V. Porto, “A low-steering piezo-driven mirror,” *Review of Scientific Instruments* **89**, 073110 (2018), <https://doi.org/10.1063/1.5035326>.
- [42] Phillip Weinberg, Marin Bukov, Luca DAlessio, Anatoli Polkovnikov, Szabolcs Vajna, and Michael Kolodrubetz, “Adiabatic perturbation theory and geometry of periodically-driven systems,” *Physics Reports* **688**, 1–35 (2017).
- [43] Viktor Novičenko, Egidijus Anisimovas, and Gediminas Juzeliūnas, “Floquet analysis of a quantum system with modulated periodic driving,” *Phys. Rev. A* **95**, 023615 (2017).
- [44] Leonardo Fallani, Luigi De Sarlo, Jessica E Lye, Michele Modugno, Robert Saers, Chiara Fort, and Massimo Inguscio, “Observation of dynamical instability for a bose-einstein condensate in a moving 1d optical lattice,” *Phys. Rev. Lett.* **93**, 140406 (2004).
- [45] E Michon, C Cabrera-Gutiérrez, A Fortun, M Berger, M Arnal, V Brunaud, J Billy, C Petitjean, P Schlagheck, and D Guéry-Odelin, “Phase transition kinetics for a bose einstein condensate in a periodically driven band system,” *New Journal of Physics* **20**, 053035 (2018).
- [46] See supplementary material.
- [47] J Näger, K Wintersperger, M Bukov, S Lellouch, E Demler, U Schneider, I Bloch, N Goldman, and M Aidelsburger, “Parametric instabilities of interacting bosons in periodically-driven 1d optical lattices,” arXiv preprint arXiv:1808.07462 (2018).
- [48] Phillip Weinberg and Marin Bukov, “QuSpin: a Python Package for Dynamics and Exact Diagonalisation of Quantum Many Body Systems part I: spin chains,” *SciPost Phys.* **2**, 003 (2017).
- [49] P. Weinberg and M Bukov, “Quspin: a python package for dynamics and exact diagonalisation of quantum many body systems. part ii: bosons, fermions and higher spins,” arXiv preprint arXiv:1804.06782 (2018).
- [50] R C Brown, R Wyllie, S B Koller, E A Goldschmidt, Michael Foss-Feig, and JV Porto, “Two-dimensional superexchange-mediated magnetization dynamics in an optical lattice,” *Science* **348**, 540 (2015).
- [51] M. Weinberg, C. Ölschläger, C. Sträter, S. Prella, A. Eckardt, K. Sengstock, and J. Simonet, “Multiphoton interband excitations of quantum gases in driven optical lattices,” *Physical Review A - Atomic, Molecular, and Optical Physics* **92**, 1–10 (2015), arXiv:arXiv:1505.02657v1.
- [52] P B Blakie, A S Bradley, M J Davis, R J Ballagh, and C W Gardiner, “Dynamics and statistical mechanics of ultra-cold Bose gases using c-field techniques,” *Adv. Phys.* **57**, 363–455 (2008).
- [53] Anatoli Polkovnikov, “Phase space representation of quantum dynamics,” *Annals of Physics* **325**, 1790–1852 (2010).

APPENDIX

Thermalization of excited atoms

The condensate fraction can decay either by direct Floquet-driven loss or by heating due to relaxation of energetic excitations. The latter mechanism occurs over a thermalization time scale, and can be probed by observing relaxation of out of equilibrium states in an undriven, static lattice. Special attention to the drive turn off is required to avoid unwanted excitation due to micro-motion during a drive period. Abruptly turning off the drive induces a kick large enough to create a significant out-of-equilibrium population by inter-band excitation.

To determine the relaxation time scale and test for this additional condensate loss mechanism, we measured the evolution of the condensed fraction when holding the atomic cloud in a static lattice, immediately after an abrupt stop of the drive. While the condensed fraction is initially unchanged, upon letting the static system evolve for a time t_{hold} , we observe a subsequent decrease of the condensed fraction as excited atoms thermalize with the rest of the sample. Fig. 6 shows an example of thermalization for a diagonal drive at $\omega = 2\pi \times 2$ kHz. When there is sufficient initial excitation (blue data on figure 6), we observe a characteristic thermalization time of order 2 ms, that does not depend on the initial non-equilibrium population.

We observe that the condensed fraction *increases* slightly with hold time in the cases where minimal atom excitation occurs (red and green data in figure 6). We attribute this to rethermalization along the tube axis: entropy added in the degree of freedom associated with the lattice direction can be redistributed along the z (tube) axis, which is visible as a slightly decreased temperature




FIG. 6. **Thermalization of the non-equilibrium population.** – The lattice is shaken for a few periods, then held static at $t_{hold} = 0$. There exist at $t_{hold} = 0$ a non-equilibrium population that thermalizes with the BEC in a few milliseconds, resulting in a clear decrease of the condensed fraction. The amplitude of this effect depends on the initial size of the non-equilibrium population. It is maximum if atoms are kicked to higher bands by an abrupt stop of the drive (blue). If no such population is created by the stopping of the drive, for example by choosing an end phase that minimizes the kick (red) or by rapidly ramping down the drive amplitude (green), no condensed fraction decay is observed.

in the x-y plane.

The amplitude of the condensed fraction decrease is indicative of the energy of the initial non-equilibrium population. We can measure this amplitude by comparing the condensed fraction at $t_{hold} = 0$ and $t_{hold} = 6$ ms, a time sufficient for the thermalization to have occurred. Fig. 7 shows the remaining condensate fraction after $t_{hold} = 6$ ms as a function of the phase at which the drive is stopped, for an abrupt stop (blue) and for a smooth but rapid (1 period) turn off (green). When the stop is abrupt, the fraction of atoms excited depends on the end phase: abruptly immobilizing the lattice can induce an effective force which depends on the lattice velocity immediately prior to immobilization. Therefore the excited fraction is maximized when $\dot{q}(t)$ is maximally discontinuous (stop phases of $0[\pi]$ rad) and minimized when smooth (stop phases of $\frac{\pi}{2}[\pi]$ rad). In all our decay rate data where we smoothly turned off the drive (in at least one period), no turnoff-induced heating was observed.

Abruptly stopping the drive is not the only potential source of non-equilibrium population. The unstable Bogoliubov modes studied in the main text could themselves produce a population that thermalizes, and that could potentially modify the measured decay rates. This can be tested with the same stop-and-hold measurement, if turnoff-induced populations are avoided. As visible in




FIG. 7. **Amplitude of the thermalization process.** – Difference between the condensed fraction at $t = t_{end}$ ($t_{hold} = 0$) and at $t = t_{end} + 6$ ms ($t_{hold} = 6$ ms), as a function of the end drive phase, for a $2\pi \times 2$ kHz diagonal (2D) drive. For an abrupt stop (blue), a large population can be transferred out of the BEC for some end phases. This results in a large thermalization event and a clear drop of the condensed fraction. The effect is minimal when stopping the drive such that $q(t)$ is a smooth function. If, however, the drive is stopped by ramping down the amplitude (green), no condensed fraction decay can be observed at any phase.

Fig. 7, this is realized for stop phases of $\frac{\pi}{2}[\pi]$ rad or when smoothly turning off the drive. For our data, avoiding turnoff-induced population when stopping the drive results in no visible decay. This is shown as the green data in Fig. 6. The result is identical for the whole region of parameter space studied here. We deduce that the energy carried by the non-equilibrium population created by the unstable modes is not enough to significantly impact the measured rates.

Experimental considerations

Accelerating the BEC to the band edge. – In order to accelerate our BEC from $\mathbf{q} = (0, 0)$ to a desired $\mathbf{q} = (q_x, q_y)$, we apply a constant force $\mathbf{F} = \dot{\mathbf{q}}$ for a fixed time in the lattice plane. The force is generated with a constant magnetic gradient, acting on the BEC in the $|F = 1, m_F = -1\rangle$ ground state. Bias coils in the three spatial directions control the gradient direction.

The BEC becomes unstable for quasimomenta about halfway to the band edge: this is the well-known static instability [44]. On the other hand, as mentioned in the main text, ramping up the drive amplitude beyond $K_0 = 2.4$ reverses the band smoothly as J_{eff} becomes negative: $\mathbf{q} \sim (0, 0)$ becomes unstable and the band edges (or corners, depending on the drive trajectory) become stable. In order keep the BEC in a stable region

(in a effectively static dynamical stability sense) at any given time, we synchronize the BEC acceleration with the ramping on of the drive, such that the BEC crosses the static instability point $q = 0.6 \pi/a$ when $K_0 = 2.4$.

We accelerate the BEC to $\mathbf{q} = (\pi, \pi)$ for the two 2D drives (diagonal and circle) and to $\mathbf{q} = (\pi, 0)$ for the x-only 1D drive, since these become stable whenever $K_0 > 2.4$.

Calibration of translation amplitude. – The piezo-actuated mirrors were each roughly calibrated offline using an interferometric technique [41]. The final calibration was performed on the atoms by measuring the tunneling-dependent magnetization decay rate of 2D staggered spin magnetization [50] as a function of drive strength K_0 . The drive strength at which the tunneling rate J_{eff} vanishes is determined by the condition $K_0 = 2.4$.

Calibration of tight-binding parameters. – The lattice depth is calibrated via Raman-Nath diffraction. In the tight-binding limit, the tunneling rate is derived through the modeled 1D dispersion as

$$J \equiv \frac{E(q = \pi/a) - E(q = 0)}{4} \quad (2)$$

The on-site interaction g is calibrated from the x -only drive cusp (see Fig. 4). The point at which the rates go from increasing to decreasing, ω_c^{Diag} , is given by

$$\omega_c^x = \sqrt{4J_{\text{eff}}(4J_{\text{eff}} + 2g)}. \quad (3)$$

Knowing J , the experimental value of ω_c^x offers a calibration for g . For $V_0 = 11 E_R$, $J = 50$ Hz and the measured value of $\omega_c^x = 444$ Hz gives $g = 700$ Hz. As an additional check, this value of g is then used to predict the diagonal drive cusp, $\omega_c^{\text{Diag}} = \sqrt{8J_{\text{eff}}(8J_{\text{eff}} + 2g)} = 655$ Hz. The observed value of ~ 650 Hz is in good agreement with this prediction.

The interaction strength g depends upon the atom number, the dipole trap and the lattice parameters. To confirm that the experimental calibration matches these known experimental parameters, we also estimate g through tight binding and Thomas-Fermi assumptions: in the lattice plane, the wavefunction $\psi(x)$ is taken to be well approximated by a Mathieu function, while we use a Thomas-Fermi profile in the tube direction. The interaction energy $g \propto \iint_{-a/2}^{a/2} |\psi(\mathbf{r})|^4 d^2\mathbf{r}$ is then calculated from the known experimental parameters, including the density profile due to the dipole trap (frequencies $\{\omega_x, \omega_y, \omega_z\} = \{11, 45, 120\}$ Hz). For $V_0 = 11 E_R$ we find $g = 850$ Hz, similar to the calibrated value of 700 Hz. Note that the systematic 20% uncertainty in the atom number can easily explain the small offset between estimation and calibration.

Bandwidths – It is important to note that ω_c is in general different from the effective bandwidth B . For a 2D (diagonal or circular) drive in our 2D lattice,

$$B^{2D} = \sqrt{8J|\mathcal{J}_0(K_0)| \times (8J|\mathcal{J}_0(K_0)| + 2g)} \quad (4)$$

and, with a 1D drive in the 2D lattice,

$$B^{1D} = \sqrt{4J(|\mathcal{J}_0(K_0)| + 1) \times (4J(|\mathcal{J}_0(K_0)| + 1) + 2g)}, \quad (5)$$

which is in general different from ω_c , as observed in the main text: only in the case of the diagonal drive do we find that the maximally unstable mode had the maximum ground band energy, and therefore $\omega_c = B$.

Background rates. – All theoretical plots take into account the background decay rate, predominantly due to lattice photon scattering. We experimentally determine this by setting $K_0 = 0$ and measuring the resulting rate with the same procedure as in the main text. This constant rate $y_0 \sim 1 \text{ s}^{-1}$ for $V_0 = 11 E_R$ is then added to the FBdG formula for comparison with the experimental data. Since multi-photon resonances to higher bands [51] could complicate the interpretation of the measured decay rates, we performed heating measurements up to drive frequencies of $\omega = 2\pi \times 21$ kHz to identify excited band resonances. Population transfer to higher bands is directly visible on the bandmapping images. The lowest frequency at which resonant higher band excitation was observed was $\omega = 2\pi \times 6.25$ kHz for $V_0 = 11 E_R$, and we therefore limited our heating measurements to below $2\pi \times 4$ kHz to avoid these effects.

Heating in the absence of Floquet drive – To rule out heating mechanisms that do not depend on the drive but are instead related to the change in J_{eff} , it would be useful to compare the heating observed at a given K_0 to the heating observed in a static lattice with a depth chosen to have an equivalent $J = J_{\text{eff}}$ (for $J_{\text{eff}} > 0$). Here, such direct comparison is experimentally challenging under the same conditions as the Floquet experiment, due to the change in the trap confinement when increasing the lattice depth. In addition to a potential change in gravitational sag (which we have minimized by aligning the optical lattice beams directly on the dipole-trapped BEC), the change in trap frequency when changing the lattice depth can excite breathing motion along the non-lattice tube direction, as well as cause a redistribution of atoms within the trap transverse to the lattice direction. These trap-changing effects do not occur for the Floquet modification of the tunneling. To avoid this motion / redistribution, one needs to increase the lattice adiabatically, i.e. on a slower timescale than the Floquet drive is turned on. Nonetheless, we did a type of comparison to the data in figure 2 by turning on the lattice slowly (over 200 ms) to a final lattice depth to obtain a given




FIG. 8. **Typical decay plot.** Each data point (blue dots) is an experimental realization, where the condensed fraction is measured after a variable shake time (all else kept constant). The resulting decay curve is then fitted with an exponential function (red thick line). This particular example used $K_0 = 2$, $\omega/2\pi = 4$ kHz and $V_0 = 11 E_R$. Each experimental rate in the manuscript is extracted from such a fit, and its error bar is given by ± 1 standard deviation.

static J , and observed the condensate fraction. As an example, we increased the lattice depth to $18.2 E_R$, which changes the tunneling from $J = 2\pi \times 50$ Hz at $11 E_R$ to $J = 2\pi \times 12$ Hz at $18.2 E_R$, corresponding in the Floquet case to $K_0 = 2$. The fact that there is still condensate remaining after the 200 ms turn on is already an indication that the condensate decay rate in the unshaken lattice is slow. Measuring the condensate decay rate at this depth gives $\Gamma/2\pi = 2 \text{ s}^{-1}$, which is slower than the equivalent Floquet rate by a factor of nine, and slower than all the data at $K_0 = 2.1$ shown in figure 2 by at least a factor of seven.

Extracting the decay rates

Rates extraction – Our data consists of series of measured condensed fractions after various driving times. A time series typically presents an exponential-looking decay. An example for such decay is given in figure 8. Since we focus on early the time decay rate, greater weight is given to earlier data points. We fit to an exponential with no offset (2 fit parameters): $f(t) = Ae^{-\Gamma_{cf}t}$, where A is the $t=0$ condensed fraction (typically $A \geq 0.5$). The rates presented in this manuscript are the extracted fit parameter Γ_{cf} , and the uncertainties correspond to ± 1 standard deviation.

A small fraction of the experimental data doesn't show an exponential decay. In such cases, the condensed fraction vs time looks linear, which is indicative of an *acceler-*




FIG. 9. **Decay rates in the critical region** – Zoomed in plot of data shown in figure 3, focusing on the region between K_0^c and $K_0 = 2.5$. The difference between circular and diagonal drive is clearly visible, with the circular drive rate as much as twice that of the diagonal drive rate.

ation relative to an exponential decay. Whenever fitting to an exponential was impossible, we took the slope, focusing on early times where the condensed fraction was more than half that of $t = 0$. In any case, the measured decay rate under all conditions corresponds to the decay rate at initial times.

For these rates to be directly compared to the BdG prediction, additional considerations must be taken. First, let us consider a maximally unstable Bogoliubov mode with an *amplitude* predicted to grow as $e^{\Gamma_{\text{mum}}t}$. An experiment will actually detect a rate $2\Gamma_{\text{mum}}$, as it measures an amplitude *squared* (typically, the number of atoms in the unstable mode). Second, since the experiment measures how many atoms leave the BEC (to populate the modes) per unit of time, it is sensitive to the number of simultaneous maximally unstable modes, as each is a decay channel. If two modes are equally and maximally unstable, as is possible in 2D, then an additional factor of 2 is needed in the theory to compare to the experiment. This multiple-modes factor is 1 for the x -only drive, and 2 for the circle and diagonal drives. All these extra factors were added to the theory plots throughout this paper: in total, the x -only drive theory is $2\times$ and the circle and diagonal drives are $4\times$ larger than the bare BdG rates predicted in [24]. These points are expanded in the derivation of the FBdG rate Γ_{mum} later in this Appendix.

Rate explosion: circular versus diagonal – We observe that the circular decay rate increases faster than the diagonal rate for $K_0 \gtrsim K_0^c$. For example, figure 9 shows




FIG. 10. **Rate comparison: FBdG theory vs. BdG simulation.** Comparison between the dynamically-extracted instability rates from solving the exact BdG equations of motion for a finite number of drive cycles (blue stars), and the FBdG prediction (red dashed line). The three plots correspond to x -only (left), 2D circular (middle) and 2D diagonal (right) drives. The numerical simulations include a tube transverse degree of freedom. The parameters are $g/J = 12$, $K_0 = 1.25$, and a momentum grid of $80 \times 80 \times 101$ modes in the x , y , and z -direction, respectively.

a zoomed-in version of figure 3, where $V_0 = 11$ Er and $\omega = 2\pi \times 2.5$ kHz, to make this observation clearer. The rates go from being approximately equal below K_0^c , to $\Gamma_{cf}^{circ} > 2 \Gamma_{cf}^{diag}$ when tunneling is suppressed. Since breaking time-reversal symmetry is necessary for many proposed schemes, this observation may be of interest to the Floquet engineering community.

Difficulties associated with dynamical rates – In the main text, we compared the decay rates measured in the experiment to the instability rates predicted by FBdG theory. Here, we elaborate on some intrinsic difficulties in the procedure which may affect the extracted values.

As explained in Ref. [24], for drive frequencies below the effective drive-renormalized Floquet-Bogoliubov bandwidth, there exists an entire manifold of resonant modes. While all of them contribute to expectation values of observables at very short times, only the maximally-unstable mode \mathbf{q}_{mum} dominates the long-time BdG dynamics, and the rate associated with \mathbf{q}_{mum} sets the parametric instability rate. Thus, at any finite time, the FBdG dynamics is in a crossover between these two regimes, which shrinks exponentially with time. Yet the time-width of this crossover also depends on the drive frequency: the higher the frequency, the smaller the instability rate and the longer it takes for the exponential behavior to become visible.

When extracting the rates from data, effects due to this crossover become relevant. To test this, we performed exact numerical simulations of the BdG equations of motion, and computed the dynamics of the excited fraction of atoms $n_{\mathbf{q}}(t)$ over a finite number of driving cycles, which increases suitably with the drive frequency. We then extracted the instability rates using least-square fitting as the slope of $\log n_{\mathbf{q}}(t)$ over the last eight driving cycles, to maximally eliminate transient effects. A comparison between the numerically-extracted rates and the analytic theory prediction is shown in Fig. 10 for the

three types of drives. Note that the agreement becomes worse at larger ω , since this decreases the rate and pushes the exponential regime to later times. This is a source of error, which is certainly relevant for the experimental determination of the rates.

Additionally, in the experiment there are strong beyond Bogoliubov effects, not captured by FBdG theory. Due to the nonlinearity of the Gross-Pitaevskii equation which leads to saturation of the condensate depletion, the exponential BdG regime mentioned above crosses over into a third, scattering-dominated regime. In this regime, the population transferred coherently to the maximally-unstable modes by the parametric resonance starts decaying into the surrounding finite-momentum modes at high energy, leading to uncontrolled irreversible heating [25]. This suggests that the instability rates can change in time. This additionally obfuscates the comparison of the experiment with the FBdG theory.

HEATING DYNAMICS OF THE TRUNCATED WIGNER APPROXIMATION

While FBdG theory is valid in the short-time regime of the dynamics, it has some serious deficiencies. Perhaps the most notable of these, when it comes to out-of-equilibrium dynamics, is the lack of particle-number conservation: the condensate is assumed to be an infinite reservoir which supplies particles to indefinitely increase the occupation of pairs of modes with finite and opposite momenta. In equilibrium, this description works well and captures the physics in the superfluid phase. Away from equilibrium, however, condensate depletion processes such as the parametric instabilities studied in this work lead to significant depletion of the BEC and the mean-field Bogoliubov description ultimately breaks down under typical observation times.

Particle conservation is obeyed in the Truncated

Wigner Approximation (TWA), which also includes nonlinear interactions modeling collisions between Bogoliubov quasiparticles [52, 53], and is capable of describing thermalization at later stages, due to the continuous

pumping of energy into the system.

The starting point for the TWA is the Gross-Pitaevskii equation which, in the co-moving real-space frame, reads ($\hbar = 1$)

$$i\partial_t a_{\mathbf{r}}(t) = -J[a_{\mathbf{r}+a\mathbf{e}_x}(t) - a_{\mathbf{r}-a\mathbf{e}_x}(t)] - J[a_{\mathbf{r}+a\mathbf{e}_y}(t) - a_{\mathbf{r}-a\mathbf{e}_y}(t)] - \frac{\partial_z^2}{2m} a_{\mathbf{r}}(t) + \omega K_0 \mathbf{r} \cdot (\sin(\omega t)\mathbf{e}_x + \kappa_d \sin(\omega t + \phi)\mathbf{e}_y) a_{\mathbf{r}}(t) + U|a_{\mathbf{r}}(t)|^2 a_{\mathbf{r}}(t), \quad (6)$$

where $a_{\mathbf{r}}(t)$ models the bosonic system at time t and position $\mathbf{r} = (x, y, z)$. The kinetic energy reflects the lattice degrees of freedom in the (x, y) -plane, and the continuous transverse mode along the z -axis. The periodic drive is in the (x, y) -plane with frequency ω and amplitude K_0 . $\kappa_d = 0$ for the x -only drive and $\kappa_d = 1$ for both 2D drives. ϕ is the relative drive phase between x and y ($\phi = 0$ for diagonal drive and $\phi = -\frac{\pi}{2}$ for circular drive). Finally, the on-site interaction strength is denoted by U (such that $U/V \sum_{i,j} \int dz |a_{\mathbf{r}}|^2 = g$).

Since at time $t = 0$ the system forms a BEC of N_0 atoms in a volume V , assuming a macroscopic occupation $n_0 = \sqrt{N_0/V}$ in the uniform ($\mathbf{q} = \mathbf{0}$) condensate, the field $a_{\mathbf{r}}$ can be decomposed as

$$a_{\mathbf{r}} = n_0 + \frac{1}{\sqrt{V}} \sum_{\mathbf{q} \neq \mathbf{0}} u_{\mathbf{q}} \gamma_{\mathbf{q}} e^{-i\mathbf{q} \cdot \mathbf{r}_j} + v_{-\mathbf{q}}^* \gamma_{-\mathbf{q}}^* e^{+i\mathbf{q} \cdot \mathbf{r}_j}, \quad (7)$$

where $u_{\mathbf{q}}$ and $v_{\mathbf{q}}$ are the Bogoliubov modes which solve the time-independent BdG equations at $t = 0$ [24]. Here, $\gamma_{\mathbf{q}}$ is a complex-valued Gaussian random variable [associated with the quantum annihilator $\hat{\gamma}_{\mathbf{q}}$ of Bogoliubov modes] with mean and variance set by the corresponding quantum expectation values in the Bogoliubov ground state [52].

Hence, in the TWA one draws multiple random realizations of $\gamma_{\mathbf{q}}$, each of which corresponds to a different initial state. One then evolves every member of this ensemble according to Eq. (6), computes the observable of interest, and takes the ensemble average $\overline{(\cdot)}$ in the end. For instance one can compute the total number of excited atoms as

$$n_{\text{ex}}(t) = \frac{1}{V} \sum_{\mathbf{q} \neq \mathbf{0}} \overline{|a_{\mathbf{q}}(t)|^2}, \quad (8)$$

which, due to particle number conservation also reflects the dynamics of the condensate depletion.

In general, we expect that the condensate depletion curve shows two types of behavior: at short times FBdG theory applies and $n_{\text{ex}}(t) \sim \exp(2\Gamma_{\text{mum}} t)$ grows exponentially in time. Hence, the condensate depletion curve $|a_{\mathbf{q}=\mathbf{0}}(t)|^2 = V(n_0 - n_{\text{ex}}(t))$ is concave. At long times, nonlinear interaction effects in the GPE become impor-

tant, leading to saturation, and the curvature of condensate depletion changes sign. Therefore, in the long-time regime the curve is concave. The opposite behavior is true for the evolution of the excitations $n_{\text{ex}}(t)$. The curvature of the experimental data, cf. Fig. 8, suggests that the system enters well into the long-time regime. Yet, the measured decay rates appear consistent with the short-time Bogoliubov theory [main text].

To shed light on this intriguing observation, we perform TWA simulations on a periodically-driven homogeneous system in (2+1)-dimensions, and extract the short-time and long-time rates from the numerical data. We use a comparison with the BdG simulations, to separate the short-time regime (where agreement between BdG equations of motion and TWA is expected) from the longer-time regime [11 left panel]. For the sake of comparison with experiments, we fit the long-time TWA growth to an exponential, even though we find that it follows a more complicated functional form.

Figure 11 [right panel] shows a scan of the TWA rates over the effective interaction parameter g . We find that both the short-time and long-time rates are of similar strength. More importantly, they do not show a quadratic scaling in g , as predicted by Fermi's Golden Rule. This behavior is consistent with the experimental observations. Note the mismatch between FBdG theory [black] and the short-time BdG simulations [blue], which arises since the most unstable mode does not yet dominate the dynamics at such short times [see Fig. 10 and corresponding discussion]. Indeed, we find an excellent agreement between BdG numerics and FBdG theory if we extract the rates from the long-time regime. The short-time BdG rates agree qualitatively with the short-time TWA rates, as expected from the agreement seen in Fig. 11 [left panel]. The rates are extracted from a least-square fit over the last 5 consecutive driving cycles of short-time region of agreement between BdG and TWA. Since the rates are dynamical, i.e. change depending on the time-window used to extract them, the curves in Fig. 11 [right panel] are not smooth.

We also did a frequency and amplitude scans of the long-time TWA rates to look for signatures of the Bessel function $\mathcal{J}_2(K_0)$, and the cusp at the critical frequency




FIG. 11. **Rate comparison: TWA vs. BdG.** Left panel: Numerical simulation of the excitations growth using TWA (solid lines) and FBdG (dashed lines). The TWA curves change curvature beyond the regime of validity of BdG. Right panel: excitations [i.e. condensate depletion] growth rate against effective interaction strength g for FBdG theory (black), BdG short-time evolution (blue), TWA short-time evolution (red) and TWA long-time evolution (green). The parameters are $K_0 = 2.1$, $\omega/J = 20.0$ and $n_0 = 50.0$. We used a system of $80 \times 80 \times 101$ momentum modes in the (x, y, z) -direction, respectively. The TWA data is averaged over 50 independent realizations and the error bars (shaded area) are computed using a bootstrapping approach.

ω_c , as expected from FBdG theory and found experimentally. Unfortunately, we do not see clear signatures of such behaviors in our TWA simulations. Thus, we cannot conclude that the TWA captures the long-time thermalization dynamics of driven bosonic cold atom systems accurately. More interestingly, the rates explosion [see main text] is also beyond the TWA dynamics, suggesting that quantum effects, such as loss of coherence, are important for describing this phenomenon. Another possible reason for disagreement is the single band approximation, as its validity for the Floquet system has not been fully understood so far.

MOST UNSTABLE MODES FOR LINEAR, DIAGONAL AND CIRCULAR LATTICE DRIVES WITHIN FBdG THEORY

In this appendix, we briefly revisit the derivation for the most unstable mode within FBdG theory [26], and extend the results to diagonal and circular drives. The take-home message is that the critical saturation frequency ω_c , which defines the position of the cusp in the instability rate curves, coincides for the linear and circular drives; for the diagonal drive the value is twice as large on the square lattice. It is straightforward to extend the analysis below to other lattice geometries.

The starting point are the BdG equations of motion for the mode functions $u_{\mathbf{q}}(t)$ and $v_{\mathbf{q}}(t)$ in the rotating

frame

$$i\partial_t \begin{pmatrix} u_{\mathbf{q}} \\ v_{\mathbf{q}} \end{pmatrix} = \begin{pmatrix} \varepsilon(\mathbf{q}, t) + g & g \\ -g & -\varepsilon(-\mathbf{q}, t) - g \end{pmatrix} \begin{pmatrix} u_{\mathbf{q}} \\ v_{\mathbf{q}} \end{pmatrix}, \quad (9)$$

where $\varepsilon(\mathbf{q}, t) \geq 0$ denotes the time-dependent free lattice dispersion relation [shifted by the chemical potential so that it is non-negative].

To analyze the effects of parametric instabilities, we first isolate the time-average term and write $\varepsilon(\mathbf{q}, t) = \varepsilon_{\text{eff}}(\mathbf{q}) + g(\mathbf{q}, t)$. This will separate the right-hand-side of the BdG EOM separate in an effective time-averaged term, and a time-periodic perturbation. Following Ref. [24], we now apply a phase rotation, followed by a static Bogoliubov transformation w.r.t. $\varepsilon_{\text{eff}}(\mathbf{q})$:

$$\begin{pmatrix} u_{\mathbf{q}} \\ v_{\mathbf{q}} \end{pmatrix} = \begin{pmatrix} \cosh(\theta_{\mathbf{q}}) & \sinh(\theta_{\mathbf{q}}) \\ \sinh(\theta_{\mathbf{q}}) & \cosh(\theta_{\mathbf{q}}) \end{pmatrix} \begin{pmatrix} e^{-2iE_{\text{eff}}^{\text{Bog}}(\mathbf{q})t} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \tilde{u}'_{\mathbf{q}} \\ \tilde{v}'_{\mathbf{q}} \end{pmatrix},$$

where $\cosh(2\theta_{\mathbf{q}}) \equiv (\varepsilon_{\text{eff}}(\mathbf{q}) + g)/E_{\text{eff}}^{\text{Bog}}(\mathbf{q})$ and $\sinh(2\theta_{\mathbf{q}}) \equiv g/E_{\text{eff}}^{\text{Bog}}(\mathbf{q})$ with

$$E_{\text{eff}}^{\text{Bog}}(\mathbf{q}) = \sqrt{\varepsilon_{\text{eff}}(\mathbf{q})(\varepsilon_{\text{eff}}(\mathbf{q}) + 2g)}$$

the time-averaged Bogoliubov dispersion. The Bogoliubov transformation diagonalizes the effective static time-average term, while the phase rotation makes it easier to identify the parametric resonant terms [see App. C in [24], and the Supplemental Material to Ref. [25] for an application to the paradigmatic parametric oscillator]. The BdG EOM now read

$$i\partial_t \begin{pmatrix} \tilde{u}'_{\mathbf{q}} \\ \tilde{v}'_{\mathbf{q}} \end{pmatrix} = \left[E_{\text{eff}}^{\text{Bog}}(\mathbf{q}) \hat{\mathbf{1}} + \hat{W}_{\mathbf{q}}(t) + \sinh(2\theta_{\mathbf{q}}) \begin{pmatrix} 0 & h_{\mathbf{q}}(t)e^{-2iE_{\text{eff}}^{\text{Bog}}(\mathbf{q})t} \\ -h_{\mathbf{q}}(t)e^{2iE_{\text{eff}}^{\text{Bog}}(\mathbf{q})t} & 0 \end{pmatrix} \right] \begin{pmatrix} \tilde{u}'_{\mathbf{q}} \\ \tilde{v}'_{\mathbf{q}} \end{pmatrix}, \quad (10)$$

where

$$W_{\mathbf{q}}(t) = \begin{pmatrix} g_{\mathbf{q}}(t) \cosh^2(\theta_{\mathbf{q}}) + g_{-\mathbf{q}}(t) \sinh^2(\theta_{\mathbf{q}}) & 0 \\ 0 & -g_{-\mathbf{q}}(t) \cosh^2(\theta_{\mathbf{q}}) - g_{\mathbf{q}}(t) \sinh^2(\theta_{\mathbf{q}}) \end{pmatrix}, \quad (11)$$

$$h_{\mathbf{q}}(t) = \frac{1}{2} [g_{\mathbf{q}}(t) + g_{-\mathbf{q}}(t)].$$

By definition of $g_{\mathbf{q}}(t)$, the diagonal matrix $W_{\mathbf{q}}(t)$ has vanishing period average, and hence does not contribute to the parametric resonance to leading order. At the same time, the off-diagonal term in Eq. (10) will be dominant, whenever $h_{\mathbf{q}}(t)$ interferes with the phase term $e^{2iE_{\text{eff}}^{\text{Bog}}(\mathbf{q})t}$ constructively. This latter condition will give the resonant frequencies. In a sense, $h_{\mathbf{q}}(t)$ can be thought of as an effective periodic drive, the amplitude of which, multiplied by the prefactor $\sinh 2\theta_{\mathbf{q}} = g/E_{\text{eff}}^{\text{Bog}}(\mathbf{q})$, determine properties of the resonance, such as the maximally unstable mode.

Let us now analyze the function $h_{\mathbf{q}}(t)$ for the three types of drives we study in the main text:

Linear Drive—using the Jacobi-Anger identity, we have

$$\varepsilon_{\mathbf{q}}^{\text{lin}}(t) = 4J \left[\sin \frac{q_x}{2} \sin \left(\frac{q_x}{2} - K_0 \sin \omega t \right) + \sin^2 \frac{q_y}{2} \right],$$

$$h_{\mathbf{q}}^{\text{lin}}(t) = 4J \left[\cos(K_0 \sin \omega t) \sin^2 \frac{q_x}{2} + \sin^2 \frac{q_y}{2} \right] - \varepsilon_{\text{eff}}(\mathbf{q})$$

$$= 8J \sin^2 \frac{q_x}{2} \sum_{l=1}^{\infty} \mathcal{J}_{2l}(K_0) \cos(2l\omega t).$$

The fact that the system is driven only along the x -direction is reflected in the momentum dependence of $h_{\mathbf{q}}(t)$. Applying the rotating wave approximation to the time-periodic off-diagonal term in Eq. (10), we conclude that the dominant resonant harmonic is $l = 1$, which gives $\omega = E_{\text{eff}}^{\text{Bog}}(\mathbf{q})$. Using the relation $\sinh 2\theta_{\mathbf{q}} = g/E_{\text{eff}}^{\text{Bog}}(\mathbf{q}) = g/\omega$, the momentum-dependent instability rate reads [24]

$$s_{\text{lin}}(\mathbf{q}) = 4J \mathcal{J}_2(K_0) \sin^2 \left(\frac{q_x}{2} \right) \frac{g}{\omega}. \quad (12)$$

Clearly, there are many resonant modes, which satisfy the condition $\omega = E_{\text{eff}}^{\text{Bog}}(\mathbf{q}_{\text{res}})$. This condition is visualized as a plane, parallel to the (q_x, q_y) -plane, cutting through the dispersion relation: the resulting set of modes defines the solutions to the resonant condition.

However, out of all resonant modes, not all have the same instability rate, because $s_{\text{lin}}(\mathbf{q})$ is a function of \mathbf{q} . We are interested in the long-time BdG dynamics, which is dominated by the most unstable mode:

$$\gamma = \max_{\mathbf{q}} s(\mathbf{q}), \quad \mathbf{q}_{\text{mum}} = \text{argmax}_{\mathbf{q}} s(\mathbf{q}). \quad (13)$$

Since $s_{\text{lin}}(\mathbf{q})$ is a monotonic function of only q_x , the most unstable modes are those resonant modes, which have the largest q_x -component. Gradually increasing the drive frequency ω , one reaches a critical saturation frequency $\omega_c = E_{\text{eff}}^{\text{Bog}}(\pi, 0)$ at the edge of the Brillouin zone. Using the same analysis as in Ref. [24] and keeping in mind the existence of a continuous degree of freedom in the z (tubes) direction, one can show that for $\omega \geq \omega_c^{\text{lin}}$, we have $\mathbf{q}_{\text{mum}}^{\text{lin}} = (\pi, 0)$ and $\gamma_{\text{lin}} = 4J \mathcal{J}_2(K_0)g/\omega$, while for $\omega \leq \omega_c^{\text{lin}}$, we find $\mathbf{q}_{\text{mum}}^{\text{lin}} = 2 \arcsin \sqrt{\frac{\sqrt{g^2 + \omega^2} - g}{4J \mathcal{J}_0(K_0)}} (\pm 1, 0)$, and the rate is independent of the hopping and given by $\gamma_{\text{lin}} = (\sqrt{g^2 + \omega^2} - g) \frac{\mathcal{J}_2(K_0)}{\mathcal{J}_0(K_0)} \frac{g}{\omega}$.

Because $\mathbf{q}_{\text{mum}}^{\text{lin}}$ is located at the side edge of the 2D Brillouin zone, the saturation frequency for linear drive ω_c^{lin} equals half the effective 2D Bogoliubov bandwidth.

Diagonal Drive—similarly, we find

$$\varepsilon_{\mathbf{q}}^{\text{diag}}(t) = 4J \left[\sin \frac{q_x}{2} \sin \left(\frac{q_x}{2} - K_0 \sin \omega t \right) + \sin^2 \frac{q_y}{2} \sin \left(\frac{q_y}{2} - K_0 \sin \omega t \right) \right], \quad (14)$$

$$h_{\mathbf{q}}^{\text{diag}}(t) = 4J \cos(K_0 \sin \omega t) \left[\sin^2 \frac{q_x}{2} + \sin^2 \frac{q_y}{2} \right] - \varepsilon_{\text{eff}}(\mathbf{q})$$

$$= 8J \left(\sin^2 \frac{q_x}{2} + \sin^2 \frac{q_y}{2} \right) \sum_{l=1}^{\infty} \mathcal{J}_{2l}(K_0) \cos(2l\omega t).$$

It follows that

$$s_{\text{diag}}(\mathbf{q}) = 4J \left(\sin^2 \frac{q_x}{2} + \sin^2 \frac{q_y}{2} \right) \mathcal{J}_2(K_0) \frac{g}{\omega}.$$

As expected, the expression is symmetric w.r.t. exchanging q_x and q_y . The critical saturation frequency ω_c^{diag} is achieved when the maximally unstable mode reaches the Brillouin zone boundary $(q_x^{\text{mum}}, q_y^{\text{mum}}) = (\pi, \pi)$. For $\omega \geq \omega_c^{\text{diag}}$, the maximally unstable mode $\mathbf{q}_{\text{mum}}^{\text{diag}} = (\pi, \pi)$ and $\gamma_{\text{diag}} = 8J \mathcal{J}_2(K_0)g/\omega$. Notice that $\gamma_{\text{diag}} = 2\gamma_{\text{lin}}$ in this regime, due to the presence of the second shaken direction which enhances the instability rate. In the other regime,

$\omega \leq \omega_c^{\text{diag}}$, we find $\mathbf{q}_{\text{mum}}^{\text{diag}} = 2 \arcsin \sqrt{\frac{\sqrt{g^2 + \omega^2} - g}{8J \mathcal{J}_0(K_0)}} \times (\pm 1, \pm 1)$ and $\gamma_{\text{diag}} = (\sqrt{g^2 + \omega^2} - g) \frac{\mathcal{J}_2(K_0)}{\mathcal{J}_0(K_0)} \frac{g}{\omega}$. Here, $\gamma_{\text{diag}} = \gamma_{\text{lin}}$.

Because $\mathbf{q}_{\text{mum}}^{\text{diag}}$ is located at the corner of the 2D Brillouin zone, the saturation frequency for diagonal drive ω_c^{diag} equals the full effective 2D Bogoliubov bandwidth.

Circular Drive— we find:

$$\begin{aligned} \varepsilon_{\mathbf{q}}^{\text{circ}}(t) &= 4J \left[\sin \frac{q_x}{2} \sin \left(\frac{q_x}{2} - K_0 \sin \omega t \right) \right. \\ &\quad \left. + \sin^2 \frac{q_y}{2} \sin \left(\frac{q_y}{2} + K_0 \sin \omega t \right) \right], \\ h_{\mathbf{q}}^{\text{circ}}(t) &= 4J \left[\sin^2 \frac{q_x}{2} \cos(K_0 \sin \omega t) \right. \\ &\quad \left. + \sin^2 \frac{q_y}{2} \cos(K_0 \cos \omega t) \right] - \varepsilon_{\text{eff}}(\mathbf{q}) \\ &= 8J \left(\sin^2 \frac{q_x}{2} - \sin^2 \frac{q_y}{2} \right) \sum_{l=1}^{\infty} \mathcal{J}_{2l}(K_0) \cos(2l\omega t). \end{aligned} \quad (15)$$

It follows that

$$s_{\text{circ}}(\mathbf{q}) = 4J \left| \sin^2 \frac{q_x}{2} - \sin^2 \frac{q_y}{2} \right| \mathcal{J}_2(K_0) \frac{g}{\omega}.$$

Note the change in signature compared to the diagonal drive. Therefore, unlike the diagonal drive, and rather similar to the linear drive, the critical saturation frequency ω_c^{circ} is achieved when the maximally unstable mode reaches the side edge of the Brillouin zone $(q_x^{\text{mum}}, q_y^{\text{mum}}) = \{(\pi, 0), (0, \pi)\}$. For $\omega \geq \omega_c^{\text{circ}}$, the maximally unstable modes are $\mathbf{q}_{\text{mum}}^{\text{circ}} = \{(\pi, 0), (0, \pi)\}$ and $\gamma_{\text{circ}} = 4J\mathcal{J}_2(K_0)g/\omega$. Notice that $\gamma_{\text{diag}} = 2\gamma_{\text{circ}}$ and $\gamma_{\text{circ}} = \gamma_{\text{lin}}$ in this regime, due to the circular form of the drive. In the other regime, $\omega \leq \omega_c^{\text{circ}}$, we find

$$\mathbf{q}_{\text{mum}}^{\text{circ}} = 2 \arcsin \sqrt{\frac{\sqrt{g^2 + \omega^2} - g}{4J\mathcal{J}_0(K_0)}} \times \{(\pm 1, 0), (0, \pm 1)\} \text{ and } \gamma_{\text{circ}} = (\sqrt{g^2 + \omega^2} - g) \frac{\mathcal{J}_2(K_0)}{\mathcal{J}_0(K_0)} \frac{g}{\omega}. \text{ Here, } \gamma_{\text{diag}} = \gamma_{\text{circ}} = \gamma_{\text{lin}}.$$

Because $\mathbf{q}_{\text{mum}}^{\text{circ}}$ is again located at the side edge of the 2D Brillouin zone, the saturation frequency for circular drive ω_c^{circ} equals half the effective 2D Bogoliubov bandwidth [similar to linear drive].

The above analysis was performed for a pure 2D lattice system. However, adding the transverse tube di-

mension is straightforward and does not change these conclusions [24], cf. Fig. 10 for a numerical simulation in the presence of a tube-like transverse direction.

Finally, note that the instability rate γ derived above corresponds to the growth rate of the bosonic operators $b_{\mathbf{q}}(t)$ and $b_{\mathbf{q}}^\dagger(t)$, which are not observable. Observables, consisting of bilinears of the bosonic operators, will thus display a parametric instability rate

$$\Gamma_{\text{mum}} = 2\gamma \quad (16)$$

for each maximally unstable mode.

Furthermore, an atom number variation per unit time (an experimental rate) will depend on how many such modes are present simultaneously. For the linear drive, there exists a single maximally unstable mode $(\pi, 0)$, while there exist two equally unstable modes for the diagonal drive $(\{(\pi, \pi), (-\pi, \pi)\})$ and for the circular drive $(\{(\pi, 0), (0, \pi)\})$. Therefore we expect an additional factor of 2 in Γ_{mum} for the two 2D drives. In total:

$$\Gamma_{\text{mum}}^{\text{lin}} = 2\gamma_{\text{lin}}, \quad (17)$$

$$\Gamma_{\text{mum}}^{\text{diag}} = 4\gamma_{\text{diag}}, \quad (18)$$

$$\Gamma_{\text{mum}}^{\text{circ}} = 4\gamma_{\text{circ}}. \quad (19)$$

As a final remark, we note that while a choice of small K_0 explores the 2D quasi-parabolic dispersion found at small BEC momenta, the above derivation shows that the problem still cannot be treated as rotationally-symmetric: the resonant modes are determined by the drive frequency through the resonance condition. Therefore, even for small drive amplitude K_0 , resonant Bogoliubov modes can be in the middle of the dispersion, where rotational symmetry is lost. Hence, the distinctions between 1D and 2D drives are expected, and observed, to hold even in the small K_0 limit.