

Global Well-Posedness and Large-Time Behavior of 1D Compressible Navier-Stokes System with Density-Depending Viscosity and Vacuum in Unbounded Domains*

Kexin Li[†] Boqiang Lü[‡] Yixuan Wang[§]

Abstract

We consider the Cauchy problem for one-dimensional (1D) barotropic compressible Navier-Stokes equations with density-dependent viscosity and large external force. Under a general assumption on the density-dependent viscosity, we prove that the Cauchy problem admits a unique global strong (classical) solution for the large initial data with vacuum. Moreover, the density is proved to be bounded from above time-independently. As a consequence, we obtain the large time behavior of the solution without external forces.

Keywords: 1D compressible Navier-Stokes equations; global well-posedness; large initial data; vacuum; density-dependent viscosity.

1 Introduction and main results

The motion of a one-dimensional viscous compressible barotropic fluid is governed by the following compressible Navier-Stokes equations:

$$\begin{cases} \rho_t + (\rho u)_x = 0, \\ (\rho u)_t + (\rho u^2)_x + [P(\rho)]_x = [\mu(\rho)u_x]_x + \rho f. \end{cases} \quad (1.1)$$

Here, $t \geq 0$ is time, $x \in \mathbb{R} = (-\infty, \infty)$ is the spatial coordinate, and $\rho(x, t) \geq 0$, $u(x, t)$ and $P(\rho) = A\rho^\gamma$ ($A > 0$, $\gamma > 1$) are the fluid density, velocity and pressure, respectively. Without loss of generality, it is assumed that $A = 1$. The viscosity $\mu(\cdot)$ is a function of the density ρ . The external force $f = f(x)$ is a known function. We look for the solutions (ρ, u) to the Cauchy problem for (1.1) with the initial condition

$$(\rho, u)|_{t=0} = (\rho_0(x), u_0(x)), \quad \text{for } x \in \mathbb{R}, \quad (1.2)$$

*K. Li is partially supported by Undergraduate Research Fund of BNU 2017-150; B. Lü is supported by NNSFC (Nos. 11601218 & 11771382), Science and Technology Project of Jiangxi Provincial Education Department (No. GJJ160719).

[†]School of Mathematical Sciences, Beijing Normal University, Beijing 100875, P. R. China (kexinli198@163.com)

[‡]College of Mathematics and Information Science, Nanchang Hangkong University, Nanchang 330063, P. R. China (lvbq86@163.com).

[§]Institute of Applied Mathematics, AMSS, Chinese Academy of Sciences, Beijing 100190, P. R. China (wangyixuan_14@163.com).

and the following far field behavior

$$u(x, t) \rightarrow 0, \quad \rho(x, t) \rightarrow \tilde{\rho} > 0, \quad \text{as } |x| \rightarrow \infty, \quad (1.3)$$

where $\tilde{\rho}$ is a given positive constant.

There is huge literature on the studies of the global existence and large time behavior of solutions to the 1D compressible Navier-Stokes equations. For constant viscosity μ and the initial data away from vacuum, the problems are addressed by Kanel [10] for sufficiently smooth data, and by Serre [19,20] and Hoff [6] for discontinuous initial data. On the other hand, if μ depends on ρ and admits a positive constant lower bound, the global well-posedness and large time behavior of solutions away from vacuum were discussed in [1, 2, 21] and the references therein. However, when it comes to the case that vacuum is allowed initially, as emphasized in many papers related to compressible fluid dynamics [3, 5, 8, 9, 16, 22], the possible presence of vacuum is one of the major difficulties in discussing the well-posedness of solutions to the compressible Navier-Stokes equations. In the presence of vacuum, Ding-Wen-Zhu [4] considered the global existence of classical solutions to 1D compressible Navier-Stokes equations in bounded domains, provided that $\mu \in C^2[0, \infty)$ satisfies

$$0 < \bar{\mu} \leq \mu(\rho) \leq C(1 + P(\rho)). \quad (1.4)$$

Recently, for general μ , we [17] establish not only the global existence but also the large-time behavior for classical solutions containing vacuum to the initial boundary value problem for 1D compressible Navier-Stokes equations. For the Cauchy problem (1.1)–(1.3) without external force ($f = 0$), Ye [23] studies the global classical large solutions under the following restriction on $\mu(\rho)$:

$$\mu(\rho) = 1 + \rho^\beta, \quad 0 \leq \beta < \gamma. \quad (1.5)$$

However, both the uniform upper bound of density and the large time behavior of solutions are not obtained in [23].

In this paper, for more general density-dependent viscosity (see (1.6)) and large external force, we will derive the uniform upper bound of density and thus prove the global well-posedness of strong (classical) large solutions containing vacuum to the Cauchy problem (1.1)–(1.3). Before stating the main results, we first explain the notations and conventions used throughout this paper. We set

$$D_t \triangleq \frac{\partial}{\partial t} + u \frac{\partial}{\partial x}, \quad \dot{v} \triangleq v_t + uv_x.$$

For $1 \leq r \leq \infty$ and $k \geq 1$, we adopt the following simplified notations:

$$L^r = L^r(\mathbb{R}), \quad W^{k,r} = W^{k,r}(\mathbb{R}), \quad H^k = W^{k,2}(\mathbb{R}).$$

The first result concerns the global existence of strong solutions to the Cauchy problem (1.1)–(1.3).

Theorem 1.1 *Suppose that $\int_{-\infty}^x f(y)dy \in H^2$ and the viscosity $\mu(\rho) \in C^1[0, \infty)$ satisfies*

$$0 < \bar{\mu} \leq \mu(\rho) \leq \lambda_0 \int_1^\rho \mu(s)ds + \lambda_1, \quad (1.6)$$

for some constants $\bar{\mu} > 0$, $\lambda_0 \geq 0$, and $\lambda_1 > 0$. Let the initial data (ρ_0, u_0) satisfy

$$\rho_0 \geq 0, \quad \rho_0 - \tilde{\rho} \in H^1, \quad u_0 \in H^1. \quad (1.7)$$

Then, there exists a unique global strong solution (ρ, u) to the Cauchy problem (1.1)–(1.3) satisfying for any $0 < T < \infty$,

$$\begin{cases} \rho - \tilde{\rho} \in C([0, T]; H^1), & \rho_t \in L^\infty(0, T; L^2), \\ u \in L^\infty(0, T; H^1) \cap L^2(0, T; H^2), \\ \sqrt{t}u \in L^\infty(0, T; H^2), & \sqrt{t}u_t \in L^2(0, T; H^1). \end{cases} \quad (1.8)$$

Moreover, the density remains uniformly bounded for all time, that is

$$\sup_{0 \leq t < \infty} \|\rho(\cdot, t)\|_{L^\infty} < \infty. \quad (1.9)$$

Similar to [17, Theorem 1.2], one can obtain directly the following result which shows that the strong solutions obtained by Theorem 1.1 become classical provided initial data (ρ_0, u_0) satisfy some additional compatibility conditions.

Theorem 1.2 *In addition to the conditions of Theorem 1.1, suppose that $\mu(\rho) \in C^2[0, \infty)$ and that the initial data (ρ_0, u_0) satisfy*

$$(\rho_0 - \tilde{\rho}, P(\rho_0) - P(\tilde{\rho})) \in H^2, \quad u_0 \in H^2, \quad (1.10)$$

and the compatibility condition:

$$[\mu(\rho_0)u_{0x}]_x - [P(\rho_0)]_x = \sqrt{\rho_0}g(x), \quad x \in \mathbb{R}, \quad (1.11)$$

for a given function $g \in L^2$. Then, the strong solution obtained in Theorem 1.1 becomes classical and satisfies for any $0 < \tau < T < \infty$,

$$\begin{cases} (\rho - \tilde{\rho}, P(\rho) - P(\tilde{\rho})) \in C([0, T]; H^2), \\ u \in C([0, T]; H^2) \cap L^2(\tau, T; H^3), \\ u_t \in L^2(0, T; H^1), & \sqrt{t}u \in L^\infty(0, T; H^3), \\ \sqrt{t}u_t \in L^\infty(0, T; H^1) \cap L^2(0, T; H^2), & \sqrt{t}\sqrt{\rho}u_{tt} \in L^2(0, T; L^2). \end{cases} \quad (1.12)$$

When there is no external force, that is $f \equiv 0$ in (1.1), we can obtain the large time behavior of the strong solutions to the Cauchy problem (1.1)–(1.3).

Theorem 1.3 *In addition to the conditions of Theorem 1.1, suppose that $f \equiv 0$. Then the following large time behavior holds for strong solution (ρ, u) to the Cauchy problem (1.1)–(1.3) obtained by Theorem 1.1:*

$$\lim_{t \rightarrow \infty} (\|\rho - \tilde{\rho}\|_{L^p} + \|u_x\|_{L^2 \cap L^p}) = 0, \quad \forall p > 2. \quad (1.13)$$

Moreover, if there exists some point $x_0 \in (-\infty, \infty)$ such that $\rho_0(x_0) = 0$, the spatial gradient of the density the unique strong solution (ρ, u) to the problem (1.1)–(1.3) has to blow up as $t \rightarrow \infty$ in the following sense,

$$\lim_{t \rightarrow \infty} \|\rho_x(\cdot, t)\|_{L^2} = \infty. \quad (1.14)$$

A few remarks are in order:

Remark 1.1 *It should be noted here that we obtain a completely new uniform upper bound of density (1.9), which is in sharp contrast to [4, 23] where the upper bound of the density are time-dependent. This is crucial for studying the large time behavior of the solutions.*

Remark 1.2 *We want to point out that our restriction on $\mu(\rho)$ in (1.6) is much more general in comparison with those of [4, 23] (see (1.5) and (1.4)). Here, we will list some special cases satisfying (1.6) as follows:*

- $\mu(\rho) = \hat{\mu}$ is a positive constant. Let $\lambda_0 = 0$ and $\lambda_1 = \hat{\mu}$. Then

$$\lambda_0 \int_1^\rho \mu(s) ds + \lambda_1 = \hat{\mu} = \mu(\rho).$$

- $\mu(\rho) = 1 + \rho^a$ for any $a \geq 0$. Choosing $\lambda_0 = 1 + a$ and $\lambda_1 = 4 + a$ leads to

$$\lambda_0 \int_1^\rho \mu(s) ds + \lambda_1 = (1 + a)\rho + \rho^{1+a} + 2 \geq 1 + \rho^a = \mu(\rho).$$

- $\mu(\rho) = e^\rho$. Letting $\lambda_0 = 1$ and $\lambda_1 = e$ yields

$$\lambda_0 \int_1^\rho \mu(s) ds + \lambda_1 = e^\rho = \mu(\rho).$$

Hence, it is clear that the density-dependent viscosity $\mu(\rho)$ in [4, 23] are all included in our results.

Remark 1.3 *To obtain the global existence of strong solutions in Theorem 1.1, we do not need the additional compatibility condition (1.11), which is required for discussing the global classical solutions in Theorem 1.2. Theorems 1.1–1.2 show that how the compatibility condition (1.11) plays its role in studying the well-posedness of solutions with initial vacuum.*

Remark 1.4 *It should be noted here that the solution (ρ, u) obtained in Theorem 1.1 is actually a classical one. Indeed, the Sobolev embedding theorem together with the regularities of (ρ, u) in (1.12) shows that*

$$(\rho, P, u) \in C([0, T]; C^{1+\frac{1}{2}}), \quad \rho_t \in C([0, T]; C^{\frac{1}{2}}). \quad (1.15)$$

Furthermore, one can deduce from (1.12) that for any $0 < \tau < T$,

$$u \in L^\infty(\tau, T; H^3), \quad u_t \in L^\infty(\tau, T; H^1) \cap L^2(\tau, T; H^2),$$

which yields that for $1 < r < 2$,

$$u \in C([\tau, T]; C^{2,r}[0, 1]), \quad u_t \in C([\tau, T]; C^r[0, 1]). \quad (1.16)$$

Hence, one can deduce from (1.15)–(1.16) that (ρ, u) is the classical solution to (1.1)–(1.3).

We now make some comments on the analysis of this paper. Since the local well-posedness result of classical solutions away from vacuum is well-known (see [3] or Lemma 2.1 below), to extend local solutions to be a global one which will eventually contain vacuum after letting the lower bound of initial density go to zero, we need some global a priori estimates which are independent of the lower bound of density. As mentioned in many papers (see [7, 12]), the main difficulty comes from initial vacuum, density-dependent viscosity, large external force, and the unboundedness of the domain. Motivated by our previous result [17], we find that the key issue is to derive both the time-independent upper bound of the density and the time-dependent derivative ones of the solutions. Motivated by [11], we localize the problem on bounded domains and use the method develop in our previous work [17] to get the upper bound of the density independently of x and thus the uniform bound of density on the whole \mathbb{R} due to the arbitrariness of x (see Lemma 2.3). Furthermore, with the help of far field behavior of ρ , we can bound the L^2 -norm of u in terms of $\|\rho^{1/2}u\|_{L^2}$ and $\|u_x\|_{L^2}$ (see (2.23)). Following the methods used in [12, 15], we use the material derivative \dot{u} instead of the usually u_t and succeed in obtaining the derivative estimates on (ρ, u) . With the desired global a priori estimates independent of the lower bound of density at hand, we can thus prove that Theorems 1.1 and 1.2 hold for any time $T > 0$ which completes the proof of Theorems 1.1 and 1.2 (see Section 3). Finally, for the case without external force ($f \equiv 0$), we can establish the time-independent lower order estimates (see (4.1) and (4.2)). Using the methods due to [9, 12] and the key time-independent a priori estimates, we prove Theorem 1.3 in Section 4.

The rest of the paper is organized as follows. Section 2 is devoted to derive the necessary a priori estimates on smooth solutions. Theorems 1.1 and 1.3 are proved in Sections 3 and 4, respectively.

2 A priori estimates

First, for the initial density strictly away from vacuum, we state the following local well-posedness theory of strong solutions, whose proof can be obtained by similar arguments as in [3, 18].

Lemma 2.1 *Let $\mu(\rho) \in C^2[0, \infty)$. Assume that $\tilde{f} \in H^2$ and the initial data (ρ_0, u_0) satisfies*

$$(\rho_0 - \tilde{\rho}, P(\rho_0) - P(\tilde{\rho}), u_0) \in H^2, \quad \inf_{x \in \mathbb{R}} \rho_0(x) > 0,$$

where $\tilde{\rho} > 0$ is a given constant. Then, there exists a small time $T_0 > 0$ such that the Cauchy problem (1.1)–(1.3) has a unique strong solution (ρ, u) on $\mathbb{R} \times (0, T_0]$ satisfying

$$\begin{cases} \rho - \tilde{\rho}, P(\rho) - P(\tilde{\rho}) \in C([0, T_0]; H^2), \\ u \in C([0, T_0]; H^2) \cap L^2(0, T_0; H^3), \\ \rho_t \in C([0, T_0]; H^1), \quad u_t \in C([0, T_0]; L^2) \cap L^2(0, T_0; H^1). \end{cases}$$

In this section, we will establish some a priori bounds for strong solutions (ρ, u) to the Cauchy problem (1.1)–(1.3) on $\mathbb{R} \times [0, T]$ whose existence is guaranteed by Lemma 2.1.

2.1 A priori estimates(I): Time-independent a priori estimates

In this subsection, we will use the convention that C denotes a generic positive constant depending on the initial data (ρ_0, u_0) and some known constants but independent of both $\inf_{x \in \mathbb{R}} \rho_0(x)$ and T , and use $C(\alpha)$ to emphasize that C depends on α .

We start with the following energy estimate for the solutions (ρ, u) .

Lemma 2.2 *There is a positive constant C depending on $\gamma, \bar{\mu}, \tilde{\rho}, \|\rho_0 - \tilde{\rho}\|_{H^1}, \|u_0\|_{H^1}$, and $\|\tilde{f}\|_{H^2}$ such that*

$$\sup_{0 \leq t \leq T} \int_{-\infty}^{+\infty} (\rho u^2 + G(\rho)) dx + \int_0^T \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx dt \leq C, \quad (2.1)$$

where and in what follows $\tilde{f} \triangleq \int_{-\infty}^x f(y) dy$ and G denotes the potential energy density given by

$$G(\rho) = \rho \int_{\tilde{\rho}}^{\rho} \frac{P(s) - P(\tilde{\rho})}{s^2} ds = \frac{1}{\gamma - 1} [P(\rho) - P(\tilde{\rho}) - \gamma \tilde{\rho}^{\gamma-1} (\rho - \tilde{\rho})]. \quad (2.2)$$

Proof. Multiplying (1.1)₁ and (1.1)₂ by $G'(\rho)$ and u respectively, we obtain after using integration by parts and the far-field condition (1.3) that

$$\frac{d}{dt} \int_{-\infty}^{+\infty} \left(\frac{1}{2} \rho u^2 + G(\rho) \right) dx + \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx = \frac{d}{dt} \int_{-\infty}^{+\infty} (\rho - \tilde{\rho}) \tilde{f} dx,$$

which yields

$$\begin{aligned} & \sup_{0 \leq t \leq T} \int_{-\infty}^{+\infty} \left(\frac{1}{2} \rho u^2 + G(\rho) \right) dx + \int_0^T \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx dt \\ & \leq C + \int_{-\infty}^{+\infty} (\rho - \tilde{\rho}) \tilde{f} dx - \int_{-\infty}^{+\infty} (\rho_0 - \tilde{\rho}) \tilde{f} dx \\ & \leq C + \int_{-\infty}^{+\infty} (\rho - \tilde{\rho}) \tilde{f} dx + C \|\rho_0 - \tilde{\rho}\|_{L^2}^2 + C \|\tilde{f}\|_{L^2}^2 \\ & \leq C + \int_{R_1} (\rho - \tilde{\rho}) \tilde{f} dx + \int_{\mathbb{R} \setminus R_1} (\rho - \tilde{\rho}) \tilde{f} dx, \end{aligned} \quad (2.3)$$

with

$$R_1 \triangleq \{x \in \mathbb{R} \mid \rho^{\gamma-1} \leq \gamma^2 (\tilde{\rho}^{\gamma-1} + 1)\}. \quad (2.4)$$

Then, on the one hand, when $0 \leq \rho \leq M$ for some positive constant $M > 0$, there are positive constants K_1 and K_2 depending only on $\tilde{\rho}$ and M such that

$$K_1 (\rho - \tilde{\rho})^2 \leq G(\rho) \leq K_2 (\rho - \tilde{\rho})^2, \quad (2.5)$$

which together with Young's inequality implies that

$$\begin{aligned} \int_{R_1} |\rho - \tilde{\rho}| |\tilde{f}| dx & \leq \varepsilon \|\rho - \tilde{\rho}\|_{L^2(R_1)}^2 + C(\varepsilon) \|\tilde{f}\|_{L^2(R_1)}^2 \\ & \leq C\varepsilon \int_{-\infty}^{+\infty} G(\rho) dx + C(\varepsilon). \end{aligned} \quad (2.6)$$

On the other hand, since $\rho > \tilde{\rho}$ in $\mathbb{R} \setminus R_1$, we have

$$\begin{aligned}
\int_{\mathbb{R} \setminus R_1} |\rho - \tilde{\rho}| |\tilde{f}| dx &\leq 2 \int_{\mathbb{R} \setminus R_1} \rho |\tilde{f}| dx \\
&\leq \varepsilon \int_{\mathbb{R} \setminus R_1} \rho^{\frac{2\gamma-1}{\gamma}} dx + C(\varepsilon) \int_{\mathbb{R} \setminus R_1} |\tilde{f}|^{\frac{2\gamma-1}{\gamma-1}} dx \\
&\leq \varepsilon \int_{\mathbb{R} \setminus R_1} \rho^{\frac{2\gamma-1}{\gamma}} dx + C(\varepsilon) \|\tilde{f}\|_{L^\infty(\mathbb{R} \setminus R_1)}^{\frac{1}{\gamma-1}} \|\tilde{f}\|_{L^2(\mathbb{R} \setminus R_1)}^2 \\
&\leq \varepsilon \int_{-\infty}^{+\infty} G(\rho) dx + C(\varepsilon),
\end{aligned} \tag{2.7}$$

where in the last inequality one has used the following fact:

$$\rho^{\frac{2\gamma-1}{\gamma}} \leq \rho \left(\frac{1}{\gamma} \rho^{\gamma-1} + \frac{\gamma-1}{\gamma} \right) \leq \frac{1}{\gamma-1} (\rho^\gamma - \gamma \tilde{\rho}^{\gamma-1} \rho) \leq G(\rho), \quad x \in \mathbb{R} \setminus R_1,$$

due to Young's inequality, (2.4), and (2.2).

Substituting (2.6) and (2.7) into (2.3), we obtain (2.1) after choosing ε suitably small. This completes the proof of Lemma 2.2. \square

Next, we derive the key uniform (in time) upper bound of the density, which is crucial for obtaining both the derivative estimates and the large time behavior of the solutions. This method is motivated by Kazhikhov [11].

Lemma 2.3 *There is a positive constant $\bar{\rho}$ depending only on γ , $\bar{\mu}$, $\tilde{\rho}$, $\|\rho_0 - \tilde{\rho}\|_{H^1}$, $\|u_0\|_{H^1}$, and $\|\tilde{f}\|_{H^2}$ such that*

$$0 \leq \rho(x, t) \leq \bar{\rho}, \quad \forall (x, t) \in \mathbb{R} \times [0, T]. \tag{2.8}$$

Proof. For any integer N , let $x \in [N, N+1]$. Integrating (1.1)₁ over (N, x) with respect to x leads to

$$-\mu(\rho)u_x + P + \left(\int_N^x \rho u dy \right)_t + \rho u^2 = (-\mu(\rho)u_x + P + \rho u^2)(N, t) + \int_N^x \rho f dy, \tag{2.9}$$

which in particular implies

$$\begin{aligned}
&(-\mu(\rho)u_x + P + \rho u^2)(N, t) \\
&= \int_N^{N+1} \left(-\mu(\rho)u_x + P + \rho u^2 + \int_N^x \rho f dy \right) dx + \left(\int_N^{N+1} \int_N^x \rho u dy dx \right)_t \\
&\leq - \int_N^{N+1} \mu(\rho)u_x dx + C + \left(\int_N^{N+1} \int_N^x \rho u dy dx \right)_t.
\end{aligned} \tag{2.10}$$

due to (2.1) and the following simple fact

$$\begin{aligned}
\int_N^{N+1} \rho^\gamma dx &\leq C + \int_N^{N+1} \mathbf{1}_{\{\rho^{\gamma-1} \geq 2\gamma \tilde{\rho}^{\gamma-1}\}} \rho^\gamma dy \\
&\leq C + 2(\gamma-1) \int_N^{N+1} G(\rho) dy \\
&\leq C.
\end{aligned} \tag{2.11}$$

Denoting

$$F(\rho) \triangleq \int_1^\rho \mu(s)s^{-1}ds,$$

one deduces from (1.1)₁ that

$$D_t F(\rho) = -\mu(\rho)u_x,$$

which together with (2.9) and (2.10) gives

$$\begin{aligned} D_t(F(\rho) + b_1(t)) + P &\leq - \int_N^{N+1} \mu(\rho)u_x dx + C \\ &\leq \int_N^{N+1} \mu(\rho)dx \int_N^{N+1} \mu(\rho)u_x^2 dx + C \\ &\leq \sup_{x \in \mathbb{R}} \frac{\mu(\rho)}{\rho + 1} \int_N^{N+1} (\rho + 1)dx \int_N^{N+1} \mu(\rho)u_x^2 dx + C \\ &\leq C \sup_{x \in \mathbb{R}} \frac{\mu(\rho)}{\rho + 1} \int_{-\infty}^{\infty} \mu(\rho)u_x^2 dx + C, \end{aligned} \tag{2.12}$$

where

$$b_1(t) \triangleq \int_N^x \rho u dy + \int_N^{N+1} \int_N^x \rho u dy dx$$

satisfies

$$\begin{aligned} |b_1(t)| &\leq C \int_N^{N+1} \rho |u| dy \\ &\leq C \int_N^{N+1} \rho dx + C \int_N^{N+1} \rho u^2 dx \leq C_1. \end{aligned} \tag{2.13}$$

due to (2.11) and (2.1).

Since μ satisfies (1.6), it holds

$$\begin{aligned} \sup_{x \in \mathbb{R}} \frac{\mu(\rho)}{\rho + 1} &\leq \lambda_0 \sup_{x \in \mathbb{R}} \left(\frac{1}{\rho + 1} \int_1^\rho \mu(s)ds \right) + \lambda_1 \\ &\leq \lambda_0 \sup_{x \in \mathbb{R}} \int_1^{\max\{\rho, 1\}} \frac{\mu(s)}{s} ds + \lambda_1, \end{aligned}$$

which together with (2.12) leads to

$$\begin{aligned} D_t(F(\rho) + b_1(t)) + P \\ \leq C_2 + C \left(\sup_{x \in \mathbb{R}} \int_1^{\max\{\rho, 1\}} \frac{\mu(s)}{s} ds + 1 \right) \int_{-\infty}^{\infty} \mu(\rho)u_x^2 dx, \end{aligned} \tag{2.14}$$

with constant $C_2 > 1$.

Choosing a constant $\nu \geq C_2^{1/\gamma}$ such that for all $\rho \geq \nu$,

$$P(\rho) - C_2 \geq 0, \tag{2.15}$$

and multiplying (2.14) by

$$H \triangleq (F(\rho) + b_1(t) - F(\nu) - C_1)_+, \tag{2.16}$$

we obtain after using (2.13) and (2.15) that

$$\begin{aligned} D_t H^2 &\leq CH \left(\sup_{x \in \mathbb{R}} \int_1^{\max\{\rho, 1\}} \frac{\mu(s)}{s} ds + 1 \right) \int_{-\infty}^{\infty} \mu(\rho) u_x^2 dx \\ &\leq C \sup_{x \in \mathbb{R}} H^2 \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx + C \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx. \end{aligned} \quad (2.17)$$

Integrating (2.17) over $(0, t)$ gives that for $x \in [N - 1, N]$,

$$H^2(x, t) \leq \bar{C} + \bar{C} \int_0^t \sup_{x \in \mathbb{R}} H^2 \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx dt, \quad (2.18)$$

where \bar{C} is a positive constant which is independent of N . Since N is arbitrary, the inequality (2.18) holds for all $x \in \mathbb{R}$, that is,

$$\sup_{x \in \mathbb{R}} H^2 \leq C + C \int_0^t \sup_{x \in \mathbb{R}} H^2 \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx dt,$$

which together with Gronwall's inequality and (2.1) yields that

$$\sup_{(x,t) \in \mathbb{R} \times (0, \infty)} H^2 \leq C. \quad (2.19)$$

Consequently, the desired (2.8) is a direct consequence of (2.16), (2.19), and (2.13). This finishes the proof of Lemma 2.3. \square

With the uniform upper bound of the density (2.8) at hand, we have the following time-independent bound on $\|\rho - \tilde{\rho}\|_{L^2}^2$.

Corollary 2.4 *There is a positive constant C depending on γ , $\bar{\mu}$, $\tilde{\rho}$, $\|\rho_0 - \tilde{\rho}\|_{H^1}$, $\|u_0\|_{H^1}$, and $\|\tilde{f}\|_{H^2}$ such that*

$$\sup_{0 \leq t \leq T} (\|\rho - \tilde{\rho}\|_{L^2}^2 + \|P(\rho) - P(\tilde{\rho})\|_{L^2}^2) \leq C. \quad (2.20)$$

Proof. It follows from (2.1) and (2.5) that

$$\begin{aligned} &\|\rho - \tilde{\rho}\|_{L^2}^2 + \|P(\rho) - P(\tilde{\rho})\|_{L^2}^2 \\ &\leq C \int_{-\infty}^{+\infty} G(\rho) dx + \|(\gamma - 1)G(\rho) + \gamma \tilde{\rho}^{\gamma-1} \rho - \gamma \tilde{\rho}^\gamma\|_{L^2}^2 \\ &\leq C \int_{-\infty}^{+\infty} G(\rho) dx \leq C, \end{aligned}$$

which completes the proof of Corollary 2.4. \square

2.2 A priori estimates(II): Time-dependent a priori estimates

In this subsection, we now proceed to derive the derivative estimates of the strong (ρ, u) to the Cauchy problem (1.1)–(1.3). In what follows, the constant C depends on T but is still independent of $\inf_{x \in \mathbb{R}} \rho_0(x)$.

Lemma 2.5 *There is a positive constant C depending on T , γ , $\bar{\mu}$, $\tilde{\rho}$, $\|\rho_0 - \tilde{\rho}\|_{H^1}$, $\|u_0\|_{H^1}$, and $\|\tilde{f}\|_{H^2}$ such that*

$$\sup_{0 \leq t \leq T} \|u\|_{H^1}^2 + \int_0^T \|\rho^{1/2} \dot{u}\|_{L^2}^2 dt \leq C(T). \quad (2.21)$$

Proof. First, multiplying (1.1)₂ by \dot{u} and integrating the resulting equation by parts yield

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx + \int_{-\infty}^{+\infty} \rho \dot{u}^2 dx \\ &= \frac{d}{dt} \left(\int_{-\infty}^{+\infty} [P(\rho) - P(\tilde{\rho})] u_x dx + \int_{-\infty}^{+\infty} \rho f u dx \right) \\ & \quad - \frac{1}{2} \int_{-\infty}^{+\infty} [\mu(\rho) + \mu'(\rho)\rho] u_x^3 dx + \gamma \int_{-\infty}^{+\infty} P(\rho) u_x^2 dx - \int_{-\infty}^{+\infty} \rho u^2 f_x dx \\ &\leq \frac{d}{dt} \left(\int_{-\infty}^{+\infty} [P(\rho) - P(\tilde{\rho})] u_x dx + \int_{-\infty}^{+\infty} \rho f u dx \right) \\ & \quad + C \|u_x\|_{L^3}^3 + C \|u_x\|_{L^2}^2 + C \|u\|_{L^\infty} \|\rho^{1/2} u\|_{L^2} \|f_x\|_{L^2} \\ &\leq \frac{d}{dt} \left(\int_{-\infty}^{+\infty} [P(\rho) - P(\tilde{\rho})] u_x dx + \int_{-\infty}^{+\infty} \rho f u dx \right) \\ & \quad + C \|u_x\|_{L^\infty} \|u_x\|_{L^2}^2 + C \|u_x\|_{L^2}^2 + C, \end{aligned} \quad (2.22)$$

where in the last inequality one has used the following fact:

$$\begin{aligned} \|u\|_{L^2}^2 + \|u\|_{L^\infty} &= \tilde{\rho}^{-1} \left(\int_{-\infty}^{+\infty} \rho u^2 dx + \int_{-\infty}^{+\infty} (\tilde{\rho} - \rho) u^2 dx \right) + \|u\|_{L^\infty} \\ &\leq C \|\rho^{1/2} u\|_{L^2}^2 + C \|\rho - \tilde{\rho}\|_{L^2} \|u\|_{L^2} \|u\|_{L^\infty} + \|u\|_{L^\infty} \\ &\leq C \|\rho^{1/2} u\|_{L^2}^2 + C \|u\|_{L^2}^{3/2} \|u_x\|_{L^2}^{1/2} + C \|u\|_{L^2}^{1/2} \|u_x\|_{L^2}^{1/2} \\ &\leq C + C \|u_x\|_{L^2}^2 + \frac{1}{2} \|u\|_{L^2}^2. \end{aligned} \quad (2.23)$$

due to (2.20), (2.1), and the Sobolev inequality. Furthermore, using (1.6), (2.1), (1.1)₂, and (2.20), we get

$$\begin{aligned} \|u_x\|_{L^\infty} &\leq C \|\mu(\rho) u_x - P(\rho) + P(\tilde{\rho})\|_{L^\infty} + C \|P(\rho) - P(\tilde{\rho})\|_{L^\infty} \\ &\leq C \|\mu(\rho) u_x - [P(\rho) - P(\tilde{\rho})]\|_{L^2} + C \|(\mu(\rho) u_x - P(\rho))_x\|_{L^2} + C \\ &\leq C \|\mu(\rho) u_x\|_{L^2} + C \|P(\rho) - P(\tilde{\rho})\|_{L^2} + C \|\rho \dot{u}\|_{L^2} + C \|\rho f\|_{L^2} + C \\ &\leq C \|\sqrt{\mu(\rho)} u_x\|_{L^2} + C \|\rho^{1/2} \dot{u}\|_{L^2} + C. \end{aligned} \quad (2.24)$$

Then, putting (2.24) into (2.22), we obtain after using Young's inequality that

$$\frac{d}{dt} B(t) + \frac{1}{2} \int_{-\infty}^{+\infty} \rho \dot{u}^2 dx \leq C + C \|\sqrt{\mu(\rho)} u_x\|_{L^2}^2 + C \|\sqrt{\mu(\rho)} u_x\|_{L^2}^4, \quad (2.25)$$

where

$$B(t) \triangleq \frac{1}{2} \int_{-\infty}^{+\infty} \mu(\rho) u_x^2 dx - \int_{-\infty}^{+\infty} [P(\rho) - P(\tilde{\rho})] u_x dx - \int_{-\infty}^{+\infty} \rho f u dx \quad (2.26)$$

satisfies

$$\frac{1}{4}\|\sqrt{\mu(\rho)}u_x\|_{L^2}^2 - C \leq B(t) \leq C\|\sqrt{\mu(\rho)}u_x\|_{L^2}^2 + C. \quad (2.27)$$

Indeed, it follows from (1.6), (2.1), and (2.20) that

$$\begin{aligned} & \int_{-\infty}^{+\infty} [P(\rho) - P(\tilde{\rho})]u_x dx + \int_{-\infty}^{+\infty} \rho f u dx \\ & \leq \frac{1}{4} \int_{-\infty}^{+\infty} \mu(\rho)u_x^2 dx + C\|P(\rho) - P(\tilde{\rho})\|_{L^2}^2 + C\|\rho^{1/2}u\|_{L^2}\|f\|_{L^2} \\ & \leq \frac{1}{4} \int_{-\infty}^{+\infty} \mu(\rho)u_x^2 dx + C, \end{aligned}$$

which along with (2.26) yields (2.27).

Finally, Gronwall's inequality combined with (2.25), (2.27), (1.6), (2.1), and (2.23) yields (2.21) and completes the proof of Lemma 2.5. \square

Lemma 2.6 *There is a positive constant C depending on T , γ , $\bar{\mu}$, $\tilde{\rho}$, $\|\rho_0 - \tilde{\rho}\|_{H^1}$, $\|u_0\|_{H^1}$, and $\|\tilde{f}\|_{H^2}$ such that*

$$\sup_{0 \leq t \leq T} (\|\rho_x\|_{L^2}^2 + \|\rho_t\|_{L^2}^2) \leq C, \quad (2.28)$$

Proof. Differentiating (1.1)₁ with respect to x gives

$$\rho_{xt} + \rho_{xx}u + 2\rho_x u_x + \rho u_{xx} = 0. \quad (2.29)$$

Multiplying (2.29) by ρ_x and integrating the resultant equation by parts yield that

$$\frac{d}{dt}\|\rho_x\|_{L^2} \leq C\|u_x\|_{L^\infty}\|\rho_x\|_{L^2} + C\|u_{xx}\|_{L^2}, \quad (2.30)$$

due to (2.8).

Then, it follows from (1.1)₂ that

$$\mu(\rho)u_{xx} = \rho\dot{u} + P_x - \rho f - \mu'(\rho)\rho_x u_x,$$

which together with (1.6), and (2.8) yields that

$$\|u_{xx}\|_{L^2} \leq C\|\rho^{1/2}\dot{u}\|_{L^2} + C\|\rho_x\|_{L^2} + C\|f\|_{L^2} + C\|\rho_x\|_{L^2}\|u_x\|_{L^\infty}. \quad (2.31)$$

Submitting (2.31) into (2.30), one obtains after using (2.24), (2.21), and Gronwall's inequality

$$\sup_{0 \leq t \leq T} \|\rho_x\|_{L^2} \leq C, \quad (2.32)$$

which along with (1.1)₁, (2.8), (2.23), and (2.21) leads to

$$\|\rho_t\|_{L^2} \leq C\|u\|_{L^\infty}\|\rho_x\|_{L^2} + C\|u_x\|_{L^2} \leq C.$$

Combining this with (2.32) shows (2.28) and finishes the proof of Lemma 2.6. \square

Lemma 2.7 *There is a positive constant C depending on T , γ , $\bar{\mu}$, $\tilde{\rho}$, $\|\rho_0 - \tilde{\rho}\|_{H^1}$, $\|u_0\|_{H^1}$, and $\|\tilde{f}\|_{H^2}$ such that*

$$\sup_{0 \leq t \leq T} \sigma \|u_{xx}\|_{L^2}^2 + \int_0^T (\|u_{xx}\|_{L^2}^2 + \sigma \|u_t\|_{H^1}^2) dt \leq C, \quad (2.33)$$

with $\sigma(t) \triangleq \min\{1, t\}$.

Proof. First, operating $\partial/\partial t + (u \cdot)_x$ to (1.1)₂ yields

$$\rho \dot{u}_t + \rho u \dot{u}_x - [\mu(\rho) \dot{u}_x]_x = -\gamma [P(\rho) u_x]_x - [(\mu(\rho) + \mu'(\rho)\rho) u_x^2]_x + \rho u f_x. \quad (2.34)$$

Multiplying (2.34) by \dot{u} , one gets after using (2.23), (2.24), and (2.21) that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{-\infty}^{+\infty} \rho |\dot{u}|^2 dx + \int_{-\infty}^{+\infty} \mu(\rho) |\dot{u}_x|^2 dx \\ &= \gamma \int_{-\infty}^{+\infty} P(\rho) u_x \dot{u}_x dx + \int_{-\infty}^{+\infty} (\mu(\rho) + \mu'(\rho)\rho) u_x^2 \dot{u}_x dx + \int_{-\infty}^{+\infty} \rho u f_x \dot{u} dx \\ &\leq C \|\dot{u}_x\|_{L^2} + C \|\dot{u}_x\|_{L^2} \|u_x\|_{L^\infty} + C \|u\|_{L^\infty} \|f_x\|_{L^2} \|\rho^{1/2} \dot{u}\|_{L^2} \\ &\leq \frac{1}{2} \|\sqrt{\mu(\rho)} \dot{u}_x\|_{L^2}^2 + C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C. \end{aligned} \quad (2.35)$$

Then, multiplying (2.35) by σ and integrating the resulting inequality over $(0, T)$, we obtain after using (2.21) that

$$\sup_{0 \leq t \leq T} \sigma \int_{-\infty}^{+\infty} \rho |\dot{u}|^2 dx + \int_0^T \sigma \int_{-\infty}^{+\infty} \mu(\rho) |\dot{u}_x|^2 dx dt \leq C(T), \quad (2.36)$$

which along with (2.23) and (2.24) implies that

$$\sup_{0 \leq t \leq T} \sigma \|u_x\|_{L^\infty}^2 \leq C(T). \quad (2.37)$$

Finally, it follows from direct calculations, (2.21), and (2.23) that

$$\begin{aligned} \|\rho^{1/2} u_t\|_{L^2}^2 &\leq C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C \|\rho\|_{L^\infty} \|u\|_{L^\infty}^2 \|u_x\|_{L^2}^2 \\ &\leq C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C, \end{aligned}$$

which gives

$$\begin{aligned} \|u_t\|_{H^1}^2 &\leq C \int |\tilde{\rho} - \rho| u_t^2 dx + C \int \rho u_t^2 dx + C \|u_{xt}\|_{L^2}^2 \\ &\leq C \|\rho - \tilde{\rho}\|_{L^2} \|u_t\|_{L^2}^{3/2} \|u_{xt}\|_{L^2}^{1/2} + C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C + C \|u_{xt}\|_{L^2}^2 \\ &\leq \frac{1}{2} \|u_t\|_{L^2}^2 + C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C + C \|u_{xt}\|_{L^2}^2 \\ &= \frac{1}{2} \|u_t\|_{L^2}^2 + C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C + \|(\dot{u} - u u_x)_x\|_{L^2}^2 \\ &\leq \frac{1}{2} \|u_t\|_{L^2}^2 + C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C + C \|\dot{u}_x\|_{L^2}^2 \\ &\quad + \|u_x\|_{L^\infty}^2 \|u_x\|_{L^2}^2 + \|u\|_{L^\infty}^2 \|u_{xx}\|_{L^2}^2 \\ &\leq \frac{1}{2} \|u_t\|_{L^2}^2 + C \|\dot{u}_x\|_{L^2}^2 + C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C, \end{aligned}$$

where in the last inequality we have used (2.23), (2.21), (2.31), (2.28), and (2.24). Combining this, (2.31), (2.23), (2.36), and (2.37) gives (2.33). The proof of Lemma 2.6 is completed. \square

3 Proof of Theorem 1.1

First, for (ρ_0, u_0, \tilde{f}) satisfying the conditions as in Theorem 1.1, we construct the smooth approximate data as follows:

$$\rho_0^{\delta, \eta} = \frac{\rho_0 * j_\delta + \eta \tilde{\rho}}{1 + \eta}, \quad u_0^{\delta, \eta} = u_0 * j_\delta, \quad \tilde{f}^{\delta, \eta} = \tilde{f} * j_\delta, \quad (3.1)$$

where $\delta \in (0, 1), \eta \in (0, 1)$, and $j_\delta(x)$ is the standard mollifier with width δ . It is easy to check that

$$\begin{cases} 0 < \frac{\eta \tilde{\rho}}{1 + \eta} \leq \rho_0^{\delta, \eta} \leq \frac{\sup_{x \in \mathbb{R}} \rho_0 + \eta \tilde{\rho}}{1 + \eta} < \infty, \\ \lim_{\delta, \eta \rightarrow 0} \left(\|\rho_0^{\delta, \eta} - \rho_0\|_{H^1} + \|u_0^{\delta, \eta} - u_0\|_{L^2} + \|\tilde{f}^{\delta, \eta} - \tilde{f}\|_{H^2} \right) = 0. \end{cases} \quad (3.2)$$

Choosing $\mu_\eta \in C^2[0, \infty)$ satisfying $\lim_{\eta \rightarrow 0} \|\mu_\eta - \mu\|_{C^1[0, M]} = 0$ for any $M > 0$, we consider the Cauchy problem (1.1)-(1.3) with μ replaced by μ_η and the data $(\rho_0^{\delta, \eta}, u_0^{\delta, \eta}, \tilde{f}^{\delta, \eta})$ satisfying (3.1)-(3.2). It follows from Lemma 2.1 that there exists a unique strong solution (ρ, u) of problem (1.1)-(1.3) on $\mathbb{R} \times [0, T_{\delta, \eta}]$. Moreover, the estimates obtained in Lemmas 2.2-2.6 show that the solution (ρ, u) satisfies for any $0 < T \leq T_{\delta, \eta}$,

$$\begin{aligned} & \sup_{0 \leq t \leq T} \left(\|(\rho - \tilde{\rho}, \mu(\rho) - \mu(\tilde{\rho}), P(\rho) - P(\tilde{\rho}))\|_{H^1} + \|\rho_t\|_{L^2} \right. \\ & \left. + \|\rho u\|_{L^2} + \|u\|_{H^1} + \sqrt{t} \|\sqrt{\rho} u_t\|_{L^2} + \sqrt{t} \|u_{xx}\|_{L^2} \right) \\ & + \int_0^T \left(\|u\|_{H^2}^2 + t \|u_{xt}\|_{L^2}^2 \right) dt \leq \bar{C}, \end{aligned} \quad (3.3)$$

where \bar{C} is independent of δ and η . Moreover, similar to [17], we can prove that there exists some \tilde{C} depending on δ and η such that

$$\sup_{0 \leq t \leq T} \left(\|\rho_{xx}\|_{L^2}^2 + \|P_{xx}\|_{L^2}^2 + \|\rho_{xt}\|_{L^2}^2 + \|P_{xt}\|_{L^2}^2 + t \|u_{xt}\|_{L^2}^2 + t \|u_{xxx}\|_{L^2}^2 \right) \leq \tilde{C},$$

which in particular implies that

$$\begin{cases} \rho - \tilde{\rho}, P(\rho) - P(\tilde{\rho}) \in C([0, T]; H^2), & u \in C([0, T]; H^2) \cap L^2(0, T; H^3), \\ u_t \in L^2(0, T; H^1), & t^{1/2} u \in L^\infty(0, T; H^3), \\ t^{1/2} u_t \in L^\infty(0, T; H^1) \cap L^2(0, T; H^2), & t^{1/2} \rho^{1/2} u_{tt} \in L^2(0, T; L^2). \end{cases} \quad (3.4)$$

Then, we will extend the existence time $T_{\delta, \eta}$ to be infinity. Indeed, let T^* be the maximal time of existence for the strong solution. Then, $T^* \geq T_{\delta, \eta}$. If $T^* < \infty$, defining

$$(\rho^*, u^*) \triangleq (\rho, u)(x, T^*) = \lim_{t \rightarrow T^*} (\rho, u)(x, t),$$

we can derive from (3.4) that (ρ^*, P^*, u^*) satisfies the initial condition (1.7) at $t = T^*$. Therefore, one can take (ρ^*, u^*) as the initial data at $t = T^*$ and then use the local existence theory (Lemma 2.1) to extend the strong solution beyond the maximum existence time T^* . This contradicts the assumption on T^* . Thus, $T^* = \infty$.

Finally, we rewrite the global strong solutions on $\mathbb{R} \times [0, \infty)$ obtained above as $(\rho^{\delta, \eta}, u^{\delta, \eta})$. With the estimates (3.3) at hand, letting first $\delta \rightarrow 0$ and then $\eta \rightarrow 0$, we find that the sequence $(\rho^{\delta, \eta}, u^{\delta, \eta})$ converges, up to the extraction of subsequences, to some limit (ρ, u) in the obvious weak sense. Then we deduce from (3.3) that (ρ, u) is a strong solution of (1.1)-(1.3) on $\mathbb{R} \times (0, T]$ (for any $0 < T < \infty$) satisfying (1.8). Moreover, the uniqueness of the strong solution (ρ, u) is guaranteed by the regularities (1.8). For the detailed proof, please see [13]. The proof of Theorem 1.1 is completed. \square

4 Proof of Theorem 1.3

We will divide the proof into three steps.

Step 1. For the case of $f \equiv 0$ in (1.1), we will derive some time-independent lower order estimates of the solutions (ρ, u) obtained in Lemmas 2.5–2.7, that is

$$\sup_{0 \leq t \leq T} \left(\|u\|_{L^2 \cap L^\infty} + \|\sqrt{\mu(\rho)}u_x\|_{L^2}^2 \right) + \int_0^T \|\rho^{1/2}\dot{u}\|_{L^2}^2 dt \leq C, \quad (4.1)$$

$$\sup_{0 \leq t \leq T} \sigma \left(\|u_x\|_{L^\infty}^2 + \|\rho^{1/2}\dot{u}\|_{L^2}^2 \right) + \int_0^T \sigma \|\sqrt{\mu}\dot{u}_x\|_{L^2}^2 dt \leq C. \quad (4.2)$$

First, multiplying (1.1)₂ with $f \equiv 0$ by \dot{u} and integrating by parts, it follows from the same arguments as the proof of (2.22) that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{-\infty}^{+\infty} \mu(\rho)u_x^2 dx + \int_{-\infty}^{+\infty} \rho \dot{u}^2 dx \\ & \leq \frac{d}{dt} \int_{-\infty}^{+\infty} [P(\rho) - P(\bar{\rho})]u_x dx + C\|u_x\|_{L^\infty}\|u_x\|_{L^2}^2 + C\|u_x\|_{L^2}^2 \\ & \leq \frac{d}{dt} \int_{-\infty}^{+\infty} [P(\rho) - P(\bar{\rho})]u_x dx + \frac{1}{2}\|\rho^{1/2}\dot{u}\|_{L^2}^2 + C\|u_x\|_{L^2}^4 + C\|u_x\|_{L^2}^2, \end{aligned} \quad (4.3)$$

where in the last inequality one has used (2.24) and Young's inequality. Since

$$\int_{-\infty}^{+\infty} [P(\rho) - P(\bar{\rho})]u_x dx \leq \frac{\bar{\mu}}{4}\|u_x\|_{L^2}^2 + \|P(\rho) - P(\bar{\rho})\|_{L^2}^2 \leq \frac{\bar{\mu}}{4}\|u_x\|_{L^2}^2 + C \quad (4.4)$$

due to (2.20), the Gronwall's inequality together with (4.4), (4.3), and (2.1) yields (4.1).

Next, operating $(\partial/\partial t + (u \cdot)_x)\dot{u}$ to (1.1) with $f \equiv 0$ and integrating the resulting equation by parts, we deduce from the same calculations as (2.34)–(2.35) that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{-\infty}^{+\infty} \rho |\dot{u}|^2 dx + \int_{-\infty}^{+\infty} \mu(\rho) |\dot{u}_x|^2 dx \\ & = \gamma \int_{-\infty}^{+\infty} P(\rho)u_x \dot{u}_x dx + \int_{-\infty}^{+\infty} (\mu(\rho) + \mu'(\rho)\rho)u_x^2 \dot{u}_x dx \\ & \leq \frac{\bar{\mu}}{2}\|\dot{u}_x\|_{L^2}^2 + C\|u_x\|_{L^2}^2 \|\rho^{1/2}\dot{u}\|_{L^2}^2 + C\|u_x\|_{L^2}^4 + C\|u_x\|_{L^2}^2. \end{aligned} \quad (4.5)$$

Multiplying (4.5) by σ , one gets after using Gronwall's inequality, (1.11), (2.1), and (4.1) that

$$\sup_{0 \leq t \leq T} \sigma \|\rho^{1/2}\dot{u}\|_{L^2}^2 + \int_0^T \sigma \|\dot{u}_x\|_{L^2}^2 dt \leq C, \quad (4.6)$$

which together with (2.23), (2.24), and (4.1) implies that

$$\sup_{0 \leq t \leq T} (\|u\|_{L^2 \cap L^\infty} + \sigma\|u_x\|_{L^\infty}) \leq C. \quad (4.7)$$

The combination of (4.7) with (4.6) gives (4.2).

Step 2. We will show that for any $p \geq 2$,

$$\lim_{t \rightarrow 0} \|u_x\|_{L^p} = 0. \quad (4.8)$$

First, we claim that

$$\lim_{t \rightarrow \infty} \int_{t-1}^t \left(\|u_x\|_{L^2}^2 + \left| \frac{d}{d\tau} \|u_x\|_{L^2}^2 \right| \right) d\tau = 0. \quad (4.9)$$

For any $t > 1$ and $s \in (t-1, t)$, it holds

$$\|u_x\|_{L^2}^2(t) - \|u_x\|_{L^2}^2(s) = \int_t^s \frac{d}{d\tau} \|u_x\|_{L^2}^2 d\tau \leq \int_{t-1}^t \left| \frac{d}{d\tau} \|u_x\|_{L^2}^2 \right| d\tau,$$

that is

$$\|u_x\|_{L^2}^2(t) \leq \int_{t-1}^t \left(\|u_x\|_{L^2}^2 + \left| \frac{d}{d\tau} \|u_x\|_{L^2}^2 \right| \right) d\tau,$$

which together with (4.9) yields

$$\lim_{t \rightarrow \infty} \|u_x\|_{L^2}^2(t) = 0. \quad (4.10)$$

It is easy to deduce from (4.7) that for any $p > 2$,

$$\|u_x\|_{L^p}^p \leq \|u_x\|_{L^2}^2 \|u_x\|_{L^\infty}^{p-2} \leq C \|u_x\|_{L^2}^2, \quad \forall t > 1. \quad (4.11)$$

The combination of (4.10) with (4.11) yields (4.8).

Now, it remains to prove (4.9). Indeed, it follows from integration by parts and (4.7) that

$$\begin{aligned} \left| \frac{d}{dt} \|u_x\|_{L^2}^2 \right| &= \left| 2 \int_{-\infty}^{+\infty} u_x u_{xt} dx \right| = \left| 2 \int_{-\infty}^{+\infty} u_x (\dot{u}_x - (uu_x)_x) dx \right| \\ &= \left| 2 \int_{-\infty}^{+\infty} u_x \dot{u}_x dx - \int_{-\infty}^{+\infty} u_x^3 dx \right| \\ &\leq C \|\dot{u}_x\|_{L^2}^2 + C(1 + \|u_x\|_{L^\infty}) \|u_x\|_{L^2}^2 \\ &\leq C \|\dot{u}_x\|_{L^2}^2 + C \|u_x\|_{L^2}^2. \end{aligned}$$

Combining this, (2.1), and (4.2) yields

$$\int_1^\infty \left(\|u_x\|_{L^2}^2 + \left| \frac{d}{dt} \|u_x\|_{L^2}^2 \right| \right) dt \leq C,$$

which yields (4.9). The proof of step 2 is completed.

Step 3. Using the methods due to [9, 12], we are now in a position to prove that for any $p > 2$,

$$\lim_{t \rightarrow \infty} \|\rho - \tilde{\rho}\|_{L^p} = 0. \quad (4.12)$$

First, we claim that

$$\int_0^\infty \|\rho - \tilde{\rho}\|_{L^6}^6 dt \leq C. \quad (4.13)$$

It follows from integration by parts, (1.1)₁, and (2.8) that

$$\begin{aligned} \frac{d}{dt} \|\rho - \tilde{\rho}\|_{L^6}^6 &= 6 \int_{-\infty}^{+\infty} (\rho - \tilde{\rho})^5 (\rho - \tilde{\rho})_t dx \\ &= -6 \int_{-\infty}^{+\infty} (\rho - \tilde{\rho})^5 (\rho - \tilde{\rho})_x u dx - 6 \int_{-\infty}^{+\infty} (\rho - \tilde{\rho})^5 \rho u_x dx \\ &\leq C \|\rho - \tilde{\rho}\|_{L^6}^3 \|u_x\|_{L^2} + C \|\rho - \tilde{\rho}\|_{L^6}^3 \|u_x\|_{L^2} \\ &\leq C \|\rho - \tilde{\rho}\|_{L^6}^6 + C \|u_x\|_{L^2}^2, \end{aligned}$$

which together with (4.13) and (2.1) gives

$$\int_0^\infty \left| \frac{d}{dt} \|\rho - \tilde{\rho}\|_{L^6}^6 \right| dt \leq C.$$

Combining this with (4.13) implies that

$$\lim_{t \rightarrow \infty} \|\rho - \tilde{\rho}\|_{L^6} = 0,$$

which along with (2.20) and (2.8) leads to the desired (4.12).

Next, we need to prove (4.13). For

$$Q \triangleq \mu(\rho)u_x - (P(\rho) - P(\tilde{\rho})), \quad (4.14)$$

it follows from (1.1)₂ with $f \equiv 0$ that

$$Q_x = \rho \dot{u},$$

which gives

$$\|Q_x\|_{L^2} = \|\rho \dot{u}\|_{L^2} \leq C \|\rho^{1/2} \dot{u}\|_{L^2}. \quad (4.15)$$

It follows from (1.1)₁ that

$$(\rho - \tilde{\rho})_t + (\rho - \tilde{\rho})_x u + (\rho - \tilde{\rho})u_x + \tilde{\rho}u_x = 0. \quad (4.16)$$

Multiplying (4.16) by $6(\rho - \tilde{\rho})^5$ and integrating by parts implies

$$\begin{aligned} & \frac{d}{dt} \|\rho - \tilde{\rho}\|_{L^6}^6 + 6 \int_{-\infty}^{+\infty} \tilde{\rho}(\mu(\rho))^{-1} (\rho - \tilde{\rho})^5 (P(\rho) - P(\tilde{\rho})) dx \\ &= -6 \int_{-\infty}^{+\infty} \tilde{\rho}(\mu(\rho))^{-1} (\rho - \tilde{\rho})^5 Q dx - 5 \int_{-\infty}^{+\infty} (\rho - \tilde{\rho})^6 u_x dx \\ &\leq C \|\rho - \tilde{\rho}\|_{L^6}^5 \|Q\|_{L^6} + C \|\rho - \tilde{\rho}\|_{L^\infty}^3 \|\rho - \tilde{\rho}\|_{L^6}^3 \|u_x\|_{L^2} \\ &\leq \varepsilon \|\rho - \tilde{\rho}\|_{L^6}^6 + C(\varepsilon) \|Q\|_{L^2}^4 \|Q_x\|_{L^2}^2 + C(\varepsilon) \|u_x\|_{L^2}^2 \\ &\leq \varepsilon \|\rho - \tilde{\rho}\|_{L^6}^6 + C(\varepsilon) \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C(\varepsilon) \|u_x\|_{L^2}^2, \end{aligned} \quad (4.17)$$

where one has used (4.14), (1.6), (2.8), (4.15), (4.2), and (2.20). Furthermore, the direct calculations combined with (1.6) and (2.8) show that for some $0 < \alpha < 1$,

$$\begin{aligned} 6\tilde{\rho}(\mu(\rho))^{-1} (\rho - \tilde{\rho})^5 (P(\rho) - P(\tilde{\rho})) &= 6\tilde{\rho}(\mu(\rho))^{-1} (\rho - \tilde{\rho})^6 P'(\alpha\rho + (1 - \alpha)\tilde{\rho}) \\ &\geq C_0 (\rho - \tilde{\rho})^6, \end{aligned} \quad (4.18)$$

where the positive constant C_0 depending only on $\gamma, \bar{\mu}, \bar{\rho}$ and $\tilde{\rho}$. Substituting (4.18) into (4.17) and choosing ε suitably small yield

$$\frac{d}{dt} \|\rho - \tilde{\rho}\|_{L^6}^6 + C_0 \|\rho - \tilde{\rho}\|_{L^6}^6 \leq C \|\rho^{1/2} \dot{u}\|_{L^2}^2 + C \|u_x\|_{L^2}^2. \quad (4.19)$$

Thus, one can derive the desired (4.13) from (4.19), (2.1), (2.20), and (4.1).

Finally, noticing that

$$\|\rho - \tilde{\rho}\|_{C(\bar{\mathbb{R}})} \leq C \|\rho - \tilde{\rho}\|_{L^6}^{3/4} \|\rho_x\|_{L^2}^{1/4},$$

the proof of (1.14) is similar as that of [14, Theorem 1.2](see also [9]). The proof of Theorem 1.3 is finished. \square

References

- [1] A. A. Amosov, A. A. Zlotnik, Global generalized solutions of the equations of the one-dimensional motion of a viscous heat-conducting gas, *Soviet Math. Dokl.*, **38**(1989), 1-5.
- [2] H. Beirão da Veiga, Long time behavior for one-dimensional motion of a general barotropic viscous fluid, *Arch. Ration. Mech. Anal.*, **108**(1989), 141-160.
- [3] Y. Cho, H. Kim, On classical solutions of the compressible Navier-Stokes equations with nonnegative initial densities. *Manuscript Math.*, **120** (2006), 91-129.
- [4] S. J. Ding, H. Y. Wen, C. J. Zhu. Global classical large solutions of 1D compressible Navier-Stokes equations with density-dependent viscosity and vacuum, *J. Differ. Eqs.*, **221**(2011), 1696-1725.
- [5] E. Feireisl, Dynamics of viscous compressible fluids. Oxford University Press, 2004.
- [6] D. Hoff, Global existence for 1D, compressible, isentropic Navier-Stokes equations with large initial data. *Trans. Amer. Math. Soc.* **303** (1987), no. 1, 169–181.
- [7] F. M. Huang, J. Li, Z. P. Xin, Convergence to equilibria and blowup behavior of global strong solutions to the Stokes approximation equations for two-dimensional compressible flows with large data. *J. Math. Pures Appl.*, 86(6)(2006), 471-491.
- [8] X. D. Huang, J. Li, Serrin-type blowup criterion for viscous, compressible, and heat conducting Navier-Stokes and Magnetohydrodynamic flows. *Comm. Math. Phys.*, **324** (2013), 147-171.
- [9] X. D. Huang, J. Li, Z. P. Xin, Global well-posedness of classical solutions with large oscillations and vacuum to the three-dimensional isentropic compressible Navier-Stokes equations, *Comm. Pure Appl. Math.*, **65**(2012), 549-585.
- [10] Ya. I. Kanel, A model system of equations for the one-dimensional motion of a gas. (Russian) *Differencial'nye Uravnenija* 4 (1968), 721-734
- [11] A. V. Kazhikhov, Cauchy problem for viscous gas equations. *Siberian Math. J.* **23** (1982), 44-49.
- [12] J. Li, J. W. Zhang, J. N. Zhao, On the global motion of viscous compressible barotropic flows subject to large external potential forces and vacuum, *SIAM J. Math. Anal.*, **47**(2)(2015), 1121-1153.
- [13] J. Li, Z. L. Liang, On classical solutions to the Cauchy problem of the two-dimensional barotropic compressible Navier-Stokes equations with vacuum, *J. Math. Pures Appl.*, (9)**102**(2014), 640-671.
- [14] J. Li, Z. P. Xin, Some uniform estimates and blowup behavior of global strong solution to the Stokes approximation equations for two-dimensional compressible flows, *J. Differential Equations*, 221(2006), 275-308.
- [15] J. Li, Z. P. Xin, Global well-posedness and large time asymptotic behavior of classical solution to the compressible Navier-Stokes equations with vacuum, <http://arxiv.org/abs/1310.1673v1>.

- [16] P. L. Lions, Mathematical topics in fluid mechanics. Vol. **2**. Compressible models. Oxford University Press, New York, 1998.
- [17] B. Q. Lü, Y. X. Wang, Y. H. Wu, On global classical solutions to 1d compressible Navier-Stokes equations with density-dependent viscosity and vacuum, arXiv:1808.03042
- [18] A. Matsumura, T. Nishida, The initial value problem for the equations of motion of viscous and heat-conductive gases, *J. Math. Kyoto Univ.*, **20**(1980), no. 1, 67-104.
- [19] D. Serre, Solutions faibles globales des equations de Navier-Stokes pour un fluide compressible, *C. R. Acad. Sci. Paris Sr. I Math.*, **303(13)**(1986), 639-642.
- [20] D. Serre, On the one-dimensional equation of a viscous, compressible, heat-conducting fluid. *C. R. Acad. Sci. Paris Sr. I Math.*, **303(14)**(1986), 703-706.
- [21] I. Straškraba, A. Zlotnik, On a decay rate for 1D-viscous compressible barotropic fluid equations, *J. Evolution Equations*, **2**(2002), 69-96.
- [22] Z. P. Xin, Blowup of smooth solutions to the compressible Navier-Stokes equation with compact density, *Comm. Pure Appl. Math.*, **51** (1998), 229-240.
- [23] Y. L. Ye, Global classical solution to 1D compressible Navier-Stokes equations with no vacuum at infinity, *Math. Meth. Appl. Sci.*, **39** (2016), 776-795.