THE BURGERS' EQUATION WITH STOCHASTIC TRANSPORT: SHOCK FORMATION, LOCAL AND GLOBAL EXISTENCE OF SMOOTH SOLUTIONS

DIEGO ALONSO-ORÁN, AYTHAMI BETHENCOURT DE LEÓN AND SO TAKAO

ABSTRACT. In this work, we examine the solution properties of the Burgers' equation with stochastic transport. First, we prove results on the formation of shocks in the stochastic equation and then obtain a stochastic Rankine-Hugoniot condition that the shocks satisfy. Next, we establish the local existence and uniqueness of smooth solutions in the inviscid case and construct a blow-up criterion. Finally, in the viscous case, we prove global existence and uniqueness of smooth solutions.

1. INTRODUCTION

We prove the well-posedness of a stochastic Burgers' equation of the form

$$du(t,x) + u(t,x)\partial_x u(t,x) dt + \sum_{k=1}^{\infty} \xi_k(x)\partial_x u(t,x) \circ dW_t^k = \nu \partial_{xx} u(t,x) dt, \qquad (1.1)$$

where $x \in \mathbb{T}$ or \mathbb{R} , $\nu \geq 0$ is constant, $\{W_t^k\}_{k \in \mathbb{N}}$ is a countable set of independent Brownian motions, $\{\xi_k(\cdot)\}_{k \in \mathbb{N}}$ is a countable set of prescribed functions depending only on the spatial variable, and \circ means that the stochastic integral is interpreted in the Stratonovich sense. If the set $\{\xi_k(\cdot)\}_{k \in \mathbb{N}}$ forms a basis of some separable Hilbert space \mathcal{H} (for example $L^2(\mathbb{T})$), then the process $dW := \sum_{k=1}^{\infty} \xi_k(x) \circ dW_t^k$ is a *cylindrical Wiener process* on \mathcal{H} , generalising the notion of a standard Wiener process to infinite dimensions.

The multiplicative noise in (1.1) makes the transport velocity stochastic, which allows the Burgers' equation to retain the form of a transport equation $\partial_t u + \tilde{u} \partial_x u = 0$, where $\tilde{u}(t,x) := u(t,x) + W$ is a stochastic vector field with noise W that is smooth in space and rough in time. Compared with the well-studied Burgers' equation with additive noise, where the noise appears as an external random forcing, this type of noise arises by taking the diffusive limit of the Lagrangian flow map regarded as a composition of a slow mean flow and a rapidly fluctuating one [CGH17]. In several recent works, this type of noise, which we call *stochastic transport*, has been used to stochastically parametrise unresolved scales in fluid models while retaining the essential physics of the system [Hol15, CCH⁺18, CCH⁺19]. On the other hand, it has also been shown to have a regularising effect on certain PDEs that are ill-posed [FGP10, FGP11, FF13, GM18]. Therefore, it is of interest to investigate how the stochastic transport in (1.1) affects the Burgers' equation, which in the inviscid case $\nu = 0$ is a prototypical model for *shock* formation. In particular, we ask whether this noise can prevent the system from developing shocks or, on the contrary, produce new shocks. We also ask whether this system is well-posed or not. In this paper, we will show that:

- (1) For $\nu = 0$, equation (1.1) has a unique solution of class H^s for s > 2 until some stopping time $\tau > 0$.
- (2) However, shock formation cannot be avoided a.s. in the case $\xi(x) = \alpha x + \beta$ and for a broader class of $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$, we can prove that it occurs in expectation.
- (3) For $\nu > 0$, we have global existence and uniqueness in H^s for s > 2.

On top of this, we prove a continuation criterion for the inviscid equation ($\nu = 0$), which generalises the result for the deterministic case. The above results are not immediately evident for reasons we will discuss below. Although we cannot prove this here, we believe that shocks in Burgers' equation are too robust and ubiquitous to be prevented by noise, regardless of what $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$ is chosen. Our results provide rigorous evidence to support this claim.

The question of whether noise can regularise PDEs is not new. In finite dimensions, it is well-known that additive noise can restore the well-posedness of ODEs whose vector fields are merely bounded and measurable (see [Ver81]). For PDEs, a general result is not known; however, there has been a significant effort in recent years to generalise this celebrated result to PDEs. In a remarkable paper, Flandoli, Gubinelli, and Priola [FGP10] demonstrated that the linear transport equation $\partial_t u + b(t, x) \cdot \nabla u = 0$, which is ill-posed if b is sufficiently irregular, can recover existence and uniqueness of L^{∞} solutions that is strong in the probabilistic sense, by the addition of a "simple" transport noise,

$$du + b(t, x) \cdot \nabla u \, dt + \nabla u \circ dW_t = 0, \tag{1.2}$$

where the drift b is bounded, measurable, Hölder continuous, and satisfies an integrability condition on the divergence $\nabla \cdot b \in L^p([0,T] \times \mathbb{R}^d)$. In a subsequent paper [FF13], the same noise was shown to retain some regularity of the initial condition, thus restoring wellposedness of strong solutions, and a selection principle based on taking the zero-noise limit as opposed to the inviscid limit was considered in [AF09].

However, for nonlinear transport equations such as Burgers', the same type of noise $du + u \partial_x u dt + \partial_x u \circ dW_t = 0$ does not help, since a simple change of variables $v(t, x) := u(t, x - W_t)$ will lead us back to the original equation $\partial_t v + v \partial_x v = 0$. Hence, if noise were to prevent shock formation, a more general class could be required, such as the cylindrical transport noise $\sum_{k=1}^{\infty} \xi_k(x) \partial_x u \circ dW_t^k$ that we consider in this paper. In [FGP11] and [DFV14], it was shown that collapse in Lagrangian point particle solutions of certain nonlinear PDEs (point vortices in 2D Euler and point charges in the Vlasov-Poisson system), can be prevented by this cylindrical transport noise with $\xi_k(x)$ satisfying a certain hypoellipticity condition, thus providing hope for regularisation of nonlinear transport equation by noise. More recently, Gess and Maurelli [GM18] showed that adding a simple stochastic transport term into a nonlinear transport equation

$$du + b(x, u(t, x))\nabla u \, dt + \nabla u \circ dW_t = 0, \tag{1.3}$$

which in the deterministic case admits non-unique entropy solutions for sufficiently irregular b, can restore uniqueness of entropy solutions, providing a first example of a nonlinear transport equation that becomes well-posed when adding a suitable noise.

We should now stress the difference between the present work and previous works. First, we acknowledge that in Flandoli [Fla11], Chapter 5.1.4, it is argued that shock formation does not occur even with the most general cylindrical transport noise, by writing the

characteristic equation as an $It\hat{o}$ SDE

$$X_t = X_0 + u(0, X_0)t + \sum_{k=1}^{\infty} \int_0^t \xi_k(X_s) \, \mathrm{d}W_s^k, \tag{1.4}$$

which is a martingale perturbation of straight lines that will cross without noise. Thus, using the property that a martingale M_t grows slower than t almost surely as $t \to \infty$, it is shown that the characteristics cross almost surely. However, the characteristic equation for the system (1.1) is in fact a *Stratonovich* SDE,

$$X_t = X_0 + u(0, X_0)t + \sum_{k=1}^{\infty} \int_0^t \xi_k(X_s) \circ \mathrm{d}W_s^k,$$
(1.5)

and therefore Flandoli's argument can be applied to the martingale term, but not to the additional drift term, which may disrupt shock formation. The techniques we use here apply to Stratonovich equations; however, due to the difficulty caused by the additional drift term, we were only able to prove that the characteristics cross almost surely in the very particular case $\xi(x) = \alpha x + \beta$, leaving the general case open for future investigation. By using a different strategy, where instead we look at how the slope $\partial_x u$ evolves along a characteristic (1.5), we manage to show that for a wider class of $\{\xi_k(\cdot)\}_{k=1}^{\infty}$ such that the infinite sum $\sum_{k \in \mathbb{N}} ((\partial_x \xi_k)^2 - \xi_k \partial_{xx} \xi_k)$ is pointwise bounded, we have that

- if $\partial_x u(0, X_0) > 0$, then $\partial_x u(t, X_t) < \infty$ almost surely for all t > 0 and
- if $\partial_x u(0, X_0)$ is sufficiently negative, then there exists $0 < t_* < \infty$ such that $\lim_{t \to t_*} \mathbb{E}[\partial_x u(t, X_t)] = -\infty$.

In summary, shock formation occurs in expectation if the initial profile has a sufficiently negative slope and no new shocks can form from a positive slope.

We finally address the question of well-posedness. We will prove that by choosing a sufficiently regular initial condition, equation (1.1) admits a unique local solution that is smooth enough, such that the arguments employed in the previous section on shock formation are valid (in fact, we show this for a noise of the type $Qu \circ dW_t$, where Qu = $a(x)\partial_x u + b(x)u$, which generalises the one considered in (1.1)). For Burgers' equation with additive space-time white noise, however, there have been many previous works showing well-posedness [BCJL94, DPDT94, DPG07, CO13]. The techniques used in these works are primarily based on reformulating the equations by a change of variable or by studying its linear part. The main difference in our work is that the multiplicative noise we consider depends on the solution and its gradient. Therefore, the effect of the noise hinges on its spatial gradient and the solution, giving rise to several complications. For instance, when deriving a priori estimates, certain high order terms appear, which need to be treated carefully. Recently, the same type of multiplicative noise has been treated for the Euler equation [CFH19, FL18] and the Boussinesq system [AOB20], whose techniques we follow closely in our proof. We note that the well-posedness analysis of a more general stochastic conservation law, which includes the inviscid stochastic Burgers' equation as a special case, has also been considered, for instance in [FG16, GS17, FGH20]. However, these works deal with the well-posedness analysis of weak kinetic and entropy solutions, in contrast to classical solutions, which we consider here. There is also the recent work [HNS19] showing the local well-posedness of weak solutions in the viscous Burgers' equation ($\nu > 0$) driven by rough paths in the transport velocity. An important contribution of this paper is showing the global well-posedness of strong solutions in the viscous case by proving that the maximum principle is retained under perturbation by stochastic transport of type $\xi(x)\partial_x u(t,x) \circ dW_t$.

1.1. Main results. Let us state here the main results of the article:

Theorem 1.1 (Shock formation in the stochastic Burgers' equation). In the following, we use the notation $\psi(x) := \frac{1}{2} \sum_{k=1}^{\infty} ((\partial_x \xi_k(x))^2 - \xi_k(x) \partial_{xx} \xi_k(x))$. The main results regarding shock formation in (1.1) are as follows:

- (1) Let $\xi_1(x) = \alpha x + \beta$, $x \in \mathbb{R}$ and $\xi_k \equiv 0$ for k = 2, 3, ... and assume that u(0, x) has a negative slope. Then, there exists two characteristics satisfying (1.5) with different initial conditions that cross in finite time almost surely.
- (2) Let X_t be a characteristic solving (1.5) with $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$ satisfying the conditions in Assumption A1 below and let $\partial_x u(0, X_0) \ge 0$. Then, if $\psi(x) < \infty$ for all $x \in \mathbb{T}$ or \mathbb{R} , we have that $\partial_t u(t, X_t) < \infty$ almost surely for all t > 0.
- (3) Again, let X_t be a characteristic solving (1.5) with $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$ satisfying the conditions in Assumption A1 and let $\partial_x u(0, X_0) < 0$. Also assume that $\partial_x u(0, X_0) < \psi(x)$ for all $x \in \mathbb{T}$ or \mathbb{R} . Then there exists $0 < t_* < \infty$ such that $\lim_{t \to t_*} \mathbb{E}[\partial_x u(t, X_t)] = -\infty$.

Theorem 1.2 (Stochastic Rankine-Hugoniot condition). The curve of discontinuity $(t, s(t)) \in [0, \infty) \times \mathbb{T}$ (or \mathbb{R}) of the stochastic Burgers' equation (1.1) satisfies the following:

$$ds_t = \frac{1}{2} \left[\left(u_-(t, s(t)) + u_+(t, s(t)) \right] dt + \sum_{k=1}^{\infty} \xi_k(s(t)) \circ dW_t^k,$$
(1.6)

where $u_{\pm}(t, s(t)) := \lim_{x \to s(t)^{\pm}} u(t, x)$ are the left and right limits of u.

Theorem 1.3 (Well-posedness in the inviscid case). Let $u_0 \in H^s(\mathbb{T})$ for some s > 2 fixed. Then there exists a pathwise unique H^s -maximal solution (τ_{max}, u) of the 1D Burgers' equation (1.1) in the sense of Definition 2.2 with initial datum u_0 . Moreover, either $\tau_{max} = \infty$ or $\limsup_{t \to \tau_{max}} |u(t)|_{H^s} = \infty$, a.s.

Theorem 1.4 (Global well-posedness in the viscous case). Let $u_0 \in H^s(\mathbb{T})$ for some s > 2 fixed. Then there exists a pathwise unique maximal global H^s -solution u of the viscous stochastic Burgers' equation (1.1) with $\nu > 0$.

Remark 1.5. Theorems 1.3 and 1.4 can be extended in a straightforward manner to the full line \mathbb{R} and to higher dimensions.

Remark 1.6. We prove Theorem 1.3 for a more general noise $Qu \circ dW_t$, where Q is a first order linear differential operator, which includes the transport noise as a special case. For the sake of clarity, our proof deals only with one noise term $Qu \circ dW_t$, however, we can readily extend this to cylindrical noise with countable set of first order linear differential operators

$$\sum_{k=1}^{\infty} \mathcal{Q}_k(u) \circ \mathrm{d} W_t^k,$$

by imposing certain smoothness and boundedness conditions for the sum of the coefficients. We also prove Theorem 1.4 for one noise term. 1.2. Structure of the paper. This manuscript is organised as follows. In Section 2 we review some classical mathematical deterministic and stochastic background. We also fix the notations we will employ and state some definitions. Section 3 contains the main results regarding shock formation in the stochastic Burgers' equation. Using a characteristic argument, we show that noise cannot prevent shocks from occurring for certain classes of $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$. Moreover, we prove that these shocks satisfy a Rankine-Hugoniot type condition in the weak formulation of the problem. In Section 4, we show local well-posedness of the stochastic Burgers' equation in Sobolev spaces and a blow-up criterion. We also establish global existence of smooth solutions of a viscous version of the stochastic Burgers' equation, which is achieved by proving a stochastic analogue of the maximum principle. In Section 5, we provide conclusions, propose possible future research lines, and comment on several open problems that are left to study.

2. Preliminaries and notation

Let us begin by reviewing some standard functional spaces and mathematical background that will be used throughout this article. Sobolev spaces are given by

$$W^{s,p} := \{ f \in L^p(\mathbb{T}, \mathbb{R}) : (I - \partial_{xx})^{s/2} f \in L^p(\mathbb{T}, \mathbb{R}) \},\$$

for any $s \ge 0$ and $p \in [1, \infty)$, equipped with the norm $||f||_{W^{s,p}} = ||(I - \partial_{xx})^{s/2}f||_{L^p}$. We will also use the notation $\Lambda^s = (-\partial_{xx})^{s/2}$. Recall that L^2 based spaces are Hilbert spaces and may alternatively be denoted by $H^s = W^{s,2}$. For s > 0, we also define $H^{-s} := (H^s)^*$, i.e. the dual space of H^s . Let us gather here some well-known Sobolev embedding inequalities:

$$\|f\|_{L^4} \lesssim \|f\|_{L^2}^{1/2} \|\partial_x f\|_{L^2}^{1/2}, \qquad (2.1)$$

$$\|\partial_x f\|_{L^4} \lesssim \|f\|_{\infty}^{1/2} \|\partial_{xx} f\|_{L^2}^{1/2}, \qquad (2.2)$$

$$\|f\|_{\infty} \lesssim \|f\|_{H^{1/2+\epsilon}}, \text{ for every } \epsilon > 0.$$

$$(2.3)$$

Let us also recall the well-known commutator estimate of Kato and Ponce:

Lemma 2.1 ([KP88]). If $s \ge 0$ and 1 , then

$$|\Lambda^{s}(fg) - f\Lambda^{s}(g)||_{L^{p}} \le C_{p} \left(\|\partial_{x}f\|_{\infty} \|\Lambda^{s-1}g\|_{L^{p}} + \|\Lambda^{s}f\|_{L^{p}} \|g\|_{\infty} \right).$$
(2.4)

We will also use the following result as main tool for proving the existence results and blow-up criterion:

Theorem 2.2 ([AOB20]). Let Q be a linear differential operator of first order

$$\mathcal{Q} = a(x)\partial_x + b(x)$$

where the coefficients are smooth and bounded. Then for $f \in H^2(\mathbb{T},\mathbb{R})$ we have

$$\langle \mathcal{Q}^2 f, f \rangle_{L^2} + \langle \mathcal{Q} f, \mathcal{Q} f \rangle_{L^2} \lesssim ||f||_{L^2}^2.$$
 (2.5)

Moreover, if $f \in H^{2+s}(\mathbb{T},\mathbb{R})$, and \mathcal{P} is a pseudodifferential operator of order s, then

$$\langle \mathcal{P}\mathcal{Q}^2 f, \mathcal{P}f \rangle_{L^2} + \langle \mathcal{P}\mathcal{Q}f, \mathcal{P}\mathcal{Q}f \rangle_{L^2} \lesssim ||f||_{H^s}^2,$$
 (2.6)

for every $s \in [1, \infty)$.

Remark 2.3. Theorem 2.2 is fundamental for closing the energy estimates when showing well-posedness of the stochastic Burgers' equation. It permits reducing the order of a sum of terms which in principle seems hopelessly singular.

Remark 2.4. As pointed out in the previous observation, in order to extend Theorem 1.3 to cylindrical noise of the form $\sum_{k=1}^{\infty} \xi_k(x) \partial_x u(t,x) \circ dW_t^k$ (or in general $\sum_{k=1}^{\infty} Q_k(u) \circ dW_t^k$), it is fundamental to show the cancellation property provided by Theorem 2.2 for such noises. This can be done under some mild Sobolev regularity assumption on the coefficients ξ_k (respectively a_k, b_k). In particular, one has to precisely compute the constants C_k hidden on the right hand-side of (2.5) and (2.6), whose sum a priori does not have to converge. We refer the reader to Lemma A.5 in [AOHR21] for a detailed calculation to deal with this extension.

Next, we briefly recall some aspects of the theory of stochastic analysis. Fix a stochastic basis $S = (\Xi, \mathcal{F}, \{\mathcal{F}_t\}_{t\geq 0}, \mathbb{P}, \{W^k\}_{k\in\mathbb{N}})$, that is, a filtered probability space together with a sequence $\{W^k\}_{k\in\mathbb{N}}$ of scalar independent Brownian motions relative to the filtration $\{\mathcal{F}_t\}_{t\geq 0}$ satisfying the usual conditions.

Given a stochastic process $X \in L^2(\Xi; L^2([0, \infty); L^2(\mathbb{T}, \mathbb{R})))$, the Burkholder-Davis-Gundy inequality is given by

$$\mathbb{E}\left[\sup_{s\in[0,T]}\left|\int_0^t X_s \,\mathrm{d}W_s\right|^p\right] \le C_p \mathbb{E}\left[\int_0^T |X_s|^2 \,\mathrm{d}t\right]^{p/2},\tag{2.7}$$

for any $p \ge 1$ and C_p an absolute constant depending on p.

We also state the celebrated Itô-Wentzell formula, which we use throughout this work.

Theorem 2.5 ([Kun81], Theorem 1.2). For $0 \le t < \tau$, let $u(t, \cdot)$ be C^3 almost surely, and $u(\cdot, x)$ be a continuous semimartingale satisfying the SPDE

$$u(t,x) = u(0,x) + \sum_{j=0}^{\infty} \int_0^t \sigma_j(s,x) \circ \mathrm{d}N_s^j,$$
(2.8)

where $\{N_t^j\}_{j=0}^{\infty}$ is a family of continuous semimartingales and $\{\sigma_j(t,x)\}_{j=0}^{\infty}$ is also a family of continuous semimartingales that are C^2 in space for $0 \leq t < \tau$. Also, let X_t be a continuous semimartingale. Then, we have the following

$$u(t, X_t) = u(0, X_0) + \sum_{j=0}^{\infty} \int_0^t \sigma_j(s, X_s) \circ \mathrm{d}N_s^j + \int_0^t \partial_x u(s, X_s) \circ \mathrm{d}X_s.$$
(2.9)

Let us also introduce three different notions of solutions:

Definition 2.1 (Local solution). An H^s -local solution (τ, u) of (1.1), $s \ge 2$, is a stochastic process $u : \Xi \times [0, \tau] \times \mathbb{T} \to \mathbb{R}$, where $\tau : \Xi \to [0, \infty)$ is a stopping time such that $u_{t \wedge \tau}$ is adapted to $(\mathcal{F}_t)_t$ and we have:

• a.s. u has paths of class $C([0, \tau]; H^s(\mathbb{T}))$.

• It holds

$$u_{\tau'} - u_0 + \int_0^{\tau'} u_s \partial_x u_s \, \mathrm{d}s + \sum_{i=1}^N \int_0^{\tau'} \xi_i \partial_x u_s \, \mathrm{d}W_s^i = \frac{1}{2} \sum_{i=1}^N \int_0^{\tau'} (\xi_i \partial_x)^2 u_s \, \mathrm{d}s,$$

as an identity in $L^2(\mathbb{T})$, a.s., for bounded stopping times $\tau' \leq \tau$.

When the stopping time is clear from the context, we simply write that u is a solution.

Definition 2.2 (Maximal solution). A maximal solution (τ_{max}, u) of (1.1) is a stochastic process $u : \Xi \times [0, \tau_{max}) \times \mathbb{T} \to \mathbb{R}$, for a stopping time $\tau_{max} : \Xi \to [0, \infty]^{-1}$, satisfying the following conditions:

- $\mathbb{P}(\tau_{max} > 0) = 1$, where $\tau_{max} = \lim_{n \to \infty} \tau_n$ for an increasing sequence of stopping times $\{\tau_n\}_{n=1}^{\infty}$.
- (τ_n, u) is a local solution in the sense of Definition 2.1, for all $n \in \mathbb{N}$.
- If (τ', u') is another pair satisfying the above two conditions and u' = u on $[0, \tau' \wedge \tau_{max})$, then $\tau' \leq \tau_{max}$, a.s.

A maximal solution is said to be global if $\tau_{max} = \infty$, a.s.

Definition 2.3 (Weak solution). A (spatially) weak solution (τ, u) of (1.1) is a stochastic process $u : \Xi \times [0, \tau] \times \mathbb{T} \to \mathbb{R}$, where $\tau : \Xi \to [0, \infty)$ is a stopping time, such that for any test function $\phi \in C^{\infty}(\mathbb{T})$, $\langle u_{t\wedge\tau}, \phi \rangle_{L^2}$ is adapted to $(\mathcal{F}_t)_t$ and we have:

- a.s. u has paths of class $C([0, \tau]; L^2(\mathbb{T}))$.
- It holds

$$\langle u_{\tau'}, \phi \rangle_{L^2} = \langle u_0, \phi \rangle_{L^2} + \frac{1}{2} \int_0^{\tau'} \langle (u_s)^2, \partial_x \phi \rangle_{L^2} \,\mathrm{d}s + \sum_{i=1}^N \int_0^{\tau'} \langle u_s, \partial_x(\xi_i \phi) \rangle_{L^2} \,\mathrm{d}W_s^i + \frac{1}{2} \sum_{i=1}^N \int_0^{\tau'} \langle u_s, \partial_x(\xi_i \partial_x(\xi_i \phi)) \rangle_{L^2} \,\mathrm{d}s,$$
(2.10)

a.s., for bounded stopping times $\tau' \leq \tau$.

Notations: Let us stress some notations that we will use throughout this work. We will denote the Sobolev L^2 based spaces by $H^s(\text{domain}, \text{target space})$. However, we will sometimes omit the domain and target space and just write H^s , when these are clear from the context. $a \leq b$ means there exists C such that $a \leq Cb$, where C is a positive universal constant that may depend on fixed parameters and constant quantities. Note also that this constant might differ from line to line. It is also important to remind that the condition "almost surely" is not always indicated, since in some cases it is obvious from the context.

3. SHOCKS IN BURGERS' EQUATION WITH STOCHASTIC TRANSPORT

Recall that we are dealing with a stochastic Burgers' equation of the form

$$\mathrm{d}u + \left(u(t,x)\,\mathrm{d}t + \sum_{k=1}^{\infty} \xi_k(x) \circ \mathrm{d}W_t^k\right) \cdot \partial_x u = \nu \partial_{xx} u\,\mathrm{d}t,$$

¹Unlike a stopping time for a local solution, a stopping time for a maximal solution is allowed to take the value infinity.

for $x \in \mathbb{T}$ or \mathbb{R} , where $\nu \geq 0$ is constant, $\{\xi_k(x)\}_{k\in\mathbb{N}}$ is an orthonormal basis of some separable Hilbert space \mathcal{H} , and \circ means that the integration is carried out in the Stratonovich sense. In this section, we study the problem of whether shocks can form in the inviscid Burgers' equation with stochastic transport. By using a characteristic argument, we prove that for some classes $\{\xi_k(x)\}_{k\in\mathbb{N}}$, the transport noise cannot prevent shock formation. We also consider a weak formulation of the problem and prove that the shocks satisfy a stochastic version of the Rankine-Hugoniot condition.

3.1. Inviscid Burgers' equation with stochastic transport. The inviscid Burgers' equation with stochastic transport is given by

$$du + \left(u(t,x) dt + \sum_{k=1}^{\infty} \xi_k(x) \circ dW_t^k\right) \cdot \partial_x u = 0, \qquad (3.1)$$

which in integral form is interpreted as

$$u(t,x) = u(0,x) - \int_0^t \left(u(s,x)\partial_x u(s,x) \, \mathrm{d}s + \sum_{k=1}^\infty \xi_k(x)\partial_x u(s,x) \circ \mathrm{d}W_s^k \right), \tag{3.2}$$

for all $x \in \mathbb{T}$ or \mathbb{R} . Also, we will assume throughout this paper that the initial condition is positive, that is, u(0, x) > 0 for all $x \in \mathbb{T}$ or \mathbb{R} .

Consider a process X_t that satisfies the *Stratonovich* SDE

$$dX_t = u(t, X_t) dt + \sum_{k=1}^{\infty} \xi_k(X_t) \circ dW_t^k,$$
 (3.3)

which in Itô form, reads

$$dX_t = \left(u(t, X_t) + \frac{1}{2} \sum_{k=1}^{\infty} \xi_k(X_t) \partial_x \xi_k(X_t)\right) dt + \sum_{k=1}^{\infty} \xi_k(X_t) dW_t^k.$$
 (3.4)

We call this process the *characteristic* of (3.1), analogous to the characteristic lines in the deterministic Burgers' equation. We assume the following conditions on $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$.

Assumption A1. ξ_k is smooth for all $k \in \mathbb{N}$ and together with the Stratonovich-to-Itô correction term $\varphi(x) := \frac{1}{2} \sum_{k=1}^{\infty} \xi_k(x) \partial_x \xi_k(x)$, satisfy the following:

• Lipschitz continuity

$$|\varphi(x) - \varphi(y)| \le C_0 |x - y|, \quad |\xi_k(x) - \xi_k(y)| \le C_k |x - y|, \quad k \in \mathbb{N},$$

$$(3.5)$$

• Linear growth condition

$$|\varphi(x)| \le D_0(1+|x|), \quad |\xi_k(x)| \le D_k(1+|x|), \quad k \in \mathbb{N}$$
(3.6)

for real constants C_0, C_1, C_2, \ldots and D_0, D_1, D_2, \ldots with

$$\sum_{k=1}^{\infty} C_k^2 < \infty, \quad \sum_{k=1}^{\infty} D_k^2 < \infty.$$
 (3.7)

Provided $u(t, \cdot)$ is sufficiently smooth and bounded (hence satisfying Lipschitz continuity and linear growth) until some stopping time τ , and $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$ satisfies the conditions in Assumption A1, the characteristic equation (3.4) is locally well-posed. One feature of the multiplicative noise in (3.1) is that u is transported along the characteristics, that is,

9

we can show that $u(t,x) = (\Phi_t)_* u_0(x)$ for $0 \le t < \tau_{max}$, where Φ_t is the stochastic flow of the SDE (3.4), $(\Phi_t)_*$ represents the pushforward by Φ_t , and (τ_{max}, X_t) is the maximal solution of (3.4). This is an easy corollary of the Itô-Wentzell formula (2.9).

Corollary 3.0.1. Let $u(t, \cdot)$ be $C^3 \cap L^{\infty}$ in space for $0 < t < \tau$. Assume also that $u(\cdot, x)$ is a continuous semimartingale satisfying (3.2), $\partial_x u(\cdot, x)$ is a continuous semimartingale satisfying the spatial derivative of (3.2), and $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$ satisfies the conditions in Assumption A1. If (τ_{max}, X_t) is a maximal solution to (3.4), then $u(t, X_t) = u(0, X_0)$ almost surely for $0 < t < \tau_{max}$.

Remark 3.1. Notice that due to our local well-posedness result (Theorem 1.3) and the maximum principle (Proposition 4.14), one has $u_t \in C^3 \cap L^{\infty}$ for $t < \tau_{max}$ provided u_0 is smooth enough and bounded. For instance, $u_0 \in H^4 \cap L^{\infty}$ is sufficient.

Proof of Corollary 3.0.1. Note that under the given assumptions, $\sigma_0(t, x) := u(t, x)\partial_x u(t, x)$, and $\sigma_k(t, x) := \xi_k(x)\partial_x u(t, x)$ for all $k \in \mathbb{N}$, satisfy the conditions in Theorem 2.5. We take $N_t^0 = t$ and $N_t^k = W_t^k$ for $k \in \mathbb{N}$. Using the Itô-Wentzell formula (2.9) for the stochastic field u(t, x) satisfying (3.2), and the semimartingale X_t , we obtain

$$u(t, X_t) = u(0, X_0) - \int_0^t \left(u(s, X_s) \partial_x u(s, X_s) \, \mathrm{d}s + \sum_{k=1}^\infty \xi_k(X_s) \partial_x u(s, X_s) \circ \mathrm{d}W_s^k \right)$$
$$+ \int_0^t \partial_x u(s, X_s) \circ \mathrm{d}X_s$$
$$= u(0, X_0) - I_1 + I_2.$$

Now, we have $I_1 = I_2$ almost surely so indeed, $u(t, X_t) = u(0, X_0)$ almost surely for $0 < t < \tau_{max}$.

3.2. Results on shock formation. In order to investigate the crossing of characteristics in the stochastic Burgers' equation (3.1) with transport noise, we define the *first crossing time* τ as

$$\tau := \inf_{\substack{a,b \in \mathbb{R} \\ a \neq b}} \left\{ \inf \left\{ t > 0 : X_t^a = X_t^b \right\} \right\},$$
(3.8)

where X_t^a, X_t^b are two characteristics that solve the SDE (3.3) with initial conditions $X_0^a = a$ and $X_0^b = b$. This gives us the first time when two characteristics intersect. In the following, we will show that in the special case $\xi_1(x) = \alpha x + \beta$ (where we only consider one noise term and the other terms ξ_k are identically zero for $k = 2, 3, \ldots$), the first crossing time is equivalent to the first hitting time of the integrated geometric Brownian motion. We note that in this case, equation (3.4) is explicitly solvable, where the general solution is given by

$$X_t^{\gamma} = e^{\alpha W_t} \left(\gamma + (u_0(\gamma) - \alpha\beta) \int_0^t e^{-\alpha W_s} \,\mathrm{d}s + \beta \int_0^t e^{-\alpha W_s} \,\mathrm{d}W_s \right). \tag{3.9}$$

Proposition 3.2. The first crossing time of the inviscid stochastic Burgers' equation (3.1) with $\xi_1(x) = \alpha x + \beta$ for constants $\alpha, \beta \in \mathbb{R}$ and $\xi_k(\cdot) \equiv 0$ for $k = 2, 3, \ldots$ is equivalent to the first hitting time for the integrated geometric Brownian motion $I_t := \int_0^t e^{-\alpha W_s} ds$.

Proof. Consider two arbitrary characteristics X_t^a and X_t^b with $X_0^a = a$ and $X_0^b = b$. From (3.9), one can check that $X_t^a = X_t^b$ if and only if

$$I_t := \int_0^t e^{-\alpha W_s} \, \mathrm{d}s = -\frac{b-a}{u_0(b) - u_0(a)}$$

Now, since the left-hand side is continuous, strictly increasing with $I_0 = 0$, and independent of a and b, we have

$$\tau = \inf_{\substack{a,b \in \mathbb{R} \\ a \neq b}} \left\{ \inf \left\{ t > 0 : I_t = -\frac{b-a}{u_0(b) - u_0(a)} \right\} \right\}$$
$$= \left\{ \inf_{\infty, t} \{t > 0 : I_t = \theta(u_0)^{-1} \}, \quad \text{if } \theta(u_0) > 0 \\ \infty, \quad \text{if } \theta(u_0) = 0 \end{cases},$$
(3.10)

where

$$\theta(u_0) := \sup_{\substack{a,b \in \mathbb{R} \\ a \neq b}} \left\{ \zeta(a,b) \right\}, \quad \zeta(a,b) = \begin{cases} \frac{|u_0(a) - u_0(b)|}{|a-b|}, & \text{if } \frac{u_0(a) - u_0(b)}{a-b} < 0\\ 0, & \text{otherwise} \end{cases}$$

is the steepest negative slope of u_0 . Hence, the first crossing time is equivalent to the first hitting time of the process I_t .

Remark 3.3. Note that the constant β does not affect the first crossing time, hence we can set $\beta = 0$ without loss of generality. Also in the following, we simply write $\xi(\cdot)$ without the index when we only consider one noise term.

As an immediate consequence of Proposition 3.2, we prove that the transport noise with $\xi(x) = \alpha x$ cannot prevent shocks from forming almost surely in the stochastic Burgers' equation (3.1).

Corollary 3.3.1. Let $\xi(x) = \alpha x$ for some $\alpha \in \mathbb{R}$. If the initial profile u_0 has a negative slope, then $\tau < \infty$ almost surely.

Proof. To prove this, it is enough to show that

$$\lim_{t \to \infty} \int_0^t e^{\alpha W_s} \, \mathrm{d}s = \infty \quad a.s.$$

where we have assumed $\alpha > 0$, without loss of generality, and $W_{\bullet} : \mathbb{R}_{\geq 0} \times \Xi \to \mathbb{R}$ is the standard Wiener process on the Wiener space $(\Xi, \mathcal{F}, \mathbb{P})$, adapted to the natural filtration \mathcal{F}_t . This implies that $\tau < \infty$ a.s. by Proposition 3.2.

First, define the set

$$A = \left\{ \omega \in \Xi : \lim_{t \to \infty} \int_0^t e^{\alpha W_s(\omega)} \, \mathrm{d}s < \infty \right\} \subset \Xi.$$

Fixing $\omega \in A$, choose $t_1, t_2, \ldots \in \mathbb{R}_{\geq 0}$ with $t_n < t_{n+1}$, such that $\lim_{n \to \infty} t_n = \infty$ and $\lim_{n \to \infty} (t_{n+1} - t_n) > 0$, and consider the sequence

$$I_n(\omega) = \int_0^{t_n} e^{\alpha W_s(\omega)} \,\mathrm{d}s, \quad n = 1, 2, \dots$$

Clearly, $\{I_n(\omega)\}_{n\in\mathbb{N}}$ is monotonic increasing, and it is also bounded since $\omega \in A$. Hence, it is convergent by the monotone convergence theorem, and in particular, it is a Cauchy sequence. Therefore we have

$$\lim_{n \to \infty} |I_{n+1}(\omega) - I_n(\omega)| = \lim_{n \to \infty} \int_{t_n}^{t_{n+1}} e^{\alpha W_s(\omega)} \,\mathrm{d}s = 0.$$

Since the integrand is strictly positive, this implies $\lim_{t\to\infty} e^{\alpha W_t(\omega)} = 0$, and hence $W_t(\omega) \to -\infty$. On the other hand, for $\omega \in \Xi$ such that $W_t(\omega) \to -\infty$, it is easy to see that $\omega \in A$. This implies that under the identification $\Xi \cong C([0,\infty);\mathbb{R})$, the set A is equivalent to the set of Wiener processes W_t with $W_t \to -\infty$, which is open in $C([0,\infty);\mathbb{R})$ endowed with the norm $\|\cdot\|_{\infty}$ and therefore measurable. In particular, for $\omega \in A$, we have

$$\limsup_{t \to \infty} W_t(\omega) = -\infty,$$

but since $\limsup_{t\to\infty} W_t = +\infty$, a.s., this implies $\mathbb{P}(A) = 0$.

In the following, we show that for a broader class of $\{\xi_k(\cdot)\}_{k\in\mathbb{N}}$, shock formation occurs in expectation provided the initial profile has a sufficiently negative slope. Moreover, no new shocks can develop from positive slopes. We show this by looking at how the slope $\partial_x u$ evolves along the characteristics X_t , which resembles the argument given in [CH18] for the stochastic Camassa-Holm equation.

Theorem 3.4. Consider a characteristic X_t , and a smooth initial profile $u(0, x) = u_0(x)$ such that $\partial_x u(0, X_0) = -\sigma < 0$. If

$$\frac{1}{2}\sum_{k=1}^{\infty} \left((\partial_x \xi_k(x))^2 - \xi_k(x) \partial_{xx} \xi_k(x) \right) > -\sigma, \quad \forall x \in \mathbb{R},$$

then there exists $0 < t_* < \infty$ such that $\lim_{t \to t_*} \mathbb{E}[\partial_x u(t, X_t)] = -\infty$. On the other hand, if $\partial_x u(0, X_0) \ge 0$ and

$$\frac{1}{2}\sum_{k=1}^{\infty} \left((\partial_x \xi_k(x))^2 - \xi_k(x) \partial_{xx} \xi_k(x) \right) < \infty, \quad \forall x \in \mathbb{R},$$

then $\partial_x u(t, X_t) < \infty$ almost surely for all t > 0.

Proof. Taking the spatial derivative of (3.2), and evaluating the stochastic field $\partial_x u(t, x)$ along the semimartingale X_t by the Ito-Wentzell formula (2.9) (again, this is valid due to the local well-posedness result, Theorem 1.3), the process $Y_t := \partial_x u(t, X_t)$ together with X_t satisfy the following coupled Stratonovich SDEs

$$dX_t = u(t, X_t) dt + \sum_{k=1}^{\infty} \xi_k(X_t) \circ dW_t^k,$$
 (3.11)

$$dY_t = -Y_t^2 dt - \sum_{k=1}^{\infty} \partial_x \xi_k(X_t) Y_t \circ dW_t^k.$$
(3.12)

In Itô form, this reads

$$dX_t = \left(u(t, X_t) + \frac{1}{2} \sum_{k=1}^{\infty} \xi_k(X_t) \partial_x \xi_k(X_t)\right) dt + \sum_{k=1}^{\infty} \xi_k(X_t) dW_t^k,$$
(3.13)

$$dY_t = \left(-Y_t^2 + \frac{1}{2}Y_t \sum_{k=1}^{\infty} \left((\partial_x \xi_k(x))^2 - \xi_k(x)\partial_{xx}\xi_k(x)\right)\right) dt - \sum_{k=1}^{\infty} \partial_x \xi_k(X_t)Y_t dW_t^k.$$
(3.14)

Taking the expectation of (3.14) on both sides, we obtain

$$\frac{\mathrm{d}\mathbb{E}[Y_t]}{\mathrm{d}t} = -\mathbb{E}[Y_t^2] + \frac{1}{2}\mathbb{E}\left[Y_t\sum_{k=1}^{\infty}\left((\partial_x\xi_k(x))^2 - \xi_k(x)\partial_{xx}\xi_k(x)\right)\right].$$
(3.15)

Now, assume that there exists a constant $C \in \mathbb{R}$ such that

$$C \le \sum_{k=1}^{\infty} \left((\partial_x \xi_k(x))^2 - \xi_k(x) \partial_{xx} \xi_k(x) \right), \qquad (3.16)$$

for all $x \in \mathbb{R}$. If $Y_0 = -\sigma < 0$, we have $Y_t < 0$ for all t > 0, since Y = 0 is a fixed line in the phase space (X, Y) and therefore cannot be crossed. Hence from (3.16), we have

$$\mathbb{E}\left[Y_t \sum_{k=1}^{\infty} \left((\partial_x \xi_k(X_t))^2 - \xi_k(X_t) \partial_{xx} \xi_k(X_t) \right) \right] \le C \mathbb{E}[Y_t],$$

and (3.15) becomes,

$$\frac{\mathrm{d}\mathbb{E}[Y_t]}{\mathrm{d}t} \leq -\mathbb{E}[Y_t^2] + \frac{C}{2}\mathbb{E}[Y_t]$$

$$= -(\mathbb{E}[Y_t^2] - \mathbb{E}[Y_t]^2) - \mathbb{E}[Y_t]^2 + \frac{C}{2}\mathbb{E}[Y_t]$$

$$\leq -\mathbb{E}[Y_t]^2 + \frac{C}{2}\mathbb{E}[Y_t],$$

$$[V]^2 = \mathbb{E}[(Y - \mathbb{E}[Y_t])^2] > 0$$

since $\mathbb{E}[Y_t^2] - \mathbb{E}[Y_t]^2 = \mathbb{E}\left[(Y_t - \mathbb{E}[Y_t])^2\right] \ge 0.$

Solving this differential inequality, we get

$$\mathbb{E}[Y_t] \le \begin{cases} \frac{-\sigma e^{Ct/2}}{1 - \frac{2\sigma}{C} \left(e^{Ct/2} - 1\right)}, & \text{if } C \neq 0\\ \frac{1}{t - \frac{1}{\sigma}}, & \text{if } C = 0 \end{cases}$$

The right-hand side tends to $-\infty$ in finite time provided $-\sigma < C/2$.

Hence, if

$$-\sigma < C/2 \le \frac{1}{2} \sum_{k=1}^{\infty} \left((\partial_x \xi_k(x))^2 - \xi_k(x) \partial_{xx} \xi_k(x) \right)$$

for all $x \in \mathbb{R}$, then there exists $t_* < \infty$ such that $\lim_{t \to t_*} \mathbb{E}[u_x(t, X_t)] = -\infty$.

Similarly, if $\frac{1}{2} \sum_{k=1}^{\infty} \left((\partial_x \xi_k(x))^2 - \xi_k(x) \partial_{xx} \xi_k(x) \right) < D$ for some $D \in \mathbb{R}$, then for $Y_0 > 0$ we have again

$$\frac{\mathrm{d}\mathbb{E}[Y_t]}{\mathrm{d}t} \le -\mathbb{E}[Y_t]^2 + D\mathbb{E}[Y_t].$$

One can check that $\mathbb{E}[Y_t] < \infty$ for all t > 0, which implies $Y_t < \infty$ almost surely. \Box

Remark 3.5. Blow-up in expectation does not imply pathwise blow-up. It is merely a necessary condition, which suggests that the law of $\partial_x u$ becomes increasingly fat-tailed with time, making it more likely for it to take extreme values. Nonetheless, it is a good indication of blow-up occurring with some probability.

Example 3.6. Consider the set $\{\xi_k(x)\}_{k\in\mathbb{N}} = \left\{\frac{1}{k^2}\sin(kx), \frac{1}{k^2}\cos(kx)\right\}_{k\in\mathbb{N}}$, which forms an orthogonal basis for $L^2(\mathbb{T})$. Then, one can easily check that

$$0 < \sum_{k=1}^{\infty} \left((\partial_x \xi_k(x))^2 - \xi_k(x) \partial_{xx} \xi_k(x) \right) < \infty,$$

for all $x \in \mathbb{T}$, so blow-up occurs in expectation for any initial profile with negative slope, but no new shocks can form from positive slopes.

3.3. Weak solutions. We saw that if the initial profile u_0 has a negative slope, then shocks may form in finite time (almost surely in the linear case $\xi(x) = \alpha x$), so solutions to (3.1) cannot exist in the classical sense. This motivates us to consider *weak solutions* to (3.1) in the sense of Definition 2.3.

Suppose that the profile u is differentiable everywhere except for a discontinuity along the curve $\gamma = \{(t, s(t)) \in [0, \infty) \times M\}$, where $M = \mathbb{T}$ or \mathbb{R} . Then the curve of discontinuity must satisfy the following for u to be a solution of the integral equation (2.10).

Proposition 3.7 (Stochastic Rankine-Hugoniot condition). The curve of discontinuity s(t) of the stochastic Burgers' equation in weak form (2.10) satisfies the following SDE

$$ds_t = \frac{1}{2} \left[\left(u_-(t, s(t)) + u_+(t, s(t)) \right] dt + \sum_{k=1}^{\infty} \xi_k(s(t)) \circ dW_t^k,$$
(3.17)

where $u_{\pm}(t, s(t)) := \lim_{x \to s(t)^{\pm}} u(t, x)$ are the left and right limits of u.

The main obstacle here is that the curve s(t) is not piecewise smooth and therefore we cannot apply the standard divergence theorem, which is how the Rankine-Hugoniot condition is usually derived. Extending classical calculus identities such as Green's theorem on domains with non-smooth boundaries is a tricky issue, but fortunately, there have been several works that extend this result to non-smooth but rectifiable boundaries in [Sha57], and to non-rectifiable boundaries in [HN92, Har93, Har99, LY06].

Lemma 3.8 (Green's theorem for non-smooth boundaries). Let Ω be a bounded domain in the (x, y)-plane such that its boundary $\partial \Omega$ is a Jordan curve and let u, v be sufficiently regular functions in Ω (see remark 3.9 below). Then

$$\int_{\Omega} div(u,v) \, \mathrm{d}x \, \mathrm{d}y = \oint_{\partial\Omega} \left(u \, \mathrm{d}y - v \, \mathrm{d}x \right), \tag{3.18}$$

where the contour integral on the right-hand side can be understood as a limit of a standard contour integral along a smooth approximation of the boundary. Here, the integral is taken in the anti-clockwise direction of the contour.

Remark 3.9. For the above to hold, there must be a pay-off between the regularity of $\partial \Omega$ and the functions u, v (i.e. the less regular the boundary, the more regular the integrand). In particular, the following condition is known:

• $\partial \Omega$ has box-counting dimension d < 2 and u, v is α -Hölder continuous for any $\alpha > d-1$ (Harrison and Norton [HN92]).

Proof of Theorem 3.7. We provide a proof in the case $M = \mathbb{T}$ with only one noise term. Extending it to the case $M = \mathbb{R}$ and countably many noise terms is straightforward. Take the atlas $\{(U_1, \varphi_1), (U_2, \varphi_2)\}$ on $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, where $U_1 := (0, 1), \varphi_1 : (0, 1) \to U_1$ and $U_2 := (-\frac{1}{2}, \frac{1}{2}), \varphi_2 : (0, 1) \to U_2$. Without loss of generality, assume that the shock $s(\cdot)$ starts at time t = 0.

Now, consider a sequence $0 = \tau_0 < \tau_1 < \tau_2 < \ldots$, with $\lim_{n\to\infty} \tau_n = \infty$ such that for all $n \in \mathbb{N}$, the curve $\gamma_n := \{s(t) : t \in [\tau_{n-1}, \tau_n)\}$ is contained in either one of the charts U_1 or U_2 . For convenience, we denote by (U_n, φ_n) to mean the chart (U_1, φ_1) or (U_2, φ_2) that contains γ_n . In local coordinates, we split the domain $\Omega_n := [\tau_{n-1}, \tau_n) \times \varphi_n^{-1}(U_n)$ into two regions (see figure 1)

$$\Omega^{n}_{-} := \{ (t, x) \in [\tau_{n-1}, \tau_n) \times (0, 1) : x \in (0, s(t)) \}, \qquad (3.19)$$

$$\Omega^{n}_{+} := \{ (t, x) \in [\tau_{n-1}, \tau_n) \times (0, 1) : x \in (s(t), 1) \}.$$
(3.20)

For $n \in \mathbb{N}$, consider the following integrals

$$\begin{split} I_n &= \iiint_{\Omega_{-}^n} \left(\left(u \partial_t \varphi + \frac{1}{2} u^2 \partial_x \varphi \right) \mathrm{d}t + u \partial_x \left(\varphi(t, x) \xi(x) \right) \circ \mathrm{d}W_t \right) \mathrm{d}x \\ &= \iint_{\Omega_{-}^n} \mathrm{div}_{x,t} \left(\frac{1}{2} \varphi u(t, x)^2, \, \varphi(t, x) u(t, x) \right) \mathrm{d}x \, \mathrm{d}t + \int_{\tau_{n-1}}^{\tau_n} \left(\int_0^{s(t)} \partial_x \left(\varphi(t, x) \xi(x) u(t, x) \right) \mathrm{d}x \right) \circ \mathrm{d}W_t \\ &- \underbrace{\iint_{\Omega_{-}^n} \varphi \left(\mathrm{d}u + u \partial_x u \, \mathrm{d}t + \xi(x) \partial_x u \circ \mathrm{d}W_t \right) \mathrm{d}x, \text{ and}}_{=0} \end{split}$$

$$\begin{split} J_n &= \iint_{\Omega^n_+} \left(\left(u \partial_t \varphi + \frac{1}{2} u^2 \partial_x \varphi \right) \mathrm{d}t + u \partial_x \left(\varphi(t, x) \xi(x) \right) \circ \mathrm{d}W_t \right) \mathrm{d}x \\ &= \iint_{\Omega^n_+} \mathrm{div}_{x,t} \left(\frac{1}{2} \varphi u(t, x)^2, \, \varphi(x, t) u(t, x) \right) \mathrm{d}x \, \mathrm{d}t + \int_{\tau_{n-1}}^{\tau_n} \left(\int_{s(t)}^1 \partial_x \left(\varphi(t, x) \xi(x) u(t, x) \right) \mathrm{d}x \right) \circ \mathrm{d}W_t \\ &- \underbrace{\iint_{\Omega^n_+} \varphi \left(\mathrm{d}u + u \partial_x u \, \mathrm{d}t + \xi(x) \partial_x u \circ \mathrm{d}W_t \right) \mathrm{d}x}_{=0} . \end{split}$$

Then by Lemma 3.8, we have

$$\begin{split} I_n &= \oint_{\partial \Omega_{-}^n} \left(\frac{1}{2} \varphi u(t,x)^2 \, \mathrm{d}t - \varphi(t,x) u(t,x) \, \mathrm{d}x \right) + \int_{\tau_{n-1}}^{\tau_n} \varphi(t,s(t)) \xi(s(t)) u_-(t,s(t)) \circ \mathrm{d}W_t \\ &= -\int_{\tau_{n-1}}^{\tau_n} \varphi(t,s(t)) \left(u_-(t,s(t)) \, \mathrm{d}s_t - \frac{1}{2} u_-(t,s(t))^2 \, \mathrm{d}t - \xi(s(t)) u_-(t,s(t)) \circ \mathrm{d}W_t \right) \\ &+ \left(\int_A - \int_B - \int_C \right) \left(\frac{1}{2} \varphi u(t,x)^2 \, \mathrm{d}t - \varphi(t,x) u(t,x) \, \mathrm{d}x \right), \end{split}$$

where

$$A := \{(\tau_{n-1}, x) : x \in (0, s(\tau_{n-1}))\}, B := \{(\tau_n, x) : x \in (0, s(\tau_n))\}, C := \{(t, 0) : t \in (\tau_{n-1}, \tau_n)\},$$
 and

$$\begin{split} J_n &= \oint_{\partial \Omega_+^n} \left(\frac{1}{2} \varphi u(t,x)^2 \, \mathrm{d}t - \varphi(x,t) u(t,x) \, \mathrm{d}x \right) - \int_{\tau_{n-1}}^{\tau_n} \varphi(t,s(t)) \xi(s(t)) u_+(t,s(t)) \circ \mathrm{d}W_t \\ &= \int_{\tau_{n-1}}^{\tau_n} \varphi(t,s(t)) \left(u_+(t,s(t)) \, \mathrm{d}s_t - \frac{1}{2} u_+(t,s(t))^2 \, \mathrm{d}t - \xi(s(t)) u_+(t,s(t)) \circ \mathrm{d}W_t \right) \\ &+ \left(\int_D + \int_E + \int_F \right) \left(\frac{1}{2} \varphi u(t,x)^2 \, \mathrm{d}t - \varphi(t,x) u(t,x) \, \mathrm{d}x \right), \end{split}$$

where

N 7

$$D := \{(\tau_{n-1}, x) : x \in (s(\tau_{n-1}), 1)\}, E := \{(\tau_n, x) : x \in (s(\tau_n), 1)\}, F := \{(t, 1) : t \in (\tau_{n-1}, \tau_n)\}$$

One can check by direct calculation that

$$\sum_{n=1}^{N} (I_n + J_n) = -\int_{\mathbb{T}} \varphi(\tau_N, x) u(\tau_N, x) \, \mathrm{d}x + \int_0^{\tau_N} \varphi(t, s(t)) \left[u_+(t, s(t)) - u_-(t, s(t)) \right] \left(\mathrm{d}s_t - \frac{1}{2} \left[u_-(t, s(t)) + u_+(t, s(t)) \right] \, \mathrm{d}t - \xi(s(t)) \circ \mathrm{d}W_t \right) + \int_0^{\tau_N} \varphi(t, s(t)) \left[u_+(t, s(t)) - u_-(t, s(t)) \right] \, \mathrm{d}t$$

where we used the assumption that $\varphi(0, \cdot) \equiv 0$.

Now, from (2.10), we have $\lim_{N\to\infty} \sum_{n=1}^{N} (I_n + J_n) = 0$ and since φ has compact support, there exists N > 0 such that $\varphi(\tau_{N'}, \cdot) \equiv 0$ for all $N' \geq N$. Hence,

$$0 = \lim_{N \to \infty} \sum_{n=1}^{N} (I_n + J_n)$$

= $\int_0^\infty \varphi(t, s(t)) \left[u_+(t, s(t)) - u_-(t, s(t)) \right] \left(\mathrm{d}s_t - \frac{1}{2} \left[u_-(t, s(t)) + u_+(t, s(t)) \right] \mathrm{d}t - \xi(s(t)) \circ \mathrm{d}W_t \right)$

and since φ is arbitrary, we have

$$ds_t = \frac{1}{2} \left[u_-(t, s(t)) + u_+(t, s(t)) \right] dt + \xi(s(t)) \circ dW_t,$$

for all t > 0.

4. Well-posedness results

4.1. Local well-posedness of a stochastic Burgers' equation. Now, we prove local well-posedness of the stochastic Burgers' equation (1.1) with one noise term and $\nu = 0$. In fact, since the techniques used in the proof are essentially the same, we prove local well-posedness of a *more general equation*, which includes (1.1) as a special case. The stochastic Burgers' equation we treat is given by

$$du + u\partial_x u \, dt + \mathcal{Q}u \, dW_t = \frac{1}{2}\mathcal{Q}^2 u \, dt.$$
(4.1)



FIGURE 1. In the proof of the stochastic Rankine-Hugoniot condition (3.17), the domain $\Omega_n := [\tau_{n-1}, \tau_n) \times (0, 1) \subset [0, \infty) \times \mathbb{T}$ is split up into two parts: Ω_{-}^n , which is on the left of the shock curve (t, s(t)) and Ω_{+}^n , which is on the right.

Here \mathcal{Q} represents a first order differential operator

$$\mathcal{Q}u = a(x)\partial_x u + b(x)u,$$

where the coefficients a(x), b(x) are smooth and bounded. We state the main result of this section:

Theorem 4.1. Let $u_0 \in H^s(\mathbb{T})$ for some s > 2 fixed. Then there exists a pathwise unique H^s -maximal solution (τ_{max}, u) of the 1D Burgers' equation (4.1) in the sense of Definition 2.2 with initial datum u_0 . Moreover, either $\tau_{max} = \infty$ or $\limsup_{t \to \tau_{max}} |u(t)|_{H^s} = \infty$, a.s.

We will provide a sketch of the proof, which follows closely the approach developed in [AOB20, CFH19]. For clarity of exposition, let us divide the argument into several steps.

• Step 1: Uniqueness of local solutions. To show pathwise uniqueness of local solutions, one argues by contradiction. More concretely, one can prove that any two different solutions to (4.1) defined up to a stopping time must coincide, as explained in the following proposition.

Proposition 4.2. Let s > 2. Let $\tau : \Xi \to [0, \infty)$ be a stopping time, and $u^1, u^2 : \Xi \times [0, \tau] \times \mathbb{T} \to \mathbb{R}$ be two H^s -local solutions of (4.1) with the same initial datum $u_0 \in H^s(\mathbb{T})$. Then a.s. $u^1 = u^2$ on $[0, \tau]$.

Proof. For this, we refer the reader to [AOB20, CFH19]. It suffices to define $\bar{u} = u^1 - u^2$, and perform standard estimates for the evolution of the L^2 norm of \bar{u} .

- Step 2: Existence and uniqueness of truncated maximal solutions. Following the techniques in [CFH19] for the Euler equation with Lie transport noise, for r > 0 we consider the truncated stochastic Burgers' equation

$$du_r + \theta_r (|\partial_x u_r|_{L^{\infty}}) u_r \partial_x u_r dt + \mathcal{Q} u_r dW_t = \frac{1}{2} \mathcal{Q}^2 u_r dt, \qquad (4.2)$$

where $\theta_r: [0,\infty) \to [0,1]$ is a smooth function such that

$$\theta_r(x) = \begin{cases} 1, & |x| \le r, \\ 0, & |x| \ge 2r. \end{cases}$$

As we explain in the following lemma, local solutions of (4.1) can be constructed by restricting global solutions of (4.2) to a certain stopping time.

Lemma 4.3. Fix r > 0 and $u_0 \in H^s(\mathbb{T})$, s > 2. Let $u_r : \Xi \times [0, \infty) \times \mathbb{T} \to \mathbb{R}$ be an H^s -global solution of (4.2) with initial datum u_0 . Consider the stopping time

$$\tau_r = \inf\left\{t \ge 0 : |u_r(t, \cdot)|_{H^s} \ge \frac{r}{C}\right\},\,$$

where C is chosen in such a way that the following inequality holds:

$$\left|\partial_x u_r\right|_{L^{\infty}} \le C \left|u_r\right|_{H^s},$$

which can be guaranteed thanks to the Sobolev embedding (2.3). Then (τ_r, u_r) is a local solution to the stochastic Burgers' equation (4.1).

Proof of Lemma 4.3. The proof is straightforward by construction. For any $t \in [0, \tau_r]$, we have that

$$\left|\partial_x u_r\right|_{L^{\infty}} \le C \left|u_r\right|_{H^s} \le r_s$$

and therefore $\theta_r(|\partial_x u_r|_{L^{\infty}}) = 1.$

Let us state the result which is the *cornerstone* for proving existence and uniqueness of maximal local solutions of the stochastic Burgers' equation (4.1).

Proposition 4.4. Given r > 0 and $u_0 \in H^s(\mathbb{T}, \mathbb{R})$ for s > 2, there exists a pathwise unique global H^s -solution u of the truncated stochastic Burgers' equation (4.2).

It is very easy to check that once Proposition 4.4 is proven, Theorem 4.1 follows immediately (cf. [CFH19]). Therefore, we focus our efforts on showing Proposition 4.4.

• Step 3: Global existence of solutions of the hyper-regularised truncated stochastic Burgers' equation. In order to show the global well-posedness of the truncated equation, we consider the following hyper-regularisation of (4.2)

$$\mathrm{d}u_r^{\nu} + \theta_r (|\partial_x u_r^{\nu}|_{L^{\infty}}) u_r^{\nu} \partial_x u_r^{\nu} \,\mathrm{d}t + \mathcal{Q}u_r^{\nu} \,\mathrm{d}W_t = \nu \partial_{xx}^{s'} u_r^{\nu} \,\mathrm{d}t + \frac{1}{2} \mathcal{Q}^2 u_r^{\nu} \,\mathrm{d}t, \tag{4.3}$$

where $\nu > 0$ is a positive parameter and s' = 2[s] + 1. Notice that we have added dissipation in order to be able to perform our computations rigorously. Equation (4.3) is understood in the *mild sense*, i.e., as a solution to an integro-differential equation, which we discuss in the next step (see (4.4)).

Proposition 4.5. For every $\nu, r > 0$ and initial datum $u_0 \in H^s(\mathbb{T})$ for s > 2, there exists a pathwise unique global strong solution u_r^{ν} of (4.3) in the class $L^2(\Xi; C([0, T]; H^s(\mathbb{T})))$, for all T > 0. Moreover, its paths gain instant extra regularity $C([\delta, T]; H^{s+2}(\mathbb{T}))$, for every $T > \delta > 0$.

Proof of Proposition 4.5. The proof is based on a classical fixed point iteration argument which employs Duhamel's principle (see [CFH19]). We omit the subscripts ν and r throughout the proof for simplicity, but it should be kept in mind that our functions depend on those parameters. Given $u_0 \in H^s(\mathbb{T})$, consider the mild formulation of the hyper-regularised truncated equation (4.3)

$$u(t) = (\Upsilon u)(t), \tag{4.4}$$

where

$$(\Upsilon u)(t) = e^{tA}u_0 - \int_0^t e^{(t-s)A}W_\theta u(s) \,\mathrm{d}s + \int_0^t e^{(t-s)A}Lu(s) \,\mathrm{d}s - \int_0^t e^{(t-s)A}Ru(s) \,\mathrm{d}W_s$$

t > 0 and we have employed the notation $A = \nu \partial_{xx}^{s'}$, $W_{\theta}u = \theta_r(|\partial_x u|_{L^{\infty}})u\partial_x u$, $Lu = \frac{1}{2}Q^2u$, and Ru = Qu. Define the space $\mathcal{W}_T = L^2(\Xi; C([0, T]; H^s(\mathbb{T})))$. One can show that Υ is a contraction on \mathcal{W}_T by following classical arguments as in [CFH19]. Therefore, by applying Picard's iteration, a local solution can be constructed. To extend it to a global one, it is sufficient to show that for any given T > 0 and initial datum $u_0 \in H^s(\mathbb{T}), s > 2$, the following bound is available

$$\mathbb{E}\left[\sup_{t\in[0,T]}\left|u(t)\right|_{H^s}^2\right] \le C(T),\tag{4.5}$$

for a finite constant $C(T) < \infty$, so that one can patch together each local solution to cover any finite time interval [0, T]. We will prove estimate (4.5) further below. Furthermore, by standard properties of the semigroup e^{tA} (cf. [Gol85]), one can prove that for positive times $T > \delta > 0$, each term in the mild equation (4.4) enjoys higher regularity, namely, $u \in L^2(\Xi; C([\delta, T]; H^{s+2}(\mathbb{T})))$. All the computations are omitted and can be carried out easily by mimicking the same ideas as in [CFH19, AOB20]. \Box

• Step 4: Limiting and compactness argument. The main objective of this step is to show that the family of solutions $\{u_r^{\nu}\}_{\nu>0}$ of the hyper-regularised stochastic Burgers' equation (4.3) is compact in a particular sense and therefore we are able to extract a subsequence converging strongly to a solution of the truncated stochastic Burgers' equation (4.2) in a convenient space. The central idea for proving this is to show compactness of the probability laws of this family. Consequently, we demonstrate that these laws are *tight* in a suitable metric space. Let T > 0 and define the Polish space E by

$$E = C([0,T]; H^{\beta}(\mathbb{T})), \quad \beta \ge 2.$$

$$(4.6)$$

Assume that the laws of $\{u_r^{\nu}\}_{\nu>0}$ are tight in E. Once this is proven, one only needs to invoke standard stochastic partial differential equations arguments based on the Skorokhod's representation and Prokhorov's theorem to conclude that there exists a subsequence of $\{u_r^{\nu}\}_{\nu>0}$ such that solutions of equation (4.3) converge to solutions of (4.2) in the weak limit in the Polish space E (4.6). A more thorough approach can be found in [CFH19, GHV14]. In the next proposition, we present the main argument to show that the sequence of laws are indeed tight.

Proposition 4.6. Assume that for some $\alpha > 0$, $M \in \mathbb{N}$, there exist constants $C_1(T)$ and $C_2(T)$ such that

$$\mathbb{E}\left[\sup_{t\in[0,T]}|u_r^{\nu}(t)|_{H^s}^4\right] \le C_1(T),\tag{4.7}$$

$$\mathbb{E}\left[\int_{0}^{T}\int_{0}^{T}\frac{|u_{r}^{\nu}(t)-u_{r}^{\nu}(s)|_{H^{-M}}^{4}}{|t-s|^{1+4\alpha}}\,\mathrm{d}t\,\mathrm{d}s\right]\leq C_{2}(T),\tag{4.8}$$

uniformly in ν . Then the sequence $\{u_r^{\nu}\}_{\nu>0}$ is tight in E.

Proof of Proposition 4.6. We employ the following lemma, which can be found in [CFH19], was originally proved in [Sim86], and constitutes a variation of the classical Aubin-Lions Lemma.

Lemma 4.7. Suppose that X, Y, Z are separable Hilbert spaces with continuous dense embedding $X \hookrightarrow Y \hookrightarrow Z$ such that there exists $\theta \in (0, 1)$ and M > 0 verifying

$$|v|_Y \le M |v|_X^{1-\theta} |v|_Z^{\theta}$$

for all $v \in X$. Assume that $X \hookrightarrow Y$ is a compact embedding. Let $\alpha > 0$. Then

$$L^{\infty}([0,T];X) \cap W^{\alpha,4}([0,T];Z) \hookrightarrow C([0,T];Y)$$

is a compact embedding.

In Lemma 4.7 we select

$$X = H^s(\mathbb{T}), \quad Y = H^\beta(\mathbb{T}), \quad Z = H^{-M}(\mathbb{T}),$$

where $\beta \geq 2$ as specified before and we impose the extra condition $\beta < s$ so that the embedding of X into Y is compact. We also choose $\alpha \in (0, 1)$. Therefore we obtain that $L^{\infty}([0, T]; H^{s}(\mathbb{T})) \cap W^{\alpha, 4}([0, T]; H^{-M}(\mathbb{T}))$ is compactly embedded in E. By hypotheses (4.7)-(4.8) [CFH19], the family of laws of $\{u_{r}^{\nu}\}_{\nu>0}$ is \mathbb{P} -a.s. supported on the space

$$E_0 = L^{\infty}([0,T]; H^s(\mathbb{T})) \cap W^{\alpha,4}([0,T]; H^{-M}(\mathbb{T})),$$

and it suffices to prove that this family is tight in E_0 . For $A_1, A_2, A_3 > 0$, define the set

$$B = \left\{ f : [0,T] \times \mathbb{T} \to \mathbb{R} : \sup_{t \in [0,T]} |f(t)|_{H^s}^4 \le A_1, \int_0^T |f(t)|_{H^{-M}}^4 \, \mathrm{d}t \le A_2, \\ \int_0^T \int_0^T \int_0^T \frac{|f(t) - f(s)|_{H^{-M}}^4}{|t - s|^{1 + 4\alpha}} \, \mathrm{d}t \, \mathrm{d}s \le A_3 \right\},$$

which is relatively compact in E_0 . It is enough to show that for every ϵ , there exist $A_1, A_2, A_3 > 0$ such that $\mathbb{P}(u_r^{\nu} \in B^c) \leq \epsilon$. Fix $\epsilon > 0$. Invoking Markov's inequality and taking into account hypothesis (4.7), we have that

$$\mathbb{P}\left(\sup_{t\in[0,T]}|u_{r}^{\nu}(t)|_{H^{s}}^{4}>A_{1}\right)\leq\frac{\mathbb{E}\left[\sup_{t\in[0,T]}|u_{r}^{\nu}(t)|_{H^{s}}^{4}\right]}{A_{1}}\leq\frac{C_{1}(T)}{A_{1}},$$

and this is smaller than $\epsilon/3$ if we choose A_1 sufficiently large. Moreover, since $|f|_{H^{-M}} \lesssim |f|_{H^s}$,

$$\mathbb{P}\left(\int_{0}^{T} |u_{r}^{\nu}(t)|_{H^{-M}}^{4} dt > A_{2}\right) \leq \mathbb{P}\left(T \sup_{t \in [0,T]} |u_{r}^{\nu}(t)|_{H^{-M}}^{4} > A_{2}\right)$$
$$\lesssim \mathbb{P}\left(T \sup_{t \in [0,T]} |u_{r}^{\nu}(t)|_{H^{s}}^{4} > A_{2}\right) \leq \frac{C_{1}(T)T}{A_{2}},$$

which can also be made arbitrarily small. A similar argument applies to the set

$$\mathbb{P}\left(\int_{0}^{T}\int_{0}^{T}\frac{|u_{r}^{\nu}(t)-u_{r}^{\nu}(s)|_{H^{-M}}^{4}}{|t-s|^{1+4\alpha}}\,\mathrm{d}t\,\mathrm{d}s>A_{3}\right),\$$

by using hypothesis (4.8). Hence we conclude that $\mathbb{P}(u_r^{\nu} \in B^c) \leq \epsilon$ if we choose A_1, A_2, A_3 large enough, as desired.

• Step 5: Hypothesis estimates. We are left to show that hypothesis (4.7)-(4.8) hold. First, we prove that condition (4.7) implies condition (4.8). By following the techniques in [CFH19] and using equation (4.3), one can obtain

$$\begin{split} \mathbb{E}\left[\left|u_{r}^{\nu}(t)-u_{r}^{\nu}(s)\right|_{H^{-M}}^{4}\right] &\lesssim (t-s)^{3} \int_{s}^{t} \mathbb{E}\left[\theta_{r}(\left|\partial_{x}u_{r}^{\nu}\right|_{L^{\infty}})^{4}\left|u_{r}^{\nu}\partial_{x}u_{r}^{\nu}\right|_{H^{-M}}^{4}\right] \mathrm{d}\gamma \\ &+ (t-s)^{3} \int_{s}^{t} \mathbb{E}\left[\nu\left|\partial_{xx}^{s'}u_{r}^{\nu}\right|_{H^{-M}}^{4}\right] \mathrm{d}\gamma + (t-s)^{3} \int_{s}^{t} \mathbb{E}\left[\left|\mathcal{Q}^{2}u_{r}^{\nu}\right|_{H^{-1}}^{4}\right] \mathrm{d}\gamma \\ &+ \mathbb{E}\left[\left|\left|\int_{s}^{t}\mathcal{Q}u_{r}^{\nu}\,\mathrm{d}W_{\gamma}\right|\right|_{L^{2}}^{4}\right], \end{split}$$

for $0 \leq s < t \leq T$. It is easy to infer that

$$\int_{s}^{t} \mathbb{E}\left[\theta_{r}(|\partial_{x}u_{r}^{\nu}|_{L^{\infty}})^{4} |u_{r}^{\nu}\partial_{x}u_{r}^{\nu}|_{H^{-M}}^{4}\right] \mathrm{d}\gamma \lesssim \int_{s}^{t} \mathbb{E}\left[|u_{r}^{\nu}|_{H^{s}}^{4}\right] \mathrm{d}\gamma \le A_{1}(t-s),$$

since

$$|u_r^{\nu}\partial_x u_r^{\nu}|_{H^{-M}} \lesssim |\partial_x u_r^{\nu}|_{L^{\infty}} |u_r^{\nu}|_{H^s},$$

where we have taken into account hypothesis (4.7). In a similar way, one can check that for M = 3[s'] + 2,

$$\int_{s}^{t} \mathbb{E}\left[\left|\nu\partial_{xx}^{s'}u_{r}^{\nu}\right|_{H^{-M}}^{4}\right] \mathrm{d}\gamma \lesssim \int_{s}^{t} \mathbb{E}\left[\left|u_{r}^{\nu}\right|_{H^{s}}^{4}\right] \mathrm{d}\gamma \leq A_{2}(t-s),$$

since

$$\left|\partial_{xx}^{s'} u_r^{\nu}\right|_{H^{-M}} \lesssim |u_r^{\nu}|_{H^s}.$$

By similar techniques, we have

$$\int_{s}^{t} \mathbb{E}\left[\left|\mathcal{Q}^{2}u_{r}^{\nu}\right|_{H^{-1}}^{4}\right] \mathrm{d}\gamma \leq A_{3}(t-s),$$

and (take into account Burkholder-Davis-Gundy as in [CFH19])

$$\mathbb{E}\left[\left\|\left|\int_{s}^{t} \mathcal{Q}u_{r}^{\nu} \,\mathrm{d}W_{\gamma}\right|\right\|_{L^{2}}^{4}\right] \leq A_{4}(t-s)^{2}.$$

Combining the above estimates, we deduce that

$$\mathbb{E}\left[|u_r^{\nu}(t) - u_r^{\nu}(s)|_{H^{-M}}^4\right] \le A_5(T)(t-s)^2.$$

Hence for $0 < \alpha < 1/2$,

$$\mathbb{E}\left[\int_{0}^{T}\int_{0}^{T}\frac{|u_{r}^{\nu}(t)-u_{r}^{\nu}(s)|_{H^{-M}}^{4}}{|t-s|^{1+4\alpha}}\,\mathrm{d}t\,\mathrm{d}s\right] \leq \mathbb{E}\left[\int_{0}^{T}\int_{0}^{T}\frac{A_{5}(T)}{|t-s|^{4\alpha-1}}\,\mathrm{d}t\,\mathrm{d}s\right] \leq C_{1}(T).$$

We are left to prove that hypothesis (4.7) holds, i.e.

$$\mathbb{E}\left[\sup_{t\in[0,T]}|u_r^{\nu}(t)|_{H^s}^4\right] \le C_2(T).$$

$$(4.9)$$

This is the most difficult part of the hypotheses estimates. We compute the evolution of the \dot{H}^s -norm of u_r^{ν} . Indeed, the equation for the L^2 -norm of $\Lambda^s u_r^{\nu}$ in (4.3) is

$$\frac{1}{2} |\Lambda^{s} u_{r}^{\nu}(t)|_{L^{2}}^{2} = \frac{1}{2} |\Lambda^{s} u_{r}^{\nu}(0)|_{L^{2}}^{2}
- \int_{0}^{t} \langle \theta_{r}(|\partial_{x} u_{r}^{\nu}|_{L^{\infty}}) \Lambda^{s}(u_{r}^{\nu} \partial_{x} u_{r}^{\nu}), \Lambda^{s} u_{r}^{\nu} \rangle_{L^{2}} d\gamma
- \int_{0}^{t} \langle \Lambda^{s} \mathcal{Q} u_{r}^{\nu}, \Lambda^{s} u_{r}^{\nu} \rangle_{L^{2}} dW_{\gamma} + \int_{0}^{t} \langle \nu \Lambda^{s} \partial_{xx}^{s'} u_{r}^{\nu}, \Lambda^{s} u_{r}^{\nu} \rangle_{L^{2}} d\gamma
+ \frac{1}{2} \int_{0}^{t} \langle \Lambda^{s} \mathcal{Q}^{2} u_{r}^{\nu}, \Lambda^{s} u_{r}^{\nu} \rangle_{L^{2}} d\gamma + \frac{1}{2} \int_{0}^{t} \langle \Lambda^{s} \mathcal{Q} u_{r}^{\nu}, \Lambda^{s} \mathcal{Q} u_{r}^{\nu} \rangle_{L^{2}} d\gamma, \qquad (4.10)$$

for $t \in [0, T]$. The nonlinear term can be estimated as

$$\int_{\mathbb{T}} \Lambda^s (u_r^{\nu} \partial_x u_r^{\nu}) \Lambda^s u_r^{\nu} \, \mathrm{d}V = \int_{\mathbb{T}} (u_r^{\nu} \Lambda^s \partial_x u_r^{\nu}) \Lambda^s u_r^{\nu} \, \mathrm{d}V + \int_{\mathbb{T}} [\Lambda^s, u_r^{\nu}] \partial_x u_r^{\nu} \Lambda^s u_r^{\nu} \, \mathrm{d}V$$
$$= I_1 + I_2.$$

Integrating by parts, we note that I_1 can be rewritten and bounded as

$$|I_1| = \frac{1}{2} \left| \int_{\mathbb{T}} \partial_x u_r^{\nu} (\Lambda^s u_r^{\nu})^2 \, \mathrm{d}V \right| \le |\partial_x u_r^{\nu}|_{L^{\infty}} |\Lambda^s u_r^{\nu}|_{L^2}^2.$$

 I_2 can be estimated via the Kato-Ponce commutator estimate (2.4)

Putting together the estimates for I_1 and I_2 , we get

$$\left| \int_{\mathbb{T}} \Lambda^{s}(u_{r}^{\nu} \partial_{x} u_{r}^{\nu}) \Lambda^{s} u_{r}^{\nu} \, \mathrm{d}V \right| \lesssim \left| \partial_{x} u_{r}^{\nu} \right|_{L^{\infty}} \left| \Lambda^{s} u_{r}^{\nu} \right|_{L^{2}}^{2}$$

We note that the dissipative term

$$\left\langle \nu \Lambda^s \partial_{xx}^{s'} u_r^{\nu}, \Lambda^s u_r^{\nu} \right\rangle_{L^2} = -\nu \left| \Lambda^{s+2[s]+1} u_r^{\nu} \right|_{L^2}^2 \le 0$$

has the "good" sign so it can be dropped. The last two terms in (4.10) can be treated in a similar fashion by taking into account estimate (2.6). The previous techniques also serve to treat the evolution of $|u_r^{\nu}|_{L^2}$ (this is a much simpler case). Denote

$$M_t = \int_0^t (\langle \mathcal{Q}u_r^{\nu}, u_r^{\nu} \rangle_{L^2} + \langle \Lambda^s \mathcal{Q}u_r^{\nu}, \Lambda^s u_r^{\nu} \rangle_{L^2}) \, \mathrm{d}W_{\gamma}, \quad t \in [0, T].$$

We drop the parameter dependence on u_r^{ν} to make the notation simpler. By putting together the previous estimates, we have

$$|u(t)|_{H^s}^2 \lesssim |u(0)|_{H^s}^2 + |M_t| + \int_0^t |u|_{H^s}^2 \,\mathrm{d}\gamma,$$

where the constant in the last inequality depends on r. Integrating the above expression against $\exp(t)$ and squaring as in [CFH19], we obtain

$$|u(t)|_{H^s}^4 \lesssim \exp(t)(|u(0)|_{H^s}^4 + |M_t|^2).$$

By applying supremum and expectation on both sides of the equation above, we have

$$\mathbb{E}\left[\sup_{\gamma\in[0,t]}|u(\gamma)|_{H^s}^4\right] \lesssim \exp(t)\left(|u(0)|_{H^s}^4 + \mathbb{E}\left[\sup_{\gamma\in[0,t]}|M_{\gamma}|^2\right]\right),\tag{4.11}$$

where we remind that $t \in [0, T]$. We apply Burkholder-Davis-Gundy inequality (2.7) in order to obtain the bound

$$\mathbb{E}\left[\sup_{\gamma\in[0,t]}|M_{\gamma}|^{2}\right] \lesssim \mathbb{E}[[M]_{t}], \qquad (4.12)$$

where

$$[M]_t = \int_0^t (\langle \mathcal{Q}u, u \rangle_{L^2} + \langle \Lambda^s \mathcal{Q}u, \Lambda^s u \rangle_{L^2})^2 \, \mathrm{d}\gamma.$$

Integrating by parts, we can derive

$$|\langle \Lambda^s \mathcal{Q} u, \Lambda^s u \rangle_{L^2}| \lesssim |u|_{H^s}^2, \qquad (4.13)$$

as in the proof of (2.6) in [AOB20] or in the appendix of [AOB20]. The constant in the last inequality depends on the $W^{s+1,\infty}$ -norm of the coefficients. Therefore,

$$\mathbb{E}[[M]_t] \lesssim \int_0^t \mathbb{E}\left[\sup_{r \in [0,t]} |u(r)|_{H^s}^4\right] \mathrm{d}\gamma.$$
(4.14)

Hence, combining (4.11)-(4.14), together with application of Grönwall's inequality yields

$$\mathbb{E}\left[\sup_{t\in[0,T]}|u(t)|_{H^s}^4\right]\leq C_2(T).$$

4.2. Blow-up criterion. We are now interested in deriving a blow-up criterion for the stochastic Burgers' equation (1.1) with $\nu = 0$. First of all, we note that for the deterministic Burgers' equation

$$u_t + u\partial_x u = 0, \tag{4.15}$$

there is a well-known blow-up criterion available. In the deterministic case, the following theorem of local existence and uniqueness of strong solutions holds.

Theorem 4.8. Let $u_0 \in H^s(\mathbb{T})$ with s > 3/2. Then there exists T > 0 and $u \in \mathbb{T}$ $C([0,T]; H^{s}(\mathbb{T})) \cap C^{1}([0,T]; H^{s-1}(\mathbb{T}))$ solving equation (4.15). Moreover, if v is another function with the same properties, necessarily u = v.

The above result is classic and numerous proofs are available in the literature (see [KNS08] for a related cutting edge result and the references therein). The blow-up criterion for equation (4.15) is the following.

Theorem 4.9 (Blow-up criterion for deterministic Burgers'). Assume that $u_0 \in H^s(\mathbb{T})$, $s > 3/2, T^* > 0$, and $u : [0, T^*) \times \mathbb{T} \to \mathbb{R}$ is a local solution of (4.15). Then the following statements are equivalent

- $\lim_{t \to T^*} |u(t, \cdot)|_{H^s} = \infty.$ $\int_0^{T^*} |\partial_x u(t, \cdot)|_{L^\infty} dt = \infty.$

In the rest of this subsection, we focus on proving the following stochastic version of Theorem 4.9.

Theorem 4.10 (Blow-up criterion for stochastic Burgers'). Assume that $u_0 \in H^s(\mathbb{T})$, s > 2, and $u : \Xi \times [0, \tau_{max}) \times \mathbb{T} \to \mathbb{R}$ is a maximal solution of (4.1). If $\tau_{max} < \infty$, then $\int_0^{\tau_{max}} |\partial_x u(t,\cdot)|_{L^{\infty}} \, \mathrm{d}t = \infty \text{ a.s. Moreover, if } 0 \le \tau < \tau_{max} \text{ is a smaller stopping time,}$ necessarily $\int_0^\tau |\partial_x u(t, \cdot)|_{L^\infty} \, \mathrm{d}t < \infty$ a.s.

Proof of Theorem 4.10. By following the argument in [CFH19], we start by noting that it is clear that if $0 \le \tau < \tau_{max}$ is a smaller stopping time, necessarily $\int_0^\tau |\partial_x u(t, \cdot)|_{L^\infty} dt < \infty$ a.s. This is guaranteed by the embedding

$$|\partial_x u|_{L^{\infty}} \lesssim |u|_{H^s} \,. \tag{4.16}$$

Proving $\int_0^{\tau_{max}} |\partial_x u(t,\cdot)|_{L^{\infty}} dt = \infty$ is more involved. For this, we consider the hyperregularised truncated Burgers' equation (4.3) and define the following stopping times:

$$\begin{aligned} \tau^s &= \lim_{n \to \infty} \tau_n^s, \qquad \tau_n^s = \inf \left\{ t \ge 0 : |u_r^{\nu}(t, \cdot)|_{H^s} \ge n \right\}, \\ \tau^{\infty} &= \lim_{n \to \infty} \tau_n^{\infty}, \qquad \tau_n^{\infty} = \inf \left\{ t \ge 0 : \int_0^t |\partial_x u_r^{\nu}(s, \cdot)|_{L^{\infty}} \, \mathrm{d}s \ge n \right\} \end{aligned}$$

We claim that $\tau^{\infty} = \tau^s$. By again making use of (4.16), it is easy to check $\tau^s \leq \tau^{\infty}$. For simplicity, we omit the subscripts ν and r throughout the proof. Imitating the techniques from the previous subsection, we arrive at

$$\frac{1}{2} \mathrm{d} |\Lambda^{s} u|_{L^{2}}^{2} + \theta(|\partial_{x} u|_{L^{\infty}}) \langle \Lambda^{s}(u \partial_{x} u), \Lambda^{s} u \rangle_{L^{2}} \mathrm{d} t + \langle \Lambda^{s} \mathcal{Q} u, \Lambda^{s} u \rangle_{L^{2}} \mathrm{d} W_{t}$$
$$= \nu \langle \Lambda^{s+2s'} u, \Lambda^{s} u \rangle_{L^{2}} \mathrm{d} t + \frac{1}{2} \langle \Lambda^{s} \mathcal{Q}^{2} u, \Lambda^{s} u \rangle_{L^{2}} \mathrm{d} t + \frac{1}{2} \langle \Lambda^{s} \mathcal{Q} u, \Lambda^{s} \mathcal{Q} u \rangle_{L^{2}} \mathrm{d} t.$$

By integration by parts and standard estimates (use (2.6) for estimating the last two terms), one gets

$$\left| \Lambda^s u \right|_{L^2}^2 + 2 \langle \Lambda^s \mathcal{Q} u, \Lambda^s u \rangle_{L^2} \, \mathrm{d} W_t \lesssim \left(1 + \left| \partial_x u \right|_{L^\infty} \right) \left| \Lambda^s u \right|_{L^2}^2 \, \mathrm{d} t.$$

By also deriving a similar estimate for s = 0 (which follows in a simpler manner by application of the techniques in the previous subsection), we obtain

$$d|u|_{H^s}^2 + 2(\langle \mathcal{Q}u, u \rangle_{L^2} + \langle \Lambda^s \mathcal{Q}u, \Lambda^s u \rangle_{L^2}) dW_t \lesssim (1 + |\partial_x u|_{L^{\infty}}) |u|_{H^s}^2 dt.$$

$$(4.17)$$

Finally, we treat the stochastic term. Without loss of generality, we assume $|u|_{H^s} \ge \epsilon > 0$ (otherwise add a positive constant to this function) and by applying Itô's formula in L^2 [KR81] to the logarithm, we obtain

$$d\log(|u|_{H^s}^2) = \frac{d|u|_{H^s}^2}{|u|_{H^s}^2} - \frac{dN_t}{2(|u|_{H^s}^2)^2},$$
(4.18)

where N_t is defined by

$$N_t = 4 \int_0^t (\langle \mathcal{Q}u, u \rangle_{L^2} + \langle \Lambda^s \mathcal{Q}u, \Lambda^s u \rangle_{L^2})^2 \, \mathrm{d}\gamma.$$

Notice that we have the bound

$$|N_t| \lesssim |u|_{H^s}^2 \,, \tag{4.19}$$

as we have already indicated before (see the proof of (2.6) in the appendix of [AOB20]), where the constant in the last inequality depends on the $W^{s+1,\infty}$ -norm of the coefficients of Q. Plug (4.17) and (4.19) into (4.18) to derive the estimate

$$d\log(|u|_{H^s}^2) \lesssim \frac{(1+|\partial_x u|_{L^{\infty}}) |u|_{H^s}^2}{|u|_{H^s}^2} dt + dM_t,$$
(4.20)

where M_t is the local martingale

$$M_t = \int_0^t \frac{\langle \mathcal{Q}u, u \rangle_{L^2} + \langle \Lambda^s \mathcal{Q}u, \Lambda^s u \rangle_{L^2}}{|u|_{H^s}^2} \, \mathrm{d}W_\gamma,$$

for any t > 0. Expressing (4.20) in integral form, we have

$$\log(|u(t)|_{H^s}^2) \lesssim \log(|u(0)|_{H^s}^2) + \int_0^t (1 + |\partial_x u|_{L^\infty}) \,\mathrm{d}\gamma + \int_0^t \mathrm{d}M_\gamma.$$
(4.21)

By applying Burkholder-Davis-Gundy inequality (see (2.7)), we can control the local martingale term by estimating

$$[M]_t = \int_0^t \frac{(\langle \mathcal{Q}u, u \rangle_{L^2} + \langle \Lambda^s \mathcal{Q}u, \Lambda^s u \rangle_{L^2})^2}{|u|_{H^s}^4} \,\mathrm{d}\gamma \lesssim t.$$

Here, we have employed again the arguments in the appendix of [AOB20] to bound the numerator in the fraction above. Burkholder-Davis-Gundy then yields

$$\mathbb{E}\left[\sup_{s\in[0,t]}\left|\int_{0}^{s}\mathrm{d}M_{\gamma}\right|\right]\lesssim\sqrt{t}.$$
(4.22)

Taking expectation in (4.21) and using estimate (4.22), we establish that for any $n, m \in \mathbb{N}$,

$$\mathbb{E}\left[\sup_{\gamma\in[0,\tau_n^{\infty}\wedge m]}\log(|u(\gamma)|_{H^s}^2)\right]\lesssim \log(|u_0|_{H^s}^2)+m(1+n)+\sqrt{m}<\infty.$$

Therefore

$$\mathbb{E}\left[\log\left(\sup_{\gamma\in[0,\tau_n^\infty\wedge m]}|u(\gamma)|_{H^s}^2\right)\right]<\infty,$$

which in particular means that for $n, m \in \mathbb{N}$, $\sup_{\gamma \in [0, \tau_n^{\infty} \wedge m]} |u(\gamma)|_{H^s}$ is finite a.s. Hence

$$\mathbb{P}\left(\sup_{\gamma\in[0,\tau_n^{\infty}\wedge m]}|u(\gamma)|_{H^s}^2<\infty\right)=1,$$
(4.23)

for every $n, m \in \mathbb{N}$, which from [CFH19] implies $\tau^{\infty} \leq \tau^s$. Therefore $\tau^{\infty} = \tau^s$. Recall that we have omitted subscripts throughout this proof but $u = u_r^{\nu}$. By application of Fatou's Lemma we also obtain (4.23) in the limit $\nu \to 0, r \to \infty$, and we can show that $\tau^{\infty} = \tau^s$ also holds in the limit, concluding our argument. The few steps missing can be checked in [CFH19].

4.3. Global well-posedness of a viscous stochastic Burgers' equation. The viscous stochastic Burgers' equation is given by

$$du + u\partial_x u \,dt + \mathcal{Q}u \,dW_t = \frac{1}{2}Q^2 u \,dt + \nu \partial_{xx} u \,dt.$$
(4.24)

We assume $Q = \xi(x)\partial_x$, s > 2.

Remark 4.11. Observe that from our assumption on Q, (4.24) is simply (1.1) with one noise term, but we wish to keep the Q-notation for convenience.

We establish the global well-posedness of strong solutions of (4.24). More concretely, we prove the following theorem.

Theorem 4.12. Let $u_0 \in H^s(\mathbb{T})$ for some s > 2. Then there exists a pathwise unique maximal H^s -global solution of the viscous stochastic Burgers' equation (4.24).

The proof follows the same strategy as the local existence proof for the Burgers' equation without viscosity (4.1) that we provided in Subsection 4.1. The most important part is proving the following a priori estimate. In this estimate, we assume that u is smooth enough, but the rigorous way to do this is to regularise the equation first as we did in Subsection 4.1.

Proposition 4.13. Let $u_0 \in H^s(\mathbb{T})$, T > 0, and $u : \Xi \times [0, T] \times \mathbb{T} \to \mathbb{R}$ be a solution to equation (4.24) that we assume to be smooth enough. Then there exists a constant such that ²

$$\mathbb{E}\left[\sup_{t\in[0,T]}|u(t)|_{H^s}^2\right] \le C_*(T).$$
(4.25)

Once the a priori estimate (4.25) is established, one can repeat the arguments in Subsection 4.1 to obtain Theorem 4.12. However, since this is repetitive and tedious, we do not explicitly carry out these arguments here. From now on, we focus on proving Proposition 4.13.

²The attentive reader might ask why the H^s -norm of the velocity in (4.25) is taken to the power of two instead of four as in (4.7), but the arguments we carry out together with the control we provide for the L^{∞} -norm of u immediately yield the bound for the fourth power.

Proof. We start by computing the evolution of the L^2 -norm of the solution u. First note that the viscosity term has the good sign since

$$\nu \langle \partial_{xx} u, u \rangle_{L^2} = -\nu \left| \partial_x u \right|_{L^2}^2.$$

By applying the same techniques as in Subsection 4.1 (take into account estimate (2.5)), we obtain

$$\mathrm{d}|u|_{L^2}^2 + 2\langle \mathcal{Q}u, u \rangle_{L^2} \,\mathrm{d}W_t + 2\nu \,|\partial_x u|_{L^2}^2 \,\mathrm{d}t \lesssim |u|_{L^2}^2 \,\mathrm{d}t,$$

and we have the bounds

$$\mathbb{E}\left[\sup_{t\in[0,T]}|u(t,\cdot)|^2_{L^2}\right] \le C_1(T),\tag{4.26}$$

$$\mathbb{E}\left[\int_0^T \left|\partial_x u(s,\cdot)\right|_{L^2}^2 \mathrm{d}s\right] \le C_2(T).$$
(4.27)

We compute the evolution of the \dot{H}^s -norm of u

$$\frac{1}{2} d|\Lambda^{s}u|_{L^{2}}^{2} + \langle\Lambda^{s}(\mathcal{Q}u), \Lambda^{s}u\rangle_{L^{2}} dW_{t}
= -\langle\Lambda^{s}(u\partial_{x}u), \Lambda^{s}u\rangle_{L^{2}} dt + \nu\langle\Lambda^{s}(\partial_{xx}u), \Lambda^{s}u\rangle_{L^{2}} dt
+ \frac{1}{2}\langle\Lambda^{s}(\mathcal{Q}u), \Lambda^{s}(\mathcal{Q}u)\rangle_{L^{2}} dt + \frac{1}{2}\langle\Lambda^{s}(\mathcal{Q}^{2}u), \Lambda^{s}u\rangle_{L^{2}} dt.$$
(4.28)

Integrating by parts the first term of the right-hand side above and observing that $u\partial_x u = (1/2)\partial_x(u^2)$, we obtain

$$-\langle \Lambda^s(u\partial_x u), \Lambda^s u \rangle_{L^2} = \frac{1}{2} \langle \Lambda^{s-1}(\partial_x(u^2)), \Lambda^{s+1} u \rangle_{L^2}.$$

Hence

$$\begin{aligned} |\langle \Lambda^{s}(u\partial_{x}u), \Lambda^{s}u \rangle_{L^{2}}| &\leq \left| \Lambda^{s-1}(\partial_{x}(u^{2})) \right|_{L^{2}} \left| \Lambda^{s+1}u \right|_{L^{2}} \\ &\lesssim \left(|u|_{L^{\infty}} |\Lambda^{s}u|_{L^{2}} \right) \left| \Lambda^{s+1}u \right|_{L^{2}} \leq \frac{1}{2\nu} \left| u \right|_{L^{\infty}}^{2} \left| \Lambda^{s}u \right|_{L^{2}}^{2} + \frac{\nu}{2} \left| \Lambda^{s+1}u \right|_{L^{2}}^{2}, \end{aligned}$$
(4.29)

where we have employed Kato-Ponce (see Lemma 2.4) in the second inequality. The second term on the right-hand side after the equality of equation (4.28) can be integrated as

$$\nu \langle \Lambda^{s}(\partial_{xx}u), \Lambda^{s}u \rangle_{L^{2}} = -\nu \langle \Lambda^{s+1}u, \Lambda^{s+1}u \rangle_{L^{2}} = -\nu \left| \Lambda^{s+1}u \right|_{L^{2}}^{2}.$$
(4.30)

As usual, applying inequality (2.6) we also have the estimate

$$|\langle \Lambda^{s}(\mathcal{Q}u), \Lambda^{s}(\mathcal{Q}u) \rangle_{L^{2}} + \langle \Lambda^{s}(\mathcal{Q}^{2}u), \Lambda^{s}u \rangle_{L^{2}}| \lesssim |\Lambda^{s}u|^{2}.$$

$$(4.31)$$

Putting together (4.29)-(4.31), one derives

$$\frac{\mathrm{d}|\Lambda^{s}u|_{L^{2}}^{2} + 2\langle\Lambda^{s}(\mathcal{Q}u),\Lambda^{s}u\rangle_{L^{2}}\,\mathrm{d}W_{t} + \nu\left|\Lambda^{s+1}u\right|_{L^{2}}^{2}\mathrm{d}t}{\lesssim \frac{1}{\nu}(1+|u|_{L^{\infty}}^{2})\left|\Lambda^{s}u\right|_{L^{2}}^{2}\mathrm{d}t}.$$

By applying expectation in the above equation and taking into account that the expectation of the Itô integral vanishes due to the martingale property, one obtains

$$\mathbb{E}\left[\sup_{t\in[0,T]}|\Lambda^{s}u(t)|_{L^{2}}^{2}\right] \lesssim \mathbb{E}\left[|\Lambda^{s}u(0)|_{L^{2}}^{2}\right] + \frac{1}{\nu}\mathbb{E}\left[\int_{0}^{T}(1+|u(\gamma)|_{L^{\infty}}^{2})\left|\Lambda^{s}u(\gamma)\right|_{L^{2}}^{2}\mathrm{d}\gamma\right], \quad (4.32)$$

and

$$\nu \mathbb{E} \left[\int_0^T \left| \Lambda^{s+1} u(\gamma) \right|_{L^2}^2 \mathrm{d}\gamma \right] \lesssim \mathbb{E} \left[\left| \Lambda^s u(0) \right|_{L^2}^2 \right] + \frac{1}{\nu} \mathbb{E} \left[\int_0^T (1 + |u(\gamma)|_{L^\infty}^2) \left| \Lambda^s u(\gamma) \right|_{L^2}^2 \mathrm{d}\gamma \right].$$
(4.33)

Now we claim that the following maximum principle holds

$$|u(t,\cdot)|_{L^{\infty}} \le |u(0,\cdot)|_{L^{\infty}}, \quad t \in [0,T],$$
(4.34)

which we show in Lemma 4.14. Once (4.34) is proven, we can infer from (4.32) and (4.33) together with Grönwall's inequality that there exist constants $A_1 = A_1(T)$, $A_2 = A_2(T)$, such that

$$\mathbb{E}\left[\sup_{t\in[0,T]}|\Lambda^{s}u(t,\cdot)|_{L^{2}}^{2}\right] \leq A_{1}, \quad \nu\mathbb{E}\left[\int_{0}^{T}\left|\Lambda^{s+1}u(s,\cdot)\right|_{L^{2}}^{2}\mathrm{d}s\right] \leq A_{2}.$$

This concludes the proof, so we are only left to show the maximum principle (4.34). \Box Lemma 4.14. The maximum principle (4.34) is satisfied.

Proof. Following the type of argument in [BFMM19], consider the SDE

$$\mathrm{d}X_t = \xi(X_t) \circ \mathrm{d}W_t,\tag{4.35}$$

and let $\psi_t(X_0)$ be the corresponding flow of (4.35) with initial condition $X_0 \in \mathbb{T}$, which is a semimartingale and a smooth diffeomorphism. By applying Itô-Wentzell in Stratonovich form (see Theorem 2.5), we evaluate u along $X_t = \psi_t(X_0)$, getting that

$$u(t, X_t) = u(0, X_0) - \int_0^t u(s, X_s) \partial_x u(s, X_s) \,\mathrm{d}s + \nu \int_0^t \partial_{xx} u(s, X_s) \,\mathrm{d}s$$
$$- \int_0^t \xi(X_s) \partial_x u(s, X_s) \circ \mathrm{d}W_s + \int_0^t \partial_x u(s, X_s) \circ \mathrm{d}X_s, \tag{4.36}$$

a.s., where the term in the last line

$$\int_0^t \xi(X_s) \partial_x u(s, X_s) \circ \mathrm{d}W_s + \int_0^t \partial_x u(s, X_s) \circ \mathrm{d}X_s = 0$$

Consider the change of variables $w(t, X_0) = u(t, \psi_t(X_0))$ and by chain rule, obtain

$$\partial_x u(t, X_t) = \frac{\partial \psi_t^{-1}}{\partial x} (\psi_t(X_0)) \frac{\partial w}{\partial X_0} (t, X_0),$$

$$\partial_{xx} u(t, X_t) = \frac{\partial^2 \psi_t^{-1}}{\partial x^2} (\psi_t(X_0)) \frac{\partial w}{\partial X_0} (t, X_0) + \left(\frac{\partial \psi_t^{-1}}{\partial x} (\psi_t(X_0))\right)^2 \frac{\partial^2 w}{\partial X_0^2} (t, X_0).$$

Therefore, (4.36) is equivalent to the following PDE with random coefficients

$$\frac{\partial w}{\partial t} + \tilde{w} \frac{\partial w}{\partial X_0} = \tilde{\nu} \frac{\partial^2 w}{\partial X_0^2},$$

where

$$\tilde{w} = \frac{\partial \psi_t^{-1}}{\partial x} (\psi_t(X_0)) u(t, \psi_t(X_0)) - \nu \frac{\partial^2 \psi_t^{-1}}{\partial x^2} (\psi_t(X_0)),$$
$$\tilde{\nu} = \nu \left(\frac{\partial \psi_t^{-1}}{\partial x} (\psi_t(X_0)) \right)^2.$$

We claim that w satisfies the maximum principle

$$|w_t|_{L^{\infty}} \le |w_0|_{L^{\infty}}, \quad t > 0.$$
(4.37)

To prove (4.37), we follow the strategy in [CC04]. If $w_0 = 0$, the claim is evident. Otherwise, for every $t \ge 0$, we define $X_t^0 \in \mathbb{T}$ to be such that $|w_t|_{L^{\infty}} = |w(t, X_t^0)|$ (i.e., the spatial point where $|w_t|_{L^{\infty}}$ is attained. Its existence is guaranteed because \mathbb{T} is compact). We start by claiming that for all t > 0, $|w_t|_{L^{\infty}}$ is differentiable in time and

$$\frac{\mathrm{d}}{\mathrm{d}t} |w_t|_{L^{\infty}} = \frac{\partial w}{\partial t}(t, X_t^0). \tag{4.38}$$

First, we check that $|w_t|_{L^{\infty}}$ is differentiable in time. For this, note that for $h \ge 0$ from the mean value theorem we have

$$||w_{t+h}|_{L^{\infty}} - |w_t|_{L^{\infty}}| \le |w_{t+h} - w_t|_{L^{\infty}} \le \max_{s \in [t,t+h]} |\partial_t w(s)|h,$$

and a similar argument can be applied if $h \leq 0$. Therefore $|w_t|_{L^{\infty}}$ is Lipschitz in time and from Rademacher's theorem a.e. differentiable.

We proceed to prove equality (4.38). Without loss of generality, we assume $w(t, X_t^0) \ge 0$, t > 0. We note that by taking small enough h > 0

$$\frac{\mathrm{d}}{\mathrm{d}t} |w_t|_{L^{\infty}} = \lim_{h \to 0} \frac{|w_{t+h}|_{L^{\infty}} - |w_t|_{L^{\infty}}}{h} = \lim_{h \to 0} \frac{w(t+h, X_{t+h}^0) - w(t, X_t^0)}{h}$$
$$= \lim_{h \to 0} \frac{w(t+h, X_{t+h}^0) - w(t+h, X_t^0)}{h} + \lim_{h \to 0} \frac{w(t+h, X_t^0) - w(t, X_t^0)}{h}$$
$$\geq \frac{\partial w}{\partial t}(t, X_t^0),$$

for t > 0. Symmetrically, by choosing small enough h > 0 we also have

$$\frac{\mathrm{d}}{\mathrm{d}t} |w_t|_{L^{\infty}} = \lim_{h \to 0} \frac{|w_{t-h}|_{L^{\infty}} - |w_t|_{L^{\infty}}}{-h} = \lim_{h \to 0} \frac{w(t-h, X_{t-h}^0) - w(t, X_t^0)}{-h}$$
$$= \lim_{h \to 0} \frac{w(t-h, X_{t-h}^0) - w(t-h, X_t^0)}{-h} + \lim_{h \to 0} \frac{w(t-h, X_t^0) - w(t, X_t^0)}{-h}$$
$$\leq \frac{\partial w}{\partial t}(t, X_t^0),$$

for t > 0. Therefore we conclude (4.38). To finish our argument, we observe that

$$\frac{\partial w}{\partial t}(t, X_t^0) = -\tilde{w}\frac{\partial w}{\partial X_0}(t, X_t^0) + \tilde{\nu}\frac{\partial^2 w}{\partial X_0^2}(t, X_t^0) \le 0.$$

Hence (4.37) is satisfied. Since ψ_t is a diffeomorphism, we have $|u_t|_{L^{\infty}} = |w_t|_{L^{\infty}}$ so the maximum principle also follows for u.

With this, we conclude the proof of Theorem 4.12.

5. Conclusion and Outlook

In this paper, we studied the solution properties of a stochastic Burgers' equation on the torus and the real line, with the noise appearing in the transport velocity. We have shown that this stochastic Burgers' equation is locally well-posed in $H^s(\mathbb{T}, \mathbb{R})$, for s > 2, and furthermore, established a blow-up criterion which extends the deterministic one to the stochastic case. We also proved that if the noise is of the form $\xi(x)\partial_x u \circ dW_t$ where $\xi(x) = \alpha x + \beta$, then shocks form almost surely from a negative slope. Moreover, for a more general type of noise, we showed that blow-up occurs in expectation, which follows from the previously mentioned stochastic blow-up criterion. Also, in the weak formulation of the problem, we provided a Rankine-Hugoniot type condition that is satisfied by the shocks, analogous to the deterministic shocks. Finally, we also studied the stochastic Burgers' equation with a viscous term, which we proved to be globally well-posed in H^s for s > 2.

Let us conclude by proposing some future research directions and open problems that have emerged during the course of this work:

- Regarding shock formation, it is natural to ask whether our results can be extended to show that shock formation occurs almost surely for more general types of noise.
- Another possible question is whether our global well-posedness result can be extended for the viscous Burgers' equation with the Laplacian replaced by a fractional Laplacian $(-\Delta)^{\alpha}$, $\alpha \in (0, 1)$. The main difficulty here is that in the stochastic case, the proof of the maximum principle (Proposition (4.14)) does not follow immediately since the pointwise chain rule for the fractional Laplacian is not available. In the deterministic case, this question has been settled and it is known that the solution exhibits a very different behaviour depending on the value of α : for $\alpha \in [1/2, 1]$, the solution is global in time, and for $\alpha \in [0, 1/2)$, the solution develops singularities in finite time [KNS08, Kis10]. Interestingly, when an Itô noise of type $\beta u \, dW_t$ is added, it is shown in [RZZ14] that the probability of solutions blowing up for small initial conditions tends to zero when $\beta > 0$ is sufficiently large. It would be interesting to investigate whether the transport noise considered in this paper can also have a similar regularising effect on the equation.
- Similar results could be derived for other one-dimensional equations with non-local transport velocity [CCF05, DG90, DG96]. For instance, the so called *CCF model* [CCF05] is also known to develop singularities in finite time, although by a different mechanism to that of Burgers'. To our knowledge, investigating these types of equations with transport noise is new.

Acknowledgements. The authors would like to thank José Antonio Carillo de la Plata, Dan Crisan, Theodore Drivas, Franco Flandoli, Darryl Holm, James-Michael Leahy, Erwin Luesink, and Wei Pan for encouraging comments and discussions that helped put together this work. DAO has been partially supported by the grant MTM2017-83496-P from the Spanish Ministry of Economy and Competitiveness, and through the Severo Ochoa Programme for Centres of Excellence in R&D (SEV-2015-0554). ABdL has been supported by the Mathematics of Planet Earth Centre of Doctoral Training (MPE CDT). ST acknowledges the Schrödinger scholarship scheme for funding during this work.

References

- [AF09] S. Attanasio and F. Flandoli. Zero-noise solutions of linear transport equations without uniqueness: an example. *Comptes Rendus Mathematique*, 347(13-14):753–756, 2009.
- [AOB20] D. Alonso-Orán and A. Bethencourt. On the Well-posedness of stochastic Boussinesq equations with transport noise. Journal of Nonlinear Science, 30(1):175–224, 2020.
- [AOHR21] D. Alonso-Orán, T. Hao, and C. Rohde. A local-in-time theory for singular SDEs with applications to fluid models with transport noise. *Journal of Nonlinear Science*, 31:31–98, 2021.
- [BCJL94] L. Bertini, N. Cancrini, and G. Jona-Lasinio. The stochastic Burgers equation. Communications in Mathematical Physics 165, 211-231, 1994.
- [BFMM19] L. Beck, F. Flandoli, Gubinelli M., and Maurelli M. Stochastic ODEs and stochastic linear PDEs with critical drift: regularity, duality and uniqueness. *Electronic Journal of Probability*, 24:1–72, 2019.
- [CC04] A. Córdoba and D. Córdoba. A Maximum Principle Applied to Quasi-Geostrophic Equations. Communications in Mathematical Physics, 249(249):511–528, 2004.
- [CCF05] A. Córdoba, D. Córdoba, and M. Fontelos. Formation of singularities for a transport equation with nonlocal velocity. *Annals of Mathematics*, 162, 1377-1389, 2005.
- [CCH⁺18] C. Cotter, D. Crisan, D. D. Holm, W. Pan, and I. Shevchenko. Modelling uncertainty using circulation-preserving stochastic transport noise in a 2-layer quasi-geostrophic model. arXiv preprint arXiv:1802.05711, 2018.
- [CCH⁺19] C. J. Cotter, D. Crisan, D. D. Holm, W. Pan, and I. Shevchenko. Numerically modelling stochastic Lie transport in fluid dynamics. *Multiscale Modelling & Simulation*, 17(1):192– 232, 2019.
- [CFH19] D. Crisan, F. Flandoli, and Darryl D. Holm. Solution Properties of a 3D Stochastic Euler Fluid Equation. Journal of Nonlinear Science, 29(3):813–870, 2019.
- [CGH17] C. J. Cotter, G. A. Gottwald, and D. D. Holm. Stochastic partial differential fluid equations as a diffusive limit of deterministic Lagrangian multi-time dynamics. Proc. R. Soc. A, 473(2205):20170388, 2017.
- [CH18] D. Crisan and D. D. Holm. Wave breaking for the stochastic Camassa–Holm equation. Physica D: Nonlinear Phenomena, 376:138–143, 2018.
- [CO13] P. Catuogno and C. Olivera. Strong solution of the stochastic Burgers equation. Applicable Analysis: An International Journal, 93:3, 646-652, DOI: 10.1080/00036811.2013.797074, 2013.
- [DFV14] F. Delarue, F. Flandoli, and D. Vincenzi. Noise prevents collapse of Vlasov-Poisson point charges. *Communications on Pure and Applied Mathematics*, 67(10):1700–1736, 2014.
- [DG90] S. De Gregorio. On a one-dimensional model for the three-dimensional vorticity equation,. J. Statist. Phys. 59, 1251-1263, 1990.
- [DG96] S. De Gregorio. A partial differential equation arising in a 1D model for the 3D vorticity equa- tion. *Math. Methods Appl. Sci.* 19, 1233-1255., 1996.
- [DPDT94] G. Da Prato, A. Debussche, and R. Temam. Stochastic Burgers' equation. Nonlinear Differential Equations and Applications 1, 389-402, 1994.
- [DPG07] G. Da Prato and D. Gatarek. Stochastic Burgers equation with correlated noise. *Stochastics and Stochastic Reports*, 52:1-2, 29-41, 2007.

- [FF13] E. Fedrizzi and F. Flandoli. Noise prevents singularities in linear transport equations. *Journal of functional analysis*, 2013.
- [FG16] P. K. Friz and B. Gess. Stochastic scalar conservation laws driven by rough paths. volume 33, pages 933–963. Elsevier, 2016.
- [FGH20] T. Funaki, Y. Gao, and D. Hilhorst. Uniqueness of the entropy solution of a stochastic conservation law with a Q-brownian motion. *Math Meth Appl Sci*, 43:5860–5886, 2020.
- [FGP10] F. Flandoli, M. Gubinelli, and E. Priola. Well-posedness of the transport equation by stochastic perturbation. *Inventiones mathematicae*, 2010.
- [FGP11] F. Flandoli, M. Gubinelli, and E. Priola. Full well-posedness of point vortex dynamics corresponding to stochastic 2D Euler equations. *Stochastic Processes and their Applications*, 2011.
- [FL18] F. Flandoli and D. Luo. Euler-Lagrangian approach to 3D stochastic Euler equations. arXiv:1803.05319, 2018.
- [Fla11] F. Flandoli. Random Perturbation of PDEs and Fluid Dynamic Models: École d'été de Probabilités de Saint-Flour XL-2010, volume 2015. Springer Science & Business Media, 2011.
- [GHV14] N. Glatt-Holtz and V. Vicol. Local and global existence of smooth solutions for the stochastic Euler equation with multiplicative noise,. The Annals of Probability, Vol.42, 80-145, DOI:10.1214/12-AOP773, 2014.
- [GM18] B. Gess and M. Maurelli. Well-posedness by noise for scalar conservation laws. Communications in Partial Differential Equations, 43(12):1702–1736, 2018.
- [Gol85] J. A. Goldstein. Semigroups of linear operators and applications. Courier Dover Publications, 1985.
- [GS17] B. Gess and P. E. Souganidis. Stochastic non-isotropic degenerate parabolic–hyperbolic equations. *Stochastic Processes and their Applications*, 127(9):2961–3004, 2017.
- [Har93] J. Harrison. Stokes' theorem for nonsmooth chains. Bulletin of the American Mathematical Society, 29(2):235–242, 1993.
- [Har99] J. Harrison. Flux across nonsmooth boundaries and fractal Gauss/Green/Stokes' theorems. Journal of Physics A, 1999.
- [HN92] J. Harrison and A. Norton. The Gauss-Green theorem for fractal boundaries. *Duke Math. J*, 67(3):575–588, 1992.
- [HNS19] A. Hocquet, T. Nilssen, and W. Stannat. Generalized Burgers' equation with rough transport noise. *Stochastic Processes and their Applications*, 2019.
- [Hol15] D. D. Holm. Variational principles for stochastic fluid dynamics. Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences, 471(2176), 2015.
- [Kis10] A. Kiselev. Regularity and blow up for active scalars. Math. Model. Nat. Phenom. Vol. 5, No. 4, 2010, pp. 225-255, 2010.
- [KNS08] A. Kiselev, F. Nazarov, and R. Shterenberg. Blow up and regularity for fractal Burgers equation. *Dynamics of PDE*, Vol.5, No.3, 211-240, 2008.
- [KP88] T. Kato and G. Ponce. Commutator estimates and the Euler and Navier-Stokes equations. Communications in Pure and Applied Mathematics 41(7), 891–907, 1988.
- [KR81] N. V. Krylov and B. L. Rozovskii. Stochastic evolution equations. Journal of Mathematical Sciences, 16(249):1233–1277, 1981.
- [Kun81] H. Kunita. Some extensions of Itô's formula. Séminaire de probabilités, 1981.
- [LY06] T. J. Lyons and P. SC Yam. On Gauss-Green theorem and boundaries of a class of Hölder domains. Journal de mathématiques pures et appliquées, 85(1):38-53, 2006.
- [RZZ14] M. Röckner, R. Zhu, and X. Zhu. Local existence and non-explosion of solutions for stochastic fractional partial differential equations driven by multiplicative noise. *Stochastic Processes* and their Applications, 124(5):1974–2002, 2014.
- [Sha57] V. L. Shapiro. The divergence theorem without differentiability conditions. Proceedings of the National Academy of Sciences of the United States of America, 1957.
- [Sim86] J. Simon. Compact sets in the space $L^p(0,T;B)$. Annali di Matematica Pura ed Applicata, 146:65–96, 1986.
- [Ver81] A. J. Veretennikov. On strong solutions and explicit formulas for solutions of stochastic integral equations. Sbornik: Mathematics, 39(3):387–403, 1981.

DA: INSTITUTO DE CIENCIAS MATEMÁTICAS CSIC-UAM-UC3M-UCM, 28049 MADRID, SPAIN. Email address: DA: diego.alonso@icmat.es

AB, ST: DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE, LONDON SW7 2AZ, UK. Email address: AB: ab1113@ic.ac.uk ST:st4312@ic.ac.uk