

# Modeling High-Dimensional Time Series: A Factor Model with Dynamically Dependent Factors and Diverging Eigenvalues

Zhaoxing Gao<sup>1</sup> and Ruey S. Tsay<sup>2</sup>

<sup>1</sup>Department of Mathematics, Lehigh University

<sup>2</sup>Booth School of Business, University of Chicago

July 2, 2020

## Abstract

This article proposes a new approach to modeling high-dimensional time series by treating a  $p$ -dimensional time series as a nonsingular linear transformation of certain common factors and idiosyncratic components. Unlike the approximate factor models, we assume that the factors capture all the non-trivial dynamics of the data, but the cross-sectional dependence may be explained by both the factors and the idiosyncratic components. Under the proposed model, (a) the factor process is dynamically dependent and the idiosyncratic component is a white noise process, and (b) the largest eigenvalues of the covariance matrix of the idiosyncratic components may diverge to infinity as the dimension  $p$  increases. We propose a white noise testing procedure for high-dimensional time series to determine the number of white noise components and, hence, the number of common factors, and introduce a projected Principal Component Analysis (PCA) to eliminate the diverging effect of the idiosyncratic noises. Asymptotic properties of the proposed method are established for both fixed  $p$  and diverging  $p$  as the sample size  $n$  increases to infinity. We use both simulated data and real examples to assess the performance of the proposed method. We also compare our method with two commonly used methods in the literature concerning the forecastability of the extracted factors and find that the proposed approach not only provides interpretable results, but also performs well in out-of-sample forecasting. Supplementary materials of the article are available online.

*Keywords:* Diverging eigenvalues, Eigen-analysis, Factor model, High dimension, Projected principal component analysis, White noise test.

# 1 Introduction

Advances in information technology make large data sets widely accessible nowadays. In many applications, the data consist naturally of high-dimensional time series. For example, the returns of a large number of assets form a high-dimensional time series and play an important role in asset pricing, portfolio allocation, and risk management. Large panel time series data are commonplace in economics and biological studies. Environmental studies often employ high-dimensional time series consisting of a large number of pollution indexes collected at many monitoring stations and over periods of time. However, modeling high-dimensional time series is always challenging because the commonly used Vector Autoregressive (VAR) or Vector-Autoregressive Moving-Average (VARMA) models are not practically applicable when the dimension is high. In particular, unregularized VARMA models often suffer the difficulties of over-parameterization and lack of identifiability as discussed in Tiao and Tsay (1989), Lütkepohl (2006), and Tsay (2014). Therefore, dimension reduction or structural specification becomes a necessity in applications of high-dimensional time series. Indeed, various methods have been developed in the literature for multivariate time series analysis, including the scalar component models of Tiao and Tsay (1989), the LASSO regularization of VAR models by Shojaie and Michailidis (2010) and Song and Bickel (2011), the sparse VAR model based on partial spectral coherence by Davis et al. (2012), and the factor modeling in Stock and Watson (2005), Bai and Ng (2002), Forni et al. (2005), Lam et al. (2011), and Lam and Yao (2012), among others. However, the complexity of the dynamical dependence in high-dimensional time series requires further investigation, as extracting dynamic information from the data plays an important role in modeling and forecasting serially dependent data.

This article marks a further development in factor modeling of high-dimensional time series. Factor models are commonly used in finance, economics, and statistics. For example, asset returns are often modeled as functions of a small number of factors, see Stock and Watson (1989) and Stock and Watson (1998). Macroeconomic variables of multiple countries are often found to have common movements, see Gregory and Head (1999) and Forni et al. (2000a). From the statistical perspective, a modeling approach that can reveal the common structure of the series and provide accurate estimation of a specified model is highly valuable in understanding the dynamic relationships of the data. To this end, we first briefly introduce the traditional approximate factor models and some existing estimation procedures under different assumptions in the literature. Let  $\mathbf{y}_t = (y_{1t}, \dots, y_{pt})'$  be a  $p$ -dimensional zero-mean time series. The approximate factor model for  $\mathbf{y}_t$  assumes the form

$$\mathbf{y}_t = \mathbf{A}\mathbf{x}_t + \boldsymbol{\varepsilon}_t, \tag{1.1}$$

where  $\mathbf{x}_t$  is a  $r$ -dimensional latent factor process,  $\mathbf{A} \in R^{p \times r}$  is an associated (full-rank) factor loading matrix,  $\boldsymbol{\varepsilon}_t$  is the idiosyncratic component, and  $\mathbf{x}_t$  and  $\boldsymbol{\varepsilon}_t$  are independent. To the best of our knowledge, there are at least three main statistical procedures to estimate the common factors and the associated loading matrix under various assumptions on the factor and idiosyncratic terms. The first procedure is based on the principal component analysis (PCA), see Bai and Ng (2002), Bai (2003), Fan et al. (2013) with pervasive factors, and Onatski (2012) with weakly influential factors, and the references therein. The second procedure is based on the eigen-analysis of the auto-covariance matrices, see Lam et al. (2011) and Lam and Yao (2012), among others. These authors assume that all components of  $\mathbf{x}_t$  are dynamically dependent, reflecting the nature of time series data, and  $\boldsymbol{\varepsilon}_t$  is a vector white noise process, which has no serial correlations. Although the model considered in Lam et al. (2011) and Lam and Yao (2012) forms a subclass of those in Bai and Ng (2002), the recovered factors are different from those in the latter because they are all dynamically dependent. The third approach is slightly different from the previous two and allows  $\mathbf{y}_t$  to depend also on factors involving lagged variables, which can also be written as Equation (1.1) with  $\mathbf{x}_t$  consisting of all dynamic factors. See Forni et al. (2000b), Forni et al. (2005), Forni et al. (2015), and the references therein. This third approach is based on the eigenvalues and principal components of spectral density matrices and, hence, is a frequency-domain analysis. But some recent studies extend the approach to time-domain analysis; see, for instance, Hallin and Lippi (2013) and Peña and Yohai (2016). For more details on the differences between the first and the third approaches, see Fan et al. (2013).

In this paper, we study the common dynamic components in high-dimensional, and possibly highly cross-sectionally correlated, time series data. We propose a new factor model under which the observed high-dimensional time series  $\mathbf{y}_t$  is a nonsingular linear transformation of a  $r$ -dimensional common factor process, which is dynamically dependent, and a  $(p - r)$ -dimensional idiosyncratic component, which is a white noise series. In other words, we assume the factors capture all the non-trivial dynamics of the data, but the cross-sectional dependence may be explained by both the factors and the idiosyncratic components. This is different from the approximate factor model, where factors capture most of the cross-sectional dependence, while the idiosyncratic terms may contain some non-trivial temporal dependence. To a large degree and in extracting the common dynamic components of the data, our approach is closer to the dynamic factor models of Forni et al. (2000a,b, 2005), even though its form is static according to, for example, Boivin and Ng (2005). In fact, the proposed new factor model is in line with that of Tiao and Tsay (1989) and Gao and Tsay (2019), and assumes that the idiosyncratic component is white noise in the sense that  $\mathbf{y}_t$  has  $(p - r)$  scalar components

of order  $(0, 0)$ . See Tiao and Tsay (1989) and Section 2 for details. However, our proposed modeling approach is different from those of the aforementioned two papers, because Tiao and Tsay (1989) assumes  $p$  is fixed and Gao and Tsay (2019) considers  $p = o(n^{1/2})$ , where  $n$  is the sample size, so that they can employ the canonical correlation analysis (CCA). In this paper, we do not employ CCA and, hence, can relax the constraints on  $p$  as  $n$  increases.

Similar to that of Lam et al. (2011) and Lam and Yao (2012), we first apply the eigen-analysis on certain auto-covariance matrices to obtain the loading matrix associated with the common factors. But we propose a projected PCA method to estimate the loading matrix associated with the idiosyncratic component in the presence of diverging noise effect. Our estimator is constructed by first projecting  $\mathbf{y}_t$  onto the directions which capture little dynamic dependencies, and the resulting projected coordinates would essentially be a vector white noise. We then perform the PCA between the original data  $\mathbf{y}_t$  and the projected white noise coordinates, and the eigenspace associated with the small eigenvalues would be able to mitigate the diverging effect of the white noise idiosyncratic components; see Section 2 for details. In addition, we propose a new method to estimate the common factors so that the resulting estimated common factors are not affected by the idiosyncratic component  $\boldsymbol{\varepsilon}_t$ . Specifically, in the presence of diverging noise components, we project the observed data into the orthogonal direction of those diverging noise components to mitigate the effect of the idiosyncratic component in estimating the common factors. Furthermore, to overcome the difficulties associated with the behavior of eigenvalues of a large random matrix, we consider a white noise testing procedure to determine the number of common factors. This testing procedure is found to be more reliable than the the ratio-based method currently used in the literature to extract the dynamically dependent factors. Details of the testing procedure are given in Section 2.3.

The idea of using white noise test to determine the number of factors has been used in Pan and Yao (2008) when the dimension is relatively small. However, their approach needs to solve a constrained optimization problem step-by-step and it cannot be extended to the high-dimensional case directly. When the factors in (1.1) have different levels of strength, Lam and Yao (2012) proposed a two-step estimation procedure to successively identify two groups of factors with top two strengths. Under an independent structure of the noises, Li et al. (2017) provided an exact description of the phase transition phenomenon whether a factor is strong enough to be detected with the observed sample singular values and proposed a new ratio-based estimator for determining the number of factors, which is shown to be more robust against possibly multiple levels of factor strengths. However, they all rely on relatively pervasive factors in the sense that their strength is (much) stronger than that of the idiosyncratic components, which is different from the setting considered in this paper. The proposed approach still works

when the factors have multiple levels of factor strengths and the theory can readily be modified accordingly. For simplicity, we only consider factors with the same strength in this paper.

We conduct simulation studies to assess the performance of the proposed modeling procedure in finite samples and to compare it with the method in Lam et al. (2011). When the largest eigenvalues of the covariance matrix of the idiosyncratic component are diverging, the results show that the method of Lam et al. (2011) may encounter estimation errors. We further apply the proposed method to two real examples, and the numerical results suggest that the factors recovered by our approach not only have reasonable interpretations but also fare well in prediction. On the other hand, even under the setting of Lam et al. (2011), the ratio-based method could fail if the covariance of the noises is large. See the simulation results in the supplementary materials. Furthermore, as the factors play an important role in diffusion index models as in Stock and Watson (2002a) and Stock and Watson (2002b), for the purpose of out-of-sample forecasting, we also compared the predictive ability of the factors extracted by different methods in an empirical example. The results suggest that the dynamically dependent factors extracted by our method fare better in predicting the U.S. Consumer Price Indexes.

The contributions of this paper are multi-fold. First, the proposed model is flexible in application. It allows a variety of structures for the common factors and the idiosyncratic components. Second, the proposed estimation method can eliminate the effect of the idiosyncratic term in estimating the common factors. This is achieved by using the projected PCA method if the dimension  $p$  is low. When the dimension is high, we assume that a few largest eigenvalues of the covariance matrix of the idiosyncratic noise term are diverging, which is a reasonable assumption in the high-dimensional setting and the diverging eigenvalues can explain some of the cross-sectional dependence. The projected PCA then helps to mitigate the effect of the diverging part of the noise covariance matrix. Third, we propose a procedure based on a white noise test for multivariate (and high-dimensional) time series to determine the number of common factors  $r$ . Under the assumption that the idiosyncratic term is a vector white noise, the limiting distribution of the test statistic used is available in closed form. This testing procedure is shown to be more reliable than the ratio-based method available in the literature in the sense that our approach can extract the dynamically dependent factors more accurately.

The rest of the paper is organized as follows. We introduce the proposed model, estimation methodology, and modeling procedure in Section 2, including a flowchart for the entire modeling procedure. In Section 3, we study the theoretical properties of the proposed model and its associated estimates. Section 4 illustrates the performance of the proposed model using both simulated and real data sets. Section 5 provides some discussions and concluding remarks.

All technical proofs and an additional real example are relegated to an online supplement. Throughout the article, we use the following notation.  $\|\mathbf{u}\|_2 = (\sum_{i=1}^p u_i^2)^{1/2}$  is the Euclidean norm of a  $p$ -dimensional vector  $\mathbf{u} = (u_1, \dots, u_p)'$ ,  $\|\mathbf{u}\|_\infty = \max_i |u_i|$ , and  $\mathbf{I}_k$  denotes the  $k \times k$  identity matrix. For a matrix  $\mathbf{H} = (h_{ij})$ ,  $\|\mathbf{H}\|_2 = \sqrt{\lambda_{\max}(\mathbf{H}'\mathbf{H})}$  is the operator norm, where  $\lambda_{\max}(\cdot)$  denotes for the largest eigenvalue of a matrix, and  $\|\mathbf{H}\|_{\min}$  is the square root of the minimum non-zero eigenvalue of  $\mathbf{H}'\mathbf{H}$ . The superscript  $'$  denotes the transpose of a vector or matrix. Finally, we use the notation  $a \asymp b$  to denote  $a = O(b)$  and  $b = O(a)$ .

## 2 The Model and Methodology

In this section, we state the proposed model, discuss the estimation methodology, and an estimation procedure when the number of common factors is known. This is followed by the proposed method for determining the number of common factors. We also provide a flowchat to summarize the proposed modeling procedure.

### 2.1 Setting

Let  $\mathbf{y}_t = (y_{1t}, \dots, y_{pt})'$  be an observable  $p$ -dimensional time series. We assume  $E(\mathbf{y}_t) = \mathbf{0}$  and  $\mathbf{y}_t$  admits a latent structure:

$$\mathbf{y}_t = \mathbf{L} \begin{bmatrix} \mathbf{f}_t \\ \boldsymbol{\varepsilon}_t \end{bmatrix} = [\mathbf{L}_1, \mathbf{L}_2] \begin{bmatrix} \mathbf{f}_t \\ \boldsymbol{\varepsilon}_t \end{bmatrix} = \mathbf{L}_1 \mathbf{f}_t + \mathbf{L}_2 \boldsymbol{\varepsilon}_t, \quad (2.1)$$

where  $\mathbf{L} \in R^{p \times p}$  is a full rank loading matrix,  $\mathbf{f}_t = (f_{1t}, \dots, f_{rt})'$  is a  $r$ -dimensional dynamically dependent factor process,  $\boldsymbol{\varepsilon}_t = (\varepsilon_{1t}, \dots, \varepsilon_{vt})'$  is a  $v$ -dimensional white noise vector, and  $r + v = p$ . For meaningful dimension reduction, we assume  $r$  is a small fixed nonnegative integer. In addition, we also assume  $\text{Cov}(\mathbf{f}_t) = \mathbf{I}_r$ ,  $\text{Cov}(\boldsymbol{\varepsilon}_t) = \mathbf{I}_v$ ,  $\text{Cov}(\mathbf{f}_t, \boldsymbol{\varepsilon}_t) = \mathbf{0}$ , and no linear combination of  $\mathbf{f}_t$  is serially uncorrelated. The last assumption is trivial, because we can reduce the dimension of  $\mathbf{f}_t$  if any such a linear combination exists. The assumptions on the covariance matrices of  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_t$  are common among factor models. The decomposition of Model (2.1) is general in the sense that any finite-order VARMA time series  $\mathbf{y}_t$  can always be written in Equation (2.1) via canonical correlation analysis between constructed vectors involving  $\mathbf{y}_t$  and its lagged values. See Tiao and Tsay (1989). Readers are referred to Anderson (2003) for details of canonical correlation analysis.

If  $v = 0$ , then no linear combination of  $\mathbf{y}_t$  is white noise, indicating no dimension reduction.

Therefore, we focus on the case of  $v > 0$  in this paper. Under Equation (2.1), the dynamic dependence of  $\mathbf{y}_t$  is driven by  $\mathbf{f}_t$  if  $r > 0$ . Thus,  $\mathbf{f}_t$  indeed consists of the common dynamically dependent factors of  $\mathbf{y}_t$ . In the terminology of Tiao and Tsay (1989), (a)  $\boldsymbol{\varepsilon}_t$  is a  $v$ -dimensional scalar component process of order (0,0) if  $v > 0$ , that is, there exists a transformation matrix  $\mathbf{V}_2 \in R^{p \times v}$  such that  $\boldsymbol{\varepsilon}_t = \mathbf{V}_2' \mathbf{y}_t$  and  $\text{Cov}(\boldsymbol{\varepsilon}_t, \mathbf{y}_{t-j}) = \mathbf{0}$  for  $j > 0$ , and (b) no linear combination of  $\mathbf{f}_t$  is a scalar component of order (0,0) if  $r > 0$ . Readers are referred to Tiao and Tsay (1989) for a formal definition of a scalar component of order (0,0). Condition (a) is equivalent to  $\boldsymbol{\varepsilon}_t$  being a white noise under the traditional factor models, where  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_t$  are assumed to be independent.

Assuming that the time series  $\mathbf{y}_t$  follows a structural model consisting of trend, seasonal component, and irregular series, Gao and Tsay (2019) employ Equation (2.1) to model the irregular series. These authors use CCA to determine the number of common factors. However, the method of CCA only works when  $p = o(n^{1/2})$ , where  $n$  is the sample size. This restricts the applicability of the modeling procedure of Gao and Tsay (2019). We relax such restrictions on  $p$  in this paper.

To study Model (2.1) in a more general setting and to provide sufficient statistical insights on the proposed factor models, we decompose  $\mathbf{L}_1$  and  $\mathbf{L}_2$  of Model (2.1) as  $\mathbf{L}_1 = \mathbf{A}_1 \mathbf{Q}_1$  and  $\mathbf{L}_2 = \mathbf{A}_2 \mathbf{Q}_2$ , respectively, where  $\mathbf{A}_1$  and  $\mathbf{A}_2$  are two half orthonormal matrices, i.e.,  $\mathbf{A}_1' \mathbf{A}_1 = \mathbf{I}_r$  and  $\mathbf{A}_2' \mathbf{A}_2 = \mathbf{I}_v$ . This can be done via the QR decomposition or singular value decomposition, and hence most of the strengths of  $\mathbf{L}_1$  and  $\mathbf{L}_2$  are retained in  $\mathbf{Q}_1$  and  $\mathbf{Q}_2$ , respectively. Furthermore, let  $\mathbf{x}_t = \mathbf{Q}_1 \mathbf{f}_t$  and  $\mathbf{e}_t = \mathbf{Q}_2 \boldsymbol{\varepsilon}_t$ , then Model (2.1) can be written as

$$\mathbf{y}_t = \mathbf{A}_1 \mathbf{x}_t + \mathbf{A}_2 \mathbf{e}_t, \quad (2.2)$$

which is close to the traditional factor model in Equation (1.1). Some remarks are in order. First, the model we are investigating is still the one in (2.1) and we rewrite it in the form of (2.2) to provide more insights to the proposed methods below. Second, even though  $\mathbf{L}$  is of full rank,  $\mathbf{A}_1$  is not orthogonal to  $\mathbf{A}_2$  in general because we performed the decomposition separately. Third,  $\mathbf{A}_1$  and  $\mathbf{x}_t$  are still not uniquely identified because we can replace  $(\mathbf{A}_1, \mathbf{x}_t)$  by  $(\mathbf{A}_1 \mathbf{H}, \mathbf{H}' \mathbf{x}_t)$  for any orthonormal matrix  $\mathbf{H} \in R^{r \times r}$ , where  $\mathbf{A}_1$  and  $\mathbf{A}_1 \mathbf{H}$  are both half orthonormal matrices, and it is also due to that the decomposition of  $\mathbf{L}_1$  is not unique. The same issue applies to  $\mathbf{A}_2$  and  $\mathbf{e}_t$ . Nevertheless, the linear space spanned by the columns of  $\mathbf{A}_1$ , denoted by  $\mathcal{M}(\mathbf{A}_1)$ , is uniquely defined, and is equal to that of  $\mathbf{L}_1$ .  $\mathcal{M}(\mathbf{A}_1)$  is called the factor loading space, and the linear space  $\mathcal{M}(\mathbf{A}_2)$  can be defined similarly for the idiosyncratic component.

## 2.2 Estimation Methodology

To begin, we provide some rationale for the proposed estimation method. Let  $\mathbf{B}_1$  and  $\mathbf{B}_2$  be the orthonormal complement of  $\mathbf{A}_1$  and  $\mathbf{A}_2$ , respectively, i.e.,  $\mathbf{B}_1 \in R^{p \times v}$  and  $\mathbf{B}_2 \in R^{p \times r}$  are two half orthonormal matrices satisfying  $\mathbf{B}_1' \mathbf{A}_1 = \mathbf{0}$  and  $\mathbf{B}_2' \mathbf{A}_2 = \mathbf{0}$ . Denote  $[\mathbf{A}_1, \mathbf{B}_1] = [\mathbf{a}_1, \dots, \mathbf{a}_r, \mathbf{b}_1, \dots, \mathbf{b}_v]$  and  $[\mathbf{A}_2, \mathbf{B}_2] = [\mathbf{a}_{r+1}, \dots, \mathbf{a}_p, \mathbf{b}_{v+1}, \dots, \mathbf{b}_p]$ , which are  $p \times p$  matrices. It follows from Model (2.2) that

$$\mathbf{B}_1' \mathbf{y}_t = \mathbf{B}_1' \mathbf{A}_2 \mathbf{e}_t, \quad (2.3)$$

and, hence,  $\mathbf{B}_1' \mathbf{y}_t$  is a  $v$ -dimensional white noise process. In other words, for any column  $\mathbf{b}_j$  of  $\mathbf{B}_1$  with  $1 \leq j \leq v$ ,  $\{\mathbf{b}_j' \mathbf{y}_t, t = 0, \pm 1, \dots\}$  is a white noise process.

Unlike the traditional factor models, which assume  $\mathbf{x}_t$  and  $\mathbf{e}_s$  are uncorrelated for any  $t$  and  $s$ , we only require  $\text{Cov}(\mathbf{x}_t, \mathbf{e}_{t+j}) = \mathbf{0}$  for  $j \geq 0$  in this paper. For  $k \geq 0$ , let

$$\boldsymbol{\Sigma}_y(k) = \text{Cov}(\mathbf{y}_t, \mathbf{y}_{t-k}), \quad \boldsymbol{\Sigma}_x(k) = \text{Cov}(\mathbf{x}_t, \mathbf{x}_{t-k}), \quad \boldsymbol{\Sigma}_{xe}(k) = \text{Cov}(\mathbf{x}_t, \mathbf{e}_{t-k}),$$

be the covariance matrices of interest. It follows from (2.2) that

$$\boldsymbol{\Sigma}_y(k) = \mathbf{A}_1 \boldsymbol{\Sigma}_x(k) \mathbf{A}_1' + \mathbf{A}_1 \boldsymbol{\Sigma}_{xe}(k) \mathbf{A}_2', \quad k \geq 1, \quad (2.4)$$

and, for  $k = 0$ ,

$$\boldsymbol{\Sigma}_y \equiv \boldsymbol{\Sigma}_y(0) = \mathbf{A}_1 \boldsymbol{\Sigma}_x \mathbf{A}_1' + \mathbf{A}_2 \boldsymbol{\Sigma}_e \mathbf{A}_2'. \quad (2.5)$$

For a pre-specified integer  $k_0 > 0$ , define

$$\mathbf{M} = \sum_{k=1}^{k_0} \boldsymbol{\Sigma}_y(k) \boldsymbol{\Sigma}_y(k)', \quad (2.6)$$

which is a  $p \times p$  semi-positive definite matrix. By  $\mathbf{B}_1' \mathbf{A}_1 = \mathbf{0}$ , we have  $\mathbf{M} \mathbf{B}_1 = \mathbf{0}$ , that is, the columns of  $\mathbf{B}_1$  are the eigenvectors associated with the zero eigenvalues of  $\mathbf{M}$ , and the factor loading space  $\mathcal{M}(\mathbf{A}_1)$  is spanned by the eigenvectors associated with the  $r$  non-zero eigenvalues of  $\mathbf{M}$ . Note that the form of  $\mathbf{M}$  in Equation (2.6) is a special case of the Orthonormalized Partial Least Squares of time series data. See the discussion in Section 5. For  $k_0 > 1$ , the summation in the definition of  $\mathbf{M}$  enables us to pool information over different lags, which is particularly helpful when the sample size is small. In practice, with a given sample size, the estimation accuracy of auto-covariance matrices of  $\mathbf{y}_t$  deteriorates as the lag  $k$  increases. Thus, some compromise in selecting  $k_0$  is needed in real applications. Limited experience suggests that a relatively small  $k_0$  is sufficient in providing useful information concerning the model



structure of  $\mathbf{y}_t$ , because, for a stationary time series, cross-correlation matrices decay to zero exponentially as  $k$  increases. Also, the choice of  $k_0$  seems to be not sensitive. See, for instance, the simulation results in Section 4 and the online supplement.

Turn to the estimation of the common factors. We observe that, from Equation (2.2),

$$\mathbf{B}'_2 \mathbf{y}_t = \mathbf{B}'_2 \mathbf{A}_1 \mathbf{x}_t, \quad (2.7)$$

which is uncorrelated with  $\mathbf{B}'_1 \mathbf{y}_t$  defined in (2.3). Therefore,

$$\mathbf{B}'_2 \boldsymbol{\Sigma}_y \mathbf{B}_1 \mathbf{B}'_1 \boldsymbol{\Sigma}_y \mathbf{B}_2 = \mathbf{0}, \quad (2.8)$$

which implies that  $\mathbf{B}_2$  consists of the last  $r$  eigenvectors corresponding to the zero eigenvalues of  $\mathbf{S} := \boldsymbol{\Sigma}_y \mathbf{B}_1 \mathbf{B}'_1 \boldsymbol{\Sigma}_y$ . From the relationship in (2.7) and the discussion of Remark 1 in Section 3 below,  $\mathbf{B}'_2 \mathbf{A}_1$  is a  $r \times r$  invertible matrix and hence  $\mathbf{x}_t = (\mathbf{B}'_2 \mathbf{A}_1)^{-1} \mathbf{B}'_2 \mathbf{y}_t$ . From Equation (2.2),  $\mathbf{x}_t$  does not include the white noise term. Moreover, the columns in  $\mathbf{A}_2$  can be treated as the eigenvectors associated with the non-zero eigenvalues of  $\mathbf{S}$ . Finally, even though  $\mathbf{B}_2$  (also  $\mathbf{A}_2$ ) is not unique and  $\mathbf{B}_2 \mathbf{H}$  is also a solution to (2.8) for any orthonormal matrix  $\mathbf{H} \in R^{r \times r}$ , this non-uniqueness does not alter the representation of  $\mathbf{x}_t = (\mathbf{B}'_2 \mathbf{A}_1)^{-1} \mathbf{B}'_2 \mathbf{y}_t$ .

### 2.3 Estimation When the Number of Common Factors is Known

For a given realization  $\{\mathbf{y}_t | t = 1, \dots, n\}$ , we discuss our estimation procedure in this section, assuming that the number of common factors  $r$  is known. The selection of  $r$  is discussed in the next section. Let  $\widehat{\boldsymbol{\Sigma}}_y(k)$  be the lag- $k$  sample auto-covariance matrix of  $\mathbf{y}_t$ . To estimate  $\mathcal{M}(\mathbf{A}_1)$ , we perform an eigen-analysis of

$$\widehat{\mathbf{M}} = \sum_{k=1}^{k_0} \widehat{\boldsymbol{\Sigma}}_y(k) \widehat{\boldsymbol{\Sigma}}_y(k)', \quad (2.9)$$

where, as before,  $k_0$  is a pre-specified positive integer. Let  $\widehat{\mathbf{A}}_1 = [\widehat{\mathbf{a}}_1, \dots, \widehat{\mathbf{a}}_r]$  and  $\widehat{\mathbf{B}}_1 = [\widehat{\mathbf{b}}_1, \dots, \widehat{\mathbf{b}}_v]$  be two half orthonormal matrices consisting of the eigenvectors of  $\widehat{\mathbf{M}}$  corresponding to the non-zero and zero eigenvalues, respectively. In view of Equation (2.8), we next perform another eigen-analysis of

$$\widehat{\mathbf{S}} = \widehat{\boldsymbol{\Sigma}}_y \widehat{\mathbf{B}}_1 \widehat{\mathbf{B}}_1' \widehat{\boldsymbol{\Sigma}}_y, \quad (2.10)$$

which is a projected PCA. That is, we project the data  $\mathbf{y}_t$  onto the direction of  $\widehat{\mathbf{B}}_1$ , then perform the PCA between the original data  $\mathbf{y}_t$  and its projected coordinates. Note that  $\widehat{\mathbf{S}} \in R^{p \times p}$  and

its rank is at most  $p - r$ . If the dimension  $p$  is small, we employ  $\widehat{\mathbf{B}}_2 = [\widehat{\mathbf{b}}_{v+1}, \dots, \widehat{\mathbf{b}}_p]$ , where  $\widehat{\mathbf{b}}_{v+1}, \dots, \widehat{\mathbf{b}}_p$  are the eigenvectors corresponding to the smallest  $r$  eigenvalues of  $\widehat{\mathbf{S}}$ . On the other hand, if  $p$  is relatively large, we expect the covariance of the idiosyncratic terms also captures some of the cross-sectional dependence of the data as the covariance of the factors does. That is, given a high-dimensional covariance matrix  $\boldsymbol{\Sigma}_e$ , we entertain that the largest  $K$  eigenvalues of  $\boldsymbol{\Sigma}_e$  are diverging. Therefore, we write  $\mathbf{A}_2 = [\mathbf{A}_{21}, \mathbf{A}_{22}]$  with  $\mathbf{A}_{21} \in R^{p \times K}$  and  $\mathbf{A}_{22} \in R^{p \times (v-K)}$  and consider the linear space  $\mathcal{M}(\mathbf{B}_2^*)$ , where  $\mathbf{B}_2^* = [\mathbf{A}_{22}, \mathbf{B}_2] \in R^{p \times (p-K)}$ . Note that  $\mathbf{B}_2^*$  consists of  $p - K$  eigenvectors corresponding to the  $p - K$  smallest eigenvalues of  $\mathbf{S} = \boldsymbol{\Sigma}_y \mathbf{B}_1 \mathbf{B}_1' \boldsymbol{\Sigma}_y$  defined before. Let  $\widehat{\mathbf{B}}_2^*$  be an estimator of  $\mathbf{B}_2^*$  consisting of the  $p - K$  eigenvectors associated with the  $p - K$  smallest eigenvalues of  $\widehat{\mathbf{S}}$ . We then estimate  $\widehat{\mathbf{B}}_2$  by  $\widehat{\mathbf{B}}_2 = \widehat{\mathbf{B}}_2^* \widehat{\mathbf{R}}$ , where  $\widehat{\mathbf{R}} = [\widehat{\mathbf{r}}_1, \dots, \widehat{\mathbf{r}}_r] \in R^{(p-K) \times r}$  with  $\widehat{\mathbf{r}}_i$  being the eigenvector associated with the  $i$ -th largest eigenvalues of  $\widehat{\mathbf{B}}_2^{*'} \widehat{\mathbf{A}}_1 \widehat{\mathbf{A}}_1' \widehat{\mathbf{B}}_2^*$ . This choice of estimator guarantees that the matrix  $(\widehat{\mathbf{B}}_2' \widehat{\mathbf{A}}_1)^{-1}$  behaves well in recovering the common factor  $\widehat{\mathbf{x}}_t$ . Although, in this case,  $\widehat{\mathbf{B}}_2$  is not a consistent estimator for  $\mathbf{B}_2$ , but  $\widehat{\mathbf{B}}_2^*$  is a consistent one for  $\mathbf{B}_2^*$ , which is sufficient to eliminate the diverging effect of  $\boldsymbol{\Sigma}_e$ . Detailed properties of the estimators are given in Section 3. Finally, we recover the factor process as  $\widehat{\mathbf{x}}_t = (\widehat{\mathbf{B}}_2' \widehat{\mathbf{A}}_1)^{-1} \widehat{\mathbf{B}}_2' \mathbf{y}_t$ .

## 2.4 Determination of the Number of Common Factors

The estimation of  $\mathbf{A}_1$  and  $\mathbf{x}_t$  in the prior sections is based on a given  $r$ , which is unknown in practice. There are several methods available in the literature to determine  $r$  for the traditional factor model in Equation (1.1). See, for example, the information criterion in Bai and Ng (2002) and Bai (2003), the random matrix theory method in Onatski (2010), and the ratio-based method in Lam and Yao (2012), Ahn and Horenstein (2013), and Li et al. (2017), among others. However, none of these methods is applicable to Model (2.1) directly. The most relevant method is the one based on testing the number of zero canonical correlations between  $\mathbf{y}_t$  and an extended vector of its lagged values employed in Gao and Tsay (2019). But this testing method only works when the dimension  $p$  is relatively small with respect to the sample size  $n$ . For a large  $p$ , some alternatives must be sought.

In this section, we propose a new approach to estimate the number of common factors based on Equation (2.3). Specifically, we perform white noise tests to determine the number of white noise components  $\widehat{v}$  and use  $\widehat{r} = p - \widehat{v}$ . Let  $\widehat{\mathbf{G}}$  be the matrix of eigenvectors (in the decreasing order of eigenvalues) of the sample matrix  $\widehat{\mathbf{M}}$  of Equation (2.9) and  $\widehat{\mathbf{u}}_t = \widehat{\mathbf{G}}' \mathbf{y}_t = (\widehat{u}_{1t}, \dots, \widehat{u}_{pt})'$  be the transformed series. We propose to test sequentially the number of white noises in  $\widehat{\mathbf{u}}_t$ , which is an estimate of  $v$ . To this end, we consider two cases depending on the

dimension  $p$ .

If the dimension  $p$  is small, we recommend using a bottom-up procedure to determine the number of white noise components. Specifically, we use the conventional test statistics, such as the well-known Ljung-Box statistic  $Q(m)$  or its rank-based variant, to test the null hypothesis that  $\hat{u}_{it}$  is a white noise series starting with  $i = p$ . If the null hypothesis is rejected, then  $\hat{v} = 0$  and  $\hat{r} = p$ ; otherwise, reduce  $i$  by one and repeat the testing process. Clearly, this testing process can only last until  $i = 1$ . If all transformed series  $\hat{u}_{it}$  are white noise, then  $\hat{v} = p$  and  $\hat{r} = 0$ . In general, if  $\hat{u}_{it}$  is not a white noise series but  $\hat{u}_{jt}$  are for  $j = i + 1, \dots, p$ , then  $\hat{v} = p - i$  and  $\hat{r} = i$ , and we have  $\hat{\mathbf{G}} = [\hat{\mathbf{A}}_1, \hat{\mathbf{B}}_1]$ , where  $\hat{\mathbf{A}}_1 \in R^{p \times \hat{r}}$  and  $\hat{\mathbf{B}}_1 \in R^{p \times \hat{v}}$ .

For a large  $p$ , the conventional white-noise test statistics are no longer adequate, but some methods have been developed in recent years to test high-dimensional white noise. See, for instance, Chang et al. (2017), Tsay (2020), and Li et al. (2019). We only consider the first two methods in this paper since the third one requires the noise to have an independent structure and its covariance is identity. The method by Chang et al. (2017) makes use of the maximum absolute auto-correlations and cross-correlations of the component series. Specifically, let  $\mathbf{w}_t = (w_{1t}, \dots, w_{dt})'$  be a  $d$ -dimensional real-valued time series. In this paper,  $1 \leq d \leq p$ . Define the lag- $k$  sample covariance matrix as  $\hat{\Sigma}_w(k) = (n - k)^{-1} \sum_{t=k+1}^n (\mathbf{w}_t - \bar{\mathbf{w}})(\mathbf{w}_{t-k} - \bar{\mathbf{w}})'$ , where  $\bar{\mathbf{w}} = n^{-1} \sum_{t=1}^n \mathbf{w}_t$  is the sample mean. The test statistic  $T_n$  of Chang et al. (2017) is

$$T_n = \max_{1 \leq k \leq \bar{k}} T_{n,k}, \quad (2.11)$$

where  $\bar{k} \geq 1$  is a pre-specified positive integer and  $T_{n,k} = \max_{1 \leq j, l \leq d} n^{1/2} |\hat{\rho}_{jl}(k)|$  with

$$\hat{\Gamma}_w(k) \equiv [\hat{\rho}_{jl}(k)]_{1 \leq j, l \leq d} = \text{diag}\{\hat{\Sigma}_w(0)\}^{-1/2} \hat{\Sigma}_w(k) \text{diag}\{\hat{\Sigma}_w(0)\}^{-1/2}.$$

The limiting distribution of  $T_n$  can be approximated by that of the  $L_\infty$ -norm of a normal random vector, i.e., there exists a random variable  $\mathbf{z}_d \sim N(\mathbf{0}, \Theta_{d,n})$  such that

$$\sup_{s \geq 0} |P(T_n > s) - P(\|\mathbf{z}_d\|_\infty > s)| = o(1),$$

where  $\Theta_{d,n}$  is close to the asymptotic covariance of the vector containing the columns of  $\hat{\Gamma}_w(1)$  to  $\hat{\Gamma}_w(\bar{k})$ , and it can be estimated from  $\{\mathbf{w}_t | t = 1, \dots, n\}$ . Therefore, the critical values of  $T_n$  can be obtained by bootstrapping from a multivariate normal distribution.

The second method of high-dimensional white noise test is introduced by Tsay (2020) using the extreme value theory. The test is robust with a closed-form limiting distribution under

some weak assumptions and is easy to use in practice. The basic idea of the test is as follows. Consider a  $d$ -dimensional time series  $\mathbf{w}_t$  with a realization of  $n$  observations  $\{\mathbf{w}_t | t = 1, \dots, n\}$ . Assume, for now, that  $d < n$ . Let  $\tilde{\mathbf{w}}_t = \Sigma_w^{-1/2} \mathbf{w}_t$  be a standardized series, where  $\Sigma_w^{1/2}$  is a square-root matrix of the covariance matrix  $\Sigma_w$ . With  $d < n$ , this standardization can be done by PCA. For simplicity, we denote the standardized realization as  $\tilde{\mathbf{w}}_t = \hat{\Sigma}_w^{-1/2} \mathbf{w}_t$ . If  $d \geq n$ ,  $\hat{\Sigma}_w$  is singular and we discuss a modification later. Note that the components of  $\tilde{\mathbf{w}}_t = (\tilde{w}_{1t}, \dots, \tilde{w}_{dt})'$  are mutually uncorrelated. Next, let  $\hat{\boldsymbol{\rho}}_t = (\hat{\rho}_{1t}, \dots, \hat{\rho}_{dt})'$  be the rank series of  $\tilde{\mathbf{w}}_t$ , where  $\hat{\rho}_{jt}$  is the rank of  $\tilde{w}_{jt}$  in  $\{\tilde{w}_{j,1}, \dots, \tilde{w}_{j,n}\}$  for  $1 \leq j \leq d$ . The lag- $\ell$  rank cross-correlation matrix is then defined as

$$\hat{\Gamma}_{w,\ell} = \frac{12}{n(n^2 - 1)} \sum_{t=\ell+1}^n (\hat{\boldsymbol{\rho}}_t - \bar{\boldsymbol{\rho}})(\hat{\boldsymbol{\rho}}_{t-\ell} - \bar{\boldsymbol{\rho}})',$$

where  $\bar{\boldsymbol{\rho}} = \frac{n+1}{2} \mathbf{1}_d$  and  $\mathbf{1}_d$  is a  $d$ -dimensional vector of ones. The test statistic of Tsay (2020) for testing that there is no serial or cross-sectional correlation in the first  $m$  lags of  $\mathbf{w}_t$  is

$$T(m) = \max\{\sqrt{n}|\hat{\Gamma}_{w,\ell}(j, k)| : 1 \leq j, k \leq d, 1 \leq \ell \leq m\}, \quad (2.12)$$

where  $\hat{\Gamma}_{w,\ell}(j, k)$  is the  $(j, k)$ -th element of  $\hat{\Gamma}_{w,\ell}$ . By the extreme-value theory, the limiting distribution of  $T(m)$  under the white noise hypothesis is a Gumbel distribution provided that the component series of  $\mathbf{w}_t$  follow a continuous distribution. Therefore, we reject the null hypothesis  $H_0 : \mathbf{w}_t$  is a vector white noise at the  $\alpha$ -level if

$$T(m) \geq c_{d,m} \times x_{1-\alpha/2} + s_{d,m},$$

where  $x_{1-\alpha/2} = -\log(-\log(1 - \alpha/2))$  is the  $(1 - \alpha/2)$ -th quantile of the standard Gumbel distribution and

$$c_{d,m} = [2 \log(d^2 m)]^{-1/2}, \text{ and } s_{d,m} = \sqrt{2 \log(d^2 m)} - \frac{\log(4\pi) + \log(\log(d^2 m))}{2(2 \log(d^2 m))^{1/2}}.$$

If  $d \geq n$ , the sample covariance matrix of  $\mathbf{w}_t$  is singular and some alternative methods must be sought to create mutually uncorrelated series. Tsay (2020) provided a method by selecting a subset series of  $\mathbf{w}_t$  to perform testing, and the method works reasonably well in simulations and some illustrative applications. In this paper, we consider a simpler method by using the relations in (2.2) and (2.3). Note that in our testing,  $\mathbf{w}_t$  is a subset of the transformed series  $\hat{\mathbf{u}}_t = \hat{\mathbf{G}}' \mathbf{y}_t$ . Since  $\hat{\mathbf{M}}$  is based on the covariance matrices of  $\mathbf{y}_t$  and its lagged values the components of  $\hat{\mathbf{u}}_t$  associated with small eigenvalues contain little information on the dynamical

dependence of  $\mathbf{y}_t$ . Therefore, we can drop the last  $(p - \varepsilon n)$  components of the transformed series  $\widehat{\mathbf{u}}_t$  without affecting the white-noise test, where  $\varepsilon \in (0, 1)$ . In other words, when  $p > n$ , we cannot start with  $\mathbf{w}_t = \widehat{\mathbf{u}}_t$ , but we can choose  $\mathbf{w}_t$  to consist of the first  $d = \varepsilon n < n$  components of  $\widehat{\mathbf{u}}_t$  to perform the white-noise test without affecting the determination of  $r$  under the assumption that  $r$  is small in applications.

Return to the determination of  $r$  when  $p$  is large. We can apply the high-dimensional white noise test of Chang et al. (2017) or Tsay (2020) to subsets of the transformed series  $\widehat{\mathbf{u}}_t$ . We choose the subset simply by dropping the last  $p - \varepsilon n$  projected coordinates in  $\widehat{\mathbf{u}}_t$  when  $p \geq n$  because those coordinates capture less dynamic dependencies by the nature of the projection method. Specifically, let  $p_* = p$  if  $p < n$  and  $p_* = \varepsilon n$  if  $p \geq n$ , where  $\varepsilon \in (0, 1)$ . Starting with  $i = 1$  and  $\mathbf{w}_t = (\widehat{u}_{i,t}, \dots, \widehat{u}_{p_*,t})'$ , we test the null hypothesis that  $\mathbf{w}_t$  has no serial or cross-sectional correlations in the first  $m$  lags using a selected test statistic. If the null hypothesis is rejected, increase  $i$  by one and repeat the testing process. Using this testing process, we select  $\widehat{r}$  as  $i - 1$  for which the  $i$ th test does not reject the null hypothesis. Note that since both test statistics considered use the maximum of absolute correlations, the computation of the testing process is trivial because we only need to compute the cross-correlation matrices of  $\mathbf{w}_t$  at each time.

## 2.5 Modeling Procedure

Figure 1 provides a flowchat that summarizes the proposed modeling procedure. Selecting a small positive integer  $k_0$ , we start with an eigenvalue-eigenvector analysis of the sample matrix  $\widehat{\mathbf{M}}$ , then apply the white-noise test to the transformed series, using the eigenvectors of  $\widehat{\mathbf{M}}$ , to determine the dimension  $\widehat{v}$  of the white noise process, which in turn provides an estimate of the number of common factors  $\widehat{r} = p - \widehat{v}$ . Next, we perform the proposed projected PCA to find the orthogonal directions of the diverging noises so that the common factors can be properly estimated. The projected PCA can mitigate the impact of white noises in recovering the common factors. Finally, with  $\widehat{\mathbf{A}}_1$  and the estimated factor process  $\widehat{\mathbf{x}}_t$ , we compute the  $h$ -step ahead prediction of the  $\mathbf{y}_t$  series using the formula  $\widehat{\mathbf{y}}_{n+h} = \widehat{\mathbf{A}}_1 \widehat{\mathbf{x}}_{n+h}$ , where  $\widehat{\mathbf{x}}_{t+h}$  is an  $h$ -step ahead forecast for  $\mathbf{x}_t$  based on the estimated past values  $\widehat{\mathbf{x}}_1, \dots, \widehat{\mathbf{x}}_n$ . This can be done, for example, by fitting a VAR model to  $\{\widehat{\mathbf{x}}_1, \dots, \widehat{\mathbf{x}}_n\}$ . Alternatively, we may also adopt the diffusion index models to do forecasting if we are interested in some particular component of  $\mathbf{y}_t$ . See Stock and Watson (2002a) and Stock and Watson (2002b) for details.




Figure 1: The flowchart of the proposed modeling procedure. ①: Applying sequential vector white noise test on  $\hat{\mathbf{G}}'y_t$ ; ②: Finding the orthogonal directions of the diverging noises.

### 3 Theoretical Properties

This section studies the asymptotic theory of the estimation method used in the paper. We start with the assumption that the number of common factors  $r$  is known, and divide the derivations into two cases depending on the value of the dimension  $p$ . The case of estimated  $r$  is discussed later.

#### 3.1 Asymptotic Properties When $p$ is Fixed, But $n \rightarrow \infty$

We consider first the asymptotic properties of the estimators when  $p$  is fixed but  $n \rightarrow \infty$ . These properties show the behavior of our estimation method when  $n$  is large and  $p$  is relatively small. We begin with the assumptions used.

**Assumption 1.** *The process  $\{(\mathbf{y}_t, \mathbf{f}_t)\}$  is  $\alpha$ -mixing with the mixing coefficient satisfying the*

condition  $\sum_{k=1}^{\infty} \alpha_p(k)^{1-2/\gamma} < \infty$  for some  $\gamma > 2$ , where

$$\alpha_p(k) = \sup_i \sup_{A \in \mathcal{F}_{-\infty}^i, B \in \mathcal{F}_{i+k}^{\infty}} |P(A \cap B) - P(A)P(B)|,$$

and  $\mathcal{F}_i^j$  is the  $\sigma$ -field generated by  $\{(\mathbf{y}_t, \mathbf{f}_t) : i \leq t \leq j\}$ .

**Assumption 2.**  $E|f_{it}|^{2\gamma} < C_1$  and  $E|\varepsilon_{jt}|^{2\gamma} < C_2$  for  $1 \leq i \leq r$  and  $1 \leq j \leq v$ , where  $C_1$  and  $C_2 > 0$  are constants, and  $\gamma$  is given in Assumption 1.

**Assumption 3.**  $\lambda_1 > \dots > \lambda_r > \lambda_{r+1} = \dots = \lambda_p = 0$ , where  $\lambda_i$  is the  $i$ -th largest eigenvalue of the  $\mathbf{M}$  matrix in Equation (2.6).

Assumption 1 is standard for dependent random processes. See Gao et al. (2019) for a theoretical justification for VAR models. The conditions in Assumption 2 imply that  $E|y_{it}|^{2\gamma} < \infty$  under the setting that  $p$  is fixed. The moment condition used is slightly stronger than the ones in Lam and Yao (2012) and Fan et al. (2013), which only assume the existence of finite fourth moments, because we use the  $\alpha$ -mixing condition on the series, which is weaker than the  $\psi$ -mixing in Lam and Yao (2012) and the exponential rate of decay of the mixing coefficient in Fan et al. (2013). In Assumption 3, if the  $r$  non-zero eigenvalues of  $\mathbf{M}$  are distinct, the eigenvector matrix  $\mathbf{A}_1$  is uniquely defined if we ignore the trivial replacement of  $\mathbf{a}_j$  by  $-\mathbf{a}_j$  for  $1 \leq j \leq r$ . In general, the choice of  $\mathbf{A}_1$  in Model (2.2) is not unique without Assumption 3, so we consider the error in estimating  $\mathcal{M}(\mathbf{A}_1)$ , the column space of  $\mathbf{A}_1$ , because  $\mathcal{M}(\mathbf{A}_1)$  is uniquely defined by (2.2) and it does not vary with different choices of  $\mathbf{A}_1$ . The same argument also applies to matrices  $\mathbf{A}_2$ ,  $\mathbf{B}_1$ , and  $\mathbf{B}_2$ . To this end, we adopt the discrepancy measure used by Pan and Yao (2008): for two  $p \times r$  half orthogonal matrices  $\mathbf{H}_1$  and  $\mathbf{H}_2$  satisfying the condition  $\mathbf{H}'_1 \mathbf{H}_1 = \mathbf{H}'_2 \mathbf{H}_2 = \mathbf{I}_r$ , the difference between the two linear spaces  $\mathcal{M}(\mathbf{H}_1)$  and  $\mathcal{M}(\mathbf{H}_2)$  is measured by

$$D(\mathbf{H}_1, \mathbf{H}_2) = \sqrt{1 - \frac{1}{r} \text{tr}(\mathbf{H}_1 \mathbf{H}'_1 \mathbf{H}_2 \mathbf{H}'_2)}. \quad (3.1)$$

Note that  $D(\mathbf{H}_1, \mathbf{H}_2) \in [0, 1]$ . It is equal to 0 if and only if  $\mathcal{M}(\mathbf{H}_1) = \mathcal{M}(\mathbf{H}_2)$ , and to 1 if and only if  $\mathcal{M}(\mathbf{H}_1) \perp \mathcal{M}(\mathbf{H}_2)$ . The following theorem establishes the consistency of the estimated loading matrix  $\widehat{\mathbf{A}}_1$ , its orthonormal complement  $\widehat{\mathbf{B}}_1$ , the matrix  $\widehat{\mathbf{B}}_2$ , and the extracted common factor  $\widehat{\mathbf{A}}_1 \widehat{\mathbf{x}}_t$ .

**Theorem 1.** *Suppose Assumptions 1 and 2 hold and  $r$  is known and fixed. Then, for fixed  $p$ ,*

$$D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = O_p(n^{-1/2}), \quad D(\widehat{\mathbf{B}}_1, \mathbf{B}_1) = O_p(n^{-1/2}), \quad \text{and} \quad D(\widehat{\mathbf{B}}_2, \mathbf{B}_2) = O_p(n^{-1/2}),$$

as  $n \rightarrow \infty$ . If Assumption 3 also holds, we further have  $\|\widehat{\mathbf{A}}_1 - \mathbf{A}_1\|_2 = O_p(n^{-1/2})$ . Therefore,

$$\|\widehat{\mathbf{A}}_1 \widehat{\mathbf{x}}_t - \mathbf{A}_1 \mathbf{x}_t\|_2 = O_p(n^{-1/2}).$$

**Remark 1.** From Theorem 1 and, as expected, the convergence rates of all estimates are standard at  $\sqrt{n}$ , which is commonly seen in the traditional statistical theory. To recover the factor process, we need to guarantee that  $\mathbf{B}'_2 \mathbf{A}_1$  is invertible. This follows from the fact that there exist  $\mathbf{R}_1 \in R^{r \times r}$  and  $\mathbf{R}_2 \in R^{v \times r}$  such that  $\mathbf{B}_2 = \mathbf{L}_1 \mathbf{R}_1 + \mathbf{L}_2 \mathbf{R}_2 = \mathbf{A}_1 \mathbf{Q}_1 \mathbf{R}_1 + \mathbf{A}_2 \mathbf{Q}_2 \mathbf{R}_2$ , i.e., each column of  $\mathbf{B}_2$  can be represented as a linear combination of the columns of  $\mathbf{L}$ . Therefore,  $\mathbf{I}_r = \mathbf{B}'_2 \mathbf{B}_2 = \mathbf{B}'_2 \mathbf{A}_1 \mathbf{Q}_1 \mathbf{R}_1$  and, hence,  $\text{rank}(\mathbf{B}'_2 \mathbf{A}_1) = r$ , which is of full rank.

Theorem 1 implies that the convergence rate does not change even when some non-zero eigenvalues of  $\mathbf{M}$  are not distinct and  $\mathbf{A}_1$  is not uniquely defined. In fact, the consistency of the column spaces  $\mathcal{M}(\mathbf{B}_1)$  and  $\mathcal{M}(\mathbf{B}_2)$  is more meaningful because their columns correspond to the zero eigenvalues of  $\mathbf{M}$  and  $\mathbf{S}$ , respectively, and cannot be uniquely characterized.

### 3.2 Asymptotic Properties When $n \rightarrow \infty$ And $p \rightarrow \infty$

Turn to the case of high-dimensional time series. Note that our framework is different from the PCA approach in Bai and Ng (2002) and Fan et al. (2013), where the consistency of the leading eigenvectors does not require any restriction between  $p$  and  $n$  by using the well-known  $\sin(\theta)$  theorem of Davis and Kahan (1970) and Yu et al. (2015), among others, because they imposed the pervasiveness assumption that all eigenvalues associated with the factors and loadings are diverging at the rate of  $p$ , which implies that the covariance has some spiked eigenvalues. See Section 2 of Fan et al. (2013) for details. However, certain developments in random matrix theory have shown that the true eigenvalues and eigenvectors might not be consistently estimated from the sample covariance matrix when  $p/n$  is not negligible in general. See, for example, Johnstone and Lu (2009) and Paul (2007). Therefore, using sample auto-covariance-based analysis without the pervasiveness assumption, we do expect some theoretical restrictions between  $p$  and  $n$ . It is well-known that if the dimension  $p$  diverges faster than  $n^{1/2}$ , the sample covariance matrix is no longer a consistent estimate of the population covariance matrix. When  $p = o(n^{1/2})$ , it is still possible to consistently estimate the factor loading matrix  $\mathbf{A}$  and the number of common factors  $r$ . See Gao and Tsay (2019) for details. Therefore, without any additional assumptions on the structure of the underlying time series,  $p$  can only be as large as  $o(n^{1/2})$ . To deal with the case of large  $p$ , we impose some conditions on the transformation matrix  $\mathbf{L}$  of Equation (2.1) and the cross dependence of time series  $\mathbf{y}_t$ .



Let  $\mathbf{L} = [\mathbf{c}_1, \dots, \mathbf{c}_p]$ , where  $\mathbf{c}_i$  is a  $p$ -dimensional column vector, and  $\mathbf{L}_1 = [\mathbf{c}_1, \dots, \mathbf{c}_r]$  and  $\mathbf{L}_2 = [\mathbf{c}_{r+1}, \dots, \mathbf{c}_p]$ .

**Assumption 4.** (i)  $\mathbf{L}_1 = [\mathbf{c}_1, \dots, \mathbf{c}_r]$  such that  $\|\mathbf{c}_j\|_2^2 \asymp p^{1-\delta_1}$ ,  $j = 1, \dots, r$  and  $\delta_1 \in [0, 1)$ ; (ii) For each  $j = 1, \dots, r$  and  $\delta_1$  given in (i),  $\min_{\theta_i \in R, i \neq j} \|\mathbf{c}_j - \sum_{1 \leq i \leq r, i \neq j} \theta_i \mathbf{c}_i\|_2^2 \asymp p^{1-\delta_1}$ .

**Assumption 5.** (i)  $\mathbf{L}_2$  admits a singular value decomposition  $\mathbf{L}_2 = \mathbf{A}_2 \mathbf{D}_2 \mathbf{V}_2'$ , where  $\mathbf{A}_2 \in R^{p \times v}$  is given in Equation (2.2),  $\mathbf{D}_2 = \text{diag}(d_1, \dots, d_v)$  and  $\mathbf{V}_2 \in R^{v \times v}$  satisfying  $\mathbf{V}_2' \mathbf{V}_2 = \mathbf{I}_v$ ; (ii) There exists a finite integer  $0 < K < v$  such that  $d_1 \asymp \dots \asymp d_K \asymp p^{(1-\delta_2)/2}$  for some  $\delta_2 \in [0, 1)$  and  $d_{K+1} \asymp \dots \asymp d_v \asymp 1$ .

**Assumption 6.** (i) For  $\gamma$  given in Assumption 1, any  $\mathbf{h} \in R^v$  and  $0 < c_h < \infty$  with  $\|\mathbf{h}\|_2 = c_h$ ,  $E|\mathbf{h}'\boldsymbol{\varepsilon}_t|^{2\gamma} < \infty$ ; (ii)  $\sigma_{\min}(\mathbf{R}'\mathbf{B}_2^* \mathbf{A}_1) \geq C_3$  for some constant  $C_3 > 0$  and some half orthogonal matrix  $\mathbf{R} \in R^{(p-K) \times r}$  satisfying  $\mathbf{R}'\mathbf{R} = \mathbf{I}_r$ , where  $\sigma_{\min}$  denotes the minimum non-zero singular value of a matrix.

The quantity  $\delta_1$  of Assumption 4 is used to quantify the strength of the factors, and the eigenvalues of  $\mathbf{L}_1 \mathbf{L}_1'$  are all of order  $p^{1-\delta_1}$ . If  $\delta_1 = 0$ , the corresponding factors are called strong factors, since it includes the case where each element of  $\mathbf{c}_i$  is  $O(1)$ . If  $\delta_1 > 0$ , the corresponding factors are weak factors and the smaller the  $\delta_1$  is, the stronger the factors are. An advantage of using index  $\delta_1$  is to link the convergence rates of the estimated factors explicitly to the strength of the factors. Assumption 4 ensures that all common factors in  $\mathbf{x}_t$  are of equal strength  $\delta_1$ . In practice, the factors may have multiple levels of strength as in Lam and Yao (2012), among others. We can make similar assumptions and the consistency of the loading matrix would then depend on the strength of the weakest factors. We do not consider this issue here to save space. There are many sufficient conditions for Assumption 5 to hold. For example, it holds if we allow  $(\mathbf{c}_{r+1}, \dots, \mathbf{c}_{r+K})$  to satisfy Assumption 4 for some  $\delta_2 \in [0, 1)$ , and the  $L_1$ - and  $L_\infty$ -norms of  $(\mathbf{c}_{r+K+1}, \dots, \mathbf{c}_p)$  are all finite. A special case is to let  $\mathbf{c}_{r+K+j}$  be a standard unit vector. The constraint between  $\delta_1$  and  $\delta_2$  will be illustrated later under different scenarios to guarantee the consistency. Assumption 6(i) is mild and includes the standard normal distribution as a special case. Together with Assumption 2 and the aforementioned sufficient condition for Assumption 5, it is not hard to show  $E|y_{it}|^{2\gamma} < \infty$ , but we do not address this issue explicitly here. Assumption 6(ii) is reasonable since  $\mathbf{B}_2$  is a subspace of  $\mathbf{B}_2^*$ , and Remark 1 implies that  $\mathbf{R}'\mathbf{B}_2^* \mathbf{A}_1$  is invertible. The choice of  $\widehat{\mathbf{R}}$  and hence  $\widehat{\mathbf{B}}_2 = \widehat{\mathbf{B}}_2^* \widehat{\mathbf{R}}$  will be discussed later.

**Remark 2.** In Assumption 5, we actually only require  $d_K \asymp p^{(1-\delta_2)/2}$  for some  $\delta_2 \in [0, 1)$  and  $K \geq 1$ , and the upper singular values  $\{d_1, \dots, d_{K-1}\}$  if  $K > 1$  can be even larger provided that the largest one  $d_1$  should be bounded by another rate  $p^{(1-\delta_3)/2}$  for some  $0 \leq \delta_3 \leq \delta_2$ . For simplicity, we assume the top singular values are of the same order.

If the dimension  $p$  is large, it is not possible to consistently estimate  $\mathbf{B}_2$  or even  $\mathcal{M}(\mathbf{B}_2)$ . Instead, we estimate  $\mathbf{B}_2^* = (\mathbf{A}_{22}, \mathbf{B}_2)$  or equivalently  $\mathcal{M}(\mathbf{B}_2^*)$ , which is the subspace spanned by the eigenvectors associated with the  $p - K$  smallest eigenvalues of  $\mathbf{S}$ . Assume  $\widehat{\mathbf{B}}_2^*$  consists of the eigenvectors corresponding to the smallest  $p - K$  eigenvalues of  $\widehat{\mathbf{S}}$ . Under some conditions, we can show that  $\mathcal{M}(\widehat{\mathbf{B}}_2^*)$  is consistent to  $\mathcal{M}(\mathbf{B}_2^*)$ . This is also the case in the literature on high-dimensional PCA with i.i.d. data. See, for example, Shen et al. (2016) and the references therein. Therefore, the choice of  $\widehat{\mathbf{B}}_2$  should be a subspace of  $\widehat{\mathbf{B}}_2^*$ , and we will discuss it before Theorem 4 below. If there exist cross-correlations between  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_{t-j}$  for  $j \geq 1$ , we assume  $\text{rank}(\boldsymbol{\Sigma}_{f\varepsilon}(k)) = r$  and define

$$\kappa_{\min} = \min_{1 \leq k \leq k_0} \|\boldsymbol{\Sigma}_{f\varepsilon}(k)\|_{\min} \text{ and } \kappa_{\max} = \max_{1 \leq k \leq k_0} \|\boldsymbol{\Sigma}_{f\varepsilon}(k)\|_2, \quad (3.2)$$

where  $\|\cdot\|_{\min}$  denotes the smallest nonzero singular value,  $\kappa_{\min}$  and  $\kappa_{\max}$  can be either finite constants or diverging rates in relation to  $p$  and  $n$ , and they control the strength of the dependence between  $\mathbf{f}_t$  and the past errors  $\boldsymbol{\varepsilon}_{t-j}$  for  $j \geq 1$ . The maximal order of  $\kappa_{\max}$  is  $p^{1/2}$  which is the Frobenius norm of  $\boldsymbol{\Sigma}_{f\varepsilon}(k)$  and  $\kappa_{\max} = 0$  (hence  $\kappa_{\min} = 0$ ) implies that  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_s$  are independent for all  $t$  and  $s$ . Throughout this article, if  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_s$  are independent for all  $t$  and  $s$ , then  $\kappa_{\min} = \kappa_{\max} = 0$  and all the conditions and expressions below concerning  $\kappa_{\min}$  and  $\kappa_{\max}$  are removed.

**Theorem 2.** *Suppose Assumptions 1-6 hold and  $r$  is known and fixed. As  $n \rightarrow \infty$ , if  $p^{\delta_1} n^{-1/2} = o(1)$  or  $\kappa_{\max}^{-1} p^{\delta_1/2 + \delta_2/2} n^{-1/2} = o(1)$ , then*

$$D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = \begin{cases} O_p(p^{\delta_1} n^{-1/2}), & \text{if } \kappa_{\max} p^{\delta_1/2 - \delta_2/2} = o(1), \\ O_p(\kappa_{\min}^{-2} p^{\delta_2} n^{-1/2} + \kappa_{\min}^{-2} \kappa_{\max} p^{\delta_1/2 + \delta_2/2} n^{-1/2}), & \text{if } r \leq K, \kappa_{\min}^{-1} p^{\delta_2/2 - \delta_1/2} = o(1), \\ O_p(\kappa_{\min}^{-2} p n^{-1/2} + \kappa_{\min}^{-2} \kappa_{\max} p^{1 + \delta_1/2 - \delta_2/2} n^{-1/2}), & \text{if } r > K, \kappa_{\min}^{-1} p^{(1 - \delta_1)/2} = o(1), \end{cases}$$

and the above results also hold for  $D(\widehat{\mathbf{B}}_1, \mathbf{B}_1)$ . Furthermore,

$$D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*) = O_p\left(p^{2\delta_2 - \delta_1} n^{-1/2} + p^{\delta_2} n^{-1/2} + (1 + p^{2\delta_2 - 2\delta_1}) D(\widehat{\mathbf{B}}_1, \mathbf{B}_1)\right).$$

**Remark 3.** (i) If  $\kappa_{\max} = \kappa_{\min} = 0$ , i.e.,  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_s$  are independent for all  $t$  and  $s$ , we have

$$D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = O_p(p^{\delta_1} n^{-1/2}) \text{ and } D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*) = O_p(p^{2\delta_2 - \delta_1} n^{-1/2} + p^{\delta_2} n^{-1/2} + p^{\delta_1} n^{-1/2}).$$

To guarantee that these estimates are consistent, we require  $p^{\delta_1} n^{-1/2} = o(1)$ ,  $p^{\delta_2} n^{-1/2} = o(1)$  and  $p^{2\delta_2 - \delta_1} n^{-1/2} = o(1)$ . When  $p \asymp n^{1/2}$ , it implies that  $0 \leq \delta_1 < 1$ ,  $0 \leq \delta_2 < 1$  and  $\delta_2 < (1 + \delta_1)/2$ , i.e., the ranges of  $\delta_1$  and  $\delta_2$  are pretty wide. On the other hand, if  $p \asymp n$ , we

see that  $0 \leq \delta_1 < 1/2$ ,  $0 \leq \delta_2 < 1/2$  and  $2\delta_2 - \delta_1 < 1/2$ , these ranges become narrower if  $p$  is large.

(ii) When  $\kappa_{\max} \neq 0$  and  $\kappa_{\min} \neq 0$ , there are many possible results. A reasonable assumption is  $\kappa_{\min} \asymp \kappa_{\max} \asymp p^{\delta/2}$  for some  $0 \leq \delta < 1$  since  $r$  is small. For example, set  $\delta = \delta_1$ ,

$$D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = \begin{cases} O_p(p^{\delta_1} n^{-1/2}), & \text{if } p^{\delta_1 - \delta_2/2} = o(1), \\ O_p(p^{\delta_2/2} n^{-1/2}), & \text{if } r \leq K, \kappa_{\min}^{-1} p^{\delta_2/2 - \delta_1/2} = o(1), \end{cases}$$

and there is no consistency result when  $r > K$ . Furthermore, we have  $D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*) = O_p(p^{2\delta_2 - \delta_1} n^{-1/2} + p^{\delta_2} n^{-1/2})$ . Thus, we require  $p^{\delta_1} n^{-1/2} = o(1)$ ,  $p^{\delta_2} n^{-1/2} = o(1)$  and  $p^{2\delta_2 - \delta_1} n^{-1/2} = o(1)$ . The ranges of  $\delta_1$  and  $\delta_2$  are the same as discussed in Remark 3(i) above, we omit the details here. On the other hand, if  $\delta > (1 - \delta_1)/2$ , it is still possible to obtain consistent estimates when  $r > K$ , the discussion is similar and is omitted for simplicity.

From Theorem 2, we see that when  $p \asymp n$ , we require  $\delta_1 < 1/2$  and  $\delta_2 < 1/2$  to guarantee the consistency of our estimation method, which rules out the cases of the presence of weaker factors with  $\delta_1 \geq 1/2$  and a slower diverging of the noise covariance matrix with  $\delta_2 \geq 1/2$ . The convergence rates in Theorem 2 are not optimal and they can be further improved under some additional assumption on  $\varepsilon_t$  below.

**Assumption 7.** For any  $\mathbf{h} \in R^v$  with  $\|\mathbf{h}\|_2 = 1$ , there exists a constant  $C_4 > 0$  such that

$$P(|\mathbf{h}'\varepsilon_t| > x) \leq 2 \exp(-C_4 x) \quad \text{for any } x > 0.$$

Assumption 7 implies that  $\varepsilon_t$  are sub-exponential, which is a larger class of distributions than sub-gaussian, and includes the uniform distribution on every convex body following the Brunn-Minkowski inequality. See, for example, Vershynin (2012) and Vershynin (2018).

**Theorem 3.** Suppose Assumptions 1-7 hold and  $r$  is known and fixed, and  $p^{\delta_1/2} n^{-1/2} = o(1)$ ,  $p^{\delta_2/2} n^{-1/2} = o(1)$ . (i) Under the condition that  $\delta_1 \leq \delta_2$ ,

$$D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = \begin{cases} O_p(p^{\delta_1/2} n^{-1/2}), & \text{if } \kappa_{\max} p^{\delta_1/2 - \delta_2/2} = o(1), \\ O_p(\kappa_{\min}^{-2} p^{\delta_2 - \delta_1/2} n^{-1/2} + \kappa_{\min}^{-2} \kappa_{\max} p^{\delta_2/2} n^{-1/2}), & \text{if } r \leq K, \kappa_{\min}^{-1} p^{\delta_2/2 - \delta_1/2} = o(1), \\ O_p(\kappa_{\min}^{-2} p^{1 - \delta_1/2} n^{-1/2} + \kappa_{\min}^{-2} \kappa_{\max} p^{1 - \delta_2/2} n^{-1/2}), & \text{if } r > K, \kappa_{\min}^{-1} p^{(1 - \delta_1)/2} = o(1), \end{cases}$$

and the above results also hold for  $D(\widehat{\mathbf{B}}_1, \mathbf{B}_1)$ , and

$$D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*) = O_p(p^{2\delta_2 - 3\delta_1/2} n^{-1/2} + p^{2\delta_2 - 2\delta_1} D(\widehat{\mathbf{B}}_1, \mathbf{B}_1)).$$

(ii) Under the condition that  $\delta_1 > \delta_2$ , if  $\kappa_{\max} = 0$  and  $p^{\delta_1 - \delta_2/2} n^{-1/2} = o(1)$ , then

$$D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = O_p(p^{\delta_1 - \delta_2/2} n^{-1/2}).$$

If  $\kappa_{\max} \gg 0$ , then

$$D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = \begin{cases} O_p(\kappa_{\min}^{-2} \kappa_{\max} p^{\delta_1/2} n^{-1/2}), & \text{if } r \leq K, \kappa_{\min}^{-1} p^{\delta_2/2 - \delta_1/2} = o(1), \\ O_p(\kappa_{\min}^{-2} \kappa_{\max} p^{1 + \delta_1/2 - \delta_2} n^{-1/2}), & \text{if } r > K, \kappa_{\min}^{-1} p^{(1 - \delta_1)/2} = o(1), \end{cases}$$

and the above results also hold for  $D(\widehat{\mathbf{B}}_1, \mathbf{B}_1)$ , and

$$D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*) = O_p(p^{\delta_2/2} n^{-1/2} + D(\widehat{\mathbf{B}}_1, \mathbf{B}_1)).$$

**Remark 4.** (i) Consider the case  $\kappa_{\min} = \kappa_{\max} = 0$ . If  $\delta_1 \leq \delta_2$ ,  $D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = O_p(p^{\delta_1/2} n^{-1/2})$  and  $D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*) = O_p(p^{2\delta_2 - 3\delta_1/2} n^{-1/2})$ . For  $p \asymp n$ , we require  $0 \leq \delta_1 \leq \delta_2 < 1$  and  $4\delta_2 - 3\delta_1 < 1$ , or equivalently  $0 \leq \delta_1 \leq \delta_2 < 3\delta_1/4 + 1/4$ . If  $\delta_1 > \delta_2$ ,  $D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = O_p(p^{\delta_1 - \delta_2/2} n^{-1/2})$  and  $D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*) = O_p(p^{\delta_1 - \delta_2/2} n^{-1/2})$ . Thus, if  $p \asymp n$ , we require  $\max\{2\delta_1 - 1, 0\} < \delta_2 < \delta_1 < 1$ . Therefore, if  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_s$  are independent and  $p \asymp n$ ,  $\delta_1$  and  $\delta_2$  need to satisfy  $0 \leq \delta_1 \leq \delta_2 < 3\delta_1/4 + 1/4$  or  $\max\{2\delta_1 - 1, 0\} < \delta_2 < \delta_1 < 1$ , which is much wider than those of Theorem 3.

(ii) If  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_s$  are correlated for  $s < t$ , we may have many consistency results depending on the strength of the dependence between  $\mathbf{f}_t$  and  $\boldsymbol{\varepsilon}_s$ . In addition, if  $\text{rank}(\boldsymbol{\Sigma}_{f\varepsilon}) < r$ , the conditions  $r \leq K$  and  $r > K$  in Theorems 2 and 3 would become  $\text{rank}(\boldsymbol{\Sigma}_{f\varepsilon}) \leq K$  and  $\text{rank}(\boldsymbol{\Sigma}_{f\varepsilon}) > K$ , respectively. We omit the details here.

(iii) When  $K$  increases with  $p$  and  $n$ , similar results can still be established, depending on the growing rates of  $K$ ,  $p$ , and  $n$ . See also Remark 2(c) in Fan et al. (2013). But we do not pursue this issue here.

By Remark 1 and Assumption 6, so long as  $\mathcal{M}(\widehat{\mathbf{A}}_1)$  and  $\mathcal{M}(\widehat{\mathbf{B}}_2^*)$  are consistent estimators for  $\mathcal{M}(\mathbf{A}_1)$  and  $\mathcal{M}(\mathbf{B}_2^*)$ , respectively, there must exist an  $\mathbf{R}$  which is the same as that in Assumption 6(ii) such that  $\mathcal{M}(\widehat{\mathbf{B}}_2^* \mathbf{R})$  is a consistent estimator for  $\mathcal{M}(\mathbf{B}_2)$ , which implies the invertibility of  $\mathbf{R}' \widehat{\mathbf{B}}_2^* \widehat{\mathbf{A}}_1$ . Therefore, once having  $\widehat{\mathbf{B}}_2^*$ , we suggest to choose  $\widehat{\mathbf{B}}_2$  as  $\widehat{\mathbf{B}}_2 = \widehat{\mathbf{B}}_2^* \widehat{\mathbf{R}}$ , where  $\widehat{\mathbf{R}} = (\widehat{\mathbf{r}}_1, \dots, \widehat{\mathbf{r}}_r) \in R^{(p-K) \times r}$ , and  $\widehat{\mathbf{r}}_i$  is the vector associated with the  $i$ -th largest eigenvalues of  $\widehat{\mathbf{B}}_2^* \widehat{\mathbf{A}}_1 \widehat{\mathbf{A}}_1' \widehat{\mathbf{B}}_2^*$ . This choice guarantees that the matrix  $(\widehat{\mathbf{B}}_2^* \widehat{\mathbf{A}}_1)^{-1}$  behaves well when recovering the factor  $\widehat{\boldsymbol{\alpha}}_i$  and is sufficient for our analysis. On the other hand, this choice could still eliminate the diverging part of the noise covariance matrix and gives prominent convergence rate, as shown in Theorem 4. There are many ways to choose the number of components  $K$  in Assumption 5 so long as  $p - K > r$ . We will discuss the choice of  $K$  in Remark 5 below

and also in Section 5. The following theorem states the convergence rate of the extracted common factors.

**Theorem 4.** *Under the conditions in Theorem 2 or 3, we have*

$$p^{-1/2} \|\widehat{\mathbf{A}}_1 \widehat{\mathbf{x}}_t - \mathbf{A}_1 \mathbf{x}_t\|_2 = O_p\{p^{-1/2} + p^{-\delta_1/2} D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) + p^{-\delta_2/2} D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*)\}. \quad (3.3)$$

Furthermore, if  $\mathbf{f}_t$  satisfies Assumption 7 with a constant  $C_5 > 0$ , we have

$$p^{-1/2} \max_{1 \leq t \leq n} \|\widehat{\mathbf{A}}_1 \widehat{\mathbf{x}}_t - \mathbf{A}_1 \mathbf{x}_t\|_2 = O_p\{p^{-1/2} \log(n) + p^{-\delta_1/2} \log(n) D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) + p^{-\delta_2/2} \log(n) D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*)\}.$$

**Remark 5.** (i) A similar result is given in Theorem 3 of Lam et al. (2011) for the approximate factor models. The above results also hold for  $\|\widehat{\mathbf{A}}_2 \widehat{\mathbf{e}}_t - \mathbf{A}_2 \mathbf{e}_t\|_2$  by a simple manipulation, where  $\widehat{\mathbf{A}}_2 \widehat{\mathbf{e}}_t = \mathbf{y}_t - \widehat{\mathbf{A}}_1 \widehat{\mathbf{x}}_t$ . In addition, if all the eigenvalues associated with the idiosyncratic covariance are bounded as in Lam et al. (2011), by the proof of Theorem 5 in the supplement,  $p^{-\delta_2/2} D(\widehat{\mathbf{B}}_2^*, \mathbf{B}_2^*)$  will disappear and (3.3) reduces to Theorem 3 in Lam et al. (2011), and we can still consistently estimate the extracted factors. When  $\delta_1 = \delta_2 = 0$ , i.e., the factors and the noise terms are all strong, the convergence rate in (3.3) is  $O_p(p^{-1/2} + n^{-1/2})$ , which is the optimal rate specified in Theorem 3 of Bai (2003) when dealing with the traditional approximate factor models.

(ii) Selecting the number of principal components is a common issue in the literature, and many approaches are available. Since it is impossible to eliminate all the noise effects in recovering the factors and we only need to guarantee that the diverging part of the noises are removed for large  $p$ , we may select  $K$  in a range of possible values. In practice, let  $\widehat{\mu}_1 \geq \dots \geq \widehat{\mu}_p$  be the sample eigenvalues of  $\widehat{\mathbf{S}}$  and define  $\widehat{K}_L$  as

$$\widehat{K}_L = \arg \min_{1 \leq j \leq \widehat{K}_U} \{\widehat{\mu}_{j+1}/\widehat{\mu}_j\}. \quad (3.4)$$

Let  $\widehat{K}_U$  be a pre-specified integer. In practice, we suggest  $\widehat{K}_U = \min\{\sqrt{p}, \sqrt{n}, p - \widehat{r}, 10\}$ . Then the estimator  $\widehat{K}$  of  $K$  can assume some value between  $\widehat{K}_L$  and  $\widehat{K}_U$ .

Next, we study the consistency of the white noise test described in Section 2. The consistency conditions depend on the test statistic used. In what follows, we only present the consistency for large  $p$  since the case of small  $p$  is trivial.

**Theorem 5.** (i) Assume that Assumptions 1-7 hold. If  $D(\widehat{\mathbf{B}}_1, \mathbf{B}_1)^2 \|\boldsymbol{\Sigma}_y\|_2 = o_p(1)$ , then the test statistic  $T_n$  defined in (2.11) can consistently estimate  $r$ , i.e.,  $P(\widehat{r} = r) \rightarrow 1$  as  $n \rightarrow \infty$ .

(ii) Suppose Assumptions 1-7 hold and  $\mathbf{f}_t$  also satisfies Assumption 7. If

$$p^{1/2} \log(np) \max\{p^{\delta_2/2-\delta_1/2} D(\widehat{\mathbf{B}}_1, \mathbf{B}_1), D(\widehat{\mathbf{B}}_1, \mathbf{B}_1), O_p(p^{\delta_2/2} n^{-1/2})\} = o_p(1),$$

then the test statistic  $T(m)$  of (2.12) can also consistently estimate  $r$ .

**Remark 6.** (i) If  $\mathbf{f}_t$  and  $\varepsilon_s$  are independent for all  $t$  and  $s$ , the conditions in Theorem 5(i) are essentially  $pn^{-1} = o(1)$  if  $\delta_1 \leq \delta_2$  and  $p^{1+2\delta_1-2\delta_2} n^{-1} = o(1)$  if  $\delta_1 > \delta_2$ . Thus, we require  $p \asymp n^\xi$  with  $0 < \xi < 1$  for both cases. As for Theorem 5(ii), the condition is  $p^{(1+\delta_2)/2} n^{-1/2} \log(np) = o(1)$  if  $\delta_1 \leq \delta_2$  and  $p^{1/2+\delta_1-\delta_2/2} n^{-1/2} \log(np) = o(1)$  if  $\delta_1 > \delta_2$ , and hence we also require  $p \asymp n^\xi$  for some  $0 < \xi < 1$ .

(ii) Even though the conditions in Theorem 5(ii) are slightly stronger, the method based on  $T(m)$  is simple and easy to use, and its performance is satisfactory when  $p$  is moderately large. See Tsay (2020) and the simulation results in Section 4 for details. We use those stronger conditions, because we require the effect of the orthogonalization using sample PCA on the rank of each component of the white noise part is asymptotically negligible uniformly in terms of  $t$  and  $p$  in the proofs.

With the estimator  $\widehat{r}$ , we may define the estimator for  $\mathbf{A}_1$  as  $\widetilde{\mathbf{A}}_1 = (\widehat{\mathbf{a}}_1, \dots, \widehat{\mathbf{a}}_{\widehat{r}})$ , where  $\widehat{\mathbf{a}}_1, \dots, \widehat{\mathbf{a}}_{\widehat{r}}$  are the orthonormal eigenvectors of  $\widehat{\mathbf{M}}$ , defined in (2.9), corresponding to the  $\widehat{r}$  largest eigenvalues. In addition, we may also replace  $r$  by  $\widehat{r}$  in the methodology described in Section 2. To see the impact of  $\widehat{r}$  on measuring the errors in estimating the loading spaces, we take  $\widetilde{\mathbf{A}}_1$  for example and use

$$\widetilde{D}(\widetilde{\mathbf{A}}_1, \mathbf{A}_1) = \sqrt{1 - \frac{1}{\max(\widehat{r}, r)} \text{tr}(\widetilde{\mathbf{A}}_1 \widetilde{\mathbf{A}}_1' \mathbf{A}_1 \mathbf{A}_1')}, \quad (3.5)$$

which is a modified version of (3.1). It takes into account the fact that the dimensions of  $\widetilde{\mathbf{A}}_1$  and  $\mathbf{A}_1$  may be different, and we use  $\max(r_1, r_2)$  instead of  $\min(r_1, r_2)$  to guarantee that the quantity in the square-root is nonnegative. We also see that  $\widetilde{D}(\widetilde{\mathbf{A}}_1, \mathbf{A}_1) = D(\widehat{\mathbf{A}}_1, \mathbf{A}_1)$  if  $\widehat{r} = r$ . We show below that  $\widetilde{D}(\widetilde{\mathbf{A}}_1, \mathbf{A}_1) \rightarrow 0$  in probability at the same rate of  $D(\widehat{\mathbf{A}}_1, \mathbf{A}_1)$ . Therefore,  $\mathcal{M}(\widetilde{\mathbf{A}}_1)$  is still a consistent estimator for  $\mathcal{M}(\mathbf{A}_1)$  even without knowing  $r$ . Let  $\rho_n$  be the convergence rate of  $D(\widehat{\mathbf{A}}_1, \mathbf{A}_1)$  as in Theorem 1, 2, or 3, that is,  $\rho_n D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) = O_p(1)$ . For any  $\varepsilon > 0$ , there exists a positive constant  $M_\varepsilon$  such that  $P(\rho_n D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) > M_\varepsilon) < \varepsilon$ . Then,

$$\begin{aligned} P(\rho_n \widetilde{D}(\widetilde{\mathbf{A}}_1, \mathbf{A}_1) > M_\varepsilon) &\leq P(\rho_n D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) > M_\varepsilon, \widehat{r} = r) + P(\rho_n \widetilde{D}(\widetilde{\mathbf{A}}_1, \mathbf{A}_1) > M_\varepsilon, \widehat{r} \neq r) \\ &\leq P(\rho_n D(\widehat{\mathbf{A}}_1, \mathbf{A}_1) > M_\varepsilon, \widehat{r} = r) + o(1) \leq \varepsilon + o(1) \rightarrow \varepsilon, \end{aligned}$$

which implies  $\rho_n \tilde{D}(\tilde{\mathbf{A}}_1, \mathbf{A}_1) = O_p(1)$ . Therefore,  $\mathcal{M}(\tilde{\mathbf{A}}_1)$  has the oracle property in estimating the factor loading space  $\mathcal{M}(\mathbf{A}_1)$  in the sense that it has the same convergence rate as  $D(\hat{\mathbf{A}}_1, \mathbf{A}_1)$ . The result also holds for  $\hat{\mathbf{B}}_1$  by a similar argument. Therefore, we still use  $\hat{\mathbf{A}}_1$  and  $\hat{\mathbf{B}}_1$  as our estimators when replacing  $r$  by  $\hat{r}$ . As for the influence of  $\hat{r}$  and  $\hat{K}$  on  $\hat{\mathbf{B}}_2$ , note that we do not have the parameter  $K$  when  $p$  is small, and hence the impact of  $\hat{r}$  on  $\hat{\mathbf{B}}_2$  remains negligible asymptotically by a similar argument. When  $p$  is large, note that we only eliminate the possible diverging effects of the noises in our approach. Thus, an accurate estimator for  $K$  is not necessary and any estimator  $\hat{K}$  which is larger than the true one is sufficient for our method to work. On the other hand, we can show  $\hat{K}_L$  is a consistent estimator for  $K$  under certain conditions by a similar argument as that of Lam and Yao (2012). Therefore, we may set an upper bound  $\hat{K}_U$  as in Remark 5(ii) and choose  $\hat{K}$  between  $\hat{K}_L$  and  $\hat{K}_U$ . The choice of  $\hat{K}_U$  is not unique but it cannot exceed  $p - \hat{r}$  since the rank of  $\hat{\mathbf{S}}$  is at most  $p - \hat{r}$ . The optimal one can be chosen via some cross-validation such as the out-of-sample testing if one is concerned with the forecasting performance of the extracted factors. For further information, see the empirical example in Section 4 and the concluding remark in Section 5.

## 4 Numerical Properties

We use simulation and a real example to assess the performance of the proposed analysis in finite samples. An additional real example is given in the online supplement.

### 4.1 Simulation

In this section, we illustrate the finite-sample properties of the proposed methodology under the scenarios when  $p$  is both small and large. As the dimensions of  $\hat{\mathbf{A}}_1$  and  $\mathbf{A}_1$  are not necessarily the same, and  $\mathbf{L}_1$  is not an orthogonal matrix in general, we first extend the discrepancy measure in Equation (3.5) to a more general one below. Let  $\mathbf{H}_i$  be a  $p \times r_i$  matrix with  $\text{rank}(\mathbf{H}_i) = r_i$ , and  $\mathbf{P}_i = \mathbf{H}_i(\mathbf{H}_i' \mathbf{H}_i)^{-1} \mathbf{H}_i'$ ,  $i = 1, 2$ . Define

$$\bar{D}(\mathbf{H}_1, \mathbf{H}_2) = \sqrt{1 - \frac{1}{\max(r_1, r_2)} \text{tr}(\mathbf{P}_1 \mathbf{P}_2)}, \quad (4.1)$$

then  $\bar{D} \in [0, 1]$ . Furthermore,  $\bar{D}(\mathbf{H}_1, \mathbf{H}_2) = 0$  if and only if either  $\mathcal{M}(\mathbf{H}_1) \subset \mathcal{M}(\mathbf{H}_2)$  or  $\mathcal{M}(\mathbf{H}_2) \subset \mathcal{M}(\mathbf{H}_1)$ , and it is 1 if and only if  $\mathcal{M}(\mathbf{H}_1) \perp \mathcal{M}(\mathbf{H}_2)$ . When  $r_1 = r_2 = r$  and  $\mathbf{H}_i' \mathbf{H}_i = \mathbf{I}_r$ ,  $\bar{D}(\mathbf{H}_1, \mathbf{H}_2)$  is the same as that in Equation (3.1). We only present the simulation results for  $k_0 = 2$  in Equation (2.9) to save space because other choices of  $k_0$  produce similar

patterns as shown in the online supplement.

**Example 1.** Consider Model (2.1) with common factors following the VAR(1) model

$$\mathbf{f}_t = \mathbf{\Phi}\mathbf{f}_{t-1} + \boldsymbol{\eta}_t,$$

where  $\boldsymbol{\eta}_t$  is a white noise process. We set the true number of factors  $r = 3$ , the dimension  $p = 5, 10, 15, 20$ , and the sample size  $n = 200, 500, 1000, 1500, 3000$ . For each realization of  $\mathbf{y}_t$ , the elements of the loading matrix  $\mathbf{L}$  are drawn independently from  $U(-2, 2)$ , and the elements of  $\mathbf{L}_2$  are then divided by  $\sqrt{p}$  to balance the accumulated variances of  $f_{it}$  and  $\varepsilon_{it}$  for each component of  $\mathbf{y}_t$ .  $\mathbf{\Phi}$  is a diagonal matrix with its diagonal elements being drawn independently from  $U(0.5, 0.9)$ ,  $\boldsymbol{\varepsilon}_t \sim N(0, \mathbf{I}_v)$  and  $\boldsymbol{\eta}_t \sim N(0, \mathbf{I}_r)$ . We use 1000 replications for each  $(p, n)$  configuration.

We first study the performance of estimating the number of factors. Since  $p$  is relatively small compared to the sample size  $n$ , we use Ljung-Box test statistics with  $m = 10$  to determine the number of factors, i.e.,  $Q(10)$ . The empirical probabilities  $P(\hat{r} = r)$  are reported in Table 1. From the table, we see that, for a given  $p$ , the performance of the proposed method improves as the sample size increases. On the other hand, for a given  $n$ , the empirical probability decreases as  $p$  increases and the probability for  $p = 20$  is roughly half of that for  $p = 5$ , which is understandable since it is harder to determine the correct number of factors when the dimension increases and the errors in the testing procedure accumulate. Overall, the Ljung-Box test works well for the case of small dimension (e.g.,  $p \leq 10$ ). However, when  $p$  is slightly larger (e.g.,  $p = 15, 20$ ), the test statistic tends to overestimate the number of factors, implying that we can still keep sufficient information of the original process  $\mathbf{y}_t$ . To illustrate, we present the boxplots of  $\bar{D}(\hat{\mathbf{A}}_1, \mathbf{L}_1)$  in Figure 2(a), where  $\bar{D}(\cdot, \cdot)$  is defined in (4.1). From Figure 2(a), for each  $p$ , the discrepancy decreases as the sample size increases and this is in agreement with the theory. The plot also shows that, as expected, the mean discrepancy increases as the dimension  $p$  increases.

Furthermore, for each  $(p, n)$ , we study the root-mean-square error (RMSE):

$$\text{RMSE} = \left( \frac{1}{n} \sum_{t=1}^n \|\hat{\mathbf{A}}_1 \hat{\mathbf{x}}_t - \mathbf{L}_1 \mathbf{f}_t\|_2^2 \right)^{1/2}, \quad (4.2)$$

which quantifies the accuracy in estimating the common factor process. Boxplots of the RMSE are shown in Figure 2(b). From the plot, we see a clear pattern that, as the sample size increases, the RMSE decreases for a given  $p$ , which is consistent with the results of Theorem



Table 1: Empirical probabilities  $P(\hat{r} = r)$  of various  $(p, n)$  configurations for the model of Example 1 with  $r = 3$ , where  $p$  and  $n$  are the dimension and the sample size, respectively. 1000 iterations are used.

		$n$					
		$p$	200	500	1000	1500	3000
$r = 3$	5	0.861	0.889	0.890	0.912	0.926	
	10	0.683	0.718	0.723	0.735	0.748	
	15	0.506	0.555	0.561	0.599	0.601	
	20	0.395	0.425	0.441	0.447	0.453	




Figure 2: (a) Boxplots of  $\bar{D}(\hat{\mathbf{A}}_1, \mathbf{L}_1)$  when  $r = 3$  under the scenario that  $p$  is relatively small in Example 1; (b) Boxplots of the RMSE defined in (4.2) when  $r = 3$  under the scenario that  $p$  is relatively small in Example 1. The sample sizes used are 200, 500, 1000, 1500, 3000, and the results are based on 1000 iterations.

1. Obviously, as expected, the RMSE increases with the dimension  $p$ . Overall, the one-by-one testing procedure for determining  $r$  works well when the dimension of  $\mathbf{y}_t$  is small, and the RMSE decreases when the sample size increases, but the performance of the test may deteriorate into overestimation of the number of the factors for larger  $p$ .

**Example 2.** In this example, we consider Model (2.1) with  $\mathbf{f}_t$  being the same as that of Example 1. We set the true number of factors  $r = 5$  and the number of the spiked components  $K = 3$  and 7 defined in Assumption 5 for the loading of white noise processes. The dimensions used are  $p = 50, 100, 300, 500$ , and the sample sizes are  $n = 300, 500, 1000, 1500, 3000$ . We consider three scenarios for the strength parameters  $\delta_1$  and  $\delta_2$ :  $(\delta_1, \delta_2) = (0, 0)$ ,  $(\delta_1, \delta_2) = (0.4, 0.5)$ , and  $(\delta_1, \delta_2) = (0.5, 0.4)$ . For each realization of  $\mathbf{y}_t$ , the elements of the loading matrix  $\mathbf{L}$  are drawn independently from  $U(-2, 2)$ , and then we divide  $\mathbf{L}_1$  by  $p^{\delta_1/2}$ , the first  $K$  columns of  $\mathbf{L}_2$  by  $p^{\delta_2/2}$ , and the remaining  $v - K$  columns by  $p$  to satisfy Assumptions 4 and

5.  $\Phi$ ,  $\varepsilon_t$  and  $\eta_t$  are drawn similarly as those of Example 1. We also use 1000 replications in each experiment.

We first study the performance of the high-dimensional white noise test. For simplicity, we only present the results of the  $T(m)$  statistics defined in (2.12) and the results for the other test are similar. When  $p \geq n$ , we only keep the upper  $\varepsilon n$  components of  $\widehat{\mathbf{G}}'\mathbf{y}_t$  with  $\varepsilon = 0.75$  in the testing and the test still works well for other choices of  $\varepsilon$  as shown in the online supplement. The testing results are reported in Table 2 for  $r = 5$  with  $K = 3$  and  $K = 7$ , respectively. From Table 2, we see that for each setting of  $(\delta_1, \delta_2)$  and fixed  $p$ , the performance of the white noise test improves as the sample size increases. The performance is also quite satisfactory for moderately larger  $p$ . Also, for  $n \geq 1000$ , the empirical probabilities of selecting the correct number of factors are high for both  $K = 3$  and  $K = 7$ .

To shed some light on the advantages of the proposed methodology, we compare it with that of Lam et al. (2011) (denoted by LYB) in selecting the number of factors. For the ratio-based method in LYB, let  $\widehat{\lambda}_1, \dots, \widehat{\lambda}_p$  be the eigenvalues of  $\widehat{\mathbf{M}}$  and define

$$\widehat{r} = \arg \min_{1 \leq j \leq R} \left\{ \frac{\widehat{\lambda}_{j+1}}{\widehat{\lambda}_j} \right\}, \quad (4.3)$$

where we choose  $R = p/2$  as suggested in their paper. Figure A1 of the online supplement presents the boxplots of  $\widehat{r}$ . We see from Figure A1 that the estimated number of factors  $\widehat{r}$  is the sum of the number of common factors and the number of spiked components of the noises. The result indicates that the ratio-based method may fail to identify the correct number of factors with dynamic dependencies if the covariance matrix of the noise has diverging eigenvalues. On the other hand, the high-dimensional white noise test considered in the paper continues to work well.

Next, we study the accuracy of the estimated loading matrices as that in Example 1. The boxplots of  $\bar{D}(\widehat{\mathbf{A}}_1, \mathbf{L}_1)$  are shown in Figure 3. Similar patterns are also obtained for the estimation of other matrices, and we omit them here. From Figure 3, there is a clear pattern that the estimation accuracy of the loading matrix improves as the sample size increases even for moderately large  $p$ , which is in line with our asymptotic theory. The results also confirm that the proposed white noise test selects  $\widehat{r}$  reasonably well even for large  $p$ .

Finally, since a key difference between the proposed method and that of LYB is that we allow some of the eigenvalues of the noise covariance to diverge, it is of interest to compare the two methodologies. Denoting the proposed method by GT, we compared it with LYB in

Table 2: Empirical probabilities  $P(\hat{r} = r)$  for Example 2 with  $r = 5$ ,  $K = 3$  or  $7$ , where  $p$  and  $n$  are the dimension and the sample size, respectively.  $\delta_1$  and  $\delta_2$  are the strength parameters of the factors and the errors, respectively. 1000 iterations are used.

$(\delta_1, \delta_2)$	$p$	$n (K = 3)$					$n (K = 7)$				
		300	500	1000	1500	3000	300	500	1000	1500	3000
(0,0)	50	0.510	0.833	0.906	0.917	0.926	0.418	0.688	0.904	0.908	0.910
	100	0.538	0.799	0.910	0.916	0.922	0.426	0.754	0.910	0.916	0.918
	300	0.582	0.907	0.916	0.924	0.932	0.406	0.686	0.914	0.925	0.926
	500	0.560	0.888	0.918	0.928	0.932	0.614	0.778	0.912	0.918	0.920
(0.4,0.5)	50	0.717	0.903	0.928	0.929	0.935	0.806	0.820	0.892	0.912	0.926
	100	0.800	0.924	0.938	0.940	0.944	0.800	0.914	0.922	0.904	0.922
	300	0.858	0.904	0.928	0.932	0.952	0.939	0.935	0.935	0.929	0.930
	500	0.834	0.922	0.932	0.933	0.948	0.898	0.904	0.926	0.930	0.933
(0.5,0.4)	50	0.420	0.890	0.910	0.916	0.920	0.332	0.856	0.900	0.928	0.938
	100	0.508	0.868	0.912	0.928	0.936	0.356	0.716	0.920	0.922	0.928
	300	0.581	0.910	0.926	0.929	0.932	0.384	0.688	0.924	0.936	0.945
	500	0.678	0.928	0.936	0.938	0.934	0.421	0.778	0.924	0.930	0.931




Figure 3: Boxplots of  $\bar{D}(\hat{\mathbf{A}}_1, \mathbf{L}_1)$  when  $r = 5$  and  $K = 3$  under the scenario that  $p$  is relatively large in Example 2. The sample sizes used are  $n = 300, 500, 1000, 1500, 3000$  and the number of iterations is 1000.

Table 3: The RMSE defined in (4.4) when  $r = 5$  and  $K = 7$  in Example 2. The sample sizes used are  $n = 300, 500, 1000, 1500, 3000$ . Standard errors are given in the parentheses and 1000 iterations are used. GT denotes the proposed method and LYB is the one in Lam et al. (2011)

Method	$p$	$n$				
		300	500	1000	1500	3000
GT	50	1.510(0.233)	1.124(0.235)	0.770(0.235)	0.627(0.224)	0.488(0.273)
LYB		3.056(0.085)	3.051(0.081)	3.056(0.075)	3.053(0.122)	2.976(0.400)
GT	100	1.490(0.179)	1.148(0.188)	0.817(0.141)	0.677(0.126)	0.519(0.191)
LYB		3.050(0.074)	3.056(0.065)	3.053(0.055)	3.046(0.159)	3.024(0.257)
GT	300	1.729(0.118)	1.463(0.107)	1.149(0.094)	1.107(0.079)	0.769(0.077)
LYB		3.052(0.047)	3.055(0.047)	3.053(0.040)	3.056(0.037)	3.056(0.034)
GT	500	1.753(0.089)	1.547(0.081)	1.285(0.052)	1.044(0.070)	0.861(0.047)
LYB		3.057(0.053)	3.050(0.042)	3.054(0.035)	3.055(0.034)	3.055(0.027)

terms of the RMSE defined below:

$$\text{RMSE} = \left( \frac{1}{np} \sum_{t=1}^n \|\widehat{\mathbf{A}}_1 \widehat{\mathbf{x}}_t - \mathbf{L}_1 \mathbf{f}_t\|_2^2 \right)^{1/2}, \quad (4.4)$$

which is different from that in Equation (4.2) because we have another factor  $p^{-1/2}$  in (4.4). This RMSE quantifies the estimation accuracy of the common factor process. In the comparison, the number of factors is obtained by the corresponding method of each methodology. The results are shown in Table 3 for  $r = 5$ ,  $K = 7$  and  $\delta_1 = \delta_2 = 0$ . The pattern is similar for the other settings of  $\delta_1$  and  $\delta_2$ . When calculating  $\widehat{\mathbf{B}}_2^*$  using our method, we choose the number of components  $\widehat{K} = 10$ , which is fixed for all the iterations. Thus,  $\widehat{\mathbf{B}}_2^*$  contains  $p - \widehat{K}$  columns corresponding to the  $p - \widehat{K}$  smallest eigenvalues of  $\widehat{\mathbf{S}}$ . From the table, we see that, because the ratio-based method tends to overestimate the number of common factors  $r$  in the presence of diverging eigenvalues in the covariance matrix of the idiosyncratic component, the RMSE of our method is much smaller than that obtained by Lam et al. (2011). Also, as expected, for a given  $p$ , the RMSE tends to decrease when the sample size increases. This is in agreement with the asymptotic theory in Theorem 4. In the online supplement, we fix the number of factors and compare our method with that of LYB using the RMSE criterion to check the need to mitigate the noise effect. Our method continues to produce smaller RMSEs in general when the noise effect is diverging. See Table A8 of the supplement for details. Overall, under the assumption that the top eigenvalues of the noise covariance matrix are diverging in high-dimensional time series, the proposed method outperforms the existing one in the literature.

## 4.2 Real data analysis

In this section, we demonstrate the application of the proposed method using U.S. monthly macroeconomic data with 108 time series and 283 observations. An additional real example is shown in the online supplement, where the dimension  $p$  is greater than the sample size  $n$ .

**Example 3.** In this example, we employ the data set of monthly U.S. macroeconomic time series, which is available from <https://www.princeton.edu/~mwatson/publi.html> and used in Stock and Watson (2009). There are 108 variables available in the data file spanning from 1959 to 2006. To remove any possible trend in the series, we take the first difference  $\mathbf{y}_t = \tilde{\mathbf{y}}_t - \tilde{\mathbf{y}}_{t-1}$ , where  $\tilde{\mathbf{y}}_t \in R^{108}$  and  $t = 2, \dots, 576$ . The differenced series are shown in Figure A2 of the online supplement, from which we see that the series continue to display some strong nonstationary patterns between later 1990s and early 2000s. This might be due to several economic and financial crises in the US and around the world in the period, including the 1997 Asian financial crisis and the 9/11 attack in 2001. Therefore, we use the data from December 1972 to July 1996 only. The resulting differenced U.S. monthly macroeconomic time series are shown in Figure A3 of the online supplement with  $n = 283$  and  $p = 108$ .

Following the proposed modeling procedure, we first applied the proposed white-noise tests of Section 2.3 to identify the number of common factors and found  $\hat{r} = 8$ . We also calculated the eigenvalues of  $\hat{\mathbf{S}}$ , and Figure A4(a) of the online supplement plots the first ten eigenvalues. We see clearly that, in this particular instance, the largest eigenvalues are extremely large because the average of the first ten eigenvalues of  $\hat{\Sigma}_y$  is  $1.0 \times 10^6$ . In our analysis, we choose  $\hat{K} = 5$ , but the results are similar for other choices. The spectral densities of the eight estimated common factors are given in Figure A5 of the supplement, and they show clearly that the estimated factors are all serially dependent processes. In addition, the variance of the first factor is extremely large compared to those of the others. We also applied the method of LYB to  $\mathbf{y}_t$ , and the estimated number of factors is  $\hat{r} = 1$ . We further present the spectral densities of the first six transformed series  $\hat{u}_{1t}, \dots, \hat{u}_{6t}$  using the eigen-analysis of LYB in Figure A6 of the supplement and find that none of these series is white noise. This marks a contradiction with the assumptions used in LYB, which assume that  $\hat{u}_{2t}$  to  $\hat{u}_{6t}$  are white noises.

Next we examine the forecasting performance of the extracted factors via different methods. We estimate the models using the data in the time span  $[1, \tau]$  with  $\tau = 183, \dots, 283 - h$  for the  $h$ -step ahead forecasts. First, we study the overall performance by employing a VAR(1) or an AR(1) model for the estimated factor processes and compute  $h$ -step ahead predictions of  $\mathbf{y}_t$

using the associated factor loadings. The forecast error is defined as

$$\text{FE}_h = \frac{1}{100 - h + 1} \sum_{\tau=183}^{283-h} E(\tau, h) \quad \text{with} \quad E(\tau, h) = \frac{1}{\sqrt{p}} \|\hat{\mathbf{y}}_{\tau+h} - \mathbf{y}_{\tau+h}\|_2, \quad (4.5)$$

where  $p = 108$ . Table 4 reports the 1-step to 3-step ahead forecast errors using VAR(1) or AR(1) models, where we also used the first 8 factors obtained by the method in Bai and Ng (2002) although it identifies 30 factors. The smallest forecast error of each step is shown in boldface. From the table, we see that, except for the 1-step ahead forecast, our method is capable of producing accurate forecasts and the associated forecast errors based on the extracted factors by our method are smaller than those based on the other two methods. The PCA method also works well in the comparison. Next, we further examine the forecastability of the extracted factors by different methods. We adopt the diffusion index model as in Stock and Watson (2002a) and Stock and Watson (2002b) and predict some particular components of  $\mathbf{y}_t$ . We focus on two important components: (a) CPI-All: the Consumer Price Index for All Urban Consumers: All Items in U.S. City Average, and (b) CPI-Core: the Consumer Price Index for All Urban Consumers: All Items Less Food and Energy in U.S. City Average. These two components are often used to measure the inflation. The difference between CPI-All and CPI-Core is that the latter does not include the more volatile categories of food and energy prices. To see the forecastability of the extracted factors, we only regress  $y_{i,t+h}$  on  $\hat{\mathbf{f}}_t$  without including the lagged variable of  $y_{it}$ , because Stock and Watson (2002b) found that the simpler diffusion index forecasts excluding the lags are even better than that with the lagged ones. Table 5 presents the 1-step to 3-step ahead mean squared forecast errors for the CPI-All and the CPI-Core, where F-DI denotes the factor-based diffusion index model as in Stock and Watson (2002b) without the lagged variables, and F-AR represents the factor-based VAR or AR approach mentioned before. From Table 5, we see that, for factor-based diffusion index models, the forecastability of the factors extracted by our method fares better than those of the other two methods, which is understandable since our extracted factors capture most of the dynamic dependencies. For the factor-based VAR or AR approach, our method is not as good as the PCA approach, but better than the LYB, in forecasting the CPI-All, but it fares better than the other two approaches in predicting the CPI-Core. In conjunction with the results of Table 4, our approach fares better in terms of the overall MSE defined in (4.5), but it may not perform uniformly better than the others for every component of  $\mathbf{y}_t$ .

To further compare the forecast ability of extracted factors by different approaches, we employ the diffusion index models and adopt the asymptotic test in Diebold and Mariano (1995) to test for equal predictive ability. The null hypothesis of interest is that the approaches

Table 4: The 1-step to 3-step ahead forecast errors. GT denotes the proposed method, BN denotes the principal component analysis as in Bai and Ng (2002), and LYB is the one in Lam et al. (2011). Boldface numbers denote the smallest one for a given model.

Step	Model	Methods		
		GT	BN	LYB
1	AR(1)	264.1	<b>260.1</b>	347.7
2	AR(1)	<b>286.1</b>	288.1	359.4
3	AR(1)	<b>306.1</b>	309.7	379.4

Table 5: The 1-step to 3-step ahead mean squared forecast errors for inflation indexes. GT denotes the proposed method, BN denotes the principal component analysis as in Bai and Ng (2002) and LYB is the one in Lam et al. (2011). Boldface numbers denote the smallest one for a given model. CPI - All denotes the Consumer Price Index for All Urban Consumers: All Items in U.S. City Average, and CPI - Core denotes the Consumer Price Index for All Urban Consumers: All Items Less Food and Energy in U.S. City Average. F-DI denoted factor-based diffusion index model and F-AR denotes factor-based VAR or AR models.

Step	Model	CPI - All			CPI - Core		
		GT	BN	LYB	GT	BN	LYB
1	F-DI	<b>0.126</b>	0.209	0.203	<b>0.111</b>	0.216	0.212
	F-AR	0.247	<b>0.218</b>	0.229	<b>0.159</b>	0.160	0.225
2	F-DI	<b>0.128</b>	0.206	0.188	<b>0.113</b>	0.214	0.218
	F-AR	0.244	<b>0.217</b>	0.228	<b>0.164</b>	0.165	0.224
3	F-DI	<b>0.135</b>	0.207	0.191	<b>0.113</b>	0.215	0.224
	F-AR	0.243	<b>0.219</b>	0.229	<b>0.167</b>	0.168	0.224

considered have equal predictive ability, and the alternative is that our proposed method performs better than the others in predicting the inflation indexes. Table 6 reports the testing results, where we use the method in Andrews (1991) to calculate the long-run covariance matrix (L-COV). In the table, GT–BN and GT–LYB denote the comparison between our approach and that in BN and LYB, respectively, and the test statistic is the difference between the forecast errors of the two methods involved, as illustrated in Diebold and Mariano (1995). From Table 6, we see that all  $p$ -values of the test statistics are small with most of them being less than or equal to 0.05, indicating that the factors extracted by our method have similar or better forecast ability than those of the other two methods in predicting the CPI indexes.

In conclusion, for U.S. monthly macroeconomic variables, the forecasting performance of the extracted factors using diffusion index models seems to favor the proposed approach, since the factors extracted by our method capture most of the dynamic dependencies of the data

Table 6: Testing for equal predictive ability of different factor-based diffusion index models, using the asymptotic test of Diebold and Mariano (1995). GT denotes the proposed method, BN denotes the principal component analysis as in Bai and Ng (2002), and LYB is the one in Lam et al. (2011). CPI - All denotes the Consumer Price Index for All Urban Consumers: All Items in U.S. City Average, and CPI - Core denotes the Consumer Price Index for All Urban Consumers: All Items Less Food and Energy in U.S. City Average.

Step		CPI - All		CPI - Core	
		GT-BN	GT-LYB	GT-BN	GT-LYB
1	L-COV	0.0026	0.0025	0.007	0.0023
	<i>p</i> -value	0.05	0.06	0.10	0.02
2	L-COV	0.0022	0.0025	0.001	0.002
	<i>p</i> -value	0.03	0.11	$\approx 0$	0.01
3	L-COV	0.002	0.002	0.001	0.002
	<i>p</i> -value	0.05	0.10	$\approx 0$	0.01

whereas those by the other methods may overlook some dynamic dependence. On the other hand, for the forecasting performance of the factor-based VAR or AR models, our method only fares better for the CPI-Core and it is not as good as the PCA approach for the CPI-All.

Finally, we state that the proposed factor model is different from those available in the literature such as those considered in Bai and Ng (2002). Therefore, the factors extracted by the proposed method are different from those by PCA and they have different interpretations. We believe that the PCA approach and the proposed one are complementary to each other rather than competitors, because they explore different aspects of the data as illustrated in the data analysis. The proposed method can be treated as another option in the toolbox for modeling high-dimensional time series and the extracted factors can be useful to practitioners who are interested in out-of-sample forecasting.

## 5 Discussion and Concluding Remarks

This article introduced an alternative factor model for high-dimensional time series analysis. We allow the largest eigenvalues of the covariance matrix of the idiosyncratic components to diverge to infinity by imposing some structure on the noise terms. The first step of the proposed analysis is an eigen-analysis of the matrix  $\widehat{\mathbf{M}}$  defined in Equation (2.9). The form of  $\widehat{\mathbf{M}}$  is a special case of the orthonormalized Partial Least Squares for time series data by assuming the covariance matrix of the data is identity. By an abuse of notation, let  $\mathbf{w}_t = (\mathbf{y}'_{t-1}, \dots, \mathbf{y}'_{t-k_0})'$  be the vector of past  $k_0$  lagged values of the time series  $\mathbf{y}_t$ , where  $k_0$  is a pre-specified positive



integer as that in (2.6) and (2.9). The orthonormalized Partial Least Squares computes the orthogonal score vectors for  $\mathbf{y}_t$  by solving the following optimization problem:

$$\max_{\mathbf{a}_i} \|E(\mathbf{a}'_i \mathbf{y}_t \mathbf{w}'_t)\|_2^2, \quad \text{subject to} \quad \mathbf{a}'_i E(\mathbf{y}_t \mathbf{y}'_t) \mathbf{a}_i = 1. \quad (5.1)$$

See, for example, Arenas-García and Camps-Valls (2008). It can be shown that the columns  $\mathbf{a}_i$  are given by the principal eigenvectors of the following generalized eigenvalue problem:

$$\boldsymbol{\Sigma}_{yw} \boldsymbol{\Sigma}'_{yw} \mathbf{a}_i = \eta \boldsymbol{\Sigma}_y \mathbf{a}_i. \quad (5.2)$$

Note that  $\mathbf{M} = \boldsymbol{\Sigma}_{yw} \boldsymbol{\Sigma}'_{yw}$ , which is just the form in (2.6). To solve the above equation, we need to obtain accurate estimates for the covariance matrix and its inverse simultaneously, which however is not easy. Instead we change the subject condition in (5.1) to  $\mathbf{a}'_i \mathbf{a}_i = 1$  and apply the eigen-analysis on  $\widehat{\mathbf{M}}$  in (2.9), and this approach remains an effective way if we assume the component variances of the data are uniformly bounded. In this case, the second step is needed.

The second step of the proposed analysis is the projected PCA on  $\widehat{\mathbf{S}}$  in (2.10) by assuming the largest  $K$  eigenvalues of the covariance matrix of the idiosyncratic component are diverging. In practice, one may assume that the largest eigenvalue is diverging whereas the remaining ones are bounded. Limited experience indicates that many real datasets exhibit such a phenomenon. If we are only concerned with the forecasting performance of the proposed analysis, we may select  $\widehat{K}$  in a range such as  $\widehat{K}_L \leq \widehat{K} \leq \widehat{K}_U$ , where  $\widehat{K}_L$  and  $\widehat{K}_U$  are defined in Remark 5(ii), via some cross-validation method like the out-of-sample testing.

The white noise test employed is an efficient way to determine the number of common factors for the proposed factor model. The one-by-one bottom-up testing procedure may not perform well when the dimension  $p$  is large. On the other hand, the limiting distribution of the test statistic of Tsay (2020) holds for large  $p$  by making use of the extreme value theory, and, hence, the test fares well for large  $p$ . If we like to use the test statistic for a wide range of dimensions and various sample sizes, we may employ empirical critical values via some simulation, which is possible because the test uses rank correlations and is robust to the underlying distribution of the data under the null hypothesis. The simulation results of Tsay (2020) show that the resulting test statistic works reasonably well.

## Supplementary Material

The supplementary material contains all technical proofs of the theorems in Section 3 and an additional real example consisting of half-hourly temperature data observed at the Adelaide Airport in Australia with  $p = 508$  and  $n = 336$ .

## Acknowledgments

We are grateful to the Editor, Associate Editor and the anonymous referees for their insightful comments and suggestions that have substantially improved the presentation and quality of the paper. This research is supported in part by the Booth School of Business, University of Chicago.

## References

- Ahn, S. C., and Horenstein, A. R. (2013). Eigenvalue ratio test for the number of factors. *Econometrica*, **81(3)**, 1203–1227.
- Anderson, T. W. (2003). *An Introduction to Multivariate Statistical Analysis*. Hoboken, NJ : John Wiley & Sons.
- Andrews, D. W. (1991). Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica*, **59**, 817–858.
- Arenas-García, J., and Camps-Valls, G. (2008). Efficient kernel orthonormalized PLS for remote sensing applications. *IEEE Transactions on Geoscience and Remote Sensing*, **46(10)**, 2872–2881.
- Bai J. (2003) Inferential theory for factor models of large dimensions. *Econometrica*, **71(1)**, 135–171.
- Bai, J., and Ng, S. (2002). Determining the number of factors in approximate factor models. *Econometrica*, **70**, 191–221.
- Boivin, J., and Ng, S. (2005). Understanding and comparing factor-based forecasts (No. w11285). *National Bureau of Economic Research*.
- Chang, J., Yao, Q., and Zhou, W. (2017). Testing for high-dimensional white noise using maximum cross-correlations. *Biometrika*, **104(1)**, 111–127.

- Davis, C., and Kahan, W. M. (1970). The rotation of eigenvectors by a perturbation. III. *SIAM Journal on Numerical Analysis*, **7(1)**, 1–46.
- Davis, R. A., Zang, P., and Zheng, T. (2012). Sparse vector autoregressive modelling. Available at *arXiv:1207.0520*.
- Diebold, F. X., and Mariano, R. S. (1995). Comparing predictive accuracy. *Journal of Business & Economic Statistics*, **13(3)**, 253–263.
- Fan, J., Liao, Y., and Mincheva, M. (2013). Large covariance estimation by thresholding principal orthogonal complements (with discussion). *Journal of the Royal Statistical Society, Series B*, **75(4)**, 603–680.
- Forni, M., Hallin, M., Lippi, M., and Reichlin, L. (2000a). Reference cycles: the NBER methodology revisited (No. 2400). Centre for Economic Policy Research.
- Forni, M., Hallin, M., Lippi, M., and Reichlin, L. (2000b). The generalized dynamic-factor model: Identification and estimation. *Review of Economics and statistics*, **82(4)**, 540–554.
- Forni, M., Hallin, M., Lippi, M., and Reichlin, L. (2005). The generalized dynamic factor model: one-sided estimation and forecasting. *Journal of the American Statistical Association*, **100(471)**, 830–840.
- Forni, M., Hallin, M., Lippi, M., and Zaffaroni, P. (2015). Dynamic factor models with infinite-dimensional factor spaces: one-sided representations. *Journal of Econometrics*, **185(2)**, 359–371.
- Gao, Z., Ma, Y., Wang, H., and Yao, Q. (2019). Banded spatio-temporal autoregressions. *Journal of Econometrics*, **208**, 211–230.
- Gao, Z., and Tsay, R. S. (2019). A structural-factor approach for modeling high-dimensional time series and space-time data. *Journal of Time Series Analysis*, **40**, 343–362.
- Gregory, A. W., and Head, A. C. (1999). Common and country-specific fluctuations in productivity, investment, and the current account. *Journal of Monetary Economics*, **44(3)**, 423–451.
- Hallin, M. and Lippi, M. (2013). Factor models in high-dimensional time series - A time-domain approach. *Stochastic Processes and Their Applications*, **123**, 2678–2695.
- Johnstone, I.M., and Lu, A.Y. (2009). On consistency and sparsity for principal components analysis in high dimensions. *Journal of the American Statistical Association*, **104(486)**, 682–693.

- Lam, C., and Yao, Q. (2012). Factor modeling for high-dimensional time series: inference for the number of factors. *The Annals of Statistics*, **40(2)**, 694–726.
- Lam, C., Yao, Q., and Bathia, N. (2011). Estimation of latent factors for high-dimensional time series. *Biometrika*, **98**, 901–918.
- Li, Z., Wang, Q., and Yao, J. (2017). Identifying the number of factors from singular values of a large sample auto-covariance matrix. *The Annals of Statistics*, **45(1)**, 257–288.
- Li, Z., Lam, C., Yao, J., and Yao, Q. (2019). On testing for high-dimensional white noise. *The Annals of Statistics* (to appear).
- Lütkepohl, H. (2006). *New Introduction to Multiple Time Series Analysis*, Springer, Berlin.
- Onatski, A. (2010). Determining the number of factors from empirical distribution of eigenvalues. *The Review of Economics and Statistics*, **92(4)**, 1004–1016.
- Onatski, A. (2012). Asymptotics of the principal components estimator of large factor models with weakly influential factors. *Journal of Econometrics*, **168(2)**, 244–258.
- Pan, J., and Yao, Q. (2008). Modelling multiple time series via common factors. *Biometrika*, **95(2)**, 365–379.
- Paul, D. (2007). Asymptotics of sample eigenstructure for a large dimensional spiked covariance model. *Statistica Sinica*, 1617–1642.
- Peña, D. and Yohai, V. J. (2016). Generalized dynamic principal components. *Journal of the American Statistical Association*, **111**, 1121–1131.
- Shen, D., Shen, H., and Marron, J. S. (2016). A general framework for consistency of principal component analysis. *Journal of Machine Learning Research*, **17(150)**, 1–34.
- Shojaie, A., and Michailidis, G. (2010). Discovering graphical Granger causality using the truncated lasso penalty. *Bioinformatics*, **26**, 517–523.
- Song, S., and Bickel, P. J. (2011). Large vector auto regressions. Available at [arXiv:1106.3519](https://arxiv.org/abs/1106.3519).
- Stock, J. H., and Watson, M. W. (1989). New indexes of coincident and leading economic indicators. *NBER Macroeconomics Annual*, **4**, 351–394.
- Stock, J. H., and Watson, M. W. (1998). Diffusion indexes. NBER Working Paper 6702.
- Stock, J. H., and Watson, M. W. (2002a), Forecasting using principal components from a large number of predictors. *Journal of the American Statistical Association*, **97**, 1167–1179.

- Stock, J. H., and Watson, M. W. (2002b). Macroeconomic forecasting using diffusion indexes. *Journal of Business and Economic Statistics*, **20(2)**, 147–162.
- Stock, J. H., and Watson, M. W. (2005). Implications of dynamic factor models for VAR analysis. Available at [www.nber.org/papers/w11467](http://www.nber.org/papers/w11467).
- Stock, J. H., and Watson, M. (2009). Forecasting in dynamic factor models subject to structural instability. *The Methodology and Practice of Econometrics. A Festschrift in Honour of David F. Hendry*, 173–205.
- Tiao, G. C., and Tsay, R. S. (1989). Model specification in multivariate time series (with discussion). *Journal of the Royal Statistical Society*, **B51**, 157–213.
- Tsay, R. S. (2014). *Multivariate Time Series Analysis*. Wiley, Hoboken, NJ.
- Tsay, R. S. (2020). Testing for serial correlations in high-dimensional time series via extreme value theory. *Journal of Econometrics*, **216(1)**, 106–117.
- Vershynin, R. (2012). How close is the sample covariance matrix to the actual covariance matrix?. *Journal of Theoretical Probability*, **25(3)**, 655–686.
- Vershynin, R. (2018). *High-Dimensional Probability: An Introduction with Applications in Data Science*. Cambridge University Press.
- Yu, Y., Wang, T., and Samworth, R. J. (2015). A useful variant of the Davis-Kahan theorem for statisticians. *Biometrika*, **102(2)**, 315–323.