

Predicting Extubation Readiness in Extreme Preterm Infants based on Patterns of Breathing*

Charles C. Onu¹, Lara J. Kanbar², Wissam Shalish³, Karen A. Brown⁴, Guilherme M. Sant'Anna³, Robert E. Kearney² and Doina Precup¹

Abstract—Extremely preterm infants commonly require intubation and invasive mechanical ventilation after birth. While the duration of mechanical ventilation should be minimized in order to avoid complications, extubation failure is associated with increases in morbidities and mortality. As part of a prospective observational study aimed at developing an accurate predictor of extubation readiness, Markov and semi-Markov chain models were applied to gain insight into the respiratory patterns of these infants, with more robust time-series modeling using semi-Markov models. This model revealed interesting similarities and differences between newborns who succeeded extubation and those who failed. The parameters of the model were further applied to predict extubation readiness via generative (joint likelihood) and discriminative (support vector machine) approaches. Results showed that up to 84% of infants who failed extubation could have been accurately identified prior to extubation.

I. INTRODUCTION

Due to immature lungs and respiratory control, extremely preterm infants (gestational age < 28 weeks) are at high risk of respiratory failure after birth. For that reason, most require endotracheal intubation and Invasive Mechanical Ventilation (MV) within the first hours or days of life [1]. As the respiratory status improves, MV is continuously weaned until the medical team deems that removal of the tube, termed *extubation*, can be performed. This is a difficult decision since if done too early, it could result in the need for re-intubation, a technically challenging intervention that has been associated with increased morbidities and mortality [2], [3]. On the other hand, delayed extubation, and hence prolonged mechanical ventilation, could increase the risk of complications. The most common complication is bronchopulmonary dysplasia (BPD), a form of chronic lung disease associated with serious long-term sequelae [4].

This project is part of an ongoing multicenter, multidisciplinary collaborative study aiming to develop an automated

prediction tool of extubation readiness using analysis of cardiorespiratory signals in extremely preterm infants (Clinicaltrials.gov identifier: NCT01909947) [5]. In the present work, we approach the task of predicting whether a patient would succeed or fail extubation based solely on respiratory patterns extracted from a 5-minute period of spontaneous breathing trial (SBT), while patient was still intubated but not receiving mechanical inflations from the ventilator.

Methods of generative modeling using Markov and semi-Markov chains were applied. The respiratory time series was processed as a sequence of five unique and mutually exclusive breathing patterns/states: Pause, Asynchronous breathing, Movement artifact, Synchronous breathing and Unknown, using an existing method of analysis of the respiratory signals called AUREA (Automated Unsupervised Respiratory Event Analysis) [6]. The models fitted to these states sequences revealed key similarities and differences between patients who succeeded and failed extubation. The characteristics were exploited in a support vector machine-based discriminative classifier and results indicated that our model could identify infants who went on failing extubation with up to 84% accuracy.

The rest of this paper is organized as follows. Section II reviews related work. Section III describes the data acquisition and preprocessing steps. Section IV describes our methodology. Section V presents results and provides a discussion on their significance, and in Section VI, we discuss future research directions.

II. RELATED WORK

The use of Markov chain models for understanding sequence data and making predictions about outcomes is popular across many domains, especially when the values of the time-series are drawn from a discrete set of states. Ye et al [7] applied Markov chain modeling to detect anomalous activities in computer networks. In another study by Gabriel et al [8], this method was successfully used to track rainfall occurrence patterns. More recently, Alinovi et al [9] applied semi-Markov chains to model the respiratory rate of infant patients experiencing breathing-related disorders such as apneas. In that study, respiratory rate was transformed from a real-valued time-series in to 5 levels by quantization. The authors demonstrated that semi-Markov chain models accurately described respiratory rate and simulated realistic sequences of respiratory rate in both normal and apneic infants.

*Research supported in part by the Canadian Institutes of Health Research. The work of L. Kanbar was supported in part by the Natural Sciences and Engineering Research Council of Canada. K. Brown was supported in part by the Queen Elizabeth Hospital of Montreal Foundation Chair in Pediatric Anesthesia.

¹C. C. Onu and D. Precup are with the School of Computer Science, McGill University, Montreal, QC H3A 2B4, Canada. (e-mail: charles.onu@mail.mcgill.ca; dprecup@cs.mcgill.ca)

²L. J. Kanbar and R. E. Kearney are with the department of Biomedical Engineering, McGill University, Montreal, QC H3A 2B4, Canada (e-mail: lara.kanbar@mail.mcgill.ca; robert.kearney@mcgill.ca)

³W. Shalish and G. M. Sant'Anna are with the division of Neonatology, McGill University Health Center, Montreal, QC H3A 2B4, Canada.

⁴K. Brown is with the department of Anesthesia, McGill University Health Center, Montreal, QC H3A 2B4, Canada.

Previous work in predicting extubation readiness of extreme preterm newborns have demonstrated that cardiorespiratory signals hold useful discriminating information between infants that succeed and those that fail extubation. In particular 2 prospective studies by Kaczmarek et al [10] and Robles-Rubio et al [11] on a small cohort of extremely preterm infants, extracted measures of heart rate variability (HRV) and respiratory variability (RV) from electrocardiogram (ECG) and respiratory inductive plethysmography (RIP) signals, respectively. Results revealed that measures of HRV and RV were excellent at detecting extubation successes (i.e. high specificity), but not as sensitive in detecting failures. In [12], using patients of the same cohort, a Support Vector Machine (SVM) was developed to predict extubation readiness directly from the combination of cardiorespiratory variability measures. This system achieved improved ability to detect extubation failures (sensitivity of 83%) while maintaining a fairly high specificity of 74%. It should however be noted that these works were carried out on fairly small datasets.

In this current work, we analyzed a larger dataset of 186 babies collected as part of a multi-institutional study. We explore the predictive potential, not of HRV or RV but, of respiratory pattern sequences of the infants. The current work: 1) empirically demonstrated that the semi-Markov modeling has a more robust capability for time-series data than the Markov chain models; 2) used semi-Markov chain models to understand the transition behavior of meaningful respiratory patterns representing actual respiratory states (not just quantized real-values); 3) showed that, in addition to generative classification via maximizing joint likelihood, the parameters of semi-Markov chains can be exploited in discriminative classifiers to improve performance. To the best of our knowledge, this is the first work that applies a time-series-based machine learning method to predict extubation readiness in extremely preterm infants.

III. DATA

A. Data Acquisition

Data from 186 infants was available for this study. Patients were enrolled and studied in five sites in Canada (Royal Victoria Hospital, Montreal Children’s Hospital, Jewish General Hospital) and the USA (Detroit Medical Center, MI, and Women and Infants Hospital of Rhode Island, RI). Ethical approval was obtained from each institutional board, and informed parental consent was obtained before recruitment.

Eligible infants were of birth weight $\leq 1250g$, and receiving MV at time of recruitment. Infants were excluded if they had any major congenital anomalies such as heart disease, or were receiving any vasopressor or sedative drugs at the time of extubation.

Respiratory signals were measured using RIP bands placed around the infant’s ribcage and abdomen. Signals were acquired for a 5-minute period of spontaneous breathing without any mechanical inflation from the ventilator, prior to extubation, at a sampling frequency of 1000Hz. Only infants which were then extubated were included in the dataset.




Fig. 1. Example of a RCG and ABD signal segment and the corresponding respiratory patterns (PAU, SYB, ASB, MVT, UNK) computed by AUREA.

Failure after extubation was defined as re-intubation within 7 days. More details on the data collection procedure for this study are available in [5].

B. Data Preprocessing

RIP signals sampled at 50Hz were analyzed using AUREA, which extracts sample-by-sample metrics of respiratory power, synchrony between the ribcage (RCG) and abdomen (ABD), and movement artifact [6]. AUREA uses k-means clustering with windowing and smoothing to assign each sample to one of the following respiratory patterns:

- Pause (PAU): A cessation of breathing.
- Synchronous Breathing (SYB): RCG and ABD are in phase.
- Asynchronous Breathing (ASB): RCG and ABD are out of phase.
- Movement Artifact (MVT): Associated with infant moving or nurse handling.
- Unknown (UNK): Ambiguous patterns not belonging to any other pattern category.

AUREA provides repeatable results with no human intervention [6]. An example of RIP signals and corresponding patterns assigned by AUREA to the different samples is shown in Fig. 1.

IV. METHODS

We applied Markov chains and Semi-Markov chains to model the time series of respiratory patterns, and then used SVM with features extracted from these chains to predict extubation readiness. We now describe the analysis methods.

A. Discrete-time Markov Chain

A Markov chain is a probabilistic graphical model in which every node in the chain is only dependent on the one preceding it. Time-series data can be represented as a Markov chain whereby the nodes of the chain are the values observed at every time step. We use discrete-time finite Markov chains, in which at time t , state x_t takes a value from a finite set of states S (in our case, the 5 respiratory patterns provided by AUREA).

A Markov chain model has 2 sets of parameters: the probability distribution over initial states (a vector) π and the

transition probabilities between states (a matrix) A . Fitting or learning the model of a Markov chain involves estimating these parameters from data. Their maximum likelihood estimates are given by [13]:

$$\pi_j = \frac{\# \text{ of sequences starting in } j}{\# \text{ of sequences}} \quad \forall j \in S \quad (1)$$

$$A_{i,j} = \frac{n_{ij}}{\sum_j n_{ij}} \quad \forall i, j \in S \quad (2)$$

where n_{ij} is the number of time steps in which a transition from state i to j occurred. Given a time-series of observations x_1, x_2, \dots, x_T , the joint likelihood of the sequence according to the Markov chain is given by:

$$P(x) = P(x_1) \prod_{t=2}^T P(x_t|x_{t-1}) = \pi_{x_1} \prod_{t=2}^T A_{x_{t-1}, x_t} \quad (3)$$

Note that in practice, the start state distribution π was not included in our model, due to an adaptation or transition phase from MV with inflations to ETT-CPAP without inflations.

In order to apply Markov chain models for classification, separate transition models A^s , A^f were first fit to the data coming from the success and failure patients, respectively. The classification for a new sequence x is done by computing its posterior likelihood with respect to both models (using Eq.3), and selecting the class, c whose model gives the higher likelihood:

$$\arg \max_c L(x|A^c) \quad (4)$$

where

$$L(x|A^c) = \prod_{t=2}^T A_{x_{t-1}, x_t}^c \quad (5)$$

B. Discrete-time Semi-Markov Chain

A Semi-Markov chain model, on the other hand, is characterised by 3 parameters: a start state distribution vector π ; the transition matrix A , which stores only cross-state transition probabilities (diagonal elements are 0); and a set of *dwell or sojourn time distributions* F , which model the duration spent in each state until a transition out of that state occurs.

This framework was useful in our application for several reasons. First, Markov chains implicitly model dwell times as an exponential distribution [14] which could introduce bias into the model if underlying data is not actually exponential. Secondly, in data characterised by very long dwell times, the transition probabilities of cross-state transitions (off-diagonal elements) go to 0, making it very difficult to get any useful information from the model. Finally, a Markov chain is highly susceptible to changes in the sampling rate of the data. Semi-Markov chains address all of these issues.

The joint likelihood of a sequence of observations under a semi-Markov chain is given by:

$$P(x) = \pi_{x_1} \prod_{t=2}^T A_{x_{t-1}, x_t} F_{x_t}(|x_t|) \quad (6)$$

where $F_{x_t}(|x_t|)$ is the probability of sojourning in the state x_t for the duration $|x_t|$. [15]

In modeling the infant respiratory pattern sequences as a semi-Markov chain, A was learned as before. To fit the dwell time distributions, all dwell times in a state (e.g., PAU) for a particular population (success or failure) were obtained. Several known probability distributions were fit to this data using MATLAB [16]. The distribution which minimized the Bayesian Information Criterion (BIC) [17] was selected. These steps were repeated for all states in both success and failure groups to obtain 10 separate dwell time distributions. Classification of a new example sequence as success or failure was done as described in the previous section by selecting the class of larger posterior likelihood.

C. Support Vector Machine (SVM)

Using the Markov model likelihood for classification can be sub-optimal if the model structure or some of the model parameters are imprecise. Discriminative models do not make probabilistic assumptions about how the inputs were generated, but rather attempt to learn a (linear or non-linear) boundary between the groups. Support vector machines (SVMs), in particular, learn a maximum margin decision boundary [18]. In order to compare our results to the Markov chain case, we derived summary statistics from the respiratory pattern sequence of each patient and used these as inputs to train an SVM. In particular, a radial basis function (RBF) SVM was used. The hyperparameters of the RBF SVM - box constraint, C (which penalises the error function to manage overfitting) and kernel scale γ (which controls the width of the Gaussian) - were optimized using leave-one-out cross-validation and the balanced loss metric, as motivated in section IV-E.

D. Symmetric KL Divergence

The Kullback-Leibler (KL) divergence [19] is a measure of how well a distribution Q is approximating another




Fig. 2. Percentage of time spent in each respiratory state by success and failure patients during a 5-min period of spontaneous breathing prior to extubation

TABLE I

THE TYPE AND PARAMETERS OF THE DISTRIBUTIONS OF BEST FIT TO THE DWELL (OR SOJOURN) TIMES IN EACH RESPIRATORY PATTERN FOR SUCCESS AND FAILURE PATIENTS.

Pause	Success	Failure
	Exponential $\mu = 2.51$	Exponential $\mu = 2.94$
Asynchrony	GeneralizedExtremeValue $k=0.63, \sigma = 1.30, \mu=1.85$	GeneralizedExtremeValue $k=0.65, \sigma = 1.36, \mu=1.81$
Movement	GeneralizedPareto $k=-0.22, \sigma = 3.62$	GeneralizedPareto $k=-0.11, \sigma = 3.31$
Synchrony	InverseGaussian $\mu =8.61, \lambda = 3.61$	InverseGaussian $\mu =7.83, \lambda = 3.41$
Unknown	GeneralizedPareto $k=-0.07, \sigma = 2.07$	GeneralizedPareto $k=-0.10, \sigma = 2.05$

distribution P . It is defined as:

$$D_{KL}(P||Q) = \sum_n P_n \log \frac{P_n}{Q_n} \quad (7)$$

The KL-divergence is non-symmetric: $D_{KL}(P||Q) \neq D_{KL}(Q||P)$, which is not desirable in our application. Hence, we use symmetrized KL-divergence to compare distributions over transitions between patterns:

$$D_{KLS}(P||Q) = D_{KL}(P||Q) + D_{KL}(Q||P) \quad (8)$$

$$D_{KLS}(P||Q) = \sum_n (P_n - Q_n) \log \frac{P_n}{Q_n} \quad (9)$$

E. Model Evaluation

Due to the small size of our dataset and class imbalance, it was necessary to give special thought to the choice of evaluation method. First, due to the relatively small number of examples, we used leave-one-out cross-validation instead of k-Fold cross-validation. Though computationally more expensive, leave-one-out is a better estimator for small datasets [20].

Secondly, the class imbalance meant that optimizing for classification accuracy could result in sub-optimal (or even degenerate) models, because it implicitly assigns higher weights to examples of the majority class. To address this, the true positive rate (sensitivity) and true negative rate (specificity) were tracked separately. As a single evaluation metric for choosing one model over another, we use the *balanced misclassification loss* (see Appendix A).

V. RESULTS

Of the 186 patients recruited, 136 succeeded extubation and 50 failed. We discuss the results of our experiments in the following sub-sections.

A. Analysis of Respiratory State Durations

As a starting point, the total time spent in each breathing pattern was obtained for the two groups of patients, shown in Fig. 2. It was normalized by the total duration of spontaneous breathing to account for the difference in

number of examples in both classes. In the 2 groups, SYB was the dominant state at least 50% of the time, with the success patients spending slightly more time in this state. The failure patients on the other hand spent a greater fraction of time in the PAU state. The time spent in the other states were essentially equivalent as confirmed by standard error estimates obtained by bootstrapping, shown in Fig. 2.

B. Modeling of Dwell Times

The optimal dwell time distributions in each state were estimated for both groups (as described in section IV-B) and are summarized in Table I. The detailed plots of the probability density functions (PDF) are shown in Appendix C. It was interesting to observe that for each pattern, the *distribution type* which best fit the dwell times in both populations were same. We suspect that this an indication of some underlying consistency in breathing behaviour of premature infants in spite of extubation outcome. It should also be noted that the dwell time was distributed exponentially only in the PAU state. As discussed earlier, the use of a Markov chain model would have implicitly taken an exponential distribution for all states. The use of semi-Markov framework has thus allowed for a more expressive and accurate representation.

C. Modeling Transitions

The respiratory pattern sequences were modeled as semi-Markov chains (Tables II and III). Each cell in the matrix represents the probability of transitioning from the state labeled on the row to that on the column. It was observed that the most probable transition given ASB and SYB (the breathing states) was the same in both infants who succeeded and those who failed extubation (shown in bold, black font). However, it differed for PAU and MVT states (shown in bold, red font). This suggests that in terms of transition behaviour, the 2 groups of infants differ more in the transitions emanating from non-breathing states than from breathing states.

Further, the symmetric KL divergence (D_{KLS}) between the 2 transition matrices for the semi-Markov model was

TABLE II

RESPIRATORY STATE TRANSITION PROBABILITIES FOR THE SUCCESS POPULATION MODELED AS A SEMI-MARKOV CHAIN.

	PAU	ASB	MVT	SYB	UNK
PAU	0	0.27	0.09	0.26	0.38
ASB	0.10	0	0.16	0.29	0.45
MVT	0.12	0.32	0	0.43	0.14
SYB	0.06	0.25	0.15	0	0.54
UNK	0.13	0.28	0.04	0.55	0

TABLE III

RESPIRATORY STATE TRANSITION PROBABILITIES FOR FAILURE POPULATION MODELED AS SEMI-MARKOV CHAIN.

	PAU	ASB	MVT	SYB	UNK
PAU	0	0.28	0.06	0.39	0.28
ASB	0.12	0	0.21	0.28	0.40
MVT	0.17	0.41	0	0.32	0.09
SYB	0.14	0.21	0.14	0	0.52
UNK	0.15	0.30	0.03	0.52	0

estimated as 0.27. When modeled as a Markov chain (see details in Appendix B), the probabilities of cross-pattern transitions and the D_{KLS} tended towards zero (0.0019), indicating that the learned models were almost identical due to relatively long dwell times. The semi-Markov transition matrix, on the other hand, being invariant to sampling frequency changes and/or length of dwell time, resulted to better numerical resolution of cross-pattern transitions and the learning of more discriminating characteristics between both groups of infants.

D. Generative Classification with Semi-Markov Chain Models

The classification performance of the learned semi-Markov transition models was evaluated using leave-one-out cross validation. As described in section IV-B, the likelihoods of each test example were computed using its entire sequence of transitions (Lk-ALL). Failure patients were identified at a rate (sensitivity) of 50% while specificity was 73%.

The predictive value of the individual patterns/states was further evaluated. In particular, to compute the likelihood of a test sequence based on one state, the product of cross-state transitions emanating from only that state are taken. As before, this likelihood is computed with respect to the transition models for the 2 classes, and a prediction is made by selecting the class whose model gave higher likelihood. Results are shown accordingly in Table IV where *Lk-STATE* represents prediction made using likelihood of the "STATE" specified. The highest sensitivity of 68% was obtained by the PAU, MVT, SYB states. Overall, the PAU pattern gave the lowest loss of 0.37.

E. Discriminative Classification with SVM

The following features, motivated by the semi-Markov chain model, were extracted from each subject and applied in an SVM classifier.

- Total dwell time in each respiratory pattern as a fraction of the total sequence duration, (*Dw-All*) - 5 features
- Number of transitions from pattern i to pattern j (where $i \neq j$) as a fraction of the total dwell time in pattern i , $\forall i \in S$, (*Tr-All*) - 20 features
- Number of occurrences of each respiratory pattern as a fraction of the number of occurrences of all patterns, (*Oc-All*) - 5 features

The features were first used individually (*Dw-ALL*, *Oc-ALL*, *Tr-ALL*) and then as an ensemble (*Dw-Oc-Tr-ALL*) of 30 features. Similar to the generative case, the predictive value of each individual pattern/state was evaluated - for each state, the dwell time in that state, *Dw-STATE* (1 feature), the cross-transitions, *Tr-STATE* (4 features) and the occurrence count, *Oc-STATE* (1 feature) were combined (*Dw-Oc-Tr-STATE*) to train the classifier. We applied SVMs with RBF/Gaussian kernels. Leave-one-out cross-validation was used in a grid search to find the best pair of hyper-parameter (box constraint C and kernel scale γ) values based on the balanced misclassification loss. This grid search was repeated

for each feature set since each would have different optimal values of C and γ .

When using features for all patterns, the sensitivity and specificity at the optimal hyper-parameter setting were 64% and 63%, respectively. The highest sensitivity of 84% was obtained when using features of only the Pause pattern, which also gave the lowest loss of 0.31. It could also be observed that whereas PAU and MVT patterns gave higher sensitivities, SYB and ASB gave higher specificities. This is likely an indication that the Pause and Movement patterns characterise better patients who may fail extubation while the breathing patterns better characterise patients who succeed. All results are summarised in Table IV. In Fig. 3, we show the receiver-operating characteristic (ROC) curve for the 2 SVM models which gave the lowest loss. ROC was obtained by fixing C at the optimal value and varying γ .

VI. DISCUSSION

We demonstrated the practical application of semi-Markov chains for modeling and classification of respiratory pattern behaviour of preterm infants in the immediate period prior to extubation. We showed that semi-Markov chain models provide more expressive and robust details about the underlying time series compared to Markov chain models. In terms of dwell time behaviour, the model revealed consistency between the success and failure groups in all respiratory states. Differences were highlighted primarily in transition behaviour arising from the Pause and Movement Artifact patterns. Prediction results confirmed that these 2 patterns provide more discriminating information (especially for patients who failed extubation) than any other pattern.

Indeed, using only features from the Pause pattern led to the greatest detection of failed extubations (with a sensitivity of 84%), but failed to recognizing nearly half of

TABLE IV
PERFORMANCE OF CLASSIFICATION METHODS. LK-ALL OR LK-STATE REFERS TO CLASSIFICATION USING LIKELIHOOD OF CHAIN CONSIDERING ALL STATES OR A SPECIFIED "STATE". DW, OC, TR REFER TO FEATURES EXTRACTED BASED ON DWELL TIME, OCCURRENCE COUNT AND TRANSITIONS IN STATES.

Approach	Sensitivity	Specificity	Loss
Generative (Semi-Markov)			
Lk-ALL	0.50	0.73	0.38
Lk-PAU	0.68	0.58	0.37
Lk-ASB	0.63	0.48	0.45
Lk-MVT	0.68	0.50	0.41
Lk-SYB	0.68	0.53	0.40
Lk-UNK	0.61	0.44	0.48
Discriminative (SVM)			
Dw-ALL	0.18	0.93	0.45
Oc-ALL	0.44	0.69	0.44
Tr-ALL	0.82	0.41	0.39
Dw-Oc-Tr-ALL	0.64	0.63	0.37
Dw-Oc-Tr-PAU	0.84	0.54	0.31
Dw-Oc-Tr-ASB	0.38	0.75	0.44
Dw-Oc-Tr-MVT	0.60	0.58	0.41
Dw-Oc-Tr-SYB	0.26	0.81	0.46
Dw-Oc-Tr-UNK	0.62	0.43	0.48




Fig. 3. Receiver-operating characteristic (ROC) curve for support vector machine trained on summary features of all respiratory patterns (AUC=0.62) and on summary features of only the Pause (PAU) pattern (AUC=0.70)

the successful extubations (specificity of 54%). This means that at the time infants were deemed ready for extubation, 8 out of 10 failures would be prevented, but at the expense of unnecessarily prolonging the course of mechanical ventilation in 5 out of 10 successes. Previous work [12] which applied SVM (on a much smaller cohort) using cardiorespiratory variables gave comparable sensitivity but a higher specificity of 74%. It will be necessary to repeat those experiments with this larger dataset. Another viable path for future work is in training a mixture of experts model (MEM) that classifies failure patients based on Pause and Movement states, and success patients on the breathing states. The use of automatically extracted respiratory patterns for prediction of extubation readiness provides an approach that unveils intuition and enhances interpretable models. We emphasize that all babies used in this study were deemed ready for extubation by an attending clinician, so these results constitute an improvement in detecting problem cases over current practice. The advantage of using an automated approach is that we can provide a quantified, repeatable and precise analysis breathing patterns to support clinical decisions.

REFERENCES

- [1] B. J. Stoll, N. I. Hansen, E. F. Bell, M. C. Walsh, W. A. Carlo, S. Shankaran, A. R. Lupton, P. J. Sánchez, K. P. Van Meurs, M. Wyckoff *et al.*, "Trends in care practices, morbidity, and mortality of extremely preterm neonates, 1993-2012," *JAMA*, vol. 314, no. 10, pp. 1039-1051, 2015.
- [2] B. J. Manley, L. W. Doyle, L. S. Owen, and P. G. Davis, "Extubating extremely preterm infants: Predictors of success and outcomes following failure," *J Pediatr*, vol. 173, pp. 45-49, 2016.
- [3] S. Chawla, G. Natarajan, S. Shankaran, B. Carper, L. P. Brion, M. Keszler, W. A. Carlo, N. Ambalavanan, M. G. Gantz, A. Das *et al.*, "Markers of successful extubation in extremely preterm infants, and morbidity after failed extubation," *The Journal of Pediatrics*, 2017.
- [4] M. C. Walsh, B. H. Morris, L. A. Wraga, B. R. Vohr, W. K. Poole, J. E. Tyson, L. L. Wright, R. A. Ehrenkranz, B. J. Stoll, A. A. Fanaroff *et al.*, "Extremely low birthweight neonates with protracted ventilation: mortality and 18-month neurodevelopmental outcomes," *The Journal of pediatrics*, vol. 146, no. 6, pp. 798-804, 2005.

- [5] W. Shalish, L. J. Kanbar, S. Rao, C. A. Robles-Rubio, L. Kovacs, S. Chawla, M. Keszler, D. Precup, K. Brown, R. E. Kearney, and G. M. Sant'Anna, "Prediction of extubation readiness in extremely preterm infants by the automated analysis of cardiorespiratory behavior: study protocol," *BMC Pediatrics*, vol. 17, no. 1, p. 167, Jul 2017. [Online]. Available: <https://doi.org/10.1186/s12887-017-0911-z>
- [6] C. A. Robles-Rubio, K. A. Brown, and R. E. Kearney, "Automated unsupervised respiratory event analysis," in *2011 Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, Aug 2011, pp. 3201-3204.
- [7] N. Ye *et al.*, "A markov chain model of temporal behavior for anomaly detection," in *Proceedings of the 2000 IEEE Systems, Man, and Cybernetics Information Assurance and Security Workshop*, vol. 166. West Point, NY, 2000, p. 169.
- [8] K. Gabriel and J. Neumann, "A markov chain model for daily rainfall occurrence at tel aviv," *Quarterly Journal of the Royal Meteorological Society*, vol. 88, no. 375, pp. 90-95, 1962.
- [9] D. Alinovi, G. Ferrari, F. Pisani, and R. Raheli, "Markov chain modeling and simulation of breathing patterns," *Biomedical Signal Processing and Control*, vol. 33, pp. 245 - 254, 2017.
- [10] J. Kaczmarek, S. Chawla, C. Marchica, M. Dwaihy, L. Grundy, and G. M. Sant'Anna, "Heart rate variability and extubation readiness in extremely preterm infants," *Neonatology*, vol. 104, no. 1, pp. 42-48, 2013.
- [11] C. A. Robles-Rubio, J. Kaczmarek, S. Chawla, L. Kovacs, K. A. Brown, R. E. Kearney, and G. M. Sant Anna, "Automated analysis of respiratory behavior in extremely preterm infants and extubation readiness," *Pediatric Pulmonology*, vol. 50, no. 5, pp. 479-486, 2015. [Online]. Available: <http://dx.doi.org/10.1002/ppul.23151>
- [12] D. Precup, C. A. Robles-Rubio, K. A. Brown, L. Kanbar, J. Kaczmarek, S. Chawla, G. M. Sant'Anna, and R. E. Kearney, "Prediction of extubation readiness in extreme preterm infants based on measures of cardiorespiratory variability," in *2012 Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, Aug 2012, pp. 5630-5633.
- [13] V. Barbu and N. Limnios, *Semi-Markov Chains and Hidden Semi-Markov Models Toward Applications: Their Use in Reliability and DNA Analysis*, 1st ed. Springer Publishing Company, Incorporated, 2008.
- [14] C. M. Bishop, *Pattern recognition and machine learning*. Springer, 2006.
- [15] M. L. Puterman, *Markov decision processes: discrete stochastic dynamic programming*. John Wiley & Sons, 1994.
- [16] M. Sheppard, "Fit all valid parametric probability distributions to data (<https://www.mathworks.com/matlabcentral/fileexchange/34943>), retrieved january 18, 2017, matlab central file exchange." 2012.
- [17] G. Schwarz *et al.*, "Estimating the dimension of a model," *The annals of statistics*, vol. 6, no. 2, pp. 461-464, 1978.
- [18] C. Cortes and V. Vapnik, "Support-vector networks," *Machine learning*, vol. 20, no. 3, pp. 273-297, 1995.
- [19] S. Kullback, *Information theory and statistics*. Courier Corporation, 1997.
- [20] T.-T. Wong, "Performance evaluation of classification algorithms by k-fold and leave-one-out cross validation," *Pattern Recognition*, vol. 48, no. 9, pp. 2839-2846, 2015.

A. *Balanced Misclassification Loss*

Accuracy is defined as:

$$acc = \frac{tp + tn}{tp + fn + tn + fp} \quad (10)$$

$$= \frac{tp + tn}{p + n} = \frac{tp}{p + n} + \frac{tn}{p + n} \quad (11)$$

$$= \frac{tp}{p} \left(\frac{p}{p + n} \right) + \frac{tn}{n} \left(\frac{n}{p + n} \right) \quad (12)$$

$$= sens. \left(\frac{p}{p + n} \right) + spec. \left(\frac{n}{p + n} \right) \quad (13)$$

where tp , tn , fp and fn are respectively the number of true positives, true negatives, false positives and false negatives; p is the number of positive examples and n the number of negative examples. $sens$ is sensitivity and $spec$ is specificity.

This means that, if the class proportions are 75:25, a classifier which simply predicts the majority class would have a misleading accuracy of 75%. The *balanced accuracy* measure, acc_b , corresponds to the average of the sensitivity and specificity measures:

$$acc_b = sensitivity * 0.5 + specificity * 0.5 \quad (14)$$

 B. *Transition Matrices for Fitted Markov chain Model*

The Markov chain transition matrices for the success and failure populations are shown in Tables V and VI, respectively. It can be seen that due to extremely long dwell times in states compared to cross-state transitions, the diagonal elements (self-transitions) account for nearly all of the probability on each row.

TABLE V
RESPIRATORY STATE TRANSITION PROBABILITIES FOR THE SUCCESS
POPULATION MODELED AS A MARKOV CHAIN

	PAU	ASB	MVT	SYB	UNK
PAU	0.9936	0.0019	0.0004	0.0022	0.0018
ASB	0.0006	0.9953	0.0010	0.0013	0.0018
MVT	0.0012	0.0029	0.9931	0.0020	0.0007
SYB	0.0003	0.0005	0.0003	0.9977	0.0012
UNK	0.0015	0.0031	0.0003	0.0055	0.9895

TABLE VI
RESPIRATORY STATE TRANSITION PROBABILITIES FOR FAILURE
POPULATION MODELED AS MARKOV CHAIN

	PAU	ASB	MVT	SYB	UNK
PAU	0.9920	0.0022	0.0007	0.0020	0.0032
ASB	0.0005	0.9955	0.0007	0.0013	0.0020
MVT	0.0007	0.0021	0.9934	0.0028	0.0010
SYB	0.0001	0.0005	0.0003	0.9978	0.0012
UNK	0.0013	0.0027	0.0004	0.0056	0.9899

 C. *Dwell/Sojourn Time Distribution Fits*

Plots of the probability density functions (PDF) of sojourn times in all states are shown in Fig 4, as well as the distributions of best fit based on the bayesian information criterion (BIC).




Fig 4: Probability Density Functions (PDF) of Dwell Time Distributions in all 5 respiratory patterns for success and failure patients