Premetric teleparallel theory of gravity and its local and linear constitutive law

Yakov Itin^{1,*}, Yuri N. Obukhov^{2,}, Jens Boos^{3,*}, Friedrich W. Hehl^{4,†}

¹The Hebrew University and Jerusalem College of Technology, Jerusalem 91160, Israel

²Institute for Nuclear Safety, Russian Academy of Sciences, 115191 Moscow, Russia

³Theor. Phys. Inst., Univ. Alberta, Edmonton, AB T6G 2E1, Canada

⁴Inst. Theor. Physics, Univ. of Cologne, 50923 Köln, Germany

*itin@math.huji.ac.il
 *obukhov@ibrae.ac.ru
 *boos@ualberta.ca
 *hehl@thp.uni-koeln.de

file teleGRchi43arXiv.tex, 06 Nov 2018

Abstract

We continue to investigate the premetric teleparallel theory of gravity (TG) with the coframe (tetrad) as gravitational potential. We start from the field equations and a local and linear constitutive law. We create a Tonti diagram of TG in order to disclose the structure of TG. Subsequently we irreducibly decompose the 6th order constitutive tensor under the linear group. Moreover, we construct the most general constitutive tensors from the metric and the totally antisymmetric Levi-Civita symbol, and we demonstrate that they encompass nontrivial axion and skewon type pieces. Using these tools, we derive for TG in the geometric-optics approximation propagating massless spin 0, 1, and 2 waves, including the special case of Einstein's general relativity.

Contents

1 Introduction						
	1.1	Field equations	4			
	1.2	Local and linear constitutive law	4			
	1.3	Reversibility	,			
2	2 Tonti diagram of the premetric teleparallel theory of gravity					

3	Decompositions of the constitutive tensor	13				
	3.1 Two forms of the constitutive tensor	13				
	3.2 $S_4 \times S_2$ decomposition of $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$	13				
	3.3 S_6 decomposition of $\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}$	16				
	3.4 Relation between the decompositions	18				
	3.5 Reversibility	19				
	3.6 Lagrangian	20				
	3.7 Parametrization of irreducible parts	21				
	3.8 Energy-momentum current	22				
	3.9 Contracting $\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$ twice: four second order tensors	22				
4	Metric-dependent constitutive tensor					
	4.1 A metric-dependent ansatz with parity conserving terms	25				
	4.2 Parity violating terms	27				
	4.3 The general case and its irreducible decomposition	31				
	4.4 Specific Lagrangians.	32				
	4.4.1 GR_{\parallel} : the teleparallel equivalent of GR	32				
	4.4.2 Von der Heyde Lagrangian	32				
	4.4.3 Torsion-square Lagrangian in nonlocal gravity (NLG)	33				
5	Propagation of gravitational waves	33				
	5.1 Geometric optics approximation	33				
	5.2 Dispersion relation—general facts	34				
	5.3 Dispersion relation decomposed	36				
	5.4 Scalar waves as a special case	37				
	5.5 A separable case	37				
	5.6 Gauge conditions	38				
6	Gravitational waves in a metric model					
	6.1 Characteristic system	38				
	6.2 Waves in $GR_{ }$, the teleparallel equivalent of GR	40				
7	Conclusions and outlook	41				
A	Young decomposition	42				
Re	References					

1 Introduction

Lately we followed in [18] the program of Kottler of 1922 to remove the metric tensor of spacetime, the gravitational potential within general relativity theory (GR), from the fundamental laws of classical electromagnetism and gravity as far as possible. In particular, we applied this to the theory of gravity [31] in that we started from a translational gauge theory of gravity, also known as teleparallel theory of gravity (TG)¹. We assume that our readers are familiar with [31]. The TG approach is reviewed in Blagojević et al. [8], see also Maluf [36], and Aldrovandi & Pereira [1]. For a somewhat related tetrad approach to gravity, in which affine symmetry is exploited, one should compare Sławianowski et al. [54].

The spacetime geometry of TG is represented by a four-dimensional (4d) manifold equipped with a coframe 1-form $\vartheta^{\alpha} = e_i^{\alpha} dx^i$ and with a linear connection 1-form $\Gamma_{\alpha}{}^{\beta} = \Gamma_{i\alpha}{}^{\beta} dx^i$, the curvature 2-form of which, $R_{\alpha}{}^{\beta} = \frac{1}{2}R_{ij\alpha}{}^{\beta} dx^i \wedge dx^j$, vanishes globally: $R_{\alpha}{}^{\beta} \equiv 0$. In the notation we follow basically the book [22]: α and β are co- and contravariant frame indices, with $\alpha, \beta, ... = \hat{0}, \hat{1}, \hat{2}, \hat{3}$, whereas i, j, ... = 0, 1, 2, 3 are coordinate indices.

Because of the vanishing curvature, we can pick suitable frames such that $\Gamma_{\alpha}{}^{\beta}$ vanishes globally:

$$\Gamma_{\alpha}{}^{\beta} \stackrel{*}{=} 0$$
 (everywhere in spacetime). (1)

In this 'teleparallel gauge,' the covariant exterior derivative taken with respect to the connection $\Gamma_{\alpha}{}^{\beta}$ is reduced to the ordinary exterior derivative. This will simplify our formalism. However, we will drop the star * over the equality sign in future since (1) is assumed to be valid throughout our paper.

The essence of the premetric approach can be formulated as follows. This universal field-theoretic scheme is based on conservation laws which hold true for the two types of variables: extensive fields ("how much?") and intensive ("how strong?") ones. These variables satisfy the *fundamental equaions* which are metric-free, whereas the metric comes in only via the *linking equations* which establish consitutive relations between the extensive and intensive variables.

Premetric electrodynamics [22] is based on the conservation laws of electric charge and magnetic flux which give rise to the fundamental equations dH = J and dF = 0. Here H is the electromagnetic excitation 2-form (extensive variable) and F the electromagnetic field strength 2-form (intensive variable). By introducing the constitutive relation $H = \kappa[F]$, one obtains a predictive physical theory.

The premetric gravity framework [31] can be constructed along the same lines by replacing the electric charge with a "gravitational charge" \rightarrow mass \rightarrow energy-

¹For the application of teleparallelism in continuum mechanics, see Delphenich [12].

momentum. In this introductory section, we provide a short overview of the premetric teleparallel approach.

1.1 Field equations

The field equations of TG will be our starting point. The inhomogeneous gravitational field equation of TG reads [31, Table I]

$$dH_{\alpha} = {}^{(\vartheta)}\Sigma_{\alpha} + {}^{(m)}\Sigma_{\alpha} \qquad (4 \times 6 \text{ components}). \tag{2}$$

It relates the twisted gravitational excitation 2-form

$$H_{\alpha} = \frac{1}{2} H_{ij\alpha} dx^{i} \wedge dx^{j} = \frac{1}{2} H_{\beta\gamma\alpha} \vartheta^{\beta\gamma} = \frac{1}{2} \check{H}^{\beta\gamma}{}_{\alpha} \varepsilon_{\beta\gamma}$$
(3)

to its source, the sum of the twisted energy-momentum 3-forms of gravity ${}^{(\vartheta)}\Sigma_{\alpha}$ and of matter ${}^{(m)}\Sigma_{\alpha}$, respectively. Here $\vartheta^{\beta\gamma} := \vartheta^{\beta} \wedge \vartheta^{\gamma}$ and $\varepsilon_{\beta\gamma} := \frac{1}{2} \varepsilon_{\beta\gamma\delta\varepsilon} \vartheta^{\delta} \wedge \vartheta^{\varepsilon}$, see [22, p.39].

As was already pointed out in [17, p.52], "...for consistency, we cannot allow spinning matter (other than as test particles) in such a T₄..." that is, in a 4d Weitzenböck spacetime. In other words, the field equation (2) is valid only for hydrodynamic and for electromagnetic matter. A careful proof of this stipulation was given by Obukhov & Pereira² [47]. For matter *with spin*, the Dirac field, for example, the Lorentz group should also be gauged, which removes the teleparallelism constraint $R_{\alpha}{}^{\beta} = 0$. Then one arrives at a Poincaré gauge theory operating in a Riemann-Cartan spacetime with torsion and with Cartan curvature $R_{\alpha}{}^{\beta} \neq 0$.

The homogeneous field equation of TG reads

$$dF^{\alpha} = 0 \qquad (4 \times 6 \text{ components}), \qquad (4)$$

with the untwisted vector-valued 2-form $F^{\alpha} := d\vartheta^{\alpha}$, the torsion of spacetime, which has the expansion

$$F^{\alpha} = \frac{1}{2} F_{ij}{}^{\alpha} dx^{i} \wedge dx^{j} = \frac{1}{2} F_{\beta\gamma}{}^{\alpha} \vartheta^{\beta\gamma}.$$
 (5)

Note that (4) is the first Bianchi identity of a linearly connected spacetime with vanishing curvature, see [53, Eq.(5.41)] or [22, Eq.(C.1.67)].

The analogy to the Maxwell equations of electrodynamics should be apparent, see [22]. Eq.(2) represents four inhomogeneous Maxwell type equations and Eq.(4) four homogeneous Maxwell type equations. Since TG is a translational gauge theory, it has the four one-form potentials $\vartheta^{\alpha} = e_i^{\alpha} dx^i$, where four is the number of generators of the translation group.

²Within the framework of TG as a translational gauge theory, the coupling to matter is achieved via the minimal coupling procedure, strictly in the sense of a bona fide gauge theory. Dispensing with the minimal coupling principle (as Maluf [35] does, e.g.) is against the spirit of gauge theory.

1.2 Local and linear constitutive law

In order to complete the two field equations of TG to a predictive system of equations, one has to adopt a constitutive law between excitation H and field strength F. The simplest assumption is that the functional $H = \kappa[F]$ is local and linear,

$$H_{\alpha} = \boldsymbol{\kappa}_{\alpha\beta} [F^{\beta}] \,. \tag{6}$$

To deduce the corresponding component representation, we remember that the functional $\kappa_{\alpha\beta}$ acts on 2-forms and creates as response other 2-forms. Since any 2-form can be decomposed with respect to the 2-form basis $\vartheta^{\nu\rho}$, it is sufficient to study the behavior of $\vartheta^{\nu\rho}$ under the application of $\kappa_{\alpha\beta}$. Because of the assumed locality and linearity, we have

$$\boldsymbol{\kappa}_{\alpha\beta}[\vartheta^{\nu\rho}] = \frac{1}{2} \kappa_{\lambda\mu\alpha}{}^{\nu\rho}{}_{\beta} \vartheta^{\lambda\mu} \,. \tag{7}$$

Let us come back to (6). We decompose H_{α} and F^{β} and, by using (7), we find:

$$H_{\alpha} = \frac{1}{2} H_{\lambda\mu\alpha} \vartheta^{\lambda\mu} = \boldsymbol{\kappa}_{\alpha\beta} [F^{\beta}] = \boldsymbol{\kappa}_{\alpha\beta} \left[\frac{1}{2} F_{\nu\rho}{}^{\beta} \vartheta^{\nu\rho} \right]$$
$$= \frac{1}{2} F_{\nu\rho}{}^{\beta} \boldsymbol{\kappa}_{\alpha\beta} [\vartheta^{\nu\rho}] = \frac{1}{4} \kappa_{\lambda\mu\alpha}{}^{\nu\rho}{}_{\beta} F_{\nu\rho}{}^{\beta} \vartheta^{\lambda\mu} . \tag{8}$$

By renaming some indices, we eventually derive the final formula

$$H_{\beta\gamma\alpha} = \frac{1}{2} \kappa_{\beta\gamma\alpha}^{\nu\rho} {}_{\mu}F_{\nu\rho}{}^{\mu} .$$
⁽⁹⁾

see [31, Eq.(47)].

If we use Schouten's [53] notation $(\alpha\beta) := \frac{1}{2} \{\alpha\beta + \beta\alpha\}$ and, moreover, $[\alpha\beta] := \frac{1}{2} \{\alpha\beta - \beta\alpha\}$, we have here the antisymmetries $H_{(\beta\gamma)\alpha} = 0$ and $F_{(\nu\rho)}^{\mu} = 0$. Thus, the constitutive tensor obeys the identities

$$\kappa_{(\beta\gamma)\alpha}{}^{\nu\rho}{}_{\mu} = 0 \quad \text{and} \quad \kappa_{\beta\gamma\alpha}{}^{(\nu\rho)}{}_{\mu} = 0.$$
 (10)

Accordingly, $\kappa_{\beta\gamma\alpha}{}^{\nu\rho}{}_{\mu}$ has $(6 \times 4)^2 = 576$ independent components. In the corresponding electrodynamics case, the constitutive law reads $H_{\beta\gamma} = \frac{1}{2} \kappa_{\beta\gamma}{}^{\nu\rho} F_{\nu\rho}$. Thus, by contrast, we have only $6^2 = 36$ independent components. As we will see further down, if we study only reversible processes, then this number is appreciably downsized in both cases.

A concise Hamiltonian formulation of teleparallel gravity was given by Ferraro and Guzmán [14]. Recently Hohmann et al. [26] studied teleparallel gravity, but instead of taking local and linear constitutive equations, they turned to local and *non*linear ones in order to incorporate f(T)-theories into the general TG formalism. These investigations [26] are very helpful since they bring order into the plethora of f(T)-theories and make them more transparent. Such nonlinear models provide an interesting development of gravitational theory based on an analogy with Born-Infeld-Plebański electrodynamics. We believe, though, that there is, at the present time, no real need to push nonlinear constitutive laws, since gravity is nonlinear anyway, due to its self-interaction—and this in spite of a *linear* constitutive law, which guarantees the quasi-linearity of the emerging field equation.

Kostelecký & Mewes [33] investigated Lorentz and diffeomorphism violations in *linearized* gravity. In this context, they introduced tensors which are of a similar type as our constitutive tensor χ . Our group-theoretical treatment of χ in Sec.3 is reminiscent of their method. However, in our article the full nonlinearity of gravity is treated in a premetric framework.

We would like to stress the following fact about teleparallelism theories: We have always a frame $e_{\alpha} = e^{i}{}_{\alpha}\partial_{i}$ and a coframe $\vartheta^{\beta} = e_{j}{}^{\beta}dx^{j}$ with us, that is, we can always change from holonomic to anholonomic indices by using $e^{i}{}_{\alpha}$ and $e_{j}{}^{\beta}$, respectively—and vice versa. This implies that all the indices in premetric TG are fundamentally equal, in particular, all those occurring in $\kappa_{\beta\gamma\alpha}{}^{\nu\rho}{}_{\mu}$ in (9). The 'group' indices α and μ are of the same quality as the 'form' indices β, γ, ν, ρ . Contractions with all indices are always allowed. Of course, at the premetric stage, raising and lowering of indices is only possible with the totally antisymmetric Levi-Civita symbols $\varepsilon_{\alpha\beta\gamma\delta} = \pm 1,0$ and $\varepsilon^{\mu\nu\rho\sigma} = \mp 1,0$, since no metric is available so far in our premetric framework.

If we follow the pattern of electrodynamics as formulated in *tensor calculus*, see Post [50], then it is obvious that we should introduce the excitation with the components $\check{H}^{\beta\gamma}{}_{\alpha} = \frac{1}{2} \varepsilon^{\beta\gamma\mu\nu} H_{\mu\nu\alpha}$, as already defined in Eq.(3). Thus,

$$\check{H}^{\beta\gamma}{}_{\alpha} = \frac{1}{2} \chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu} F_{\nu\rho}{}^{\mu}, \qquad (11)$$

with

$$\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu} := \frac{1}{2} \varepsilon^{\beta\gamma\delta\varepsilon} \kappa_{\delta\varepsilon\alpha}{}^{\nu\rho}{}_{\mu}.$$
(12)

The constitutive tensor density³ $\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu}$ is equivalent to $\kappa_{\beta\gamma\alpha}{}^{\nu\rho}{}_{\mu}$, in particular,

$$\chi^{(\beta\gamma)}{}_{\alpha}{}^{\nu\rho}{}_{\mu} = 0 \quad \text{and} \quad \chi^{\beta\gamma}{}_{\alpha}{}^{(\rho\nu)}{}_{\mu} = 0.$$
 (13)

But there is still a third useful version of the constitutive tensor available. For the purpose of the irreducible decomposition under the $GL(4,\mathbb{R})$, it is optimal to have the indices of the constitutive tensor all exclusively either in lower or in

³This tensor density emerged already in [21], see also [1, 9].

upper position. Since so far we have no metric at our disposal, we can only move the antisymmetric sets of indices. If we have a look at (9), we can rewrite it with $\check{F}^{\nu\rho\mu} = \frac{1}{2} \varepsilon^{\nu\rho\nu'\rho'} F_{\nu'\rho'}{}^{\mu}$ as

$$H_{\beta\gamma\alpha} = \frac{1}{2} \check{\chi}_{\beta\gamma\alpha\nu\rho\mu} \check{F}^{\nu\rho\mu} , \qquad (14)$$

that is,

$$\check{\chi}_{\beta\gamma\alpha\nu\rho\mu} := \frac{1}{2} \varepsilon_{\nu\rho\nu'\rho'} \kappa_{\beta\gamma\alpha}{}^{\nu'\rho'}{}_{\mu} = \frac{1}{4} \varepsilon_{\beta\gamma\beta'\gamma'} \varepsilon_{\nu\rho\nu'\rho'} \chi^{\beta'\gamma'}{}_{\alpha}{}^{\nu'\rho'}{}_{\mu} \,. \tag{15}$$

We have the symmetries

$$\check{\chi}_{(\beta\gamma)\alpha\nu\rho\mu} = 0 \quad \text{and} \quad \check{\chi}_{\beta\gamma\alpha(\nu\rho)\mu} = 0.$$
 (16)

1.3 Reversibility

A process which can in no way be completely reversed is termed irreversible, all other processes reversible. That a process may be irreversible, it is not sufficient that it cannot be directly reversed. This is the case with many mechanical processes which are not irreversible... ... The full requirement is, that it be impossible, even with the assistance of all agents in nature, to restore everywhere the exact initial state when the process has once taken place... the generation of heat by friction, the expansion of a gas without the performance of external work and the absorption of external heat, the conduction of heat, etc., are irreversible processes... Max Planck [49, p.84]⁴

The merit of formulating a field theory only in terms of its field equations here Eqs.(2) and (4)—and an associated constitutive law—here Eq.(6)—is that it covers both processes, irreversible and reversible ones. At first, like in the conventional treatment of general relativity [13], we turn our attention to reversible processes. From their definition it is clear that, for instance, periodic processes and those the equations of motion of which are formulated in a time symmetric way, are reversible. Time symmetry means that we can substitute in the field equations t by -t without changing them.

Thus dissipation is not allowed in reversible processes and we can define for each such process an energy function and, by a Legendre transformation, a Lagrangian. Accordingly, reversible processes can always be formulated by means of an action principle. For TG, the twisted Lagrange 4-form reads

$${}^{(\vartheta)}\Lambda = -\frac{1}{2}F^{\alpha} \wedge H_{\alpha} = -\frac{1}{2}F^{\alpha} \wedge \boldsymbol{\kappa}_{\alpha\beta}[F^{\beta}].$$
⁽¹⁷⁾

⁴Similarly, we have: Suppose that when a system under consideration changes from a state, α , to another state, α' , the environment changes from β to β' . If in some way it is possible to return the system from α' to α and at the same time to return the environment from β' to β , the process $(\alpha, \beta) \rightarrow (\alpha', \beta')$ is said to be reversible. Ryogo Kubo [34, p.61]

We substitute (5) and (8) into (17) and find:

With the definition (12) and the volume 4-form vol, Eq.(18) can be rewritten as

$${}^{(\vartheta)}\Lambda = -\frac{1}{8}\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu}F_{\beta\gamma}{}^{\alpha}F_{\nu\rho}{}^{\mu} \operatorname{vol}$$
(19)

or, by renaming the indices of the F's, equivalently as

$${}^{(\vartheta)}\Lambda = -\frac{1}{8}\chi^{\nu\rho}{}_{\mu}{}^{\beta\gamma}{}_{\alpha}F_{\beta\gamma}{}^{\alpha}F_{\nu\rho}{}^{\mu} \text{ vol.}$$
(20)

Consequently, only those components of the constitutive tensor enter the Lagrangian that satisfy the relations

$$\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu} = \chi^{\nu\rho}{}_{\mu}{}^{\beta\gamma}{}_{\alpha}.$$
⁽²¹⁾

If we assume that the model is completely specified by the Lagrangian (17), we restrict our considerations to reversible processes. Then the relations (21) are necessary conditions, which correspond to the symmetry of a 24×24 matrix. Thus, the set of the 576 independent components of $\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu}$ reduces to only 300 ones.

It is possible to rewrite the constitutive law in a more compact way. We introduce the 6-dimensional co-basis $\vartheta^{\alpha\beta} \rightarrow \vartheta^I$ in the space of 2-forms, with the collective indices $I, J, \dots = (01, 02, 03; 23, 31, 12) = (1, 2, \dots, 6)$, see [22, p.40]. Then excitation and field strength decompose as $H_{\alpha} = H_{I\alpha}\vartheta^I$ and $F^{\beta} = F_J{}^{\beta}\vartheta^J$, respectively, and the constitutive law reads,

$$H_{I\alpha} = \kappa_{I\alpha}{}^{J}{}_{\beta}F_{J}{}^{\beta}, \quad \check{H}^{I}{}_{\alpha} = \frac{1}{2}\chi^{I}{}_{\alpha}{}^{J}{}_{\beta}F_{J}{}^{\beta}, \qquad (22)$$

with the 300 independent components

$$\chi^{I}{}_{\alpha}{}^{J}{}_{\beta} = \chi^{J}{}_{\beta}{}^{I}{}_{\alpha} \,. \tag{23}$$

Thus, for the Lagrangian we find

$${}^{(\vartheta)}\Lambda = -\frac{1}{2}\chi^{I}{}_{\alpha}{}^{J}{}_{\beta}F_{I}{}^{\alpha}F_{J}{}^{\beta}\operatorname{vol}.$$
(24)

This closes our short introduction to TG. Like all classical theories, whose form is established, we can put TG into a Tonti type diagram [56] in order to

clearly display its structure. This will be done for the first time in Sec.2. Then we turn to a closer examination of the constitutive tensor of TG. In Sec.3, we decompose it into smaller pieces, in particular into the irreducible pieces with respect to the linear group $GL(4, \mathbb{R})$. In Sec.4, metric dependent constitutive tensors will be addressed, in particular those which relate TG to general relativity. In Sec.5, we will study the propagation of gravitational waves in TG within the geometric optics approximation. We will follow the procedure that we developed for electrodynamics [2, 22, 29]. Since in TG we have four generators of the gauge group, things become a bit more complicated than in electrodynamics. In Sec.6, we will specialize these considerations on gravitational waves to metric dependent models.

2 Tonti diagram of the premetric teleparallel theory of gravity

Over the past decades, Tonti [56] developed a general classification program for classical and relativistic theories in physics, such as, e.g., for particle dynamics, electromagnetism, the mechanics of deformable media, fluid mechanics, thermodynamics, and gravitation. Here we will display in Figure 1 for the first time an appropriate and consistent diagram of the teleparallel theory of gravity (TG).

If a theory is well-understood, its configuration and its source variables can be clearly identified and their interrelationships displayed in the form of a Tonti diagram. Such a diagram defines what one may call the skeleton of a theory. In Tonti's book [56], for all classical theories, including the relativistic ones, a corresponding framework was established—and this step by step, based on an operational definition of the quantities involved.

Tonti [56, p.402] has also displayed a diagram for relativistic gravitation. In Tonti's own words, it was an "attempt" of a diagram based on an ansatz for a tetrad theory of gravity by Kreisel & Treder, see [57, pp.60–67, pp.71–91]. Due to our enhanced knowledge of TG, see [8, Chapters 5 and 6], we can now improve on Tonti's attempt, see [23] and our Figure 1. The notation in Figure 1 is based on our recent paper [31] and on the present one.

Let us have a look at our new diagram. The configuration variables of TG are the coordinates x^i of the 4-dimensional spacetime (four 0-forms) and the coframes ϑ^{α} (four 1-forms). By differentiation, we find the torsion F^{β} (four 2-forms) and by further differentiation the homogeneous field equation of gravity $dF^{\beta} = 0$ (four 3-forms), the right hand side of which vanishes.

The round boxes on the left column depict geometrical objects and the square boxes interrelate these geometrical objects. The formula $dF^{\beta} = 0$, for example,

corresponds to the first Bianchi identity of a teleparallel spacetime. In a Riemann-Cartan space, we have $DF^{\beta} = \vartheta^{\alpha} \wedge R_{\alpha}{}^{\beta}$. In teleparallelism, the curvature vanishes, $R_{\alpha}{}^{\beta} = 0$, and, in the teleparallel gauge, $dF^{\beta} = 0$.

For global variables, Tonti distinguishes spatial domains, such as *volumes* V, *surfaces* S, *lines* L, and *points* P, with respect to time he introduces *instants* I and *intervals* T. Furthermore, for the respective domains, he has *inner* (interior) – and outer (exterior) \sim orientation.

Hence $[\mathbf{I} \times \overline{\mathbf{S}}]$, for example, refers to a time domain \mathbf{I} with exterior orientation and a space domain $\overline{\mathbf{S}}$ with interior orientation. And this domain $[\mathbf{I} \times \overline{\mathbf{S}}]$ supports the torsion two-form. The holonomic coordinates x^j , to take another example, depend on an instant of time \mathbf{I} and a spatial point \mathbf{P} . i.e. $[\mathbf{I} \times \mathbf{P}]$. Then we have to add the orientation. All exterior forms on the left columns are forms without twist, see [22], those on the right columns all carry twist. This can be read off from the corresponding orientations. This comes about as follows:

For configuration variables the associated space elements are endowed always with an interior orientation, whereas in the case of source variables it is the exterior orientation which plays a role. This behavior is found by phenomenologically examining the different theories. According to Tonti, the underlying theoretical reason for this correspondence is not clear, but phenomenology does not allow any other attributions. The configuration variables are related to the theory of chains of algebraic topology, whereas the source variables are associated to co-chains. For more details, we refer to the exhaustive monograph of Tonti [56].

Since the gravitational field itself carries energy-momentum, it is also the source of a new gravitational field, which likewise carries energy-momentum, etc.. Thus, like general relativity, TG is an intrinsically nonlinear theory, even it carries a linear constitutive law. Within the field equation of TG, $dH_{\alpha} - {}^{(\vartheta)}\Sigma_{\alpha} = {}^{(m)}\Sigma_{\alpha}$, the gravitational energy-momentum 3-form

$${}^{(\vartheta)}\Sigma_{\alpha} := \frac{1}{2} [F^{\beta} \wedge (e_{\alpha} \rfloor H_{\beta}) - H_{\beta} \wedge (e_{\alpha} \rfloor F^{\beta})]$$
⁽²⁵⁾

shows up explicitly and is a manifestation of this nonlinearity. By differentiation of the field equation, we find

$$ddH_{\alpha} = 0 = d\left({}^{(\vartheta)}\Sigma_{\alpha} + {}^{(m)}\Sigma_{\alpha}\right).$$
(26)

Thus, $d^{(m)}\Sigma_{\alpha} = -d^{(\vartheta)}\Sigma_{\alpha}$ is *nonvanishing* in general, which clearly shows up in our Tonti diagram.

The Tonti diagram displayed in Fig. 1 was constructed for the *premetric* version of teleparallelism, no metric is involved at all. We developed the corresponding formalism in our previous paper in [31, Sec.II]. Above, in Sec.1.1, we discussed already that teleparallel gravity is only equivalent to GR as long as the mat-

ter which is involved does *not* carry intrinsic spin. If this is the case, the energymomentum of matter is described by the 3-form ${}^{(m)}\sigma_{\alpha}$, with $\vartheta_{[\alpha} \wedge {}^{(m)}\sigma_{\beta]} = 0$, i.e., the corresponding energy-momentum tensor is symmetric. Since $\vartheta_{\alpha} := g_{\alpha\gamma} \vartheta^{\gamma}$, we need a metric for the specification of such an energy-momentum ${}^{(m)}\sigma_{\alpha}$. And the metric $g_{\alpha\beta}$ induces an associated Levi-Civita connection 1-form $\widetilde{\Gamma}_{\alpha}{}^{\beta}$.

The energy-momentum law, in the teleparallel gauge, reads $d^{(m)}\Sigma_{\alpha} = (e_{\alpha} \rfloor F^{\beta})$ $\wedge^{(m)}\Sigma_{\beta}$. If we dispense with the teleparallel gauge for the moment, the derivative $d^{(m)}\Sigma_{\alpha}$ can be substituted by $D^{(m)}\Sigma_{\alpha}$, with D as the covariant exterior derivative operator with respect to the teleparallel connection $\Gamma_{\alpha}{}^{\beta}$. Having now a metric available, we can assume that the teleparallel connection is metric compatible, that is, $Dg_{\alpha\beta} = 0$. For the symmetric energy-momentum, we have then the energy-momentum law

$$D^{(m)}\sigma_{\alpha} = (e_{\alpha} \rfloor F^{\beta}) \wedge^{(m)}\sigma_{\beta}.$$
⁽²⁷⁾

Due to a lemma of Meyer [38], the right side of (27) can be absorbed by the left side. This can be demonstrated by expanding the covariant derivative:

$$d^{(m)}\sigma_{\alpha} - (\Gamma_{\alpha}{}^{\beta} + e_{\alpha} \rfloor F^{\beta}) \wedge^{(m)} \sigma_{\beta} = 0.$$
⁽²⁸⁾

Let us now introduce a tilde to denote the Riemannian objects and operators: $\Gamma_{\alpha}{}^{\beta}$ is the Christoffel connection, for example, and \widetilde{D} the Riemannian covariant derivative. Since $\Gamma_{\alpha}{}^{\beta} = \widetilde{\Gamma}_{\alpha}{}^{\beta} - K_{\alpha}{}^{\beta}$, with the contortion 1-form $K_{\alpha}{}^{\beta}$, we can rewrite (28) as

$$\widetilde{D}^{(m)}\sigma_{\alpha} - (-K_{\alpha}{}^{\beta} + e_{\alpha} \rfloor F^{\beta}) \wedge^{(m)}\sigma_{\beta} = 0.$$
⁽²⁹⁾

The contortion, is related to the torsion via $F^{\beta} = K^{\beta}{}_{\gamma} \wedge \vartheta^{\gamma}$. If we substitute this into (29), it can be recast into

$$\widetilde{D}^{(m)}\sigma_{\alpha} + \left[K_{\alpha}{}^{\beta} - (e_{\alpha} \rfloor K^{\beta}{}_{\gamma}) \vartheta^{\gamma} + K^{\beta}{}_{\gamma}\delta^{\gamma}_{\alpha}\right] \wedge^{(m)}\sigma_{\beta} = 0.$$
(30)

Because of the antisymmetry of the contortion, $K_{(\alpha\beta)} = 0$, Eq.(30) simplifies to

$$\widetilde{D}^{(m)}\sigma_{\alpha} - (e_{\alpha} \rfloor K^{\beta\gamma}) \vartheta_{[\gamma} \wedge^{(m)} \sigma_{\beta]} = 0.$$
(31)

Now we recall that ${}^{(m)}\sigma_{\alpha}$ is symmetric, $\vartheta_{[\gamma} \wedge {}^{(m)}\sigma_{\beta]} = 0$. Consequently,

$$\widetilde{D}^{(m)}\sigma_{\alpha} = 0.$$
(32)

This is the energy-momentum law of GR. Accordingly, our entry $d^{(m)}\Sigma_{\alpha} = (e_{\alpha} \rfloor F^{\beta}) \wedge {}^{(m)}\Sigma_{\beta}$ in the Tonti diagram becomes, provided a symmetric energy-momentum tensor is prescribed, $\widetilde{D}^{(m)}\sigma_{\alpha} = 0$, well in accordance with the vanishing divergences in analogous Tonti diagrams.

Tonti diagram of the premetric teleparallel theory of gravity^{*} (TG)



Figure 1: • Patterned after E. Tonti: *The Mathematical Structure of Classical and Relativistic Physics*, A general classification diagram (Birkhäuser-Springer, New York, 2013) pages 402 and 315.

• Notation: Y. Itin, F. W. Hehl, Yu. N. Obukhov, Phys. Rev. D **95**, 084020 (2017), arXiv:1611.05759. We denoted *here* the torsion 2-form with F^{β} in order to underline its function as a field strength. Usually, however, we use for torsion T^{β} .

• We chose everywhere the 'teleparallel gauge' such that the connection 1-form vanishes globally: $\Gamma^{\alpha\beta}(x) \stackrel{*}{=} 0$.

• For the reversible case, we have the gravitational Lagrangian as ${}^{(\vartheta)}\Lambda = -\frac{1}{2}F^{\alpha} \wedge H_{\alpha}$ and $\Lambda = {}^{(\vartheta)}\Lambda + {}^{(m)}\Lambda$. The gravitational constant is denoted by \varkappa .

*Also known as translation gauge theory of gravity.

3 Decompositions of the constitutive tensor

In this section we characterize the constitutive tensor due to its symmetry transformations relative to the general linear group $GL(4,\mathbb{R})$ and the permutation (symmetry) group. We refer to Hamermesh [15] and to Barut & Raczka [5] as background information for group theory.

3.1 Two forms of the constitutive tensor

According to (11), the constitutive tensor density $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ is a order- $\begin{pmatrix} 4\\2 \end{pmatrix}$ tensor density that is skew-symmetric in the two pairs of upper indices, see (13):

$$\chi^{(\alpha\beta)}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = 0, \quad \chi^{\alpha\beta}{}_{\mu}{}^{(\gamma\delta)}{}_{\nu} = 0.$$
(33)

In *n* dimensions, we have $n\binom{n}{2}n\binom{n}{2} = \frac{1}{4}n^4(n-1)^2$ independent components. For n = 4, it gives 576 relevant components. Due to this large number, it is instructive to consider decomposition of the constitutive tensor into smaller pieces. We apply Young's decomposition technique that yields irreducible decomposition under the group GL(4, \mathbb{R}). For our conventions and the details of the Young decomposition, see A. For the tensor $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$, separate S_4 -permutations of four upper indices and S_2 -permutations of two lower indices are available, where S_n denotes the *n*-dimensional permutation group. Correspondingly, we are dealing with the Cartesian product group $S_4 \times S_2$.

Although the metric tensor is not available in our construction, we can, as we have shown in (15), lower two pairs of skew-symmetric indices by the Levi-Civita symbol. Thus, we have a order- $\begin{pmatrix} 0 \\ 6 \end{pmatrix}$ tensor

$$\check{\chi}_{\alpha\beta\mu\gamma\delta\nu} := \frac{1}{4} \varepsilon_{\alpha\beta\omega\lambda} \varepsilon_{\gamma\delta\rho\sigma} \chi^{\omega\lambda}{}_{\mu}{}^{\rho\sigma}{}_{\nu}, \qquad (34)$$

with the identities

$$\check{\chi}_{(\alpha\beta)\mu\gamma\delta\nu} = 0, \quad \check{\chi}_{\alpha\beta\mu(\gamma\delta)\nu} = 0.$$
(35)

This tensor is naturally decomposed under the permutation group S_6 . Both types of decompositions are invariant under the action of the basis transformation group $GL(4,\mathbb{R})$.

3.2 $S_4 \times S_2$ decomposition of $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$

Treating the covariant and contravariant indices as belonging to two separate tensor spaces, the irreducible decomposition of $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ is defined as a product of the irreducible decompositions. The corresponding Young diagrams are expanded as

Here, the three first diagrams relate to the upper indices. This decomposition repeats the known irreducible decomposition of the electromagnetic constitutive tensor. The remaining two diagrams represent the symmetric and antisymmetric parts of the second order tensor. Collecting all possible combinations, we obtain the decomposition of the tensor $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ into 6 independent pieces:

In tensor notation, the above decomposition reads

$$\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = \sum_{I=1}^{6}{}^{[I]}\mathbb{P}^{\alpha\beta}{}_{\omega\lambda\mu\rho\sigma\nu}{}^{\gamma\delta}{}_{\nu}\chi^{\omega\lambda}{}_{\kappa}{}^{\rho\sigma}{}_{\varepsilon} = \sum_{I=1}^{6}{}^{[I]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}.$$
(38)

Here we used the set of six projection operators $[I]\mathbb{P}$. We chose the labeling of I = 1, ..., 6 such that it corresponds to the same sequence of Young diagrams as depicted in the second equality of Eq.(38). Explicitly, the $S_4 \times S_2$ -irreducible pieces of the constitutive tensor are expressed as

$${}^{[1]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} - {}^{[3]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} - {}^{[5]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}, \tag{39}$$

$${}^{[2]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} - {}^{[4]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} - {}^{[6]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}, \tag{40}$$

$$^{[3]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \frac{1}{2} \left(\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} - \chi^{\gamma\delta}{}_{(\mu}{}^{\alpha\beta}{}_{\nu)} \right), \tag{41}$$

$${}^{[4]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \frac{1}{2} \left(\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} - \chi^{\gamma\delta}{}_{[\mu}{}^{\alpha\beta}{}_{\nu]} \right), \tag{42}$$

$$^{[5]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \chi^{[\alpha\beta}{}_{(\mu}{}^{\gamma\delta]}{}_{\nu)}, \tag{43}$$

$${}^{[6]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \chi^{[\alpha\beta}{}_{[\mu}{}^{\gamma\delta]}{}_{\nu]}. \tag{44}$$

In terms of independent components in n = 4 dimensions, this decomposition corresponds to

$$576 = 200 \oplus 120 \oplus 150 \oplus 90 \oplus 10 \oplus 6.$$
 (45)

It is straightforward to show that the above operators $[I]\mathbb{P} = [I]\mathbb{P}_{\omega\lambda\mu\rho\sigma\nu}^{\alpha\beta\kappa\gamma\delta\varepsilon}$ are orthogonal projectors:

$${}^{[I]}\mathbb{P} \circ {}^{[J]}\mathbb{P} = {}^{[I]}\mathbb{P}\delta^{IJ}, \quad \delta^{IJ} = \begin{cases} 1 & \text{for } I = J \\ 0 & \text{for } I \neq J \end{cases}.$$
(46)

In (39) to (44), we first decompose the gravitational constitutive tensor into three independent pieces relative to its four upper indices. This S_4 decomposition is done completely similar to electrodynamics. Then we extract relative to the pair of the lower indices the symmetric and antisymmetric parts. Since the principal and axion part of the electromagnetic constitutive tensor are reversible, the corresponding parts of the gravitational constitutive tensor are reversible when symmetrized in the lower indices. The electrodynamics the skewon part is irreversible. Thus its gravitational analog is reversible when antisymmetrized in the lower indices. The remaining three parts are irreversible (change their sign under permutation of triads of indices).

In order to demonstrate this behavior explicitly, we decompose the constitutive tensor into its reversible and irreversible parts,

$$\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = \chi^{+\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + \bar{\chi}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}, \qquad (47)$$

with

$$\chi^{\alpha\beta}{}_{\nu}{}^{\gamma\delta}{}_{\nu} := \frac{1}{2} (\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + \chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu}) = \chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu}, \qquad (48)$$

$$\bar{\chi}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \frac{1}{2} (\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} - \chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu}) = -\bar{\chi}^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu}.$$
(49)

The first part $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ describes *reversible* processes, it can be derived from a Lagrangian. In contrast, the second part $\bar{\chi}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ corresponds to irreversibility, and it does not enter the Lagrangian. Following the nomenclature in electrodynamics, we will call $\bar{\chi}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ the *skewon* part.

In the next step, we can decompose each of these parts into the symmetric/antisymmetric piece in the lower indices:

$$\overset{\pm}{\chi}{}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = \overset{\pm}{\chi}{}^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} + \overset{\pm}{\chi}{}^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]}.$$
(50)

Two immediate consequences are

$$\overset{+}{\chi}{}^{[\alpha\beta}{}_{[\mu}{}^{\gamma\delta]}{}_{\nu]} \equiv 0, \qquad \overset{-}{\chi}{}^{[\alpha\beta}{}_{(\mu}{}^{\gamma\delta]}{}_{\nu)} \equiv 0.$$
(51)

Thus, axion pieces can be extracted only from $\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)}$ and from $\bar{\chi}^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]}$.

After these preliminaries, we can recast the irreducible decompositions (39) to (44) into a more transparent form,

$$^{[1]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \chi^{+\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} - \chi^{+[\alpha\beta}{}_{(\mu}{}^{\gamma\delta]}{}_{\nu)}, \tag{52}$$

$$^{[2]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \bar{\chi}^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} - \bar{\chi}^{[\alpha\beta}{}_{[\mu}{}^{\gamma\delta]}{}_{\nu]}, \tag{53}$$

$$^{[3]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \bar{\chi}^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)}, \tag{54}$$

$${}^{[4]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \chi^{+\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]}, \tag{55}$$

$${}^{[5]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \chi^{+[\alpha\beta}{}_{(\mu}{}^{\gamma\delta]}{}_{\nu)}, \tag{56}$$

$${}^{[6]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \bar{\chi}^{[\alpha\beta}{}_{[\mu}{}^{\gamma\delta]}{}_{\nu]}.$$

$$(57)$$

In other words, we have a decomposition of the reversible and skewon parts into, respectively,

$$\chi^{+\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = {}^{[1]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[4]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[5]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}, \tag{58}$$

$$\bar{\chi}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = {}^{[2]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[3]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[6]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}.$$
(59)

We may call ${}^{[1]}\chi$ a reversible symmetric principal part (principal-1), ${}^{[4]}\chi$ a reversible antisymmetric principal part (principal-2), ${}^{[5]}\chi$ a reversible axion (axion-1); likewise we may call ${}^{[2]}\chi$ a skewon antisymmetric principal part (skewon-1), ${}^{[3]}\chi$ a skewon symmetric principal part (skewon-2), and ${}^{[6]}\chi$ a skewon axion (axion-2). This is summarized in Table 1.

For completeness, we present here the results about the contributions of the irreducible pieces into the dispersion relation for the gravitational waves. These facts will be derived in Sec.6.

3.3 S_6 decomposition of $\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}$

Now we consider the decomposition of the constitutive tensor under the permutation of its six lower indices. We follow Hamermesh's [15] Eqs. (7-159) to (7-162). Then, for n = 4 dimensions, the symmetries (35) imply the Young decomposition

Irr. Parts	Nomenclature	Lagr.	En-mom.	Disp.
$^{[1]}\chi^{lphaeta}\gamma^{\mu u} ho$	principal-1	yes	yes	yes
$^{[2]}\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$	skewon-1	no	yes	yes
$^{[3]}\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$	skewon-2	no	yes	yes
${}^{[4]}\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$	principal-2	yes	yes	yes
$^{[5]}\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$	axion-1	yes	no	no
$^{[6]}\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$	axion-2	no	yes	no

Table 1: Physical identifications of the irreducible parts of the constitutive tensor $\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$.

of $\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}$:



Here we omitted vanishing diagrams in n = 4 (i.e. diagrams that contain antisymmetrization with respect to more than four indices). In terms of independent components, we have

$$576 = 126 \oplus 70 \oplus 50 \oplus 4 \times 64 \oplus 2 \times 10$$
$$\oplus 3 \times 10 \oplus 4 \times 6. \tag{61}$$

In tensorial language, this corresponds to

$$\check{\chi}_{\alpha\beta\mu\gamma\delta\nu} = \sum_{I=1}^{7} {}^{(I)} \mathbb{P}_{\alpha\beta\mu\gamma\delta\nu}^{\omega\lambda\kappa\rho\sigma\varepsilon} \check{\chi}_{\omega\lambda\kappa\rho\sigma\varepsilon} = \sum_{I=1}^{7} {}^{(I)} \check{\chi}_{\alpha\beta\mu\gamma\delta\nu}, \qquad (62)$$

where again the expressions ${}^{(I)}\mathbb{P}^{\omega\lambda\kappa\rho\sigma\varepsilon}_{\alpha\beta\mu\gamma\delta\nu}$ denote orthogonal projectors,

$${}^{(I)}\mathbb{P} \circ {}^{(J)}\mathbb{P} = {}^{(I)}\mathbb{P}\delta^{IJ}, \quad \delta^{IJ} = \begin{cases} 1 & \text{for } I = J \\ 0 & \text{for } I \neq J \end{cases}.$$
(63)

The explicit expressions of the seven terms ${}^{(I)}\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}$ are quite involved and we do not display them here; they can be found in A.

3.4 **Relation between the decompositions**

In order to relate the two inequivalent decompositions discussed above, it is useful to define

$${}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} := \frac{1}{4} \varepsilon^{\alpha\beta\omega\lambda} \varepsilon^{\gamma\delta\rho\sigma\,(I)} \check{\chi}_{\omega\lambda\mu\rho\sigma\nu}. \tag{64}$$

By using Eqs. (241) to (247), one may verify that the tensors ${}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ satisfy the relations

$${}^{\{I\}}\chi^{[\alpha\beta}{}_{\mu}{}^{\gamma\delta]}{}_{\nu} = 0 \text{ for } I = 1, 2, 3, 4, 5, \tag{65}$$

$${}^{\{I\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{V]} = 0 \text{ for } I = 1, 2, \tag{66}$$

$${}^{\{I\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} = 0 \text{ for } I = 1, 2,$$

$${}^{\{I\}}\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} = 0 \text{ for } I = 3,$$
(66)
(67)

as well as

$${}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = + {}^{\{I\}}\chi^{\gamma\delta}{}_{\mu}{}^{\alpha\beta}{}_{\nu} \quad \text{for } I = 1,3,$$
(68)

$${}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = -{}^{\{I\}}\chi^{\gamma\delta}{}_{\mu}{}^{\alpha\beta}{}_{\nu} \quad \text{for } I = 2, \tag{69}$$

$${}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = + {}^{\{I\}}\chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu} \quad \text{for } I = 1, 5,$$
(70)

$${}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = -{}^{\{I\}}\chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu} \quad \text{for } I = 2,3.$$

$$(71)$$

Eventually, we find

$${}^{\{6\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} + {}^{\{6\}}\chi^{\gamma\delta}{}_{[\mu}{}^{\alpha\beta}{}_{\nu]} = 0,$$
(72)

$${}^{\{6\}}\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} + {}^{\{6\}}\chi^{\gamma\delta}{}_{(\mu}{}^{\alpha\beta}{}_{\nu)} = {}^{\{6\}}\chi^{[\alpha\beta}{}_{(\mu}{}^{\gamma\delta]}{}_{\nu)}, \tag{73}$$

$${}^{\{7\}}\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} + {}^{\{7\}}\chi^{\gamma\delta}{}_{(\mu}{}^{\alpha\beta}{}_{\nu)} = 0.$$
(74)

We can apply directly the $S_4 \times S_2$ decomposition to these order- $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$ tensors and compare the results to the tensors

 $^{[I]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$. By extensive use of computer algebra, we find the following relations:

$$\begin{aligned} & {}^{[2]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = {}^{\{3\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} - {}^{\{7\}}\chi^{[\alpha\beta}{}_{[\mu}{}^{\gamma\delta]}{}_{\nu]} \\ & + \frac{1}{2}\sum_{I=4.7} \left[{}^{\{I\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} + {}^{\{I\}}\chi^{\gamma\delta}{}_{[\mu}{}^{\alpha\beta}{}_{\nu]} \right], \end{aligned}$$
(76)

$${}^{[6]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = {}^{\{7\}}\chi^{[\alpha\beta}{}_{[\mu}{}^{\gamma\delta]}{}_{\nu]}.$$

$$(80)$$

3.5 Reversibility

For reversible processes, as shown in (21), the constitutive tensor satisfies

$$\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = \chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu}. \tag{81}$$

As we demonstrated above, the reversible and irreversible parts of the constitutive tensor are presented, respectively, as

$$\begin{aligned} \chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} &= \frac{1}{2} \left(\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + \chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu} \right) \\ &= {}^{[1]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[4]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[5]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}, \\ \bar{\chi}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} &= \frac{1}{2} \left(\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} - \chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu} \right) \\ &= {}^{[2]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[3]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{[6]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}. \end{aligned} \tag{82}$$

This corresponds to $576 = 300 \oplus 276$, that is, the reversible constitutive tensor comprises 300 independent components. Employing relations Eqs. (75)–(80), one

finds

$$\begin{aligned} \chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} &= {}^{\{1\}}\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} + {}^{\{5\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} & (84) \\ &+ \frac{1}{2}\sum_{I=4,6,7} \left[{}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} + {}^{\{I\}}\chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu}{}^{\mu} \right], \\ \bar{\chi}^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} &= {}^{\{2\}}\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} + {}^{\{3\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} \\ &+ \frac{1}{2}\sum_{I=4,6,7} \left[{}^{\{I\}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} - {}^{\{I\}}\chi^{\gamma\delta}{}_{\nu}{}^{\alpha\beta}{}_{\mu}{}^{\mu} \right]. \end{aligned}$$

Let us emphasize that Eqs. (66) and (67) imply

$${}^{\{1\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} = 0, \; {}^{\{2\}}\chi^{\alpha\beta}{}_{[\mu}{}^{\gamma\delta}{}_{\nu]} = 0, \; {}^{\{3\}}\chi^{\alpha\beta}{}_{(\mu}{}^{\gamma\delta}{}_{\nu)} = 0.$$
(86)

In summary, we see that the reversible piece of the constitutive tensor $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ is comprised of the order- $\begin{pmatrix} 0 \\ 6 \end{pmatrix}$ irreducible pieces I = 1,5 as well as the appropriately (anti-)symmetrized parts of I = 4,6,7.

3.6 Lagrangian

Let us first consider how these independent pieces contribute to the gravitational Lagrangian ${}^{(\vartheta)}\Lambda = -\frac{1}{2}F^{\alpha} \wedge H_{\alpha}$. In components, we have

$${}^{(\vartheta)}\Lambda = -\frac{1}{4}F_{\beta\gamma}{}^{\alpha}\check{H}^{\beta\gamma}{}_{\alpha}\text{vol}.$$
(87)

Substitution of the constitutive law (11) yields

$${}^{(\vartheta)}\Lambda = -\frac{1}{8}\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu}F_{\beta\gamma}{}^{\alpha}F_{\nu\rho}{}^{\mu} \operatorname{vol}.$$
(88)

Consequently only those parts of the constitutive tensor contribute to the Lagrangian that satisfy the "pair commutation" symmetries

$$\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu} = \chi^{\nu\rho}{}_{\mu}{}^{\beta\gamma}{}_{\alpha}.$$
(89)

Accordingly, only the following terms are left over in the Lagrangian:

$${}^{(\vartheta)}\Lambda = -\frac{1}{8} \sum_{I=1,4,5} {}^{[I]} \chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu}F_{\beta\gamma}{}^{\alpha}F_{\nu\rho}{}^{\mu} \operatorname{vol}.$$
(90)

3.7 Parametrization of irreducible parts

In premetric electrodynamics, there exists a convenient parametrization of the axion and skewon parts by a pseudo-scalar and a traceless second order tensor, see Secs.D.1.4 and D.1.5 of the book [22].

One may wish to develop a similar formalism for the gravitational case under consideration. By analogy with electrodynamics, let us introduce the following new objects:

$$\mathscr{J}_{\gamma\rho\sigma}{}^{\nu} := \frac{1}{4} \varepsilon_{\alpha\beta\mu\sigma}{}^{[3]} \chi^{\alpha\beta}{}^{\mu\nu}{}_{\rho}, \qquad (91)$$

$$\mathcal{Q}_{\gamma\rho\sigma}{}^{\nu} := \frac{1}{4} \varepsilon_{\alpha\beta\mu\sigma}{}^{[4]} \chi^{\alpha\beta}{}^{\mu\nu}{}_{\rho}.$$
(92)

By construction (hint: use the definitions of ${}^{[3]}\chi$ and ${}^{[4]}\chi$), these tensors obey

$$\mathscr{J}_{[\alpha\beta]\mu}{}^{\nu} = 0, \qquad \mathscr{J}_{\alpha\beta\nu}{}^{\nu} = 0; \tag{93}$$

$$\mathcal{Q}_{(\alpha\beta)\mu}{}^{\nu} = 0, \qquad \mathcal{Q}_{\alpha\beta\nu}{}^{\nu} = 0.$$
(94)

Accordingly, the number of independent components for $\beta_{\alpha\beta\mu}{}^{\nu}$ is $10 \times 4 \times 4 - 10 = 150$ and for $\mathcal{Q}_{\alpha\beta\mu}{}^{\nu} 6 \times 4 \times 4 - 6 = 90$, respectively. It is straightforward to see that

$${}^{[3]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = \varepsilon^{\alpha\beta\lambda[\mu}\,\mathcal{S}_{\gamma\rho\lambda}{}^{\nu]} - \varepsilon^{\mu\nu\lambda[\alpha}\,\mathcal{S}_{\gamma\rho\lambda}{}^{\beta]}, \tag{95}$$

$${}^{[4]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = \varepsilon^{\alpha\beta\lambda[\mu}\mathcal{Q}_{\gamma\rho\lambda}{}^{\nu]} - \varepsilon^{\mu\nu\lambda[\alpha}\mathcal{Q}_{\gamma\rho\lambda}{}^{\beta]}.$$
(96)

Thereby, we found a complete parametrization of ${}^{[3]}\chi$ and ${}^{[4]}\chi$. This construction obviously generalizes the representation of a skewon in electrodynamics in terms of the traceless second order tensor [22]. In the context of the gravitational theory, however, only (95) refers to the skewon, whereas (96) parametrizes the *reversible* irreducible part.

Now, let us turn to the irreducible parts ${}^{[3]}\chi$ and ${}^{[4]}\chi$ for which we introduce

$${}^{[5]}A_{\gamma\rho} := \frac{1}{4!} \varepsilon_{\alpha\beta\mu\nu} {}^{[5]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = {}^{[5]}A_{\rho\gamma}, \qquad (97)$$

$${}^{[6]}A_{\gamma\rho} := \frac{1}{4!} \varepsilon_{\alpha\beta\mu\nu} {}^{[6]}\chi^{\alpha\beta}{}_{\gamma}{}^{\mu\nu}{}_{\rho} = -{}^{[6]}A_{\rho\gamma}.$$
(98)

The reciprocal relations read

$${}^{[5]}\chi^{\alpha\beta}\gamma^{\mu\nu}\rho = {}^{[5]}A_{\gamma\rho}\varepsilon^{\alpha\beta\mu\nu}, \qquad (99)$$

$${}^{[6]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = {}^{[6]}A_{\gamma\rho}\varepsilon^{\alpha\beta\mu\nu}.$$
(100)

As a result, we find for the axion type part of the constitutive tensor

$$^{(\mathrm{ax})}\chi^{\alpha\beta}{}_{\gamma}{}^{\mu\nu}{}_{\rho} = \mathscr{A}_{\gamma\rho}\,\varepsilon^{\alpha\beta\mu\nu}, \ \mathscr{A}_{\alpha\beta} := {}^{[5]}A_{\alpha\beta} + {}^{[6]}A_{\alpha\beta}. \tag{101}$$

Summarizing, we demonstrated that the four irreducible parts ${}^{[3]}\chi$, ${}^{[4]}\chi$, ${}^{[5]}\chi$, and ${}^{[6]}\chi$ of the constitutive tensor can be conveniently parametrized, see Eqs.(95), (96), (99), and (100). They are expressed in terms of the tensors (91), (92), (97), and (98) with the correct number of independent components: for $\beta_{\alpha\beta\mu}{}^{\nu}$ 150, for $\mathcal{Q}_{\alpha\beta\mu}{}^{\nu}$ 90, for ${}^{[5]}A_{\alpha\beta}$ 10, and for ${}^{[6]}A_{\alpha\beta}$ 6. The two remaining pieces ${}^{[1]}\chi$ and ${}^{[2]}\chi$ do not admit simple parametrizations, though.

3.8 Energy-momentum current

The energy-momentum current of the coframe field is defined as [31]

$${}^{(\vartheta)}\Sigma_{\alpha} = \frac{1}{2} \left[F^{\beta} \wedge (e_{\alpha} \rfloor H_{\beta}) - H_{\beta} \wedge (e_{\alpha} \rfloor F^{\beta}) \right].$$
(102)

The constitutive law for the pure axion piece (101) is expressed as

$${}^{(\mathrm{ax})}H_{\alpha} = \mathscr{A}_{\alpha\beta}F^{\beta} \,. \tag{103}$$

In this case, the energy-momentum current takes the form

$${}^{(\mathrm{ax})}\Sigma_{\alpha} = \mathscr{A}_{[\beta\gamma]}F^{\beta} \wedge (e_{\alpha} \rfloor F^{\gamma}).$$
(104)

Consequently, the symmetric combination of the axionic part ${}^{[5]}\chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu}$ does *not* contribute to the energy-momentum current.

3.9 Contracting $\chi^{\alpha\beta}\gamma^{\mu\nu}\rho$ twice: four second order tensors

In electrodynamics, we can extract from the constitutive tensor $\chi^{\alpha\beta\gamma\delta}$ a pseudoscalar density, the axion field, by contracting it with the Levi-Civita symbol: ${}^{(ax)}\alpha := \frac{1}{4!} \varepsilon_{\alpha\beta\gamma\delta} \quad \chi^{\alpha\beta\gamma\delta}$. In gravity, we can only reduce the constitutive tensor $\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho}$ of type $\binom{4}{2}$ to four different 2nd order tensors:

$$m^{\alpha\beta} := \chi^{\alpha\mu}{}_{\mu}{}^{\beta\nu}{}_{\nu}, \qquad (105)$$

$$n^{\alpha\beta} := \chi^{\alpha\mu}{}_{\nu}{}^{\beta\nu}{}_{\mu}, \qquad (106)$$

$$k^{\alpha\beta} := \chi^{\mu\nu}{}_{\mu}{}^{\alpha\beta}{}_{\nu}, \qquad (107)$$

$$l^{\alpha\beta} := \chi^{\alpha\beta}{}_{\mu}{}^{\mu\nu}{}_{\nu}. \tag{108}$$

The construction of scalars is impossible at this premetric stage.

The tensors $k^{\alpha\beta} = -k^{\beta\alpha}$ and $l^{\alpha\beta} = -l^{\beta\alpha}$ are antisymmetric, that is, $k^{(\alpha\beta)} = 0$, $l^{(\alpha\beta)} = 0$, whereas $m^{\alpha\beta}$ and $n^{\alpha\beta}$ are asymmetric. However, in the reversible case, $m^{\alpha\beta}$ and $n^{\alpha\beta}$ turn out to be symmetric, whereas the two other antisymmetric tensors turn out to be transposed to each other, $k^{\alpha\beta} = l^{\beta\alpha}$. This can be recognized more clearly by collecting all the traces into the specific constitutive tensor

$$\chi^{\alpha\beta}{}_{\gamma}{}^{\mu\nu}{}_{\rho} = \mathscr{L}^{\alpha\beta}\,\delta^{[\mu}{}_{\gamma}\delta^{\nu]}{}_{\rho} + \mathscr{K}^{\mu\nu}\,\delta^{[\alpha}{}_{\rho}\delta^{\beta]}{}_{\gamma} + \delta^{[\alpha}{}_{\gamma}\mathscr{M}^{\beta][\mu}\delta^{\nu]}{}_{\rho} + \delta^{[\alpha}{}_{\rho}\mathscr{N}^{\beta][\mu}\delta^{\nu]}{}_{\gamma}.$$
(109)

This tensor is determined by four tensors of 2nd order: two of them are antisymmetric, $\mathscr{L}^{\alpha\beta} = -\mathscr{L}^{\beta\alpha}$ and $\mathscr{K}^{\alpha\beta} = -\mathscr{K}^{\beta\alpha}$, that is, $\mathscr{L}^{(\alpha\beta)} = 0$, $\mathscr{K}^{(\alpha\beta)} = 0$, and the two other ones are general 2nd order tensors $\mathscr{M}^{\alpha\beta}$ and $\mathscr{N}^{\alpha\beta}$.

It is straightforward to establish one-to-one correspondences between these objects and the traces (105) to (108):

$$m^{\alpha\beta} = \frac{3}{2} \left(\mathscr{L}^{\alpha\beta} - \mathscr{K}^{\alpha\beta} \right) - \frac{1}{4} \left(9 \mathscr{M}^{\alpha\beta} + 2 \mathscr{N}^{\alpha\beta} + \mathscr{N}^{\beta\alpha} \right), \qquad (110)$$

$$n^{\alpha\beta} = \frac{3}{2} \left(\mathscr{K}^{\alpha\beta} - \mathscr{L}^{\alpha\beta} \right) - \frac{1}{4} \left(9 \mathscr{N}^{\alpha\beta} + 2 \mathscr{M}^{\alpha\beta} + \mathscr{M}^{\beta\alpha} \right), \qquad (111)$$

$$k^{\alpha\beta} = \mathscr{L}^{\alpha\beta} - 6\mathscr{K}^{\alpha\beta} - \frac{3}{2}\mathscr{M}^{[\alpha\beta]} + \frac{3}{2}\mathscr{N}^{[\alpha\beta]}, \qquad (112)$$

$$l^{\alpha\beta} = 6\mathscr{L}^{\alpha\beta} - \mathscr{K}^{\alpha\beta} - \frac{3}{2}\mathscr{M}^{[\alpha\beta]} + \frac{3}{2}\mathscr{N}^{[\alpha\beta]}.$$
(113)

The inverse relations read

$$\mathscr{L}^{\alpha\beta} = \frac{1}{10} \left(3l^{\alpha\beta} + k^{\alpha\beta} - 3m^{[\alpha\beta]} + 3n^{[\alpha\beta]} \right), \qquad (114)$$

$$\mathscr{K}^{\alpha\beta} = \frac{1}{10} \left(-l^{\alpha\beta} - 3k^{\alpha\beta} + 3m^{[\alpha\beta]} - 3n^{[\alpha\beta]} \right), \qquad (115)$$

$$\mathcal{M}^{\alpha\beta} = \frac{1}{10} \left(3l^{\alpha\beta} + 3k^{\alpha\beta} - 9m^{[\alpha\beta]} + 5n^{[\alpha\beta]} \right) \\ -\frac{1}{6} \left(3m^{(\alpha\beta)} - n^{(\alpha\beta)} \right), \qquad (116)$$

$$\mathcal{N}^{\alpha\beta} = \frac{1}{10} \left(-3l^{\alpha\beta} - 3k^{\alpha\beta} + 5m^{[\alpha\beta]} - 9n^{[\alpha\beta]} \right) \\ + \frac{1}{6} \left(m^{(\alpha\beta)} - 3n^{(\alpha\beta)} \right).$$
(117)

Decomposing the specific constitutive tensor (109) into reversible and irreversible (skewon) parts yields

$$\overset{\pm}{\chi}^{\alpha\beta}{}_{\gamma}{}^{\mu\nu}{}_{\rho} = \overset{\pm}{\mathscr{L}}{}^{\alpha\beta}\delta^{[\mu}{}_{\gamma}\delta^{\nu]}{}_{\rho} \pm \overset{\pm}{\mathscr{L}}{}^{\mu\nu}\delta^{[\alpha}{}_{\rho}\delta^{\beta]}{}_{\gamma} + \delta^{[\alpha}{}_{\gamma}{}^{\pm}{}_{\beta}{}^{\beta][\mu}\delta^{\nu]}{}_{\gamma} + \delta^{[\alpha}{}_{\rho}{}^{\pm}{}_{\gamma}{}^{\beta][\mu}\delta^{\nu]}{}_{\gamma}.$$
(118)

Here we introduced

$$\overset{\pm}{\mathscr{L}}{}^{\alpha\beta} = \frac{1}{2} \left(\mathscr{L}^{\alpha\beta} \pm \mathscr{K}^{\alpha\beta} \right), \qquad (119)$$

$$\overset{+}{\mathscr{M}}^{\alpha\beta} = \mathscr{M}^{(\alpha\beta)}, \quad \overset{-}{\mathscr{M}}^{\alpha\beta} = \mathscr{M}^{[\alpha\beta]}, \quad (120)$$

$$\overset{+}{\mathscr{N}}^{\alpha\beta} = \mathscr{N}^{(\alpha\beta)}, \quad \overset{-}{\mathscr{N}}^{\alpha\beta} = \mathscr{N}^{[\alpha\beta]}.$$
(121)

This confirms that the irreversible, the skewon part vanishes, $\bar{\chi}^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = 0$, provided $\bar{\mathscr{L}}^{\alpha\beta} = 0$, $\bar{\mathscr{M}}^{\alpha\beta} = 0$, and $\bar{\mathscr{N}}^{\alpha\beta} = 0$, i.e., when $m^{\alpha\beta}$ and $n^{\alpha\beta}$ are symmetric and $k^{\alpha\beta} = l^{\beta\alpha}$.

In the general case, the specific constitutive trace tensor (109) is characterized by 5 nontrivial irreducible parts:

$${}^{[1]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = \delta^{[\alpha}_{(\gamma}{}^{s}_{\mathcal{R}}{}^{\beta][\mu}\delta^{\nu]}_{\rho)}, \qquad (122)$$

$${}^{[2]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = \bar{\mathscr{L}}^{\alpha\beta}\delta^{[\mu}_{\gamma}\delta^{\nu]}_{\rho} - \bar{\mathscr{L}}^{\mu\nu}\delta^{[\alpha}_{\rho}\delta^{\beta]}_{\gamma} + \delta^{[\alpha}_{[\gamma}{}^{\alpha}_{\mathcal{S}}{}^{\beta][\mu}\delta^{\nu]}_{\rho]} - \mathscr{A}^{[\alpha\beta}\delta^{\mu}_{\gamma}\delta^{\nu]}_{\rho}, \qquad (123)$$

$$^{[3]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = \delta^{[\alpha} \mathscr{S}^{\beta][\mu} \delta^{\nu]}_{\rho)}, \qquad (124)$$

$${}^{[4]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = \overset{+}{\mathscr{L}}{}^{\alpha\beta}\delta^{[\mu}\gamma^{\nu]} + \overset{+}{\mathscr{L}}{}^{\mu\nu}\delta^{[\alpha}\delta^{\beta]} + \delta^{[\alpha}_{[\gamma}\overset{a}{\mathscr{R}}{}^{\beta][\mu}\delta^{\nu]}_{\rho]}, \qquad (125)$$

$${}^{[5]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = 0, \tag{126}$$

$${}^{[6]}\chi^{\alpha\beta}\gamma^{\mu\nu}{}_{\rho} = \mathscr{A}^{[\alpha\beta}\delta^{\mu}\gamma\delta^{\nu]}_{\rho}.$$
(127)

Here we denoted

$$\mathscr{A}^{\alpha\beta} := 2\mathscr{L}^{\alpha\beta} + \mathscr{M}^{\alpha\beta} - \mathscr{N}^{\alpha\beta}, \qquad (128)$$

$$\overset{s}{\mathscr{R}}^{\alpha\beta} := \overset{+}{\mathscr{M}}^{\alpha\beta} + \overset{+}{\mathscr{N}}^{\alpha\beta}, \qquad (129)$$

$$\overset{a}{\mathscr{R}}^{\alpha\beta} := \overset{+}{\mathscr{M}}^{\alpha\beta} - \overset{+}{\mathscr{N}}^{\alpha\beta}, \qquad (130)$$

$$\mathscr{S}^{\alpha\beta} := \mathscr{M}^{\alpha\beta} + \mathscr{N}^{\alpha\beta}, \qquad (131)$$

$$\mathscr{J}^{\alpha\beta} := \widetilde{\mathcal{M}}^{\alpha\beta} - \widetilde{\mathcal{N}}^{\alpha\beta}.$$
(132)

The first object (128) represents a skewonic axion, whereas the four other objects (129) to (132) describe contributions to the principal reversible and irreversible (skewon) parts. In the reversible case, only the first irreducible piece of Eq.(122) survives and the fourth one of Eq.(125).

4 Metric-dependent constitutive tensor

In general relativity as well as in teleparallelism, the existence of a metric tensor g is conventionally assumed. We choose the metric signature (+, -, -, -). The question is then how the constitutive tensor density $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ can be expressed in terms of the metric, namely in terms of its covariant and/or contravariant components $g_{\alpha\beta}$ and $g^{\gamma\delta}$, respectively. Since χ is a 6th order tensor of type $\binom{4}{2}$, it seems reasonable to start with an ansatz of a purely contravariant dimensionless 6th order tensor $K^{\alpha\beta\mu\gamma\delta\nu}$. Then the contravariant metric components $g^{\gamma\delta}$ are the only metric components that have to be taken into account. In general, one can come up with a polynomial ansatz of arbitrary order in the metric tensor. However, since the resulting constitutive tensor is of 6th order, all indices of the metric factors except six should be necessarily contracted. Recalling that a contraction of the covariant and contravariant metric yields a Kronecker delta, we then find that a polynomial of an arbitrary order is automatically reduced to a general polynomial expression in the metric which is just *cubic* in $g^{\alpha\beta}$.

4.1 A metric-dependent ansatz with parity conserving terms

If we look only for parity conserving (even) pieces in χ , the totally antisymmetric unit tensor is not allowed in the polynomial expression. Thus, for the 6th order tensor $K^{\alpha\beta\mu\gamma\delta\nu}$, the most general cubic expression in the metric, omitting terms explicitly symmetric in the index pairs $\alpha\beta$ and $\gamma\delta$, reads

$$K^{\alpha\beta\mu\gamma\delta\nu} = \alpha_{1}g^{\alpha\mu}g^{\beta\gamma}g^{\delta\nu} + \alpha_{2}g^{\alpha\mu}g^{\beta\delta}g^{\gamma\nu} + \alpha_{3}g^{\alpha\gamma}g^{\beta\mu}g^{\delta\nu} + \alpha_{4}g^{\alpha\gamma}g^{\beta\delta}g^{\mu\nu} + \alpha_{5}g^{\alpha\gamma}g^{\beta\nu}g^{\mu\delta} + \alpha_{6}g^{\alpha\delta}g^{\beta\mu}g^{\gamma\nu} + \alpha_{7}g^{\alpha\delta}g^{\beta\gamma}g^{\mu\nu} + \alpha_{8}g^{\alpha\delta}g^{\beta\nu}g^{\gamma\mu} + \alpha_{9}g^{\alpha\nu}g^{\beta\gamma}g^{\mu\delta} + \alpha_{10}g^{\alpha\nu}g^{\beta\delta}g^{\mu\gamma},$$
(133)

where $\alpha_1, \ldots, \alpha_{10}$ are constants.

In order to tailor the ansatz (133) for our constitutive tensor $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$, we lower the indices μ and ν and, at the same time, we reorder terms with only one metric tensor such that the metric tensor is sandwiched in between two Kronecker deltas:

$$K^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = \alpha_{1}\delta^{\alpha}{}_{\mu}g^{\beta\gamma}\delta^{\delta}{}_{\nu} + \alpha_{2}\delta^{\alpha}{}_{\mu}g^{\beta\delta}\delta^{\gamma}{}_{\nu} + \alpha_{3}\delta^{\beta}{}_{\mu}g^{\alpha\gamma}\delta^{\delta}{}_{\nu} + \alpha_{4}g^{\alpha\gamma}g^{\beta\delta}g_{\mu\nu} + \alpha_{5}\delta^{\beta}{}_{\nu}g^{\alpha\gamma}\delta^{\delta}{}_{\mu} + \alpha_{6}\delta^{\beta}{}_{\mu}g^{\alpha\delta}\delta^{\gamma}{}_{\nu} + \alpha_{7}g^{\alpha\delta}g^{\beta\gamma}g_{\mu\nu} + \alpha_{8}\delta^{\beta}{}_{\nu}g^{\alpha\delta}\delta^{\gamma}{}_{\mu} + \alpha_{9}\delta^{\alpha}{}_{\nu}g^{\beta\gamma}\delta^{\delta}{}_{\mu} + \alpha_{10}\delta^{\alpha}{}_{\nu}g^{\beta\delta}\delta^{\gamma}{}_{\mu}.$$
(134)

Since $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$ is antisymmetric in the index pairs $\alpha\beta$ and $\gamma\delta$, we have to antisymmetrize *K* correspondingly by subsequently putting brackets around the indices $\alpha\beta$ and $\gamma\delta$, respectively. Also recall that the metric is symmetric, hence its index ordering is optional:

$$K^{[\alpha\beta]}{}_{\mu}{}^{[\gamma\delta]}{}_{\nu} = \alpha_{1}\delta^{[\alpha}{}_{\mu}g^{\beta][\gamma}\delta^{\delta]}_{\nu} + \alpha_{2}\delta^{[\alpha}{}_{\mu}g^{\beta][\delta}\delta^{\gamma]}_{\nu} + \alpha_{3}\delta^{[\beta}{}_{\mu}g^{\alpha][\gamma}\delta^{\delta]}_{\nu} + \alpha_{4}g^{\gamma[\alpha}{}_{g}{}^{\beta]\delta}{}_{g\mu\nu} + \alpha_{5}\delta^{[\beta}{}_{\nu}g^{\alpha][\gamma}\delta^{\delta]}_{\mu} + \alpha_{6}\delta^{[\beta}{}_{\mu}g^{\alpha][\delta}\delta^{\gamma]}_{\nu} + \alpha_{7}g^{\delta[\alpha}{}_{g}{}^{\beta]\gamma}{}_{g\mu\nu} + \alpha_{8}\delta^{[\beta}{}_{\nu}g^{\alpha][\delta}\delta^{\gamma]}_{\mu} + \alpha_{9}\delta^{[\alpha}{}_{\nu}g^{\beta][\gamma}\delta^{\delta]}_{\mu} + \alpha_{10}\delta^{[\alpha}{}_{\nu}g^{\beta][\delta}\delta^{\gamma]}_{\mu}.$$
(135)

Note that the α_4 and the α_7 terms are already antisymmetric in $\gamma\delta$. Thus, there is no need to put brackets [] around $\gamma\delta$ or $\delta\gamma$, respectively. We can now collect all terms by recalling the antisymmetry in both index pairs $\alpha\beta$ and $\gamma\delta$. We find, without any further intermediary step, the compact equation

$$K^{[\alpha\beta]}{}_{\mu}{}^{[\gamma\delta]}{}_{\nu} = (\alpha_4 - \alpha_7) g^{\gamma[\alpha} g^{\beta]\delta} g_{\mu\nu} + (\alpha_1 - \alpha_2 - \alpha_3 + \alpha_6) \delta^{[\alpha}{}_{\mu} g^{\beta][\gamma} \delta^{\delta]}_{\nu} + (-\alpha_5 + \alpha_8 + \alpha_9 - \alpha_{10}) \delta^{[\alpha}{}_{\nu} g^{\beta][\gamma} \delta^{\delta]}_{\mu}.$$
(136)

We know that χ is a density, like $\sqrt{-g}$, with $g := \det g_{\mu\nu}$. Accordingly, we can identify the parity even part of the constitutive tensor density as follows,

$$^{\text{even}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{\sqrt{-g}}{\varkappa}K^{[\alpha\beta]}{}_{\mu}{}^{[\gamma\delta]}{}_{\nu}, \qquad (137)$$

with \varkappa as the gravitational constant. In our formalism up to now, including the Tonti diagram, \varkappa was put to 1 for simplicity. Here we do display it for clarity. Thus, we find, with the three constants,

$$\beta_{1} := \alpha_{4} - \alpha_{7} \beta_{2} := \alpha_{1} - \alpha_{2} - \alpha_{3} + \alpha_{6} \beta_{3} := -\alpha_{5} + \alpha_{8} + \alpha_{9} - \alpha_{10},$$

$$(138)$$

the expression

$$^{\text{even}} \chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{\sqrt{-g}}{\varkappa} \left(\beta_1 g^{\gamma[\alpha} g^{\beta]\delta} g_{\mu\nu} + \beta_2 \delta^{[\alpha}{}_{\mu} g^{\beta][\gamma} \delta^{\delta]}_{\nu} + \beta_3 \delta^{[\alpha}{}_{\nu} g^{\beta][\gamma} \delta^{\delta]}_{\mu} \right).$$
(139)

This proves our ansatz with three independent constants in [31, Eq.(80)].⁵ It is, indeed, the most general expression which, up to a factor of $\sqrt{-g}$, turns out to be a *cubic* polynomial in the metric.

4.2 Parity violating terms

Already for quite some time, see Pellegrini & Plebanski [48] and Müller-Hoissen & Nitsch [40], also parity violating (odd) constitutive tensors for torsion square Lagrangians have been considered in the literature; for recent reviews, see [3, 7, 32, 45]. There it has been shown that *two* parity odd terms occur, which are both linear in the totally antisymmetric Levi-Civita symbol $\varepsilon^{\alpha\beta\gamma\delta}$, which is, as we may recall, a tensor density:

$${}^{\text{odd}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{1}{\varkappa} \left\{ \beta_4 \,\varepsilon^{\alpha\beta\gamma\delta} \,g_{\mu\nu} + \beta_5 \,\varepsilon^{\alpha\beta[\gamma}{}_{[\mu} \,\delta^{\delta]}_{\nu]} \right\}. \tag{140}$$

One can also take into account such additional parity violating terms in the framework of the Poincaré gauge theory of gravity. In that more general theory, it leads to interesting new cosmological models [4, 25] and to new gravitational wave solutions [6, 43].

⁵In [31], there was a typo in Eq.(80): the last plus sign + in this formula should have been a minus sign -. Moreover, for simplification, we changed now our conventions with respect to the β 's slightly: $4\beta_1^{\text{old}} = \beta_1$, $8\beta_2^{\text{old}} = -\beta_2$, $8\beta_3^{\text{old}} = -\beta_3$.

We would like to understand the generality of the result in (140) in a similar way as we did it for the parity even case. For parity reasons, the Levi-Civita symbol has to be linearly or in odd powers in the corresponding ansatz. Since again, as in the even part of $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}$, we start with an tensor expression of type $\begin{pmatrix} 6\\0 \end{pmatrix}$, we need additionally a tensor of type $\begin{pmatrix} 2\\0 \end{pmatrix}$. Clearly the contravariant components of the metric $g^{\alpha\beta}$ qualify for such a purpose. Thus, the simplest possible ansatz, see also (140), is to start with a 6th order tensor density as follows:

$$M^{\alpha\beta\mu\gamma\delta\nu} = \mu_{1}g^{\alpha\mu}\varepsilon^{\beta\gamma\delta\nu} + \mu_{2}g^{\alpha\gamma}\varepsilon^{\beta\mu\delta\nu} + \mu_{3}g^{\alpha\delta}\varepsilon^{\beta\mu\gamma\nu} + \mu_{4}g^{\alpha\nu}\varepsilon^{\beta\mu\gamma\delta} + \mu_{5}g^{\beta\mu}\varepsilon^{\alpha\gamma\delta\nu} + \mu_{6}g^{\beta\gamma}\varepsilon^{\alpha\mu\delta\nu} + \mu_{7}g^{\beta\delta}\varepsilon^{\alpha\mu\gamma\nu} + \mu_{8}g^{\beta\nu}\varepsilon^{\alpha\mu\gamma\delta} + \mu_{9}g^{\mu\gamma}\varepsilon^{\alpha\beta\delta\nu} + \mu_{10}g^{\mu\delta}\varepsilon^{\alpha\beta\gamma\nu} + \mu_{11}g^{\mu\nu}\varepsilon^{\alpha\beta\gamma\delta} + \mu_{12}g^{\gamma\nu}\varepsilon^{\alpha\beta\mu\delta} + \mu_{13}g^{\delta\nu}\varepsilon^{\alpha\beta\mu\gamma}.$$
(141)

The $\mu_1, \mu_2, \dots, \mu_{13}$ are arbitrary constants. We now lower again the indices μ and ν and antisymmetrize at the same time with respect to the index pairs $\alpha\beta$ and $\gamma\delta$, respectively:

$$M^{[\alpha\beta]}{}_{\mu}{}^{[\gamma\delta]}{}_{\nu} = \mu_{11}g_{\mu\nu}\varepsilon^{\alpha\beta\gamma\delta} + (\mu_{1} - \mu_{5}) \,\delta^{[\alpha}{}_{\mu}\varepsilon^{\beta]\gamma\delta}{}_{\nu} + (\mu_{4} - \mu_{8}) \,\delta^{[\alpha}{}_{\nu}\varepsilon^{\beta]\gamma\delta}{}_{\mu} + (\mu_{9} - \mu_{10}) \,\delta^{[\gamma}{}_{\mu}\varepsilon^{\delta]\alpha\beta}{}_{\nu} - (\mu_{12} - \mu_{13}) \,\delta^{[\gamma}{}_{\nu}\varepsilon^{\delta]\alpha\beta}{}_{\mu} + \frac{1}{2}(-\mu_{2} + \mu_{3} + \mu_{6} - \mu_{7}) \left(g^{\gamma[\alpha}{}_{\nu}\varepsilon^{\beta]\delta}{}_{\mu\nu} - g^{\delta[\alpha}{}_{\nu}\varepsilon^{\beta]\gamma}{}_{\mu\nu}\right).$$
(142)

Having established the required antisymmetries, we can now identify the odd piece of the constitutive tensor as

$${}^{\text{odd}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{1}{\varkappa} M^{[\alpha\beta]}{}_{\mu}{}^{[\gamma\delta]}{}_{\nu}.$$
(143)

Here we have to recall an algebraic trick that we had already applied in discussing the electromagnetic 4th order constitutive tensor, see [22, Eq.(D.1.65)]. Since in four dimensions the indices have always four values, any expression antisymmetrized over five indices vanishes identically. Thus, for an arbitrary tensor $T^{\alpha\beta\gamma\delta\varepsilon\cdots}_{\cdots}$, we have $T^{[\alpha\beta\gamma\delta\varepsilon]\cdots}_{\cdots} \equiv 0$. Accordingly, we have, for example, $T^{[\rho}\varepsilon^{\alpha\beta\gamma\delta]} = 0$ or, if we evaluate the brackets,

$$T^{\rho}\varepsilon^{\alpha\beta\gamma\delta} \equiv T^{\alpha}\varepsilon^{\rho\beta\gamma\delta} + T^{\beta}\varepsilon^{\alpha\rho\gamma\delta} + T^{\gamma}\varepsilon^{\alpha\beta\rho\delta} + T^{\delta}\varepsilon^{\alpha\beta\gamma\rho}.$$
(144)

Looking now at (142), we have typically $\delta^{[\alpha}_{\mu} \varepsilon^{\beta]\gamma\delta\nu}$. With $T^{\alpha}{}_{\mu} = \delta^{\alpha}_{\mu}$, we find

$$\delta^{[\alpha}_{\mu}\varepsilon^{\beta]\gamma\delta\nu} \equiv \delta^{[\beta}_{\mu}\varepsilon^{\alpha]\gamma\delta\nu} + \delta^{\gamma}_{\mu}\varepsilon^{\beta\alpha\delta\nu} + \delta^{\delta}_{\mu}\varepsilon^{\beta\gamma\alpha\nu} + \delta^{\nu}_{\mu}\varepsilon^{\beta\gamma\delta\alpha} \,. \tag{145}$$

By such methods, we can derive several identities for the Levi-Civita symbol in 4 dimensions:

$$\varepsilon^{\alpha\beta[\gamma}{}_{(\mu}\,\delta^{\delta]}_{\nu)} + \varepsilon^{\gamma\delta[\alpha}{}_{(\mu}\,\delta^{\beta]}_{\nu)} \equiv \frac{1}{2}\,\varepsilon^{\alpha\beta\gamma\delta}\,g_{\mu\nu},\tag{146}$$

$$\varepsilon^{\alpha\beta[\gamma}{}_{[\mu}\,\delta^{\delta]}_{\nu]} + \varepsilon^{\gamma\delta[\alpha}{}_{[\mu}\,\delta^{\beta]}_{\nu]} \equiv 0, \qquad (147)$$

$$g^{\gamma[\alpha}\varepsilon^{\beta]\delta}{}_{\mu\nu} - g^{\delta[\alpha}\varepsilon^{\beta]\gamma}{}_{\mu\nu} \equiv 2\varepsilon^{\alpha\beta[\gamma}{}_{[\mu}\delta^{\delta]}{}_{\nu]}.$$
(148)

Let us come back to (143) with (142). We decompose the $\delta\varepsilon$ -terms in symmetric and antisymmetric pieces with respect to the lower indices:

$$\delta^{[\alpha}_{\mu}\varepsilon^{\beta]\gamma\delta}_{\nu} = \delta^{[\alpha}_{(\mu}\varepsilon^{\beta]\gamma\delta}_{\nu)} + \delta^{[\alpha}_{[\mu}\varepsilon^{\beta]\gamma\delta}_{\nu]}, \qquad (149)$$

$$\delta^{[\gamma}_{\mu}\varepsilon^{\delta]\alpha\beta}{}_{\nu} = \delta^{[\gamma}_{(\mu}\varepsilon^{\delta]\alpha\beta}{}_{\nu)} + \delta^{[\gamma}_{[\mu}\varepsilon^{\delta]\alpha\beta}{}_{\nu]}.$$
(150)

As a result, the $\delta \varepsilon$ -terms in *the second and third lines of* (142) can be rearranged as follows:

$$(\mu_{1} - \mu_{5} + \mu_{4} - \mu_{8}) \,\delta^{[\alpha}_{(\mu} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} + (\mu_{9} - \mu_{10} - \mu_{12} + \mu_{13}) \,\delta^{[\gamma}_{(\mu} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} + (\mu_{1} - \mu_{5} - \mu_{4} + \mu_{8}) \,\delta^{[\alpha}_{[\mu} \varepsilon^{\beta]\gamma\delta}{}_{\nu]} + (\mu_{9} - \mu_{10} + \mu_{12} - \mu_{13}) \,\delta^{[\gamma}_{[\mu} \varepsilon^{\delta]\alpha\beta}{}_{\nu]}.$$
(151)

Next, we write

$$\delta_{(\mu}^{[\alpha} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} = \frac{1}{2} \left(\delta_{(\mu}^{[\alpha} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} + \delta_{(\mu}^{[\gamma} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} \right) + \frac{1}{2} \left(\delta_{(\mu}^{[\alpha} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} - \delta_{(\mu}^{[\gamma} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} \right), \qquad (152)$$

$$\delta_{(\mu}^{[\gamma} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} = \frac{1}{2} \left(\delta_{(\mu}^{[\gamma} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} + \delta_{(\mu}^{[\alpha} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} \right) + \frac{1}{2} \left(\delta_{(\mu}^{[\gamma} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} - \delta_{(\mu}^{[\alpha} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} \right).$$
(153)

Consequently, the expression (151) transforms into

$$(\mu_{1} - \mu_{5} - \mu_{4} + \mu_{8} - \mu_{9} + \mu_{10} - \mu_{12} + \mu_{13}) \,\delta^{[\alpha}_{[\mu} \varepsilon^{\beta]\gamma\delta}{}_{\nu]} + \frac{1}{2} (\mu_{1} - \mu_{5} + \mu_{4} - \mu_{8} - \mu_{9} + \mu_{10} + \mu_{12} - \mu_{13}) \times \\ \left(\delta^{[\alpha}_{(\mu} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} - \delta^{[\gamma}_{(\mu} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} \right) \\ + \frac{1}{2} (\mu_{1} - \mu_{5} + \mu_{4} - \mu_{8} + \mu_{9} - \mu_{10} - \mu_{12} + \mu_{13}) \times \\ \left(\delta^{[\alpha}_{(\mu} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} + \delta^{[\gamma}_{(\mu} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} \right).$$
(154)

where we used the identity (147). We recall that (154) belongs to the second and third line of (142).

In the last step, it remains to use (146) in the last line of (154) and to substitute (148) into the last line of (142). Then, with the constants

$$\begin{split} \beta_4 &:= \frac{1}{4} \left(\mu_1 - \mu_5 + \mu_4 - \mu_8 + \mu_9 - \mu_{10} - \mu_{12} + \mu_{13} + 4\mu_{11} \right) ,\\ \beta_5 &:= \mu_1 - \mu_5 - \mu_4 + \mu_8 - \mu_9 + \mu_{10} \\ &+ \mu_{12} - \mu_{13} + \mu_2 - \mu_6 - \mu_3 + \mu_7 \,,\\ \beta_6 &:= \frac{1}{2} \left(\mu_1 - \mu_5 + \mu_4 - \mu_8 - \mu_9 + \mu_{10} + \mu_{12} - \mu_{13} \right) , \end{split}$$

the result eventually reads

$$^{\text{odd}} \chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{\alpha}{\varkappa} \left[\beta_{4} \varepsilon^{\alpha\beta\gamma\delta} g_{\mu\nu} + \beta_{5} \varepsilon^{\alpha\beta[\gamma}{}_{[\mu} \delta^{\delta]}_{\nu]} + \beta_{6} \left(\delta^{[\alpha}_{(\mu} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} - \delta^{[\gamma}_{(\mu} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} \right) \right].$$
(155)

Here α is dimensionless pseudoscalar. It is necessary in order to take into account the transformation law of this part under improper reflections.

Therefore, the most general parity-odd part of the constitutive tensor in the end boils down to just the three independent terms with β_4 , β_5 , and β_6 . The β_4 -term, being totally antisymmetric in $\alpha\beta\gamma\delta$, represents an β_4 -axion, whereas the β_5 -term describes a β_5 -axion. They both represent reversible processes. In contrast, the β_6 -term corresponds to *irreversibility*. When contracted by $F_{\alpha\beta}{}^{\mu}$ and $F_{\gamma\delta}{}^{\nu}$, this term drops out from the general teleparallel Lagrangian. Moreover, one can reduce the number of independent terms in the Lagrangian by making use of the Nieh-Yan topological invariant [41, 42, 46].

4.3 The general case and its irreducible decomposition

The general constitutive tensor

$$\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = {}^{\text{even}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) + {}^{\text{odd}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g)$$
(156)

encompasses the parity-even piece of Eq.(139) and the parity-odd piece of Eq.(155),

Computing the irreducible parts, we find

$${}^{[2]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = {}^{[6]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = 0, \qquad (157)$$

whereas the nontrivial pieces read

$$^{[1]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{\sqrt{-g}}{\varkappa} \Big[\beta_1 g^{\gamma[\alpha} g^{\beta]\delta} g_{\mu\nu} + (\beta_2 + \beta_3) \,\delta^{[\alpha}_{(\mu} g^{\beta][\gamma} \delta^{\delta]}_{\nu)}\Big], \tag{158}$$

$$^{[3]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{\beta_{6}}{\varkappa} \left(\delta^{[\alpha}_{(\mu} \varepsilon^{\beta]\gamma\delta}{}_{\nu)} - \delta^{[\gamma}_{(\mu} \varepsilon^{\delta]\alpha\beta}{}_{\nu)} \right), \tag{159}$$

$$\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{1}{2} \left[(\beta_2 - \beta_2) \sqrt{-g} \delta^{[\alpha}{}_{\beta}{}^{\beta][\gamma} \delta^{\delta]} + \beta_5 \varepsilon^{\alpha\beta[\gamma}{}_{\mu} \delta^{\delta]} \right], \tag{160}$$

$$\frac{1}{\varkappa} \Big[(\beta_2 - \beta_3) \sqrt{-g} \delta^{[\alpha}_{[\mu} g^{\beta][\gamma} \delta^{o]}_{\nu]} + \beta_5 \varepsilon^{\alpha\beta[\gamma}_{[\mu} \delta^{o]}_{\nu]} \Big], \qquad (160)$$

$${}^{[5]}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g) = \frac{p_4}{\varkappa}\varepsilon^{\alpha\beta\gamma\delta}g_{\mu\nu}.$$
(161)

Summarizing, a general metric-dependent constitutive tensor encompasses two principal ${}^{[1]}\chi$ and ${}^{[4]}\chi$ parts, one axion ${}^{[5]}\chi$ part, and one skewon ${}^{[3]}\chi$ part.

Calculating the traces (105)–(108) of the constitutive tensor (156), we obtain

$$m^{\alpha\beta} = \frac{3\sqrt{-g}}{4\varkappa} (2\beta_1 - 3\beta_2 - \beta_3) g^{\alpha\beta}, \qquad (162)$$

and

$$n^{\alpha\beta} = \frac{3\sqrt{-g}}{4\varkappa} (2\beta_1 - \beta_2 - 3\beta_3)g^{\alpha\beta}.$$
(163)

The antisymmetric tensors vanish, $k^{\alpha\beta} = l^{\alpha\beta} = 0$. The parity odd terms drops out altogether in the trace building.

4.4 Specific Lagrangians.

4.4.1 GR_{\parallel}: the teleparallel equivalent of GR

We found in the framework of the *teleparallel equivalent* GR_{\parallel} of general relativity (GR) for the β 's in [31, Eq.(88)]⁶,

$$\begin{cases} \beta_1 = -1, & \beta_2 = -4, & \beta_3 = 2\\ \beta_4 = 0, & \beta_5 = 0, & \beta_6 = 0 \end{cases}$$
 GR_{||}. (164)

Accordingly, the three β 's are related to GR in a quite definite way, and we get a feeling for their interpretation.

The GR_{\parallel} Lagrangian is distinguished from the other teleparallelism Lagrangians as being *locally* Lorentz invariant, see, e.g., Cho [11] and Müller-Hoissen & Nitsch [40]. Any other additional term of even parity in the gravitational Lagrangian removes this local invariance.

Ferrarro and Guzmán [14] called the constitutive tensor $\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu}(g)$ of (156) together with the parameters of (164) the *supermetric*. In fact, up to a conventional factor of -2, our result agrees with their Eq.(17) in [14].

4.4.2 Von der Heyde Lagrangian

Earlier, different parity even pieces of torsion square Lagrangians were compared by Muench et al. [39], see also the literature cited therein. A particular role played in these considerations the torsion square piece of a Lagrangian of *von der Heyde* [24]. Its constitutive tensor [17] reads

$${}^{\mathrm{vdH}}\chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} = -\frac{2\sqrt{-g}}{\varkappa} \left(g^{\gamma[\alpha}g^{\beta]\delta}g_{\mu\nu} + 2\delta^{[\alpha}_{\mu}g^{\beta][\gamma}\delta^{\delta]}_{\nu} \right).$$
(165)

For the corresponding constitutive law, we have

$$\check{H}^{\alpha\beta}{}_{\mu} = \frac{1}{2} {}^{\nu dH} \chi^{\alpha\beta}{}_{\mu}{}^{\gamma\delta}{}_{\nu} F_{\gamma\delta}{}^{\nu}
= -\frac{\sqrt{-g}}{\varkappa} \left(F^{\alpha\beta}{}_{\mu} + 2\delta^{[\alpha}{}_{\mu}F^{\beta]}{}_{\gamma}{}^{\gamma} \right).$$
(166)

This law, in the teleparallel case, leads already to the correct Newtonian approximation. The constitutive tensor (165) carries the following β -values:

$$\begin{cases} \beta_1 = -2, & \beta_2 = -4, & \beta_3 = 0\\ \beta_4 = 0, & \beta_5 = 0, & \beta_6 = 0 \end{cases}$$
 vdH (167)

As we saw, the teleparallel equivalent $GR_{||}$ has a slightly different constitutive law (164).

⁶There occurred an computing error. The correct values in the old conventions are $\beta_1^{\text{old}} = -\frac{1}{4}$, $\beta_2^{\text{old}} = \frac{1}{2}$, $\beta_3^{\text{old}} = -\frac{1}{4}$.

4.4.3 Torsion-square Lagrangian in nonlocal gravity (NLG)

In a nonlocal extension of GR—*Nonlocal Gravity (NLG)* —the starting point [19, 20] was the parameter set of GR_{||}, namely (164). Later, Mashhoon found it necessary, to enlarge it by adding a supplementary piece proportional to the axial⁷ torsion ${}^{(3)}F_{\alpha\beta}{}^{\mu}$, see [37, Eq.(6.109)]. This yields, with an unknown dimensionless parameter \check{p} ,

$$\beta_{1} = -1, \quad \beta_{2} = -4, \qquad \beta_{3} = 2, \\ \beta_{4} = -\frac{\check{p}}{6}, \quad \beta_{5} = -4\frac{\check{p}}{6}, \quad \beta_{6} = 2\frac{\check{p}}{6}.$$
 Mashh. (168)

It is amazing that $(\beta_4, \beta_5, \beta_6) = \frac{p}{6}(\beta_1, \beta_2, \beta_3)$. Probably this means that Mashhoon's additional parity odd term is locally Lorentz invariant. Incidentally, there is a conventional factor between Mashhoon's p and ours.

According to Mashhoon,⁸ "...the mere fact of postulating a *nonlocal* constitutive relation violates local Lorentz invariance." Still, the version of NLG of 2009 is based on the reversible and locally Lorentz invariant parameter set (164) and Mashhoon's version corresponds to an irreversible and possibly also locally Lorentz invariant parameter set, see the 2nd line of (168).

5 Propagation of gravitational waves

In this section, we discuss the wave propagation in our premetric teleparallel theory (TG) in linear approximation.

5.1 Geometric optics approximation

We start with the source-free field equations

$$dH_{\alpha} = 0$$
 and $dF^{\alpha} = 0.$ (169)

Notice that the source term Σ in (2) contains two independent expressions: the matter energy-momentum current ${}^{(m)}\Sigma$ and the gravitational energy-momentum current ${}^{(\vartheta)}\Sigma$. In matter-free regions, we have ${}^{(m)}\Sigma = 0$. Similar to standard GR, we will assume, in addition, ${}^{(\vartheta)}\Sigma = 0$. This requirement is applicable for small waves and means linearization of the field equations.

In the geometric optics approximation (and, equivalently, in Hadamard's approach, see [22]), one derives the linear system

$$\check{H}^{\mu\nu}{}_{\alpha}q_{\nu} = 0 \quad \text{and} \quad \varepsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}{}^{\alpha}q_{\nu} = 0.$$
 (170)

⁷In exterior calculus, we have ${}^{(3)}F^{\alpha} = \frac{1}{3}e^{\alpha} \rfloor (F^{\beta} \wedge \vartheta_{\beta}).$

⁸Private communication, August 2018.

Here the components q_v of the wave covector are determined by the differential of the phase φ function of the wave: $d\varphi = q_v \vartheta^v$.

The second equation of (170) has the solution

$$F_{\rho\sigma}{}^{\alpha} = A_{\rho}{}^{\alpha} q_{\sigma} - A_{\sigma}{}^{\alpha} q_{\rho} , \qquad (171)$$

with an arbitrary tensor A_{β}^{α} . It is a gravitational analog of the electromagnetic potential. We observe the *gauge invariance* of this gravitational potential. An expression of the form $A_{\beta}^{\alpha} = q_{\beta}C^{\alpha}$ does not contribute to the field strength $F_{\rho\sigma}^{\alpha}$. In other words, the model is invariant under the transformations

$$A_{\beta}{}^{\alpha} \to A_{\beta}{}^{\alpha} + q_{\beta}C^{\alpha} \,, \tag{172}$$

with an arbitrary vector C^{α} .

We substitute (171) into (170) and use the constitutive relation

$$\check{H}^{\mu\nu}{}_{\alpha} = \frac{1}{2} \chi^{\mu\nu}{}_{\alpha}{}^{\rho\sigma}{}_{\beta} F_{\rho\sigma}{}^{\beta} .$$
(173)

This yields the characteristic equation

$$M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta}A_{\nu}{}^{\beta}=0, \qquad (174)$$

with fourth order characteristic tensor

$$M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta} := \chi^{\mu\rho}{}_{\alpha}{}^{\nu\sigma}{}_{\beta} q_{\rho} q_{\sigma}.$$
(175)

Inserting here the decomposition (38) of the constitutive tensor, we observe that the axion-type parts ${}^{[5]}\chi^{\mu\rho}{}_{\alpha}{}^{\nu\sigma}{}_{\beta}$ and ${}^{[6]}\chi^{\mu\rho}{}_{\alpha}{}^{\nu\sigma}{}_{\beta}$ do not enter the tensor $M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta}$. Consequently, these parts do not contribute to the wave propagation as it was already outlined in Table 1.

In the reversible case, the characteristic tensor $M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta}$ satisfies the symmetry relations

$$M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta} = M^{\nu}{}_{\beta}{}^{\mu}{}_{\alpha}.$$
(176)

A generalized Fresnel equation can be derived as the condition for the solvability of the equation (174) along the lines of the algebraic computations of Itin [30].

5.2 Dispersion relation—general facts

Observe that the characteristic equation

$$E^{\mu}{}_{\alpha} = 0, \quad \text{with} \quad E^{\mu}{}_{\alpha} := M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta}A_{\nu}{}^{\beta}, \qquad (177)$$

represents 16 equations for the 16 variables A_{ν}^{β} .

For a compact representation of this system, we denote a pair of upper and lower indices by a multi-index

$${{}^{\mu}}_{\alpha} = I \qquad I = 1, \cdots 16.$$
 (178)

In this notation, the system (177) reads

$$E^{I} = M^{IJ}A_{J} = 0. (179)$$

The gauge transformation (172) can be rewritten as

$$A_J \to A_J + Q_J$$
, where $Q_J = q_\beta C^\alpha$. (180)

The 4 linearly independent vectors C^{α} imply that there are likewise 4 linearly independent vectors Q_J . The identities

$$M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta}\,q_{\mu} = M^{\nu}{}_{\alpha}{}^{\mu}{}_{\beta}\,q_{\mu} = 0 \tag{181}$$

translate into

$$M^{IJ}Q_J = M^{JI}Q_J = 0. (182)$$

These equations can be understood as 4 linear relations between the rows (and the columns) of the matrix M^{IJ} . Consequently, the matrix M^{IJ} has rank 12 = 16 - 4. The condition for the existence of non-trivial solutions of Eq.(179) reads now as

$$^{(4)}\mathrm{Adj}(M) = 0. \tag{183}$$

We used here the fourth order adjoint of the matrix M^{IJ} . It can be constructed by evaluating the determinants of the matrices that are left in the 16 × 16 matrix M^{IJ} by removing four rows and four columns. Consequently ⁽⁴⁾Adj(M) is a set of 12th order polynomials of the entries M^{IJ} . Accordingly, it is a set of 12th order polynomials of the variables q_{α} . The 4th order adjoint can be written as a tensor with eight free indices:

⁽⁴⁾Adj(
$$M$$
) _{$I_1 \cdots I_4 J_1 \cdots J_4 = 0$. (184)}

In electromagnetism with only one gauge invariance constraint, there appears the 1st order adjoint matrix $\operatorname{Adj}(M)_{\alpha\beta}$. This matrix is expressed as a scalar function multiplied by a tensor product of q_{α} , namely $\operatorname{Adj}(M)_{\alpha\beta} = \lambda(q)q_{\alpha}q_{\beta}$. Thus, the electromagnetic dispersion relation takes the form $\lambda(q) = 0$.

Similarly, in the case of the gravitational equation (184), we have

⁽⁴⁾Adj
$$(M)_{I_1\cdots I_4J_1\cdots J_4} = \Lambda(Q)Q_{I_1}\cdots Q_{I_4}Q_{J_1}\cdots Q_{J_4} = 0.$$
 (185)

Thus, the gravitational dispersion relation takes the form

$$\Lambda(q) = 0. \tag{186}$$

Here $\Lambda(q)$ is a homogeneous polynomial of the order 16 = 24 - 8 in the variable q_{α} . We recall that in the electromagnetic case, the corresponding form is of 4th order. This fact yields birefringence in the wave propagation. In 3-dimensional elasticity theory, the dispersion relation is of 6th order, with 3 different waves in general. In our generalized gravitational model, there are 8 different waves in general.

In the simplest case, Eq.(186) reads $(g^{\alpha\beta}q_{\alpha}q_{\beta})^8 = 0$. We discuss the metrical case in Sec.6.

5.3 Dispersion relation decomposed

In order to clarify the nature of the equation (177), we decompose the solution under the GL(4, \mathbb{R}) irreducibly into the scalar $A\delta^{\beta}_{\alpha}$ and the traceless part $A\alpha^{\beta}$,

$$A_{\alpha}{}^{\beta} = \mathcal{A}_{\alpha}{}^{\beta} + A\delta_{\alpha}^{\beta}$$
, with $A := \frac{1}{4}A_{\gamma}{}^{\gamma}$ and $\mathcal{A}_{\alpha}{}^{\alpha} = 0$. (187)

Consequently, Eq.(177) decomposes into

$$E^{\mu}{}_{\alpha} = M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta} \mathcal{A}_{\nu}{}^{\beta} + M^{\mu}{}_{\alpha}{}^{\beta}{}_{\beta}A = 0.$$
(188)

A similar decomposition can be performed with $E^{\mu}{}_{\alpha}$. Thus, we find the 1+15 equations

$$E^{\alpha}{}_{\alpha} = 0 \quad \text{and} \quad E^{\mu}{}_{\alpha} - \frac{1}{4}E^{\gamma}{}_{\gamma}\delta^{\mu}_{\alpha} = 0 \quad (189)$$

or, explicitly, one equation

$$M^{\alpha}{}_{\alpha}{}^{\nu}{}_{\beta} \mathcal{A}_{\nu}{}^{\beta} + M^{\alpha}{}_{\alpha}{}^{\beta}{}_{\beta}A = 0$$
(190)

and 15 equations

$$\begin{pmatrix}
M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta} - \frac{1}{4}\delta^{\mu}_{\alpha}M^{\gamma}{}_{\gamma}{}^{\nu}{}_{\beta} \\
+ \left(M^{\mu}{}_{\alpha}{}^{\beta}{}_{\beta} - \frac{1}{4}\delta^{\mu}_{\alpha}M^{\gamma}{}_{\gamma}{}^{\beta}{}_{\beta} \right)A = 0.$$
(191)

Substituting (190) into (191), we derive the algebraic system for the traceless variable:

$$N^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta} \mathcal{A}_{\nu}{}^{\beta} = 0, \tag{192}$$

where we introduced

$$N^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta} := M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta}M^{\rho}{}_{\rho}{}^{\sigma}{}_{\sigma} - M^{\mu}{}_{\alpha}{}^{\rho}{}_{\rho}M^{\sigma}{}_{\sigma}{}^{\nu}{}_{\beta}.$$
(193)

The latter tensor is apparently traceless for both pairs of indices:

$$N^{\mu}{}_{\alpha}{}^{\rho}{}_{\rho} = 0, \qquad N^{\sigma}{}_{\sigma}{}^{\nu}{}_{\beta} = 0.$$
 (194)

Accordingly, we find a system of 15 algebraic equations for 15 variables A_v^{μ} . After one solves the homogeneous equation (192), one immediately finds the scalar A from (190).

5.4 Scalar waves as a special case

We consider now a special case of pure scalar waves. Let the field A_{ν}^{β} be identically zero: $A_{\nu}^{\beta} = 0$. Hence, we are left with a system of two scalar equations

$$\left(M^{\mu}{}_{\alpha}{}^{\beta}{}_{\beta}-\frac{1}{4}\delta^{\mu}_{\alpha}M^{\gamma}{}_{\gamma}{}^{\beta}{}_{\beta}\right)A=0\,,\quad M^{\alpha}{}_{\alpha}{}^{\beta}{}_{\beta}A=0\,.$$
(195)

The second equation has a non-trivial solution only if $M^{\alpha}{}_{\alpha}{}^{\beta}{}_{\beta} = 0$. Consequently we have a dispersion relation

$$\chi^{\alpha\rho}{}_{\alpha}{}^{\beta\sigma}{}_{\beta}q_{\rho}q_{\sigma} = 0.$$
 (196)

Using the double-trace tensor (105), we can write it as

$$m^{\alpha\beta}q_{\alpha}q_{\beta} = 0. \tag{197}$$

In the metric-dependent case (162), the tensor $m^{\alpha\beta}$, up to a factor, coincides with the Lorentz metric tensor $g^{\alpha\beta}$. Thus we recover the standard light cone of general relativity,

$$g^{\alpha\beta}q_{\alpha}q_{\beta} = 0. \tag{198}$$

5.5 A separable case

We assume now that two terms in the left-hand side of Eqs. (190) and (191) vanish independently. In other words, we assume that the system separates into two independent subsystems. In [28], a similar type of consideration allowed to extract the teleparallel equivalent of GR from a set of metric based models. Thus we have 15 equations for 15 variables

$$\left(M^{\mu}{}_{\alpha}{}^{\nu}{}_{\beta}-\frac{1}{4}\delta^{\mu}_{\alpha}M^{\gamma}{}_{\gamma}{}^{\nu}{}_{\beta}\right)\mathcal{A}_{\nu}{}^{\beta}=0.$$
(199)

and 1 equation for 1 variable

$$M^{\alpha}{}_{\alpha}{}^{\beta}{}_{\beta}A = 0. \tag{200}$$

5.6 Gauge conditions

In electromagnetism, gauge invariance can be restricted by applying a gauge fixing condition. The Lorenz gauge condition is the unique diffeomorphism invariant expression. For wave solutions, it takes the form $A_{\alpha}q^{\alpha} = 0$. Note that this condition can be formulated only on a metric manifold.

In teleparallel theory, we are able to formulate a similar condition in a premetric form:

$$A_{\nu}{}^{\beta}q_{\beta} = 0. \tag{201}$$

A more general gauge condition can be proposed in the form of

$$K_1 A_{\alpha}{}^{\beta} q_{\beta} + K_2 A_{\beta}{}^{\beta} q_{\alpha} = 0.$$
⁽²⁰²⁾

Here K_1 and K_2 are arbitrary constants. The exceptional case for $K_1 = 1$ and $K_2 = -1$, namely

$$A_{\alpha}{}^{\beta}q_{\beta} - A_{\beta}{}^{\beta}q_{\alpha} = 0, \qquad (203)$$

is invariant under the gauge transformation $A_{\alpha}{}^{\beta} \rightarrow A_{\alpha}{}^{\beta} + q_{\alpha}C^{\beta}$ and, accordingly, does not represent a gauge condition.

6 Gravitational waves in a metric model

Let us now specialize to the metric-dependent constitutive tensor (156). The analysis of the wave propagation in the geometric optics approximation is straightforward. Here we will confine our attention to the *parity-even* case (139). Recently a related paper was published by Hohmann et al. [27].

6.1 Characteristic system

We substitute the constitutive tensor (139) with its parity even pieces into (175). As a result, $M^{\beta} {}_{\alpha}{}^{\nu}{}_{\mu}(g) = \chi^{\beta\gamma}{}_{\alpha}{}^{\nu\rho}{}_{\mu}(g) \times q_{\gamma}q_{\rho}$ takes the form

$$M^{\beta}{}_{\alpha}{}^{\nu}{}_{\mu}(g) = \frac{\sqrt{-g}}{4\varkappa} \Big\{ 2\beta_{1}g_{\alpha\mu} \left(g^{\beta\nu}q^{2} - q^{\nu}q^{\beta} \right) \\ -\beta_{2} \left[\left(q^{2}\delta^{\beta}_{\alpha} - q^{\beta}q_{\alpha} \right) \delta^{\nu}_{\mu} - \left(q^{\nu}\delta^{\beta}_{\alpha} - g^{\beta\nu}q_{\alpha} \right) q_{\mu} \right] \\ -\beta_{3} \left[\left(q^{2}\delta^{\beta}_{\mu} - q^{\beta}q_{\mu} \right) \delta^{\nu}_{\alpha} - \left(q^{\nu}\delta^{\beta}_{\mu} - g^{\beta\nu}q_{\mu} \right) q_{\alpha} \Big] \Big\}.$$
(204)

We can immediately derive the contractions

$$M^{\beta}{}_{\alpha}{}^{\nu}{}_{\nu} = M^{\nu}{}_{\nu}{}^{\beta}{}_{\alpha} = \frac{\sqrt{-g}}{4\varkappa} \left(2\beta_1 - 3\beta_2 - \beta_3\right) \left(q^2 \delta^{\beta}_{\alpha} - q^{\beta} q_{\alpha}\right), \tag{205}$$

where $q^2 := g^{\alpha\beta} q_{\alpha} q_{\beta}$. Consequently,

$$M^{\alpha}{}_{\alpha}{}^{\nu}{}_{\nu} = \frac{3\sqrt{-g}}{4\varkappa} \left(2\beta_1 - 3\beta_2 - \beta_3\right) q^2.$$
(206)

Furthermore, we verify that the tensor (204) has the following properties:

$$M^{\beta}{}_{\alpha}{}^{\nu}{}_{\mu}q_{\nu} = 0, \qquad M^{\beta}{}_{\alpha}{}^{\nu}{}_{\mu}q_{\beta} = 0, \tag{207}$$

$$M^{\beta}{}_{\alpha}{}^{\nu}{}_{\mu}q^{\mu} = \frac{\sqrt{-g}}{4\varkappa} (2\beta_1 - \beta_2 - \beta_3) \left(q^2 g^{\beta\nu} - q^{\beta} q^{\nu}\right) q_{\alpha}, \qquad (208)$$

$$M^{\beta}{}_{\alpha}{}^{\nu}{}_{\mu}q^{\alpha} = \frac{\sqrt{-g}}{4\varkappa} (2\beta_1 - \beta_2 - \beta_3) \left(q^2 g^{\beta\nu} - q^{\beta} q^{\nu}\right) q_{\mu}.$$
 (209)

Interestingly, for the GR_{||} case of (164), the right-hand sides of (208) and (209) vanish, i.e., the tensor $M^{\beta}{}_{\alpha}{}^{\nu}{}_{\mu}$ turns out to be transversal to the wave covector q_{μ} in all four indices.

Taking into account these properties, the wave propagation system (177) yields

$$(2\beta_1 - \beta_2 - \beta_3) \left(q^2 g^{\beta \nu} - q^{\beta} q^{\nu} \right) q_{\mu} A_{\nu}{}^{\mu} = 0.$$
(210)

In the generic case, when $2\beta_1 - \beta_2 - \beta_3 \neq 0$, we find the relation

$$q^{2}A_{\beta}{}^{\mu}q_{\mu} = q_{\beta}A_{\nu}{}^{\mu}q_{\mu}q^{\nu}, \qquad (211)$$

which implies

$$q^2 \mathcal{A}_{\beta}{}^{\mu}q_{\mu} = q_{\beta} \mathcal{A}_{\nu}{}^{\mu}q_{\mu}q^{\nu}.$$
(212)

We will use this in subsequent derivations.

Let us now turn to the decomposed equations (191) and (192), Using (205), (206), and $A_v^v = 0$, we can recast (191) into

$$3q^2A = \mathcal{A}_{\nu}{}^{\mu}q_{\mu}q^{\nu}. \tag{213}$$

This is derived for the generic case by assuming $2\beta_1 - 3\beta_2 - \beta_3 \neq 0$, otherwise (191) is trivial.

Finally, we turn to the traceless part (192) of the wave propagation equation. Substituting (204)-(206) into (193), and then making use of (212), we can rewrite (192) in the equivalent form

$$6\beta_{1}q^{2}\left(q^{2}\mathcal{A}^{\beta}_{\alpha}-q^{\beta}q_{\gamma}\mathcal{A}^{\gamma}_{\alpha}\right)$$

$$-3\beta_{3}q^{2}\left(q^{2}\mathcal{A}_{\alpha}^{\beta}-q_{\alpha}q^{\gamma}\mathcal{A}_{\gamma}^{\beta}\right)$$

$$+\left(2\beta_{1}-\beta_{3}\right)\left(q^{2}\delta_{\alpha}^{\beta}-q^{\beta}q_{\alpha}\right)\mathcal{A}_{\nu}^{\mu}q_{\mu}q^{\nu}=0.$$

$$(214)$$

Quite remarkably, the coupling constant β_2 does not contribute.

Decomposing (214) into symmetric and antisymmetric parts, we obtain the two equations

$$(2\beta_1 - \beta_3) \left\{ 3 \left(q^2 \delta^{\mu}_{\alpha} - q^{\mu} q_{\alpha} \right) \left(q^2 \delta^{\nu}_{\beta} - q^{\nu} q_{\beta} \right) - \left(q^2 g_{\alpha\beta} - q_{\alpha} q_{\beta} \right) q^{\mu} q^{\nu} \right\} \mathcal{A}_{(\mu\nu)} = 0,$$
(215)

$$(2\beta_1 + \beta_3) \left(q^2 \delta^{\mu}_{\alpha} - q^{\mu} q_{\alpha} \right) \left(q^2 \delta^{\nu}_{\beta} - q^{\nu} q_{\beta} \right) \mathcal{A}_{[\mu\nu]} = 0.$$
(216)

Two important observations are in order. Firstly, we see that the symmetric $A_{(\mu\nu)}$ and skew-symmetric $A_{[\mu\nu]}$ variables decouple from each other and are governed by two separate dynamical equations. Secondly, the symmetric $A_{(\mu\nu)}$ mode remains the only one when coupling constants satisfy $2\beta_1 = -\beta_3$ and, similarly, the antisymmetric $A_{[\mu\nu]}$ mode remains the only one when the coupling constants satisfy $2\beta_1 = \beta_3$.

In the generic case (when $2\beta_1 \neq -\beta_3$ and $2\beta_1 \neq \beta_3$), both modes are dynamical. In principle, one can proceed by deriving the Fresnel equations for each mode in a way similar to classical electrodynamics. However, one can choose an alternative way and analyze the propagation equations by using the method of spinprojection operators. The corresponding tools were earlier developed in metric gravity [52] and in its teleparallel formulation [40]. Their application is straightforward. Ultimately we conclude that the wave equations describe the massless spin-0, spin-2 and spin-1 modes propagating along the light cones $q^2 = 0$.

6.2 Waves in $GR_{||}$, the teleparallel equivalent of GR

In order to understand more clearly what happens in $GR_{||}$, it is instructive to analyze the wave propagation equation from scratch. Plugging (204) into (175), we find for (174), after some straightforward algebra and lowering the index μ ,

$$(2\beta_{1} - \beta_{2} - \beta_{3}) \times \left\{ 4A(q^{2}g_{\alpha\beta} - q_{\alpha}q_{\beta}) - g_{\alpha\beta}q^{\mu}q^{\nu}A_{\mu\nu} + q_{\alpha}A_{\beta\gamma}q^{\gamma} \right\} - (2\beta_{1} + \beta_{3}) \left\{ q^{2}A_{[\alpha\beta]} - q_{\alpha}q^{\gamma}A_{[\gamma\beta]} - q_{\beta}q^{\gamma}A_{[\alpha\gamma]} \right\} + (2\beta_{1} - \beta_{3}) \left\{ q^{2}A_{(\alpha\beta)} - q_{\alpha}q^{\gamma}A_{(\gamma\beta)} - q_{\beta}q^{\gamma}A_{(\alpha\gamma)} - 4A(q^{2}g_{\alpha\beta} - q_{\alpha}q_{\beta}) + g_{\alpha\beta}q^{\mu}q^{\nu}A_{\mu\nu} \right\} = 0.$$
(217)

Contracting with $g^{\alpha\beta}$, we find

$$(2\beta_1 - 3\beta_2 - \beta_3) \left(4Aq^2 - q^{\mu}q^{\nu}A_{\mu\nu} \right) = 0, \qquad (218)$$

whereas contraction with q^{α} yields

$$(2\beta_1 - \beta_2 - \beta_3) \left(-q_\beta q^\mu q^\nu A_{\mu\nu} + q^2 A_{\beta\gamma} q^\gamma \right) = 0.$$
 (219)

In the generic case, we recover the earlier findings (211) and (210). Making use of this, we see that the first line in (217) vanishes, whereas the remaining equation reduces to the system (215) and (216).

However, $GR_{||}$ represents a very special case when the coupling constants (164) are such that both $2\beta_1 - \beta_2 - \beta_3 = 0$ and $2\beta_1 + \beta_3 = 0$. As a result, the first two lines of (217) disappear, and the propagation equation has only a symmetric traceless part. The bottom line is that in $GR_{||}$ Eq.(217) reduces to

$$q^{2}h_{\alpha\beta} - q_{\alpha}q^{\gamma}h_{\gamma\beta} - q_{\beta}q^{\gamma}h_{\alpha\gamma} = 0, \qquad (220)$$

after we introduce the familiar variable

$$h_{\alpha\beta} := A_{(\alpha\beta)} - 2Ag_{\alpha\beta} = A_{(\alpha\beta)} - \frac{1}{2}g_{\alpha\beta}A_{\gamma}^{\gamma}.$$
 (221)

Accordingly, we indeed recover the result of GR with the massless spin-2 graviton propagating along the light cone. In other words, $GR_{||}$ is completely consistent with Einstein's theory with respect to the propagation of gravitational waves.

7 Conclusions and outlook

Recently, in Ref.[29], we developed a novel framework for a premetric teleparallel theory of gravity (TG). Here we continue this study on TG by paying attention specifically to TG models with a general *local and linear* constitutive law.

In Sec.2, we constructed the Tonti diagram of TG which explicitly displays the generic structure of the theory.

Our main new results are presented in Sec.3 where the irreducible decomposition of the premetric constitutive tensor is established in full detail. This issue was not analysed in the earlier literature. Here we considered two types of such decompositions with respect to the permutation group, namely one related to $S_4 \times S_2$ and the other one with respect to S_6 . The relations between these two decompositions are explicitly derived and the physical meaning of the different irreducible pieces clarified.

After establishing for the constitutive tensor these premetric results, we turned in Sec.4 to a special case: The spacetime continuum is supposed to carry a metric tensor. We constructed the most general metric dependent constitutive tensor that is cubic in the metric tensor. It includes both, parity even and parity odd parts. Thereby extending earlier results, we for the first time obtained the most general family of teleparallel consitutive tensors. In particular, a parity odd Lagrangian was newly found, and its physical interpretation fixed.

In Sec.5, the propagation of gravitational waves are derived for the premetric case in the geometrical optics approximation. Additional propagating modes of spin 1 and 0 are extracted. Our results are generally consistent with the recent findings reported in [27].

Subsequently, in Sec.6, the metric case is addressed. The GR limit is naturally embedded in our formalism.

The results presented here can serve as a basis for the study of expanded gravitational models including axion and skewon effects. In particular, violation of Lorentz invariance can be meaningfully addressed in our premetric set-ups.

Acknowledgments

We are grateful to Enzo Tonti (Trieste) for help in creating our Tonti diagram. Moreover, we thank Bahram Mashhoon (Tehran) for most helpful remarks in the context of nonlocal gravity (NLG). Remarks of María José Guzmán (Buenos Aires), Manuel Hohmann (Tartu), José Maluf (Brasilia), and James Nester (Chung-Li) are gratefully acknowledged. For Y.N.O. this work was partially supported by the Russian Foundation for Basic Research (Grant No. 16-02-00844-A). J.B. is grateful for a Vanier Canada Graduate Scholarship administered by the Natural Sciences and Engineering Research Council of Canada as well as for the Golden Bell Jar Graduate Scholarship in Physics by the University of Alberta.

A Young decomposition

Here we briefly explain how to determine the $GL(4,\mathbb{R})$ decomposition of the order- $\begin{pmatrix} 0\\6 \end{pmatrix}$ constitutive tensor $\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}$. The decomposition rules for tensors of other valence, say order-2 or order-4, can easily be derived. In four dimensions, the decomposition of $\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}$ with the symmetry properties

$$\check{\chi}_{(\alpha\beta)\mu\gamma\delta\nu} = 0, \quad \check{\chi}_{\alpha\beta\mu(\gamma\delta)\nu} = 0, \tag{35}$$

consists of the following seven pieces:



Let us label them from λ_1 to λ_7 , respectively. Each diagram λ_i (*i* = 1...7) has corresponding dimensions according to

$$\dim V^{\lambda} = \prod_{(\alpha,\beta)\in\lambda} \frac{n+\beta-\alpha}{\operatorname{hook}(\alpha,\beta)},$$

$$\dim S_{6}^{\lambda} = f^{\lambda} = \frac{6!}{\prod_{x\in\lambda} \operatorname{hook}(x)}.$$
(223)

We find

$$\dim V^{\lambda_1} = 126, \qquad \dim V^{\lambda_2} = 70, \qquad \dim V^{\lambda_3} = 50, \\ \dim V^{\lambda_4} = 64, \qquad \dim V^{\lambda_5} = 10, \qquad \dim V^{\lambda_6} = 10,$$
(224)
$$\dim V^{\lambda_7} = 6,$$

as well as

$$\dim S_6^{\lambda_1} = 9, \qquad \dim S_6^{\lambda_2} = 10, \qquad \dim S_6^{\lambda_3} = 5, \\ \dim S_6^{\lambda_4} = 16, \qquad \dim S_6^{\lambda_5} = 5, \qquad \dim S_6^{\lambda_6} = 10,$$
(225)
$$\dim S_6^{\lambda_7} = 9.$$

With the relevant Young diagrams now given, we can extract the projection operators in the standard fashion and arrive at the irreducible decomposition of $\chi_{\alpha\beta\gamma\mu\nu\rho}$.

For each λ_i we may extract projection operators in the standard way: we can fill f^{λ_i} copies of the λ_i Young diagram with numbers to create all allowed Young tableaux, and then create the corresponding projection operator onto that subspace by summing over the induced Young symmetrizers:

$$\mathbb{P}_{p}^{\lambda_{i}} := \frac{f^{\lambda_{i}}}{p!} \sum_{j=1}^{f^{\lambda_{i}}} \mathbb{P}\left(Y_{j}^{\lambda_{i}}\right), \qquad (226)$$

$${}^{(I)}\check{\chi}_{\alpha\beta\mu\gamma\delta\nu} := \mathbb{P}_{p}^{\lambda_{I}}\left(\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}\right), \qquad (227)$$

where $\mathbb{P}\left(Y_{j}^{\lambda_{i}}\right)$ is the Young symmetrizer corresponding to the *j*-th Young tableaux of λ_{i} , and ${}^{(I)}\check{\chi}_{\alpha\beta\mu\gamma\delta\nu}$ denotes the *I*-th irreducible piece of the constitutive tensor.

The Young tableaux with non-trivial symmetrizers are as follows:

$$\lambda_1: \begin{array}{cccc} 1 & 3 & 5 & 6 \\ 2 & 4 & & \\ \end{array}, \begin{array}{ccccc} 1 & 3 & 4 & 6 \\ 2 & 5 & & \\ \end{array}, \tag{228}$$

$$\lambda_3: \begin{array}{cccc} 1 & 3 & 5 \\ 2 & 4 & 6 \end{array}, \begin{array}{ccccc} 1 & 3 & 4 \\ 2 & 5 & 6 \end{array},$$
(230)

$$\lambda_{4}: \begin{array}{c} 1 & 3 & 4 \\ 2 & 6 \\ 5 \\ 5 \\ \end{array}, \begin{array}{c} 1 & 3 & 5 \\ 2 & 4 \\ 6 \\ \end{array}, \begin{array}{c} 1 & 3 & 5 \\ 2 & 6 \\ 4 \\ \end{array}, \begin{array}{c} 1 & 3 & 5 \\ 2 & 6 \\ 6 \\ \end{array}, \begin{array}{c} 1 & 3 & 4 \\ 2 & 5 \\ 6 \\ \end{array},$$
(231)

$$\begin{array}{c}
1 & 4 & 6 \\
2 & 5 \\
3 \\
\end{array}, \begin{array}{c}
1 & 3 & 6 \\
2 & 4 \\
5 \\
\end{array}, \begin{array}{c}
1 & 3 & 6 \\
2 & 5 \\
4 \\
\end{array}, \begin{array}{c}
1 & 3 & 6 \\
2 & 5 \\
4 \\
\end{array},$$
(232)

For example, $\mathbb{P}\left(Y_6^{\lambda_4}\right)$ can be computed as follows:

$$\mathbb{P}\left(Y_{6}^{\lambda_{4}}\right) = \mathbb{P}\left(\frac{136}{2}\right) = S_{136} \circ S_{24} \circ A_{125} \circ A_{34} \qquad (236)$$

$$= [1 + (13) + (16) + (36) + (136) + (631)] \circ [1 + (24)]$$

$$\circ [1 - (12) - (15) - (25) + (125) + (521)] \circ [1 - (34)],$$

where S_{136} denotes symmetrization between the 1st, 3rd, 6th index; similarly, A_{34} denotes antisymmetrization between the 3rd and 4th index, and so on. In the last lines we switched to the more convenient cycle notation. Note that in our conventions the Young symmetrizers are not normalized. The exact expressions are lengthy, so we make some comments:

 $\mathbb{1}_{T_{\check{\mathbf{x}}}} = \text{diag}({}^{(1)}\mathbb{P}, {}^{(2)}\mathbb{P}, {}^{(3)}\mathbb{P}, {}^{(4)}\mathbb{P}, {}^{(5)}\mathbb{P}, {}^{(6)}\mathbb{P}, {}^{(7)}\mathbb{P})$



Figure 2: The resolution of the identity on the 576-dimensional vector space $T_{\check{\chi}}$ has a block-diagonal structure in terms of the canonical Young projection operators ${}^{\{I\}}\mathbb{P}: T_{\check{\chi}} \to T_{\check{\chi}}^{I}$ with I = 1, ..., 7. We depict these operators by 1×1 blocks, with the relative width of $f^{\lambda_{I}} \times d_{I}/576$. Provided $f^{\lambda_{I}} > 1$, these operators can be further decomposed into *non-canonical* operators ${}^{\{I\}}\mathbb{P}: T_{\check{\chi}}^{I} \to T_{\check{\chi}}^{I}$ with $j = 1, ..., f^{\lambda_{I}}$. We depict this internal ambiguity by inserting $f^{\lambda_{I}} \times f^{\lambda_{I}}$ smaller blocks.

(i) Using the Young decomposition technique, the 576-dimensional tensor space of $\check{\chi}_{\alpha\beta\gamma\mu\nu\rho}$, call it $T_{\check{\chi}}$, can be decomposed into the direct sum of lower-dimensional vector spaces $T^{I}_{\check{\chi}}$ with I = 1, ..., 7. Denoting the projection operators onto these lower-dimensional subspaces via ${}^{(I)}\mathbb{P}: T_{\check{\chi}} \to T^{I}_{\check{\chi}}$, the identity element on $T_{\check{\chi}}$ can be written as

$$\mathbb{1}_{T_{\check{\chi}}} = \bigoplus_{I=1}^{\prime} {}^{(I)}\mathbb{P} \qquad \Rightarrow \qquad {}^{(I)}\check{\chi} := {}^{(I)}\mathbb{P}(\check{\chi}). \tag{237}$$

(ii) In a suitable matrix representation of $T_{\check{\chi}}$, call it $\rho(T_{\check{\chi}}) = \mathbb{R}^{576}$, the operators ${}^{(I)}\mathbb{P}$ can be thought of as $(f^{\lambda_I}d_I) \times (f^{\lambda_I}d_I)$ matrices, where f^{λ_I} denotes the

degeneracy of $T^{I}_{\check{\gamma}}$.

(iii) On $T^{I}_{\check{\chi}}$, various auxiliary projection operators ${}^{(I,j)}\mathbb{P}$ can be defined such that

$${}^{(I)}\mathbb{P} = \bigoplus_{j=1}^{f^{\lambda_I}} {}^{(I,j)}\mathbb{P}.$$
(238)

We emphasize that the operators ${}^{(I,j)}\mathbb{P}$ are *not canonical*, that is, they are only determined up to arbitrary linear transformations $G: \rho(T^I_{\check{\chi}}) \to \rho(T^I_{\check{\chi}})$. For a graphical representation of statements (i)–(iii), see Fig. 2.

We now use computer algebra [10, 16] to obtain explicit expressions which are a bit lengthy. To compress our results, we define a projector **C** (its index representation $C^{\rho\sigma\omega\lambda}_{\alpha\beta\gamma\delta}$) by

$$\mathbf{C} := \frac{1}{4} \left[1 - (12) - (45) + (12)(45) \right],$$

$$C^{\rho\sigma\omega\lambda}_{\alpha\beta\gamma\delta} := \frac{1}{4} \delta^{\rho\sigma}_{\alpha\beta} \delta^{\omega\lambda}_{\gamma\delta}, \quad \delta^{\rho\sigma}_{\alpha\beta} := \delta^{\rho}_{\alpha} \delta^{\sigma}_{\beta} - \delta^{\sigma}_{\alpha} \delta^{\rho}_{\beta}.$$
(239)

Observe that $\mathbf{C}^2 = \mathbf{C}$, indeed. This projector antisymmetrizes in the two index pairs $\alpha\beta$ and $\gamma\delta$ such that any constitutive tensor satisfying the relation $\check{\chi}_{\alpha\beta\mu\gamma\delta\nu} = \check{\chi}_{[\alpha\beta]\mu[\gamma\delta]\nu}$ has eigenvalue 1 with respect to this operator:

$$C^{\rho\sigma\omega\lambda}_{\alpha\beta\gamma\delta}\check{\chi}_{\rho\sigma\mu\omega\lambda\nu} = \check{\chi}_{\alpha\beta\mu\gamma\delta\nu}.$$
(240)

In general, $C^{\rho\sigma\omega\lambda}_{\alpha\beta\gamma\delta}$ projects a general order-6 tensor without any symmetries, call it $T_{\alpha\beta\mu\gamma\delta\nu}$, onto a tensor that has the symmetries of the constitutive tensor. In that sense we make use of this projector to shorten our exact expressions for the irreducible decomposition of the constitutive tensor by a factor of 4. We obtain:

$$^{(5)}\check{\chi}_{\alpha\beta\mu\gamma\delta\nu} = \frac{1}{36} C^{\rho\sigma\omega\lambda}_{\alpha\beta\gamma\delta} \left(\check{\chi}_{\rho\sigma\omega\lambda\mu\nu} + \check{\chi}_{\rho\sigma\omega\lambda\nu\mu} - 2\check{\chi}_{\rho\sigma\omega\mu\nu\lambda} \right)$$

$$+ 2\check{\chi}_{\rho\sigma\mu\omega\lambda\nu} - \check{\chi}_{\rho\sigma\mu\omega\lambda} - \check{\chi}_{\rho\sigma\nu\omega\mu\lambda} + \check{\chi}_{\rho\omega\lambda\sigma\mu\nu} + \check{\chi}_{\rho\omega\mu\sigma\lambda\nu} + 2\check{\chi}_{\rho\omega\nu\sigma\lambda\mu} + \check{\chi}_{\rho\omega\nu\sigma\lambda\mu} + 3\check{\chi}_{\rho\mu\omega\sigma\lambda\nu} + 2\check{\chi}_{\rho\mu\omega\sigma\lambda\nu} - 3\check{\chi}_{\rho\mu\nu\sigma\nu\lambda} + 2\check{\chi}_{\rho\mu\nu\sigma\lambda\nu} - \check{\chi}_{\rho\mu\nu\sigma\lambda\nu} + 5\check{\chi}_{\rho\mu\nu\sigma\nu\lambda} + 3\check{\chi}_{\rho\nu\nu\sigma\mu\lambda} + 3\check{\chi}_{\rho\nu\mu\sigma\lambda} - \check{\chi}_{\rho\nu\mu\omega\lambda\sigma} - \check{\chi}_{\rho\nu\omega\lambda\mu\sigma} + \check{\chi}_{\rho\nu\sigma\nu\mu} + 3\check{\chi}_{\rho\nu\mu\sigma\nu\lambda} + 2\check{\chi}_{\omega\lambda\rho\mu\nu\sigma} + 2\check{\chi}_{\omega\lambda\rho\sigma\mu\nu} + 3\check{\chi}_{\omega\nu\rho\sigma\mu} + 2\check{\chi}_{\omega\nu\mu\rho\sigma\lambda} + 3\check{\chi}_{\omega\nu\rho\sigma\mu} + 2\check{\chi}_{\omega\nu\rho\sigma\mu\lambda} + 3\check{\chi}_{\omega\nu\rho\sigma\mu\lambda} + 3\check{\chi}_{\omega\nu\rho\sigma\mu\lambda} + 3\check{\chi}_{\omega\nu\rho\sigma\mu\lambda} + 2\check{\chi}_{\omega\nu\rho\sigma\mu\lambda} + 2\check{\chi}_{\omega\nu\rho\mu\nu\nu} + 2\check{\chi}_{\omega\nu\rho\mu\nu} + 2\check{\chi}_{\omega\nu\rho\mu\nu} + 2\check{\chi}_{\omega\nu\rho\nu\rho\sigma\mu\lambda} + 2\check{\chi}_{\omega\nu\rho\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\sigma\mu\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\sigma\mu\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\sigma\mu\lambda} + 2\check{\chi}_{\omega\nu\rho\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\rho\mu\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\rho\mu\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\rho\mu\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\nu\rho\sigma\lambda} + 2\check{\chi}_{\omega\nu\nu\rho\nu\rho\lambda} + 2\check{\chi}_{\omega\nu\nu\rho}$$

It has been verified by computer algebra that the above expressions indeed belong to orthogonal subspaces.

References

- [1] R. Aldrovandi and J. G. Pereira, *Teleparallel Gravity: An Introduction* (Springer, Dordrecht, 2013).
- [2] P. Baekler, A. Favaro, Y. Itin and F. W. Hehl, The Kummer tensor density in electrodynamics and in gravity, Annals Phys. (NY) 349, 297–324 (2014), arXiv:1403.3467.
- [3] P. Baekler and F. W. Hehl, Beyond Einstein-Cartan gravity: Quadratic torsion and curvature invariants with even and odd parity including all boundary terms, Class. Quant. Grav. **28**, 215017 (2011), arXiv:1105.3504.
- [4] P. Baekler, F. W. Hehl and J. M. Nester, Poincaré gauge theory of gravity: Friedman cosmology with even and odd parity modes. Analytic part, Phys. Rev. D 83, 024001 (2011), arXiv:1009.5112.
- [5] A. O. Barut and R. Raczka, *Theory of Group Representations and Applications* (PWN, Warszawa, 1977, and World Scientific, Singapore, 1986).

- [6] M. Blagojević, B. Cvetković and Y. N. Obukhov, Generalized plane waves in Poincaré gauge theory of gravity, Phys. Rev. D 96, 064031 (2017), arXiv:1708.08766.
- [7] M. Blagojević and B. Cvetković, General Poincaré gauge theory: Hamiltonian structure and particle spectrum, Phys. Rev. D 98, 024014 (2018), arXiv:1804.05556.
- [8] M. Blagojević and F. W. Hehl (eds.), *Gauge Theories of Gravitation: A Reader with Commentaries* (Imperial College Press, London, 2013), see arXiv:1210.3775.
- [9] C. G. Böhmer and R. J. Downes, *From continuum mechanics to general relativity*, Int. J. Mod. Phys. D **23** (2014) 1442015, arXiv:1405.4728.
- [10] J. Boos, Irreducible decomposition of the rank-6 constitutive tensor (online computer algebra code), see http://spintwo.net/static/2018.07.17/, last retrieved: 17 Jul 2018.
- [11] Y. M. Cho, Einstein Lagrangian as the Translational Yang-Mills Lagrangian, Phys. Rev. D 14, 2521 (1976).
- [12] D. H. Delphenich, *The use of the teleparallelism connection in continuum mechanics*, Mathematics and Mechanics of Solids **21**, 1260–1275 (2016), arXiv:1305.3477.
- [13] A. Einstein, *The Meaning of Relativity, 5th ed.* (Princeton University Press, Princeton, 1955).
- [14] R. Ferraro and M. J. Guzmán, Hamiltonian formulation of teleparallel gravity, Phys. Rev. D 94, 104045 (2016), arXiv:1609.06766.
- [15] M. Hamermesh, *Group Theory and its Applications to Physical Problems* (Addison-Wesley, Reading, 1962).
- [16] A. C. Hearn, *REDUCE User's Manual, Version 3.5*, RAND Publication CP78 (Rev. 10/93). The RAND Corporation, Santa Monica, CA 90407-2138, USA (1993). Nowadays Reduce is freely available for download; for details see [reduce-algebra.com] and [sourceforge.net].
- [17] F. W. Hehl, Four lectures on Poincaré gauge field theory, in: Proc. of the 6th Course of the School of Cosmology and Gravitation on *Spin, Torsion, Rotation, and Supergravity*, held at Erice, Italy, May 1979, P. G. Bergmann, V. de Sabbata, eds. (Plenum, New York 1980) pp. 5–61; see also the author's homepage: http://www.thp.uni-koeln.de/gravitation/mitarbeiter/hehl.html

- [18] F. W. Hehl, Y. Itin and Y. N. Obukhov, On Kottler's path: origin and evolution of the premetric program in gravity and in electrodynamics, arXiv:1607.06159; a condensed version appeared in the Int. J. Mod. Phys. D 25, 1640016 (2016).
- [19] F. W. Hehl and B. Mashhoon, Nonlocal Gravity Simulates Dark Matter, Phys. Lett. B 673, 279 (2009), arXiv:0812.1059.
- [20] F. W. Hehl and B. Mashhoon, A Formal framework for a nonlocal generalization of Einstein's theory of gravitation, Phys. Rev. D 79, 064028 (2009), arXiv:0902.0560.
- [21] F. W. Hehl, J. Nitsch, and P. von der Heyde, Gravitation and the Poincaré gauge field theory with quadratic Lagrangian, in: *General Relativity and Gravitation: One Hundred Years After the Birth of Albert Einstein*, A. Held, ed. (Plenum Press, New York, 1980) Vol.1, pp.329–355.
- [22] F. W. Hehl and Yu. N. Obukhov, *Foundations of Classical Electrodynamics: Charge, Flux, and Metric* (Birkhäuser, Boston, MA, 2003).
- [23] F.W. Hehl and E. Tonti, *Tonti diagrams for the teleparallelism theory of gravity (TG) and for the Poincaré gauge theory (PG)*, Verhandlungen DPG (German Physical Society) (VI) 52, 3/p.56 (2017).
- [24] P. von der Heyde: Is gravitation mediated by the torsion of spacetime? Z. Naturforsch. **31a**, 1725–1726 (1976).
- [25] F. H. Ho, H. Chen, J. M. Nester and H. J. Yo, General Poincaré Gauge Theory Cosmology, Chin. J. Phys. 53, 110109 (2015), arXiv:1512.01202.
- [26] M. Hohmann, L. Järv, M. Krššák and C. Pfeifer, Teleparallel theories of gravity as analogue of non-linear electrodynamics, Phys. Rev. D 97, 104042 (2018), arXiv:1711.09930.
- [27] M. Hohmann, M. Krššák, C. Pfeifer, and U. Ualikhanova, Propagation of gravitational waves in teleparallel gravity theories, to be published (2018), arXiv:1807.04580.
- [28] Y. Itin, Weak field reduction in teleparallel coframe gravity: Vacuum case, J.Math. Phys. 46, 012501 (2005), arXiv:gr-qc/0409021.
- [29] Y. Itin, On light propagation in premetric electrodynamics: the covariant dispersion relation, J. Phys. A 42, 475402 (2009), arXiv:0903.5520.

- [30] Y. Itin, Skewon modification of the light cone structure, Phys. Rev. D **91**, 085002 (2015), arXiv:1407.6722.
- [31] Y. Itin, F. W. Hehl and Y. N. Obukhov, Premetric equivalent of general relativity: Teleparallelism, Phys. Rev. D 95, 084020 (2017), arXiv:1611.05759.
- [32] G. K. Karananas, The particle spectrum of parity-violating Poincaré gravitational theory, Class. Quant. Grav. 32, 055012 (2015), arXiv:1411.5613; corrigendum: ibid. 32, 089501 (2015).
- [33] V. A. Kostelecký and M. Mewes, Lorentz and diffeomorphism violations in linearized gravity, Phys. Lett. B 779, 136–142 (2018), arXiv:1712.10268.
- [34] R. Kubo (in cooperation with H. Ichimura, T. Usui, N. Hashitsume), *Thermodynamics*, 2nd printing (North Holland, Amsterdam, 1976).
- [35] J. W. Maluf, Dirac spinor fields in the teleparallel gravity: Comment on 'Metric affine approach to teleparallel gravity', Phys. Rev. D 67, 108501 (2003), arXiv:gr-qc/0304005.
- [36] J. W. Maluf, The teleparallel equivalent of general relativity, Annalen Phys. (Berlin) 525, 339 (2013), arXiv:1303.3897.
- [37] B. Mashhoon, Nonlocal Gravity (Oxford Univ. Press, Oxford, UK, 2017).
- [38] H. Meyer, Møller's tetrad theory of gravitation as a special case of a Poincaré gauge theory—a coincidence? Gen. Rel. Grav. 14 (1982) 531–547.
- [39] U. Muench, F. Gronwald and F. W. Hehl, A Small guide to variations in teleparallel gauge theories of gravity and the Kaniel-Itin model, Gen. Rel. Grav. 30, 933 –961 (1998), arXiv:gr-qc/9801036.
- [40] F. Müller-Hoissen and J. Nitsch, Teleparallelism—A viable theory of gravity? Phys. Rev. D 28, 718–728 (1983).
- [41] H. T. Nieh and M. L. Yan, An identity in Riemann-Cartan geometry, J. Math. Phys. 23, 373 (1982).
- [42] H. T. Nieh, A torsional topological invariant, Int. J. Mod. Phys. A 22, 5237-5244 (2007), arXiv:1309.0915.
- [43] Y. N. Obukhov, Poincaré gauge gravity: Selected topics, Int. J. Geom. Meth. Mod. Phys. 3, 95 (2006), arXiv:gr-qc/0601090.
- [44] Y. N. Obukhov, Gravitational waves in Poincaré gauge gravity theory, Phys. Rev. D 95, 084028 (2017), arXiv:1702.05185.

- [45] Y. N. Obukhov, Poincaré gauge gravity: An overview, Lecture in Tartu, Estonia (2017), Int. J. Geom. Methods Mod. Phys. 15, 1840005 (2018), arXiv:1805.07385.
- [46] Y. N. Obukhov, E. W. Mielke, J. Budczies, and F. W. Hehl, On the chiral anomaly in non-Riemannian spacetimes, Found. Phys. 27, 1221–1236 (1997), arXiv:gr-qc/9702011.
- [47] Y. N. Obukhov and J. G. Pereira, Lessons of spin and torsion: Reply to 'Consistent coupling to Dirac fields in teleparallelism,' Phys. Rev. D 69, 128502 (2004), arXiv:gr-qc/0406015.
- [48] C. Pellegrini and J. Plebanski, Tetrad fields and gravitational fields, Mat. Fys. Skr. Dan. Vid. Selsk. 2, no. 4, 1–39 (1963).
- [49] M. Planck, *Treatise on Thermodynamics*, 3rd edition 1926, translated from the 7th German edition (Dover, New York, 1945).
- [50] E. J. Post, Formal Structure of Electromagnetics General Covariance and Electromagnetics (North Holland, Amsterdam, 1962, and Dover, Mineola, NY, 1997).
- [51] I. Prigogine, *Thermodynamics of Irreversible Processes*, 2nd, rev. ed. (Interscience, New York, 1965).
- [52] R. J. Rivers, Lagrangian theory for neutral massive spin-2 fields, Nuovo Cim. 34, 386–403 (1964).
- [53] J. A. Schouten, *Ricci Calculus*, 2nd ed. (Springer, Berlin 1954).
- [54] J. J. Sławianowski, V. Kovalchuk, B. Gołubowska, A. Martens, E. E. Rożko, Space-time as a structured relativistic continuum, Mathematical Methods in the Applied Sciences (Wiley) 41, 5404–5422 (2018).
- [55] R. Stanley, *Hooks and Contents*, Talk at Penn. State, 05 Dec 2008, http://www-math.mit.edu/~rstan/ transparencies/hooks.pdf
- [56] E. Tonti, *The Mathematical Structure of Classical and Relativistic Physics*, A general classification diagram (Birkhäuser-Springer, New York, 2013).
- [57] H.-J. Treder (editor, with contributions by H. H. v. Borzeszkowski, U. Kasper, E. Kreisel, D.-E. Liebscher, H.-J. Treder), *Gravitationstheorie* und Äquivalenzprinzip (in German) (Akademie-Verlag, Berlin, 1971).