

# Cardinality of product sets in torsion-free groups and applications in group algebras

*Alireza Abdollahi and Fatemeh Jafari*

Let  $G$  be a unique product group, i.e., for any two finite subsets  $A, B$  of  $G$  there exists  $x \in G$  which can be uniquely expressed as a product of an element of  $A$  and an element of  $B$ . We prove that, if  $C$  is a finite subset of  $G$  containing the identity element such that  $\langle C \rangle$  is not abelian, then for all subsets  $B$  of  $G$  with  $|B| \geq 7$ ,  $|BC| \geq |B| + |C| + 2$ . Also, we prove that if  $C$  is a finite subset containing the identity element of a torsion-free group  $G$  such that  $|C| = 3$  and  $\langle C \rangle$  is not abelian, then for all subsets  $B$  of  $G$  with  $|B| \geq 7$ ,  $|BC| \geq |B| + 5$ . Moreover, if  $\langle C \rangle$  is not isomorphic to the Klein bottle group, i.e., the group with the presentation  $\langle x, y \mid xyx = y \rangle$ , then for all subsets  $B$  of  $G$  with  $|B| \geq 5$ ,  $|BC| \geq |B| + 5$ . The support of an element  $\alpha = \sum_{x \in G} \alpha_x x$  in a group algebra  $\mathbb{F}[G]$  ( $\mathbb{F}$  is any field), denoted by  $\text{supp}(\alpha)$ , is the set  $\{x \in G \mid \alpha_x \neq 0\}$ . By the latter result, we prove that if  $\alpha\beta = 0$  for some non-zero  $\alpha, \beta \in \mathbb{F}[G]$  such that  $|\text{supp}(\alpha)| = 3$ , then  $|\text{supp}(\beta)| \geq 12$ . Also, we prove that if  $\alpha\beta = 1$  for some  $\alpha, \beta \in \mathbb{F}[G]$  such that  $|\text{supp}(\alpha)| = 3$ , then  $|\text{supp}(\beta)| \geq 10$ . These results improve a part of results in Schweitzer [J. Group Theory, 16 (2013), no. 5, 667-693] and Dykema et al. [Exp. Math., 24 (2015), 326-338] to arbitrary fields, respectively.

## 1. Introduction and Results

Let  $G$  be a torsion-free group written multiplicatively, and let  $|\cdot|$  denote the cardinality of a finite set. One of the basic problems in Additive Number Theory is to obtain lower bounds for the cardinality of  $BC = \{bc \mid b \in B, c \in C\}$  of two finite subsets  $B$  and  $C$  in terms of  $|B|$  and  $|C|$ . There are several related results if  $G$  is abelian (see [9, 10, 11, 13, 26, 30, 32], for instance). But, in the non-abelian case the situation is much less understood. By the main result of [22],

$$(1.1) \quad |BC| \geq |B| + |C| - 1.$$

In [4], it is proved that, for  $|B|, |C| \geq 2$ , the equality in Equation 1.1 holds if there exist  $b \in B^{-1}$  and  $c \in C^{-1}$  such that both  $bB$  and  $Cc$  are progressions with common ratio that is, a set of the form  $\{a, ar, \dots, ar^{n-1}\}$ , for some commuting elements  $a$  and  $r$  of  $G$  and some integer  $n$  ( $r$  will be called the ratio of the progression and  $n$  its length). It is proved in [17] that if  $C$  is a finite subset containing

---

2010 Mathematics Subject Classification. Primary: 20D60; 20C05; Secondary: 11P70; 16S34.

Keywords and Phrases: Torsion-free group; Unique product group; Group algebra; Sumset; Product of sets.

the identity element of a torsion-free group  $G$  such that  $\langle C \rangle$  is not abelian, then for all subsets  $B$  with  $|B| \geq 4$ ,

$$(1.2) \quad |BC| \geq |B| + |C| + 1.$$

Also, in [21], it is proved that if  $B$  is not contained in the left coset of a cyclic subgroup, and  $|C| \geq 32(3+k)^6$ , for  $k \geq 1$ , then

$$|BC| \geq |B| + |C| + k.$$

By using the so-called isoperimetric method, see [14], [15] or [16], and a multigraph  $S(B, C)$  associated with a pair  $(B, C)$  of subsets in a group (see Definition 2.11, below), we prove that the lower bound of  $|BC|$  in 1.2 can be improved for unique product groups. Recall that a group  $G$  has the unique product property, if for every pair of finite non-empty sets  $A, B \subseteq G$ , there exists an element  $x \in AB$  such that  $|\{(a, b) \in A \times B \mid x = ab\}| = 1$ . Note that every unique product group is torsion-free. Obvious examples of unique product groups are right or left linearly orderable groups and so including torsion-free nilpotent groups. In fact the class of unique product groups are very vast as it is closed under taking extensions, subdirect products and being local graded (see [27, pp. 111, Theorem 26.3], [31] and [19]); it was first shown in [29] that not all torsion-free groups have the unique product property by constructing a group with the use of a generalization of small cancellation theory so-called graphical small cancellation theory. Later in [28] a non-unique product group as a subgroup of the direct product of 3 infinite dihedral groups was found. The notion of unique product groups was first considered to settle the existence of non-zero zero divisors in group algebras of such groups (see Higman's PhD Thesis [18]); the unit conjecture of group algebras of torsion-free groups cannot be easily settled in the class of unique product groups and to solve the problem a similar slightly stronger notation of "two unique product group" has been considered (a two unique product group is defined as a unique product group, where in the definition, one replaces "an element" by "two distinct elements" and consider the obvious necessary condition " $|A| > 1$  or  $|B| > 1$ "). However these two notations are the same [35].

Our main results are as follows:

**Theorem 1.1.** *Let  $G$  be a unique product group and  $C$  be a finite subset of  $G$  containing the identity element such that  $\langle C \rangle$  is not abelian. Then for all subsets  $B$  of  $G$  with  $|B| \geq 7$ ,  $|BC| \geq |B| + |C| + 2$ .*

In Lemmas 3.2, 3.4 and 3.7 (see below), we determine the structure of  $B$  and  $C$  with  $|C| = 3$  and  $|B| \in \{4, 5, 6\}$  satisfying the equality of 1.2. By Lemmas 3.4 and 3.7 (see below), if  $|B| \in \{5, 6\}$ , then  $G$  is isomorphic to the Klein bottle group i.e., the group with the presentation  $\langle x, y \mid xyx = y \rangle$ .

**Theorem 1.2.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 3$  and  $\langle C \rangle$  is not abelian. Then for all subsets  $B$  of  $G$  with  $|B| \geq 7$ ,  $|BC| \geq |B| + 5$ . In particular, if  $\langle C \rangle$  is not isomorphic to the Klein bottle group, then for all subsets with  $|B| \geq 5$ ,  $|BC| \geq |B| + 5$ .*

Table 1 summarizes some results for the lower bound of  $|BC|$ , where  $B$  and  $C$  are two finite subsets of a torsion-free group.

$G$	$\langle C \rangle$	$ C $	$ B $	Lower bound for $ BC $	Reference
Torsion-free	Arbitrary	$\geq 2$	$\geq 2$	$ B  +  C  - 1$	[22]
	Non-abelian	$\geq 3$	$\geq 4$	$ B  +  C  + 1$	[17]
		$= 3$	$\geq 7$	$ B  + 5$	Theorem 1.2
	Non-abelian, $\not\cong$ Klein bottle group	$= 3$	$\geq 5$	$ B  + 5$	Theorem 1.2
Unique product	Non-abelian	$\geq 3$	$\geq 7$	$ B  +  C  + 2$	Theorem 1.1

TABLE 1. Results for the lower bound of  $|BC|$ , where  $B$  and  $C$  are two finite subsets of a group  $G$  and  $1 \in C$ . The gray rows show the result in this paper.

We note that the support of an element  $\alpha = \sum_{x \in G} \alpha_x x$  in a group algebra, denoted by  $\text{supp}(\alpha)$ , is the set  $\{x \in G \mid \alpha_x \neq 0\}$ . The following are corollaries of Theorem 1.2.

**Corollary 1.3.** *Let  $\alpha$  and  $\beta$  be non-zero elements of the group algebra of any torsion-free group over an arbitrary field. If  $|\text{supp}(\alpha)| = 3$  and  $\alpha\beta = 0$ , then  $|\text{supp}(\beta)| \geq 12$ .*

**Corollary 1.4.** *Let  $\delta$  and  $\gamma$  be elements of the group algebra of any torsion-free group over an arbitrary field. If  $|\text{supp}(\delta)| = 3$  and  $\delta\gamma = 1$ , then  $|\text{supp}(\gamma)| \geq 11$ .*

Corollaries 1.4 and 1.3 improve the lower bounds of 10 and 9 in [2, Theorem 1.7] and [2, Theorem 1.4], respectively.

## 2. Preliminaries

Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element. For  $X \subseteq G$ ,  $\partial_C(X) := XC \setminus X$ . For every positive integer  $k$ , the  $k$ -isoperimetric number of  $C$  is

$$\kappa_k(C) := \min\{|\partial_C(X)| \mid X \subseteq G, k \leq |X| < \infty\}.$$

A finite subset  $X$  of  $G$  is a  $k$ -critical set of  $C$  if  $|X| \geq k$  and  $|\partial_C(X)| = \kappa_k(C)$ . A  $k$ -atom of  $C$  is a  $k$ -critical set of  $C$  with minimal cardinality. The cardinality of a  $k$ -atom of  $C$  will be denoted by  $\alpha_k(C)$ . The minimality of the cardinality of atoms directly yields the following lemma that we repeatedly use in the paper.

**Lemma 2.1.** ([17, Lemma 4]) *Let  $1 \in C$  be a finite generating subset of a torsion-free group  $G$ . Let  $A$  be a  $k$ -atom of  $C$  such that  $|A| > k$ . Then  $|zC^{-1} \cap A| \geq 2$  for all  $z \in AC$ . Moreover,  $|A|(|C| - 2) \geq 2\kappa_k(C)$ .*

**Definition 2.2.** *Let  $B$  and  $C$  be finite non-empty subsets of  $G$ . For each element  $x \in BC$ , denote by  $R_{BC}(x)$  the set  $\{(b, c) \in B \times C \mid x = bc\}$  and let  $r_{BC}(x) := |R_{BC}(x)|$ . Clearly,  $r_{BC}(x) \leq \min\{|B|, |C|\}$  for all  $x \in BC$ .*

In [16], Y.O. Hamidoune proposed the following conjecture:

**Conjecture 2.3.** *Let  $G$  be a torsion-free group,  $C$  a finite subset of  $G$  containing the identity element and  $n$  a positive integer. Then  $|\alpha_n(C)| = n$ .*

In view of [21, p. 5, lines 27-31], Conjecture 2.3 holds valid for unique product groups. Conjecture 2.3 is also valid for  $n = 1$  (see the proof of [14, Proposition 2.8] or [17, Corollary 2]). Conjecture 2.3 is still open and attempts to confirm it even for  $n = 2$  were unsuccessful see e.g. [17, 21].

It was known that some problems in “product of finite subsets of groups” are related to the problem of “absence of zero divisors in the group ring of a torsion-free group over an integral domain” (see [4, p. 463, lines 5-8]). Here we show that Conjecture 2.3 is related to zero divisor and unit conjectures on group algebras of torsion-free groups. Let us first state the latter conjectures.

In 1940, Irving Kaplansky [20] stated his well known conjecture so-called zero divisor conjecture, as follows:

**Conjecture 2.4.** *Let  $\mathbb{F}$  be any field and  $G$  any torsion-free group. Then  $\mathbb{F}[G]$  contains no zero divisor, where a zero divisor is a non-zero element  $\alpha \in \mathbb{F}[G]$  such that  $\alpha\beta = 0$  for some non-zero element  $\beta \in \mathbb{F}[G]$ .*

Another famous conjecture of Kaplansky on group algebras so-called unit conjecture, is the following [20]:

**Conjecture 2.5.** *Let  $\mathbb{F}$  be any field and  $G$  any torsion-free group. Then  $\mathbb{F}[G]$  contains no non-trivial units, where trivial units are non-zero scalar multiples of group elements.*

Conjecture 2.5 is actually stronger than Conjecture 2.4 so that the affirmative solution to Conjecture 2.5 implies the positive one for Conjecture 2.4 [27, Lemma 13.1.2]. Partial results have been obtained on Conjectures 2.5 and 2.4 [5, 7, 8, 23, 24, 25, 27, 34].

Now we show connections of Conjectures 2.5 and 2.4 to Conjecture 2.3 in the following two propositions.

**Proposition 2.6.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|\alpha_k(C)| > k$ . If  $A$  is a  $k$ -atom of  $C$  and  $|AC| = \frac{|A||C|}{2}$ , then there exists a counterexample to Conjecture 2.4.*

*Proof.* Since  $|\alpha_k(C)| > k$ ,  $r_{AC}(x) \geq 2$  for all  $x \in AC$  (see Lemma 2.1). Now it follows from  $|AC| = \frac{|A||C|}{2}$  that  $r_{AC}(x) = 2$  for all  $x \in AC$ . We now define  $\alpha$  and  $\beta$ , respectively, as the elements  $\alpha := \sum_{a \in A} a$  and  $\beta := \sum_{c \in C} c$  in the group algebra  $\mathbb{F}_2[G]$  of  $G$  over the field  $\mathbb{F}_2$  with 2 elements. Therefore  $\alpha\beta = \sum_{x \in AC} r_{AC}(x)x = 0$ . This completes the proof.  $\square$

**Proposition 2.7.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|\alpha_k(C)| > k$ . If  $A$  is a  $k$ -atom of  $C$  and  $|AC| = \frac{|A||C|-1}{2}$ , then there exists a counterexample to Conjecture 2.5.*

*Proof.* Since  $|\alpha_k(C)| > k$ ,  $r_{AC}(x) \geq 2$  for all  $x \in AC$  (see Lemma 2.1). Now it follows from  $|AC| = \frac{|A||C|-1}{2}$  that there exists  $x \in AC$  such that  $r_{AC}(x) = 3$  and  $r_{AC}(x') = 2$ , for all  $x' \in AC \setminus \{x\}$ . Let  $(a_0, c_0) \in R_{AC}(x)$  and define  $\alpha$  and  $\beta$ , respectively, as the elements  $\alpha = \sum_{a \in A} a$  and  $\beta = \sum_{c \in C} c$  in the group algebra  $\mathbb{F}_2[G]$ . Then

$$a_0^{-1}\alpha\beta c_0^{-1} = a_0^{-1} \left( \sum_{y \in AC} r_{AC}(y)y \right) c_0^{-1} = a_0^{-1}(3x)c_0^{-1} = a_0^{-1}(a_0c_0)c_0^{-1} = 1.$$

This completes the proof.  $\square$

In what follows, we present two lemmas which we use to prove Theorem 2.10.

**Lemma 2.8.** ([17, Lemma 5]) *Let  $C$  be a finite generating subset of a torsion-free group  $G$  such that  $|C| \geq 3$  and  $1 \in C$ . Let  $A$  be a 2-atom of  $C$ . Then  $|A| \leq |C| - 1$ .*

**Lemma 2.9.** ([21, Lemma 5]) *For a torsion-free group  $G$  and  $n \geq 2$ , if  $A$  is an  $n$ -atom for  $C \subseteq G$  and  $g \in G \setminus \{1\}$ , then*

$$|A \cap Ag| \leq \frac{n-2}{n-1}|A| + \frac{1}{n-1} \leq \frac{n}{n-1}|A|.$$

Using Propositions 2.6 and 2.7, we give a partial answer to Conjecture 2.3.

**Theorem 2.10.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element. If  $|C| \leq 5$ , then  $\alpha_2(C) = 2$ . Moreover, if  $|C| \leq 7$ , then  $\alpha_2(C) \neq 3$ .*

*Proof.* Observe that if  $|C| = 1$ , then each subset with two elements of  $G$  is a 2-atom of  $C$  and also if  $|C| = 2$ , then according to 1.1,  $C$  is a 2-atom of itself. Now, suppose that  $|C| \geq 3$  and  $\alpha_2(C) > 2$ . Let  $A$  be a 2-atom of  $C$  containing the identity element. It follows from Lemma 2.8 that  $|A| \leq |C| - 1$ . Hence,  $|C| \geq 4$ . We first show that  $\kappa_2(C) \geq |C|$ . In view of 1.1,  $\kappa_2(C) \geq |C| - 1$ . If  $\kappa_2(C) = |C| - 1$ , then by the main result of [4], there exist  $r \in G \setminus \{1\}$  and  $a \in G$  such that  $A = \{a, ar, \dots, ar^l\}$  for some non-negative integer  $l$  and therefore  $|A \cap Ar| \geq 2$ , contradicting Lemma 2.9. So,  $\kappa_2(C) \geq |C|$ . Now since  $|A| > 2$ , Lemma 2.1 implies that  $r_{AC}(x) \geq 2$  for all  $x \in AC$ . It follows that  $|AC| \leq \frac{|A||C|}{2}$ . Thus,

$$(2.1) \quad |C| + |A| \leq |AC| \leq \frac{|A||C|}{2}.$$

If  $|C| = 4$ , then  $|A| = 3$  which contradicts 2.1. So far we have proved that if  $|C| \leq 4$  then  $\alpha_2(C) = 2$ ; we will use the latter in the rest of the proof.

Now, suppose that  $|C| = 5$ . Then  $|A| \in \{3, 4\}$ . Now 2.1 implies that  $|A| \neq 3$  and so  $|A| = 4$ . So, 2.1 implies  $|AC| \in \{9, 10\}$ . Suppose first that  $|AC| = 10$ . Thus, according to the proof of Proposition 2.6, there are  $\alpha, \beta \in \mathbb{F}_2[G]$  such that  $\text{supp}(\alpha) = A$ ,  $\text{supp}(\beta) = C$  and  $\alpha\beta = 0$ , contradicting [33, Theorem 1.3]. Now, suppose that  $|AC| = 9$ . Therefore,  $|C^{-1}A^{-1}| = 9$  leads to  $\kappa_2(A^{-1}) \leq |A^{-1}|$ . Since  $|A^{-1}| = 4$ , by the above,  $\alpha_2(A^{-1}) = 2$ . Let  $A' = \{1, r\}$  be a 2-atom of  $A^{-1}$ . From  $|A'A^{-1}| \leq 6$  and 1.1 it follows that  $|A \cap Ar| \geq 2$  contradicting Lemma 2.9. This completes the proof of the first part of the theorem. If  $|C| = 6$  and  $|A| = 3$ , then by 2.1,  $|AC| = 9$ . Thus, according to the proof of Proposition 2.6, there are  $\alpha, \beta \in \mathbb{F}_2[G]$  such that  $\text{supp}(\alpha) = A$ ,  $\text{supp}(\beta) = C$  and  $\alpha\beta = 0$ , contradicting [33, Theorem 1.3]. Now, suppose that  $|C| = 7$  and  $|A| = 3$ . Then, 2.1 implies  $|AC| = 10$ . Hence, according to the proof of Proposition 2.7, there are  $\alpha, \beta \in \mathbb{F}_2[G]$  such that  $|\text{supp}(\alpha)| = |A|$ ,  $|\text{supp}(\beta)| = |C|$  and  $\alpha\beta = 1$ , contradicting [6, Proposition 4.12]. This completes the proof.  $\square$

**2.1. Product set graph.** We follow the definitions and notations on graphs as in [2]. Note that by a graph we mean a triple  $(\mathcal{V}, \mathcal{E}, \psi)$ , where  $\psi$  is a function which appears if the edge set  $\mathcal{E}$  is non-empty (for details see [2, p. 2, paragraph 3]). In [2, Definitions 2.1 and 2.7], two multigraphs  $Z(\alpha, \beta)$  and  $U(a, b)$  are associated with a pair  $(\alpha, \beta)$  of zero divisors (i.e.  $\alpha\beta = 0$ ) and a pair  $(a, b)$  of unit elements (i.e.  $ab = 1$ ) in a group algebra, respectively. The definitions of the latter graphs are independent of any field and the conditions on  $\alpha$  and  $\beta$  or  $a$  and  $b$ ; in fact, the definitions only depend on the product of two pair of subsets  $(\text{supp}(\alpha), \text{supp}(\beta))$  and  $(\text{supp}(a), \text{supp}(b))$ . Here, we extend these definitions to the product of any pair of non-empty finite subsets of a group as follows:

**Definition 2.11.** *Let  $G$  be a group and  $B, C$  be two finite non-empty subsets of  $G$ . We assign a graph  $P(B, C)$  to  $(B, C)$  called the product set graph of  $(B, C)$  as follows: the vertex set is  $B$ , the edge set is*

$$\{ \{(b, b', c, c'), (b', b, c', c)\} \mid c, c' \in C, b, b' \in B, b \neq b', bc = b'c' \},$$

and if  $\mathcal{E}_{P(B, C)} \neq \emptyset$ , the function  $\psi_{P(B, C)} : \mathcal{E}_{P(B, C)} \rightarrow \mathcal{V}_{P(B, C)}^2$  is defined by

$$\psi_{P(B, C)}(\{(b, b', c, c'), (b', b, c', c)\}) = \{b, b'\},$$

for all  $\{(b, b', c, c'), (b', b, c', c)\} \in \mathcal{E}_{P(B, C)}$ .

**Remark 2.12.** *Notice that in general the graph  $P(B, C)$  may not be a simple graph. In fact  $P(B, C)$  is an undirected graph with no loops but it may happen that it has a multi-edge. For example two multi-graphs  $Z(\alpha, \beta)$  and  $U(a, b)$  associated with a pair  $(\alpha, \beta)$  of zero divisors or a pair  $(a, b)$  of unit elements in a group algebra [2, 1] are isomorphic to the graphs  $P(\text{supp}(\beta)^{-1}, \text{supp}(\alpha)^{-1})$  and  $P(\text{supp}(b)^{-1}, \text{supp}(a)^{-1})$ , respectively.*

**Lemma 2.13.** *Let  $B, C$  be two finite non-empty subsets of a group  $G$ . Then  $P(B, C) \cong P(xB, Cy)$  for all  $x, y \in G$ .*

*Proof.* The proof is similar to [2, Lemma 2.4]. □

**Theorem 2.14.** *Let  $G$  be a torsion-free group and  $B, C$  be two finite non-empty subsets of  $G$  such that  $1 \in C$  and  $|C| = 3$ . If  $\langle C \rangle$  is not cyclic, then  $P(B, C)$  is the induced subgraph of the Cayley graph of  $G$  with respect to  $S := \{hh'^{-1} \mid h \neq h', h, h' \in C\}$  on the set  $B$ .*

*Proof.* Suppose that there are two edges between distinct vertices  $g$  and  $g'$  of  $P(B, C)$ . Hence, there exist  $h_1, h_2, h'_1, h'_2 \in C$  such that  $\{(g, g', h_1, h'_1), (g', g, h'_1, h_1)\}$  and  $\{(g, g', h_2, h'_2), (g', g, h'_2, h_2)\}$  are distinct elements of  $\mathcal{E}_{P(B, C)}$ . Clearly,  $h_1 \neq h'_1, h_2 \neq h'_2, h_1 \neq h_2, h'_1 \neq h'_2$  and  $h_1 h'_1{}^{-1} = g^{-1} g' = h_2 h'_2{}^{-1}$  which implies  $h_2^{-1} h_1 = h'_2{}^{-1} h'_1$ . Then,  $|\{h^{-1} h' \mid h \neq h', h, h' \in C\}| < 6$ . Now, since  $1 \in C$ , it follows from [2, Lemma 3.5] that  $\langle C \rangle$  is an infinite cyclic group that is a contradiction. □

In the following, we need some definitions that are very similar to the definitions in [2, p. 11]. For the reader's convenience, all necessary definitions are given below.

**Definition 2.15.** Let  $G$  be a torsion-free group and  $B$  and  $C$  be two non-empty finite subsets of  $G$  such that  $1 \in C$ ,  $|C| = 3$  and  $\langle C \rangle$  is not cyclic. Suppose that  $\mathcal{C}$  is a cycle of length  $n$  in  $P(B, C)$  with the vertex set  $\{g_1, g_2, \dots, g_n\} \subseteq B$  such that  $g_i \sim g_{i+1}$  for all  $i \in \{1, \dots, n-1\}$  and  $g_1 \sim g_n$ . By an arrangement  $l$  of the vertex set  $\mathcal{C}$ , we mean a sequence of all vertices as  $x_1, \dots, x_k$  such that  $x_i \sim x_{i+1}$  for all  $i \in \{1, \dots, n-1\}$  and  $x_1 \sim x_n$ . There exist unique distinct elements  $h_i, h'_i \in C$ ,  $i \in \{1, \dots, n\}$ , satisfying the following relations:

$$(2.2) \quad R : \begin{cases} g_1 h_1 = g_2 h'_1 \\ g_2 h_2 = g_3 h'_2 \\ \vdots \\ g_n h_n = g_1 h'_n. \end{cases}$$

We assign the  $2n$ -tuple  $T_C^l = [h_1, h'_1, h_2, h'_2, \dots, h_n, h'_n]$  to  $\mathcal{C}$  corresponding to the above arrangement  $l$  of the vertex set of  $\mathcal{C}$ . We denote by  $R(T_C^l)$  the above set  $R$  of relations. It can be derived from the relations 2.2 that  $r(T_C^l) := (h_1 h'_1)^{-1} (h_2 h'_2)^{-1} \dots (h_n h'_n)^{-1}$  is equal to 1. It follows from Lemma 2.14 that if  $T_C^{l'}$  is the  $2n$ -tuple of  $\mathcal{C}$  corresponding to another arrangement  $l'$  of the vertex set of  $\mathcal{C}$ , then  $T_C^{l'}$  is one of the following  $2n$ -tuples:

$$\begin{aligned} & [h_1, h'_1, h_2, h'_2, \dots, h_n, h'_n], \\ & [h_2, h'_2, h_3, h'_3, \dots, h_n, h'_n, h_1, h'_1], \\ & \vdots \\ & [h_n, h'_n, h_1, h'_1, \dots, h_{n-2}, h'_{n-2}, h_{n-1}, h'_{n-1}], \\ & [h'_n, h_n, h'_{n-1}, h_{n-1}, \dots, h'_2, h_2, h'_1, h_1], \\ & \vdots \\ & [h'_1, h_1, h'_n, h_n, \dots, h'_3, h_3, h'_2, h_2]. \end{aligned}$$

The set of all such  $2n$ -tuples will be denoted by  $\mathcal{T}(\mathcal{C})$ .

**Definition 2.16.** Let  $G$  be a torsion-free group and  $B$  and  $C$  be two finite non-empty subsets of  $G$  such that  $1 \in C$ ,  $|C| = 3$  and  $\langle C \rangle$  is not cyclic. Let  $\mathcal{C}$  be a cycle of length  $n$  in  $P(B, C)$ . Since  $r(T_1) = 1$  if and only if  $r(T_2) = 1$ , for all  $T_1, T_2 \in \mathcal{T}(\mathcal{C})$ , a member of  $\{r(T) \mid T \in \mathcal{T}(\mathcal{C})\}$  is given as a representative and denoted by  $r(\mathcal{C})$ . Also,  $r(\mathcal{C}) = 1$  is called the relation of  $\mathcal{C}$ .

**Definition 2.17.** Let  $G$  be a torsion-free group and  $B$  and  $C$  be two finite non-empty subsets of  $G$  such that  $1 \in C$ ,  $|C| = 3$  and  $\langle C \rangle$  is not cyclic. Let  $\mathcal{C}$  and  $\mathcal{C}'$  be two cycles of length  $n$  in  $P(B, C)$ . We say that these two cycles are non-equivalent, if  $\mathcal{T}(\mathcal{C}) \cap \mathcal{T}(\mathcal{C}') = \emptyset$ .

**Remark 2.18.** Suppose that  $\mathcal{C}$  and  $\mathcal{C}'$  are two cycles of length  $n$  in  $P(B, C)$ . Then  $\mathcal{C}$  and  $\mathcal{C}'$  are equivalent if  $\mathcal{T}(\mathcal{C}) = \mathcal{T}(\mathcal{C}')$ .

**Remark 2.19.** Let  $G$  be a torsion-free group and  $B$  and  $C$  be two finite non-empty subsets of  $G$  such that  $1 \in C$ ,  $|C| = 3$  and  $\langle C \rangle$  is not cyclic. Suppose that  $\mathcal{C}$  is a cycle of length 3 (a triangle) in  $P(B, C)$  with the vertex set  $\mathcal{V}_{\mathcal{C}} = \{g_1, g_2, g_3\} \subseteq B$ . Suppose further that  $T \in \mathcal{T}(\mathcal{C})$  and  $T = [h_1, h'_1, h_2, h'_2, h_3, h'_3]$ , where  $h_i, h'_i \in C$  for all  $i \in \{1, 2, 3\}$ . Then  $T$  satisfies exactly one of the following conditions:

- (i)  $h_1 \neq h'_1 = h_2 \neq h'_2 = h_3 \neq h'_3 = h_1$ .
- (ii)  $h_1 \neq h'_1 \neq h_2 \neq h'_2 \neq h_3 \neq h'_3 \neq h_1$ .

More precisely, we shall speak of a triangle  $\mathcal{C}$  in  $P(B, C)$  of type (j) if  $T \in \mathcal{T}(\mathcal{C})$  satisfies the condition (j) in the above list (j being i or ii).

**Lemma 2.20.** *Let  $G$  be a torsion-free group and  $B$  and  $C = \{1, x, y\}$  be two finite non-empty subsets of  $G$  such that  $\langle C \rangle$  is not abelian. If  $P(B, C)$  contains a triangle of type (ii), then  $x, y$  satisfy exactly one of the relations as follows: (1)  $xyx = x$ ; (2)  $xyx = y$ ; (3)  $x^2 = y^2$ . Moreover, every two triangles of  $P(B, C)$  are equivalent.*

*Proof.* Suppose that  $\mathcal{C}$  is a triangle of type (ii) in  $P(B, C)$ . It can be seen that there are 13 non-equivalent cases for  $\mathcal{C}$  and  $r(\mathcal{C})$  corresponding to such cases are the elements of the set

$$A = \{x^{-3}, (y^{-1}x)^2y^{-1}, x^{-2}y^{-1}, x^{-1}y^{-2}, x^{-1}y^{-1}xy^{-1}, x^{-1}yxy^{-1}, x^{-1}yx^{-1}y^{-1}, \\ (x^{-1}y)^2x^{-1}, y^{-3}, y^{-2}xy^{-1}, x^{-1}y^2x^{-1}, (x^{-1}y)^3, x^{-2}yx^{-1}\}.$$

If  $r(\mathcal{C})$  is  $i$ -th element of the set  $A$ , where  $i \in \{2, 3, 4, 6, 8, 10, 13\}$ , then  $r(\mathcal{C}) = 1$  implies that  $\langle C \rangle$  is abelian that is a contradiction. If  $r(\mathcal{C})$  is 1-th or 9-th element of the set  $A$ , then  $G$  has a non-trivial torsion element, a contradiction. Hence,  $r(\mathcal{C})$  is one of the elements 5-th, 7-th or 11-th of the set  $A$ . On the other hand, in a torsion-free group, at most one of these three relations can hold. Indeed,  $xyx = x$  and  $xyx = y$  imply  $x = xyx^2y$  and therefore  $x^2 = y^{-2}$ . So,  $y^{-2} = x^2 = yxy^2xy = y^2$ , a contradiction. If  $xyx = x$  and  $x^2 = y^2$ , then  $y^2 = y^6$ , a contradiction. Hence, two triangles of type (ii) are equivalent. Observe that two triangles of type (i) are equivalent. This completes the proof.  $\square$

**Lemma 2.21.** *Let  $G$  be a torsion-free group and  $B$  and  $C$  be two non-empty finite subsets of  $G$  such that  $1 \in C$ ,  $|C| = 3$  and  $\langle C \rangle$  is not abelian. Suppose that  $P(B, C)$  contains a subgraph isomorphic to the graph in Figure 1. Then there exist  $h \in C$  and  $a' \in G$  such that  $Ch^{-1} = \{1, a, b\}$ ,  $a^2 = b^2$  and one of the sets  $a'\{1, a^{-1}, b^{-1}, b^{-1}a\}$  and  $a'\{1, b^{-1}, b, ba^{-1}\}$  is a subset of  $B$ .*

*Proof.* Let  $C = \{1, x, y\}$ . Suppose that  $\mathcal{C}$  and  $\mathcal{C}'$  are two triangles with the vertex sets  $\{g_1, g_2, g_3\}$  and  $\{g_1, g_2, g_4\}$  in  $P(B, C)$  as Figure 1. Suppose further that  $T = [h_1, h'_1, h_2, h'_2, h_3, h'_3]$  is the 6-tuple to  $\mathcal{C}$  corresponding to the arrangement  $g_1, g_2, g_3$  and also  $T' = [h_1, h'_1, t_2, t'_2, t_3, t'_3]$  is the 6-tuple to  $\mathcal{C}'$  corresponding to the arrangement  $g_1, g_2, g_4$ , where the first two components of  $T$  and  $T'$  are related to the common edge between these triangles. It is clear that either  $h_2 \neq t_2$  or  $h_2 = t_2$  and  $h'_2 \neq t'_2$  and also either  $h'_3 \neq t'_3$  or  $h'_3 = t'_3$  and  $h_3 \neq t_3$  since otherwise  $g_3 = g_4$ . Hence,  $T \neq T'$  and therefore it is impossible that the triangles  $\mathcal{C}$  and  $\mathcal{C}'$  are both of type (i). So, without loss of generality, we may assume that  $\mathcal{C}$  is a triangle of type (ii). Then Lemma 2.20 implies that  $r(\mathcal{C}) \in \{x^{-1}y^{-1}xy^{-1}, x^{-1}yx^{-1}y^{-1}, x^{-1}y^2x^{-1}\}$ . If  $r(\mathcal{C}) = x^{-1}y^{-1}xy^{-1}$ , then

$$\mathcal{T}(\mathcal{C}) = \{[1, x, 1, y, x, y], [1, y, x, y, 1, x], [x, y, 1, x, 1, y], [y, x, y, 1, x, 1], [y, 1, x, 1, y, x], [x, 1, y, x, y, 1]\},$$

and if  $r(\mathcal{C}) = x^{-1}y^2x^{-1}$ , then

$$\mathcal{T}(\mathcal{C}) = \{[1, x, y, 1, y, x], [y, 1, y, x, 1, x], [y, x, 1, x, y, 1], [x, y, 1, y, x, 1], [1, y, x, 1, x, y], [x, 1, x, y, 1, y]\}.$$






FIGURE 1. Two triangles with one edge in common.

Now, if  $\mathcal{C}'$  is a triangle of type (ii), then by Lemma 2.20,  $\mathcal{C}$  and  $\mathcal{C}'$  are equivalent and therefore  $\mathcal{T}(\mathcal{C}) = \mathcal{T}(\mathcal{C}')$ . But in this case, by every choice of  $r(\mathcal{C})$  as above, there are no two elements of  $\mathcal{T}(\mathcal{C})$  satisfying the mentioned conditions for  $T$  and  $T'$ . Thus,  $\mathcal{C}'$  must be a triangle of type (i). Hence,  $g_1h_1 = g_2h'_1 = g_4t_3$ , where  $\{h_1, h'_1, t_3\} = \mathcal{C}$ . Note that by Lemma 2.13, we may assume that  $g_1 = 1$ . Suppose first that  $r(\mathcal{C}) = x^{-1}y^{-1}xy^{-1}$ . Then if we let  $a = yx^{-1}$  and  $b = x^{-1}$ , then  $Cx^{-1} = \{1, a, b\}$ ,  $a^2 = b^2$  and if  $T$  is the 1-th and 6-th elements of  $\mathcal{T}(\mathcal{C})$ , then  $h = y = ab^{-1} = a^{-1}b$  and  $\{g_1, g_2, g_3, g_4\}$  is equal to  $b\{1, a^{-1}, b^{-1}, b^{-1}a\}$  and  $\{1, a^{-1}, b^{-1}, b^{-1}a\}$ , respectively; if  $T$  is the 2-th and 5-th elements of  $\mathcal{T}(\mathcal{C})$ , then  $h = x = b^{-1}$  and  $\{g_1, g_2, g_3, g_4\}$  is equal to  $\{1, b^{-1}, b, b^{-1}a\}$  and  $\{1, a^{-1}, a, a^{-1}b\}$ ; if  $T$  is the 3-th and 4-th elements of  $\mathcal{T}(\mathcal{C})$ , then  $h = 1$  and  $\{g_1, g_2, g_3, g_4\}$  is equal to  $\{1, a^{-1}, b^{-1}, a^{-1}b\}$  and  $a\{1, a^{-1}, b^{-1}, a^{-1}b\}$ , respectively. Now, suppose that  $r(\mathcal{C}) = x^{-1}y^2x^{-1}$ . Then if  $T$  is the 1-th and 6-th elements of  $\mathcal{T}(\mathcal{C})$ , then  $h = y$  and  $\{g_1, g_2, g_3, g_4\}$  is equal to  $\{1, x^{-1}, y^{-1}, x^{-1}y\}$  and  $x\{1, x^{-1}, y^{-1}, x^{-1}y\}$ , respectively; if  $T$  is the 2-th and 5-th elements of  $\mathcal{T}(\mathcal{C})$ , then  $h = x$  and  $\{g_1, g_2, g_3, g_4\}$  is equal to  $y\{1, x^{-1}, y^{-1}, y^{-1}x\}$  and  $\{1, x^{-1}, y^{-1}, y^{-1}x\}$ ; if  $T$  is the 3-th and 4-th elements of  $\mathcal{T}(\mathcal{C})$ , then  $h = 1$  and  $\{g_1, g_2, g_3, g_4\}$  is equal to  $\{1, y^{-1}, y, y^{-1}x\}$  and  $\{1, x^{-1}, x, x^{-1}y\}$ , respectively. Note that if  $r(\mathcal{C}) = x^{-1}yx^{-1}y^{-1}$ , then by interchanging  $x$  and  $y$  in the case  $r(\mathcal{C}) = x^{-1}y^{-1}xy^{-1}$  and with the same discussion, the statement is true. This completes the proof.  $\square$

**Lemma 2.22.** *Let  $G$  be a torsion-free group and  $B$  and  $C$  be two non-empty finite subsets of  $G$  such that  $1 \in C$ ,  $|C| = 3$  and  $\langle C \rangle$  is not abelian. Then  $P(B, C)$  contains no subgraph isomorphic to the complete graph with 4 vertices.*

*Proof.* In view of the proof of Lemma 2.21, if  $P(B, C)$  contains two triangles which have an edge in common, then one of them should be of type (i) and the other of type (ii). But, if  $P(B, C)$  contains a subgraph isomorphic to the complete graph with 4 vertices, then clearly  $P(B, C)$  contains two triangles which have an edge in common and both of them are of type (i) or (ii) that is a contradiction. This completes the proof.  $\square$

**Remark 2.23.** *Let  $G$  be a torsion-free group and  $B$  and  $C$  be two non-empty finite subsets of  $G$  such that  $1 \in C$ ,  $|C| = 3$  and  $\langle C \rangle$  is not abelian. Suppose that  $\mathcal{C}$  is a cycle of length 4 (a square) in  $P(B, C)$ . Suppose further that  $T \in \mathcal{T}(\mathcal{C})$  and  $T = [h_1, h'_1, h_2, h'_2, h_3, h'_3, h_4, h'_4]$ , where  $h_i, h'_i \in C$  for all  $i \in \{1, 2, 3, 4\}$ , then by Lemma 2.13 and Remark 2.2,  $T$  satisfies exactly one of the following conditions:*

(i)  $h_{i_1} \neq h'_{i_1} = h_{i_2} \neq h'_{i_2} \neq h_{i_3} \neq h'_{i_3} \neq h_{i_4} \neq h'_{i_4} \neq h_{i_1}$ , where

$$(i_1, i_2, i_3, i_4) \in \{(1, 2, 3, 4), (2, 3, 4, 1), (3, 4, 1, 2), (4, 1, 2, 3)\}.$$

(ii)  $h_1 \neq h'_1 \neq h_2 \neq h'_2 \neq h_3 \neq h'_3 \neq h_4 \neq h'_4 \neq h_1$ .

More precisely, we shall speak of a square  $\mathcal{C}$  in  $P(B, C)$  of type (j) if  $T \in \mathcal{T}(\mathcal{C})$  satisfies the condition (j) in the above list (j being i or ii). Note that if  $\mathcal{C}$  is a square of type (i), then  $\mathcal{C}$  is a square obtained from two triangles with one edge in common as Figure 1.

**Remark 2.24.** Let  $G$  be a torsion-free group and  $B$  and  $C = \{1, x, y\}$  be two non-empty finite subsets of  $G$  such that  $\langle C \rangle$  is not abelian. Suppose that  $\mathcal{C}$  is a square of type (ii) in  $P(B, C)$ . By using GAP [12], we know that there are 36 non-equivalent cases for  $\mathcal{C}$ . The relations of such non-equivalent cases are listed in the column labelled by  $R$  of Table 2. It is easy to see that each of the cases 2, 3, 10, 12, 24, 30, 32, 34, 35 and 36 leads to being  $\langle C \rangle$  is an abelian group; each of the cases 1 and 31 implies  $G$  has a non-trivial torsion element. Thus,  $r(\mathcal{C}) = 1$  satisfies one of the relations marked by “\*” s in the column labelled by  $E$  of Table 2. We note that each relation which leads to being  $\langle C \rangle$  an abelian group or  $G$  having a non-trivial torsion element is marked by an  $A$  or a  $T$  in the column labelled by  $E$ , respectively.

**Remark 2.25.** The Klein bottle group has the presentation  $\langle x, y \mid xyx = y \rangle$ , also  $\langle x, y \mid x^2 = y^2 \rangle$  is another presentation of the Klein bottle group. We note that by the first part of the proof of [33, Theorem 3.1], every torsion-free quotient of the Klein bottle group is either abelian or it is isomorphic to the Klein bottle group itself.

TABLE 2. Non-equivalent relations of a cycle of length 4 in  $P(B, C)$ .

n	R	E	n	R	E
1	$x^{-4} = 1$	T	19	$x^{-1}y(xy^{-1})^2 = 1$	*
2	$x^{-3}y^{-1} = 1$	A	20	$x^{-1}y^2xy^{-1} = 1$	*
3	$x^{-3}yx^{-1} = 1$	A	21	$x^{-1}y^3x^{-1} = 1$	*
4	$x^{-2}y^{-2} = 1$	*	22	$x^{-1}y^2x^{-1}y^{-1} = 1$	*
5	$x^{-2}y^{-1}xy^{-1} = 1$	*	23	$x^{-1}y(yx^{-1})^2 = 1$	*
6	$x^{-2}yxy^{-1} = 1$	*	24	$(x^{-1}yx^{-1})^2 = 1$	A
7	$x^{-2}y^2x^{-1} = 1$	*	25	$x^{-1}yx^{-1}y^{-2} = 1$	*
8	$x^{-2}yx^{-1}y^{-1} = 1$	*	26	$x^{-1}yx^{-1}y^{-1}xy^{-1} = 1$	*
9	$x^{-1}(x^{-1}y)^2x^{-1} = 1$	*	27	$(x^{-1}y)^2xy^{-1} = 1$	*
10	$(x^{-1}y^{-1})^2 = 1$	A	28	$(x^{-1}y)^2yx^{-1} = 1$	*
11	$x^{-1}y^{-1}x^{-1}yx^{-1} = 1$	*	29	$(x^{-1}y)^2x^{-1}y^{-1} = 1$	*
12	$x^{-1}y^{-3} = 1$	A	30	$(x^{-1}y)^3x^{-1} = 1$	A
13	$x^{-1}y^{-2}xy^{-1} = 1$	*	31	$y^{-4} = 1$	T
14	$x^{-1}y^{-1}x^2y^{-1} = 1$	*	32	$y^{-3}xy^{-1} = 1$	A
15	$x^{-1}y^{-1}xyx^{-1} = 1$	*	33	$y^{-1}(y^{-1}x)^2y^{-1} = 1$	*
16	$x^{-1}y^{-1}xy^{-2} = 1$	*	34	$(y^{-1}xy^{-1})^2 = 1$	A
17	$x^{-1}(y^{-1}x)^2y^{-1} = 1$	*	35	$(y^{-1}x)^3y^{-1} = 1$	A
18	$x^{-1}yxy^{-2} = 1$	*	36	$(xy^{-1})^4 = 1$	A

**Lemma 2.26.** *Let  $G$  be a torsion-free group and  $B, C$  be two non-empty finite subsets of  $G$  such that  $1 \in C$ . If  $\langle C \rangle$  is neither abelian nor isomorphic to the Klein bottle group, then every two squares of  $P(B, C)$  are equivalent.*


*Proof.* Let  $C = \{1, x, y\}$ . Suppose that there are two squares in  $P(B, C)$ . Since  $\langle C \rangle$  is not isomorphic to the Klein bottle group, Lemma 2.21 and Remarks 2.23 and 2.25 imply that both squares are of type (ii). Hence, it follows from Remark 2.24 that the relation corresponding to each of these two squares must be one of the 24 non-equivalent cases with the relations marked by “\*”s unless the cases 4, 9 and 33 in Table 2 (note that by Remark 2.25, each of the cases 4, 9 and 33 leads to  $\langle C \rangle$  is isomorphic to the Klein bottle group). When choosing two relations different from each other, there are 210 cases. Using GAP [12], we see in 201 cases among these 210 cases, a free group with generators  $x, y$  and relations of each of such cases is finite or abelian that is a contradiction. If the ordered pair of integers  $(i, j)$  shows the numbers of two relations corresponding to each of 210 cases in Table 2, then for the other 9 cases

$$(i, j) \in \{(6, 11), (6, 15), (8, 15), (13, 18), (16, 20), (18, 20), (19, 27), (19, 28), (23, 27)\}.$$

It is easy to see that if  $(i, j)$  is 1-th or 3-th element of the set, then  $x^5 = 1$ ; if  $(i, j)$  is 2-th element of the set, then  $x^3 = 1$ ; if  $(i, j)$  is 4-th or 5-th element of the set, then  $y^5 = 1$ ; if  $(i, j)$  is 6-th element of the set, then  $y^3 = 1$ ; if  $(i, j)$  is 7-th element of the set, then  $(x^{-1}y)^3 = 1$ ; if  $(i, j)$  is 8-th element of the set, then  $(yx^{-1})^3 = 1$ ; if  $(i, j)$  is 9-th element of the set, then  $(xy^{-1})^3 = 1$ . Thus, each of these 9 cases leads to  $G$  having a non-trivial torsion element, a contradiction. This completes the proof.  $\square$

**Lemma 2.27.** *Let  $G$  be a torsion-free group and  $B, C$  be two non-empty finite subsets of  $G$  such that  $1 \in C$ . If  $\langle C \rangle$  is neither abelian nor isomorphic to the Klein bottle group, then  $P(B, C)$  contains no subgraph isomorphic to one of the graphs in Figure 2.*

*Proof.* Let  $C = \{1, x, y\}$ . We note first that since  $\langle C \rangle$  is not isomorphic to the Klein bottle group, Lemmas 2.20 and 2.21 and Remark 2.25 imply that all triangles and squares in  $P(B, C)$  must be of type (i) and (ii), respectively. Suppose that  $P(B, C)$  contains the graph  $\Gamma_1$  in Figure 2. Suppose that  $\mathcal{C}$  and  $\mathcal{C}'$  are two cycles of length 4 in  $\Gamma_1$  which have exactly two consecutive edges in common. Further suppose that  $T \in \mathcal{T}(\mathcal{C})$ ,  $T' \in \mathcal{T}(\mathcal{C}')$ ,  $T = [h_1, h'_1, h_2, h'_2, h_3, h'_3, h_4, h'_4]$  and  $T' = [h_1, h'_1, h_2, h'_2, t_3, t'_3, t_4, t'_4]$ , where the first four components are related to the common edges between these cycles. Lemma 2.26 and Remark 2.18 imply  $\mathcal{T}(\mathcal{C}) = \mathcal{T}(\mathcal{C}')$  and therefore  $r(\mathcal{C}) = r(\mathcal{C}')$ . Also, since  $\langle C \rangle$  is not isomorphic to the Klein bottle group, Lemma 2.21 implies  $h_3 \neq t_3$  and  $h'_4 \neq t'_4$  and hence  $T \neq T'$ . Also, by Lemma 2.13, we may assume that  $h_1 = 1$ . Clearly,  $r(\mathcal{C}) = r(\mathcal{C}') = 1$  is one of the relations marked by “\*”s in Table 2 unless the cases 4, 9 and 33 (21 cases). With every choice of  $r(\mathcal{C}) = 1$  of such cases, it is easy to see that there are no two elements of  $\mathcal{T}(\mathcal{C})$  that satisfy the mentioned conditions for  $T$  and  $T'$ . So,  $P(B, C)$  contains no subgraph isomorphic to the graph  $\Gamma_1$ . Now suppose that  $P(B, C)$  contains two cycles  $\mathcal{C}, \mathcal{C}'$  of length 4 which have exactly one edge in common. Further suppose that  $T \in \mathcal{T}(\mathcal{C})$ ,  $T' \in \mathcal{T}(\mathcal{C}')$ ,  $T = [h_1, h'_1, h_2, h'_2, h_3, h'_3, h_4, h'_4]$  and  $T' = [h_1, h'_1, t_2, t'_2, t_3, t'_3, t_4, t'_4]$ , where the first two components are related to the common edge between these cycles. Lemma 2.26 and Remark 2.18 imply  $\mathcal{T}(\mathcal{C}) = \mathcal{T}(\mathcal{C}')$  and therefore  $r(\mathcal{C}) = r(\mathcal{C}') = 1$ . It is easy to see that either  $h_2 \neq t_2$  or  $h_2 = t_2$  and  $h'_2 \neq t'_2$  and also

FIGURE 2. Some forbidden subgraphs of  $P(B, C)$ .

either  $h'_4 \neq t'_4$  or  $h'_4 = t'_4$  and  $h_4 \neq t_4$  and so  $T \neq T'$ . On the other hand,  $r(C) = r(C') = 1$  is one of the relations marked by “\*”s in Table 2 unless the cases 4, 9 and 33 (21 cases). It can be seen that just in the cases 5,6,7,8,11,13,14,15,16,18,19,20,21 and 22 there are two elements of  $\mathcal{T}(C)$  that satisfy the mentioned conditions for  $T$  and  $T'$ . So, there are 14 cases for the existence of two squares which have exactly one edge in common in  $P(B, C)$ . The graphs  $\Gamma_2$ ,  $\Gamma_3$  and  $\Gamma_4$  obtained from adding a square to two squares which have exactly one edge in common and also the graph  $\Gamma_5$  obtains from adding two triangles to the two squares which have exactly one edge in common. By a same argument as above it can be seen that in each of 14 cases it is impossible that adding a square with properties of graphs  $\Gamma_2$ ,  $\Gamma_3$  and  $\Gamma_4$  or adding two triangles of type (i) with properties of the graph  $\Gamma_5$  to the two squares which have exactly one edge in common. So,  $P(B, C)$  contains no subgraph isomorphic to one of the graphs  $\Gamma_2$ ,  $\Gamma_3$ ,  $\Gamma_4$  and  $\Gamma_5$  in Figure 2.  $\square$

**Remark 2.28.** Let  $G$  be a torsion-free group,  $C$  be a finite subset of  $G$  containing the identity element and  $B$  be a  $k$ -atom of  $C$ , for some positive integer number  $k$ . Since  $xB$  is a  $k$ -atom of  $Cy$  for all  $x \in G$  and  $y \in C^{-1}$ , we may consider the graph  $P(xB, Cy)$  which by Lemma 2.13 is isomorphic to the graph  $P(B, C)$ .

### 3. Small product sets

**Lemma 3.1.** Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $\langle C \rangle$  is not abelian. If  $|C| = 3$  and  $\kappa_4(C) = 4$ , then  $\alpha_4(C) = 4$ .

*Proof.* Let  $A$  be a 4-atom of  $C$ . So,  $|AC| = |A| + 4$ . Suppose, for a contradiction, that  $|A| > 4$ . It follows from Lemma 2.1 that  $|A| \geq 8$  and  $r_{AC}(x) \geq 2$ , for all  $x \in AC$ . If  $|A| = 8$ , then  $|AC| = \frac{|A||C|}{2}$  that according to the proof of Proposition 2.6 and [33, Theorem 1.3], is a contradiction. Hence,  $|A| \geq 9$ . By Lemma 2.1, for every  $a \in A$ , there are  $s_a, t_a, r_a \in C^{-1} \setminus \{1\}$  such that  $A_a = \{a, as_a, as_at_a, as_at_ar_a\} \subseteq A$ . Since  $\langle C \rangle$  is a non-abelian group and the group has no element of order 2 and 3 as it is torsion-free, it follows that  $s_at_a \neq 1$ ,  $t_ar_a \neq 1$  and  $s_at_ar_a \neq 1$ . Hence, for every  $a \in A$ ,  $|A_a| = 4$ . Since  $|C^{-1} \setminus \{1\}| = 2$ , there are 8 choices for the ordered triple  $(s_a, t_a, r_a)$ . So, as  $|A| \geq 9$ , there exist distinct elements  $a, b \in A$  such that  $A_a = A_b$ . Therefore,  $ba^{-1}A_a = A_b \subseteq ba^{-1}A \cap A$ , contradicting [17, Lemma 1]. Hence,  $|A| = 4$ . This completes the proof.  $\square$

**Lemma 3.2.** Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 3$  and  $\langle C \rangle$  is not abelian. If  $\kappa_4(C) = 4$  and  $B$  is a 4-atom of  $C$ , then there exists  $h \in C$  such that  $Ch^{-1} = \{1, a, b\}$  and exactly one of the following statements holds:




FIGURE 3

- (1)  $a^3 = b^2$  and there exists  $a' \in G$  such that  $B = a'\{1, b, a, a^2\}$ .
- (2)  $aba^{-2}b = 1$  and there exists  $a' \in G$  such that  $B = a'\{1, a, b^{-1}, b^{-1}a\}$ .
- (3)  $ab^{-1}aba^{-1}b = 1$  and there exists  $a' \in G$  such that  $B = a'\{1, a, ba^{-1}, aba^{-1}\}$ .
- (4)  $ba^2b^{-1}a = 1$  and there exists  $a' \in G$  such that  $B = a'\{1, a, b^{-1}, b^{-1}a^{-1}\}$ .
- (5)  $b^{-1}a^2ba = 1$  and there exists  $a' \in G$  such that  $B = a'\{1, a^{-1}, a^{-2}, b\}$ .
- (6)  $ba^{-2}b^{-1}a = 1$  and there exists  $a' \in G$  such that  $B = a'\{1, a, b^{-1}, b^{-1}a\}$ .
- (7)  $a^{-2}bab^{-1} = 1$  and there exists  $a' \in G$  such that  $B = a'\{1, a^{-1}, a^{-2}, ba^{-1}\}$ .
- (8)  $a^{-2}b^{-2} = 1$  and there exists  $a' \in G$  such that  $B = a'\{1, a^{-1}, a^{-2}, b\}$ .
- (9)  $a^{-2}b^2 = 1$  and there exists  $a' \in G$  such that  $a'B$  is one of the sets  $\{1, a^{-1}, b^{-1}, ab^{-1}\}$ ,  $\{1, a^{-1}, b^{-1}a, ab^{-1}\}$ ,  $\{1, a, b, a^{-1}\}$ ,  $\{1, a^{-1}, a, ab^{-1}\}$  and  $\{1, a, b^{-1}, ba^{-1}\}$ .

*Proof.* It follows from Lemma 3.1 that  $|B| = 4$ . Let  $B = \{g_1, g_2, g_3, g_4\}$ . As  $\kappa_4(C) = 4$ ,  $|BC| = 8$ . In view of Definition 2.11,  $P(B, C)$  must have at least  $|B||C| - |BC| = 4$  edges. Therefore, it follows from Lemma 2.14 that  $P(B, C)$  contains one of the graphs in Figure 3.

Let  $C = \{1, x, y\}$ . Suppose that  $P(B, C)$  contains an square  $\mathcal{C}$  as Figure 3. Suppose first that  $\mathcal{C}$  is of type (ii). Hence, according to Remark 2.24  $r(\mathcal{C}) = 1$  satisfies one of the relations marked by “\*”s (24 cases) in Table 2. In the following we show that the cases (5), (7), (17), (21), (25) and (29) satisfy the part (1):

- (5)  $x^{-2}y^{-1}xy^{-1} = 1$ : In this case  $[1, x, 1, x, 1, y, x, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3x$ ,  $g_3 = g_4y$  and  $g_4x = g_1y$ . In view of Remark 2.28, we may assume  $g_1 = 1$  and we can replace  $C$  with  $Cx^{-1}$ . Now, if we let  $a = x^{-1}$  and  $b = yx^{-1}$ , then  $Cx^{-1} = \{1, a, b\}$ ,  $a^3 = b^2$  and  $B = \{1, a, a^2, b\}$ .
- (7)  $x^{-2}y^2x^{-1} = 1$ : In this case  $[1, x, 1, x, y, 1, y, x] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3x$ ,  $g_3y = g_4$  and  $g_4y = g_1x$ . Now, by assuming that  $g_1 = 1$ ,  $x^2B = \{1, x, x^2, y\}$ .
- (17)  $x^{-1}(y^{-1}x)^2y^{-1} = 1$ : In this case  $[1, x, 1, y, x, y, x, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3y$ ,  $g_3x = g_4y$  and  $g_4x = g_1y$ . According to Remark 2.28, we may assume  $g_1 = 1$  and we can replace  $C$  with  $Cx^{-1}$ . Now, if we let  $b = x^{-1}$  and  $a = yx^{-1}$ , then we have  $Cx^{-1} = \{1, a, b\}$ ,  $a^3 = b^2$  and  $B = \{1, a, a^2, b\}$ .
- (21)  $x^{-1}y^3x^{-1} = 1$ : In this case, since  $[1, x, y, 1, y, 1, y, x] \in \mathcal{T}(\mathcal{C})$ , by the same argument as above, it can be seen that  $xB = \{1, y, y^2, x\}$ .
- (25)  $x^{-1}yx^{-1}y^{-2} = 1$ : By interchanging  $x$  and  $y$  in (5) and with the same discussion, it can be seen that this case satisfies the part (1).
- (29)  $(x^{-1}y)^2x^{-1}y^{-1} = 1$ : By interchanging  $x$  and  $y$  in (17) and with the same argument, it can be seen that this case satisfies the part (1).

In the following we show that the cases (14) and (22) in Table 2 satisfy the part (2):

- (14)  $x^{-1}y^{-1}x^2y^{-1} = 1$ : In this case  $[1, x, 1, y, x, 1, x, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3y$ ,  $g_3x = g_4$  and  $g_4x = g_1y$ . Now, by assuming that  $g_1 = 1$ ,  $xB = \{1, x, y^{-1}, y^{-1}x\}$ .
- (22)  $x^{-1}y^2x^{-1}y^{-1} = 1$ : By interchanging  $x$  and  $y$  in (14) and with the same argument, it can be seen that this case satisfies the part (2).

With the same argument as above, it can be seen that if  $r(\mathcal{C}) = 1$  satisfies the case (26) in Table 2, then since  $[1, x, y, x, 1, y, x, y] \in \mathcal{T}(\mathcal{C})$ , this case satisfies the part (3). In the following we show that the cases (11), (16) and (28) satisfy the part (4):

- (11)  $x^{-1}y^{-1}x^{-1}yx^{-1} = 1$ : In this case  $[1, x, 1, y, 1, x, y, x] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3y$ ,  $g_3 = g_4x$  and  $g_4y = g_1x$ . Now, by assuming that  $g_1 = 1$ ,  $xB = \{1, x, y^{-1}x^{-1}, y^{-1}\}$ .
- (16)  $x^{-1}y^{-1}xy^{-2} = 1$ : By interchanging  $x$  and  $y$  in (11) and with the same discussion, it can be seen that this case satisfies the part (4).
- (28)  $(x^{-1}y)^2yx^{-1} = 1$ : In this case  $[1, x, y, x, y, 1, y, x] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2y = g_3x$ ,  $g_3 = g_4y$  and  $g_4y = g_1x$ . Now, by assuming that  $g_1 = 1$ , if we let  $b = x^{-1}$  and  $a = yx^{-1}$ , then  $Cx^{-1} = \{1, a, b\}$ ,  $ba^2b^{-1}a = 1$  and  $b^{-1}B = \{1, b^{-1}, a, b^{-1}a^{-1}\}$ .

In the following we show that the cases (8), (13) and (23) in Table 2 satisfy the part (5):

- (8)  $x^{-2}yx^{-1}y^{-1} = 1$ : In this case  $[1, x, 1, x, y, x, 1, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3x$ ,  $g_3y = g_4x$  and  $g_4 = g_1y$ . Now, by assuming that  $g_1 = 1$ ,  $B = \{1, x^{-1}, x^{-2}, y\}$ .
- (13)  $x^{-1}y^{-2}xy^{-1} = 1$ : By interchanging  $x$  and  $y$  in (8) and with the same discussion, it can be seen that this case satisfies the part (5).
- (23)  $x^{-1}y(yx^{-1})^2 = 1$ : In this case  $[1, x, y, 1, y, x, y, x] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2y = g_3$ ,  $g_3y = g_4x$  and  $g_4y = g_1x$ . Now, by assuming that  $g_1 = 1$ , if we let  $b = x^{-1}$  and  $a = yx^{-1}$ , then we have  $Cx^{-1} = \{1, a, b\}$ ,  $a^2bab^{-1} = 1$  and  $B = \{1, a^{-1}, a^{-2}, b\}$ .

In the following we show that the cases (15), (18) and (27) in Table 2 satisfy the part (6):

- (15)  $x^{-1}y^{-1}xyx^{-1} = 1$ : In this case  $[1, x, 1, y, x, 1, y, x] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3y$ ,  $g_3x = g_4$  and  $g_4y = g_1x$ . Now, by assuming that  $g_1 = 1$ ,  $xB = \{1, x, y^{-1}, y^{-1}x\}$ .
- (18)  $x^{-1}yxy^{-2} = 1$ : By interchanging  $x$  and  $y$  in (15) and with the same discussion, it can be seen that this case satisfies the part (6).
- (27)  $(x^{-1}y)^2xy^{-1} = 1$ : In this case  $[1, x, y, x, y, 1, x, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2y = g_3x$ ,  $g_3y = g_4$  and  $g_4x = g_1y$ . Now, by assuming that  $g_1 = 1$ , if we let  $b = y^{-1}$  and  $a = xy^{-1}$ , then  $Cy^{-1} = \{1, a, b\}$ ,  $ba^{-2}b^{-1}a = 1$  and  $a^2b^{-1}B = \{1, a, b^{-1}, b^{-1}a\}$ .

In the following we show that the cases (6), (19) and (20) in Table 2 satisfy the part (7):

- (6)  $x^{-2}yxy^{-1} = 1$ : In this case  $[1, x, 1, x, y, 1, x, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3x$ ,  $g_3y = g_4$  and  $g_4x = g_1y$ . Now, by assuming that  $g_1 = 1$ ,  $B = \{1, x^{-1}, x^{-2}, yx^{-1}\}$ .
- (19)  $x^{-1}y(xy^{-1})^2 = 1$ : In this case  $[1, x, y, 1, x, y, x, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2y = g_3$ ,  $g_3x = g_4y$  and  $g_4x = g_1y$ . Now, by assuming that  $g_1 = 1$ , if we let  $b = y^{-1}$  and  $a = xy^{-1}$ , then  $Cy^{-1} = \{1, a, b\}$ ,  $a^2ba^{-1}b^{-1} = 1$  and  $B = \{1, a^{-1}, a^{-2}, ba^{-1}\}$ .
- (20)  $x^{-1}y^2xy^{-1} = 1$ : By interchanging  $x$  and  $y$  in (6) and with the same discussion, it can be seen that this case satisfies the part (7).

In the following we show that the cases (4), (9) and (33) in Table 2 satisfy the part (8):

- (4)  $x^{-2}y^{-2} = 1$ : In this case  $[1, x, 1, x, 1, y, 1, y] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3x$ ,  $g_3 = g_4y$  and  $g_4 = g_1y$ . Now, by assuming that  $g_1 = 1$ ,  $B = \{1, x^{-1}, x^{-2}, y\}$ .
- (9)  $x^{-1}(x^{-1}y)^2x^{-1} = 1$ : In this case  $[1, x, 1, x, y, x, y, x] \in \mathcal{T}(\mathcal{C})$  and therefore we may assume  $g_1 = g_2x$ ,  $g_2 = g_3x$ ,  $g_3y = g_4x$  and  $g_4y = g_1x$ . Now, by assuming that  $g_1 = 1$ , if we let  $b = x^{-1}$  and  $a = yx^{-1}$ , then  $Cx^{-1} = \{1, a, b\}$ ,  $b^2a^2 = 1$  and  $b^{-2}B = \{1, a^{-1}, a^{-2}, b\}$ .
- (33)  $y^{-1}(y^{-1}x)^2y^{-1} = 1$ : By interchanging  $x$  and  $y$  in (9) and with the same discussion, it can be seen that this case satisfies the part (8).

On the other hand, by Lemma 2.21, if the square  $\mathcal{C}$  is of type (i), then the part (9) holds. Now, suppose that  $P(B, C)$  contains the graph  $\Gamma$  as Figure 3. If the triangle which is included in the graph  $\Gamma$  is of type (i), then there exists  $x \in BC$  such that  $r_{BC}(x) = 3$  and therefore since  $|BC| = 8$ ,  $P(B, C)$  must have at least 5 edges and therefore  $P(B, C)$  contains a graph isomorphic to the graph in Figure 1. So, we may assume that the triangle which is included in the graph  $\Gamma$  is of type (ii). Then lemma 2.20 implies  $r(\mathcal{C}) \in \{x^{-1}y^{-1}xy^{-1}, x^{-1}yx^{-1}y^{-1}, x^{-1}y^2x^{-1}\}$ . Suppose that  $T = [h_1, h'_1, h_2, h'_2, h_3, h'_3]$  is the 6-tuple to  $\mathcal{C}$  corresponding to the arrangement  $g_1, g_2, g_3$ . Clearly, there exist  $h, h' \in C$  such that  $g_1h = g_4h'$ . Note that we may assume  $g_1 = 1$  (by Remark 2.28) and  $h \neq h_1$  and  $h \neq h'_3$  since otherwise  $P(B, C)$  contains a graph isomorphic to the graph in Figure 1. It is easy to see that every choice for  $r(\mathcal{C})$  leads to holding the part (9). We will prove the case  $r(\mathcal{C}) = x^{-1}y^{-1}xy^{-1}$  and  $T = [1, x, 1, y, x, y]$  only as the proof of the other cases are similar. In this case,  $g_1 = g_2x$ ,  $g_2 = g_3y$ ,  $g_3x = g_1y$  and  $g_1x = g_4h'$ , where  $h' \in \{1, y\}$ . Hence,  $B$  is one of the sets  $\{1, x, x^{-1}, yx^{-1}\}$  and  $\{1, xy^{-1}, x^{-1}, yx^{-1}\}$ . Now, if we let  $b = yx^{-1}$  and  $a = x^{-1}$ , then  $Cx^{-1} = \{1, a, b\}$ ,  $a^2 = b^2$  and  $B$  is one of the sets  $\{1, a, a^{-1}, b\}$  and  $\{1, b, b^{-1}, a\}$ . This completes the proof.  $\square$

**Lemma 3.3.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 3$  and  $\langle C \rangle$  is not abelian. If  $\kappa_5(C) = 4$ , then  $\alpha_5(C) = 5$ .*

*Proof.* If  $\langle C \rangle$  is isomorphic to the Klein bottle group, then  $\langle C \rangle$  is a unique product group [27] and so there is nothing to prove. So, we may assume that  $\langle C \rangle$  is not isomorphic to the Klein bottle group. Let  $A$  be a 5-atom of  $C$ . Suppose, for a contradiction, that  $|A| > 5$ . Thus, Lemma 2.1 and Remark 2.2 imply  $|AC| \geq 8$  and  $r_{AC}(t) \in \{2, 3\}$ , for all  $t \in AC$ . If  $|A| = 8$  and  $|A| = 9$ , then  $|AC| = \frac{3|A|}{2}$  and  $|AC| = \frac{3|A|-1}{2}$ , respectively, that according to the proof of Propositions 2.6 and 2.7, [33, Theorem 1.3] and [6, Proposition 4.12] are contradictions. Thus,  $|A| \geq 10$ . Now, suppose that  $|A| \geq 13$ . Let  $C = \{1, x, y\}$ . Observe that, since  $r_{AC}(ay) \geq 2$ , for all  $a \in A$ ,  $|AC| = |A\{1, x\}|$ . Hence,  $|A \cap Ax| = |A| - 4 \geq 9$ . Therefore, by Lemma 2.1, for every  $a \in A \cap Ax$ , there are  $s_a, t_a, r_a \in C^{-1} \setminus \{1\}$  such that  $A_a = \{a, ax^{-1}, ax^{-1}s_a, ax^{-1}s_at_a, ax^{-1}s_at_ar_a\} \subseteq A$ . Since  $\langle C \rangle$  is a non-abelian torsion-free group and also  $\langle C \rangle$  is not isomorphic to the Klein bottle group, for every  $a \in A \cap Ax$ ,  $|A_a| = 5$ . Since  $|C^{-1} \setminus \{1\}| = 2$ , there are 8 choices for the ordered triple  $(s_a, t_a, r_a)$ . So, as  $|A \cap Ax| \geq 9$ , there exist distinct elements  $a, b \in A$  such that  $A_a = A_b$ . Therefore,  $ba^{-1}A_a = A_b \subseteq (ba^{-1})A \cap A$ , contradicting [17, Lemma 1]. Hence,  $10 \leq |A| \leq 12$ . Since for each  $t \in AC$ ,  $r_{AC}(t) \in \{2, 3\}$ , if  $T = \{t \in AC \mid r_{AC}(t) = 3\}$ , then  $|T| = |A||C| - 2|AC|$  and therefore  $|T| = |A| - 8$ . It is clear that, for each element  $t \in T$ , there exists

a triangle of type (i) in  $S(A, C)$ . Since  $\langle C \rangle$  is not isomorphic to the Klein bottle group, it follows from Lemma 2.20 that  $S(A, C)$  contains no triangle of type (ii). So, if  $|A| = n$ , then  $S(A, C)$  contains exactly  $n - 8$  triangles. On the other hand, by Lemma 2.14 and Lemma 2.21, each two triangles in  $S(A, C)$  have at most one vertex in common. Also, since for each  $g \in A$  and  $h \in C$ ,  $r_{AC}(gh) \in \{2, 3\}$ , it is easy to see that the degree of each vertex  $g$  in  $S(A, C)$  is equal to  $3 + s$ , where  $s = |\{h \mid h \in C, r_{AC}(gh) = 3\}|$ . So, it is clear that the degree of every vertex in  $S(A, C)$  is equal to 3, 4, 5 or 6. Then,  $S(A, C)$  has the following properties: (1) contains exactly  $n - 8$  triangles each two of which triangles have at most one vertex in common; (2) every vertex of it has degree 3, 4, 5 or 6; (3) the degree of each its vertex  $g$  is equal to  $3 + s'$ , where  $s'$  is the number of triangles which  $g$  is a common vertex between them; (4) contains no subgraph isomorphic to one of the graphs in Figure 2. For each  $n \in \{10, 11, 12\}$ , we check and see that there is no graph with  $n$  vertices and with above properties. Thus,  $|A| \notin \{10, 11, 12\}$  and therefore  $\alpha_5(C) \neq 5$ . This completes the proof.  $\square$

**Theorem 3.4.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 3$  and  $\langle C \rangle$  is not abelian. If  $\kappa_5(C) \leq 4$  and  $B$  is a 5-atom of  $C$ , then there exist  $h \in C$  and  $a' \in G$  such that  $Ch^{-1} = \{1, a, b\}$ ,  $a^2 = b^2$  and  $a'B$  is one of the sets  $\{1, a^{-1}, b^{-1}, b^{-1}a, ab^{-1}\}$ ,  $\{1, a, b, a^{-1}, ab^{-1}\}$  and  $\{1, a^{-1}, a, ab^{-1}, aba^{-1}\}$ .*

*Proof.* Let  $C = \{1, x, y\}$ . Since  $\kappa_5(C) \leq 4$ , it follows from [17, Lemma 8] that  $\kappa_4(C) = \kappa_5(C) = 4$ . Let  $B$  be a 5-atom of  $C$ . By Lemma 3.3,  $|B| = 5$ . Also, since  $\kappa_5(C) = 4$ ,  $|BC| = 9$ . So, there exist  $b \in B$  and  $c \in C$  such that  $r_{BC}(bc) = 1$  since otherwise we must have  $|BC| \leq \frac{|B||C|}{2}$ , a contradiction. By the choice of  $b$ , if  $B' = B \setminus \{b\}$ , then clearly  $\kappa_4(C) \leq |\partial_C(B')| \leq |\partial_C(B)| = 4$ . Therefore,  $B'$  is a 4-atom of  $C$  and  $|\partial_C(B')| = 4$ . So, one of the 9 parts of Lemma 3.2 holds. Suppose first that the part (9) holds. In other word, we may assume that  $x, y$  satisfy  $x^2 = y^2$  and  $B$  is one of sets  $\{1, x^{-1}, y^{-1}, xy^{-1}, b\}$ ,  $\{1, x^{-1}, y^{-1}x, xy^{-1}, b\}$ ,  $\{1, x, y, x^{-1}, b\}$ ,  $\{1, x^{-1}, x, xy^{-1}, b\}$  and  $\{1, x, y^{-1}, yx^{-1}, b\}$ . Since  $|B'C| = 8$  and  $|BC| = 9$ , there exists an element  $a' \in \{b, bx, by\}$  such that  $BC = B'C \cup \{a'\}$  and  $T = \{b, bx, by\} \setminus \{a'\} \subseteq B'C$ . We checked all possible choices for  $a'$ , many of them lead to either  $\langle C \rangle$  has a non-trivial torsion element or  $\langle C \rangle$  is an abelian group, that are contradictions. Just in the following cases, we have no contradiction: if  $B = \{1, x^{-1}, y^{-1}, xy^{-1}, b\}$ , then  $B'C = \{1, x^{-1}, y^{-1}, xy^{-1}, x, y^{-1}x, xy^{-1}x, y\}$  and there are two possible cases: (1)  $b = x^{-2}$  which leads  $\{bx, by\} = \{x^{-1}, y^{-1}\} \subseteq B'C$ ,  $BC = B'C \cup \{x^{-2}\}$ ; (2)  $b = y^{-1}x$  which leads  $\{bx, b\} = \{y, y^{-1}x\} \subseteq B'C$ ,  $BC = B'C \cup \{y^{-1}xy\}$ ; if  $B = \{1, x^{-1}, y^{-1}x, xy^{-1}, b\}$ , then  $B'C = \{1, x^{-1}, y^{-1}x, xy^{-1}, x, y^{-1}xy, xy^{-1}x, y\}$  and the only possibility is  $b = y^{-1}$  which leads to  $\{bx, by\} = \{1, y^{-1}x\} \subseteq B'C$ ,  $BC = B'C \cup \{y^{-1}\}$ ; if  $B = \{1, x, y, x^{-1}, b\}$ , then  $B'C = \{1, x, y, x^{-1}, xy, x^{-1}y, x^2, yx\}$  and the only possibility is  $b = x^{-1}y$  which leads to  $\{b, by\} = \{x^{-1}y, x\} \subseteq B'C$ ,  $BC = B'C \cup \{x^{-1}yx\}$ ; if  $B = \{1, x^{-1}, x, xy^{-1}, b\}$ , then  $B'C = \{1, x^{-1}, x, xy^{-1}, x^2, xy^{-1}x, xy, y\}$  and there are two possible cases: (1)  $b = y$  which leads to  $\{b, by\} = \{y, x^2\} \subseteq B'C$ ,  $BC = B'C \cup \{yx\}$ ; (2)  $b = xyx^{-1}$  which leads to  $\{b, bx\} = \{xy^{-1}x, xy\} \subseteq B'C$ ,  $BC = B'C \cup \{xyx^{-1}y\}$ ; if  $B = \{1, x, y^{-1}, yx^{-1}, b\}$ , then  $B'C = \{1, y^{-1}, x, yx^{-1}, x^2, yx^{-1}y, xy, y\}$  and the only possibility is  $b = y$  which leads to  $\{b, by\} = \{x^2, y\} \subseteq B'C$ ,  $BC = B'C \cup \{yx\}$ . By a same argument as above we checked all other parts (1) to (8) of Lemma 3.2, in every of these parts, every



choice for  $b$  implies that either  $\langle C \rangle$  has a non-trivial torsion element or  $\langle C \rangle$  is an abelian group, that are contradictions. This completes the proof.  $\square$

The following Corollary follows from Theorem 3.4 and Remark 2.25.

**Corollary 3.5.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 3$  and  $\langle C \rangle$  is neither abelian nor isomorphic to the Klein bottle group. Then for all subsets  $B \subseteq G$  with  $|B| \geq 5$ ,  $|BC| \geq |B| + 5$ .*

**Lemma 3.6.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 3$  and  $\langle C \rangle$  is not abelian. If  $\kappa_6(C) \leq 4$  and  $B$  is a 6-atom of  $C$ , then there exist  $h \in C$  and  $a' \in G$  such that  $Ch^{-1} = \{1, a, b\}$ ,  $a^2 = b^2$  and  $a'B = \{1, a^{-1}, b^{-1}, b^{-1}a, ab^{-1}, a^{-2}\}$ .*

*Proof.* Let  $C = \{1, x, y\}$ . Without loss of generality, we may assume that  $1 \in B$ . Since  $\kappa_6(C) \leq 4$ ,  $|BC| \leq |B| + 4$  and therefore by Corollary 3.5 we may assume that  $\langle C \rangle$  is isomorphic to the Klein bottle group. Hence,  $\langle C \rangle$  is a unique product group [27]. Thus,  $|B| = 6$  and there exist  $b \in B$  and  $c \in C$  such that  $r_{BC}(bc) = 1$ . By the choice of  $b$ , if  $B' = B \setminus \{b\}$ , then by [17, Lemma 8],  $4 \leq \kappa_4(C) \leq |\partial_C(B')| \leq |\partial_C(B)| = 4$ . Therefore,  $B'$  is a 5-atom of  $C$  and  $|\partial_C(B')| = 4$ . So, in view of Lemma 3.4 we may assume that  $x, y$  satisfy  $x^2 = y^2$  and  $B$  is one of sets  $\{1, x^{-1}, y^{-1}, y^{-1}x, xy^{-1}, b\}$ ,  $\{1, x, y, x^{-1}, xy^{-1}, b\}$  and  $\{1, x^{-1}, x, xy^{-1}, xyx^{-1}, b\}$ . Since  $|B'C| = 9$  and  $|BC| = 10$ , there exists an element  $a' \in \{b, bx, by\}$  such that  $BC = B'C \cup \{a'\}$  and  $T = \{b, bx, by\} \setminus \{a'\} \subseteq B'C$ . We checked all possible choices for  $a'$ , many of them lead to either  $G$  has a non-trivial torsion element or being  $\langle C \rangle$  is an abelian group, that are contradictions. Just in the following cases, we have no contradiction: if  $B = \{1, x^{-1}, y^{-1}, y^{-1}x, xy^{-1}, b\}$ , then  $B'C = \{1, x^{-1}, y^{-1}, xy^{-1}, x, y^{-1}x, xy^{-1}x, y, yx^{-1}y\}$  and the only possibility is  $b = x^{-2}$  which leads to  $\{bx, by\} = \{x^{-1}, y^{-1}\} \subseteq B'C$ ,  $BC = B'C \cup \{x^{-2}\}$ ; if  $B = \{1, x, y, x^{-1}, xy^{-1}, b\}$ , then  $B'C = \{1, x^{-1}, y, xy^{-1}, x, x^2, xy, yx, xy^{-1}x\}$  and the only possibility is  $b = xy^{-1}x$  which leads to  $\{bx, b\} = \{xy, xy^{-1}x\} \subseteq B'C$ ,  $BC = B'C \cup \{xy^{-1}xy\}$ ; if  $B = \{1, x^{-1}, x, xy^{-1}, xyx^{-1}, b\}$ , then  $B'C = \{1, x^{-1}, y, xy^{-1}, x, x^2, xy, xy^{-1}xy, xy^{-1}x\}$  and the only possibility is  $b = y$  which leads to  $\{by, b\} = \{y, y^2\} \subseteq B'C$ ,  $BC = B'C \cup \{yx\}$ . Hence, in every case, there is  $a \in G$  such that  $aB$  is equal to  $\{1, x^{-1}, y^{-1}, y^{-1}x, xy^{-1}, x^{-2}\}$ . This completes the proof.  $\square$

**Corollary 3.7.** *Let  $G$  be a torsion-free group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 3$  and  $\langle C \rangle$  is not abelian. Then for all subsets  $B \subseteq G$  with  $|B| \geq 7$ ,  $|BC| \geq |B| + 5$ .*

*Proof.* Let  $C = \{1, x, y\}$ . By Corollary 3.5, if  $\langle C \rangle$  is not isomorphic to the Klein bottle group, then there is nothing to prove. Hence, we may assume that  $\langle C \rangle$  is isomorphic to the Klein bottle group. It is sufficient to prove  $\kappa_7(C) \geq 5$ . Suppose, for a contradiction, that  $\kappa_7(C) \leq 4$ . Let  $A$  be a 7-atom of  $C$ . Since the Klein bottle group is a unique product group [27],  $|A| = 7$  and also there exist  $a \in A$  and  $c \in C$  such that  $r_{AC}(ac) = 1$ . By the choice of  $a$ , if  $A' = A \setminus \{a\}$ , then by [17, Lemma 8],  $4 \leq \kappa_6(C) \leq |\partial_C(A')| \leq |\partial_C(A)| \leq 4$ . Therefore,  $|\partial_C(A')| = 4$ . So, in view of Lemma 3.6 we may assume that  $x, y$  satisfy  $x^2 = y^2$  and  $A = \{1, x^{-1}, y^{-1}, y^{-1}x, xy^{-1}, x^{-2}, a\}$ . Since  $|A'C| = 10$  and  $|AC| = 11$ , there exists an element  $a' \in \{a, ax, ay\}$  such that  $AC = A'C \cup \{a'\}$  and  $\{a, ax, ay\} \setminus \{a'\} \subseteq A'C$ . It is

not hard to see that every choice for  $a'$  leads to either  $\langle C \rangle$  has a non-trivial torsion element or  $\langle C \rangle$  is an abelian group, that are contradictions. Hence,  $\kappa_7(C) \geq 5$ . This completes the proof.  $\square$

**Lemma 3.8.** *Let  $G$  be a unique product group and  $C$  be a finite subset of  $G$  containing the identity element such that  $|C| = 4$  and  $\langle C \rangle$  is not abelian. Then  $\kappa_7(C) \geq 6$ .*

*Proof.* Suppose, for a contradiction, that  $\kappa_7(C) \leq 5$ . So, [17, Lemma 8] implies  $\kappa_7(C) = 5$ . Let  $A$  be a 7-atom of  $C$  containing the identity element. Since  $G$  is a unique product group,  $|A| = 7$ . Suppose that  $\langle C \rangle \neq \langle A \rangle$ . Hence, either  $C$  intersects at least two right cosets of  $\langle A \rangle$  or  $A$  intersects at least two left cosets of  $\langle C \rangle$ . Suppose first that  $C$  intersects at least two right cosets of  $\langle A \rangle$ . Let  $C_1$  be one of these intersections. Thus, 1.1 implies  $12 = |AC| = |AC_1| + |A(C \setminus C_1)| \geq 2|A| + |C| - 2 = 16$ , a contradiction. Now suppose that  $A$  intersects at least two left cosets of  $\langle C \rangle$ . Let  $A_1$  be one of these intersections. Thus, 1.1 implies  $12 = |AC| = |A_1C| + |(A \setminus A_1)C| \geq |A| + 2|C| - 2 = 13$ , a contradiction. So,  $\langle A \rangle = \langle C \rangle$ .

Since  $G$  is a unique product group, there exist  $a \in A$  and  $c \in C$  such that  $r_{AC}(ac) = 1$ . Let  $C' = C \setminus \{c\}$ . Suppose that  $c = 1$ . Then if we replace  $C$  with  $Ct^{-1}$ , where  $t \in C \setminus \{1\}$ , then  $\langle Ct^{-1} \rangle$  is not abelian,  $1 \in Ct^{-1}$ ,  $A$  is a 7-atom of  $Ct^{-1}$  and  $r_{AC}(at^{-1}) = 1$ . So, without loss of generality, we may assume that  $1 \in C'$ . It is clear that  $ac \notin Ac \cap AC'$  and therefore

$$(3.1) \quad |AC| = |AC'| + |Ac| - |AC' \cap Ac| \geq |AC'| + 1.$$

Thus,  $|AC'| \leq 11$  and therefore Corollary 3.7 implies  $\langle C' \rangle \neq \langle C \rangle$ . Since  $\langle A \rangle = \langle C \rangle$ ,  $A$  intersects at least two left cosets of  $\langle C' \rangle$ . Partition  $A = A_1 \cup A_2 \cup \dots \cup A_t$ , where each  $A_i$  is the nonempty intersection of  $A$  with some left coset of  $\langle C' \rangle$ . Then  $|AC'| = \sum_{i=1}^t |A_i C'|$ . By 1.1,  $|A_i C'| \geq |A_i| + 2$ , for all  $i \in \{1, \dots, t\}$ , and by the main result of [4], if  $|A_i C'| = |A_i| + 2$ , then  $A_i$  and  $C'$  are left and right progressions with common ratio, respectively. According to the above condition, the only possibility is  $t = 2$  and  $|A_i C'| = |A_i| + 2$  for each  $i \in \{1, 2\}$ . Therefore, without loss of generality, we may assume that  $A = \{1, x, \dots, x^{(i-1)}\} \cup \{r, rx, \dots, rx^{(j-1)}\}$  and  $C' = \{1, x, x^2\}$ , where  $r, x \in G \setminus \{1\}$ ,  $i + j = 7$  and  $i > j$ . Since  $\langle C \rangle = \langle A \rangle$  is not abelian,  $rx \neq xr$  and  $xc \neq cx$ . Note that we may assume that for each  $a' \in A$ ,  $r_{AC}(a') \geq 2$  since otherwise by the choice  $c = 1$ ,  $\langle C' \rangle$  is not abelian which leads to a contradiction. It is clear that  $(i, j) \in \{(6, 1), (5, 2), (4, 3)\}$ . Suppose first that  $(i, j) = (6, 1)$ . In this case  $AC' = \{1, x, \dots, x^7, r, rx, rx^2\}$ . Since  $|AC| = 12$ , there exists  $h \in Ac$  such that  $AC = AC' \cup \{h\}$  and  $Ac \setminus \{h\} \subseteq AC'$ . So, since  $\langle C \rangle$  is not abelian, there exists  $B \subseteq \{c, xc, x^2c, x^3c, x^4c, x^5c\}$  such that  $|B| \geq 4$  and  $B \subseteq \{r, rx, rx^2\}$ . Thus, there are distinct elements  $i', j' \in \{0, 1, 2, 3, 4, 5\}$  such that  $x^{i'}c = x^{j'}c$  which leads to  $G$  has a non-trivial torsion-element, a contradiction. Now, suppose that  $(i, j) = (5, 2)$ . In this case,  $AC' = \{1, x, \dots, x^6, r, rx, rx^2, rx^3\}$  and there exists  $h \in Ac$  such that  $AC = AC' \cup \{h\}$  and so  $Ac \setminus \{h\} \subseteq AC'$ . Since  $\langle C \rangle$  is not abelian, there exists  $B \subseteq \{c, xc, x^2c, x^3c, x^4c\}$  such that  $|B| \geq 4$  and  $B \subseteq \{r, rx, rx^2, rx^3\}$ . Hence, we must have  $|B| = 4$  since otherwise  $G$  has a non-trivial torsion element, a contradiction. Thus,  $h \in \{c, xc, x^2c, x^3c, x^4c\}$ . Therefore,  $\{rc, rxc\} \subseteq AC'$  and therefore  $\{rc, rxc\} \subseteq \{1, x, \dots, x^6\}$ . Thus, there are distinct elements  $i', j' \in \{0, 1, \dots, 6\}$  such that  $rc = x^{i'}$  and  $rxc = x^{j'}$ . On the other hand, since  $r_{AC}(r) \geq 2$ ,  $G$  is a torsion-free group and  $\langle A \rangle$  is not abelian, there exists  $s \in \{0, 1, 2, 3, 4\}$  such that  $r = x^s c$ . Then, we have  $c^2 = x^{(i'-s)}$ ,  $xcx = x^{(j'-s)}$  and  $c^{-1}xc = x^{(j'-i')}$ . It is clear that  $s \notin \{i', j'\}$  since otherwise we have contradiction with  $\langle C \rangle$  is a non-abelian torsion-free

group. So,  $c^{-1}x^{(i'-s)}c = x^{(j'-i')(i'-s)}$  which leads to  $x^{(j'-i'-1)(i'-s)} = 1$ . Hence, we must have  $j' - i' = 1$  implies  $xc = cx$  that is a contradiction. For the last case, suppose that  $(i, j) = (4, 3)$ . In this case,  $AC' = \{1, x, \dots, x^5, r, rx, rx^2, rx^3, rx^4\}$  and there exists  $h \in Ac$  such that  $AC = AC' \cup \{h\}$  and therefore  $Ac \setminus \{h\} \subseteq AC'$ . Consider two cases: (1)  $h = c$ : in this case  $\{rc, rxc, rx^2c\} \subseteq AC'$  and by the same argument as the latter case, we get into a contradiction; (2)  $h \neq c$ : so,  $c \in AC'$  and since  $\langle C \rangle$  is not abelian,  $c \in \{r, rx, rx^2, rx^3, rx^4\}$ . Thus, there exists  $i' \in \{0, 1, 2, 3, 4\}$  such that  $c = rx^{i'}$ . On the other hand, since  $r_{AC}(1)$  and  $r_{AC}(r)$  are greater than or equal to 2,  $\langle C \rangle$  is a non-abelian torsion-free group and  $\langle A \rangle$  is not abelian, there exists  $s \in \{0, 1, 2, 3\}$  and  $s' \in \{0, 1, 2\}$  such that  $r = x^s c$  and  $1 = rx^{s'} c$ . Then we have  $c^2 = x^{(i'-s')}$ ,  $x^s c x^{s'} c = 1$  and  $cx^{i'} c^{-1} = x^{-s}$ . It is clear that  $i' \neq s'$  since otherwise  $G$  has a non-trivial torsion element, a contradiction. Hence,  $cx^{i'(i'-s')} c^{-1} = x^{-s(i'-s')}$  which leads to  $x^{(-s-i')(i'-s')} = 1$ . Then,  $s = i' = 0$  and therefore  $r = c$  and  $r^2 = x^{-s'}$ . So, the only possibility is  $s' = 2$  which implies that  $\{r^2, xr\} \subseteq Ac$  and  $\{r^2, xr\} \not\subseteq AC'$  since otherwise  $G$  has a non-trivial torsion element. Hence,  $|AC| \geq 13$ , a contradiction. Thus,  $\kappa_7(C) \geq 6$  and this completes the proof.  $\square$

**Theorem 3.9.** *Let  $G$  be a unique product group and  $C$  be a finite subset of  $G$  containing the identity element such that  $\langle C \rangle$  is not abelian. Then for all subsets  $B$  of  $G$  with  $|B| \geq 7$ ,  $|BC| \geq |B| + |C| + 2$ .*

*Proof.* It is sufficient to prove  $\kappa_7(C) \geq |C| + 2$ . Suppose the contrary and choose a counter-example with minimal  $|C|$ . Since  $\langle C \rangle$  is not abelian and by Lemma 3.8 and Corollary 3.7,  $|C| \geq 5$ . Let  $A$  be a 7-atom of  $C$  containing the identity element. Since  $G$  is a unique product group,  $|A| = 7$ . By the same argument as the proof of Lemma 3.8, we have  $\langle C \rangle = \langle A \rangle$ . Since  $G$  is a unique product group, there exists an element  $x \in AC$  that can be represented in a unique way in the form  $ac$  with  $a \in A$  and  $c \in C$ . Let  $C' = C \setminus \{c\}$ . Then  $x \notin Ac \cap AC'$  and therefore

$$(3.2) \quad |AC| = |AC'| + |Ac| - |AC' \cap Ac| \geq |AC'| + 1.$$

Without loss of generality, we may assume that  $1 \in C'$ . Consider two cases: (1)  $\langle C \rangle = \langle C' \rangle$ : then the minimality of  $|C|$  implies that  $|AC'| \geq |A| + |C'| + 2$  and by 3.2,  $|AC| \geq |A| + |C| + 2$  that is a contradiction; (2)  $\langle C \rangle \neq \langle C' \rangle$ : then  $A$  intersects at least two left cosets of  $\langle C' \rangle$ . Let  $A_1$  be one of these intersections. Hence, 1.1 implies  $|AC'| = |A_1 C'| + |(A \setminus A_1)C'| \geq |A| + 2|C'| - 2 \geq |A| + |C'| + 2$ , and by 3.2,  $|AC| \geq |A| + |C| + 2$  that is a contradiction. This completes the proof.  $\square$

**Corollary 3.10.** *Let  $\alpha$  and  $\beta$  be non-zero elements of  $\mathbb{F}[G]$ , the group algebra of any torsion-free group over an arbitrary field. If  $|supp(\alpha)| = 3$  and  $\alpha\beta = 0$ , then  $|supp(\beta)| \geq 12$ .*

*Proof.* By [2, Theorem 1.4], it is sufficient to prove  $|supp(\beta)| \notin \{10, 11\}$ . Since  $\alpha\beta = 0$ ,  $\beta^* \alpha^* = 0$ , where  $supp(\alpha^*) = supp(\alpha)^{-1}$  and  $supp(\beta^*) = supp(\beta)^{-1}$ . Let  $B = supp(\beta^*)$  and  $C = supp(\alpha^*)$ . By [1, Lemma 2.7] and since  $\beta^* \alpha^* h^{-1} = 0$ , for all  $h \in C$ , we may assume that  $1 \in C$  and  $G = \langle C \rangle$ . Since  $\beta^* \alpha^* = 0$ ,  $r_{BC}(x) \geq 2$ , for all  $x \in BC$ . Hence,  $|BC| \leq \frac{3|B|}{2}$ . On the other hand, since  $G$  is not abelian (see [27, Theorem 26.2], Corollary 3.7 implies  $|BC| \geq |B| + 5$ . Thus,  $|B| + 5 \leq |BC| \leq \frac{3|B|}{2}$ . Suppose first that  $|B| = 10$ . Then  $|BC| = 15$  and therefore  $r_{BC}(x) = 2$ , for all  $x \in BC$ . Hence, if we let  $\delta = \sum_{b \in B} b$  and  $\gamma = \sum_{c \in C} c$ , then  $\delta, \gamma \in \mathbb{F}_2[G]$  and  $\delta\gamma = 0$  which by [33, Theorem 1.3] is a contradiction. Now, suppose that  $|B| = 11$ . Hence,  $|BC| = 16$  and therefore there exist  $x \in BC$  such that  $r_{BC}(x) = 3$  and

$r_{BC}(x') = 2$ , for all  $x' \in BC \setminus \{x\}$ . Suppose that  $(b, c) \in R_{BC}(x)$ . Hence, if we let  $\delta = \sum_{b \in B} b$  and  $\gamma = \sum_{c \in C} c$ , then  $\delta, \gamma \in \mathbb{F}_2[G]$  and  $b^{-1}\delta\gamma c^{-1} = 1$  which by [6, Proposition 4.12] is a contradiction.  $\square$

**Corollary 3.11.** *Let  $\delta$  and  $\gamma$  be elements of the group algebra of any torsion-free group over an arbitrary field. If  $|\text{supp}(\delta)| = 3$  and  $\delta\gamma = 1$ , then  $|\text{supp}(\gamma)| \geq 10$ .*

*Proof.* By [2, Theorem 1.7], it is sufficient to prove  $|B| \neq 9$ . Suppose, for a contradiction, that  $|B| = 9$ . Let  $B = \text{supp}(\gamma)$  and  $C = \text{supp}(\delta)$ . Without loss of generality, we may assume that  $\gamma\delta = 1$ . Since  $\gamma\delta = 1$ , there exist  $b \in B$  and  $c \in C$  such that  $bc = 1$  and also  $r_{BC}(x) \geq 2$ , for all  $x \in BC \setminus \{1\}$ . Therefore,  $|BC| \leq 14$ . Also, since  $b^{-1}\gamma\delta c^{-1} = 1$ , by [1, Lemma 2.9] we may assume that  $1 \in C$  and  $G = \langle C \rangle$ . Now since  $G$  is not abelian (see [27, Theorem 26.2], Corollary 3.7 implies  $|BC| \geq 14$ . Then,  $|BC| = 14$ . Thus, the only possibility is  $r_{BC}(1) = 1$  and  $r_{BC}(x) = 2$ , for all  $x \in BC \setminus \{1\}$ . So, if we let  $\alpha = \sum_{b \in B} b$  and  $\beta = \sum_{c \in C} c$ , then  $\alpha, \beta \in \mathbb{F}_2[G]$  and  $\alpha\beta = 1$  which by [6, Proposition 4.12] is a contradiction. This completes the proof.  $\square$

## REFERENCES

- [1] A. Abdollahi and F. Jafari, Zero divisor and unit elements with supports of size 4 in group algebras of torsion-free groups, to appear in Comm. Algebra. arXiv:1709.08204 [math.GR].
- [2] A. Abdollahi and Z. Taheri, Zero divisors and units with small supports in group algebras of torsion-free groups, Comm. Algebra, 46 (2018), no. 2, 887-925.
- [3] V. G. Bardakov and M. S. Petukhova, On potential counterexamples to the problem of zero divisors, J. Math. Sci. (N.Y.), 221 (2017), no. 6, 778-797.
- [4] L. V. Brailovsky and G. A. Freiman, On a product of finite subsets in a torsion-free group, J. Algebra, 130 (1990), 462-476.
- [5] K. A. Brown, On zero divisors in group rings, Bull. London. Math. Soc., 8 (1976), no. 3, 251- 256.
- [6] K. Dykema, T. Heister and K. Juschenko, Finitely presented groups related to Kaplansky's direct finiteness conjecture, Exp. Math., 24 (2015), 326-338.
- [7] D. R. Farkas and R. L. Snider,  $K_0$  and Noetherian group rings, J. Algebra, 42 (1976), no. 1, 192-198.
- [8] E. Formanek, The zero divisor question for supersolvable groups, Bull. Austral. Math. Soc., 9 (1973), 69-71.
- [9] G.A. Freiman, Foundations of a structural theory of set addition, Translations of Mathematical Monographs, Vol 37. American Mathematical Society, Providence, R. I., 1973.
- [10] G.A. Freiman, Inverse additive number theory, XI. Long arithmetic progressions in sets with small sumsets, Acta Arith., 137 (2009), 325-331.
- [11] R.J. Gardner and P. Gronchi, A Brunn-Minkowski inequality for the integer lattice, Trans. Amer. Math. Soc., 353 (2001), 3995-4024.
- [12] The GAP Group, GAP-Groups, Algorithms, and Programming, version 4.6.4, 2013. (<http://www.gap-system.org>).
- [13] B. Green and T. Tao, Compressions, convex geometry and the Freiman-Bilu theorem, Quarterly J. Math., 57 (2006), 495-504.
- [14] Y.O. Hamidoune, An isoperimetric method in additive theory, J. Algebra, 179 (1996), 622-630.
- [15] Y.O. Hamidoune, Some additive applications of the isoperimetric approach, Ann. Inst. Fourier (Grenoble), 58 (2008), 2007-2036.
- [16] Y.O. Hamidoune, The isoperimetric method, In: Combinatorial number theory and additive group theory, Adv. Courses Math. CRM Barcelona, Birkhäuser Verlag, Basel, 2009, 241-252.
- [17] Y. O. Hamidoune, A. S. Lladó and O. Serra, On subsets with small product in torsion-free groups, Combinatorica, 18 (1998), no. 4, 529-540.

- 
- [18] G. Higman, Units in group rings, D. Phil. Thesis, Oxford Univ., (1940).
- [19] G. Higman, The units of group rings, Proc. London Math. Soc., 46 (1940), 231-248.
- [20] I. Kaplansky, "Problems in the theory of rings" revisited, Amer. Math. Monthly, 77 (1970), 445-454.
- [21] K. J. Böröczky, P. P. Pálffy and O. Serra, On the cardinality of sumsets in torsion-free groups, Bull. London Math. Soc., 44 (2012), 1034-1041.
- [22] J. H. B. Kemperman, On complexes in a semigroup, Indag. Math., 18 (1956), 247-254.
- [23] P. H. Kropholler, P. A. Linnell and J. A. Moody, Applications of a new  $K$ -theoretic theorem to soluble group rings, Proc. Amer. Math. Soc., 104 (1988), no. 3, 675-684.
- [24] J. Lewin, A note on zero divisors in group-rings, Proc. Amer. Math. Soc., 31 (1972), 357-359.
- [25] J. Lewin and T. Lewin, An embedding of the group algebra of a torsion-free one relator group in a field, J. Algebra, 52 (1978), 39-74.
- [26] M. Matolcsi and I.Z. Ruzsa, Sumsets and the convex hull, In: D. Chudnovsky, G. Chudnovsky (eds.), Additive Number Theory: Festschrift In Honor of the Sixtieth Birthday of Melvyn B. Nathanson, Springer-Verlag, (2010), 221-227.
- [27] D. S. Passman, Infinite group rings, Dekker, New York, 1971.
- [28] S. D. Promislow, A simple example of a torsion-free non unique product group, Bull. London Math. Soc. 20 (1998), 302-304.
- [29] E. Rips and Y. Segev, Torsion-free group without unique product property, J. Algebra, 108 (1987), 116-126.
- [30] D. J. S. Robinson, A course in the theory of groups, 2nd ed., Springer, 1996.
- [31] W. Rudin and H. Schneider, Idempotents in group rings, Duke Math. J., 31 (1964), 585-602.
- [32] I.Z. Ruzsa, Sum of sets in several dimensions, Combinatorica, 14 (1994), no. 4, 485-490.
- [33] P. Schweitzer, On zero divisors with small support in group rings of torsion-free groups, J. Group Theory, 16 (2013), no. 5, 667-693.
- [34] R. L. Snider, The zero divisor conjecture for some solvable groups, Pacific J. Math., 90 (1980), no. 1, 191-196.
- [35] A. Strojnowsky, A note on u.p. groups, Comm. Algebra, 8 (1980), no. 3, 231-234.

**Alireza Abdollahi**

Department of Mathematics, University of Isfahan, Isfahan 81746-73441, Iran,

and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O.Box 19395-5746, Tehran, Iran

E-mail address: [a.abdollahi@math.ui.ac.ir](mailto:a.abdollahi@math.ui.ac.ir)

**Fatemeh Jafari**

Department of Mathematics, University of Isfahan, Isfahan 81746-73441, Iran.

E-mail address: [f\\_jafari@sci.ui.ac.ir](mailto:f_jafari@sci.ui.ac.ir)