

Corrigendum to "Coupling the SO(2) supergravity through dimensional reduction" [Phys. Lett. B 96 (1-2) (1980) 89-93]

A. Chamseddine ^{1,2}, H. Nicolai ³

¹Physics Department, American University of Beirut, Lebanon

²I.H.E.S. F-91440 Bures-sur-Yvette, France

³Max-Planck-Institut fuer Gravitationsphysik (Albert-Einstein-Institute) Muehlenberg 1, D-14476, Potsdam Germany

Emails chams@aub.edu.lb and hermann.nicolai@aei.mpg.de

Due to renewed interest in $D = 5$ supergravity and its quartic fermionic terms, fuelled by very recent work on supersymmetric quantum cosmology [1], we would like to correct an error in eq.(4) of the above paper [2] (to which we also refer for our notations and further explanations¹). In 1.5 order formalism [3] the correct Lagrangian of pure $D = 5$ supergravity reads

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4}eR(\omega) - \frac{i}{2}e \left[\bar{\psi}_M \Gamma^{MNP} \vec{D}_N \left(\frac{\omega + \hat{\omega}}{2} \right) \psi_P - \bar{\psi}_M \overleftarrow{D}_N \left(\frac{\omega + \hat{\omega}}{2} \right) \Gamma^{MNP} \psi_P \right] \quad (0.1) \\ & - \frac{1}{4}e F^{MN} F_{MN} - \frac{\sqrt{3}}{8} i e \bar{\psi}_M X^{MNPQ} \psi_N (F_{PQ} + \hat{F}_{PQ}) - \frac{1}{6\sqrt{3}} \epsilon^{MNPQR} F_{MN} F_{PQ} A_R \end{aligned}$$

with the 4-component Dirac vector spinor ψ_M . The $(3\omega - \hat{\omega})/2$ in eq.(4) of [2] must thus be replaced by $(\omega + \hat{\omega})/2$. The terms with $\hat{\omega}$ and \hat{F} account for all higher order fermionic terms in 1.5 order formalism, as they do in $D = 11$ supergravity [4]. Here the spin connection ω is determined by its equation of motion

$$\omega_{MAB} = \overset{\circ}{\omega}_{MAB}(e) + \kappa_{MAB} \quad (0.2)$$

with the contorsion tensor

$$\begin{aligned} \kappa_{MAB} = & -\frac{i}{2} \bar{\psi}_Q \Gamma^{QR}{}_{MAB} \psi_R + \frac{i}{2} (\bar{\psi}_M \Gamma_B \psi_A - \bar{\psi}_A \Gamma_B \psi_M) + \\ & + \frac{i}{2} (\bar{\psi}_B \Gamma_M \psi_A - \bar{\psi}_A \Gamma_M \psi_B) - \frac{i}{2} (\bar{\psi}_M \Gamma_A \psi_B - \bar{\psi}_B \Gamma_A \psi_M) \quad (0.3) \end{aligned}$$

The supercovariant spin connection and field strength are

$$\hat{\omega}_{MAB} \equiv \omega_{MAB} + \frac{i}{2} \bar{\psi}_Q \Gamma^{QR}{}_{MAB} \psi_R, \quad \hat{F}_{MN} \equiv F_{MN} + \frac{\sqrt{3}i}{2} (\bar{\psi}_M \psi_N - \bar{\psi}_N \psi_M) \quad (0.4)$$

From (0.1) one obtains the supercovariant Rarita-Schwinger equation (\equiv eq. (14) of [2])

$$\Gamma^{MNP} D_N(\hat{\omega}) \psi_P + \frac{\sqrt{3}}{4} X^{MNPQ} \psi_N \hat{F}_{PQ} = 0 \quad (0.5)$$

The Lagrangian in second order formalism, which was not explicitly written out in [2], is obtained in the usual way by substituting the second order spin connection into the Lagrangian

¹In particular, our metric signature is $(+ - - -)$, and thus $\bar{\psi}_M \equiv \psi_M^\dagger \Gamma^0$. Furthermore, $\Gamma_{\alpha\beta}^{ABCDE} = \epsilon^{ABCDE} \delta_{\alpha\beta}$ with $\epsilon^{01234} = \epsilon_{01234} = 1$.

(0.1), with the result

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4}eR(\overset{\circ}{\omega}(e)) - \frac{i}{2}e \left[\bar{\psi}_M \Gamma^{MNP} \vec{D}_N \left(\overset{\circ}{\omega}(e) \right) \psi_P - \bar{\psi}_M \overleftarrow{D}_N \left(\overset{\circ}{\omega}(e) \right) \Gamma^{MNP} \psi_P \right] \\ & - \frac{1}{4}e F^{MN} F_{MN} - \frac{\sqrt{3}}{4} i e \bar{\psi}_M X^{MNPQ} \psi_N F_{PQ} - \frac{1}{6\sqrt{3}} \epsilon^{MNPQR} F_{MN} F_{PQ} A_R + \mathcal{L}_{\text{quartic}} \end{aligned} \quad (0.6)$$

where

$$\begin{aligned} \mathcal{L}_{\text{quartic}} = & \frac{1}{4}e \left[(\bar{\psi}_M \Gamma^N \psi_N - \bar{\psi}_N \Gamma^N \psi_M) (\bar{\psi}^M \Gamma^P \psi_P - \bar{\psi}_P \Gamma^P \psi^M) \right. \\ & - \frac{1}{4} (\bar{\psi}_M \Gamma_N \psi_P - \bar{\psi}_P \Gamma_N \psi_M) (\bar{\psi}^M \Gamma^N \psi^P - \bar{\psi}^P \Gamma^N \psi^M) \\ & - \frac{1}{2} (\bar{\psi}_M \Gamma_N \psi_P - \bar{\psi}_P \Gamma_N \psi_M) (\bar{\psi}^M \Gamma^P \psi^N - \bar{\psi}^N \Gamma^P \psi^M) \\ & \left. + \bar{\psi}_M \Gamma^{MNPQ} \psi_N \bar{\psi}_P \psi_Q + \frac{3}{2} (\bar{\psi}_M \psi_N - \bar{\psi}_N \psi_M) \bar{\psi}^M \psi^N \right] \end{aligned}$$

This can now be compared to [5] (which, however, uses symplectic Majorana spinors rather than Dirac spinors).

Acknowledgments: We would like to thank T. Damour, P. Spindel and A. Van Proeyen for discussions and correspondence.

References

- [1] T. Damour and P. Spindel, Phys.Rev. **D90** (2014) 103509; Phys.Rev. **D95** (2017) 126011
- [2] A. Chamseddine and H. Nicolai, Phys. Lett. **96B** (1980) 89
- [3] A. Chamseddine and P. West, Nucl. Phys. **B129** (1977) 39;
P.K. Townsend and P. van Nieuwenhuizen, Phys. Lett. **B67** (1977) 439
- [4] E. Cremmer, B. Julia and J. Scherk, Phys. Lett. **76B** (1978) 409
- [5] E. Cremmer, *Supergravities in 5 Dimensions*, published in A. Salam, E. Sezgin (eds.):
Supergravities in diverse dimensions, Cambridge University Press (1980)