

CANONICAL BASES FOR TENSOR PRODUCTS AND SUPER KAZHDAN-LUSZTIG THEORY

HUANCHEN BAO, WEIQIANG WANG, AND HIDEYA WATANABE

ABSTRACT. We generalize a construction in [BW18b] by showing that, for a quantum symmetric pair $(\mathbf{U}, \mathbf{U}^\iota)$ of finite type, the tensor product of a based \mathbf{U}^ι -module and a based \mathbf{U} -module is a based \mathbf{U}^ι -module. This is then used to formulate a Kazhdan-Lusztig theory for an *arbitrary* parabolic BGG category \mathcal{O} of the ortho-symplectic Lie superalgebras, extending a main result in [BW18a].

1. INTRODUCTION

1.1. Following Lusztig [Lu94], we shall refer to (M, \mathbf{B}) , which consists of a \mathbf{U} -module M and its canonical basis \mathbf{B} , as a based \mathbf{U} -module, where \mathbf{U} is a Drinfeld-Jimbo quantum group of finite type. Examples of such based \mathbf{U} -modules include any finite-dimensional simple \mathbf{U} -module or a tensor product of several such simple \mathbf{U} -modules. The canonical basis on a tensor product of several finite-dimensional simple \mathbf{U} -modules was constructed by Lusztig [Lu92], and it has found applications to the Kazhdan-Lusztig theory for general linear Lie superalgebra $\mathfrak{gl}(m|n)$ of type A [Br03, CLW15].

1.2. As a generalization of canonical bases for quantum groups, a theory of canonical basis arising from quantum symmetric pairs (QSP, for short) $(\mathbf{U}, \mathbf{U}^\iota)$ of finite type is systematically developed in [BW18b]. For any finite-dimensional based \mathbf{U} -module (M, \mathbf{B}) , a new bar involution ψ_ι on M was formulated and a ψ_ι -invariant basis \mathbf{B}^ι of M (called an ι -canonical basis for M) was constructed (see [BW18b, Theorem 5.7]), which satisfies some specific properties when expanded with respect to \mathbf{B} ; we shall call (M, \mathbf{B}^ι) a based \mathbf{U}^ι -module.

The first examples of based \mathbf{U}^ι -modules were constructed in [BW18a] for quasi-split QSP $(\mathbf{U}, \mathbf{U}^\iota)$ of type AIII. The ι -canonical basis on a tensor product of \mathbf{U} -modules was used to formulate a Kazhdan-Lusztig theory on the full BGG category for ortho-symplectic Lie superalgebra of type B [BW18a] (for type D see [Bao17]); also see [CLW11] on some parabolic BGG category via a super duality approach.

1.3. This paper is intended to supplement the two earlier papers [BW18a, BW18b] of the first two authors on canonical bases arising from quantum symmetric pairs and applications to super Kazhdan-Lusztig theory; it extends two principal results on ι -canonical basis and super Kazhdan-Lusztig theory therein to full generalities.

1.4. By definition of a QSP $(\mathbf{U}, \mathbf{U}^\iota)$, \mathbf{U}^ι is a coideal subalgebra of \mathbf{U} [Le99]; that is, the comultiplication Δ on \mathbf{U} when restricting to \mathbf{U}^ι satisfies $\Delta : \mathbf{U}^\iota \rightarrow \mathbf{U}^\iota \otimes \mathbf{U}$. Hence $M \otimes N$ is a \mathbf{U}^ι -module for any \mathbf{U}^ι -module M and \mathbf{U} -module N . In the first main theorem (see Theorem 4) we show that, for a based \mathbf{U}^ι -module (M, \mathbf{B}^ι) and a based \mathbf{U} -module (N, \mathbf{B}) , there exists a ψ_ι -invariant basis $\mathbf{B}^\iota \diamond_\iota \mathbf{B}$ on the \mathbf{U}^ι -module $M \otimes N$ such that $(M \otimes N, \mathbf{B}^\iota \diamond_\iota \mathbf{B})$ is a based \mathbf{U}^ι -module. This generalizes a main result in [BW18b] on the ι -canonical basis on a tensor product of \mathbf{U} -modules, since a based \mathbf{U}^ι -module which is not a \mathbf{U} -module exists (cf. [BW18a]).

The construction of the new bar involution ψ_ι on $M \otimes N$ above uses a certain element Θ^ι in a completion of $\mathbf{U}^\iota \otimes \mathbf{U}^+$, which was due to [BW18a] for quasi-split QSP of type AIII/IV and then established in Kolb [Ko17] in full generality with an elegant new proof. We establish the integrality of Θ^ι by using the integrality of the quasi- \mathcal{R} matrix in [Lu94] and the integrality of the quasi- \mathcal{K} matrix in [BW18b]. To construct the ι -canonical basis on $M \otimes N$, we use crucially a partial order, which is different from and simpler than the old one used in [BW18b] even when M is a \mathbf{U} -module; the old partial order does not make much sense in our new setting.

1.5. For the quasi-split QSP $(\mathbf{U}, \mathbf{U}^\iota)$ of type AIII/AIV, the quantum group \mathbf{U} is of type A; we let \mathbb{V} and \mathbb{W} denote the natural representation of \mathbf{U} and its dual. In this case, the ι -canonical basis on a based \mathbf{U} -module was first constructed in [BW18a] when a certain parameter $\kappa = 1$ (also see [Bao17] with parameter $\kappa = 0$). The super Kazhdan-Lusztig theory for the *full* BGG category $\mathcal{O}_{\mathfrak{b}}$ of an ortho-symplectic Lie superalgebra \mathfrak{g} of type B in [BW18a] (for type D see [Bao17]) of integer or half-integer weights was formulated via the ι -canonical basis on a mixed tensor \mathbf{U} -module with m copies of \mathbb{V} and n copies of \mathbb{W} , where the order of the tensor product depends on the choice of a Borel subalgebra \mathfrak{b} in \mathfrak{g} .

As a consequence, a Kazhdan-Lusztig theory for parabolic categories $\mathcal{O}_{\mathfrak{b}}^{\mathfrak{l}}$, where the Levi subalgebra \mathfrak{l} in \mathfrak{g} is a product of Lie algebras of type A, can be formulated and established via the ι -canonical basis on a tensor product \mathbf{U} -module \mathbb{T} of various exterior powers of \mathbb{V} and of \mathbb{W} . Note however not all the parabolic categories of \mathfrak{g} -modules arise in this way; indeed a general Levi subalgebra of \mathfrak{g} is isomorphic to a product of several Lie subalgebras of type A and a Lie subalgebra of type B.

Theorem 4, when specialized for the QSP $(\mathbf{U}, \mathbf{U}^\iota)$ of quasi-split type AIII/AIV, provides an ι -canonical basis for a \mathbf{U}^ι -module on the tensor product of the form $\wedge^a \mathbb{V}_- \otimes \mathbb{T}$. Here \mathbb{T} is a tensor product \mathbf{U} -module of various exterior powers of \mathbb{V} and of \mathbb{W} , while $\wedge^a \mathbb{V}_-$ (for $a > 0$) is a “type B” exterior power, which is a \mathbf{U}^ι -module but not a \mathbf{U} -module. These new ι -canonical bases are used to formulate the super Kazhdan-Lusztig theory for an *arbitrary* parabolic BGG category \mathcal{O} of the ortho-symplectic Lie superalgebras of type B and D; see Theorem 10. The super Kazhdan-Lusztig polynomials $t_{gf}^{\mathfrak{b}}(q)$ admit a positivity property; see Theorem 12.

1.6. This paper is organized as follows. Theorem 4 and its proof are presented in Section 2, and we shall follow notations in [BW18b] throughout Section 2. The formulation of Theorem 10 is given in Section 3; its proof basically follows the proof for the Kazhdan-Lusztig theory for the full category \mathcal{O} in [BW18a, Part 2] once we have

Theorem 4 available to us. We shall follow notations in [BW18a] throughout Section 3. To avoid much repetition, we refer precisely and freely to the two earlier papers [BW18a, BW18b].

Acknowledgements. The research of WW is partially supported by an NSF grant DMS-1702254.

2. TENSOR PRODUCT MODULES AS BASED \mathbf{U}^ι -MODULES

2.1. We shall follow the notations in [BW18b] throughout this section.

Let \mathbf{U} denote a quantum group of finite type over the field $\mathbb{Q}(q)$ associated to a root datum of type (\mathbb{I}, \cdot) , and let Δ denote its comultiplication as in [Lu94]. We denote the bar involution on \mathbf{U} or its based module by ψ .

Let $\mathbf{U}^\iota \subset \mathbf{U}$ be a coideal subalgebra associated to a Satake diagram such that $(\mathbf{U}, \mathbf{U}^\iota)$ forms a quantum symmetric pair [Le99]. Let $\mathcal{A} := \mathbb{Z}[q, q^{-1}]$. Let $\dot{\mathbf{U}}^\iota$ be the modified version of \mathbf{U}^ι and let ${}_{\mathcal{A}}\dot{\mathbf{U}}^\iota$ be its \mathcal{A} -form, respectively; see [BW18b, §3.7]. Let ψ_ι be the bar involution on \mathbf{U}^ι , $\dot{\mathbf{U}}^\iota$ and ${}_{\mathcal{A}}\dot{\mathbf{U}}^\iota$. Let X_ι be the ι -weight lattice [BW18b, (3.3)]. A weight (i.e., X_ι -weight) module of \mathbf{U}^ι can be naturally regarded as a $\dot{\mathbf{U}}^\iota$ -module.

We introduce based \mathbf{U}^ι -modules generalizing [Lu94, §27.1.2]. Let $\mathbf{A} = \mathbb{Q}[[q^{-1}]] \cap \mathbb{Q}(q)$. We write $-\otimes - = -\otimes_{\mathbb{Q}(q)} -$ whenever the base ring is $\mathbb{Q}(q)$.

Definition 1. Let M be a finite-dimensional weight \mathbf{U}^ι -module over $\mathbb{Q}(q)$ with a given $\mathbb{Q}(q)$ -basis \mathbf{B}^ι . The pair (M, \mathbf{B}^ι) is called a based \mathbf{U}^ι -module if the following conditions are satisfied:

- (1) $\mathbf{B}^\iota \cap M_\nu$ is a basis of M_ν , for any $\nu \in X_\iota$;
- (2) The \mathcal{A} -submodule ${}_{\mathcal{A}}M$ generated by \mathbf{B}^ι is stable under ${}_{\mathcal{A}}\dot{\mathbf{U}}^\iota$;
- (3) The \mathbb{Q} -linear involution $\psi_\iota : M \rightarrow M$ defined by $\psi_\iota(q) = q^{-1}$, $\psi_\iota(b) = b$ for all $b \in \mathbf{B}^\iota$ is compatible with the $\dot{\mathbf{U}}^\iota$ -action, i.e., $\psi_\iota(um) = \psi_\iota(u)\psi_\iota(m)$, for all $u \in \dot{\mathbf{U}}^\iota$, $m \in M$;
- (4) Let $L(M)$ be the \mathbf{A} -submodule of M generated by \mathbf{B}^ι . Then the image of \mathbf{B}^ι in $L(M)/q^{-1}L(M)$ forms a \mathbb{Q} -basis in $L(M)/q^{-1}L(M)$.

We shall denote by $\mathcal{L}(M)$ the $\mathbb{Z}[q^{-1}]$ -span of \mathbf{B}^ι ; then \mathbf{B}^ι forms a $\mathbb{Z}[q^{-1}]$ -basis for $\mathcal{L}(M)$. (There are similar constructions for a based \mathbf{U} -module in similar notations.)

2.2. Let $\Upsilon = \sum_{\mu} \Upsilon_{\mu}$ (with $\Upsilon_0 = 1$ and $\Upsilon_{\mu} \in \mathbf{U}_{\mu}^+$) be the intertwiner (also called quasi- \mathcal{K} matrix) of the quantum symmetric pair $(\mathbf{U}, \mathbf{U}^\iota)$ introduced in [BW18a, Theorem 2.10]; for full generality see [BK19, Theorem 6.10], [BW18b, Theorem 4.8, Remark 4.9]. It follows from [BW18b, Theorem 5.7] (also cf. [BW18a, Theorem 4.25]) that a based \mathbf{U} -module (M, \mathbf{B}) becomes a based \mathbf{U}^ι -module with a new basis \mathbf{B}^ι (which is uni-triangular relative to \mathbf{B}) with respect to the involution $\psi_\iota := \Upsilon \circ \psi$.

Let $\widehat{\mathbf{U} \otimes \mathbf{U}}$ be the completion of the $\mathbb{Q}(q)$ -vector space $\mathbf{U} \otimes \mathbf{U}$ with respect to the descending sequence of subspaces

$$\mathbf{U} \otimes \mathbf{U}^{-\mathbf{U}^0} \left(\sum_{\text{ht}(\mu) \geq N} \mathbf{U}_{\mu}^+ \right) + \mathbf{U}^+ \mathbf{U}^0 \left(\sum_{\text{ht}(\mu) \geq N} \mathbf{U}_{\mu}^- \right) \otimes \mathbf{U}, \text{ for } N \geq 1, \mu \in \mathbb{Z}\mathbb{I}.$$

We have the obvious embedding of $\mathbf{U} \otimes \mathbf{U}$ into $\widehat{\mathbf{U} \otimes \mathbf{U}}$. By continuity the $\mathbb{Q}(q)$ -algebra structure on $\mathbf{U} \otimes \mathbf{U}$ extends to a $\mathbb{Q}(q)$ -algebra structure on $\widehat{\mathbf{U} \otimes \mathbf{U}}$. We know the quasi- \mathcal{R} matrix Θ lies in $\widehat{\mathbf{U} \otimes \mathbf{U}}$ by [Lu94, Theorem 4.1.2]. It follows from [BW18b, Theorem 4.8] and [BK19, Theorem 6.10] that $\Upsilon^{-1} \otimes \text{id}$ and $\Delta(\Upsilon)$ are both in $\widehat{\mathbf{U} \otimes \mathbf{U}}$.

We define

$$(2.1) \quad \Theta^i = \Delta(\Upsilon) \cdot \Theta \cdot (\Upsilon^{-1} \otimes \text{id}) \in \widehat{\mathbf{U} \otimes \mathbf{U}}.$$

We can write

$$(2.2) \quad \Theta^i = \sum_{\mu \in \mathbb{N}\mathbb{I}} \Theta_\mu^i, \quad \text{where } \Theta_\mu^i \in \mathbf{U} \otimes \mathbf{U}_\mu^+.$$

The following result first appeared in [BW18a, Proposition 3.5] for the quantum symmetric pairs of (quasi-split) type AIII/AIV.

Lemma 2. [Ko17, Proposition 3.10] *We have $\Theta_\mu^i \in \mathbf{U}^i \otimes \mathbf{U}_\mu^+$, for all μ . (The element Θ_μ^i is denoted by R_μ^θ in [Ko17].)*

Another basic ingredient which we shall need is the integrality property of Θ^i .

Lemma 3. *We have $\Theta_\mu^i \in {}_A\mathbf{U} \otimes_A {}_A\mathbf{U}_\mu^+$, for all μ .*

Proof. By a result of Lusztig [Lu94, 24.1.6], we have $\Theta = \sum_{\nu \in \mathbb{N}\mathbb{I}} \Theta_\nu$ is integral, i.e., $\Theta_\nu \in {}_A\mathbf{U}_\nu^- \otimes_A {}_A\mathbf{U}_\nu^+$. By [BW18b, Theorem 5.3] we have $\Upsilon = \sum_{\mu \in \mathbb{N}\mathbb{I}} \Upsilon_\mu$ is integral, i.e., $\Upsilon_\mu \in {}_A\mathbf{U}_\mu^+$ for each μ ; it follows that $\Upsilon^{-1} = \psi(\Upsilon)$ is integral too thanks to [BW18b, Corollary 4.11]. It is well known that the comultiplication Δ preserves the \mathcal{A} -form, i.e., $\Delta({}_A\mathbf{U}) \subset {}_A\mathbf{U} \otimes_A {}_A\mathbf{U}$. The lemma follows now by the definition of Θ^i in (2.1). \square

2.3. Define a partial order $<$ on X by setting $\mu' < \mu$ if $\mu' - \mu \in \mathbb{N}\mathbb{I}$. Denote by $|b| = \mu$ if an element b in a \mathbf{U} -module is of weight μ . Now we are ready to prove the first main result of this paper.

Theorem 4. *Let (M, \mathbf{B}^i) be a based \mathbf{U}^i -module and (N, \mathbf{B}) be a based \mathbf{U} -module.*

- (1) *For $b_1 \in \mathbf{B}^i, b_2 \in \mathbf{B}$, there exists a unique element $b_1 \diamond_i b_2$ which is ψ_i -invariant such that $b_1 \diamond_i b_2 \in b_1 \otimes b_2 + q^{-1}\mathbb{Z}[q^{-1}]\mathbf{B}^i \otimes \mathbf{B}$.*
- (2) *We have $b_1 \diamond_i b_2 \in b_1 \otimes b_2 + \sum_{(b'_1, b'_2) \in \mathbf{B}^i \times \mathbf{B}, |b'_2| < |b_2|} q^{-1}\mathbb{Z}[q^{-1}]b'_1 \otimes b'_2$.*
- (3) *$\mathbf{B}^i \diamond_i \mathbf{B} := \{b_1 \diamond_i b_2 \mid b_1 \in \mathbf{B}^i, b_2 \in \mathbf{B}\}$ forms a $\mathbb{Q}(q)$ -basis for $M \otimes N$, an \mathcal{A} -basis for ${}_A M \otimes_A {}_A N$, and a $\mathbb{Z}[q^{-1}]$ -basis for $\mathcal{L}(M) \otimes_{\mathbb{Z}[q^{-1}]} \mathcal{L}(N)$. (This is called the i -canonical basis for $M \otimes N$.)*
- (4) *$(M \otimes N, \mathbf{B}^i \diamond_i \mathbf{B})$ is a based \mathbf{U}^i -module.*

Proof. It follows by Lemma 2 that the element Θ^i gives rise to a well-defined operator on the tensor product $M \otimes N$. Following [BW18a, (3.17)], we define a new bar involution on $M \otimes N$ (still denoted by ψ_i) by letting

$$\psi_i := \Theta^i \circ (\psi_i \otimes \psi) : M \otimes N \longrightarrow M \otimes N.$$

Recall from [Lu94] that $\Delta(E_i) = E_i \otimes 1 + \tilde{K}_i \otimes E_i$. It follows that

$$\Delta(\Upsilon) \in \Upsilon \otimes 1 + \sum_{0 \neq \mu \in \mathbb{N}^I} \mathbf{U} \otimes \mathbf{U}_\mu^+.$$

Recalling (2.2), we have

$$(2.3) \quad \Theta_0^i = (\Upsilon \otimes 1) \cdot (1 \otimes 1) \cdot (\Upsilon^{-1} \otimes 1) = 1 \otimes 1.$$

Let $b_1 \in \mathbf{B}^i$ and $b_2 \in \mathbf{B}$. By (2.3) and Lemma 3, we have

$$(2.4) \quad \psi_i(b_1 \otimes b_2) \in b_1 \otimes b_2 + \sum_{\substack{(b'_1, b'_2) \in \mathbf{B}^i \times \mathbf{B} \\ |b'_2| < |b_2|}} \mathcal{A} b'_1 \otimes b'_2.$$

Applying [Lu94, Lemma 24.2.1], there exists a ψ_i -invariant element $(b_1 \otimes b_2)^i \in M \otimes N$ such that

$$b_1 \diamond_i b_2 \in b_1 \otimes b_2 + \sum_{\substack{(b'_1, b'_2) \in \mathbf{B}^i \times \mathbf{B} \\ |b'_2| < |b_2|}} q^{-1} \mathbb{Z}[q^{-1}] b'_1 \otimes b'_2.$$

This proves (2), and Part (3) follows immediately.

A by now standard argument shows the uniqueness of $b_1 \diamond_i b_2$ as stated in (1); note a weaker condition than (2) is used in (1).

It remains to see that $(M, \mathbf{B}^i \diamond_i \mathbf{B})$ is a based \mathbf{U}^i -module. The item (3) in the definition of a based \mathbf{U}^i -module is proved in the same way as for [BW18a, Proposition 3.13], while the remaining items are clear.

This completes the proof. \square

Remark 5. (1) An elementary but key new ingredient in Theorem 4 above is the use of a (coarser) partial order $<$, which is different from the partial order $<_i$ used in [BW18b, (5.2)].

(2) Assume that M is a based \mathbf{U} -module. Then the i -canonical basis for $M \otimes N$ in Theorem 4 coincides with the one in [BW18b, Theorem 5.7], thanks to the uniqueness in Theorem 4(1).

Remark 6. Theorem 4 would be valid whenever we can establish the (weaker) integrality of Θ^i acting on $M \otimes N$. This might occur when we consider more general parameters for \mathbf{U}^i than [BW18b] or when we consider quantum symmetric pairs of Kac-Moody type in a forthcoming work of the first two authors.

3. APPLICATIONS TO SUPER KAZHDAN-LUSZTIG THEORY

3.1. In this section, we shall apply Theorem 4 to formulate and establish the (super) Kazhdan-Lusztig theory for an arbitrary parabolic category \mathcal{O} of modules of integer or half-integer weights for ortho-symplectic Lie superalgebras, generalizing [BW18a, Part 2] (also see [Bao17]). We shall present only the details on an arbitrary parabolic category \mathcal{O} consisting of modules of integer weights for Lie superalgebra $\mathfrak{osp}(2m+1|2n)$.

3.2. All relevant notations throughout this section shall be consistent with [BW18a, Part 2]. In particular, we use a comultiplication for \mathbf{U} different from [Lu94]; this leads to a version of the intertwiner $\Upsilon = \sum_{\mu} \Upsilon_{\mu}$ with $\Upsilon_{\mu} \in \widehat{\mathbf{U}}^{-}$ (compare with §2.2), and a version of Theorem 4 in which the opposite partial order and the lattice $\mathbb{Z}[q]$ are used. To further match notations with [BW18a] in this section, we denote the \mathcal{A} -form of any based \mathbf{U} or \mathbf{U}^{ι} -modules M , as $M_{\mathcal{A}}$ (instead of ${}_{\mathcal{A}}M$).

We consider the infinite-rank quantum symmetric pair $(\mathbf{U}, \mathbf{U}^{\iota})$ as defined in [BW18a, Section 8] (where the parameter is chosen to be $\kappa = 1$ in the notation of [BW18b]). It is a direct limit of quantum symmetric pairs of type AIII, $(\mathbf{U}(\mathfrak{sl}_N), \mathbf{U}^{\iota}(\mathfrak{sl}_N))$, for N even. We denote by \mathbb{V} the natural representation of \mathbf{U} , and by \mathbb{W} the restricted dual of \mathbb{V} .

Associated to any given $0^m 1^n$ -sequence $\mathbf{b} = (b_1, \dots, b_{m+n})$ starting with 0, we have a fundamental system of $\mathfrak{osp}(2m+1|2n)$, denoted by $\Pi_{\mathbf{b}} = \{-\epsilon_1^{b_1}, \epsilon_i^{b_i} - \epsilon_{i+1}^{b_{i+1}} \mid 1 \leq i \leq m+n-1\}$; here $\epsilon_i^0 = \epsilon_x$ for some $1 \leq x \leq m$ and $\epsilon_j^1 = \epsilon_{\bar{y}}$ for some $1 \leq y \leq n$ so that $\{\epsilon_i^{b_i} \mid 1 \leq i \leq m+n\}$ form a permutation of $\{\epsilon_a, \epsilon_b \mid 1 \leq a \leq m, 1 \leq b \leq n\}$.

3.3. Let W_{B_s} and $W_{A_{s-1}}$ be the Weyl group of type B_s and type A_{s-1} with unit e , respectively. We denote their corresponding Hecke algebras by \mathcal{H}_{B_s} and $\mathcal{H}_{A_{s-1}}$, with Kazhdan-Lusztig bases by $\{\underline{H}_w \mid w \in W_{B_s}\}$ and $\{\underline{H}_w \mid w \in W_{A_{s-1}}\}$, respectively. Both algebras act naturally on the right on $\mathbb{V}^{\otimes s}$ and $\mathbb{W}^{\otimes s}$; cf. [BW18a, Section 5]. We define

$$\begin{aligned} \wedge^s \mathbb{V}_- &= \mathbb{V}^{\otimes s} / \sum_{e \neq w \in W_{B_s}} \mathbb{V}^{\otimes s} \cdot \underline{H}_w, \\ \wedge^s \mathbb{V} &= \mathbb{V}^{\otimes s} / \sum_{e \neq w \in W_{A_{s-1}}} \mathbb{V}^{\otimes s} \cdot \underline{H}_w. \end{aligned}$$

We similarly define $\wedge^s \mathbb{W}_-$ and $\wedge^s \mathbb{W}$. Note $\wedge^s \mathbb{V}_-, \wedge^s \mathbb{V}, \wedge^s \mathbb{W}_-$ and $\wedge^s \mathbb{W}$ are all based \mathbf{U}^{ι} -modules by [BW18a, Theorem 5.8]. We shall denote

$$\mathbb{V}^c := \begin{cases} \mathbb{V} & \text{if } c = 0, \\ \mathbb{W} & \text{if } c = 1. \end{cases}$$

The following corollary is a direct consequence of Theorem 4.

Corollary 7. *Let $c_1, \dots, c_k \in \{0, 1\}$ and $a_0, a_1, \dots, a_k \in \mathbb{N}$. Then (a suitable completion of) the tensor product*

$$\mathbb{T}^{\mathbf{b}, \mathbf{l}} = \wedge^{a_0} \mathbb{V}_- \otimes \wedge^{a_1} \mathbb{V}^{c_1} \otimes \dots \otimes \wedge^{a_k} \mathbb{V}^{c_k}$$

is a based \mathbf{U}^{ι} -module.

The completion above arises since we deal with quantum symmetric pairs of infinite rank, and it is a straightforward generalization of the B -completion studied in [BW18a, Section 9]. Note that the ι -canonical basis lives in $\mathbb{T}^{\mathbf{b}, \mathbf{l}}$ (instead of its completion) by Theorem 12 below.

3.4. Associated to the fundamental system $\Pi_{\mathfrak{b}}$ are the set of positive roots $\Phi_{\mathfrak{b}}^+$ and the Borel subalgebra $\mathfrak{b}_{\mathfrak{b}}$ of $\mathfrak{osp}(2m+1|2n)$. Let $\Pi_{\mathfrak{l}} \subset \Pi_{\mathfrak{b}}$ be a subset of even simple roots. We introduce the corresponding Levi subalgebra \mathfrak{l} and parabolic subalgebra \mathfrak{p} of $\mathfrak{osp}(2m+1|2n)$:

$$\mathfrak{l} = \mathfrak{h}_{m|n} \bigoplus \bigoplus_{\alpha \in \mathbb{Z}\Pi_{\mathfrak{l}} \cap \Phi_{\mathfrak{b}}} \mathfrak{osp}(2m+1|2n)_{\alpha}, \quad \mathfrak{p} = \mathfrak{l} + \mathfrak{b}_{\mathfrak{b}}.$$

Recall [BW18a, §7] the weight lattice $X(m|n) = \sum_{i=1}^m \mathbb{Z}\epsilon_i + \sum_{j=1}^n \mathbb{Z}\epsilon_{\bar{j}}$. We denote

$$X_{\mathfrak{b}}^{\mathfrak{l},+} = \{\lambda \in X(m|n) \mid (\lambda|\alpha) \geq 0, \forall \alpha \in \Pi_{\mathfrak{l}}\}.$$

Let $L_0(\lambda)$ be the irreducible \mathfrak{l} -module with highest weight λ , which is extended trivially to a \mathfrak{p} -module. We form the parabolic Verma module

$$M_{\mathfrak{b}}^{\mathfrak{l}}(\lambda) := \text{Ind}_{\mathfrak{p}}^{\mathfrak{osp}(2m+1|2n)} L_0(\lambda).$$

Definition 8. Let $\mathcal{O}_{\mathfrak{b}}^{\mathfrak{l}}$ be the category of $\mathfrak{osp}(2m+1|2n)$ -modules M such that

- (i) M admits a weight space decomposition $M = \bigoplus_{\mu \in X(m|n)} M_{\mu}$, and $\dim M_{\mu} < \infty$;
- (ii) M decomposes over \mathfrak{l} into a direct sum of $L_{\mathfrak{l}}(\lambda)$ for some $\lambda \in X_{\mathfrak{b}}^{\mathfrak{l},+}$;
- (iii) there exist finitely many weights ${}^1\lambda, {}^2\lambda, \dots, {}^k\lambda \in X_{\mathfrak{b}}^{\mathfrak{l},+}$ (depending on M) such that if μ is a weight in M , then $\mu \in {}^i\lambda - \sum_{\alpha \in \Pi_{\mathfrak{b}}} \mathbb{N}\alpha$, for some i .

The morphisms in $\mathcal{O}_{\mathfrak{b}}^{\mathfrak{l}}$ are all (not necessarily even) homomorphisms of $\mathfrak{osp}(2m+1|2n)$ -modules.

For $\lambda \in X_{\mathfrak{b}}^{\mathfrak{l},+}$, we shall denote by $L_{\mathfrak{b}}^{\mathfrak{l}}(\lambda)$ the simple quotient of the parabolic Verma module $M_{\mathfrak{b}}^{\mathfrak{l}}(\lambda)$ in $\mathcal{O}_{\mathfrak{b}}^{\mathfrak{l}}$ with highest weight λ . Following [BW18a, Definition 7.4], we can define the tilting modules $T_{\mathfrak{b}}^{\mathfrak{l}}(\lambda)$ in $\mathcal{O}_{\mathfrak{b}}^{\mathfrak{l}}$, for $\lambda \in X_{\mathfrak{b}}^{\mathfrak{l},+}$. We denote by $\mathcal{O}_{\mathfrak{b}}^{\mathfrak{l},\Delta}$ the full subcategory of $\mathcal{O}_{\mathfrak{b}}^{\mathfrak{l}}$ generated by all modules possessing finite parabolic Verma flags.

3.5. Recall the bijection $X(m|n) \leftrightarrow I^{m+n}$ [BW18a, §8.4], where an element $f \in I^{m+n}$ is understood as a ρ -shifted weight. We consider the restriction $X_{\mathfrak{b}}^{\mathfrak{l},+} \leftrightarrow I_{\mathfrak{l},+}^{m+n}$, where the index set $I_{\mathfrak{l},+}^{m+n}$ is defined as the image under the bijection.

Let $W_{\mathfrak{l}}$ be the Weyl group of \mathfrak{l} with the corresponding Hecke algebra $\mathcal{H}_{\mathfrak{l}}$. Recall that $\Pi_{\mathfrak{l}} \subset \Pi_{\mathfrak{b}}$ is a subset of even simple roots. Hence we have the natural right action of $\mathcal{H}_{\mathfrak{l}}$ on the \mathcal{A} -module $\mathbb{T}_{\mathcal{A}}^{\mathfrak{b}} := \mathbb{V}_{\mathcal{A}}^{b_1} \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} \mathbb{V}_{\mathcal{A}}^{b_{m+n}}$ with a standard basis $M_f^{\mathfrak{b}} \in \mathbb{T}_{\mathcal{A}}^{\mathfrak{b}}$, for $f \in I_{\mathfrak{l},+}^{m+n}$; cf. [BW18a, §8.2]. We define

$$\mathbb{T}_{\mathcal{A}}^{\mathfrak{b},\mathfrak{l}} = \mathbb{T}_{\mathcal{A}}^{\mathfrak{b}} / \sum_{e \neq w \in W_{\mathfrak{l}}} \mathbb{T}_{\mathcal{A}}^{\mathfrak{b}} \cdot \underline{H}_w.$$

The quotient space is an \mathcal{A} -form $\mathbb{T}_{\mathcal{A}}^{\mathfrak{b},\mathfrak{l}}$ of the $\mathbb{Q}(q)$ -space $\mathbb{T}^{\mathfrak{b},\mathfrak{l}}$ appearing in Corollary 7:

$$(3.1) \quad \mathbb{T}_{\mathcal{A}}^{\mathfrak{b},\mathfrak{l}} = \wedge^{a_0} \mathbb{V}_{-, \mathcal{A}} \otimes_{\mathcal{A}} \wedge^{a_1} \mathbb{V}_{\mathcal{A}}^{c_1} \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} \wedge^{a_k} \mathbb{V}_{\mathcal{A}}^{c_k}, \quad \text{for } c_i \in \{0, 1\}, \quad a_i \in \mathbb{N},$$

where c_i and a_i are determined as follows. Let W' denote a subgroup of the Weyl group of $\mathfrak{osp}(2m+1|2n)$, $W' = W_{B_m} \times S_n = \langle s_0, s_1, \dots, s_{m-1}, s_{m+1}, \dots, s_{m+n-1} \rangle$, where $s_i = s_{\alpha_i}$, and $\alpha_0 = -\epsilon_1^{b_0}$, $\alpha_i = \epsilon_i^{b_i} - \epsilon_{i+1}^{b_{i+1}}$ for $1 \leq i \leq m+n-1$. Then, $W_{\mathfrak{l}}$ is the

parabolic subgroup of W' generated by $\{s_i \mid \alpha_i \in \Pi_I\}$. Let us write $\{0, 1, \dots, m+n\} \setminus \{i \mid \alpha_i \in \Pi_I\} = \{j_1 < j_2 < \dots < j_{k+1}\}$. Then, $a_i = j_{i+1} - j_i$ and $c_{i+1} = b_{j_i}$, where it is understood that $j_0 = 0$.

For any standard basis element $M_f^{\mathbf{b}} \in \mathbb{T}_{\mathcal{A}}^{\mathbf{b}}$ with $f \in I_{l,+}^{m+n}$, we denote by $M_f^{\mathbf{b},l}$ its image in $\mathbb{T}_{\mathcal{A}}^{\mathbf{b},l}$. Then $\{M_f^{\mathbf{b},l} \mid f \in I_{l,+}^{m+n}\}$ forms an \mathcal{A} -basis of $\mathbb{T}_{\mathcal{A}}^{\mathbf{b},l}$. Let

$$\mathbb{T}_{\mathbb{Z}}^{\mathbf{b},l} = \mathbb{T}_{\mathcal{A}}^{\mathbf{b},l} \otimes_{\mathcal{A}} \mathbb{Z}$$

be the specialization of $\mathbb{T}_{\mathcal{A}}^{\mathbf{b},l}$ at $q = 1$. Let $\widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b},l}$ be the B -completion of $\mathbb{T}_{\mathbb{Z}}^{\mathbf{b},l}$ following [BW18a, Section 9]. It follows from Corollary 7 the space $\widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b},l}$ admits the ι -canonical basis $\{T_f^{\mathbf{b},l} \mid f \in I_{l,+}^{m+n}\}$. We can similarly define the dual ι -canonical basis $\{L_f^{\mathbf{b},l} \mid f \in I_{l,+}^{m+n}\}$ of $\widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b},l}$ following [BW18a, Theorem 9.9].

3.6. We denote by $[\mathcal{O}_{\mathbf{b}}^{l,\Delta}]$ the Grothendieck group of the category $\mathcal{O}_{\mathbf{b}}^{l,\Delta}$. We have the following isomorphism of \mathbb{Z} -modules:

$$\begin{aligned} \Psi : [\mathcal{O}_{\mathbf{b}}^{l,\Delta}] &\longrightarrow \mathbb{T}_{\mathbb{Z}}^{\mathbf{b},l} \\ M_{\mathbf{b}}^l(\lambda) &\mapsto M_{f_{\lambda}^{\mathbf{b}}}^{\mathbf{b},l}(1), \quad \text{for } \lambda \in X_{\mathbf{b}}^{l,+}. \end{aligned}$$

We define $[[\mathcal{O}_{\mathbf{b}}^{l,\Delta}]]$ as the completion of $[\mathcal{O}_{\mathbf{b}}^{l,\Delta}]$ such that the extension of Ψ ,

$$\Psi : [[\mathcal{O}_{\mathbf{b}}^{l,\Delta}]] \longrightarrow \widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b},l},$$

is an isomorphism of \mathbb{Z} -modules.

The following proposition is a reformulation of the Kazhdan-Lusztig theory for the parabolic category \mathcal{O} of the Lie algebra $\mathfrak{so}(2m+1)$ (theorems of Brylinski-Kashiwara, Beilinson-Bernstein).

Proposition 9. *Let $\mathbf{b} = (0^m)$ (that is $n = 0$). The isomorphism $\Psi : [[\mathcal{O}_{\mathbf{b}}^{l,\Delta}]] \longrightarrow \mathbb{T}_{\mathbb{Z}}^{\mathbf{b},l}$ sends*

$$\Psi([L_{\mathbf{b}}^l(\lambda)]) = L_{f_{\lambda}^{\mathbf{b}}}^{\mathbf{b},l}(1), \quad \Psi([T_{\mathbf{b}}^l(\lambda)]) = T_{f_{\lambda}^{\mathbf{b}}}^{\mathbf{b},l}(1), \quad \text{for } \lambda \in X_{\mathbf{b}}^{l,+}.$$

Note $\widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b},l} = \mathbb{T}_{\mathbb{Z}}^{\mathbf{b},l}$ in this case, i.e. no completion is needed.

Proof. Thanks to [BW18a, Theorem 5.8], the ι -canonical basis on $\mathbb{T}^{\mathbf{b}}$ can be identified with the Kazhdan-Lusztig basis (of type B) on $\mathbb{T}^{\mathbf{b}}$. Note by [BW18a, Theorem 5.4] that $\mathbb{S}^l := \sum_{e \neq w \in W_l} \mathbb{T}^{\mathbf{b}} \cdot \underline{H}_w$ is a \mathbf{U}^l -submodule of $\mathbb{T}^{\mathbf{b}}$, and it is actually a based \mathbf{U}^l -submodule of $\mathbb{T}^{\mathbf{b}}$ with its Kazhdan-Lusztig basis. Therefore the ι -canonical basis on $\mathbb{T}_{\mathcal{A}}^{\mathbf{b},l}$ in Theorem 4 can be identified with the basis in the based quotient $\mathbb{T}^{\mathbf{b}}/\mathbb{S}^l$, which is exactly the parabolic Kazhdan-Lusztig basis. The proposition follows now from the classical Kazhdan-Lusztig theory (cf. [BW18a]). \square

Now we can formulate the super Kazhdan-Lusztig theory for $\mathcal{O}_{\mathbf{b}}^l$.

Theorem 10. *The isomorphism $\Psi : [[\mathcal{O}_{\mathbf{b}}^{l,\Delta}]] \longrightarrow \widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b},l}$ sends*

$$\Psi([L_{\mathbf{b}}^l(\lambda)]) = L_{f_{\lambda}^{\mathbf{b}}}^{\mathbf{b},l}(1), \quad \Psi([T_{\mathbf{b}}^l(\lambda)]) = T_{f_{\lambda}^{\mathbf{b}}}^{\mathbf{b},l}(1), \quad \text{for } \lambda \in X_{\mathbf{b}}^{l,+}.$$

Proof. Let us briefly explain the idea of the proof from [BW18a]. The crucial new ingredient of this paper (cf. Remark 11 below) is the existence of the ι -canonical basis and dual ι -canonical basis on $\widehat{\mathbb{T}}^{\mathbf{b},\mathfrak{l}}$ thanks to Theorem 4. Here the dual ι -canonical basis refers to a version of canonical basis where the lattice $\mathbb{Z}[q]$ is replaced by $\mathbb{Z}[q^{-1}]$; see [BW18a].

We have already established the version of the theorem for the full category \mathcal{O} of the Lie superalgebra $\mathfrak{osp}(2m+1|2n)$ in [BW18a, Theorem 11.13]. We have the following commutative diagram of \mathbb{Z} -modules:

$$\begin{array}{ccc} [[\mathcal{O}_{\mathbf{b}}^{\mathfrak{l},\Delta}]] & \longrightarrow & \widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b},\mathfrak{l}} \\ \downarrow & & \downarrow \\ [[\mathcal{O}_{\mathbf{b}}^{\Delta}]] & \longrightarrow & \widehat{\mathbb{T}}_{\mathbb{Z}}^{\mathbf{b}} \end{array}$$

(Note that the vertical arrow on the right is not a based embedding of \mathbf{U}^{ι} -modules.) Then the theorem follows from comparison of characters entirely similar to [BW18a, §11.2]. Note that this comparison uses only the classical Kazhdan-Lusztig theory. \square

Remark 11. In the case of the full category \mathcal{O} (i.e., \mathfrak{l} is the Cartan subalgebra), the theorem goes back to [BW18a, Theorem 11.13]. Following [BW18a, Remark 11.16], the Kazhdan-Lusztig theory for the parabolic category $\mathcal{O}_{\mathbf{b}}^{\mathfrak{l}}$ with $\alpha_0 \neq \Pi_{\mathfrak{l}}$ was a direct consequence of [BW18a, Theorem 11.13], via the ι -canonical basis in [BW18a, Theorem 4.25] in the ${}_{\mathcal{A}}\mathbf{U}$ -module $\mathbb{T}_{\mathcal{A}}^{\mathbf{b},\mathfrak{l}}$ in (3.1) with $a_0 = 0$.

When $a_0 > 0$ (which corresponds to the condition $\alpha_0 \in \Pi_{\mathfrak{l}}$ on the Levi \mathfrak{l}), the space $\mathbb{T}_{\mathcal{A}}^{\mathbf{b},\mathfrak{l}}$ in (3.1) is a ${}_{\mathcal{A}}\mathbf{U}^{\iota}$ -module but not a ${}_{\mathcal{A}}\mathbf{U}$ -module, and hence Theorem 4 is needed.

Denote

$$T_f^{\mathbf{b},\mathfrak{l}} = M_f^{\mathbf{b},\mathfrak{l}} + \sum_g t_{gf}^{\mathbf{b}}(q) M_g^{\mathbf{b},\mathfrak{l}}, \quad \text{for } t_{gf}^{\mathbf{b}}(q) \in \mathbb{Z}[q].$$

By Theorem 10, $t_{gf}^{\mathbf{b}}(q)$ plays the role of Kazhdan-Lusztig polynomials for $\mathcal{O}_{\mathbf{b}}^{\mathfrak{l}}$. The following positivity and finiteness results generalize [BW18a, Theorem 9.11] and follow by the same proof.

Theorem 12.

- (1) We have $t_{gf}^{\mathbf{b},\mathfrak{l}}(q) \in \mathbb{N}[q]$.
- (2) The sum $T_f^{\mathbf{b},\mathfrak{l}} = M_f^{\mathbf{b},\mathfrak{l}} + \sum_g t_{gf}^{\mathbf{b}}(q) M_g^{\mathbf{b},\mathfrak{l}}$ is finite, for all f .

Remark 13. To formulate a super Kazhdan-Lusztig theory for the parabolic category \mathcal{O} consisting of modules of half-integer weights for $\mathfrak{osp}(2m+1|2n)$, we use the quantum symmetric pair $(\mathbf{U}, \mathbf{U}^{\iota})$ which is a direct limit of $(\mathbf{U}(\mathfrak{sl}_N), \mathbf{U}^{\iota}(\mathfrak{sl}_N))$ for N odd; cf. [BW18a, Sections 6, 12]. Theorem 10 holds again in this setting.

Remark 14. Following [Bao17], a simple conceptual modification allows us to formulate a super (type D) Kazhdan-Lusztig theory for the parabolic category \mathcal{O} consisting of modules of integer (respectively, half-integer) weights for $\mathfrak{osp}(2m|2n)$. To that end, we use the ι -canonical basis of the module (3.1) for the quantum symmetric pair $(\mathbf{U}, \mathbf{U}^{\iota})$, where the parameter is now chosen to be $\kappa = 0$ in the notation of [BW18b]. Theorem 10

holds again in this setting, where the cases new to this paper correspond to the cases $a_0 > 0$.

REFERENCES

- [Bao17] H. Bao, *Kazhdan-Lusztig theory of super type D and quantum symmetric pairs*, Represent. Theory **21** (2017), 247–276, [arXiv:1603.05105](#).
- [Br03] J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra $\mathfrak{gl}(m|n)$* , J. Amer. Math. Soc. **16** (2003), 185–231.
- [BK19] M. Balagovic and S. Kolb, *Universal K-matrix for quantum symmetric pairs*, J. Reine Angew. Math. **747** (2019), 299–353, [arXiv:1507.06276v2](#).
- [BW18a] H. Bao and W. Wang, *A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs*, Astérisque **402**, 2018, vii+134pp, [arXiv:1310.0103](#).
- [BW18b] ———, *Canonical bases arising from quantum symmetric pairs*, Invent. Math. **213** (2018), 1099–1177, [arXiv:1610.09271v2](#).
- [CLW11] S.-J. Cheng, N. Lam and W. Wang, *Super duality and irreducible characters of ortho-symplectic Lie superalgebras*, Invent. Math. **183** (2011), 189–224.
- [CLW15] S.-J. Cheng, N. Lam and W. Wang, *Brundan-Kazhdan-Lusztig conjecture for general linear Lie superalgebras*, Duke J. Math. **164** (2015), 617–695.
- [Ko17] S. Kolb, *Braided module categories via quantum symmetric pairs*, [arXiv:1705.04238v2](#).
- [Le99] G. Letzter, *Symmetric pairs for quantized enveloping algebras*, J. Algebra **220** (1999), 729–767.
- [Lu92] G. Lusztig, *Canonical bases in tensor products*, Proc. Nat. Acad. Sci. **89** (1992), 8177–8179.
- [Lu94] G. Lusztig, *Introduction to Quantum Groups*, Modern Birkhäuser Classics, Reprint of the 1993 Edition, Birkhäuser, Boston, 2010.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076, SINGAPORE.

E-mail address: huanchen@nus.edu.sg

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904, UNITED STATES.

E-mail address: ww9c@virginia.edu

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN

E-mail address: hideya@kurims.kyoto-u.ac.jp