

# SHARP HARDY-LERAY AND RELICH-LERAY INEQUALITIES FOR CURL-FREE VECTOR FIELDS

NAOKI HAMAMOTO AND FUTOSHI TAKAHASHI

ABSTRACT. In this paper, we prove Hardy-Leray and Rellich-Leray inequalities for curl-free vector fields with sharp constants. This complements the former work by Costin-Maz'ya [2] on the sharp Hardy-Leray inequality for axisymmetric divergence-free vector fields.

## 1. INTRODUCTION

In this paper, we concern the classical functional inequalities called Hardy-Leray and Rellich-Leray inequality for smooth vector fields and study how the best constants will change according to the pointwise constraints on their differentials.

Let  $N \in \mathbb{N}$  be an integer with  $N \geq 2$  and put  $\mathbf{x} = (x_1, \dots, x_N) \in \mathbb{R}^N$ . In the following,  $C_c^\infty(\mathbb{R}^N)^N$  denotes a set of smooth vector fields

$$\mathbf{u} = (u_1, u_2, \dots, u_N) : \mathbb{R}^N \ni \mathbf{x} \mapsto \mathbf{u}(\mathbf{x}) \in \mathbb{R}^N$$

having compact supports on  $\mathbb{R}^N$ . Let  $\gamma \neq 1 - N/2$ . Then it is well known that

$$(1) \quad \left( \gamma + \frac{N}{2} - 1 \right)^2 \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx$$

holds for any vector field  $\mathbf{u} \in C_c^\infty(\mathbb{R}^N)^N$  with  $\mathbf{u}(\mathbf{0}) = \mathbf{0}$  if  $\gamma < 1 - N/2$ . This is a higher dimensional extension of 1-dimensional inequality by G. H. Hardy, see [6], also [10], and was first proved by J. Leray [8] in 1933 when the weight  $\gamma = 0$ , see also the book by Ladyzhenskaya [7]. It is also known that the constant  $\left( \gamma + \frac{N}{2} - 1 \right)^2$  is sharp and never attained. In [2], Costin and Maz'ya proved that if the smooth vector fields are axisymmetric and subject to the divergence-free constraint  $\operatorname{div} \mathbf{u} = 0$ , then the constant in (1) can be improved and replaced by a larger one. More precisely, they proved the following:

**Theorem A.** (Costin-Maz'ya [2]) *Let  $N \geq 3$ . Let  $\gamma \neq 1 - N/2$  be a real number and  $\mathbf{u} \in C_c^\infty(\mathbb{R}^N)^N$  be an axisymmetric divergence-free vector field. Assume that  $\mathbf{u}(\mathbf{0}) = \mathbf{0}$  if  $\gamma < 1 - N/2$ . Then*

$$C_{N,\gamma} \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx$$

---

*Date:* August 30, 2018.

*2010 Mathematics Subject Classification.* Primary 35A23; Secondary 26D10.

*Key words and phrases.* Hardy-Leray inequality, Rellich-Leray inequality, curl-free vector fields.

holds with the optimal constant  $C_{N,\gamma}$  given by

$$C_{N,\gamma} = \begin{cases} \left(\gamma + \frac{N}{2} - 1\right)^2 \frac{N+1 + \left(\gamma - \frac{N}{2}\right)^2}{N-1 + \left(\gamma - \frac{N}{2}\right)^2} & (\gamma \leq 1), \\ \left(\gamma + \frac{N}{2} - 1\right)^2 + 2 + \min_{\kappa \geq 0} \left( \kappa + \frac{4(N-1)(\gamma-1)}{\kappa + N - 1 + \left(\gamma - \frac{N}{2}\right)^2} \right) & (N \geq 4, \gamma > 1), \\ \left(\gamma + \frac{1}{2}\right)^2 + 2 & (N = 3, \gamma > 1), \end{cases}$$

Note that the expression of the best constant  $C(N, \gamma)$  is slightly different from that in [2] when  $N \geq 4$ , but a careful checking the proof in [2] leads to the above formula in Theorem A. Choosing  $\gamma = 0$  in Theorem A, we see that the best constant in (1) is actually improved for axisymmetric divergence-free vector fields in the sense that

$$C_{N,0} \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 dx$$

holds with the optimal constant  $C_{N,0} = \left(\frac{N}{2} - 1\right)^2 \frac{N^2 + 4N + 4}{N^2 + 4N - 4} > \left(\frac{N-2}{2}\right)^2$ .

In 2-dimensional case, the result in [2] reads as follows:

**Theorem B.** (Costin-Maz'ya [2]) *Let  $\gamma \neq 0$  be a real number and  $\mathbf{u} \in C_c^\infty(\mathbb{R}^2)^2$  be a divergence-free vector field. We assume that  $\mathbf{u}(\mathbf{0}) = \mathbf{0}$  if  $\gamma < 0$ . Then*

$$C_{2,\gamma} \int_{\mathbb{R}^2} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^2} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx$$

holds with the optimal constant  $C_{2,\gamma}$  given by

$$C_{2,\gamma} = \begin{cases} \gamma^2 \frac{3 + (\gamma - 1)^2}{1 + (\gamma - 1)^2} & \text{for } |\gamma + 1| \leq \sqrt{3}, \\ \gamma^2 + 1 & \text{otherwise.} \end{cases}$$

When  $N = 2$ ,  $\mathbf{u}$  in Theorem B need not be axisymmetric. Furthermore if we consider  $\mathbf{u}^\perp = (-u_2, u_1)$  for  $\mathbf{u} = (u_1, u_2)$  in Theorem B, then the condition  $\operatorname{div} \mathbf{u} = 0$  is replaced by  $\operatorname{curl} \mathbf{u}^\perp = 0$  and also  $|\nabla \mathbf{u}|^2 = |\nabla \mathbf{u}^\perp|^2$ . Thus the above inequality in Theorem B holds also for curl-free vector fields with the same constant.

Motivated by this observation, our aim in this paper is to generalize Costin-Maz'ya's result for curl-free vector fields when  $N = 2$  to higher-dimensional cases. In addition, we also consider the Rellich type inequality involving higher-order derivative,  $\Delta \mathbf{u}$ , for curl-free vector fields. We refer to [5] for the Rellich-Leray inequality for divergence-free vector fields.

Now, main results of this paper are as follows:

**Theorem 1.** *Let  $\gamma \neq 1 - N/2$  be a real number and let  $\mathbf{u} \in C_c^\infty(\mathbb{R}^N)^N$  be a curl-free vector field. We assume that  $\mathbf{u}(\mathbf{0}) = \mathbf{0}$  if  $\gamma < 1 - N/2$ . Then*

$$(2) \quad H_{N,\gamma} \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx$$

with the optimal constant  $H_{N,\gamma}$  given by

$$(3) \quad H_{N,\gamma} = \begin{cases} \left(\gamma + \frac{N}{2} - 1\right)^2 \frac{3(N-1) + \left(\gamma + \frac{N}{2} - 2\right)^2}{N-1 + \left(\gamma + \frac{N}{2} - 2\right)^2} & \text{if } \left|\gamma + \frac{N}{2}\right| \leq \sqrt{N+1}, \\ \left(\gamma + \frac{N}{2} - 1\right)^2 + N - 1 & \text{otherwise.} \end{cases}$$

We remark that no symmetry assumption for  $\mathbf{u}$  is needed. Theorem 1 corresponds to the higher-dimensional analogue of Theorem B in the sense that  $C_{2,\gamma} = H_{2,\gamma}$ .

For curl-free vector fields  $\mathbf{u}$ , Poincaré's lemma implies that there exists a smooth  $\phi$  such that  $\mathbf{u} = \nabla\phi$ . Thus by using the potential function  $\phi$ , Theorem 1 is equivalent to the following corollary.

**Corollary 2.** *Let  $\gamma \neq 1 - N/2$  be a real number and let  $\phi \in C_c^\infty(\mathbb{R}^N)$ . We assume that  $\nabla\phi(\mathbf{0}) = \mathbf{0}$  if  $\gamma < 1 - N/2$ . Then*

$$H_{N,\gamma} \int_{\mathbb{R}^N} \frac{|\nabla\phi|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |D^2\phi|^2 |\mathbf{x}|^{2\gamma} dx$$

with the optimal constant  $H_{N,\gamma}$  given in (3). Here  $D^2\phi(\mathbf{x}) = \left(\frac{\partial^2\phi}{\partial x_i \partial x_j}(\mathbf{x})\right)_{1 \leq i, j \leq N}$  denotes the Hessian matrix of  $\phi$ .

By similar arguments, we prove the following Rellich-Leray inequality for curl-free vector fields.

**Theorem 3.** *Let  $\gamma \neq 2 - N/2$  be a real number and let  $\mathbf{u} \in C_c^\infty(\mathbb{R}^N)^N$  be a curl-free vector field. We assume that  $\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\mathbf{u}|^2 dx < \infty$ . Then*

$$(4) \quad R_{N,\gamma} \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^4} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\Delta\mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx$$

with the optimal constant  $R_{N,\gamma}$  given by

$$(5) \quad R_{N,\gamma} = \min_{\nu \in \mathbb{N} \cup \{0\}} \frac{\left(1 - \frac{N}{2} - \gamma\right)^2 + \alpha_\nu}{\left(3 - \frac{N}{2} - \gamma\right)^2 + \alpha_\nu} (\alpha_{3 - \frac{N}{2} - \gamma} - \alpha_\nu)^2$$

for  $\gamma \neq 3 - N/2$ , where we put

$$\alpha_s = s(s + N - 2), \quad s \in \mathbb{R},$$

and

$$(6) \quad R_{N,3-N/2} = \begin{cases} 4(N-2)^2 & \text{for } N = 2, 3, 4, \\ (N+3)(N-1) & \text{for } N \geq 5. \end{cases}$$

**Corollary 4.** *Let  $\gamma \neq 2 - N/2$  be a real number and let  $\phi \in C_c^\infty(\mathbb{R}^N)$  be a potential function such that  $\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\nabla\phi|^2 dx < \infty$ . Then*

$$R_{N,\gamma} \int_{\mathbb{R}^N} \frac{|\nabla\phi|^2}{|\mathbf{x}|^4} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla\Delta\phi|^2 |\mathbf{x}|^{2\gamma} dx$$

holds with the optimal constant  $R_{N,\gamma}$  as in (5) and (6).

**Remark 5.** *We do not know that the optimal constants  $H_{N,\gamma}$  and  $R_{N,\gamma}$  are attained or not in the class of vector fields in Theorem 1 and Theorem 3.*

*Also in Theorem 1 and Theorem 3, if we restrict ourselves only on vector fields in  $C_c^\infty(\mathbb{R}^N \setminus \{\mathbf{0}\})$ , then the additional assumptions  $\mathbf{u}(\mathbf{0}) = \mathbf{0}$  if  $\gamma < 1 - N/2$ , or  $\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\mathbf{u}|^2 dx < \infty$  are not needed and the same conclusions hold true.*

Concerning Corollary 2 which is equivalent to Theorem 1, we should remark that the similar results already exist by [11], [3]; see also [4] Chapter 6.5. More precisely, improving the work by Tertikas and Zographopoulos [11], Ghoussoub and Moradifam ([3]: Appendix B) prove the following: Let  $C_c^\infty(B_R)$  denote the set of smooth functions having compact supports in a ball  $B_R \subset \mathbb{R}^N$  with radius  $R$ . Define

$$A_{N,\gamma}(R) = \inf \left\{ \frac{\int_{B_R} |\Delta\phi|^2 |\mathbf{x}|^{2\gamma} dx}{\int_{B_R} \frac{|\nabla\phi|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx} : \phi \in C_c^\infty(B_R) \right\}.$$

Assume  $\gamma \geq 1 - N/2$ . Then  $A_{N,\gamma}(R)$  is independent of  $R$ , and is equal to

$$A_{N,\gamma} = \min_{\nu \in \mathbb{N} \cup \{0\}} \left\{ \frac{\left( \frac{(N-4+2\gamma)(N-2\gamma)}{4} + \alpha_\nu \right)^2}{\left( \frac{N-4+2\gamma}{2} \right)^2 + \alpha_\nu} \right\},$$

where  $\alpha_\nu = \nu(N+\nu-2)$  ( $\nu \in \mathbb{N} \cup \{0\}$ ) is the  $\nu$ -th eigenvalue of the Laplace-Beltrami operator on the unit sphere  $\mathbb{S}^{N-1}$  in  $\mathbb{R}^N$ . Note that by a simple formula

$$|D^2\phi|^2 = \sum_{i,j=1}^N \left( \frac{\partial^2\phi}{\partial x_i \partial x_j} \right)^2 = \operatorname{div} \left( \frac{1}{2} \nabla |\nabla\phi|^2 - (\Delta\phi) \nabla\phi \right) + (\Delta\phi)^2,$$

for  $\phi \in C_c^\infty(B_R)$ , we have  $\int_{B_R} |D^2\phi|^2 dx = \int_{B_R} |\Delta\phi|^2 dx$  which implies  $H_{N,0} = A_{N,0}$ . However, in weighted cases, it holds  $\int_{B_R} |D^2\phi|^2 |\mathbf{x}|^{2\gamma} dx \neq \int_{B_R} |\Delta\phi|^2 |\mathbf{x}|^{2\gamma} dx$ , and in general we have  $H_{N,\gamma} \neq A_{N,\gamma}$ . Also the inequality in Corollary 4 seems new.

The organization of this paper is as follows: In §2, we recall the method by Costin-Maz'ya in [2] and derive the equivalent curl-free condition in polar coordinates. In §3, we prove Theorem 1 and the sharpness of the constant (3). In §4, we prove Theorem 3 and the sharpness of the constants (5) and (6). Since the test vector fields introduced in [2] may not have compact supports, we will use different test vector fields for the proof of the sharpness of the constants.

## 2. PREPARATION: COSTIN-MAZ'YA'S SETTING

In this section, we recall the method of Costin-Maz'ya [2] and derive the polar coordinate representation of the curl-free condition.

**Spherical polar coordinate.** We introduce the spherical polar coordinates

$$(\rho, \theta_1, \theta_2, \dots, \theta_{N-2}, \theta_{N-1}) \in (0, \infty) \times (0, \pi)^{N-2} \times [0, 2\pi)$$

whose relation to the standard Euclidean coordinates  $\mathbf{x} = (x_1, \dots, x_N) \in \mathbb{R}^N$  is given by

$$\mathbf{x} = \rho(\cos\theta_1, \pi_1 \cos\theta_2, \pi_2 \cos\theta_3, \dots, \pi_{k-1} \cos\theta_k, \dots, \pi_{N-2} \cos\theta_{N-1}, \pi_{N-1}),$$

hereafter we use the notation

$$\pi_0 = 1, \quad \pi_k = \prod_{j=1}^k \sin\theta_j, \quad (k = 1, 2, \dots, N-1)$$

for simplicity. Also we use the notation

$$\partial_\rho = \frac{\partial}{\partial\rho}, \quad \partial_{\theta_k} = \frac{\partial}{\partial\theta_k}, \quad (k = 1, 2, \dots, N-1)$$

for the partial derivatives, and

$$dx = \prod_{k=1}^N dx_k = dx_1 dx_2 \cdots dx_N, \quad d\sigma = \prod_{k=1}^{N-1} (\sin \theta_k)^{N-k-1} d\theta_k$$

for the volume elements on  $\mathbb{R}^N$  and  $\mathbb{S}^{N-1}$ .

The orthonormal basis vector fields  $\mathbf{e}_\rho, \mathbf{e}_{\theta_1}, \mathbf{e}_{\theta_2}, \dots, \mathbf{e}_{\theta_{N-1}}$  along the polar coordinates are given by

$$(7) \quad \begin{cases} \mathbf{e}_\rho = \frac{\partial_\rho \mathbf{x}}{|\partial_\rho \mathbf{x}|} = (\cos \theta_1, \pi_1 \cos \theta_2, \pi_2 \cos \theta_3, \dots, \pi_{N-2} \cos \theta_{N-1}, \pi_{N-1}), \\ \mathbf{e}_{\theta_k} = \frac{\partial_{\theta_k} \mathbf{x}}{|\partial_{\theta_k} \mathbf{x}|} = \frac{1}{\pi_{k-1}} \partial_{\theta_k} \mathbf{e}_\rho, \quad (k = 1, 2, \dots, N-1) \end{cases}$$

that are clearly independent of the variable  $\rho$ . Note that we can rewrite

$$\begin{aligned} \mathbf{e}_\rho &= (\cos \theta_1, \pi_1 \cos \theta_2, \pi_2 \cos \theta_3, \dots, \pi_{k-1} \cos \theta_k, \pi_k \boldsymbol{\varphi}_k), \\ \mathbf{e}_{\theta_k} &= \underbrace{(0, 0, \dots, 0)}_{k-1}, -\sin \theta_k, \cos \theta_k \boldsymbol{\varphi}_k, \end{aligned}$$

where

$$\boldsymbol{\varphi}_k = \left( \cos \theta_{k+1}, \frac{\pi_{k+1}}{\pi_k} \cos \theta_{k+2}, \frac{\pi_{k+2}}{\pi_k} \cos \theta_{k+3}, \dots, \frac{\pi_{N-2}}{\pi_k} \cos \theta_{N-1}, \frac{\pi_{N-1}}{\pi_k} \right) \in \mathbb{S}^{N-k-1}$$

is a  $(N-k)$ -vector, which depends only on  $\theta_{k+1}, \dots, \theta_{N-1}$ . From these expressions, we can easily check the orthonormality of  $\mathbf{e}_\rho, \mathbf{e}_{\theta_1}, \mathbf{e}_{\theta_2}, \dots, \mathbf{e}_{\theta_{N-1}}$ .

For any smooth vector field  $\mathbf{u} = (u_1, u_2, \dots, u_N) : \mathbb{R}^N \rightarrow \mathbb{R}^N$ , its polar components  $u_\rho, u_{\theta_1}, u_{\theta_2}, \dots, u_{\theta_{N-1}}$  as  $\mathbb{R}$ -valued smooth functions are defined by

$$\mathbf{u} = u_\rho \mathbf{e}_\rho + \sum_{k=1}^{N-1} u_{\theta_k} \mathbf{e}_{\theta_k}.$$

The second term of the right-hand side is denoted as

$$\mathbf{u}_\sigma = \sum_{k=1}^{N-1} u_{\theta_k} \mathbf{e}_{\theta_k}$$

and we call this the spherical component of  $\mathbf{u}$ . Thus we have the polar decomposition of  $\mathbf{u}$ :

$$(8) \quad \mathbf{u} = u_\rho \mathbf{e}_\rho + \mathbf{u}_\sigma$$

which gives the decomposition of  $\mathbf{u}$  into the radial and the spherical parts. Also by using the chain rules together with (7), we have

$$\partial_\rho = \mathbf{e}_\rho \cdot \nabla, \quad \text{and} \quad \frac{1}{\rho} \partial_{\theta_k} = \pi_{k-1} \mathbf{e}_{\theta_k} \cdot \nabla, \quad (k = 1, \dots, N-1),$$

which give the polar decomposition of the gradient operator  $\nabla$ :

$$(9) \quad \nabla = \mathbf{e}_\rho \partial_\rho + \frac{1}{\rho} \nabla_\sigma,$$

where

$$(10) \quad \nabla_\sigma = \sum_{k=1}^{N-1} \frac{\mathbf{e}_{\theta_k}}{\pi_{k-1}} \partial_{\theta_k}$$

is the gradient operator on  $\mathbb{S}^{N-1}$ .

Moreover, it is well-known that the polar representation of the Laplace operator  $\Delta = \sum_{k=1}^N \partial^2 / \partial x_k^2$  is given by

$$(11) \quad \Delta = \frac{1}{\rho^{N-1}} \partial_\rho (\rho^{N-1} \partial_\rho) + \frac{1}{\rho^2} \Delta_\sigma ,$$

where

$$(12) \quad \Delta_\sigma = \sum_{k=1}^{N-1} \frac{(\sin \theta_k)^{k+1-N}}{\pi_{k-1}^2} \partial_{\theta_k} ((\sin \theta_k)^{N-k-1} \partial_{\theta_k}) = \sum_{k=1}^{N-1} \frac{1}{\pi_{k-1}^2} D_{\theta_k} \partial_{\theta_k}$$

is the Laplace-Beltrami operator on  $\mathbb{S}^{N-1}$  and for every  $k = 1, \dots, N-1$

$$D_{\theta_k} = \partial_{\theta_k} + (N-k-1) \cot \theta_k$$

is the adjoint operator of  $-\partial_{\theta_k}$  in  $L^2(d\sigma, \mathbb{S}^{N-1})$ : it holds that

$$-\int_{\mathbb{S}^{N-1}} f (\partial_{\theta_k} g) d\sigma = \int_{\mathbb{S}^{N-1}} (D_{\theta_k} f) g d\sigma$$

for any smooth functions  $f, g$  on  $\mathbb{S}^{N-1}$ .

We also introduce the deformed radial coordinate  $t \in \mathbb{R}$  by the Emden transformation

$$(13) \quad t = \log \rho.$$

Note that (13) leads to the transformation law of the differential operators  $\rho \partial_\rho = \partial_t$ . By this transformation, it is easy to check that the polar decomposition of  $\nabla$ ,  $\Delta$  in (9), (11) are also given by

$$(14) \quad \rho \nabla = \mathbf{e}_\rho \partial_t + \nabla_\sigma,$$

$$(15) \quad \rho^2 \Delta = \partial_t^2 + (N-2) \partial_t + \Delta_\sigma.$$

For the later use, we prove the following lemma.

**Lemma 6.** *Let  $\nabla_\sigma$  and  $\Delta_\sigma$  are defined by (10) and (12) respectively. Then for any  $f \in C^\infty(\mathbb{S}^{N-1})$ ,  $\sigma = \mathbf{e}_\rho \in \mathbb{S}^{N-1}$  and  $\alpha \in \mathbb{C}$ , there hold that*

$$(16) \quad \Delta_\sigma(\mathbf{e}_\rho f) - \mathbf{e}_\rho \Delta_\sigma f = (2\nabla_\sigma - (N-1)\mathbf{e}_\rho) f,$$

$$(17) \quad \Delta_\sigma \nabla_\sigma f - \nabla_\sigma \Delta_\sigma f = ((N-3)\nabla_\sigma - 2\mathbf{e}_\rho \Delta_\sigma) f,$$

$$(18) \quad \Delta_\sigma (f \mathbf{e}_\rho + \alpha \nabla_\sigma f) = \mathbf{e}_\rho ((1-2\alpha)\Delta_\sigma f - (N-1)f) \\ + (2 + (N-3)\alpha) \nabla_\sigma f + \alpha \nabla_\sigma \Delta_\sigma f.$$

*Proof.* Take any  $f \in C^\infty(\mathbb{S}^{N-1})$ . We identify  $f$  and the function  $\tilde{f} \in C^\infty(\mathbb{R}^N \setminus \{\mathbf{0}\})$  defined by  $\tilde{f}(\mathbf{x}) = f(\mathbf{x}/|\mathbf{x}|)$ . Since  $f = \tilde{f}$  does not depend on the radial variable  $\rho$ , we have  $\nabla_\sigma f = \rho \nabla f$  by (9) and  $\Delta_\sigma f = \rho^2 \Delta f$  by (11). Thus we compute

$$\begin{aligned} \Delta_\sigma(\mathbf{e}_\rho f) - \mathbf{e}_\rho \Delta_\sigma f &= \rho^2 \Delta \left( \frac{\mathbf{x} f}{\rho} \right) - \frac{\mathbf{x}}{\rho} \rho^2 \Delta f \\ &= \rho^2 \left( \frac{\Delta(\mathbf{x} f)}{\rho} + 2((\nabla \rho^{-1}) \cdot \nabla)(\mathbf{x} f) + (\Delta \rho^{-1}) \mathbf{x} f \right) - \rho \mathbf{x} \Delta f \\ &= 2\rho(\nabla f \cdot \nabla) \mathbf{x} - 2(\nabla \rho \cdot \nabla)(\mathbf{x} f) + \rho^3 (\Delta \rho^{-1}) \mathbf{e}_\rho f \\ &= 2\rho \nabla f - 2\partial_\rho(\rho \mathbf{e}_\rho f) - (N-3) \mathbf{e}_\rho f \\ &= (2\nabla_\sigma - (N-1)\mathbf{e}_\rho) f, \end{aligned}$$

here we have used  $\nabla\rho \cdot \nabla = \partial_\rho$  and  $\Delta\rho^{-1} = -(N-3)\rho^{-3}$ . This proves (16). Similarly, also noting the commutativity  $\Delta\nabla = \nabla\Delta$  and using  $\Delta\rho = (N-1)\rho^{-1}$ , we have

$$\begin{aligned} (\Delta_\sigma \nabla_\sigma - \nabla_\sigma \Delta_\sigma) f &= \rho^2 \Delta \nabla_\sigma f - \rho \nabla \Delta_\sigma f \\ &= \rho^2 \Delta (\rho \nabla f) - \rho \nabla (\rho^2 \Delta f) \\ &= \rho^2 ((\Delta \rho) \nabla f + 2(\nabla \rho \cdot \nabla) \nabla f) - \rho (\nabla \rho^2) \Delta f \\ &= (N-1)\rho \nabla f + 2\rho^2 \partial_\rho \rho^{-1} \nabla_\sigma f - 2\rho^2 e_\rho \Delta f \\ &= (N-3) \nabla_\sigma f - 2e_\rho \Delta_\sigma f. \end{aligned}$$

This proves (17). Finally, by (16) and (17), we see

$$\begin{aligned} \Delta_\sigma (f e_\rho + \alpha \nabla_\sigma f) &= \Delta_\sigma (e_\rho f) + \alpha \Delta_\sigma \nabla_\sigma f \\ &= (e_\rho \Delta_\sigma + 2\nabla_\sigma - (N-1)e_\rho) f + \alpha (\nabla_\sigma \Delta_\sigma + (N-3)\nabla_\sigma - 2e_\rho \Delta_\sigma) f \\ &= e_\rho ((1-2\alpha)\Delta_\sigma f - (N-1)f) + (2 + (N-3)\alpha) \nabla_\sigma f + \alpha \nabla_\sigma \Delta_\sigma f. \end{aligned}$$

This proves (18).  $\square$

**Representing the curl-free condition in polar coordinates.** In the following, let “ $\cdot$ ” denote the standard inner product in  $\mathbb{R}^N$ , “ $\wedge$ ” the wedge product for differential forms and “ $d$ ” the exterior derivative operator. For a vector field  $\mathbf{a} = (a_1, a_2, \dots, a_N) : \mathbb{R}^N \rightarrow \mathbb{R}^N$ , we define the vector valued 1-form  $d\mathbf{a} = (da_1, da_2, \dots, da_N)$ . If  $\mathbf{u} = (u_1, u_2, \dots, u_N)$  is a vector field, then  $\mathbf{u} \cdot d\mathbf{a}$  denotes the 1-form  $\sum_{k=1}^N u_k da_k$ . Now, for any  $C^\infty$  vector field  $\mathbf{u} : \mathbb{R}^N \rightarrow \mathbb{R}^N$  with variable  $\mathbf{x} = (x_1, \dots, x_N)$ , we define its curl as the differential 2-form

$$\text{curl } \mathbf{u} = d(\mathbf{u} \cdot d\mathbf{x}).$$

This can be expressed in terms of the standard Euclidean coordinates, according to the usual manipulations for the exterior derivative  $d$  and the wedge product  $\wedge$  :

$$d(\mathbf{u} \cdot d\mathbf{x}) = \sum_{k=1}^N du_k \wedge dx_k = \sum_{k=1}^N \sum_{j=1}^N \frac{\partial u_k}{\partial x_j} dx_j \wedge dx_k = \sum_{j < k} \sum \left( \frac{\partial u_k}{\partial x_j} - \frac{\partial u_j}{\partial x_k} \right) dx_j \wedge dx_k.$$

As well as the standard representation, we want to find a representation of  $d(\mathbf{u} \cdot d\mathbf{x})$  in terms of the polar coordinates  $(\rho, \theta_1, \dots, \theta_{N-1})$ . For this purpose, first we differentiate the unit vector field  $e_\rho$  given by (7) and expand it in the spherical-coordinate basis:

$$de_\rho = \sum_{k=1}^{N-1} \frac{\partial e_\rho}{\partial \theta_k} d\theta_k = \sum_{k=1}^{N-1} e_{\theta_k} \pi_{k-1} d\theta_k.$$

Then, taking the inner product with the vector field  $\mathbf{u} = u_\rho e_\rho + \sum_{k=1}^{N-1} u_{\theta_k} e_{\theta_k}$  and also taking its exterior derivative, we see that

$$\begin{aligned} \mathbf{u} \cdot de_\rho &= \sum_{k=1}^{N-1} u_{\theta_k} \pi_{k-1} d\theta_k, \\ d(\mathbf{u} \cdot de_\rho) &= d\rho \wedge \sum_{k=1}^{N-1} (\partial_\rho u_{\theta_k}) \pi_{k-1} d\theta_k + \sum_{j=1}^{N-1} \sum_{k=1}^{N-1} \partial_{\theta_j} (\pi_{k-1} u_{\theta_k}) d\theta_j \wedge d\theta_k. \end{aligned}$$

Also we have

$$\mathbf{u} \cdot d\mathbf{x} = \mathbf{u} \cdot \left( (d\rho)\mathbf{e}_\rho + \rho d\mathbf{e}_\rho \right) = u_\rho d\rho + \rho \mathbf{u} \cdot d\mathbf{e}_\rho.$$

From these relations, we obtain the polar representation of the curl of  $\mathbf{u}$ :

$$\begin{aligned} d(\mathbf{u} \cdot d\mathbf{x}) &= d(u_\rho d\rho + \rho \mathbf{u} \cdot d\mathbf{e}_\rho) \\ &= du_\rho \wedge d\rho + d\rho \wedge (\mathbf{u} \cdot d\mathbf{e}_\rho) + \rho d(\mathbf{u} \cdot d\mathbf{e}_\rho) \\ &= d\rho \wedge \left( -du_\rho + \sum_k u_{\theta_k} \pi_{k-1} d\theta_k + \sum_k \rho \partial_\rho u_{\theta_k} \pi_{k-1} d\theta_k \right) \\ &\quad + \rho \sum_j \sum_k \partial_{\theta_j} (\pi_{k-1} u_{\theta_k}) d\theta_j \wedge d\theta_k \\ &= d\rho \wedge \sum_k \left( \pi_{k-1} \partial_\rho (\rho u_{\theta_k}) - \partial_{\theta_k} u_\rho \right) d\theta_k \\ &\quad + \rho \sum_{j < k} \left( \partial_{\theta_j} (\pi_{k-1} u_{\theta_k}) - \partial_{\theta_k} (\pi_{j-1} u_{\theta_j}) \right) d\theta_j \wedge d\theta_k. \end{aligned}$$

Therefore, the curl-free condition  $d(\mathbf{u} \cdot \mathbf{x}) = 0$  for the vector field  $\mathbf{u}$  is represented by

$$(19) \quad \begin{cases} \partial_\rho (\rho \pi_{k-1} u_{\theta_k}) = \partial_{\theta_k} u_\rho \\ \partial_{\theta_j} (\pi_{k-1} u_{\theta_k}) = \partial_{\theta_k} (\pi_{j-1} u_{\theta_j}) \end{cases}, \quad (j, k = 1, 2, \dots, N-1).$$

We claim that the second relations in (19) are actually the consequences of the first. Indeed, by integrating the first equation in (19) on any interval  $(0, r] \subset \mathbb{R}$  with respect to the variable  $\rho$ , we have  $r\pi_{k-1}u_{\theta_k} = \partial_{\theta_k} \int_0^r u_\rho d\rho$  for every  $k$ . Thus the function  $\phi \in C^\infty(\mathbb{R}^N \setminus \{\mathbf{0}\})$  defined by  $\phi(\mathbf{x}) = \frac{1}{|\mathbf{x}|} \int_0^{|\mathbf{x}|} u_\rho(\rho \mathbf{x}/|\mathbf{x}|) d\rho$  satisfies  $\pi_{k-1}u_{\theta_k} = \partial_{\theta_k} \phi$  for all  $k$ . Then the second relation in (19) is the same as  $\partial_{\theta_j} \partial_{\theta_k} \phi = \partial_{\theta_k} \partial_{\theta_j} \phi$ , which holds trivially. This proves the claim.

Consequently we have proved that a vector field  $\mathbf{u} \in C^\infty(\mathbb{R}^N)^N$  is curl-free if and only if

$$\partial_\rho (\rho u_{\theta_k}) = \frac{1}{\pi_{k-1}} \partial_{\theta_k} u_\rho, \quad k = 1, \dots, N-1.$$

That is, using vector notation as in (8) and (9), we have

$$(20) \quad \partial_\rho (\rho \mathbf{u}_\sigma) = \nabla_\sigma u_\rho \quad (\rho, \sigma) \in \mathbb{R}_+ \times \mathbb{S}^{N-1}.$$

In what follows we also call (20) the curl-free condition for  $\mathbf{u}$ .

**Brezis-Vazquez, Maz'ya transformation.** Let  $\varepsilon \neq 1$  be a real number. As in [1], [9], we introduce a new vector field  $\mathbf{v}$  by the formula

$$(21) \quad \mathbf{v}(\mathbf{x}) = \rho^{1-\varepsilon} \mathbf{u}(\mathbf{x}).$$

Then the curl-free condition (20) is transformed into

$$\nabla_\sigma (\rho^{\varepsilon-1} v_\rho) = \partial_\rho (\rho^\varepsilon \mathbf{v}_\sigma),$$

that is,

$$(22) \quad (\varepsilon + \rho \partial_\rho) \mathbf{v}_\sigma = \nabla_\sigma v_\rho.$$



**Fourier transformation in radial direction.** In the following, let us denote  $\mathbf{x} = \rho\boldsymbol{\sigma}$  and we abbreviate  $\mathbf{v}(t, \boldsymbol{\sigma}) = \mathbf{v}(e^t\boldsymbol{\sigma})$  for a vector field  $\mathbf{v}(\mathbf{x}) = \mathbf{v}(\rho\boldsymbol{\sigma})$ , where  $t = \log \rho$  is the Emden transformation given in (13). As in [2], we apply the one-dimensional Fourier transformation

$$\mathbf{v}(t, \boldsymbol{\sigma}) \mapsto \widehat{\mathbf{v}}(\lambda, \boldsymbol{\sigma}) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-i\lambda t} \mathbf{v}(t, \boldsymbol{\sigma}) dt$$

with respect to the variable  $t$ . By the transformation law between the derivative operators

$$\rho \partial_\rho = \partial_t \mapsto \widehat{\partial}_t = i\lambda \cdot ,$$

the curl-free condition (22) is changed into the equation

$$(\varepsilon + i\lambda) \widehat{\mathbf{v}}_\sigma = \nabla_\sigma \widehat{v}_\rho ,$$

that is,

$$\widehat{\mathbf{v}}_\sigma = \frac{1}{\varepsilon + i\lambda} \nabla_\sigma f \quad \text{where } f = \widehat{v}_\rho .$$

Thus we see that  $\widehat{\mathbf{v}}_\sigma$  is expressed by the spherical gradient of some function  $f = \widehat{v}_\rho$ . In this sense, we may consider  $f$  as a kind of scalar potential of  $\widehat{\mathbf{v}}$ , corresponding to the fact that the curl-free vector field  $\mathbf{u}$  has a scalar potential.

Now we have proved the following proposition:

**Proposition 7.** *Let  $\varepsilon \neq 1$  and let  $\mathbf{u}$  be a smooth vector field on  $\mathbb{R}^N$ . Then  $\mathbf{u}$  is curl-free if and only if its Brezis-Vázquez, Maz'ya transformation  $\mathbf{v}(t, \boldsymbol{\sigma}) = e^{t(1-\varepsilon)} \mathbf{u}(e^t\boldsymbol{\sigma})$  satisfies*

$$(23) \quad (\varepsilon + \partial_t) \mathbf{v}_\sigma = \nabla_\sigma v_\rho .$$

*In particular, if  $\mathbf{u}$  is curl-free and has a compact support on  $\mathbb{R}^N$ , then the Fourier transformation of  $\mathbf{v}$  satisfies*

$$(24) \quad \widehat{\mathbf{v}}(\lambda, \boldsymbol{\sigma}) = f \mathbf{e}_\rho + \frac{1}{\varepsilon + i\lambda} \nabla_\sigma f$$

*for some complex-valued scalar function  $f = f(\lambda, \boldsymbol{\sigma}) \in C^\infty(\mathbb{R} \times \mathbb{S}^{N-1})$ .*

We list up some formulae for  $\widehat{\mathbf{v}}$  and its differentials. The square length of  $\widehat{\mathbf{v}}$  is

$$|\widehat{\mathbf{v}}|^2 = |f|^2 + \frac{1}{\varepsilon^2 + \lambda^2} |\nabla_\sigma f|^2 .$$

By using Lemma 6, we also see that

$$-\Delta_\sigma \widehat{\mathbf{v}} = \mathbf{e}_\rho \left( (N-1)f + \left( \frac{2}{\varepsilon + i\lambda} - 1 \right) \Delta_\sigma f \right) - \left( \frac{N-3}{\varepsilon + i\lambda} + 2 \right) \nabla_\sigma f - \frac{1}{\varepsilon + i\lambda} \nabla_\sigma \Delta_\sigma f .$$

Then integrating  $|\widehat{\mathbf{v}}|^2$ ,  $-\overline{\widehat{\mathbf{v}}} \cdot \Delta_\sigma \widehat{\mathbf{v}}$  and  $|\Delta_\sigma \widehat{\mathbf{v}}|^2$  over  $\mathbb{S}^{N-1}$ , we find that

$$\begin{aligned} \int_{\mathbb{S}^{N-1}} |\widehat{\mathbf{v}}|^2 d\sigma &= \int_{\mathbb{S}^{N-1}} \overline{f} \left( 1 + \frac{1}{\varepsilon^2 + \lambda^2} (-\Delta_\sigma) \right) f d\sigma, \\ \int_{\mathbb{S}^{N-1}} |\nabla_\sigma \widehat{\mathbf{v}}|^2 d\sigma &= \int_{\mathbb{S}^{N-1}} \overline{f} \left( N-1 + \left( 1 + \frac{3-4\varepsilon-N}{\varepsilon^2 + \lambda^2} \right) (-\Delta_\sigma) + \frac{1}{\varepsilon^2 + \lambda^2} (-\Delta_\sigma)^2 \right) f d\sigma, \\ \int_{\mathbb{S}^{N-1}} |\Delta_\sigma \widehat{\mathbf{v}}|^2 d\sigma &= \int_{\mathbb{S}^{N-1}} \overline{f} \left( (N-1)^2 + \left( 2N+2 + \frac{(N-3)^2 - 8\varepsilon}{\varepsilon^2 + \lambda^2} \right) (-\Delta_\sigma) \right. \\ &\quad \left. + \left( 1 + \frac{10-8\varepsilon-2N}{\varepsilon^2 + \lambda^2} \right) (-\Delta_\sigma)^2 + \frac{1}{\varepsilon^2 + \lambda^2} (-\Delta_\sigma)^3 \right) f d\sigma. \end{aligned}$$

Thus, we have proved the following lemma.

**Lemma 8.** *Let  $\widehat{\mathbf{v}} = f\mathbf{e}_\rho + \frac{1}{\varepsilon+i\lambda}\nabla_\sigma f$  as in (24). Then*

$$\begin{aligned}\int_{\mathbb{S}^{N-1}} |\widehat{\mathbf{v}}|^2 d\sigma &= \int_{\mathbb{S}^{N-1}} \bar{f} P_1(\lambda, -\Delta_\sigma) f d\sigma, \\ \int_{\mathbb{S}^{N-1}} |\nabla_\sigma \widehat{\mathbf{v}}|^2 d\sigma &= \int_{\mathbb{S}^{N-1}} \bar{f} P_2(\lambda, -\Delta_\sigma) f d\sigma, \\ \int_{\mathbb{S}^{N-1}} |\Delta_\sigma \widehat{\mathbf{v}}|^2 d\sigma &= \int_{\mathbb{S}^{N-1}} \bar{f} P_3(\lambda, -\Delta_\sigma) f d\sigma\end{aligned}$$

where the three polynomials  $\alpha \mapsto P_k(\lambda, \alpha)$  ( $k = 1, 2, 3$ ) are given by

$$\begin{aligned}P_1(\lambda, \alpha) &= 1 + \frac{1}{\varepsilon^2 + \lambda^2} \alpha, \\ P_2(\lambda, \alpha) &= N - 1 + \left(1 + \frac{3 - 4\varepsilon - N}{\varepsilon^2 + \lambda^2}\right) \alpha + \frac{1}{\varepsilon^2 + \lambda^2} \alpha^2, \\ P_3(\lambda, \alpha) &= (N - 1)^2 + \left(2N + 2 + \frac{(N - 3)^2 - 8\varepsilon}{\varepsilon^2 + \lambda^2}\right) \alpha \\ &\quad + \left(1 + \frac{10 - 8\varepsilon - 2N}{\varepsilon^2 + \lambda^2}\right) \alpha^2 + \frac{1}{\varepsilon^2 + \lambda^2} \alpha^3.\end{aligned}$$

### 3. PROOF OF THEOREM 1

Let  $\gamma \neq 1 - N/2$  be a real number and put  $\varepsilon = 2 - N/2 - \gamma \neq 1$ . If the right-hand side of (2) diverges, there is nothing to prove. When the right-hand side of (2) is bounded, the smoothness of  $\mathbf{u}$  implies the existence of an integer  $m > \varepsilon - 2$  such that  $\nabla \mathbf{u}(\mathbf{x}) = O(|\mathbf{x}|^m)$  as  $|\mathbf{x}| \rightarrow +0$ . If  $\varepsilon < 1$ , then the Brezis-Vázquez, Maz'ya transformation  $\mathbf{v}(\mathbf{x})$  in (21) is Hölder continuous at  $\mathbf{x} = \mathbf{0}$  and satisfies  $\mathbf{v}(\mathbf{0}) = \mathbf{0}$ . When  $\varepsilon > 1$ , again the assumption  $\mathbf{u}(\mathbf{0}) = \mathbf{0}$  implies  $\mathbf{u}(\mathbf{x}) = O(|\mathbf{x}|^{m+1})$  and  $\mathbf{v}(\mathbf{x}) = O(|\mathbf{x}|^{m+2-\varepsilon})$ , thus the same properties hold for  $\mathbf{v}$ . Also since

$$\begin{aligned}\nabla \mathbf{u} &= \nabla(\rho^{\varepsilon-1} \mathbf{v}) = \rho^{\varepsilon-2} ((\varepsilon - 1)\mathbf{e}_\rho \otimes \mathbf{v} + \rho \nabla \mathbf{v}) \\ &= \rho^{\varepsilon-2} (\mathbf{e}_\rho \otimes (\varepsilon - 1 + \partial_t) \mathbf{v} + \nabla_\sigma \mathbf{v})\end{aligned}$$

by (14), and since  $\iint_{\mathbb{S}^{n-1} \times \mathbb{R}} \partial_t \mathbf{v} \cdot \mathbf{v} d\sigma dt$  vanishes, we calculate

$$\begin{aligned}(25) \quad \int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\nabla \mathbf{u}|^2 dx &= \int_{\mathbb{R}^N} |\mathbf{x}|^{4-2\varepsilon-N} |\nabla \mathbf{u}|^2 dx \\ &= \int_{\mathbb{S}^{N-1}} d\sigma \int_{\mathbb{R}} |\mathbf{e}_\rho \otimes (\varepsilon - 1 + \partial_t) \mathbf{v} + \nabla_\sigma \mathbf{v}|^2 dt \\ &= \iint_{\mathbb{S}^{N-1} \times \mathbb{R}} ((\varepsilon - 1)^2 |\mathbf{v}|^2 + |\partial_t \mathbf{v}|^2 + |\nabla_\sigma \mathbf{v}|^2) d\sigma dt \\ &= \iint_{\mathbb{S}^{N-1} \times \mathbb{R}} (((\varepsilon - 1)^2 + \lambda^2) |\widehat{\mathbf{v}}|^2 + |\nabla_\sigma \widehat{\mathbf{v}}|^2) d\sigma d\lambda \\ &= \iint_{\mathbb{S}^{N-1} \times \mathbb{R}} \bar{f} (((\varepsilon - 1)^2 + \lambda^2) P_1(\lambda, -\Delta_\sigma) + P_2(\lambda, -\Delta_\sigma)) f d\sigma d\lambda\end{aligned}$$

and

$$\begin{aligned}
(26) \quad \int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-2} |\mathbf{u}|^2 dx &= \int_{\mathbb{R}^N} |\mathbf{x}|^{2-2\varepsilon-N} |\mathbf{u}|^2 dx \\
&= \int_{\mathbb{S}^{N-1}} d\sigma \int_0^\infty |v|^2 \frac{d\rho}{\rho} = \iint_{\mathbb{R} \times \mathbb{S}^{N-1}} |v|^2 dt d\sigma \\
&= \iint_{\mathbb{R} \times \mathbb{S}^{N-1}} |\widehat{v}|^2 d\lambda d\sigma = \iint_{\mathbb{R} \times \mathbb{S}^{N-1}} \overline{f} P_1(\lambda, -\Delta_\sigma) f d\lambda d\sigma
\end{aligned}$$

by Lemma 8. Therefore, by (25) and (26), the optimal constant in (2) can be expressed as

$$(27) \quad H_{N,\gamma} = \inf_{\mathbf{u} \neq 0, \operatorname{curl} \mathbf{u} = 0} \frac{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\nabla \mathbf{u}|^2 dx}{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-2} |\mathbf{u}|^2 dx} = \inf_{f \neq 0} \frac{\int_{\mathbb{R} \times \mathbb{S}^{N-1}} \overline{f} Q_1(\lambda, -\Delta_\sigma) f d\lambda d\sigma}{\int_{\mathbb{R} \times \mathbb{S}^{N-1}} \overline{f} P_1(\lambda, -\Delta_\sigma) f d\lambda d\sigma},$$

where  $Q_1(\lambda, \cdot)$  is the polynomial defined by

$$(28) \quad Q_1(\lambda, \alpha) = ((\varepsilon - 1)^2 + \lambda^2) P_1(\lambda, \alpha) + P_2(\lambda, \alpha).$$

**Calculation of a lower bound.** In the same manner as Costin-Maz'ya [2], we expand  $f$  in  $L^2(\mathbb{S}^{N-1})$  by eigenfunctions  $\{\psi_\nu\}_{\nu \in \{0\} \cup \mathbb{N}}$  of  $-\Delta_\sigma$  as

$$(29) \quad f(\lambda, \sigma) = \sum_{\nu=0}^{\infty} f_\nu(\lambda) \psi_\nu(\sigma), \quad \begin{cases} -\Delta_\sigma \psi_\nu = \alpha_\nu \psi_\nu, \\ \alpha_\nu = \nu(\nu + N - 2) \quad (\nu = 0, 1, 2, \dots). \end{cases}$$

Then we find that (27) is estimated from below as

$$H_{N,\gamma} = \inf_{f \neq 0} \frac{\sum_{\nu \in \mathbb{N} \cup \{0\}} \int_{\mathbb{R}} Q_1(\lambda, \alpha_\nu) |f_\nu(\lambda)|^2 d\lambda}{\sum_{\nu \in \mathbb{N} \cup \{0\}} \int_{\mathbb{R}} P_1(\lambda, \alpha_\nu) |f_\nu(\lambda)|^2 d\lambda} \geq \inf_{\lambda \in \mathbb{R} \setminus \{0\}} \inf_{\nu \in \mathbb{N} \cup \{0\}} \frac{Q_1(\lambda, \alpha_\nu)}{P_1(\lambda, \alpha_\nu)},$$

where  $P_1, Q_1$  is as in Lemma 8, (28) and where in the last inequality we have used Lemma 9 in Appendix, applied to  $X = \{(\nu, \lambda) \in (\mathbb{N} \cup \{0\}) \times \mathbb{R}\}$ ,  $\mu = \left(\sum_{\nu \in \mathbb{N} \cup \{0\}} \delta_\nu\right) \times d\lambda$  and  $g(\nu, \lambda) = |f_\nu(\lambda)|^2$ . Therefore, we have

$$(30) \quad H_{N,\gamma} \geq \inf_{\kappa > 0} \inf_{\nu \in \mathbb{N} \cup \{0\}} F(\kappa, \alpha_\nu)$$

with  $F(\kappa, \cdot)$  defined by

$$(31) \quad F(\kappa, \alpha) = \frac{Q_1(\sqrt{\kappa}, \alpha)}{P_1(\sqrt{\kappa}, \alpha)} = (\varepsilon - 1)^2 + N - 1 + \kappa + \alpha - 2\alpha \frac{2\varepsilon + N - 2}{\varepsilon^2 + \kappa + \alpha}$$

for  $\kappa > 0$  and  $\alpha \geq 0$ . Here we also define  $F(0, \alpha)$  by

$$\begin{aligned}
(32) \quad F(0, \alpha) &= \lim_{|\lambda| \searrow +0} \frac{Q_1(\lambda, \alpha)}{P_1(\lambda, \alpha)} = \lim_{\kappa \searrow +0} F(\kappa, \alpha) \\
&= \begin{cases} (\varepsilon - 1)^2 + N - 1 + \alpha - 2\alpha \frac{2\varepsilon + N - 2}{\varepsilon^2 + \alpha} & \text{for } \alpha > 0 \\ (\varepsilon - 1)^2 + N - 1 & \text{for } \alpha = 0. \end{cases}
\end{aligned}$$

In this setting, we calculate the right-hand side of (30). In the case  $\varepsilon < 1 - N/2$ , by differentiating (31) directly with respect to  $\alpha$ , we see

$$\frac{\partial}{\partial \alpha} F(\kappa, \alpha) = 1 - 2(2\varepsilon + N - 2) \frac{\varepsilon^2 + \kappa}{(\varepsilon^2 + \kappa + \alpha)^2} > 0 .$$

Thus  $0 \leq \alpha \mapsto F(\kappa, \alpha)$  is monotone increasing for each  $\kappa > 0$ , and

$$F(\kappa, \alpha) \geq F(\kappa, 0) = (\varepsilon - 1)^2 + N - 1 + \kappa > F(0, 0) = F(0, \alpha_0) ,$$

that implies

$$\inf_{\kappa > 0} \inf_{\nu \in \mathbb{N} \cup \{0\}} F(\kappa, \alpha_\nu) = F(0, \alpha_0) \quad \text{when } \varepsilon < 1 - N/2 .$$

In the case  $\varepsilon \geq 1 - N/2$ , by (31) we see that  $F(\kappa, \alpha)$  is increasing in  $\kappa > 0$  for each  $\alpha \geq 0$ . Thus  $F(\kappa, \alpha) \geq F(0, \alpha)$  and

$$\inf_{\kappa > 0} \inf_{\nu \in \mathbb{N} \cup \{0\}} F(\kappa, \alpha_\nu) = \inf_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu) .$$

To evaluate the right-hand side, we compute

$$\begin{aligned} \frac{\partial}{\partial \alpha} F(0, \alpha) &= 1 - 2(2\varepsilon + N - 2) \frac{\varepsilon^2}{(\varepsilon^2 + \alpha)^2} = \frac{\varepsilon^4 - 4\varepsilon^3 + 2(\alpha - (N - 2))\varepsilon^2 + \alpha^2}{(\varepsilon^2 + \alpha)^2} \\ &\geq \frac{\varepsilon^2(\varepsilon + 2)^2 + \alpha^2}{(\varepsilon^2 + \alpha)^2} > 0 \quad \text{if } \alpha \geq N . \end{aligned}$$

Thus we have  $F(0, \alpha) > F(0, N)$  for any  $\alpha \geq N$ , which implies  $F(0, \alpha_\nu) \geq F(0, \alpha_2) = F(0, 2N)$  for all  $\nu \geq 2$ . This in turn implies

$$\inf_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu) = \min_{\nu \in \{0, 1, 2\}} F(0, \alpha_\nu) .$$

Moreover, by computing

$$\begin{aligned} F(0, \alpha_2) - F(0, \alpha_1) &= F(0, 2N) - F(0, N - 1) \\ &= \frac{(N + 1)\varepsilon^2((\varepsilon - 2)^2 + N - 1) + 2N(N - 1)}{(\varepsilon^2 + N - 1)(\varepsilon^2 + 2N)} > 0 , \end{aligned}$$

we see that

$$\inf_{\nu \in \{0, 1, 2\}} F(0, \alpha_\nu) = \min_{\nu \in \{0, 1\}} F(0, \alpha_\nu) .$$

Therefore, by calculating

$$F(0, \alpha_1) - F(0, \alpha_0) = F(0, N - 1) - F(0, 0) = (N - 1) \frac{(\varepsilon - 2)^2 - (N + 1)}{\varepsilon^2 + N - 1} ,$$

it turns out that

$$(33) \quad \begin{aligned} \inf_{\kappa > 0} \inf_{\nu \in \mathbb{N} \cup \{0\}} F(\kappa, \alpha_\nu) &= \min_{\nu \in \{0, 1\}} F(0, \alpha_\nu) \\ &= \begin{cases} F(0, \alpha_1) & \text{for } (\varepsilon - 2)^2 \leq N + 1 , \\ F(0, \alpha_0) & \text{for } (\varepsilon - 2)^2 > N + 1 \end{cases} \end{aligned}$$

when  $\varepsilon \geq 1 - N/2$ . The expression (33) holds true even for  $\varepsilon < 1 - N/2$  since  $\varepsilon < 1 - N/2$  implies  $(\varepsilon - 2)^2 > N + 1$ .

Inserting this result into (30), we have

$$\begin{aligned} H_{N,\gamma} &\geq \min_{\nu \in \{0,1\}} F(0, \alpha_\nu) \\ &= \begin{cases} F(0, \alpha_1) = (\varepsilon - 1)^2 \frac{\varepsilon^2 + 3(N-1)}{\varepsilon^2 + N - 1} & \text{for } |\varepsilon - 2| \leq \sqrt{N+1}, \\ F(0, \alpha_0) = (\varepsilon - 1)^2 + N - 1 & \text{otherwise.} \end{cases} \end{aligned}$$

Returning to  $\varepsilon = 2 - \frac{N}{2} - \gamma$ , we arrive at the desired infimum value in Theorem 1.

**Optimality for  $H_{N,\gamma}$ .** In this subsection, we prove that the former lower bound of  $H_{N,\gamma}$  is indeed realized as equality:

$$H_{N,\gamma} = \min_{\nu \in \{0,1\}} F(0, \alpha_\nu) = \min_{\nu \in \{0,1\}} \lim_{|\lambda| \searrow +0} \frac{Q_1(\lambda, \alpha_\nu)}{P_1(\lambda, \alpha_\nu)}.$$

For that purpose, let  $\nu_0 \in \{0, 1\}$  be such that

$$\min_{\nu \in \{0,1\}} F(0, \alpha_\nu) = F(0, \alpha_{\nu_0}).$$

By the argument in the last subsection, it is enough to prove that there exists a sequence of vector fields  $\{\mathbf{u}_n\}_{n \in \mathbb{N}} \subset C_c^\infty(\mathbb{R}^N)^N$  such that

$$(34) \quad \lim_{n \rightarrow \infty} \frac{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\nabla \mathbf{u}_n|^2 dx}{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-2} |\mathbf{u}_n|^2 dx} = \lim_{|\lambda| \searrow +0} \frac{Q_1(\lambda, \alpha_{\nu_0})}{P_1(\lambda, \alpha_{\nu_0})}.$$

For the construction of  $\{\mathbf{u}_n\}_{n \in \mathbb{N}}$ , take any nonnegative  $h \in C_c^\infty(\mathbb{R})$ ,  $h \not\equiv 0$  and put  $h_n(t) = h(t/n)$  for  $n \in \mathbb{N}$ . Set

$$(35) \quad \mathbf{v}_n(\rho, \boldsymbol{\sigma}) = \mathbf{e}_\rho (\varepsilon h_n(t) + h'_n(t)) \psi_{\nu_0}(\boldsymbol{\sigma}) + h_n(t) \nabla_{\boldsymbol{\sigma}} \psi_{\nu_0}(\boldsymbol{\sigma})$$

where  $\rho = e^t$  and  $\psi_{\nu_0}$  denotes an eigenfunction of  $-\Delta_{\boldsymbol{\sigma}}$  associated with the eigenvalue  $\alpha_{\nu_0} = \nu_0(\nu_0 + N - 2)$ . Then it is clear that  $\mathbf{v}_n$  satisfies (23). Define

$$(36) \quad \mathbf{u}_n(\rho, \boldsymbol{\sigma}) = \rho^{\varepsilon-1} \mathbf{v}_n(\rho, \boldsymbol{\sigma})$$

for  $\varepsilon = 2 - N/2 - \gamma$ . Then  $\{\mathbf{u}_n\}_{n \in \mathbb{N}}$  is a sequence of curl-free vector fields having compact supports in  $\mathbb{R}^N \setminus \{\mathbf{0}\}$ . Put

$$f_n(\lambda, \boldsymbol{\sigma}) = (\widehat{v_n})_\rho(\lambda, \boldsymbol{\sigma}) = (\varepsilon + i\lambda) \widehat{h_n}(\lambda) \psi_{\nu_0}(\boldsymbol{\sigma})$$

and compute the Hardy-Leray quotient for  $\mathbf{u}_n$  by using (25) and (26). We see

$$\begin{aligned} \frac{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\nabla \mathbf{u}_n|^2 dx}{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-2} |\mathbf{u}_n|^2 dx} &= \frac{\iint_{\mathbb{S}^{N-1} \times \mathbb{R}} \overline{f_n} Q_1(\lambda, -\Delta_{\boldsymbol{\sigma}}) f_n d\boldsymbol{\sigma} d\lambda}{\iint_{\mathbb{S}^{N-1} \times \mathbb{R}} \overline{f_n} P_1(\lambda, -\Delta_{\boldsymbol{\sigma}}) f_n d\boldsymbol{\sigma} d\lambda} \\ &= \frac{\int_{\mathbb{R}} (\varepsilon^2 + \lambda^2) Q_1(\lambda, \alpha_{\nu_0}) |\widehat{h_n}(\lambda)|^2 d\lambda}{\int_{\mathbb{R}} (\varepsilon^2 + \lambda^2) P_1(\lambda, \alpha_{\nu_0}) |\widehat{h_n}(\lambda)|^2 d\lambda} \\ &= \frac{\int_{\mathbb{R}} Q_{01}(\lambda, \alpha_{\nu_0}) |\widehat{h_n}(\lambda)|^2 d\lambda}{\int_{\mathbb{R}} P_{01}(\lambda, \alpha_{\nu_0}) |\widehat{h_n}(\lambda)|^2 d\lambda}, \end{aligned}$$

here

$$(37) \quad \begin{aligned} P_{01}(\lambda, \alpha) &= (\varepsilon^2 + \lambda^2) P_1(\lambda, \alpha) = \varepsilon^2 + \alpha + \lambda^2, \\ Q_{01}(\lambda, \alpha) &= (\varepsilon^2 + \lambda^2) Q_1(\lambda, \alpha) \end{aligned}$$

are polynomials of  $\lambda$ . Note that  $\widehat{h}_n(\lambda) = \widehat{h}(t/n)(\lambda) = n\widehat{h}(n\lambda)$ . Thus if  $\varepsilon^2 + \alpha_{\nu_0} \neq 0$ , then we have

$$\begin{aligned} \frac{\iint_{\mathbb{R}^N} |\nabla \mathbf{u}_n|^2 |x|^{2\gamma} dx}{\iint_{\mathbb{R}^N} |\mathbf{u}_n|^2 |x|^{2\gamma-2} dx} &= \frac{\int_{\mathbb{R}} Q_{01}(\lambda, \alpha_{\nu_0}) |\widehat{h}(n\lambda)|^2 d\lambda}{\int_{\mathbb{R}} P_{01}(\lambda, \alpha_{\nu_0}) |\widehat{h}(n\lambda)|^2 d\lambda} \\ &\rightarrow \frac{Q_{01}(0, \alpha_{\nu_0})}{P_{01}(0, \alpha_{\nu_0})} = \lim_{|\lambda| \rightarrow +0} \frac{Q_1(\lambda, \alpha_{\nu_0})}{P_1(\lambda, \alpha_{\nu_0})} \end{aligned}$$

as  $n \rightarrow \infty$ . In the case  $\varepsilon = 0 = \alpha_{\nu_0}$ , by using

$$P_{01}(\lambda, 0) = \lambda^2, \quad Q_{01}(\lambda, 0) = N\lambda^2 + \lambda^4,$$

we can check that

$$\begin{aligned} \frac{\iint_{\mathbb{R}^N} |\nabla \mathbf{u}_n|^2 |x|^{2\gamma} dx}{\iint_{\mathbb{R}^N} |\mathbf{u}_n|^2 |x|^{2\gamma-2} dx} &= \frac{\int_{\mathbb{R}} Q_{01}(\lambda, 0) |\widehat{h}(n\lambda)|^2 d\lambda}{\int_{\mathbb{R}} P_{01}(\lambda, 0) |\widehat{h}(n\lambda)|^2 d\lambda} = \frac{\int_{\mathbb{R}} (N\lambda^2 + \lambda^4) |\widehat{h}(n\lambda)|^2 d\lambda}{\int_{\mathbb{R}} \lambda^2 |\widehat{h}(n\lambda)|^2 d\lambda} \\ &\rightarrow N = \lim_{|\lambda| \rightarrow +0} \frac{Q_1(\lambda, 0)}{P_1(\lambda, 0)} \end{aligned}$$

as  $n \rightarrow \infty$ . Thus we have proved (34) which shows the optimality of  $H_{N,\gamma}$  in the class of curl-free vector fields in  $C_c^\infty(\mathbb{R}^N)^N$ .  $\square$

#### 4. PROOF OF THEOREM 3

Let  $\gamma \neq 2 - N/2$  be a real number and put  $\varepsilon = 3 - N/2 - \gamma \neq 1$ . Under the transformation  $\mathbf{v} = \rho^{1-\varepsilon} \mathbf{u}$  in (21), the gradient vector field is transformed as

$$\nabla \mathbf{v} = \nabla(\rho^{1-\varepsilon} \mathbf{u}) = (1-\varepsilon)\rho^{-\varepsilon} \mathbf{e}_\rho \otimes \mathbf{u} + \rho^{1-\varepsilon} \nabla \mathbf{u},$$

which leads to

$$(38) \quad |\rho \nabla \mathbf{v}|^2 = (1-\varepsilon)^2 |\rho^{1-\varepsilon} \mathbf{u}|^2 + 2(1-\varepsilon)\rho^{2-2\varepsilon} \mathbf{u} \cdot \rho \partial_\rho \mathbf{u} + \rho^{2-2\varepsilon} |\rho \nabla \mathbf{u}|^2.$$

On the other hand, the assumption  $\int_{\mathbb{R}^N} |\mathbf{x}|^{2-2\varepsilon-N} |\mathbf{u}|^2 dx < \infty$  and the smoothness of  $\mathbf{u}$  imply that

$$\mathbf{u}(\mathbf{x}) = O(|\mathbf{x}|^m), \quad \nabla \mathbf{u}(\mathbf{x}) = O(|\mathbf{x}|^{m-1}) \quad \text{as } |\mathbf{x}| \searrow 0$$

for some integer  $m > \varepsilon - 1$  if  $\varepsilon > 1$ . Therefore, we see that  $\mathbf{v}$  must satisfy

$$(39) \quad |\mathbf{v}(0)| = \lim_{\rho \searrow 0} |\rho \nabla \mathbf{v}| = 0$$

by (38) when  $\varepsilon > 1$ .

Next, we see the  $\Delta \mathbf{u}$  is written in terms of  $\mathbf{v}$  as follows:

$$(40) \quad \Delta \mathbf{u} = \Delta(\rho^{\varepsilon-1} \mathbf{v}) = \rho^{\varepsilon-3} (\alpha_{\varepsilon-1} \mathbf{v} + (2\varepsilon + N - 4) \partial_t \mathbf{v} + \partial_t^2 \mathbf{v} + \Delta_\sigma \mathbf{v}),$$

here we have used (15) and  $\Delta \rho^{\varepsilon-1} = \alpha_{\varepsilon-1} \rho^{\varepsilon-3}$ . Note that  $\iint_{\mathbb{S}^{n-1} \times \mathbb{R}} \partial_t \mathbf{v} \cdot \mathbf{v} d\sigma dt = \iint_{\mathbb{S}^{n-1} \times \mathbb{R}} \partial_t^2 \mathbf{v} \cdot \partial_t \mathbf{v} d\sigma dt = 0$  and  $\iint_{\mathbb{S}^{n-1} \times \mathbb{R}} \mathbf{v} \cdot \partial_t^2 \mathbf{v} d\sigma dt = -\iint_{\mathbb{S}^{n-1} \times \mathbb{R}} |\partial_t \mathbf{v}|^2 d\sigma dt$  by (39). Thus by using (40), Lemma 8, and noting  $(2\varepsilon + N - 4)^2 - 2\alpha_\varepsilon = (N - 2)^2 + 2\alpha_\varepsilon$ ,

we find that the both sides of the Rellich-Leray inequality (4) are written as

$$\begin{aligned}
(41) \quad \int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\Delta \mathbf{u}|^2 dx &= \int_{\mathbb{R}^N} |\mathbf{x}|^{6-2\varepsilon-N} |\Delta \mathbf{u}|^2 dx \\
&= \int_{\mathbb{S}^{N-1}} d\sigma \int_0^\infty |\alpha_{\varepsilon-1} \mathbf{v} + (2\varepsilon + N - 4) \partial_t \mathbf{v} + \partial_t^2 \mathbf{v} + \Delta_\sigma \mathbf{v}|^2 \frac{d\rho}{\rho} \\
&= \int_{\mathbb{S}^{N-1}} d\sigma \int_{\mathbb{R}} \left( \alpha_{\varepsilon-1}^2 |\mathbf{v}|^2 + ((N-2)^2 + 2\alpha_{\varepsilon-1}) |\partial_t \mathbf{v}|^2 + |\partial_t^2 \mathbf{v}|^2 \right. \\
&\quad \left. - 2\alpha_{\varepsilon-1} |\nabla_\sigma \mathbf{v}|^2 + 2|\partial_t \nabla_\sigma \mathbf{v}|^2 + |\Delta_\sigma \mathbf{v}|^2 \right) dt \\
&= \iint_{\mathbb{R} \times \mathbb{S}^{N-1}} \left( \left( \alpha_{\varepsilon-1}^2 + ((N-2)^2 + 2\alpha_{\varepsilon-1}) \lambda^2 + \lambda^4 \right) |\widehat{\mathbf{v}}|^2 \right. \\
&\quad \left. + 2(\lambda^2 - \alpha_{\varepsilon-1}) |\nabla_\sigma \widehat{\mathbf{v}}|^2 + |\Delta_\sigma \widehat{\mathbf{v}}|^2 \right) d\lambda d\sigma \\
&= \iint_{\mathbb{R} \times \mathbb{S}^{N-1}} \overline{f} \left( \left( \alpha_{\varepsilon-1}^2 + ((N-2)^2 + 2\alpha_{\varepsilon-1}) \lambda^2 + \lambda^4 \right) P_1(\lambda, -\Delta_\sigma) \right. \\
&\quad \left. + 2(\lambda^2 - \alpha_{\varepsilon-1}) P_2(\lambda, -\Delta_\sigma) + P_3(\lambda, -\Delta_\sigma) \right) f d\lambda d\sigma,
\end{aligned}$$

and

$$\begin{aligned}
(42) \quad \int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\mathbf{u}|^2 dx &= \int_{\mathbb{R}^N} |\mathbf{x}|^{2-2\varepsilon-N} |\mathbf{u}|^2 dx \\
&= \iint_{\mathbb{R} \times \mathbb{S}^{N-1}} \overline{f} P_1(\lambda, -\Delta_\sigma) f d\lambda d\sigma.
\end{aligned}$$

Therefore, by (41) and (42), the optimal constant in (4) can be expressed as

$$(43) \quad R_{N,\gamma} = \inf_{\mathbf{u} \neq 0, \operatorname{curl} \mathbf{u} = \mathbf{0}} \frac{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\Delta \mathbf{u}|^2 dx}{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\mathbf{u}|^2 dx} = \inf_{f \neq 0} \frac{\iint_{\mathbb{R} \times \mathbb{S}^{N-1}} \overline{f} Q_2(\lambda, -\Delta_\sigma) f d\lambda d\sigma}{\iint_{\mathbb{R} \times \mathbb{S}^{N-1}} \overline{f} P_1(\lambda, -\Delta_\sigma) f d\lambda d\sigma}$$

with the polynomial  $Q_2(\lambda, \alpha)$  given by

$$\begin{aligned}
(44) \quad Q_2(\lambda, \alpha) &= \left( \alpha_{\varepsilon-1}^2 + ((N-2)^2 + 2\alpha_{\varepsilon-1}) \lambda^2 + \lambda^4 \right) P_1(\lambda, \alpha) \\
&\quad + 2(\lambda^2 - \alpha_{\varepsilon-1}) P_2(\lambda, \alpha) + P_3(\lambda, \alpha).
\end{aligned}$$

**Calculation of a lower bound.** As in (29), we expand  $f$  in terms of eigenfunctions of  $-\Delta_\sigma$ . Then by (43), (44), and Lemma 9, we find

$$R_{N,\gamma} \geq \inf_{\nu \in \mathbb{N} \cup \{0\}} \inf_{\lambda \in \mathbb{R} \setminus \{0\}} \frac{Q_2(\lambda, \alpha_\nu)}{P_1(\lambda, \alpha_\nu)} = \inf_{\nu \in \mathbb{N} \cup \{0\}} \inf_{\kappa > 0} F(\kappa, \alpha_\nu),$$

where for  $\kappa > 0$  and  $\alpha \geq 0$ ,  $F(\kappa, \alpha)$  is defined as

$$\begin{aligned}
F(\kappa, \alpha) &= \frac{Q_2(\sqrt{\kappa}, \alpha)}{P_1(\sqrt{\kappa}, \alpha)} \\
&= \alpha_{\varepsilon-1}^2 + ((N-2)^2 + 2\alpha_{\varepsilon-1}) \kappa + \kappa^2 + \frac{2(\kappa - \alpha_{\varepsilon-1}) P_2(\sqrt{\kappa}, \alpha) + P_3(\sqrt{\kappa}, \alpha)}{P_1(\sqrt{\kappa}, \alpha)}.
\end{aligned}$$

By directly calculating further, we can check that

$$F(\kappa, \alpha) = \kappa^2 + \frac{4\alpha(1-\varepsilon)(N+2\varepsilon-2)^2\kappa}{(\varepsilon^2+\alpha)(\kappa+\varepsilon^2+\alpha)} \\ + \left( \frac{N^2}{2} + 2 \left( \varepsilon + \frac{N-4}{2} \right)^2 + 2\alpha \right) \kappa + \frac{(\varepsilon-2)^2+\alpha}{\varepsilon^2+\alpha} (\alpha_\varepsilon - \alpha)^2$$

for  $\varepsilon = 3 - N/2 - \gamma \neq 0$ , and

$$F(\kappa, \alpha) = \kappa^2 + \frac{4(N-2)^2\kappa}{\kappa+\alpha} + ((N-2)^2 + 4 + 2\alpha) \kappa + (4+\alpha)\alpha$$

for  $\varepsilon = 0$ . We also define  $F(0, \alpha)$  as

$$(45) \quad F(0, \alpha) = \lim_{|\lambda| \searrow +0} \frac{Q_2(\lambda, \alpha)}{P_1(\lambda, \alpha)} = \lim_{\kappa \searrow +0} F(\kappa, \alpha) \\ = \begin{cases} \frac{(\varepsilon-2)^2+\alpha}{\varepsilon^2+\alpha} (\alpha_\varepsilon - \alpha)^2, & \text{for } \varepsilon \neq 0, \alpha \geq 0, \\ (4+\alpha)\alpha, & \text{for } \varepsilon = 0, \alpha > 0, \\ 4(N-2)^2, & \text{for } \varepsilon = 0, \alpha = 0. \end{cases}$$

In these settings, from now on we evaluate the expression

$$\inf_{\nu \in \mathbb{N} \cup \{0\}} \inf_{\kappa > 0} F(\kappa, \alpha_\nu).$$

If  $\varepsilon < 1$ , it is clear that the map  $0 < \kappa \mapsto F(\kappa, \alpha)$  is increasing for any fixed  $\alpha \geq 0$ . Also, if  $\varepsilon > 1$ , estimating  $\partial_\kappa F(\kappa, \alpha)$  from below as

$$\frac{\partial F(\kappa, \alpha)}{\partial \kappa} = 2\kappa - \frac{4\alpha(\varepsilon-1)(N+2\varepsilon-2)^2}{(\kappa+\varepsilon^2+\alpha)^2} + \frac{N^2}{2} + 2 \left( \varepsilon + \frac{N-4}{2} \right)^2 + 2\alpha \\ \geq -\frac{4\alpha(\varepsilon-1)(N+2\varepsilon-2)^2}{(\varepsilon^2+\alpha)^2} + \frac{N^2}{2} + 2 \left( \varepsilon + \frac{N-4}{2} \right)^2 + 2\alpha \\ \geq -\frac{\varepsilon-1}{\varepsilon^2} (N+2\varepsilon-2)^2 + \frac{N^2}{2} + 2 \left( \varepsilon + \frac{N-4}{2} \right)^2 + 2\alpha \\ \geq -\frac{1}{4} (N+2\varepsilon-2)^2 + \frac{N^2}{2} + 2 \left( \varepsilon + \frac{N-4}{2} \right)^2 + 2\alpha \\ = \left( \varepsilon + \frac{N}{2} - 3 \right)^2 + \frac{N^2-4}{2} + 2\alpha \geq 0,$$

we see again  $F(\kappa, \alpha)$  is increasing with respect to  $\kappa > 0$  for any  $\alpha \geq 0$ . Therefore we have

$$\inf_{\kappa > 0} F(\kappa, \alpha) = F(0, \alpha)$$

for all  $\varepsilon \neq 1$ , which implies

$$\inf_{\nu \in \mathbb{N} \cup \{0\}} \inf_{\kappa > 0} F(\kappa, \alpha_\nu) = \inf_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu).$$

Moreover, we can check that

$$\frac{\partial F(0, \alpha)}{\partial \alpha} \geq 0, \quad \alpha \geq \max\{\alpha_1, \alpha_\varepsilon\},$$



see Lemma 10. This implies  $\inf_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu)$  is attained. Therefore, we have the desired estimate:

$$(46) \quad R_{N,\gamma} \geq \min_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu) \quad \text{with} \quad F(0, \alpha_\nu) = \lim_{|\lambda| \searrow +0} \frac{Q_2(\lambda, \alpha_\nu)}{P_1(\lambda, \alpha_\nu)}.$$

Furthermore, we see that  $\min_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu)$  is given by

$$\min_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu) = \min_{\nu \in \mathbb{N} \cup \{0\}} \frac{(\varepsilon - 2)^2 + \alpha_\nu}{\varepsilon^2 + \alpha_\nu} (\alpha_\varepsilon - \alpha_\nu)^2$$

for  $\varepsilon = 3 - N/2 - \gamma \neq 0$ , and

$$\begin{aligned} \min_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu) &= \min \left\{ 4(N-2)^2, (4 + \alpha_1)\alpha_1 \right\} \\ &= \begin{cases} 4(N-2)^2 = F(0, \alpha_0) & \text{for } N = 2, 3, 4, \\ (N+3)(N-1) = F(0, \alpha_1) & \text{for } N \geq 5 \end{cases} \end{aligned}$$

for  $\varepsilon = 3 - N/2 - \gamma = 0$ . This gives the lower bound of  $R_{N,\gamma}$ . In the next subsection we will show that the above inequality is indeed the equality.

**Optimality for  $R_{N,\gamma}$ .** To show that the inequality (46) is indeed the equality, let  $\nu_0 \in \mathbb{N} \cup \{0\}$  be such that  $F(0, \alpha_{\nu_0}) = \min_{\nu \in \mathbb{N} \cup \{0\}} F(0, \alpha_\nu)$  is satisfied. We use the sequence of curl-free vector fields  $\{\mathbf{u}_n\}_{n \in \mathbb{N}}$  in (36) again with (35), however for  $\varepsilon = 3 - N/2 - \gamma$ . Then, as in the proof of Theorem 1, we obtain the following expression:

$$\begin{aligned} \frac{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\Delta \mathbf{u}_n|^2 dx}{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\mathbf{u}_n|^2 dx} &= \frac{\iint_{\mathbb{S}^{N-1} \times \mathbb{R}} \overline{f_n} Q_2(\lambda, -\Delta \boldsymbol{\sigma}) f_n d\sigma d\lambda}{\iint_{\mathbb{S}^{N-1} \times \mathbb{R}} \overline{f_n} P_1(\lambda, -\Delta \boldsymbol{\sigma}) f_n d\sigma d\lambda} \\ &= \frac{\int_{\mathbb{R}} (\varepsilon^2 + \lambda^2) Q_2(\lambda, \alpha_{\nu_0}) |\widehat{h}_n(\lambda)|^2 d\lambda}{\int_{\mathbb{R}} (\varepsilon^2 + \lambda^2) P_1(\lambda, \alpha_{\nu_0}) |\widehat{h}_n(\lambda)|^2 d\lambda} \\ &= \frac{\int_{\mathbb{R}} Q_{02}(\lambda, \alpha_{\nu_0}) |\widehat{h}_n(\lambda)|^2 d\lambda}{\int_{\mathbb{R}} P_{01}(\lambda, \alpha_{\nu_0}) |\widehat{h}_n(\lambda)|^2 d\lambda}, \end{aligned}$$

where  $P_{01}(\lambda, \alpha)$  is as in (37) and

$$Q_{02}(\lambda, \alpha) = (\varepsilon^2 + \lambda^2) Q_2(\lambda, \alpha)$$

is a polynomial of  $\lambda$ . When  $\varepsilon = 0$  and  $\alpha_{\nu_0} = 0$ , by using the facts

$$Q_{02}(\lambda, 0) = 4(N-2)^2 \lambda^2 + (N^2 - 4N + 8) \lambda^4 + \lambda^6$$

and  $P_{01}(\lambda, 0) = \lambda^2$ , we prove that

$$\lim_{n \rightarrow \infty} \frac{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\Delta \mathbf{u}_n|^2 dx}{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\mathbf{u}_n|^2 dx} = 4(N-2)^2 = F(0, 0).$$

Thus as in the proof of Theorem 1, we can show that

$$\lim_{n \rightarrow \infty} \frac{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma} |\Delta \mathbf{u}_n|^2 dx}{\int_{\mathbb{R}^N} |\mathbf{x}|^{2\gamma-4} |\mathbf{u}_n|^2 dx} = F(0, \alpha_{\nu_0})$$

for all cases  $\varepsilon^2 + \alpha_{\nu_0} \neq 0$  and  $\varepsilon^2 + \alpha_{\nu_0} = 0$ . This leads to the optimality of  $R_{N,\gamma}$ .  $\square$

## 5. APPENDIX.

In this appendix, we prove technical lemmas.

**Lemma 9.** *Let  $(X, \mathcal{M}, \mu)$  be a measure space and let  $\xi, \eta : X \rightarrow \mathbb{R}$  be a  $\mu$ -measurable function such that  $\xi \neq 0$   $\mu$ -a.e. Suppose  $g : X \rightarrow \mathbb{R}$  is a  $\mu$ -measurable function satisfying,  $\xi g \geq 0$   $\mu$ -a.e.,  $0 < \int_X \xi g d\mu < \infty$ , and  $\int_X |\eta g| d\mu < \infty$ . Then we have*

$$\frac{\int_X \eta g d\mu}{\int_X \xi g d\mu} \geq \operatorname{ess\,inf}_{x \in X} \frac{\eta(x)}{\xi(x)}.$$

*Proof.* Let  $I = \operatorname{ess\,inf}_{x \in X} \frac{\eta(x)}{\xi(x)}$ . Then  $\frac{\eta}{\xi} \geq I$   $\mu$ -a.e. Multiply the both sides by  $\xi g \geq 0$ , we have  $\eta g = \frac{\eta}{\xi} \xi g \geq I \xi g$   $\mu$ -a.e.. By integrating over  $X$ , we obtain

$$\int_X \eta g d\mu \geq I \int_X \xi g d\mu$$

which leads the result.  $\square$

**Lemma 10.** *Let  $F(0, \alpha)$  be defined by (45). Then we have*

$$\frac{\partial F(0, \alpha)}{\partial \alpha} \geq 0 \quad \text{for } \alpha \geq \max\{\alpha_1, \alpha_\varepsilon\}.$$

*Proof.* Recall  $\alpha_1 = N - 1$  and  $\alpha_\varepsilon = \varepsilon(\varepsilon + N - 2)$ . It is enough to show Lemma when  $\varepsilon \neq 0$  and  $F(0, \alpha) = \frac{(\varepsilon-2)^2 + \alpha}{\varepsilon^2 + \alpha}(\alpha_\varepsilon - \alpha)^2$ . A direct computation shows that

$$\begin{aligned} \frac{\partial F(0, \alpha)}{\partial \alpha} &= \frac{2(\alpha - \alpha_\varepsilon)}{(\alpha + \varepsilon^2)^2} f_\varepsilon(\alpha), \quad \text{where} \\ f_\varepsilon(\alpha) &= \alpha^2 + 2(\varepsilon^2 - \varepsilon + 1)\alpha + \varepsilon^2(\varepsilon - 1)^2 + 2\alpha_\varepsilon(1 - \varepsilon). \end{aligned}$$

Since  $\varepsilon^2 - \varepsilon + 1 > 0$  for any  $\varepsilon \in \mathbb{R}$ , we see that  $f_\varepsilon$  is strictly increasing for  $\alpha \geq 0$ . Thus if we show (i)  $f_\varepsilon(\alpha_\varepsilon) \geq 0$  if  $\alpha_\varepsilon \geq \alpha_1$ , and (ii)  $f_\varepsilon(\alpha_1) \geq 0$  if  $\alpha_1 \geq \alpha_\varepsilon$ , then  $f_\varepsilon(\alpha) \geq 0$  for any  $\alpha \geq \max\{\alpha_1, \alpha_\varepsilon\}$ , which concludes Lemma.

To prove (i), we observe that  $f_\varepsilon(\alpha_\varepsilon) = (\alpha_\varepsilon + \varepsilon^2)(\alpha_\varepsilon + (\varepsilon - 2)^2)$ . Thus if  $\alpha_\varepsilon \geq \alpha_1 = N - 1 > 0$ , clearly we have  $f_\varepsilon(\alpha_\varepsilon) > 0$ .

To prove (ii), we observe that  $f_\varepsilon(\alpha_1) = f_\varepsilon(N - 1) = \varepsilon^4 - 6\varepsilon^3 + 8\varepsilon^2 - 2\varepsilon + N^2 - 1$ . We need to prove this quartic function is nonnegative for  $\varepsilon \in \mathbb{R}$  such that  $\alpha_1 \geq \alpha_\varepsilon$ , i.e.,  $-(N - 1) \leq \varepsilon \leq 1$ . However, this is an elementary fact.  $\square$

### Acknowledgments.

The second author (F.T.) was supported by JSPS Grant-in-Aid for Scientific Research (B), No.15H03631.

### REFERENCES

- [1] H. Brezis, and J. L. Vázquez: *Blow-up solutions of some nonlinear elliptic problems*, Rev. Mat. Univ. Complut. Madrid., **10**, (1997), No. 2, 443–469.
- [2] O. Costin, and V. Maz'ya: *Sharp Hardy-Leray inequality for axisymmetric divergence-free fields*, Calculus of Variations and Partial Diff. Eq., **32** (2008), no. 4, 523–532.
- [3] N. Ghoussoub, and A. Moradifam: *Bessel pairs and optimal Hardy and Hardy-Rellich inequalities*, Math. Ann., **349**, (2011), 1–57.
- [4] N. Ghoussoub, and A. Moradifam: *Functional inequalities: new perspectives and new applications*, Mathematical Surveys and Monographs, 187. American Mathematical Society, Providence, RI, 2013. xxiv+299.

- [5] N. Hamamoto: *Rellich-Leray inequality for divergence free vector fields*, In preparation.
- [6] G. H. Hardy, J. E. Littlewood, and G. Polya: *Inequalities*, Cambridge University Press, 1952.
- [7] O. A. Ladyzhenskaya: *The mathematical theory of viscous incompressible flow, Second edition, revised and enlarged*, Mathematics and its Applications, Vol. 2 Gordon and Breach, Science Publishers, New York-London-Paris, (1969).
- [8] J. Leray: *Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l'hydrodynamique*, J. Math. Pures Appl., **12**, 1933. 1–82.
- [9] V. Maz'ya: *Sobolev spaces with applications to elliptic partial differential equations. Second, revised and augmented edition*, Grundlehren der Mathematischen Wissenschaften, **342**. Springer, Heidelberg, 2011. xxviii+866 pp.
- [10] B. Opic, and A. Kufner: *Hardy-type inequalities*, Pitman Res. Notes Math. Ser., **219**, Longman Sci. Tech., Harlow, 1990.
- [11] A. Tertikas, and N. B. Zographopoulos: *Best constants in the Hardy-Rellich inequalities and related improvements*, Adv. Math., **209** 407–459, (2007)

DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, 3-3-138 SUGIMOTO, SUMIYOSHI-KU, OSAKA 558-8585, JAPAN

*E-mail address*: yhjyoe@yahoo.co.jp (N.Hamamoto)

*E-mail address*: futoshi@sci.osaka-cu.ac.jp (F.Takahashi)