

ENERGY CONSERVATION FOR INHOMOGENEOUS INCOMPRESSIBLE AND COMPRESSIBLE EULER EQUATIONS

QUOC-HUNG NGUYEN, PHUOC-TAI NGUYEN, AND BAO QUOC TANG

ABSTRACT. Energy conservations are studied for inhomogeneous incompressible and compressible Euler equations with general pressure law in a torus or a bounded domain. We provide sufficient conditions for a weak solution to conserve the energy. By exploiting a suitable test function, the spatial regularity for the density is only required to be of order $2/3$ in the incompressible case, and of order $1/3$ in the compressible case. When the density is constant, we recover the existing results for classical incompressible Euler equation.

CONTENTS

| | |
|----------------------------------|----|
| 1. Introduction and Main results | 1 |
| 2. Proof of Theorem 1.1 | 7 |
| 2.1. Estimate of (A) | 8 |
| 2.2. Estimate of (B) | 10 |
| 2.3. Estimate of (C) | 12 |
| 2.4. Conclusion | 13 |
| 3. Proof of Theorem 1.3 | 14 |
| 3.1. Estimate of (D) | 14 |
| 3.2. Estimate of (E) | 16 |
| 3.3. Estimate of (F) | 17 |
| 3.4. Estimate of (D3) + (E3) | 18 |
| 3.5. Conclusion | 20 |
| 4. Proof of Theorem 1.5 | 20 |
| 4.1. Conclusion | 24 |
| 5. Proof of Theorem 1.7 | 24 |
| Appendix A. Appendix | 25 |
| References | 27 |

1. INTRODUCTION AND MAIN RESULTS

Let Ω be either \mathbb{T}^d or a bounded and connected domain in \mathbb{R}^d with C^2 boundary $\partial\Omega$, with $d \geq 2$. This paper studies the conservation of energy for weak solutions to the *inhomogeneous*

Date: September 23, 2019.

2010 Mathematics Subject Classification. 35Q31, 76B03.

Key words and phrases. Inhomogeneous incompressible Euler equation; Compressible isentropic Euler equation; Energy conservation; Onsager's conjecture.

incompressible Euler equation

$$\begin{cases} \partial_t \varrho + \nabla \cdot (\varrho \mathbf{u}) = 0, & \text{in } \Omega \times (0, T), \\ \partial_t (\varrho \mathbf{u}) + \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla P = 0, & \text{in } \Omega \times (0, T), \\ \nabla \cdot \mathbf{u} = 0, & \text{in } \Omega \times (0, T), \\ \mathbf{u}(x, t) \cdot n(x) = 0, & \text{on } \partial\Omega \times (0, T), \\ (\varrho \mathbf{u})(x, 0) = \varrho_0(x) \mathbf{u}_0(x), & \text{in } \Omega, \\ \varrho(x, 0) = \varrho_0(x), & \text{in } \Omega, \end{cases} \quad (\text{E})$$

as well as the *compressible Euler equation with general pressure law*

$$\begin{cases} \partial_t \varrho + \nabla \cdot (\varrho \mathbf{u}) = 0, & \text{in } \Omega \times (0, T), \\ \partial_t (\varrho \mathbf{u}) + \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla p(\varrho) = 0, & \text{in } \Omega \times (0, T), \\ \mathbf{u}(x, t) \cdot n(x) = 0, & \text{on } \partial\Omega \times (0, T), \\ (\varrho \mathbf{u})(x, 0) = \varrho_0(x) \mathbf{u}_0(x), & \text{in } \Omega, \\ \varrho(x, 0) = \varrho_0(x), & \text{in } \Omega, \end{cases} \quad (\text{Ec})$$

where $T > 0$ is the time horizon, $n(x)$ denotes the outward unit normal vector field to the boundary $\partial\Omega^1$, $\varrho : \Omega \times (0, T) \rightarrow \mathbb{R}_+$ is the scalar density of a fluid, and $\mathbf{u} : \Omega \times (0, T) \rightarrow \mathbb{R}^d$ denotes its velocity. In system (E), $P : \Omega \times (0, T) \rightarrow \mathbb{R}$ stands for the scalar pressure, while the general pressure law $p : [0, \infty) \rightarrow \mathbb{R}$ in (Ec) satisfies some conditions, which will be specified later. Note that when $p(\varrho) = \varrho^\gamma$ for some $\gamma > 1$, (Ec) is the well-known isentropic compressible Euler equation.

In his celebrated paper [Ons49] Onsager conjectured that there is dissipation of energy for homogeneous Euler equation (namely $\varrho \equiv 1$ in (E)) for weak solutions with low regularity. More precisely, if a weak solution is in C^α for $\alpha > 1/3$ then the energy is conserved while the energy is dissipated if $\alpha < 1/3$. The first landmark result concerning the loss (or gain) of energy is due to Scheffer [Sch93] in which he proved the existence of a weak solution having compact support both in time and space. This was later also recovered by Shnirelman [Shn97] for the torus \mathbb{T}^d . This direction of research has been greatly pushed forward by a series of works of De Lellis and Székelyhidi in e.g. [DS12, DS13, DS14, BDIS15]. The Onsager's conjecture has been recently settled by Isett in [Ise18a, Ise18]. The other direction, i.e. the conservation of energy, was first proved by Constant-E-Titi [CET94] for the torus \mathbb{T}^3 . The case of bounded domains is studied only recently in [BT18, BTW] in the context of Hölder spaces and in [DN18, NN19] in the context of Besov spaces.

Much less works concerning the inhomogeneous incompressible Euler equation (E) and the compressible equation (Ec) have been published in the literature and so far only the case of a bounded domain with periodic boundary condition, namely \mathbb{T}^d , has been treated (see the recent papers [LS16, FGGW17, CY, ADSW]). More precisely, in [FGGW17], Feireisl *et al.* provided sufficient conditions in terms of Besov regularity both in time and space of the density ϱ , the velocity \mathbf{u} and the momentum $\mathbf{m} = \varrho \mathbf{u}$ to guarantee the conservation of the energy. Their method relies on the idea in [CET94] and requires also regularity conditions on the pressure. Recently, by using a different approach, Chen and Yu [CY] obtained the energy conservation, under a different set of regularity conditions. Their method has the advantage of dealing with vacuum. However, since the authors used the convolution both in time and space, additional regularity in time for the density ϱ is required.

In this paper, we provide modest sufficient conditions for a weak solution to conserve the energy for both inhomogeneous incompressible Euler equation (E) and compressible Euler equation (Ec).

¹Naturally, the boundary condition $\mathbf{u} \cdot n = 0$ is neglected when $\Omega = \mathbb{T}^d$.

An attempt to avoid the time regularity of the density is to use the test function $\frac{1}{\varrho^\varepsilon}(\varrho\mathbf{u})^\varepsilon$ instead of \mathbf{u}^ε , where the convolution is *only taken in spatial variable* (see the definition of \mathbf{u}^ε and ϱ^ε in the list of notations). This choice of test functions was used in [LS16] for density-dependent Navier-Stokes equations and in [FGGW17, Remark 4.2] for Euler equations, and it allows to avoid any regularity in time for the density ϱ . However, by way of compensation, one has to exclude vacuum, or at least assume that the inverse density is essentially bounded, i.e. $\varrho^{-1} \in L^\infty(\Omega \times (0, T))$. In this work, we improve this by using the test function $\frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon}(\varrho\mathbf{u})^\varepsilon$ where $\phi_{\eta,\varepsilon}$ is the cut-off function which vanishes when $\varrho^\varepsilon \leq \eta$ (see the definition of $\phi_{\eta,\varepsilon}$ in (30)). This helps us to weaken the condition $\varrho^{-1} \in L^\infty(\Omega \times (0, T))$ to $\varrho^{-1} \in L^{r_1}(\Omega \times (0, T))$ for some $r_1 > 0$, with the price that the density needs to be smoother near vacuum (see e.g. Theorem 1.1). Naturally, in case of no vacuum, these additional regularity assumptions are not any more necessary. Moreover, we are able to reduce the spatial smoothness required for the density. More precisely, $\varrho \in L^\infty(0, T; B_\infty^{\alpha,\infty}(\mathbb{T}^d))$ (with $\alpha = \frac{2}{3}$ for (E) and $\alpha = \frac{1}{3}$ for (Ec)) instead of $\varrho \in L^p(0, T; W^{1,p}(\mathbb{T}^d))$ as in [CY]. Our techniques are also suitable to deal with both the case of a torus or a bounded domain.

To state our results, we need first the definition of weak solutions for (E) and (Ec).

Definition 1.1. *A triple (ϱ, \mathbf{u}, P) is called a weak solution to (E) if*

(i)

$$\int_0^T \int_\Omega (\varrho \partial_t \varphi + \varrho \mathbf{u} \nabla \varphi) dx dt = 0$$

for every test function $\varphi \in C_0^\infty(\Omega \times (0, T))$.

(ii)

$$\int_0^T \int_\Omega (\varrho \mathbf{u} \cdot \partial_t \psi + \varrho \mathbf{u} \otimes \mathbf{u} : \nabla \psi + P \nabla \cdot \psi) dx dt = 0 \quad (1)$$

for every test vector field $\psi \in C_0^\infty(\Omega \times (0, T))^d$.

(iii) $\varrho(\cdot, t) \rightharpoonup \varrho_0$ in $\mathcal{D}'(\Omega)$ as $t \rightarrow 0$, i.e.

$$\lim_{t \rightarrow 0} \int_\Omega \varrho(x, t) \varphi(x) dx = \int_\Omega \varrho_0(x) \varphi(x) dx \quad (2)$$

for every test function $\varphi \in C_0^\infty(\Omega)$.

(iv) $(\varrho \mathbf{u})(\cdot, t) \rightharpoonup \varrho_0 \mathbf{u}_0$ in $\mathcal{D}'(\Omega)$ as $t \rightarrow 0$, i.e.

$$\lim_{t \rightarrow 0} \int_\Omega (\varrho \mathbf{u})(x, t) \psi(x) dx = \int_\Omega (\varrho_0 \mathbf{u}_0)(x) \psi(x) dx \quad (3)$$

for every test vector field $\psi \in C_0^\infty(\Omega)^d$.

A weak solution for (Ec) can be defined in the same way where in (1), the scalar pressure P is replaced by the general pressure law $p(\varrho)$.

For $\beta > 0$, $\delta > 0$ and $p \geq 1$, we define, in the case of the torus \mathbb{T}^d , the quantity²

$$\|f\|_{\mathcal{V}_\delta^{\beta,p}(\mathbb{T}^d)} := \sup_{|h| < \delta} |h|^{-\beta} \|f(\cdot + h) - f(\cdot)\|_{L^p(\mathbb{T}^d)},$$

and in the case of a bounded domain Ω the quantity

$$\|f\|_{\mathcal{V}_\delta^{\beta,p}(K)} := \sup_{|h| < \delta} |h|^{-\beta} \|f(\cdot + h) - f(\cdot)\|_{L^p(K)},$$

²Clearly here is a slight abuse of notation since the definition is not a norm but only a seminorm.

for any $K \subset\subset \Omega$ with $d(K, \partial\Omega) > 2\delta$. For $T > 0$ we denote by

$$\|f\|_{L^q(0,T;\mathcal{V}_\delta^{\beta,p}(\mathbb{T}^d))} = \left(\int_0^T \|f(t)\|_{\mathcal{V}_\delta^{\beta,p}(\mathbb{T}^d)}^q dt \right)^{1/q}, \quad \text{for } 1 \leq q < \infty,$$

$$\|f\|_{L^\infty(0,T;\mathcal{V}_\delta^{\beta,p}(\mathbb{T}^d))} = \operatorname{ess\,sup}_{t \in (0,T)} \|f(t)\|_{\mathcal{V}_\delta^{\beta,p}(\mathbb{T}^d)},$$

and similarly for $\|f\|_{L^q(0,T;\mathcal{V}_\delta^{\beta,p}(K))}$ with $K \subset\subset \Omega$.

For simplicity, we denote by $Q_T = \Omega \times (0, T)$ both for Ω to be \mathbb{T}^d or a bounded domain with boundary, as it should not lead to any confusion.

Our first main result reads as follows.

Theorem 1.1 (Conservation of energy for (E) in a torus). *Let (ϱ, \mathbf{u}, P) be a weak solution to (E) in the case $\Omega = \mathbb{T}^d$. Assume that*

$$\mathbf{u} \in L^3(Q_T) \cap L^3(0, T; \mathcal{V}_{\delta_0}^{\frac{1}{3},3}(\mathbb{T}^d)), \quad \limsup_{\delta \rightarrow 0} \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d))} = 0, \quad P \in L^{\frac{3}{2}}(Q_T), \quad (4)$$

$$0 \leq \varrho \in L^\infty(Q_T) \cap L^\infty(0, T; \mathcal{V}_{\delta_0}^{\frac{2}{3},\infty}(\mathbb{T}^d)), \quad (5)$$

$$\varrho^{-1} \in L^{r_1}(Q_T) \quad \text{and} \quad \mathbf{1}_{\{\varrho \leq \delta_0\}} \partial_t \varrho \in L^{r_2}(Q_T), \quad (6)$$

for some $\delta_0 > 0$ and $r_1 > 0, r_2 \geq 3$ such that $\frac{1}{r_1} + \frac{1}{r_2} = \frac{1}{3}$.

Then the energy for (E) conserves for all time, i.e.

$$\int_\Omega (\varrho |\mathbf{u}|^2)(x, t) dx = \int_\Omega (\varrho_0 |\mathbf{u}_0|^2)(x) dx \quad \forall t \in (0, T). \quad (7)$$

Remark 1.2.

- In the case of no vacuum, i.e. $\varrho(x, t) \geq \underline{\varrho} > 0$, then (6) is obviously satisfied by choosing $\delta_0 < \underline{\varrho}$, and therefore we require only the assumption (5) on the density.
- Though we allow the density ϱ to be zero at some places, the condition $\|\varrho^{-1}\|_{L^{r_1}(Q_T)} < +\infty$ in particular prevents ϱ to be zero in a set with positive measure. Consequently, we have the following useful limit

$$\limsup_{\eta \rightarrow 0} |\{(x, t) \in Q_T : \varrho(x, t) \leq \eta\}| = 0. \quad (8)$$

- It is easy to check that if $f \in W^{\frac{1}{3},3}(\mathbb{T}^d)$, then $f \in \mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)$ for all $\delta > 0$ and

$$\limsup_{\delta \rightarrow 0} \|f\|_{\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)} = 0.$$

Thus, condition (4) holds if $\mathbf{u} \in L^3(0, T; W^{\frac{1}{3},3}(\mathbb{T}^d))$ and $P \in L^{\frac{3}{2}}(Q_T)$.

- When $\varrho \equiv \text{const}$, we recover the classical assumption $\mathbf{u} \in L^3(0, T; W^{\frac{1}{3},3}(\mathbb{T}^d))$ for the homogeneous incompressible Euler equation (see e.g. [CET94]). Note that the condition $P \in L^{\frac{3}{2}}(Q_T)$ is only needed when the density is not constant.

Our results in Theorem 1.1 improve that of [CY] where the authors assume in particular $\varrho \in L^p(0, T; W^{1,p}(\mathbb{T}^d))$ and $\mathbf{u} \in L^q(0, T; B_q^{\alpha,\infty}(\mathbb{T}^d))$ with $\frac{1}{p} + \frac{3}{q} \leq 1$ and $\alpha > \frac{1}{3}$. Here we are able to reduce the spatial regularity of the density ϱ to the order $\frac{2}{3}$ and keep the original order $\frac{1}{3}$ of the velocity. When $\|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\delta^{\alpha,3}(\mathbb{T}^d))} < \infty$ for some $\alpha > \frac{1}{3}$, the limit condition in (4) is automatically satisfied. It's remarked that [CY] also studies the case where ϱ only belongs to $L^\infty((0, T) \times \mathbb{T}^d)$ (where vacuum is allowed to happen), but they need to impose Besov regularity for the velocity both in time and space $\mathbf{u} \in B_p^{\beta,\infty}(0, T; B_q^{\alpha,\infty}(\mathbb{T}^d))$ with $2\alpha + \beta > 1$ and $\alpha + 2\beta > 1$.

In comparison to [LS16] or [FGGW17, Remark 4.2], where the author also used the test function $\frac{1}{\varrho^\varepsilon}(\varrho \mathbf{u})^\varepsilon$, we are able to allow vacuum, though not "too much", i.e. $\varrho^{-1} \in L^1(Q_T)$, by choosing the test function $\frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon}(\varrho \mathbf{u})^\varepsilon$ where the cut-off $\phi_{\eta,\varepsilon}$ vanishes when $\varrho^\varepsilon \leq \eta$. The price to pay is that $\partial_t \varrho$ near vacuum needs to be L^2 -integrable. An integrability of the inverse of the density is also imposed in the recent work [ADSW], in which the authors again require time regularity of both the density and the velocity.

For the case of a bounded domain with smooth boundary Ω , we need additionally some behavior of \mathbf{u} and P near the boundary. In the sequel, $\Omega_r := \{x \in \Omega : d(x, \partial\Omega) > r\}$ for any $r \geq 0$ and $\int_E f dx := \frac{1}{\mathcal{L}^d(E)} \int_E f dx$ for any Borel set $E \subset \mathbb{R}^d$. Since Ω is a bounded, connected domain with C^2 boundary, we find $r_0 > 0$ and a unique C_b^1 -vector function $n : \Omega \setminus \Omega_{r_0} \rightarrow S^{d-1}$ such that the following holds true: for any $r \in [0, r_0)$, $x \in \Omega_r \setminus \Omega_{r_0}$ there exists a unique $x_r \in \partial\Omega_r$ such that $d(x, \partial\Omega_r) = |x - x_r|$ and $n(x)$ is the outward unit normal vector field to the boundary $\partial\Omega_r$ at x_r .

Theorem 1.3 (Conservation of energy for (E) in a bounded domain). *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with C^2 boundary $\partial\Omega$. Let (ϱ, \mathbf{u}, P) be a weak solution to (E). Assume that*

$$0 \leq \varrho \in L^\infty(Q_T), \quad \mathbf{u} \in L^3(Q_T), \quad P \in L^{\frac{3}{2}}(Q_T), \quad (9)$$

$$\|\varrho\|_{L^\infty(0,T;V_{\frac{2}{3},\infty}(\Omega_{2\delta}))} + \|\mathbf{u}\|_{L^3(0,T;V_{\frac{1}{3},3}(\Omega_{2\delta}))} < \infty, \quad \limsup_{\varepsilon \rightarrow 0} \|\mathbf{u}\|_{L^3(0,T;V_{\frac{1}{3},3}(\Omega_\delta))} = 0 \quad \forall \delta > 0, \quad (10)$$

$$\varrho^{-1} \in L^1(Q_T) \quad \text{and} \quad \mathbf{1}_{\{\varrho \leq \delta_0\}} \partial_t \varrho \in L^2(Q_T), \quad (11)$$

for some $\delta_0 > 0$ and $r_1, r_2 > 0$ such that $\frac{1}{r_1} + \frac{1}{r_2} = \frac{1}{3}$. Moreover, we assume the following boundary layer conditions

$$\left(\int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t)|^3 dx dt \right)^{\frac{2}{3}} \left(\int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t) \cdot n(x)|^3 dx dt \right)^{\frac{1}{3}} = o(1) \quad \text{as} \quad \varepsilon \rightarrow 0, \quad (12)$$

$$\left(\int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |P(x,t)|^{\frac{3}{2}} dx dt \right)^{2/3} \left(\int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t) \cdot n(x)|^3 dx dt \right)^{\frac{1}{3}} = o(1) \quad \text{as} \quad \varepsilon \rightarrow 0. \quad (13)$$

Then the energy for (E) conserves for all time, i.e.

$$\int_\Omega (\varrho |\mathbf{u}|^2)(x,t) dx = \int_\Omega (\varrho_0 |\mathbf{u}_0|^2)(x) dx \quad \forall t \in (0, T). \quad (14)$$

Remark 1.4.

- Note that for the case of bounded domain, we only require the density and velocity belong locally to a Besov space.
- When $\varrho \equiv \text{const}$ we recover (and slightly improved) the recent results in [NN19] with the remark that the integrability of the pressure P in (9) is only needed in the inhomogeneous incompressible case.
- Conditions (12) and (13) can be replaced by the following conditions

$$\limsup_{\varepsilon \rightarrow 0} \int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t)|^3 dx dt + \limsup_{\varepsilon \rightarrow 0} \int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |P(x,t)|^{\frac{3}{2}} dx dt < \infty$$

and

$$\liminf_{\varepsilon \rightarrow 0} \int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t) \cdot n(x)|^3 dx dt = 0.$$

- As [NN19], we put $v(x, t) = \mathbf{u}(x, t) \cdot \mathbf{n}(x)$ for any $x \in \Omega \setminus \Omega_{r_0}$. If the function

$$\phi : \varepsilon \mapsto \phi(\varepsilon) = \|v\|_{L^3((\Omega \setminus \Omega_\varepsilon) \times (0, T))}$$

satisfies $\phi(\varepsilon) \leq C\varepsilon^{2/3}$ for every $\varepsilon \in (0, r_0)$ with some $C > 0$, then conditions (12) and (13) are fulfilled.

- The conditions in (10) are satisfied if for each $\delta > 0$, $\varrho \in L^\infty(0, T; W^{\frac{1}{3}, \infty}(\Omega_{2\delta}))$ and $\mathbf{u} \in L^3(0, T; W^{\frac{1}{3}, 3}(\Omega_{2\delta}))$.

We now turn to the energy conservation for the compressible Euler equation (Ec). We will assume the following conditions of the pressure law $p : (0, \infty) \rightarrow \mathbb{R}$,

$$p \in C([0, \infty)) \cap C^2((0, \infty)) \quad \text{and} \quad \limsup_{z \rightarrow 0} \frac{|p(z)| + |p_2(z)|}{|z|} < +\infty, \quad (15)$$

where

$$p_2(z) = z \int_1^z \frac{p(\ell)}{\ell^2} d\ell. \quad (16)$$

We remark that, in the isentropic case, i.e. $p(\varrho) = \varrho^\gamma$ for $\gamma > 1$, the conditions in (15) are obviously satisfied.

Theorem 1.5 (Conservation of energy for (Ec) in a torus). *Let (ϱ, \mathbf{u}) be a weak solution to (Ec) in the case $\Omega = \mathbb{T}^d$. Assume that the pressure law satisfies (15), and the density ϱ and the velocity satisfy*

$$0 \leq \varrho \in L^\infty(Q_T), \quad \mathbf{u} \in L^3(Q_T), \quad (17)$$

$$\|\varrho\|_{L^\infty(0, T; \mathcal{V}_{\delta_0}^{\frac{1}{3}, \infty}(\mathbb{T}^d))} + \|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_{\delta_0}^{\frac{1}{3}, 3}(\mathbb{T}^d))} < \infty, \quad \limsup_{\delta \rightarrow 0} (\|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_{\delta}^{\frac{1}{3}, 3}(\mathbb{T}^d))} + \|\varrho\|_{L^\infty(0, T; \mathcal{V}_{\delta}^{\frac{1}{3}, \infty}(\mathbb{T}^d)}) = 0, \quad (18)$$

$$\mathbf{1}_{\{\varrho \leq \delta_0\}} \partial_t \varrho \in L^3(Q_T), \quad \mathbf{1}_{\{\varrho \leq \delta_0\}} \nabla \varrho \in L^\infty(Q_T) \quad \text{and} \quad \lim_{\eta \rightarrow 0} |\{(x, t) \in Q_T : \varrho(x, t) \leq \eta\}| = 0, \quad (19)$$

for some $\delta_0 > 0$. Then the energy for (Ec) conserves for all time, i.e.

$$\int_{\mathbb{T}^d} \left(\frac{1}{2} (\varrho |\mathbf{u}|^2)(x, t) + p_2(\varrho)(x, t) \right) dx = \int_{\mathbb{T}^d} \left(\frac{1}{2} (\varrho_0 |\mathbf{u}_0|^2)(x) + p_2(\varrho_0)(x) \right) dx \quad \forall t \in (0, T). \quad (20)$$

Remark 1.6. *The conditions (17) and (18) are satisfied if*

$$\varrho \in L^\infty(0, T; W^{\frac{1}{3}, \infty}(\mathbb{T}^d)) \quad \text{and} \quad \mathbf{u} \in L^3(0, T; W^{\frac{1}{3}, 3}(\mathbb{T}^d)).$$

In comparison to the incompressible case in Theorem 1.1, we require in (18) the spatial regularity of ϱ to be *only* $\frac{1}{3}$ instead of $\frac{2}{3}$. This is due to the fact that the general pressure p depends on the density, which allows a finer analysis using smoothness of ϱ itself (see estimate (80)). We are also able to eliminate any condition on ϱ^{-1} , though we have to impose the spatial smoothness of ϱ near vacuum in (19). Yet, we still need to assume that the vacuum set has zero measure. The result of Theorem 1.5 is applicable for a large class of pressure law satisfying (15), which includes the isentropic case $p(\varrho) = \varrho^\gamma$, $\gamma > 1$, as a special situation.

We have the following parallel result for the case of bounded domains.

Theorem 1.7 (Conservation of energy for (Ec) in a bounded domain). *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with C^2 boundary $\partial\Omega$. Let (ϱ, \mathbf{u}) be a weak solution to (Ec). Assume that the pressure law satisfies (15), and the density and the velocity satisfy*

$$0 \leq \varrho \in L^\infty(Q_T), \quad \mathbf{u} \in L^3(Q_T), \quad (21)$$

and for some $\delta_0 > 0$,

$$\|\varrho\|_{L^\infty(0,T;\mathcal{V}_\delta^{\frac{1}{3},\infty}(\Omega_{2\delta}))} + \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\delta^{\frac{1}{3},3}(\Omega_{2\delta}))} < \infty, \quad \limsup_{\varepsilon \rightarrow 0} \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\varepsilon^{\frac{1}{3},3}(\Omega_\delta))} + \|\varrho\|_{L^\infty(0,T;\mathcal{V}_\varepsilon^{\frac{1}{3},\infty}(\Omega_\delta))} = 0, \quad (22)$$

for any $0 < \delta < \delta_0$, and

$$\mathbf{1}_{\{\varrho \leq \delta_0\}} \partial_t \varrho \in L^3(Q_T), \quad \mathbf{1}_{\{\varrho \leq \delta_0\}} \nabla \varrho \in L^\infty(Q_T) \quad \text{and} \quad \lim_{\eta \rightarrow 0} |\{(x,t) \in Q_T : \varrho(x,t) \leq \eta\}| = 0. \quad (23)$$

Moreover,

$$\left(\int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t)|^3 dx dt \right)^{\frac{2}{3}} \left(\int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t) \cdot n(x)|^3 dx dt \right)^{\frac{1}{3}} = o(1) \quad \text{as} \quad \varepsilon \rightarrow 0, \quad (24)$$

$$\int_0^T \int_{\Omega \setminus \Omega_\varepsilon} |\mathbf{u}(x,t) \cdot n(x)| dx dt = o(1) \quad \text{as} \quad \varepsilon \rightarrow 0. \quad (25)$$

Then the energy for (E) conserves for all time, i.e.

$$\int_\Omega \left(\frac{1}{2} (\varrho |\mathbf{u}|^2)(x,t) + p_2(\varrho)(x,t) \right) dx = \int_\Omega \left(\frac{1}{2} (\varrho_0 |\mathbf{u}_0|^2)(x) + p_2(\varrho_0)(x) \right) dx \quad \forall t \in (0,T). \quad (26)$$

The paper is organized as follows: The proofs of Theorems 1.1, 1.3, 1.5 and 1.7 will be presented in the next four Sections respectively. In Appendix A, we collect some useful estimates which will be used in the proofs.

Notation:

- For convenience, we simply write $\|f\|_{L^p}$ for $\|f\|_{L^p(Q_T)}$ for $1 \leq p \leq \infty$, recalling that $Q_T = \Omega \times (0,T)$ where either $\Omega = \mathbb{T}^d$ or $\Omega \subset \mathbb{R}^d$ is a bounded domain with smooth boundary.
- We denote by C a generic constant, whose value can change from line to line or even the same line. Sometimes we write $C(\lambda)$ to emphasize the dependence on $\lambda > 0$.
- For simplicity, we write \limsup for $\limsup_{\varepsilon_n; \dots; \varepsilon_1 \rightarrow 0} \limsup_{\varepsilon_n \rightarrow 0} \dots \limsup_{\varepsilon_1 \rightarrow 0}$.
- Let $\omega : \mathbb{R}^d \rightarrow \mathbb{R}$ be a standard mollifier, i.e. $\omega(x) = c_0 e^{-\frac{1}{1-|x|^2}}$ for $|x| < 1$ and $\omega(x) = 0$ for $|x| \geq 1$, where c_0 is a constant such that $\int_{\mathbb{R}^d} \omega(x) dx = 1$. For any $\varepsilon > 0$, we define the rescaled mollifier $\omega_\varepsilon(x) = \varepsilon^{-d} \omega(\frac{x}{\varepsilon})$. For any function $f \in L^1_{loc}(\Omega)$, its mollified version is defined as

$$f^\varepsilon(x) = (f \star \omega_\varepsilon)(x) = \int_{\mathbb{R}^d} f(x-y) \omega_\varepsilon(y) dy, \quad x \in \Omega_\varepsilon,$$

where $\Omega_\varepsilon = \{x \in \Omega : d(x, \partial\Omega) > \varepsilon\}$. Throughout this paper, we will use the notations $\mathbf{u}^\varepsilon(x,t) = \int_{\mathbb{R}^d} \mathbf{u}(x-y,t) \omega_\varepsilon(y) dy$ and $\varrho^\varepsilon(x,t) = \int_{\mathbb{R}^d} \varrho(x-y,t) \omega_\varepsilon(y) dy$.

2. PROOF OF THEOREM 1.1

In this section we write \mathbb{T}^d instead of Ω . By smoothing (E), we obtain

$$\partial_t \varrho^\varepsilon + \nabla \cdot (\varrho \mathbf{u})^\varepsilon = 0 \quad (27)$$

and

$$\partial_t (\varrho \mathbf{u})^\varepsilon + \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon + \nabla P^\varepsilon = 0. \quad (28)$$

for any $0 < \varepsilon < 1$.

Let $\phi \in C^\infty(\mathbb{R}_+)$ such that

$$0 \leq \phi \leq 1, \quad \phi = 0 \text{ in } [0,1], \quad \phi = 1 \text{ in } [2,\infty) \quad \text{and} \quad |\phi'| + |\phi''| \leq C \text{ in } [0,\infty), \quad (29)$$

for some constant C . For $\eta > 0$, put

$$\phi_{\eta,\varepsilon}(x, t) = \phi\left(\frac{\varrho^\varepsilon(x, t)}{\eta}\right). \quad (30)$$

We see that $\phi_{\eta,\varepsilon} = 0$ in $\{\varrho^\varepsilon \leq \eta\}$, $\phi_{\eta,\varepsilon} = 1$ in $\{\varrho^\varepsilon \geq 2\eta\}$ and $0 \leq \phi_{\eta,\varepsilon} \leq 1$ in Q_T .

Let δ_0 be the positive constant in (4) and (6). Let M be a positive constant such that

$$\|\varrho\|_{L^\infty} + \|\varrho\|_{L^\infty(0,T;\mathcal{V}_{\delta_0}^{\frac{2}{3}}(\mathbb{T}^d))} + \|\varrho^{-1} \mathbf{1}_{\{\varrho \leq \delta_0\}}\|_{L^1} + \|\partial_t \varrho \mathbf{1}_{\{\varrho \leq \delta_0\}}\|_{L^2} \leq M. \quad (31)$$

Take $\varepsilon > 0$ and $\eta > 0$ such that $0 < \varepsilon^{\frac{2}{3}} < \frac{1}{3M}\eta < \frac{1}{12M}\delta_0$.

Multiplying (28) by $\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon}$ then integrating on $(\tau, t) \times \mathbb{T}^d$, for $0 < \tau < t < T$, we get

$$\int_\tau^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \partial_t (\varrho \mathbf{u})^\varepsilon dx ds + \int_\tau^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon dx ds + \int_\tau^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \nabla P^\varepsilon dx ds = 0. \quad (32)$$

Denote by (A), (B) and (C) the terms on the left-hand side of (32) respectively. We will estimate their convergence separately in the next subsections.

2.1. Estimate of (A). We rewrite (A) as, using $\partial_t \varrho^\varepsilon = \nabla \cdot (\varrho \mathbf{u})^\varepsilon$,

$$\begin{aligned} (A) &= \frac{1}{2} \int_\tau^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \partial_t \left(\frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} \right) dx ds - \frac{1}{2} \int_\tau^t \int_{\mathbb{T}^d} \frac{\phi_{\eta,\varepsilon}}{(\varrho^\varepsilon)^2} \nabla \cdot [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] |(\varrho \mathbf{u})^\varepsilon|^2 dx ds \\ &\quad - \frac{1}{2} \int_\tau^t \int_{\mathbb{T}^d} \frac{\phi_{\eta,\varepsilon}}{(\varrho^\varepsilon)^2} \nabla \cdot (\varrho^\varepsilon \mathbf{u}^\varepsilon) |(\varrho \mathbf{u})^\varepsilon|^2 dx ds \\ &=: (A1) + (A2) + (A3). \end{aligned} \quad (33)$$

By integration by parts, we have

$$\begin{aligned} (A1) &= \frac{1}{2} \partial_t \int_\tau^t \int_{\mathbb{T}^d} \left(\phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} \right) (x, s) dx ds - \frac{1}{2} \int_\tau^t \int_{\mathbb{T}^d} \partial_t \phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} dx ds \\ &=: (A11) + (A12). \end{aligned} \quad (34)$$

From (31), Lemma A.3, the fact that $\text{supp } \partial_t \phi_{\eta,\varepsilon} = \text{supp } (\phi'_{\eta,\varepsilon} \partial_t \varrho^\varepsilon) \subset \{\eta \leq \varrho^\varepsilon \leq 2\eta\}$ and Hölder inequality, we obtain

$$\begin{aligned} |(A12)| &\leq \frac{1}{2} \eta^{-1} \int_\tau^t \int_{\mathbb{T}^d} \left| \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \partial_t \varrho^\varepsilon \right| \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} dx ds \\ &\leq C \eta^{-2} \int_\tau^t \int_{\mathbb{T}^d} |\partial_t \varrho^\varepsilon| |(\varrho \mathbf{u})^\varepsilon|^2 \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} dx ds \\ &\leq C \eta^{-2} \|\partial_t \varrho^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^3} \|(\varrho \mathbf{u})^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^3}^2. \end{aligned} \quad (35)$$

By Hölder inequality,

$$\begin{aligned} |\partial_t \varrho^\varepsilon(x, t)| &= \left| \int_{|y| < \varepsilon} \partial_t \varrho(x - y, t) \omega_\varepsilon(y) dy \right| \\ &\leq \left(\int_{|y| < \varepsilon} |\partial_t \varrho(x - y)|^3 dy \right)^{\frac{1}{3}} \left(\int_{|y| < \varepsilon} \omega_\varepsilon(y)^{\frac{3}{2}} dy \right)^{\frac{2}{3}} \\ &\leq C \varepsilon^{-\frac{d}{3}} \left(\int_{|y| < \varepsilon} |\partial_t \varrho(x - y)|^3 dy \right)^{\frac{1}{3}}. \end{aligned}$$

This, together with Lemma A.3 and (31), implies

$$\begin{aligned}
\int_{\mathbb{T}^d} |\partial_t \varrho^\varepsilon(x, t) \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}(x, t)|^3 dx &\leq C\varepsilon^{-d} \int_{|y| < \varepsilon} \int_{\mathbb{T}^d} |\partial_t \varrho(x-y, t) \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}(x, t)|^3 dx dy \\
&\leq C\varepsilon^{-d} \int_{|y| < \varepsilon} \int_{\mathbb{T}^d} |\partial_t \varrho(x-y, t) \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}(x-y, t)|^3 dx dy \\
&= C\varepsilon^{-d} \int_{|y| < \varepsilon} \int_{\mathbb{T}^d} |\partial_t \varrho(x, t) \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}(x, t)|^3 dx dy \\
&\leq CM^3.
\end{aligned} \tag{36}$$

Also, by Hölder's inequality,

$$\begin{aligned}
|(\varrho \mathbf{u})^\varepsilon(x, t)| &= \left| \int_{|y| < \varepsilon} \varrho(x-y, t) \mathbf{u}(x-y) \omega_\varepsilon(y) dy \right| \\
&\leq \left(\int_{|y| < \varepsilon} \varrho(x-y)^3 |\mathbf{u}(x-y)|^3 dy \right)^{\frac{1}{3}} \left(\int_{|y| < \varepsilon} \omega_\varepsilon(y)^{\frac{3}{2}} dy \right)^{\frac{2}{3}} \\
&= C\varepsilon^{-\frac{d}{3}} \left(\int_{|y| < \varepsilon} \varrho(x-y)^3 |\mathbf{u}(x-y)|^3 dy \right)^{\frac{1}{3}}.
\end{aligned}$$

Consequently, by Lemma A.3 (ii),

$$\begin{aligned}
&\int_{\mathbb{T}^d} |(\varrho \mathbf{u})^\varepsilon(x, t) \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}(x, t)|^3 dx \\
&\leq C\varepsilon^{-d} \eta^3 \int_{|y| < \varepsilon} \int_{\mathbb{T}^d} |\mathbf{u}(x-y, t) \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}(x-y, t)|^3 dx dy \\
&\leq C\varepsilon^{-d} \eta^3 \int_{|y| < \varepsilon} \int_{\mathbb{T}^d} |\mathbf{u}(x-y, t) \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}(x-y, t)|^3 dx dy \\
&= C\eta^3 \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3(\mathbb{T}^d)}^3.
\end{aligned} \tag{37}$$

Combining (35), (36) and (37) implies

$$|(A12)| \leq CM \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3(\mathbb{T}^d)}^2. \tag{38}$$

Therefore, by the dominated convergence theorem, we deduce that

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(A12)| = 0.$$

The term (A11) is the desired term. The term (A3) will be canceled with the term (B22) when estimating (B) (see (51)). Therefore, for (A) it remains to estimate (A2).

By integration by parts and Hölder's inequality,

$$|(A2)| \leq \frac{1}{2} \int_\tau^t \left\| \nabla \left(\phi_{\eta, \varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \|((\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon)\|_{L^3(\mathbb{T}^d)} ds. \tag{39}$$

For any $\delta \in (\varepsilon, \delta_0)$, by Lemma A.2 (2), we estimate

$$\|(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon\|_{L^3(\mathbb{T}^d)} \leq C(M) \varepsilon^{\frac{2}{3}} \|\mathbf{u}\|_{V^{\frac{1}{3}, 3}(\mathbb{T}^d)}. \tag{40}$$

On the other hand

$$\left\| \nabla \left(\phi_{\eta, \varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \leq \left\| \nabla \phi_{\eta, \varepsilon} \left(\frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} + \left\| \phi_{\eta, \varepsilon} \nabla \left(\frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)}. \tag{41}$$

From (31), Lemma A.2 (1) and Lemma A.3, we see that

$$\begin{aligned}
\left\| \nabla \phi_{\eta,\varepsilon} \left(\frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} &= \eta^{-1} \left\| \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) |\nabla \varrho^\varepsilon| \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \\
&\leq C\eta^{-3} \|\nabla \varrho^\varepsilon\|_{L^\infty(\mathbb{T}^d)} \|(\varrho \mathbf{u})^\varepsilon\|_{L^3(\mathbb{T}^d)}^2 \\
&\leq CM\eta^{-3} \varepsilon^{-\frac{1}{3}} \|\varrho\|_{\mathcal{V}_\delta^{\frac{2}{3},\infty}(\mathbb{T}^d)} \|\mathbf{u}\|_{L^3(\mathbb{T}^d)}^2 \\
&\leq C(M)\eta^{-3} \varepsilon^{-\frac{1}{3}} \|\mathbf{u}\|_{L^3(\mathbb{T}^d)}^2.
\end{aligned} \tag{42}$$

Next, since $\text{supp } \phi_{\eta,\varepsilon} \subset \{\varrho^\varepsilon \geq \eta\}$, $0 \leq \phi_{\eta,\varepsilon} \leq 1$, by Lemma A.3, together with Lemma A.2, we can estimate

$$\begin{aligned}
\left\| \phi_{\eta,\varepsilon} \nabla \left(\frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} &\leq 2 \left\| \phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon \nabla (\varrho \mathbf{u})^\varepsilon}{(\varrho^\varepsilon)^2} \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} + 2 \left\| \phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2 \nabla \varrho^\varepsilon}{(\varrho^\varepsilon)^3} \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \\
&\leq C\eta^{-2} \|(\varrho \mathbf{u})^\varepsilon\|_{L^3(\mathbb{T}^d)} \|\nabla (\varrho \mathbf{u})^\varepsilon\|_{L^3(\mathbb{T}^d)} + C\eta^{-3} \|\nabla \varrho^\varepsilon\|_{L^\infty(\mathbb{T}^d)} \|(\varrho \mathbf{u})^\varepsilon\|_{L^3(\mathbb{T}^d)}^2 \\
&\leq C(M)\eta^{-2} \varepsilon^{-\frac{2}{3}} \|\mathbf{u}\|_{L^3(\mathbb{T}^d)} (\|\mathbf{u}\|_{\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)} + \|\mathbf{u}\|_{L^3(\mathbb{T}^d)}) + C\eta^{-3} \varepsilon^{-\frac{1}{3}} \|\varrho\|_{\mathcal{V}_\delta^{\frac{2}{3},\infty}(\mathbb{T}^d)} \|\mathbf{u}\|_{L^3(\mathbb{T}^d)}^2.
\end{aligned} \tag{43}$$

Combining (41)–(43) leads to

$$\left\| \nabla \left(\phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \leq C(M) \varepsilon^{-\frac{2}{3}} \eta^{-3} \|\mathbf{u}\|_{L^3(\mathbb{T}^d)} \left(\|\mathbf{u}\|_{\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)} + \|\mathbf{u}\|_{L^3(\mathbb{T}^d)} \right). \tag{44}$$

Inserting (40) and (44) into (39) yields

$$|(A2)| \leq C(M) \eta^{-3} \|\mathbf{u}\|_{L^3} \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d))} \left(\|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d))} + \|\mathbf{u}\|_{L^3} \right). \tag{45}$$

Therefore, by assumption (4),

$$\limsup_{\eta,\delta,\varepsilon;\tau \rightarrow 0} |(A2)| = 0. \tag{46}$$

We remark that, though there is a factor η^{-3} in (45), the limit in (46) is already 0 after taking the limit $\delta \rightarrow 0$.

2.2. Estimate of (B). By integration by parts we have

$$\begin{aligned}
(B) &= - \int_\tau^t \int_{\mathbb{T}^d} [(\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon - (\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon] \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) dx ds \\
&\quad - \int_\tau^t \int_{\mathbb{T}^d} (\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) dx ds \\
&=: (B1) + (B2).
\end{aligned} \tag{47}$$

It can be checked by using Lemma A.2 (2) that

$$\|(\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon - (\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \leq C(M) \varepsilon^{\frac{2}{3}} \left(\|\mathbf{u}\|_{\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)} + \|\mathbf{u}\|_{L^3(\mathbb{T}^d)} \right) \|\mathbf{u}\|_{\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)}.$$

By similar (even simpler) arguments in obtaining (44), we can estimate

$$\left\| \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) \right\|_{L^3(\mathbb{T}^d)} \leq C(M) \eta^{-2} \varepsilon^{-\frac{2}{3}} \left(\|\mathbf{u}\|_{\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)} + \|\mathbf{u}\|_{L^3(\mathbb{T}^d)} \right). \tag{48}$$

Thus, by Hölder's inequality,

$$\begin{aligned} |(B1)| &\leq C \int_{\tau}^t \left\| (\varrho \mathbf{u} \otimes \mathbf{u})^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon} \otimes \mathbf{u}^{\varepsilon} \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \left\| \nabla \cdot \left(\phi_{\eta, \varepsilon} \frac{(\varrho \mathbf{u})^{\varepsilon}}{\varrho^{\varepsilon}} \right) \right\|_{L^3(\mathbb{T}^d)} ds \\ &\leq C(M) \eta^{-2} \left(\|\mathbf{u}\|_{L^3}^2 + \|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_{\delta}^{\frac{1}{3}, 3}(\mathbb{T}^d))}^2 \right) \|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_{\delta}^{\frac{1}{3}, 3}(\mathbb{T}^d))}. \end{aligned} \quad (49)$$

Therefore, by (4),

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(B1)| = 0. \quad (50)$$

We now turn to the estimate of (B2). Using integration by parts we have

$$\begin{aligned} (B2) &= \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \frac{(\varrho \mathbf{u})^{\varepsilon}}{\varrho^{\varepsilon}} \nabla \cdot [(\varrho \mathbf{u})^{\varepsilon} \otimes \mathbf{u}^{\varepsilon}] dx ds = \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \frac{(\varrho \mathbf{u})^{\varepsilon}}{\varrho^{\varepsilon}} [\nabla(\varrho \mathbf{u})^{\varepsilon} \mathbf{u}^{\varepsilon} + (\varrho \mathbf{u})^{\varepsilon} \nabla \mathbf{u}^{\varepsilon}] dx ds \\ &= -\frac{1}{2} \int_{\tau}^t \int_{\mathbb{T}^d} |(\varrho \mathbf{u})^{\varepsilon}|^2 \nabla \cdot \left(\phi_{\eta, \varepsilon} \frac{\mathbf{u}^{\varepsilon}}{\varrho^{\varepsilon}} \right) dx ds + \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \frac{|(\varrho \mathbf{u})^{\varepsilon}|^2}{\varrho^{\varepsilon}} \nabla \cdot \mathbf{u}^{\varepsilon} dx ds \\ &= \int_{\tau}^t \int_{\mathbb{T}^d} \nabla \phi_{\eta, \varepsilon} |(\varrho \mathbf{u})^{\varepsilon}|^2 \frac{\mathbf{u}^{\varepsilon}}{\varrho^{\varepsilon}} dx ds + \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} |(\varrho \mathbf{u})^{\varepsilon}|^2 \left[\frac{1}{\varrho^{\varepsilon}} \nabla \cdot \mathbf{u}^{\varepsilon} - \frac{1}{2} \nabla \cdot \frac{\mathbf{u}^{\varepsilon}}{\varrho^{\varepsilon}} \right] dx ds \\ &= \int_{\tau}^t \int_{\mathbb{T}^d} \nabla \phi_{\eta, \varepsilon} |(\varrho \mathbf{u})^{\varepsilon}|^2 \frac{\mathbf{u}^{\varepsilon}}{\varrho^{\varepsilon}} dx ds + \frac{1}{2} \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \frac{|(\varrho \mathbf{u})^{\varepsilon}|^2}{(\varrho^{\varepsilon})^2} \nabla \cdot (\varrho^{\varepsilon} \mathbf{u}^{\varepsilon}) dx ds \\ &=: (B21) + (B22). \end{aligned} \quad (51)$$

It is remarked that (B22) = -(A3), and therefore we only need to estimate the term (B21). We rewrite (B21), using the divergence free condition, as

$$\begin{aligned} (B21) &= \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \frac{\mathbf{u}^{\varepsilon} \nabla \varrho^{\varepsilon}}{\varrho^{\varepsilon}} |(\varrho \mathbf{u})^{\varepsilon}|^2 dx ds \\ &= \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \frac{\nabla(\varrho^{\varepsilon} \mathbf{u}^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon})}{\varrho^{\varepsilon}} |(\varrho \mathbf{u})^{\varepsilon}|^2 dx ds + \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \frac{\partial_t \varrho^{\varepsilon}}{\varrho^{\varepsilon}} |(\varrho \mathbf{u})^{\varepsilon}|^2 dx ds \\ &= -\eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} (\varrho^{\varepsilon} \mathbf{u}^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon}) \nabla \cdot \left(\phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \frac{|(\varrho \mathbf{u})^{\varepsilon}|^2}{\varrho^{\varepsilon}} \right) dx ds \\ &\quad + \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \frac{\partial_t \varrho^{\varepsilon}}{\varrho^{\varepsilon}} |(\varrho \mathbf{u})^{\varepsilon}|^2 dx ds. \end{aligned}$$

It then follows from Hölder's inequality that

$$\begin{aligned} |(B21)| &\leq \eta^{-1} \int_{\tau}^t \|\varrho^{\varepsilon} \mathbf{u}^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon}\|_{L^3(\mathbb{T}^d)} \left\| \nabla \cdot \left(\phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \frac{|(\varrho \mathbf{u})^{\varepsilon}|^2}{\varrho^{\varepsilon}} \right) \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \\ &\quad + C \eta^{-2} \|\partial_t \varrho^{\varepsilon} \mathbf{1}_{\eta \leq \varrho^{\varepsilon} \leq 2\eta}\|_{L^3} \|(\varrho \mathbf{u})^{\varepsilon} \mathbf{1}_{\eta \leq \varrho^{\varepsilon} \leq 2\eta}\|_{L^3}^2 \\ &=: (B211) + (B212). \end{aligned}$$

The term (B211) can be estimated similarly to (A2) in (39),

$$|(B211)| \leq C(M) \eta^{-3} \|\mathbf{u}\|_{L^3} \|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_{\delta}^{\frac{1}{3}, 3}(\mathbb{T}^d))} \left(\|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_{\delta}^{\frac{1}{3}, 3}(\mathbb{T}^d))} + \|\mathbf{u}\|_{L^3} \right).$$

Therefore, thanks to assumption (4),

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(B211)| = 0.$$

The term (B212) can be estimated similarly to (A12), by using (36) and (37),

$$|(B212)| \leq CM \|\mathbf{u}\mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3(\mathbb{T}^d)}^2.$$

Hence, by the dominated convergence theorem, we obtain

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(B212)| = 0.$$

Remark 2.1. *We remark that we only use the free divergence free condition when estimating (B21), and all the terms related to the pressure. This will be useful in the proof of Theorem 1.5 for compressible Euler equations.*

2.3. **Estimate of (C).** ³ Using the divergence free condition ⁴, we estimate

$$\begin{aligned} (C) &= \int_{\tau}^t \int_{\mathbb{T}^d} \frac{\phi_{\eta, \varepsilon}}{\varrho^{\varepsilon}} ((\varrho \mathbf{u})^{\varepsilon} - \varrho^{\varepsilon} \mathbf{u}^{\varepsilon}) \nabla P^{\varepsilon} dx ds - \int_{\tau}^t \int_{\mathbb{T}^d} \nabla \phi_{\eta, \varepsilon} \mathbf{u}^{\varepsilon} P^{\varepsilon} dx ds \\ &=: (C1) + (C2). \end{aligned} \quad (52)$$

From Lemma A.3 (i), the fact $0 \leq \phi_{\eta, \varepsilon} \leq 1$, Hölder's inequality and Lemma A.2 (2), we get

$$\begin{aligned} |(C1)| &\leq C \eta^{-1} \int_0^T \|(\varrho \mathbf{u})^{\varepsilon} - \varrho^{\varepsilon} \mathbf{u}^{\varepsilon}\|_{L^3(\mathbb{T}^d)} \|\nabla P^{\varepsilon}\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds \\ &\leq C \eta^{-1} \int_0^T \varepsilon^{\frac{2}{3}} \|\varrho\|_{\mathcal{V}_{\delta}^{\frac{2}{3}, \infty}(\mathbb{T}^d)} \varepsilon^{\frac{1}{3}} \|\mathbf{u}\|_{\mathcal{V}_{\delta}^{\frac{1}{3}, 3}(\mathbb{T}^d)} \varepsilon^{-1} \|P\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds \\ &\leq C \eta^{-1} \|\varrho\|_{L^{\infty}(0, T; \mathcal{V}_{\delta}^{\frac{2}{3}, \infty}(\mathbb{T}^d))} \|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_{\delta}^{\frac{1}{3}, 3}(\mathbb{T}^d))} \|P\|_{L^{\frac{3}{2}}}. \end{aligned} \quad (53)$$

By assumption (4), we have

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(C1)| = 0.$$

Next we estimate the term (C2). By integration by parts,

$$\begin{aligned} (C2) &= \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \nabla \varrho^{\varepsilon} \mathbf{u}^{\varepsilon} P^{\varepsilon} dx ds \\ &= \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \nabla (\varrho^{\varepsilon} \mathbf{u}^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon}) P^{\varepsilon} dx ds + \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \partial_t \varrho^{\varepsilon} P^{\varepsilon} dx ds \\ &\leq -\eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} (\varrho^{\varepsilon} \mathbf{u}^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon}) \nabla \cdot \left(\phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) P^{\varepsilon} \right) dx ds \\ &\quad + 2 \int_{\tau}^t \int_{\mathbb{T}^d} \left| \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \frac{\partial_t \varrho^{\varepsilon}}{\varrho^{\varepsilon}} P^{\varepsilon} \mathbf{1}_{\{\eta \leq \varrho^{\varepsilon} \leq 2\eta\}} \right| dx ds. \end{aligned}$$

Hence, using Holder inequality, we deduce

$$\begin{aligned} |(C2)| &\leq \eta^{-2} \int_{\tau}^t \|\varrho^{\varepsilon} \mathbf{u}^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon}\|_{L^3(\mathbb{T}^d)} \left\| \phi'' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \nabla \varrho^{\varepsilon} P^{\varepsilon} \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds \\ &\quad + \eta^{-1} \int_{\tau}^t \|\varrho^{\varepsilon} \mathbf{u}^{\varepsilon} - (\varrho \mathbf{u})^{\varepsilon}\|_{L^3(\mathbb{T}^d)} \left\| \phi' \left(\frac{\varrho^{\varepsilon}}{\eta} \right) \nabla P^{\varepsilon} \right\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds \\ &\quad + C \|(\varrho^{\varepsilon})^{-1} \mathbf{1}_{\{\eta \leq \varrho^{\varepsilon} \leq 2\eta\}}\|_{L^{r_1}} \|\partial_t \varrho^{\varepsilon} \mathbf{1}_{\{\eta \leq \varrho^{\varepsilon} \leq 2\eta\}}\|_{L^{r_2}} \|P^{\varepsilon} \mathbf{1}_{\{\eta \leq \varrho^{\varepsilon} \leq 2\eta\}}\|_{L^{\frac{3}{2}}} \end{aligned}$$

³When $\varrho \equiv \text{const}$, the term (C) vanishes, and therefore the assumption $P \in L^{\frac{3}{2}}(Q_T)$ is not needed.

⁴We observe that except for the term (B21), the divergence free is only used in the proof of Theorem 1.1 to treat the term involving the pressure. This observation will become helpful when treating compressible Euler equation in the next sections.

$$=: (C21) + (C22) + (C23),$$

where r_1, r_2 are in (6). Since ϕ'' is bounded, due to Lemma A.2 (2), we get

$$\begin{aligned} |(C21)| &\leq C\eta^{-2} \int_{\tau}^t \varepsilon^{\frac{2}{3}} \|\varrho\|_{\mathcal{V}_{\delta}^{\frac{2}{3},\infty}(\mathbb{T}^d)} \varepsilon^{\frac{1}{3}} \|\mathbf{u}\|_{\mathcal{V}_{\delta}^{\frac{1}{3},3}(\mathbb{T}^d)} \varepsilon^{-\frac{1}{3}} \|\varrho\|_{\mathcal{V}_{\delta}^{\frac{2}{3},\infty}(\mathbb{T}^d)} \|P\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} dt \\ &\leq C\eta^{-2} \varepsilon^{\frac{2}{3}} \|\varrho\|_{L^{\infty}(0,T;\mathcal{V}_{\delta}^{\frac{2}{3},\infty}(\mathbb{T}^d))}^2 \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_{\delta}^{\frac{1}{3},3}(\mathbb{T}^d))} \|P\|_{L^{\frac{3}{2}}}. \end{aligned}$$

Thus

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} |(C21)| = 0.$$

The term (C22) can be estimated similarly to (C1) in (53) using the fact that ϕ' is uniformly bounded (29), and thus we get

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} |(C22)| = 0.$$

To estimate (C23), we observe, due to Lemma A.3, that

$$\|(\varrho^{\varepsilon})^{-1} \mathbf{1}_{\{\eta \leq \varrho^{\varepsilon} \leq 2\eta\}}\|_{L^{r_1}} \leq C \left\| \varrho^{-1} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}} \right\|_{L^{r_1}} \leq C \|\varrho^{-1} \mathbf{1}_{\{\varrho \leq \delta_0\}}\|_{L^{r_1}} \leq CM \quad (54)$$

and (by using the argument leading to (36) and replacing 3 by r_2)

$$\|\partial_t \varrho^{\varepsilon} \mathbf{1}_{\{\eta \leq \varrho^{\varepsilon} \leq 2\eta\}}\|_{L^{r_2}} \leq CM. \quad (55)$$

Similarly

$$\|P^{\varepsilon} \mathbf{1}_{\{\eta \leq \varrho^{\varepsilon} \leq 2\eta\}}\|_{L^{\frac{3}{2}}} \leq C \|P \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^{\frac{3}{2}}}. \quad (56)$$

From the above estimates, we obtain

$$(C23) \leq C(M) \|P \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^{\frac{3}{2}}}.$$

By dominated convergence theorem,

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} |(C23)| = 0.$$

Collecting the above estimates, we eventually deduce that

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} |(C)| = 0.$$

Remark 2.2. *The condition $\|\varrho^{-1}\|_{L^{r_1}} < +\infty$ is only needed to control (C23) resulting from the pressure term. In the compressible equation, this condition is not needed. Removing this technical condition in the incompressible case remains as an interesting open problem.*

2.4. Conclusion. From the previous estimates we have

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} \left| \int_{\tau}^t \int_{\mathbb{T}^d} \partial_t \left[\phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^{\varepsilon}|^2}{\varrho^{\varepsilon}} \right] (x, s) dx ds \right| = 0.$$

By (2) and (3), $\varrho(\cdot, \tau) \rightarrow \varrho_0$ and $(\varrho \mathbf{u})(\cdot, \tau) \rightarrow \varrho_0 \mathbf{u}_0$ in $\mathcal{D}'(\Omega)$, so for every $x \in \mathbb{T}^d$,

$$\varrho^{\varepsilon}(x, \tau) \rightarrow \varrho_0^{\varepsilon}(x), \quad (\varrho \mathbf{u})^{\varepsilon}(x, \tau) \rightarrow (\varrho_0 \mathbf{u}_0)^{\varepsilon}(x),$$

as $\tau \rightarrow 0$. Thanks to dominated convergence theorem, it yields

$$\limsup_{\eta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \left| \int_{\mathbb{T}^d} \left(\phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^{\varepsilon}|^2}{\varrho^{\varepsilon}} \right) (x, t) dx - \int_{\mathbb{T}^d} \left(\phi_{\eta,\varepsilon} \frac{|(\varrho_0 \mathbf{u}_0)^{\varepsilon}|^2}{\varrho_0^{\varepsilon}} \right) (x) dx \right| = 0.$$

Thus, by the standard property of convolution, we derive

$$\limsup_{\eta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \left| \int_{\mathbb{T}^d} (\phi_{\eta,\varepsilon} \varrho |\mathbf{u}|^2) (x, t) dx - \int_{\mathbb{T}^d} (\phi_{\eta,\varepsilon} \varrho_0 |\mathbf{u}_0|^2) (x) dx \right| = 0.$$

This implies (7) by using the dominated convergence theorem. The proof is complete. \square

3. PROOF OF THEOREM 1.3

The proof of Theorem 1.3 is similar to that of Theorem 1.1, except that we have to take care of the boundary layers when taking integration by parts. Recalling the smooth version of (E) as

$$\partial_t \varrho^\varepsilon + \nabla \cdot (\varrho \mathbf{u})^\varepsilon = 0 \quad \text{in} \quad \Omega_{2\varepsilon}, \quad (57)$$

and

$$\partial_t (\varrho \mathbf{u})^\varepsilon + \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon + \nabla P^\varepsilon = 0 \quad \text{in} \quad \Omega_{2\varepsilon}. \quad (58)$$

Take $0 < \varepsilon < \varepsilon_1/10 < \varepsilon_2/10 < \delta_0/100$. Note that if $\varepsilon_2 \rightarrow 0$ then $\varepsilon, \varepsilon_1 \rightarrow 0$. Multiplying (58) by $\frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon$ then integrating on $(\tau, t) \times \Omega_{\varepsilon_2}$, with $0 < \tau < t < T$, yield

$$\begin{aligned} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \partial_t (\varrho \mathbf{u})^\varepsilon dx ds + \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon dx ds \\ + \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla P^\varepsilon dx ds = 0. \end{aligned} \quad (59)$$

For $\varepsilon_3 > 0$ small, we integrate (59) with respect to ε_2 on $(\varepsilon_1, \varepsilon_1 + \varepsilon_3)$ to get

$$\begin{aligned} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \partial_t (\varrho \mathbf{u})^\varepsilon dx ds d\varepsilon_2 + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon dx ds d\varepsilon_2 \\ + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla P^\varepsilon dx ds d\varepsilon_2 = 0. \end{aligned} \quad (60)$$

Denote by (D), (E) and (F) the three terms on the left-hand side of (60), which will be estimated separately in the following subsections. Let M_{ε_1} be a constant such that

$$\|\varrho\|_{L^\infty} + \|\varrho\|_{L^\infty(0,T;V_{\frac{2}{\varepsilon_1/4},\infty}(\Omega_{\varepsilon_1/2}))} + \|\varrho^{-1} \mathbf{1}_{\{\varrho \leq \delta_0\}}\|_{L^{r_1}} + \|\partial_t \varrho \mathbf{1}_{\{\varrho \leq \delta_0\}}\|_{L^{r_2}} \leq M_{\varepsilon_1}.$$

It's worth mentioning that in the following, we let successively $\tau \rightarrow 0$, $\varepsilon \rightarrow 0$, $\varepsilon_1 \rightarrow 0$, $\varepsilon_3 \rightarrow 0$ and then $\eta \rightarrow 0$. Therefore, at some estimates, after letting $\varepsilon \rightarrow 0$, constants depending on M_{ε_1} (usually denoted by $C(M_{\varepsilon_1})$) vanishes for each $\varepsilon_1 > 0$, therefore we will be able to send $\varepsilon_1 \rightarrow 0$ afterwards without encountering any issue.

3.1. Estimate of (D). The term (D) is rewritten as

$$\begin{aligned} (D) &= \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \phi_{\eta,\varepsilon} \partial_t \left(\frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} \right) dx ds d\varepsilon_2 \\ &\quad - \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{(\varrho^\varepsilon)^2} \nabla \cdot [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] |(\varrho \mathbf{u})^\varepsilon|^2 dx ds d\varepsilon_2 \\ &\quad - \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{1}{(\varrho^\varepsilon)^2} \nabla \cdot (\varrho^\varepsilon \mathbf{u}^\varepsilon) |(\varrho \mathbf{u})^\varepsilon|^2 dx ds d\varepsilon_2 \\ &=: (D1) + (D2) + (D3). \end{aligned} \quad (61)$$

For (D1) we have

$$(D1) = \frac{1}{2} \frac{1}{\varepsilon_3} \int_\tau^t \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_{\Omega_{\varepsilon_2}} \partial_t \left[\phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon}(x, s) \right] dx ds d\varepsilon_2$$

$$\begin{aligned}
& -\frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \partial_t \phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} dx ds d\varepsilon_2 \\
& =: (D11) + (D12).
\end{aligned}$$

Since (D11) is the desired term, we only estimate (D12). Using similar arguments to estimate (A12), we have

$$\begin{aligned}
|(D12)| & \leq C\eta^{-2} \int_{\tau}^t \int_{\Omega} |\partial_t \varrho^\varepsilon| |(\varrho \mathbf{u})^\varepsilon|^2 \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} dx ds \\
& \leq C\eta^{-2} \|\partial_t \varrho^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^3} \|(\varrho \mathbf{u})^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^3}^2 \\
& \leq C \|\partial_t \varrho \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3} \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3}^2,
\end{aligned}$$

thus

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(D12)| = 0.$$

It remains to estimate (D2) since (D3) will be estimated together with (E3) later (see (73)). Using integration by parts we have

$$\begin{aligned}
|(D2)| & \leq \frac{1}{2} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\partial\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{(\varrho^\varepsilon)^2} |(\varrho \mathbf{u})^\varepsilon|^2 [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] \cdot n(\theta) d\mathcal{H}^{d-1}(\theta) ds d\varepsilon_2 \right| \\
& + \frac{1}{2} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] \nabla \left(\phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) dx ds d\varepsilon_2 \right| \\
& =: (D21) + (D22).
\end{aligned}$$

The term (D22) can be estimated similarly to (A2) as

$$\begin{aligned}
(D22) & \leq \frac{1}{2} \int_{\tau}^t \int_{\Omega_{\varepsilon_1}} \left| [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] \nabla \left(\phi_{\eta,\varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{(\varrho^\varepsilon)^2} \right) \right| dx ds \\
& \leq C(M_{\varepsilon_1}) \eta^{-3} \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_{\varepsilon}^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))} \left[\|\mathbf{u}\|_{L^3}^2 + \|\mathbf{u}\|_{L^3} \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_{\varepsilon}^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))} \right].
\end{aligned} \tag{62}$$

Therefore, by using (10),

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} (D22) = 0. \tag{63}$$

For (D21) we use the coarea formula: for any $0 < r_1 < r_2 < 1$, and $g \in L^1(\Omega_{r_1} \setminus \Omega_{r_2})$,

$$\int_{\Omega_{r_1} \setminus \Omega_{r_2}} g(x) dx = \int_{r_1}^{r_2} \int_{\partial\Omega_\nu} g(\theta) d\mathcal{H}^{d-1}(\theta) d\nu, \tag{64}$$

and $\varepsilon_3 \approx \mathcal{L}^d(\Omega_{\varepsilon_1} \setminus \Omega_{\varepsilon_1+\varepsilon_3})$, and $\phi_{\eta,\varepsilon} = 0$ if $\varrho^\varepsilon \leq \eta$, to get

$$\begin{aligned}
(D21) & \leq C \left| \int_{\tau}^t \int_{\Omega_{\varepsilon_1} \setminus \Omega_{\varepsilon_1+\varepsilon_3}} \frac{\phi_{\eta,\varepsilon}}{(\varrho^\varepsilon)^2} |(\varrho \mathbf{u})^\varepsilon|^2 [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] \cdot n(x) dx ds \right| \\
& \leq C\eta^{-2} \left| \int_{\tau}^t \int_{\Omega_{\varepsilon_1} \setminus \Omega_{\varepsilon_1+\varepsilon_3}} |(\varrho \mathbf{u})^\varepsilon|^2 [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] \cdot n(x) dx ds \right|.
\end{aligned}$$

Since $\varrho \in L^\infty(Q_T)$ and $\mathbf{u} \in L^3(Q_T)$, we have

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(D21)| = 0. \tag{65}$$

3.2. **Estimate of (E) .** Using integration by parts we have

$$\begin{aligned}
(E) &= \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\partial\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon n(\theta) d\mathcal{H}^{d-1}(\theta) ds d\varepsilon_2 \\
&\quad - \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) dx ds d\varepsilon_2 \\
&=: (E1) - \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} [(\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon - (\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon] \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) dx ds d\varepsilon_2 \\
&\quad - \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} (\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) dx ds d\varepsilon_2 \\
&=: (E1) + (E2) + (E3).
\end{aligned} \tag{66}$$

By the coarea formula (64), we have

$$(E1) = \frac{1}{\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_1} \setminus \Omega_{\varepsilon_1+\varepsilon_3}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon n(x) dx ds.$$

Therefore, letting successively $\tau \rightarrow 0$, $\varepsilon \rightarrow 0$ and $\varepsilon_1 \rightarrow 0$, then using the fact $\varepsilon_3 \approx \mathcal{L}^d(\Omega \setminus \Omega_{\varepsilon_3})$, $\phi_{\eta,\varepsilon} = 0$ for $\varrho^\varepsilon \leq \eta$, and Hölder's inequality, we obtain

$$\begin{aligned}
\limsup_{\varepsilon_1; \varepsilon; \tau \rightarrow 0} |(E1)| &\leq \frac{C(\|\varrho\|_{L^\infty})}{\varepsilon_3} \eta^{-1} \int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u}|^2 |\mathbf{u} \cdot n| dx ds \\
&\leq C(\|\varrho\|_{L^\infty}) \eta^{-1} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u}|^3 dx ds \right)^{\frac{2}{3}} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u} \cdot n|^3 dx ds \right)^{\frac{1}{3}}.
\end{aligned}$$

Thus, by assumption (12), we derive

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(E1)| = 0.$$

By proceeding as in estimating (B1) we derive

$$\begin{aligned}
|(E2)| &\leq C \int_{\tau}^t \int_{\Omega_{\varepsilon_1}} |(\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon - (\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon| \left| \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) \right| dx ds \\
&\leq C \int_0^T \|(\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon - (\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon\|_{L^{\frac{3}{2}}(\Omega_{\varepsilon_1})} \left\| \nabla \cdot \left(\phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \right) \right\|_{L^3(\Omega_{\varepsilon_1})} ds \\
&\leq C(M_{\varepsilon_1}) \eta^{-2} \left(\|\mathbf{u}\|_{L^3}^2 + \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_\varepsilon^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))}^2 \right) \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_\varepsilon^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))}.
\end{aligned} \tag{67}$$

Therefore, by assumption (10), it follows that

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(E2)| = 0. \tag{68}$$

3.3. Estimate of (F) . By using the divergence free condition we can split (F) as

$$\begin{aligned}
(F) &= \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} ((\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon) \nabla P^\varepsilon dx ds d\varepsilon_2 + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \phi_{\eta,\varepsilon} \mathbf{u}^\varepsilon \nabla P^\varepsilon dx ds d\varepsilon_2 \\
&=: (F1) + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\partial\Omega_{\varepsilon_2}} \phi_{\eta,\varepsilon} P^\varepsilon \mathbf{u}^\varepsilon \cdot n(\theta) d\mathcal{H}^{d-1}(\theta) ds d\varepsilon_2 \\
&\quad - \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \nabla \phi_{\eta,\varepsilon} \mathbf{u}^\varepsilon P^\varepsilon dx ds d\varepsilon_2 \\
&=: (F1) + (F2) + (F3).
\end{aligned} \tag{69}$$

The term $(F1)$ is estimated similarly to the term $(C1)$ in the case $\Omega = \mathbb{T}^d$, thus by Lemma A.1 (2), we obtain

$$\begin{aligned}
|(F1)| &\leq \int_0^T \int_{\Omega_{\varepsilon_1}} \left| \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} \right| |(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon| |\nabla P^\varepsilon| dx ds \\
&\leq C \eta^{-1} \|\varrho\|_{L^\infty(0,T; \mathcal{V}_{\frac{3}{4}}^{\frac{2}{3},\infty}(\Omega_{\frac{\varepsilon_1}{2}}))} \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_{\frac{1}{2}}^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))} \|P\|_{L^{\frac{3}{2}}}.
\end{aligned} \tag{70}$$

Hence, by assumption (10),

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(F1)| = 0. \tag{71}$$

For the term $(F2)$ we use the coarea formula (64) to obtain

$$(F2) = \frac{1}{\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_1} \setminus \Omega_{\varepsilon_1+\varepsilon_3}} \phi_{\eta,\varepsilon} P^\varepsilon \mathbf{u}^\varepsilon \cdot n(x) dx ds.$$

By using the fact $\varepsilon_3 \approx \mathcal{L}^d(\Omega \setminus \Omega_{\varepsilon_3})$ and Hölder's inequality, we obtain

$$\begin{aligned}
\limsup_{\varepsilon_1; \varepsilon; \tau \rightarrow 0} |(F2)| &= \left| \frac{1}{\varepsilon_3} \int_{\tau}^t \int_{\Omega \setminus \Omega_{\varepsilon_3}} \phi_{\eta} P(x, s) \mathbf{u}(x, s) \cdot n(x) dx ds \right| \\
&\leq C \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |P(x, s)|^{\frac{3}{2}} dx ds \right)^{\frac{2}{3}} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u}(x, s) \cdot n(x)|^3 dx ds \right)^{\frac{1}{3}}.
\end{aligned}$$

By the assumption (13) we get

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(F2)| = 0. \tag{72}$$

It remains to estimate $(F3)$. We use similar arguments in the estimation of $(C2)$. More precisely,

$$\begin{aligned}
|(F3)| &\leq \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \nabla (\varrho^\varepsilon \mathbf{u}^\varepsilon - (\varrho \mathbf{u})^\varepsilon) P^\varepsilon dx ds d\varepsilon_2 \right| \\
&\quad + \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \partial_t \varrho^\varepsilon P^\varepsilon dx ds d\varepsilon_2 \right| \\
&\leq \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_0^t \int_{\partial\Omega_{\varepsilon_2}} \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) (\varrho^\varepsilon \mathbf{u}^\varepsilon - (\varrho \mathbf{u})^\varepsilon) P^\varepsilon d\mathcal{H}^{d-1}(\theta) dt d\varepsilon_2 \right| \\
&\quad + \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} (\varrho^\varepsilon \mathbf{u}^\varepsilon - (\varrho \mathbf{u})^\varepsilon) \left[\phi'' \left(\frac{\varrho^\varepsilon}{\eta} \right) \frac{\nabla \varrho^\varepsilon}{\eta} P^\varepsilon + \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \nabla P^\varepsilon \right] dx ds d\varepsilon_2 \right|
\end{aligned}$$

$$\begin{aligned}
& + \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \partial_t \varrho^\varepsilon P^\varepsilon dx ds d\varepsilon_2 \right| \\
& =: (F31) + (F32) + (F33).
\end{aligned}$$

We estimate (F32) similarly to (C1) and (C21) to get

$$\begin{aligned}
|(F32)| & \leq C \eta^{-1} \|\varrho\|_{L^\infty(0,T;\mathcal{V}_\varepsilon^{\frac{2}{3},\infty}(\Omega_{\frac{\varepsilon_1}{2}}))} \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\varepsilon^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))} \|P\|_{L^{\frac{3}{2}}} \\
& \quad + C \eta^{-2} \varepsilon^{\frac{2}{3}} \|\varrho\|_{L^\infty(0,T;\mathcal{V}_\varepsilon^{\frac{2}{3},\infty}(\Omega_{\frac{\varepsilon_1}{2}}))}^2 \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\varepsilon^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))} \|P\|_{L^{\frac{3}{2}}},
\end{aligned}$$

and thus

$$\limsup_{\eta;\varepsilon_3;\varepsilon_1;\varepsilon;\tau \rightarrow 0} |(F32)| = 0.$$

Using similar arguments in estimating (F2), taking into account that $\varrho \in L^\infty(Q_T)$, we have

$$\limsup_{\varepsilon_1;\varepsilon;\tau \rightarrow 0} |(F31)| \leq C \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |P(x,s)|^{\frac{3}{2}} dx ds \right)^{\frac{2}{3}} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u}(x,s) \cdot n(x)|^3 dx ds \right)^{\frac{1}{3}}$$

and therefore, thanks to assumption (13)

$$\limsup_{\eta;\varepsilon_3;\varepsilon_1;\varepsilon;\tau \rightarrow 0} |(F31)| = 0.$$

Finally, (F33) is estimated similarly to (C23) as

$$|(F33)| \leq C \|\varrho^{-1} \mathbf{1}_{\{\varrho \leq \frac{\varepsilon}{3}\eta}\} \|_{L^1} \|\partial_t \varrho \mathbf{1}_{\{\varrho \leq \frac{\varepsilon}{3}\eta}\} \|_{L^2} \|P \mathbf{1}_{\{\varrho \leq \frac{\varepsilon}{3}\eta}\} \|_{L^{\frac{3}{2}}},$$

thus

$$\limsup_{\eta;\varepsilon_3;\varepsilon_1;\varepsilon;\tau \rightarrow 0} |(F33)| = 0.$$

3.4. Estimate of (D3)+(E3). Using integration by parts we compute (similarly to (A3)+(B2) = 0)

$$\begin{aligned}
(E3) & = -\frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\partial\Omega_{\varepsilon_2}} \phi_{\eta,\varepsilon}(\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \cdot n(\theta) d\mathcal{H}^{d-1}(\theta) ds d\varepsilon_2 \\
& \quad + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} \nabla \cdot [(\varrho \mathbf{u})^\varepsilon \otimes \mathbf{u}^\varepsilon] dx ds d\varepsilon_2 \\
& =: (E31) + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \phi_{\eta,\varepsilon} \frac{(\varrho \mathbf{u})^\varepsilon}{\varrho^\varepsilon} [\nabla \cdot (\varrho \mathbf{u})^\varepsilon \mathbf{u}^\varepsilon + (\varrho \mathbf{u})^\varepsilon \nabla \cdot \mathbf{u}^\varepsilon] dx ds d\varepsilon_2 \\
& = (E31) + \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\partial\Omega_{\varepsilon_2}} \phi_{\eta,\varepsilon} |(\varrho \mathbf{u})^\varepsilon|^2 \frac{\mathbf{u}^\varepsilon}{\varrho^\varepsilon} \cdot n(\theta) d\mathcal{H}^{d-1}(\theta) ds d\varepsilon_2 \\
& \quad - \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} |(\varrho \mathbf{u})^\varepsilon|^2 \nabla \cdot \left(\frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} \mathbf{u}^\varepsilon \right) dx ds d\varepsilon_2 \\
& \quad + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} |(\varrho \mathbf{u})^\varepsilon|^2 \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} \nabla \cdot \mathbf{u}^\varepsilon dx ds d\varepsilon_2 \\
& =: (E31) + (E32) - \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} |(\varrho \mathbf{u})^\varepsilon|^2 \nabla \phi_{\eta,\varepsilon} \frac{\mathbf{u}^\varepsilon}{\varrho^\varepsilon} dx ds d\varepsilon_2
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} |(\varrho \mathbf{u})^\varepsilon|^2 \frac{\nabla \cdot (\varrho^\varepsilon \mathbf{u}^\varepsilon)}{(\varrho^\varepsilon)^2} dx ds d\varepsilon_2 \\
& =: (E31) + (E32) - (E33) - (D3).
\end{aligned} \tag{73}$$

Therefore, it remains only to estimate (E31), (E32), and (E33). By using a similar argument as in estimating (E1), together with the coarea formula, Hölder's inequality and the fact $\varepsilon_3 \approx \mathcal{L}^d(\Omega \setminus \Omega_{\varepsilon_3})$, we obtain

$$\limsup_{\varepsilon_1; \varepsilon; \tau \rightarrow 0} (|(E31)| + |(E32)|) \leq C \eta^{-1} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u}|^3 dx ds \right)^{\frac{2}{3}} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u} \cdot n|^3 dx ds \right)^{\frac{1}{3}}.$$

By assumption (12), we deduce

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} (|(E31)| + |(E32)|) = 0. \tag{74}$$

For (E33), similar arguments used in estimating (B21) can be reused

$$\begin{aligned}
|(E33)| & \leq \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} (\varrho^\varepsilon \mathbf{u}^\varepsilon - (\varrho \mathbf{u})^\varepsilon) \nabla \cdot \left(\phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} \right) dx ds d\varepsilon_2 \right| \\
& + \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \frac{\partial_t \varrho^\varepsilon}{\varrho^\varepsilon} |(\varrho \mathbf{u})^\varepsilon|^2 dx ds d\varepsilon_2 \right| \\
& + \eta^{-1} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_0^t \int_{\partial \Omega_{\varepsilon_2}} (\varrho^\varepsilon \mathbf{u}^\varepsilon - (\varrho \mathbf{u})^\varepsilon) \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon} d\mathcal{H}^{d-1}(\theta) dt d\varepsilon_2 \right| \\
& =: (E331) + (E332) + (E333).
\end{aligned}$$

The terms (E331) and (E332) are estimated similarly to (B211) and (B212), respectively,

$$|(E331)| \leq C(M_{\varepsilon_1}) \eta^{-3} \|\mathbf{u}\|_{L^3} \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_\varepsilon^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))} (\|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_\varepsilon^{\frac{1}{3},3}(\Omega_{\frac{\varepsilon_1}{2}}))} + \|\mathbf{u}\|_{L^3})$$

and

$$|(E332)| \leq C \|\partial_t \varrho \mathbf{1}_{\{\varrho \leq \delta_0\}}\|_{L^3} \|\mathbf{u} \mathbf{1}_{\{\varrho \leq \frac{8}{3}\eta}\|_{L^3}^2,$$

while the term (E333) is estimated similarly to (E1)

$$\limsup_{\varepsilon_1; \varepsilon; \tau \rightarrow 0} |(E333)| \leq C \eta^{-2} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u}|^3 dx ds \right)^{\frac{2}{3}} \left(\int_0^T \int_{\Omega \setminus \Omega_{\varepsilon_3}} |\mathbf{u} \cdot n|^3 dx ds \right)^{\frac{1}{3}}.$$

Combining these estimates one gets

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(E33)| = 0. \tag{75}$$

It then follows from (73), (74) and (75) that

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(D3) + (E3)| = 0.$$

3.5. **Conclusion.** From the estimate of (D), (E) and (F) we have

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} \left| \frac{1}{2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \partial_t \left[\phi_{\eta, \varepsilon} \frac{|(\varrho \mathbf{u})^\varepsilon|^2}{\varrho^\varepsilon}(x, s) \right] dx ds \varepsilon_2 \right| = 0.$$

Arguing similarly to the case of torus $\Omega = \mathbb{T}^d$ we obtain finally the results of Theorem 1.3,

$$\int_{\Omega} (\varrho |\mathbf{u}|^2)(x, t) dx = \int_{\Omega} (\varrho_0 |\mathbf{u}_0|^2)(x) dx \quad \forall t \in (0, T).$$

4. PROOF OF THEOREM 1.5

From the proof of Theorem 1.1, we have for any $0 < \tau < t < T$,

$$\begin{aligned} \int_{\tau}^t \int_{\mathbb{T}^d} \frac{\phi_{\eta, \varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \partial_t (\varrho \mathbf{u})^\varepsilon dx ds + \int_{\tau}^t \int_{\mathbb{T}^d} \frac{\phi_{\eta, \varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon dx ds \\ + \int_{\tau}^t \int_{\mathbb{T}^d} \frac{\phi_{\eta, \varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla (p(\varrho))^\varepsilon dx ds = 0. \end{aligned} \quad (76)$$

Let M be a constant such that

$$\|\varrho\|_{L^\infty} + \|\varrho\|_{L^\infty(0, T; \mathcal{V}_{\delta_0}^{\frac{1}{3}, \infty}(\mathbb{T}^d))} \leq M.$$

In view of the proof of Theorem 1.1 and Remark 2.1, we only need to estimate the term (B21) (now without the divergence free condition) and the last term on the left-hand side of (76).

For (B21) we have

$$\begin{aligned} |(B21)| &= \left| \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \nabla \varrho^\varepsilon \cdot \frac{\mathbf{u}^\varepsilon}{\varrho^\varepsilon} |(\varrho \mathbf{u})^\varepsilon|^2 dx ds \right| \\ &\leq \eta^{-2} \|\mathbf{u}^\varepsilon\|_{L^1} \| |(\varrho \mathbf{u})^\varepsilon|^2 \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} \|_{L^1} \|\nabla \varrho^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^\infty} \\ &\leq C \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3}^3 \|\nabla \varrho \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^\infty}. \end{aligned}$$

Therefore, thanks to (19),

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(B21)| = 0.$$

It therefore remains to estimate the last term on the left-hand side of (76). Put

$$p_1(z) := \int_1^z \frac{p'(\ell)}{\ell} d\ell. \quad (77)$$

We can then estimate

$$\begin{aligned} (G) &:= \int_{\tau}^t \int_{\mathbb{T}^d} \frac{\phi_{\eta, \varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla (p(\varrho))^\varepsilon dx ds \\ &= \int_{\tau}^t \int_{\mathbb{T}^d} \frac{\phi_{\eta, \varepsilon}}{\varrho^\varepsilon} [(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon] \nabla (p(\varrho))^\varepsilon dx ds + \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \mathbf{u}^\varepsilon \nabla (p(\varrho))^\varepsilon dx ds \\ &=: (G1) + \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \mathbf{u}^\varepsilon \nabla [(p(\varrho))^\varepsilon - p(\varrho^\varepsilon)] dx ds + \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \mathbf{u}^\varepsilon p'(\varrho^\varepsilon) \nabla \varrho^\varepsilon dx ds \\ &=: (G1) + (G2) + \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} \mathbf{u}^\varepsilon \varrho^\varepsilon \underbrace{\frac{p'(\varrho^\varepsilon)}{\varrho^\varepsilon} \nabla \varrho^\varepsilon}_{=\nabla p_1(\varrho^\varepsilon)} dx ds \\ &= (G1) + (G2) + \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} [\mathbf{u}^\varepsilon \varrho^\varepsilon - (\varrho \mathbf{u})^\varepsilon] \nabla p_1(\varrho^\varepsilon) dx ds - \int_{\tau}^t \int_{\mathbb{T}^d} p_1(\varrho^\varepsilon) \nabla (\phi_{\eta, \varepsilon} (\varrho \mathbf{u})^\varepsilon) dx ds \end{aligned}$$

$$\begin{aligned}
& =: (G1) + (G2) + (G3) - \int_{\tau}^t \int_{\mathbb{T}^d} p_1(\varrho^\varepsilon)(\varrho \mathbf{u})^\varepsilon \nabla \phi_{\eta, \varepsilon} dx ds - \int_{\tau}^t \int_{\mathbb{T}^d} p_1(\varrho^\varepsilon) \phi_{\eta, \varepsilon} \nabla \cdot (\varrho \mathbf{u})^\varepsilon dx ds \\
& =: (G1) + (G2) + (G3) - (G4) - (G5).
\end{aligned} \tag{78}$$

To estimate (G1), we observe from the fact $\int_{\mathbb{R}^d} \nabla \omega_\varepsilon(y) dy = 0$ and Hölder's inequality that

$$|\nabla(p(\varrho(x, t))^\varepsilon)| \leq C\varepsilon^{-1-\frac{2d}{3}} \left(\int_{|y|<\varepsilon} |p(\varrho(x-y, t)) - p(\varrho(x))|^{\frac{3}{2}} dy \right)^{\frac{2}{3}}.$$

It follows that

$$\begin{aligned}
& \int_{\mathbb{T}^d} |\nabla(p(\varrho(x, t))^\varepsilon)|^{\frac{3}{2}} \mathbf{1}_{\{\varrho \geq \eta\}}(x, t) dx \\
& \leq C\varepsilon^{-\frac{3}{2}-d} \int_{|y|<\varepsilon} \int_{\mathbb{T}^d} |p(\varrho(x-y, t)) - p(\varrho(x, t))|^{\frac{3}{2}} \mathbf{1}_{\{\varrho \geq \eta\}}(x, t) dx dy \\
& = C\varepsilon^{-\frac{3}{2}-d} \int_{|y|<\varepsilon} \int_{\mathbb{T}^d} |p'(\xi)|^{\frac{3}{2}} |\varrho(x-y, t) - \varrho(x, t)|^{\frac{3}{2}} \mathbf{1}_{\{\varrho \geq \eta\}}(x, t) dx dy,
\end{aligned}$$

where ξ is between $\varrho(x)$ and $\varrho(x-y, t)$. Since $\varrho \in L^\infty(0, T; \mathcal{V}_\delta^{\frac{1}{3}, \infty}(\mathbb{T}^d))$, if $\varrho(x, t) \geq \eta$ then $\varrho(x-y, t) > \frac{2}{3}\eta$ for any $|y| < \varepsilon$ (see (89)). Therefore $\xi \in [\frac{2}{3}\eta, \|\varrho\|_{L^\infty}]$ and hence $|p'(\xi)|$ is bounded in $\{\varrho \geq \eta\}$. Consequently,

$$\begin{aligned}
& \int_{\mathbb{T}^d} |\nabla(p(\varrho(x, t))^\varepsilon)|^{\frac{3}{2}} \mathbf{1}_{\{\varrho \geq \eta\}}(x, t) dx \\
& \leq C(\eta)\varepsilon^{-\frac{3}{2}-d} \int_{|y|<\varepsilon} \int_{\mathbb{T}^d} |\varrho(x-y, t) - \varrho(x)|^{\frac{3}{2}} \mathbf{1}_{\{\varrho \geq \eta\}}(x, t) dx dy \\
& \leq C(\eta)\varepsilon^{-\frac{3}{2}-d} \int_{|y|<\varepsilon} |y|^{\frac{1}{2}} \left(|y|^{-\frac{1}{3}} \|\varrho(x-y, t) - \varrho(x, t)\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \right)^{\frac{3}{2}} dy.
\end{aligned}$$

This implies

$$\|\nabla(p(\varrho(x, t))^\varepsilon) \mathbf{1}_{\{\varrho \geq \eta\}}\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \leq C(\eta)\varepsilon^{-\frac{2}{3}} \|\varrho(t)\|_{\mathcal{V}_\delta^{\frac{1}{3}, \frac{3}{2}}(\mathbb{T}^d)}. \tag{79}$$

Using (79) and arguments in estimating (C1) in (53), we have

$$\begin{aligned}
|(G1)| & \leq \int_{\tau}^t \int_{\mathbb{T}^d} \left| \frac{\phi_{\eta, \varepsilon}}{\varrho^\varepsilon} \right| |(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon| |\nabla(p(\varrho))^\varepsilon| dx ds \\
& \leq C\eta^{-1} \int_0^T \|(\varrho \mathbf{u})^\varepsilon - \varrho^\varepsilon \mathbf{u}^\varepsilon\|_{L^3(\mathbb{T}^d)} \|\nabla(p(\varrho))^\varepsilon \mathbf{1}_{\{\varrho \geq \eta\}}\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds \\
& \leq C(\eta)\eta^{-1} \int_0^T \varepsilon^{\frac{2}{3}} \|\varrho\|_{\mathcal{V}_\delta^{\frac{1}{3}, \infty}(\mathbb{T}^d)} \|\mathbf{u}\|_{\mathcal{V}_\delta^{\frac{1}{3}, 3}(\mathbb{T}^d)} \varepsilon^{-\frac{2}{3}} \|\varrho\|_{\mathcal{V}_\delta^{\frac{1}{3}, \frac{3}{2}}(\mathbb{T}^d)} ds \\
& \leq C(M, \eta)\eta^{-1} \|\mathbf{u}\|_{L^3(0, T; \mathcal{V}_\delta^{\frac{1}{3}, 3}(\mathbb{T}^d))}.
\end{aligned} \tag{80}$$

Therefore,

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(G1)| = 0.$$

Similarly,

$$\begin{aligned}
|(G3)| & \leq \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta, \varepsilon} |\varrho^\varepsilon \mathbf{u}^\varepsilon - (\varrho \mathbf{u})^\varepsilon| \frac{|p'(\varrho^\varepsilon)|}{\varrho^\varepsilon} |\nabla \varrho^\varepsilon| dx ds \\
& \leq C(\eta)\eta^{-1} \int_0^T \|\varrho^\varepsilon \mathbf{u}^\varepsilon - (\varrho \mathbf{u})^\varepsilon\|_{L^3(\mathbb{T}^d)} \|\nabla \varrho^\varepsilon\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds
\end{aligned}$$

$$\begin{aligned}
&\leq C(\eta)\eta^{-1} \int_0^T \varepsilon^{\frac{2}{3}} \|\varrho\|_{\mathcal{V}_\delta^{\frac{1}{3},\infty}(\mathbb{T}^d)} \|\mathbf{u}\|_{\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d)} \varepsilon^{-\frac{2}{3}} \|\varrho\|_{\mathcal{V}_\delta^{\frac{1}{3},\frac{3}{2}}(\mathbb{T}^d)} ds \\
&\leq C(\eta, M)\eta^{-1} \|\mathbf{u}\|_{L^3(0,T;\mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d))}.
\end{aligned}$$

Therefore

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} |(G3)| = 0.$$

For (G4), we use (37) and (19) to obtain

$$\begin{aligned}
|(G4)| &\leq \int_\tau^t \int_{\mathbb{T}^d} \left| p_1(\varrho^\varepsilon) (\varrho \mathbf{u})^\varepsilon \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \frac{\nabla \varrho^\varepsilon}{\eta} \right| dx ds \\
&\leq C\eta^{-1} \|p_1(\varrho^\varepsilon) \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^\infty} \|(\varrho \mathbf{u})^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^1} \|\nabla \varrho^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^\infty} \\
&\leq C \|p_1(\varrho^\varepsilon) \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^\infty} \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta}\|_{L^1}.
\end{aligned}$$

We notice that

$$p_1(\varrho^\varepsilon) = \frac{p(\varrho^\varepsilon) + p_2(\varrho^\varepsilon)}{\varrho^\varepsilon} - p(1),$$

hence by (15), $\|p_1(\varrho^\varepsilon) \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^\infty} < C$, where C is independent of ε and η . Therefore,

$$|(G4)| \leq C \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta}\|_{L^1}$$

and consequently,

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} |(G4)| = 0.$$

We use $\partial_t \varrho^\varepsilon = \nabla \cdot (\varrho \mathbf{u})^\varepsilon$ and the fact that

$$\partial_t p_2(\varrho^\varepsilon) = \partial_t \varrho^\varepsilon (p_1(\varrho^\varepsilon) + p(1))$$

(recalling p_2 defined in (16)) to estimate (G5),

$$\begin{aligned}
(G5) &= \int_\tau^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \partial_t p_2(\varrho^\varepsilon) dx ds - p(1) \int_\tau^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \partial_t \varrho^\varepsilon dx ds \\
&= \int_\tau^t \int_{\mathbb{T}^d} \partial_t (\phi_{\eta,\varepsilon} p_2(\varrho^\varepsilon))(x, s) dx ds - \int_\tau^t \int_{\mathbb{T}^d} p_2(\varrho^\varepsilon) \partial_t \phi_{\eta,\varepsilon} dx ds - p(1) \int_\tau^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \partial_t \varrho^\varepsilon dx ds \\
&=: (G51) - (G52) - (G53).
\end{aligned}$$

Since (G51) is a desired term, we only need to estimate (G52) and (G53). We have

$$\begin{aligned}
|(G52)| &\leq \eta^{-1} \int_\tau^t \int_{\mathbb{T}^d} \left| p_2(\varrho^\varepsilon) \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \partial_t \varrho^\varepsilon \right| dx ds \\
&\leq C \left\| \left| \frac{p_2(\varrho^\varepsilon)}{\varrho^\varepsilon} \right| \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} \partial_t \varrho^\varepsilon \right\|_{L^1} \\
&\leq C \|\mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} \partial_t \varrho^\varepsilon\|_{L^1} \\
&\leq C \|\mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} \partial_t \varrho^\varepsilon\|_{L^3} |\{(x, t) \in Q_T : \eta \leq \varrho^\varepsilon \leq 2\eta\}|^{\frac{2}{3}} \\
&\leq CM |\{(x, t) \in Q_T : \eta/3 \leq \varrho \leq 8\eta/3\}|^{\frac{2}{3}}.
\end{aligned}$$

for small enough $\eta > 0$. Here we have used (15) for the third estimate, Hölder's inequality for the fourth estimate and (36) for the last estimate. Thus, thanks to (19),

$$\limsup_{\eta;\delta;\varepsilon;\tau \rightarrow 0} |(G52)| = 0.$$

Next we estimate (G53). By integration by parts,

$$\begin{aligned}
(G53) &= p(1) \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \partial_t \varrho^\varepsilon dx ds \\
&= p(1) \int_{\tau}^t \int_{\mathbb{T}^d} (\varrho \mathbf{u})^\varepsilon \nabla \phi_{\eta,\varepsilon} dx ds \\
&= \eta^{-1} p(1) \int_{\tau}^t \int_{\mathbb{T}^d} (\varrho \mathbf{u})^\varepsilon \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \nabla \varrho^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} dx ds
\end{aligned}$$

Therefore, by Hölder inequality, (37) and (19),

$$\begin{aligned}
|(G53)| &\leq C \eta^{-1} \int_{\tau}^t \int_{\mathbb{T}^d} |(\varrho \mathbf{u})^\varepsilon| |\nabla \varrho^\varepsilon| \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}} dx ds \\
&\leq C \eta^{-1} \|(\varrho \mathbf{u})^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^3} \|\nabla \varrho^\varepsilon \mathbf{1}_{\{\eta \leq \varrho^\varepsilon \leq 2\eta\}}\|_{L^{\frac{3}{2}}} \\
&\leq CM \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3} \|\nabla \varrho \mathbf{1}_{\{\varrho \leq \delta_0\}}\|_{L^{\frac{3}{2}}} \\
&\leq CM \|\mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta\}}\|_{L^3}.
\end{aligned}$$

It follows that

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(G53)| = 0.$$

It therefore remains to estimate (G2). Integration by parts leads to

$$\begin{aligned}
(G2) &= - \int_{\tau}^t \int_{\mathbb{T}^d} \nabla \phi_{\eta,\varepsilon} \mathbf{u}^\varepsilon [(p(\varrho))^\varepsilon - p(\varrho^\varepsilon)] dx ds - \int_{\tau}^t \int_{\mathbb{T}^d} \phi_{\eta,\varepsilon} \nabla \mathbf{u}^\varepsilon [(p(\varrho))^\varepsilon - p(\varrho^\varepsilon)] dx ds \\
&=: (G21) + (G22).
\end{aligned}$$

We will show that

$$\|((p(\varrho))^\varepsilon - p(\varrho^\varepsilon)) \mathbf{1}_{\{\eta \leq \varrho^\varepsilon\}}\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \leq C(\eta) \varepsilon \quad (81)$$

where the constant $C(\eta)$ might blow up as $\eta \rightarrow 0$. Indeed, first, thanks to Taylor's expansion and the continuity of p'' , we have

$$|p(a+b) - p(a) - p'(a)b| = |p''(\xi)| |b|^2 \leq \Lambda(\eta) |b|^2 \quad (82)$$

for all a, b such that $\frac{1}{3}\eta \leq a, a+b \leq \|\varrho\|_{L^\infty}$, where $\Lambda(\eta) = \sup\{|p''(\xi)| : \frac{1}{3}\eta \leq \xi \leq \|\varrho\|_{L^\infty}\}$. Now by writing

$$\varrho(x-y, s) = \varrho(x, s) + [\varrho(x-y, s) - \varrho(x, s)]$$

and

$$\int_{\mathbb{T}^d} \varrho(x-y, s) \omega_\varepsilon(y) dy = \varrho(x, s) + \int_{\mathbb{T}^d} (\varrho(x-y, s) - \varrho(x, s)) \omega_\varepsilon(y) dy$$

we can apply (82) with $a = \varrho(x, s)$ and $b = \varrho(x-y, s) - \varrho(x, s)$, and then $a = \varrho(x, s)$ and $b = \int_{\mathbb{T}^d} (\varrho(x-y, s) - \varrho(x, s)) \omega_\varepsilon(y) dy$ respectively, to obtain

$$\begin{aligned}
&|(p(\varrho))^\varepsilon(x, s) - p(\varrho^\varepsilon)(x, s)| \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho\}}(x, s) \\
&\leq \left| \int_{\mathbb{T}^d} [p(\varrho(x-y, s)) - p(\varrho(x, s)) - p'(\varrho(x, s))(\varrho(x-y, s) - \varrho(x, s))] \omega_\varepsilon(y) dy \right| \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho\}}(x, s) \\
&+ \left| p \left(\int_{\mathbb{T}^d} \varrho(x-y, s) \omega_\varepsilon(y) dy \right) - p(\varrho(x, s)) - p'(\varrho(x, s)) \int_{\mathbb{T}^d} (\varrho(x-y, s) - \varrho(x, s)) \omega_\varepsilon(y) dy \right| \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho\}}(x, s) \\
&\leq 2\Lambda(\eta) \int_{\mathbb{T}^d} |\varrho(x-y, s) - \varrho(x, s)|^2 \omega_\varepsilon(y) dy \\
&\leq C(\eta) \varepsilon^{\frac{2}{3}}.
\end{aligned}$$

Therefore,

$$\|((p(\varrho))^\varepsilon(\cdot, s) - p(\varrho^\varepsilon)(\cdot, s))\mathbf{1}_{\{\eta \leq \varrho^\varepsilon\}}\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} \leq C(\eta)\varepsilon^{\frac{2}{3}}.$$

Using (81) we now can estimate

$$\begin{aligned} |(G21)| &\leq \eta^{-1} \int_\tau^t \int_{\mathbb{T}^d} \left| \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \nabla \varrho^\varepsilon \mathbf{u}^\varepsilon [(p(\varrho))^\varepsilon - p(\varrho^\varepsilon)] \right| dx ds \\ &\leq C\eta^{-1} \int_0^T \|\nabla \varrho^\varepsilon\|_{L^\infty(\mathbb{T}^d)} \|\mathbf{u}^\varepsilon\|_{L^3(\mathbb{T}^d)} \|((p(\varrho))^\varepsilon - p(\varrho^\varepsilon))\mathbf{1}_{\{\eta \leq \varrho^\varepsilon\}}\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds \\ &\leq C\eta^{-1} \varepsilon^{-\frac{2}{3}} \|\varrho\|_{L^\infty(0,T; \mathcal{V}_\delta^{\frac{1}{3},\infty}(\mathbb{T}^d))} \|\mathbf{u}\|_{L^3} C(\eta)\varepsilon^{\frac{2}{3}} \\ &\leq C(\eta)\eta^{-1} \|\varrho\|_{L^\infty(0,T; \mathcal{V}_\delta^{\frac{1}{3},\infty}(\mathbb{T}^d))} \|\mathbf{u}\|_{L^3} \end{aligned}$$

and thus by (18),

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(G21)| = 0.$$

Similarly

$$\begin{aligned} |(G22)| &\leq \int_\tau^t \int_{\mathbb{T}^d} |\nabla \mathbf{u}^\varepsilon| \mathbf{1}_{\{\eta \leq \varrho^\varepsilon\}} |(p(\varrho))^\varepsilon - p(\varrho^\varepsilon)| dx ds \\ &\leq \int_0^T \|\nabla \mathbf{u}^\varepsilon\|_{L^3(\mathbb{T}^d)} \|\mathbf{1}_{\{\eta \leq \varrho^\varepsilon\}} [(p(\varrho))^\varepsilon - p(\varrho^\varepsilon)]\|_{L^{\frac{3}{2}}(\mathbb{T}^d)} ds \\ &\leq C\varepsilon^{-\frac{2}{3}} \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d))} C\Lambda(\eta)\varepsilon^{\frac{2}{3}} \\ &\leq C\Lambda(\eta) \|\mathbf{u}\|_{L^3(0,T; \mathcal{V}_\delta^{\frac{1}{3},3}(\mathbb{T}^d))}, \end{aligned}$$

hence by (18),

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} |(G22)| = 0.$$

4.1. Conclusion. From the previous estimates we have

$$\limsup_{\eta; \delta; \varepsilon; \tau \rightarrow 0} \left| \int_\tau^t \int_{\mathbb{T}^d} \partial_t \left[\frac{1}{2} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} |(\varrho \mathbf{u})^\varepsilon|^2 + \phi_{\eta,\varepsilon} p_2(\varrho^\varepsilon) \right] dx ds \right| = 0.$$

By proceeding as in Section 2.4 with an additional argument concerning the term $p_2(\varrho)$, we derive the energy conservation (20). \square

5. PROOF OF THEOREM 1.7

The proof of Theorem 1.7 is similar to that of Theorem 1.5, except that we have to take care of boundary layers when taking integration by parts. Take $0 < \varepsilon < \varepsilon_1/10 < \varepsilon_3/10 < \delta_0/100$, as proof of Theorem 1.3, we have

$$\begin{aligned} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \partial_t (\varrho \mathbf{u})^\varepsilon dx ds d\varepsilon_2 + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla \cdot (\varrho \mathbf{u} \otimes \mathbf{u})^\varepsilon dx ds d\varepsilon_2 \\ + \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1+\varepsilon_3} \int_\tau^t \int_{\Omega_{\varepsilon_2}} \frac{\phi_{\eta,\varepsilon}}{\varrho^\varepsilon} (\varrho \mathbf{u})^\varepsilon \nabla (p(\varrho))^\varepsilon dx ds d\varepsilon_2 = 0. \end{aligned} \tag{83}$$

In view of the proof of Theorem 1.3, we only have to deal with the term (E33) (now without the divergence free condition) and the last term on the left-hand side of (83).

For (E33) we have, using arguments similar to (37),

$$\begin{aligned}
|(E33)| &= \frac{1}{2} \eta^{-1} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} |(\varrho \mathbf{u})^\varepsilon|^2 \phi' \left(\frac{\varrho^\varepsilon}{\eta} \right) \nabla \varrho^\varepsilon \frac{\mathbf{u}^\varepsilon}{\varrho^\varepsilon} dx ds d\varepsilon_2 \\
&\leq C \eta^{-2} \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} |\mathbf{u}^\varepsilon| |(\varrho \mathbf{u})^\varepsilon|^2 \mathbf{1}_{\eta \leq \varrho^\varepsilon} |\nabla \varrho^\varepsilon| dx ds d\varepsilon_2 \\
&\leq C \|\mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta}\} \nabla \varrho^\varepsilon\|_{L^\infty} \left\| \mathbf{u} \mathbf{1}_{\{\frac{1}{3}\eta \leq \varrho \leq \frac{8}{3}\eta}\} \right\|_{L^3}^3,
\end{aligned}$$

and therefore

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} |(E33)| = 0$$

thanks to (23). For the last term on the left-hand side of (83), we can repeat all the arguments in estimating (G) in the proof of Theorem 1.5. The only extra effort is the boundary terms when integrating by parts. More precisely, for the sixth line of (78), (G2) and (G53), which are

$$\begin{aligned}
(H1) &= \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_0^t \int_{\partial \Omega_{\varepsilon_2}} p_1(\varrho^\varepsilon) \phi_{\eta, \varepsilon}(\varrho \mathbf{u})^\varepsilon \cdot n(\theta) d\mathcal{H}^{d-1}(\theta) dt d\varepsilon_2, \\
(H2) &= \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_0^t \int_{\partial \Omega_{\varepsilon_2}} \phi_{\eta, \varepsilon} \mathbf{u}^\varepsilon [(p(\varrho))^\varepsilon - p(\varrho^\varepsilon)] \cdot n(\theta) d\mathcal{H}^{d-1}(\theta) dt d\varepsilon_2,
\end{aligned}$$

and

$$(H3) = p(1) \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_0^t \int_{\partial \Omega_{\varepsilon_2}} \phi_{\eta, \varepsilon}(\varrho \mathbf{u})^\varepsilon \cdot n(\theta) d\mathcal{H}^{d-1}(\theta) dt d\varepsilon_2,$$

respectively. Using the assumptions (24) and (25) we can easily estimate (H1), (H2) and (H3) similarly to (E31) and (E32) to obtain

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} (|(H1)| + |(H2)| + |(H3)|) = 0.$$

Therefore, we conclude

$$\limsup_{\eta; \varepsilon_3; \varepsilon_1; \varepsilon; \tau \rightarrow 0} \left| \frac{1}{\varepsilon_3} \int_{\varepsilon_1}^{\varepsilon_1 + \varepsilon_3} \int_{\tau}^t \int_{\Omega_{\varepsilon_2}} \partial_t \left[\frac{1}{2} \frac{\phi_{\eta, \varepsilon}}{\varrho^\varepsilon} |(\varrho \mathbf{u})^\varepsilon|^2 + \phi_{\eta, \varepsilon} p_2(\varrho^\varepsilon) \right] dx ds \right| = 0.$$

This gives the energy conservation (26). The proof is complete. \square

APPENDIX A. APPENDIX

In this section, we collect some lemmata and estimates which are used for the proofs of the main results.

We recall that, for a function $f : \Omega \rightarrow \mathbb{R}$ we denote by $f^\varepsilon = f \star \omega_\varepsilon$ the smoothing version of f , where $\omega_\varepsilon(x) = (1/\varepsilon^d) \omega(x/\varepsilon)$ is a standard mollifier in \mathbb{R}^d .

Lemma A.1. *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with C^2 boundary $\partial \Omega$.*

(1) *Let $\beta \in (0, 1)$ and $1 \leq p \leq \infty$. Then for any function $f : \Omega \rightarrow \mathbb{R}$ and small $0 < \varepsilon \leq \frac{\delta}{2}$, there holds*

$$\|\nabla f^\varepsilon\|_{L^p(\Omega_\delta)} \leq C \varepsilon^{-1} \|f\|_{L^p(\Omega_{\frac{\delta}{2}})}, \quad (84)$$

$$\|\nabla f^\varepsilon\|_{L^p(\Omega_\delta)} \leq C \varepsilon^{-1+\beta} \|f\|_{\mathcal{V}_\delta^{\beta, p}(\Omega)}. \quad (85)$$

(2) Let $\beta_1, \beta_2 \in (0, 1)$, $p \geq 1, p_1, p_2 \geq 1$ such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$ and $\delta_1 > 0, \delta_2 > 0, \delta \geq \max\{\delta_1, \delta_2\}$. Then for any functions $g_1, g_2 : \Omega \rightarrow \mathbb{R}$ and small $0 < \varepsilon \leq \frac{\min\{\delta_1, \delta_2\}}{2}$, there holds

$$\|(g_1 g_2)^\varepsilon - g_1^\varepsilon g_2^\varepsilon\|_{L^p(\Omega_\delta)} \leq C \varepsilon^{\beta_1 + \beta_2} \|g_1\|_{\mathcal{V}_{\delta_1}^{\beta_1, p_1}(\Omega)} \|g_2\|_{\mathcal{V}_{\delta_2}^{\beta_2, p_2}(\Omega)}. \quad (86)$$

(3) Let $\beta \in (0, 1)$ and $p \geq 1$. Then for any functions $g_1, g_2 : \Omega \rightarrow \mathbb{R}$ and small $\delta \in (0, 1)$, there holds

$$\|g_1 g_2\|_{\mathcal{V}_\varepsilon^{\beta, p}(\Omega_\delta)} \leq C \left(\|g_1\|_{L^\infty(\Omega_{\frac{\delta}{2}})} \|g_2\|_{\mathcal{V}_\varepsilon^{\beta, p}(\Omega_{\frac{\delta}{2}})} + \|g_1\|_{\mathcal{V}_\varepsilon^{\beta, \infty}(\Omega_{\frac{\delta}{2}})} \|g_2\|_{L^p(\Omega_{\frac{\delta}{2}})} \right), \quad (87)$$

for any $0 < \varepsilon < \delta/4$.

Proof. 1) Since $\int_{\mathbb{R}^d} \nabla \omega_\varepsilon(y) dy = 0$, it follows that, for a.e. $x \in \Omega_\delta$,

$$|\nabla f^\varepsilon(x)| = \left| \int_{\mathbb{R}^d} [f(x-y) - f(x)] \nabla \omega_\varepsilon(y) dy \right| \leq \varepsilon^{-1-\frac{d}{p}} \left(\int_{|y| < \varepsilon} |f(x-y) - f(x)|^p dy \right)^{\frac{1}{p}} \|\nabla \omega\|_{L^{\frac{p}{p-1}}}.$$

This yields (84) and (85).

2) It is not hard to see that

$$\begin{aligned} |(g_1 g_2)^\varepsilon(x) - g_1^\varepsilon(x) g_2^\varepsilon(x)| &\leq \int_{\mathbb{R}^d} |g_1(x-y) - g_1(x)| |g_2(x-y) - g_2(x)| \omega_\varepsilon(y) dy \\ &\quad + |g_1^\varepsilon(x) - g_1(x)| |g_2^\varepsilon(x) - g_2(x)|. \end{aligned}$$

Thus, using Holder's inequality, we get (86).

3) For every $x \in \Omega_\delta$ and $h \in \mathbb{R}^d$ such that $|h| < \delta/2$, we have

$$|(fg)(x+h) - (fg)(x)| \leq |f(x+h)| |g(x+h) - g(x)| + |f(x+h) - f(x)| |g(x)|.$$

Clearly, this gives (87). \square

Lemma A.2.

(1) Let $\beta \in (0, 1)$ and $1 \leq p \leq \infty$. Then for any function $f : \mathbb{T}^d \rightarrow \mathbb{R}$ and $0 < \varepsilon < \delta$, there holds

$$\begin{aligned} \|\nabla f^\varepsilon\|_{L^p(\mathbb{T}^d)} &\leq C \varepsilon^{-1} \|f\|_{L^p(\mathbb{T}^d)}, \\ \|\nabla f^\varepsilon\|_{L^p(\mathbb{T}^d)} &\leq C \varepsilon^{-1+\beta} \|f\|_{\mathcal{V}_\varepsilon^{\beta, p}(\mathbb{T}^d)}. \end{aligned}$$

(2) Let $\beta_1, \beta_2 \in (0, 1)$ and $p \geq 1, p_1, p_2 \geq 1$ such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Then for any functions $g_1, g_2 : \mathbb{T}^d \rightarrow \mathbb{R}$ and $0 < \varepsilon < \delta$, there holds

$$\|(g_1 g_2)^\varepsilon - g_1^\varepsilon g_2^\varepsilon\|_{L^p(\mathbb{T}^d)} \leq C \varepsilon^{\beta_1 + \beta_2} \|\omega\|_{L^{\frac{p}{p-1}}(\mathbb{T}^d)} \|g_1\|_{\mathcal{V}_\varepsilon^{\beta_1, p_1}(\mathbb{T}^d)} \|g_2\|_{\mathcal{V}_\varepsilon^{\beta_2, p_2}(\mathbb{T}^d)}.$$

(3) Let $\beta \in (0, 1)$ and $p \geq 1$. Then for any functions $g_1, g_2 : \mathbb{T}^d \rightarrow \mathbb{R}$ and $\varepsilon > 0$, there holds

$$\|g_1 g_2\|_{\mathcal{V}_\varepsilon^{\beta, p}(\mathbb{T}^d)} \leq C \left(\|g_1\|_{L^\infty(\mathbb{T}^d)} \|g_2\|_{\mathcal{V}_\varepsilon^{\beta, p}(\mathbb{T}^d)} + \|g_1\|_{\mathcal{V}_\varepsilon^{\beta, \infty}(\mathbb{T}^d)} \|g_2\|_{L^p(\mathbb{T}^d)} \right).$$

Proof. The proof is similar to that of Lemma A.1 and we omit it. \square

Lemma A.3. Assume that there exists $\delta_0 > 0$ and $M > 0$ such that

$$\|\varrho\|_{L^\infty(0, T; \mathcal{V}_{\delta_0}^{\frac{1}{3}, \infty}(\mathbb{T}^d))} < M. \quad (88)$$

For any $0 < \varepsilon^{\frac{1}{3}} < \frac{1}{3M} \eta < \frac{1}{12M} \delta_0$, the followings hold.

(i) If $\varrho^\varepsilon(x, t) \geq \eta$ then $\varrho(x, t) > \frac{2}{3} \eta$. In addition, for any $y \in B_\varepsilon(0)$, $\varrho(x-y, t) > \frac{1}{3} \eta$.

(ii) If $\varrho^\varepsilon(x, t) \leq 2\eta$ then $\varrho(x, t) < \frac{7}{3} \eta$. In addition, for any $y \in B_\varepsilon(0)$, $\varrho(x-y, t) < \frac{8}{3} \eta$.

Proof. From (88), we see that

$$|\varrho(x-y, t) - \varrho(x, t)| < M|y|^{\frac{1}{3}} \quad \text{for a.e. } x \in \mathbb{T}^d, y \in B_{\delta_0}(0), t \in (0, T). \quad (89)$$

We write

$$\varrho^\varepsilon(x, t) = \int_{B_\varepsilon(0)} (\varrho(x-y, t) - \varrho(x, t)) \omega_\varepsilon(y) dy + \varrho(x, t).$$

Therefore, if $\varrho^\varepsilon(x, t) \geq \eta$, by (89), we obtain

$$\varrho(x, t) \geq \varrho^\varepsilon(x, t) - \int_{B_\varepsilon(0)} |\varrho(x-y, t) - \varrho(x, t)| \omega_\varepsilon(y) dy \geq \eta - M\varepsilon^{\frac{1}{3}} > \frac{2}{3}\eta.$$

This, together with (89), implies

$$\varrho(x-y, t) > \varrho(x, t) - M\varepsilon^{\frac{1}{3}} > \frac{1}{3}\eta.$$

Similarly, if $\varrho(x, t) \leq 2\eta$, by (89), we obtain

$$\varrho(x, t) \leq \varrho^\varepsilon(x, t) + \int_{B_\varepsilon(0)} |\varrho(x-y, t) - \varrho(x, t)| \omega_\varepsilon(y) dy < 2\eta + M\varepsilon^{\frac{1}{3}} < \frac{7}{3}\eta.$$

This, together with (89), implies

$$\varrho(x-y, t) \leq \varrho(x, t) + M\varepsilon^{\frac{1}{3}} < \frac{8}{3}\eta.$$

□

Acknowledgements: Q.-H. Nguyen has been supported by the Centro De Giorgi, Scuola Normale Superiore, Pisa, Italy. P.-T. Nguyen has been supported by Czech Science Foundation, project GJ19 – 14413Y. B. Q. Tang gratefully acknowledges the support of the Hausdorff Research Institute for Mathematics (Bonn), through the Junior Trimester Program on Kinetic Theory. This work is partially supported by the International Training Program IGDK 1754 and NAWI Graz.

REFERENCES

- [ADSW] I. Akramov, T. Debiec, J. W. D. Skipper, E. Wiedemann. Energy Conservation for the Compressible Euler and Navier-Stokes Equations with Vacuum. *arXiv:1808.05029*.
- [BT18] C. Bardos, E.S. Titi. Onsager’s conjecture for the incompressible Euler equations in bounded domains. *Arch. Ration. Mech. Anal.* 228 (2018), no. 1, 197-207.
- [BTW] C. Bardos, E.S. Titi, E. Wiedemann. Onsager’s Conjecture with Physical Boundaries and an Application to the Vanishing Viscosity Limit. *arXiv:1803.04939*.
- [CET94] P. Constantin, W. E, E.S. Titi. Onsager’s conjecture on the energy conservation for solutions of Euler’s equation. *Comm. Math. Phys.* 165 (1994), no. 1, 207-209.
- [CY] R.M. Chen, C. Yu. Onsager’s energy conservation for inhomogeneous Euler equations. *to appear in J. Math. Pure. Appl.*
- [DN18] T.D. Drivas, H.Q. Nguyen. Onsagers conjecture and anomalous dissipation on domains with boundary. *SIAM J. Math. Anal.* 50(5) (2018) 4785–4811.
- [DS12] C. De Lellis, L.J. Székelyhidi, The h -principle and the equations of fluid dynamics. *Bull. Amer. Math. Soc. (N.S.)* 49 (2012), no. 3, 347-375.
- [DS13] C. De Lellis, L.J. Székelyhidi. Dissipative continuous Euler flows. *Invent. Math.* 193 (2013), no. 2, 377-407.
- [DS14] C. De Lellis, L.J. Székelyhidi. Dissipative Euler flows and Onsager’s conjecture. *J. Eur. Math. Soc.* 16 (2014), no. 7, 1467-1505.
- [Fei04] E. Feireisl. Dynamics of viscous compressible fluids. Oxford Lecture Series in Mathematics and its Applications, 26. Oxford Science Publications. *The Clarendon Press, Oxford University Press, New York* (2004).
- [FGGW17] E. Feireisl, P. Gwiazda, A. Swierczewska-Gwiazda, and E. Wiedemann. Regularity and energy conservation for the compressible Euler equations. *Arch. Ration. Mech. Anal.* 223 (2017), 1375-1395.
- [Ise18] P. Isett, On the endpoint regularity in Onsagers conjecture. Preprint, 2017. *arXiv:1706.01549*.

- [Ise18a] P. Isett. A proof of Onsager's conjecture. *Ann. Math.* 188 (2018) 871–963.
- [BDIS15] T. Buckmaster, C. De Lellis, P. Isett, L.J. Székelyhidi. Anomalous dissipation for $1/5$ -Hölder Euler flows. *Ann. of Math. (2)* 182 (2015), no. 1, 127-172.
- [LVV18] I. Lacroix-Violet, A. Vasseur. Global weak solutions to the compressible quantum Navier-Stokes equation and its semi-classical limit. *J. Math. Pures Appl.* 114 (2018) 191-210.
- [LS16] T.M. Leslie, R. Shvydkoy. The energy balance relation for weak solutions of the density-dependent Navier-Stokes equations. *J. Differ. Equ.* 261 (2016), 3719-3733.
- [Lio96] P.-L. Lions. Mathematical topics in fluid mechanics. Volume 1: Incompressible models. *Clarendon Press - Oxford* (1996).
- [NN19] Q.-H. Nguyen, P.-T. Nguyen. Onsager's conjecture on the energy conservation for solutions of Euler equations in bounded domains. *Journal of Nonlinear Science* 29.1 (2019) 207–213.
- [Ons49] L. Onsager, Statistical Hydrodynamics. *Nuovo Cimento (Supplemento)*, 6 (1949), 279-287
- [Sch93] V. Scheffer, An inviscid flow with compact support in space-time. *J. Geom. Anal.* 3 (1993), 343-401
- [Shn97] A. Shnirelman, On the nonuniqueness of weak solution of the Euler equation, *Comm. Pure Appl. Math.* 50.12 (1997) 1261-1286.

QUOC-HUNG NGUYEN

SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI 7, 56100 PISA, ITALY;
 DEPARTMENT OF MATHEMATICS, NEW YORK UNIVERSITY ABU DHABI, ABU DHABI, UNITED ARAB EMIRATES
E-mail address: quochung.nguyen@sns.it, qn2@nyu.edu

PHUOC-TAI NGUYEN

DEPARTMENT OF MATHEMATICS AND STATISTICS, MASARYK UNIVERSITY, 61137 BRNO, CZECH REPUBLIC
E-mail address: ptnguyen@math.muni.cz, nguyenphuoc tai.hcmup@gmail.com

BAO QUOC TANG

INSTITUTE OF MATHEMATICS AND SCIENTIFIC COMPUTING, UNIVERSITY OF GRAZ, HEINRICHSTRASSE 36, 8010
 GRAZ, AUSTRIA
E-mail address: quoc.tang@uni-graz.at