

Kaluza–Klein reduction on a maximally non-Riemannian space is moduli-free

Kyoungcho Cho,^{*} Kevin Morand,[†] and Jeong-Hyuck Park[‡]

Department of Physics, Sogang University, 35 Baekbeom-ro, Mapo-gu, Seoul 04107, Korea

We propose a novel Kaluza–Klein scheme which assumes the internal space to be maximally non-Riemannian, meaning that no Riemannian metric can be defined for any subspace. Its description is only possible through Double Field Theory but not within supergravity. We spell out the corresponding Scherk–Schwarz twistable Kaluza–Klein ansatz, and point out that the internal space prevents rigidly any graviscalar moduli. Plugging the same ansatz into higher-dimensional pure Double Field Theory and also to a known doubled-yet-gauged string action, we recover heterotic supergravity as well as heterotic worldsheet action. In this way, we show that 1) supergravity and Yang–Mills theory can be unified into higher-dimensional pure Double Field Theory, free of moduli, and 2) heterotic string theory may have a higher-dimensional non-Riemannian origin.

I. Introduction

Kaluza–Klein theory attempts to unify General Relativity and electromagnetism into higher-dimensional pure gravity. Yet, the (aesthetically unpleasing) cylindrical extra dimension brings along an unwanted additional massless scalar field, *i.e.* radion or graviscalar modulus, which is not observed in nature: it would spoil the Equivalence Principle by appearing on the right-hand side of the geodesic equation. This moduli stabilization problem is essentially rooted in the fact that there is no natural scale in pure gravity which would fix or stabilize the radius of the cylinder. The problem persists in modern string compactifications, in view of the arbitrary size and shape of an internal space (mani/coni/orbifold, compact or not). Turning on fluxes or non-perturbative corrections might promise to solve the problem, but such scenarios require generically subtle analyses regarding both the validity of the effective-field-theory approximation and calculational control [1] (*c.f.* [2] and references therein for related current controversies).

In this paper we propose a novel Kaluza–Klein scheme to unify Stringy Gravity and Yang–Mills (including Maxwell), which postulates a non-Riemannian internal space and consequently does not suffer from any moduli problem. By Stringy Gravity, we mean the string theory effective action of the NS-NS (or purely bosonic) closed-string massless sector, conventionally represented by the three fields, $\{g_{\mu\nu}, B_{\mu\nu}, \phi\}$. They transform into one another under T-duality and hence form a (reducible) $\mathbf{O}(D, D)$ multiplet [3, 4]. Furthermore, within the framework of Double Field Theory (DFT) initiated in [5–9], $\mathbf{O}(D, D)$ T-duality becomes the manifest principal symmetry and the effective action itself is to be identified as an integral of a stringy scalar curvature beyond Riemann. The whole closed-string massless NS-NS sector may then be viewed as stringy graviton fields consisting of the DFT-dilaton, d , and the DFT-metric, \mathcal{H}_{AB} [10].

The latter satisfies two defining properties:

$$\mathcal{H}_{AB} = \mathcal{H}_{BA}, \quad \mathcal{H}_A{}^C \mathcal{H}_B{}^D \mathcal{J}_{CD} = \mathcal{J}_{AB}, \quad (1)$$

where $\mathcal{J}_{AB} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ is the $\mathbf{O}(D, D)$ invariant constant metric which can freely raise and lower the $\mathbf{O}(D, D)$ vector indices, A, B, \dots (capital letters). A pair of symmetric projection matrices are then defined,

$$P_{AB} = \frac{1}{2}(\mathcal{J}_{AB} + \mathcal{H}_{AB}), \quad \bar{P}_{AB} = \frac{1}{2}(\mathcal{J}_{AB} - \mathcal{H}_{AB}), \quad (2)$$

while their ‘square roots’ give twofold DFT-vielbeins,

$$P_{AB} = V_A{}^p V_B{}^q \eta_{pq}, \quad \bar{P}_{AB} = \bar{V}_A{}^{\bar{p}} \bar{V}_B{}^{\bar{q}} \bar{\eta}_{\bar{p}\bar{q}}, \quad (3)$$

which satisfy their own defining properties,

$$V_{Ap} V^A{}_q = \eta_{pq}, \quad \bar{V}_{A\bar{p}} \bar{V}^A{}_{\bar{q}} = \bar{\eta}_{\bar{p}\bar{q}}, \quad V_{Ap} \bar{V}^A{}_{\bar{q}} = 0, \quad (4)$$

or equivalently [11]

$$V_A{}^p V_B{}^q \eta_{pq} + \bar{V}_A{}^{\bar{p}} \bar{V}_B{}^{\bar{q}} \bar{\eta}_{\bar{p}\bar{q}} = \mathcal{J}_{AB}. \quad (5)$$

Here η_{pq} and $\bar{\eta}_{\bar{p}\bar{q}}$ are local Lorentz invariant metrics characterizing the twofold spin groups which have dimensions $\text{Tr}(P)$ and $\text{Tr}(\bar{P})$ separately. They are distinguished by unbarred and barred small letter indices.

The above stringy graviton fields constitute the diffeomorphic DFT-Christoffel symbols [12–14],

$$\Gamma_{CAB} = 2(P\partial_C P\bar{P})_{[AB]} + 2(\bar{P}_{[A}{}^D \bar{P}_{B]}{}^E - P_{[A}{}^D P_{B]}{}^E) \partial_D P_{EC} - 4\left(\frac{P_{C[A} P_{B]}{}^D}{P_F{}^{F-1}} + \frac{\bar{P}_{C[A} \bar{P}_{B]}{}^D}{\bar{P}_F{}^{F-1}}\right) (\partial_D d + (P\partial^E P\bar{P})_{[ED]}),$$

and, consequently with $\nabla_A := \partial_A + \Gamma_A$, twofold local Lorentz spin connections [15–17],

$$\begin{aligned} \Phi_{Apq} &= V^B{}_p \nabla_A V_{Bq} = V^B{}_p (\partial_A V_{Bq} + \Gamma_{AB}{}^C V_{Cq}), \\ \bar{\Phi}_{A\bar{p}\bar{q}} &= \bar{V}^B{}_{\bar{p}} \nabla_A \bar{V}_{B\bar{q}} = \bar{V}^B{}_{\bar{p}} (\partial_A \bar{V}_{B\bar{q}} + \Gamma_{AB}{}^C \bar{V}_{C\bar{q}}). \end{aligned} \quad (6)$$

^{*} khcho23@sogang.ac.kr

[†] morand@sogang.ac.kr

[‡] park@sogang.ac.kr

These connections form covariant curvatures, Ricci and scalar [13, 17], of which the latter can be constructed as

$$\begin{aligned} S_{(0)} &= (P^{AC}P^{BD} - \bar{P}^{AC}\bar{P}^{BD})S_{ABCD}, \\ \mathcal{G}_{(0)} &= \mathcal{F}_{ABpq}V^{Ap}V^{Bq} + \frac{1}{2}\Phi_{Apq}\Phi^{Apq}, \\ \bar{\mathcal{G}}_{(0)} &= \bar{\mathcal{F}}_{AB\bar{p}\bar{q}}\bar{V}^{A\bar{p}}\bar{V}^{B\bar{q}} + \frac{1}{2}\bar{\Phi}_{A\bar{p}\bar{q}}\bar{\Phi}^{A\bar{p}\bar{q}}, \end{aligned} \quad (7)$$

where, with $R_{CDAB} = \partial_A\Gamma_{BCD} + \Gamma_{AC}{}^E\Gamma_{BED} - (A \leftrightarrow B)$,

$$\begin{aligned} S_{ABCD} &:= \frac{1}{2}(R_{ABCD} + R_{CDAB} - \Gamma^E{}_{AB}\Gamma_{ECD}), \\ \mathcal{F}_{ABpq} &:= \nabla_A\Phi_{Bpq} - \nabla_B\Phi_{Apq} + \Phi_{Ap}{}^r\Phi_{Brq} - \Phi_{Bp}{}^r\Phi_{Arq}, \\ \bar{\mathcal{F}}_{AB\bar{p}\bar{q}} &:= \nabla_A\bar{\Phi}_{B\bar{p}\bar{q}} - \nabla_B\bar{\Phi}_{A\bar{p}\bar{q}} + \bar{\Phi}_{A\bar{p}}{}^{\bar{r}}\bar{\Phi}_{B\bar{r}\bar{q}} - \bar{\Phi}_{B\bar{p}}{}^{\bar{r}}\bar{\Phi}_{A\bar{r}\bar{q}}. \end{aligned}$$

While $S_{(0)}$ coincides exactly with the well-known expression of scalar curvature written in terms of d, \mathcal{H}_{AB} [18],

$$\begin{aligned} S_{(0)} &= \frac{1}{8}\mathcal{H}^{AB}\partial_A\mathcal{H}_{CD}\partial_B\mathcal{H}^{CD} + \frac{1}{2}\mathcal{H}^{AB}\partial^C\mathcal{H}_{AD}\partial^D\mathcal{H}_{BC} \\ &\quad - \partial_A\partial_B\mathcal{H}^{AB} + 4\mathcal{H}^{AB}(\partial_A\partial_B d - \partial_A d\partial_B d) + 4\partial_A\mathcal{H}^{AB}\partial_B d, \end{aligned}$$

the other two accommodate the vielbeins and read [19]

$$\begin{aligned} +\mathcal{G}_{(0)} &= \frac{1}{2}S_{(0)} + 2\partial_A\partial^A d - 2\partial_A d\partial^A d + \frac{1}{2}\partial_A V_{Bp}\partial^A V^{Bp}, \\ -\bar{\mathcal{G}}_{(0)} &= \frac{1}{2}S_{(0)} - 2\partial_A\partial^A d + 2\partial_A d\partial^A d - \frac{1}{2}\partial_A \bar{V}_{B\bar{p}}\partial^A \bar{V}^{B\bar{p}}. \end{aligned} \quad (8)$$

Clearly their differences would vanish upon the section condition, $\partial_A\partial^A \equiv 0$, but they provide precisely the known ‘missing’ pieces in the Scherk–Schwarz reduction of DFT while relaxing the section condition on the internal space, as previously added by hand in [20–23]. Hence, the above vielbein formalism generates these crucial terms in a natural manner. In particular, $\mathcal{G}_{(0)} - \bar{\mathcal{G}}_{(0)}$ matches the action adopted in [22]. Below we focus on computing $+\mathcal{G}_{(0)}$ and $-\bar{\mathcal{G}}_{(0)}$ separately in higher dimensions with the Scherk–Schwarz twisted non-Riemannian Kaluza–Klein ansatz.

II. Moduli-free non-Riemannian Kaluza–Klein ansatz

With $\hat{D} = D' + D$, the DFT Kaluza–Klein ansatz [14] breaks $\mathbf{O}(\hat{D}, \hat{D})$ into $\mathbf{O}(D', D') \times \mathbf{O}(D, D)$ and takes the form:

$$\hat{\mathcal{H}} = \exp[\hat{W}] \begin{pmatrix} \mathcal{H}' & 0 \\ 0 & \mathcal{H} \end{pmatrix} \exp[\hat{W}^T], \quad \hat{\mathcal{J}} = \begin{pmatrix} \mathcal{J}' & 0 \\ 0 & \mathcal{J} \end{pmatrix}. \quad (9)$$

In our notation, hatted, primed, and unprimed symbols refer to the ambient, internal, and external spaces respectively. In particular, the ambient doubled coordinates

split into the internal and external ones as

$$z^{\hat{A}} = (y^{A'}, x^A), \quad \partial_{\hat{A}} = (\partial_{A'}, \partial_A). \quad (10)$$

In (9), \hat{W} is an off-block-diagonal $\mathfrak{so}(\hat{D}, \hat{D})$ element which satisfies

$$\hat{W}\hat{\mathcal{J}} + \hat{\mathcal{J}}\hat{W}^T = 0, \quad (11)$$

and takes the form,

$$\hat{W} = \begin{pmatrix} 0 & -W^c \\ W & 0 \end{pmatrix}, \quad W^c{}_{A'}{}^A := W^A{}_{A'} = \mathcal{J}^{AB}\mathcal{J}'_{A'B'}W_B{}^{B'}, \quad (12)$$

where the $2D \times 2D'$ block, $W_A{}^{A'}$, should meet [24]

$$W_A{}^{A'}W^{AB'} = 0, \quad W^A{}_{A'}\partial_A = 0. \quad (13)$$

This condition sets half of the components to be trivial, truncates the exponential,

$$\exp[\hat{W}] = 1 + \hat{W} + \frac{1}{2}\hat{W}^2,$$

and makes the above ansatz consistent with the ordinary Kaluza–Klein ansatz in supergravity. Moreover, crucially for the purpose of the present work, the ansatz (9) can accommodate non-Riemannian geometry in which the Riemannian metric cannot be defined, see [14] for classification and [25–27] for earlier examples including the attainment of the Gomis–Ooguri non-relativistic string [28, 29].

Henceforth, we focus on a specific internal space of which the DFT-metric is fully $\mathbf{O}(D', D')$ symmetric and maximally non-Riemannian, namely $(D', 0)$ type as classified in [14]:

$$\mathcal{H}'_{A'B'} \equiv \mathcal{J}'_{A'B'}. \quad (14)$$

In general, from the defining relations (1), the infinitesimal variation of any DFT-metric should satisfy

$$\delta\mathcal{H}' = P'\delta\mathcal{H}'\bar{P}' + \bar{P}'\delta\mathcal{H}'P'. \quad (15)$$

Meanwhile, the particular choice of the internal space (14) implies $P' = \mathcal{J}'$ and $\bar{P}' = 0$. Thus, the fluctuation must be trivial: no graviscalar modulus can be generated and the non-Riemannian internal space is *rigid*,

$$\delta\mathcal{H}'_{A'B'} = 0. \quad (16)$$

In fact, (14) sets the ‘‘twofold’’ internal spin group to be $\mathbf{O}(D', D') \times \mathbf{O}(0, 0)$, such that the coset structures of the internal and the ambient DFT-metrics, \mathcal{H}' , $\hat{\mathcal{H}}$, are ‘trivial’ and ‘heterotic’ respectively, if \mathcal{H} is Riemannian (27),

$$\frac{\mathbf{O}(D', D')}{\mathbf{O}(D', D') \times \mathbf{O}(0, 0)} = \mathbf{1}, \quad \frac{\mathbf{O}(\hat{D}, \hat{D})}{\mathbf{O}(D'+1, \hat{D}-1) \times \mathbf{O}(D-1, 1)}. \quad (17)$$

The latter has dimension $D^2 + 2DD'$, which matches the total degrees of the external DFT-metric, \mathcal{H}_{AB} (27), and the gravivector, $W_A{}^{A'}$ (13) [30], *c.f.* [31].

The corresponding DFT-vielbeins are [14]

$$\hat{V}_{\hat{A}\hat{p}} = \exp[\hat{W}] \begin{pmatrix} V'_{A'p'} & 0 \\ 0 & V_{Ap} \end{pmatrix}, \quad \hat{\bar{V}}_{\hat{A}\hat{p}} = \exp[\hat{W}] \begin{pmatrix} 0 \\ \bar{V}_{A\bar{p}} \end{pmatrix}, \quad (18)$$

where $V'_{A'p'}$ is now an invertible ‘square’ matrix,

$$V'_{A'p'} V'_{B'p'} = \mathcal{J}'_{A'B'}.$$

III. Reduction to heterotic DFT

Before inserting the ansatz (9), (18) into the ambient scalar curvatures (8), or $\hat{\mathcal{G}}_{(0)}, \hat{\bar{\mathcal{G}}}_{(0)}$ (all hatted), we perform a Scherk–Schwarz twist over the internal space. Following [19], we introduce a twisting matrix, $U_{\hat{A}}{}^{A'}(y)$, which depends on the internal coordinates only and is an $\mathbf{O}(D', D')$ element satisfying

$$U_{\hat{A}}{}^{A'} U_{\hat{B}}{}^{B'} \mathcal{J}'_{A'B'} = \dot{\mathcal{J}}_{\hat{A}\hat{B}}, \quad (19)$$

where $\dot{\mathcal{J}}_{\hat{A}\hat{B}}$ coincides with $\mathcal{J}'_{A'B'}$ numerically: both are $\mathbf{O}(D', D')$ invariant constant metrics. Essentially the twist converts all the primed indices to dotted ones,

$$W_A{}^{A'}(z) = W_{\hat{A}}{}^{\hat{A}}(x) U_{\hat{A}}{}^{A'}(y), \quad V'^{A'}{}_{p'}(z) = \dot{V}^{\hat{A}}{}_{p'}(x) U_{\hat{A}}{}^{A'}(y), \quad (20)$$

where now $\dot{V}^{\hat{A}}{}_{p'} \dot{V}^{\hat{B}}{}_{p'} = \dot{\mathcal{J}}_{\hat{A}\hat{B}}$. We further put

$$\hat{d}(z) = \lambda(y) + d(x), \quad V_{Ap}(x), \quad \bar{V}_{A\bar{p}}(x), \quad (21)$$

such that the external fields are independent of the internal coordinates: $\partial_{A'} d = 0$, $\partial_{A'} V_{Bp} = 0$, $\partial_{A'} \bar{V}_{B\bar{p}} = 0$.

Finally, we impose the standard section condition on the external space, $\partial_A \partial^A = 0$, and the twistability conditions on the internal space separately [22],[19]:

$$\partial_{A'} \lambda - \frac{1}{2} U^{\hat{A}}{}_{A'} \partial_{B'} U_{\hat{A}}{}^{B'} = 0, \quad f_{[\hat{A}\hat{B}}{}^{\hat{E}} f_{\hat{C}]\hat{D}\hat{E}} = 0, \quad (22)$$

where $f_{\hat{A}\hat{B}\hat{C}} = 3(\partial_{B'} U_{[\hat{A}}{}^{A'}) U_{\hat{B}}{}^{B'} U_{\hat{C}]}{}^{A'}$ which we require to be constant.

After straightforward yet tedious computation – assisted by a computer algebra system [32] and through intermediate expressions like (4.14) in [19] – we obtain our main result: with the above Scherk–Schwarz twisted Kaluza–Klein ansatz substituted, the higher-dimensional scalar curvatures (8) reduce precisely to (*c.f.* [22, 33–38])

$$\begin{aligned} +2\hat{\mathcal{G}}_{(0)} &= -2\hat{\bar{\mathcal{G}}}_{(0)} + \frac{1}{3} f_{\hat{A}\hat{B}\hat{C}} f^{\hat{A}\hat{B}\hat{C}}, \\ -2\hat{\bar{\mathcal{G}}}_{(0)} &= S_{(0)} - \frac{1}{4} \mathcal{H}^{AC} \mathcal{H}^{BD} F_{AB}{}^{\hat{A}} F_{CD\hat{A}} \\ &\quad - \frac{1}{12} \mathcal{H}^{AD} \mathcal{H}^{BE} \mathcal{H}^{CF} \omega_{ABC} \omega_{DEF} \\ &\quad + \frac{1}{2} \mathcal{H}^{AD} \mathcal{H}^{BE} \mathcal{H}^{CF} \omega_{ABC} \mathcal{H}_{[D}{}^G \partial_E \mathcal{H}_{F]G}, \end{aligned} \quad (23)$$

where we set Yang–Mills and Chern–Simons terms,

$$\begin{aligned} F_{AB}{}^{\hat{C}} &= \partial_A W_B{}^{\hat{C}} - \partial_B W_A{}^{\hat{C}} + f_{\hat{A}\hat{B}}{}^{\hat{C}} W_A{}^{\hat{A}} W_B{}^{\hat{B}}, \\ \omega_{ABC} &= 3W_{[\hat{A}}{}^{\hat{A}} \partial_B W_{\hat{C}]\hat{A}} + f_{\hat{A}\hat{B}\hat{C}} W_A{}^{\hat{A}} W_B{}^{\hat{B}} W_C{}^{\hat{C}}, \end{aligned} \quad (24)$$

of which the $\mathbf{O}(D, D)$ indices are totally skew-symmetric, $F_{AB}{}^{\hat{C}} = F_{[AB]}{}^{\hat{C}}$, $\omega_{ABC} = \omega_{[ABC]}$, and further from (13),

$$F^{AB}{}_{\hat{C}} \partial_A = 0, \quad \omega^{ABC} \partial_A = 0. \quad (25)$$

The higher \hat{D} -dimensional diffeomorphisms generated by the standard DFT Lie derivative give rise, with a twisted parameter, $\hat{\xi}^{\hat{A}} = (\Lambda^{\hat{A}} U_{\hat{A}}{}^{A'}, \xi^A)$, to the D -dimensional diffeomorphisms plus Yang–Mills gauge symmetry, *c.f.* [19, 22],

$$\begin{aligned} \delta W_A{}^{\hat{A}} &= \xi^C \partial_C W_A{}^{\hat{A}} + (\partial_A \xi^B - \partial^B \xi_A) W_B{}^{\hat{A}} \\ &\quad + \partial_A \Lambda^{\hat{A}} + f^{\hat{A}}{}_{\hat{B}\hat{C}} W_A{}^{\hat{B}} \Lambda^{\hat{C}}, \\ \delta \mathcal{H}_{AB} &= \xi^C \partial_C \mathcal{H}_{AB} + 2\partial_{[A} \xi_{C]} \mathcal{H}^C{}_B + 2\partial_{[B} \xi_{C]} \mathcal{H}_A{}^C \\ &\quad + (W_{[A}{}^{\hat{A}} \partial_{C]} \Lambda_{\hat{A}}) \mathcal{H}^C{}_B + (W_{[B}{}^{\hat{A}} \partial_{C]} \Lambda_{\hat{A}}) \mathcal{H}_A{}^C, \\ \delta d &= \xi^C \partial_C d - \frac{1}{2} \partial_C \xi^C. \end{aligned} \quad (26)$$

In each transformation above, the first line with $\xi^{\hat{A}}$ is the (external) diffeomorphic DFT Lie derivative and the second with $\Lambda^{\hat{A}}$ is the (internal) Yang–Mills gauge symmetry. In fact, every single term in (23) is (external) diffeomorphism-invariant [39].

It is worth while to note the only difference between $2\hat{\mathcal{G}}_{(0)}$ and $-2\hat{\bar{\mathcal{G}}}_{(0)}$: the former contains a DFT-cosmological constant term [13], $\frac{1}{3} e^{-2d} f_{\hat{A}\hat{B}\hat{C}} f^{\hat{A}\hat{B}\hat{C}}$, but the latter does not. As anticipated in [36, 40], our result (23) is manifestly symmetric for $\mathbf{O}(D, D)$ as well as any subgroup of $\mathbf{O}(D', D')$ which stabilizes the structure constant, $f_{\hat{A}\hat{B}\hat{C}}$.

Finally, if we adopt the well-known Riemannian parametrization of the DFT-metric,

$$\mathcal{H}_{AB} = \begin{pmatrix} g^{-1} & -g^{-1}B \\ Bg^{-1} & g - Bg^{-1}B \end{pmatrix}, \quad e^{-2d} = \sqrt{-g} e^{-2\phi}, \quad (27)$$

and solve both the external section condition, $\partial_A \partial^A = 0$, and (13), by letting $\partial_A = (\tilde{\partial}^\mu, \partial_\nu) \equiv (0, \partial_\nu)$ and $W_A{}^{\hat{A}} \equiv (0, W_\nu{}^{\hat{A}})$, our main result (23) reproduces the heterotic supergravity action [41, 42], up to total derivatives,

$$\begin{aligned} &\int -2e^{-2d} \hat{\bar{\mathcal{G}}}_{(0)} \\ &= \int \sqrt{-g} e^{-2\phi} \left(R + 4\partial_\mu \phi \partial^\mu \phi - \frac{1}{12} \tilde{H}_{\lambda\mu\nu} \tilde{H}^{\lambda\mu\nu} - \frac{1}{4} F_{\mu\nu\hat{C}} F^{\mu\nu\hat{C}} \right), \end{aligned} \quad (28)$$

where

$$\tilde{H}_{\lambda\mu\nu} = 3\partial_{[\lambda}B_{\mu\nu]} - \omega_{\lambda\mu\nu}, \quad (29)$$

which is invariant under the Yang–Mills gauge symmetry (26), or specifically in components,

$$\delta W_{\mu}^{\hat{A}} = \partial_{\mu}\Lambda^{\hat{A}} + f^{\hat{A}}_{\hat{B}\hat{C}}W_{\mu}^{\hat{B}}\Lambda^{\hat{C}}, \quad \delta B_{\mu\nu} = W_{[\mu}^{\hat{A}}\partial_{\nu]}\Lambda_{\hat{A}}.$$

However, we have not pinned down the gauge group to be either $\mathbf{E}_8 \times \mathbf{E}_8$ or $\mathbf{SO}(32)$. Further, as the starting scalar curvatures (8) are at most two-derivative, our final result (23) lacks the higher-derivative gravitational Chern–Simons term [43], *c.f.* [44–46].

IV. Reduction to heterotic string

Henceforth we discuss the worldsheet aspect of the DFT background (9) involving the maximally non-Riemannian internal space. The analysis of the string propagating on the most general non-Riemannian background was carried out in [14], with the conclusion that string becomes generically (anti-)chiral. To review this and apply it to the current non-Riemannian background (9), we recall the doubled-yet-gauged string action [47], [25] (*c.f.* [27, 48, 49]),

$$S_{\text{string}} = \frac{1}{4\pi\alpha'} \int d^2\sigma \mathcal{L}_{\text{string}}, \quad (30)$$

which contains a generic DFT-metric,

$$\mathcal{L}_{\text{string}} = -\frac{1}{2}\sqrt{-\hbar}h^{\alpha\beta}D_{\alpha}z^{\hat{A}}D_{\beta}z^{\hat{B}}\hat{\mathcal{H}}_{\hat{A}\hat{B}} - \epsilon^{\alpha\beta}D_{\alpha}z^{\hat{A}}\mathcal{A}_{\beta\hat{A}}.$$

With ambient doubled coordinates, $z^{\hat{A}} = (\tilde{z}_{\hat{\mu}}, z^{\hat{\nu}})$, and an auxiliary gauge potential, $\mathcal{A}_{\alpha}^{\hat{A}}$, a covariant derivative is introduced,

$$D_{\alpha}z^{\hat{A}} := \partial_{\alpha}z^{\hat{A}} - \mathcal{A}_{\alpha}^{\hat{A}}, \quad \mathcal{A}_{\alpha}^{\hat{A}}\partial_{\hat{A}} = 0. \quad (31)$$

While this action is completely covariant with respect to the desired symmetries like $\mathbf{O}(D, D)$ T-duality, Weyl symmetry, and worldsheet as well as target-spacetime diffeomorphisms, it also realizes concretely the assertion that doubled coordinates in DFT are actually gauged and each gauge orbit in the doubled coordinate system corresponds to a single physical point [50]. Specifically, with the choice of the section, $\partial_{\hat{A}} = (\tilde{\partial}^{\hat{\mu}}, \partial_{\hat{\nu}}) \equiv (0, \partial_{\hat{\nu}})$, the condition on the auxiliary gauge potential (31) is solved by $\mathcal{A}_{\alpha}^{\hat{A}} = (\mathcal{A}_{\alpha\hat{\mu}}, 0)$. Therefore, half of the doubled coordinates, namely the tilde coordinates, are gauged,

$$D_{\alpha}z^{\hat{A}} = (\partial_{\alpha}\tilde{z}_{\hat{\mu}} - \mathcal{A}_{\alpha\hat{\mu}}, \partial_{\alpha}z^{\hat{\nu}}).$$

Upon the Riemannian background (27), the auxiliary gauge potential appears quadratically in the action (30). Then after integrating it out, one recovers the standard undoubled string action with $g_{\mu\nu}$ and $B_{\mu\nu}$. On the

other hand, upon a generic non-Riemannian DFT background which is characterized by two non-negative integers, (n, \bar{n}) , the components of the auxiliary field, $\mathcal{A}_{\alpha}^{\hat{A}}$, appear linearly for n and \bar{n} directions, playing the role of Lagrange multipliers. Consequently, string becomes chiral and anti-chiral over the n and \bar{n} directions respectively [14].

Now, for the present non-Riemannian background (9), in accordance with the decomposition of the ambient space (10), we put $y^{A'} = (\tilde{y}_{\mu'}, y^{\nu'})$ for the internal doubled coordinates, $x^A = (\tilde{x}_{\mu}, x^{\nu})$ for the external ones, and $W_{\mu}^{A'} = (W_{\mu\nu'}, \tilde{W}_{\mu}^{\nu'})$ for the gravivector. Further, for the ambient gauge potential, $\mathcal{A}_{\alpha}^{\hat{A}} = (\mathcal{A}_{\alpha}^{A'}, \mathcal{A}_{\alpha}^A)$, we set the internal and the external ones as $\mathcal{A}_{\alpha}^{A'} = (A_{\alpha\mu'}, 0)$ and $\mathcal{A}_{\alpha}^A = (A_{\alpha\mu}, 0)$. With this preparation and (9), (27), the string action (30) becomes quadratic in the combination of the gauge potential components, $A_{\alpha\mu} - \tilde{W}_{\mu}^{\mu'}A_{\alpha\mu'}$, and linear in $A_{\alpha\mu}$. Hence, the former leads to a Gaussian integral and the latter plays the role of a Lagrange multiplier. After all, the doubled-yet-gauged string action (30) reduces to $\frac{1}{2\pi\alpha'} \int d^2\sigma \mathcal{L}_{\text{Het}}$, with

$$\begin{aligned} \mathcal{L}_{\text{Het}} = & -\frac{1}{2}\sqrt{-\hbar}h^{\alpha\beta}\partial_{\alpha}x^{\mu}\partial_{\beta}x^{\nu}g_{\mu\nu} + \frac{1}{2}\epsilon^{\alpha\beta}\partial_{\alpha}x^{\mu}\partial_{\beta}x^{\nu}B_{\mu\nu} \\ & + \frac{1}{2}\epsilon^{\alpha\beta}(\partial_{\alpha}\tilde{y}_{\mu'} + \partial_{\alpha}x^{\mu}W_{\mu\nu'}) (\partial_{\beta}y^{\mu'} + \partial_{\beta}x^{\lambda}\tilde{W}_{\lambda}^{\mu'}) \\ & + \frac{1}{2}\epsilon^{\alpha\beta}(\partial_{\alpha}\tilde{x}_{\mu} - W_{\mu A'}\partial_{\alpha}y^{A'})\partial_{\beta}x^{\mu}. \end{aligned} \quad (32)$$

Furthermore, $\partial_{\alpha}y^{\mu'} + \partial_{\alpha}x^{\mu}\tilde{W}_{\mu}^{\mu'}$ must be *chiral*:

$$\left(h^{\alpha\beta} + \frac{1}{\sqrt{-\hbar}}\epsilon^{\alpha\beta}\right)\left(\partial_{\beta}y^{\mu'} + \partial_{\beta}x^{\mu}\tilde{W}_{\mu}^{\mu'}\right) = 0. \quad (33)$$

Especially when the gravivector, $W_{\mu}^{\hat{A}}$, is trivial, the second and third lines in (32) merge into a known topological term, $\frac{1}{2}\epsilon^{\alpha\beta}\partial_{\alpha}\tilde{z}_{\hat{\mu}}\partial_{\beta}z^{\hat{\mu}}$ [47, 51], while (33) gets simplified to make the internal coordinates chiral:

$$y^{\mu'}(\tau, \sigma) = y^{\mu'}(0, \tau + \sigma). \quad (34)$$

This implies – at least classically [14, 26] – that a closed string subject to a periodic boundary condition cannot vibrate in the internal space. In summary, upon the DFT background (9) with the maximally non-Riemannian internal space and trivial gravivector, the external fields, x^{μ} , are described by the usual string action corresponding to the first line in (32), and consist of the equal number of left-moving and right-moving sectors. On the other hand, the internal fields, $y^{\mu'}$, become all chiral, which agrees with the rigidity (16).

V. Conclusion

The maximally non-Riemannian DFT background specified by the DFT-metric, $\mathcal{H}'_{A'B'} = \mathcal{J}'_{A'B'}$ (14), is singled out to be completely $\mathbf{O}(D', D')$ symmetric and rigid: it does not admit any linear fluctuation, $\delta\mathcal{H}'_{A'B'} = 0$ (16),

nor graviscalar moduli, and the coset structure is trivial (17). It is the most symmetric vacuum in DFT.

For the DFT Kaluza–Klein ansatz, (9) and (18), we set the internal space to be maximally non-Riemannian, performed a Scherk–Schwarz twist (20), (22), and computed the ambient higher $(D' + D)$ -dimensional DFT scalar curvatures, $+\hat{\mathcal{G}}_{(0)}$ and $-\hat{\mathcal{G}}_{(0)}$ (8), which lead to the $\mathbf{O}(D, D)$ -manifest formulation of the non-Abelian heterotic supergravity (23). Only the former, $+\hat{\mathcal{G}}_{(0)}$, contains a DFT-cosmological constant. In this way, *supergravity and Yang–Mills theory can be unified into a higher-dimensional pure Stringy Gravity, free of moduli.*

Plugging the *same* non-Riemannian Kaluza–Klein ansatz (9) into the doubled-yet-gauged string action (30) may reproduce the usual heterotic string action,

$$\frac{1}{2} \int_{\Sigma} (-\sqrt{-\hbar} h^{\alpha\beta} g_{\mu\nu} + \epsilon^{\alpha\beta} B_{\mu\nu}) \partial_{\alpha} x^{\mu} \partial_{\beta} x^{\nu} + \epsilon^{\alpha\beta} \partial_{\alpha} \tilde{z}_{\bar{\mu}} \partial_{\beta} z^{\bar{\mu}},$$

with chiral internal coordinates, $(h^{\alpha\beta} + \frac{\epsilon^{\alpha\beta}}{\sqrt{-\hbar}}) \partial_{\beta} y^{\mu} = 0$. This is indeed the case when the Yang–Mills sector is trivial. Therefore, *heterotic string theory may have a higher-dimensional origin with non-Riemannian internal space.*

Our analyses have been focused on the bosonic sectors, both on the target-spacetime and on the world-sheet. Inclusion of fermions is highly desired, *c.f.* [16, 17, 27].

We also leave the exploration of the worldsheet aspect of the somewhat mysterious ‘relaxed’ section condition designed for the Scherk–Schwarz twist [19–23] for future work with nontrivial gravivector, $W_A{}^{A'} \neq 0$, for the action (32). Uplift of the Standard Model of particle physics coupled to DFT [24] to higher dimensions (bottom-up) would be also of interest, as well as applications to string compactifications (top-down), possibly on other types of non-Riemannian internal spaces [52].

Note added.

Motivated by the first version of this work on arXiv, Berman, Blair, and Otsuki explored non-Riemannian geometries in M-theory [53]. In particular, they pointed out that the maximally non-Riemannian E_8 background naturally realizes the topological (hence ‘moduli-free’) phase of Exceptional Field Theory [54].

Acknowledgments.

We wish to thank Stephen Angus, Kanghoon Lee for useful comments, and Wonyoung Cho for valuable help at the early stage of the project. This work was supported by the National Research Foundation of Korea through the Grants, NRF-2016R1D1A1B01015196 and NRF-2018H1D3A1A01030137 (Korea Research Fellowship Program).

-
- [1] S. Kachru, R. Kallosh, A. D. Linde and S. P. Trivedi, “De Sitter vacua in string theory,” *Phys. Rev. D* **68** (2003) 046005 [hep-th/0301240].
- [2] G. Obied, H. Ooguri, L. Spodyneiko and C. Vafa, “De Sitter Space and the Swampland,” arXiv:1806.08362 [hep-th].
- [3] T. H. Buscher, “A Symmetry of the String Background Field Equations,” *Phys. Lett. B* **194** (1987) 59.
- [4] T. H. Buscher, “Path Integral Derivation of Quantum Duality in Nonlinear Sigma Models,” *Phys. Lett. B* **201** (1988) 466.
- [5] W. Siegel, “Two vierbein formalism for string inspired axionic gravity,” *Phys. Rev. D* **47** (1993) 5453 [hep-th/9302036].
- [6] W. Siegel, “Superspace duality in low-energy superstrings,” *Phys. Rev. D* **48** (1993) 2826 [hep-th/9305073].
- [7] C. Hull and B. Zwiebach, “Double Field Theory,” *JHEP* **0909** (2009) 099 [arXiv:0904.4664 [hep-th]].
- [8] C. Hull and B. Zwiebach, *JHEP* **0909** (2009) 090 [arXiv:0908.1792 [hep-th]].
- [9] O. Hohm, C. Hull and B. Zwiebach, “Background independent action for double field theory,” *JHEP* **1007** (2010) 016 [arXiv:1003.5027 [hep-th]].
- [10] All the equations of motion of the closed-string massless sector coupled to matter in an $\mathbf{O}(D, D)$ -covariant manner can be unified into a single expression, $G_{AB} = 8\pi G T_{AB}$: S. Angus, K. Cho and J. H. Park, “Einstein Double Field Equations,” *Eur. Phys. J. C* **78** (2018) no.6, 500 [arXiv:1804.00964 [hep-th]].
- [11] Equations (4) and (5) imply respectively that $(V_A{}^p, \bar{V}_A{}^{\bar{p}})$ is the right and left inverse of $(V^B{}_q, \bar{V}^B{}_{\bar{q}})^T$. Since left inverse is the same as right inverse in general, (4) and (5) are equivalent.
- [12] I. Jeon, K. Lee and J. H. Park, “Differential geometry with a projection: Application to double field theory,” *JHEP* **1104** (2011) 014 [arXiv:1011.1324 [hep-th]].
- [13] I. Jeon, K. Lee and J. H. Park, “Stringy differential geometry, beyond Riemann,” *Phys. Rev. D* **84** (2011) 044022 [arXiv:1105.6294 [hep-th]].
- [14] K. Morand and J. H. Park, “Classification of non-Riemannian doubled-yet-gauged spacetime,” *Eur. Phys. J. C* **77** (2017) no.10, 685 [arXiv:1707.03713 [hep-th]].
- [15] I. Jeon, K. Lee and J. H. Park, “Incorporation of fermions into double field theory,” *JHEP* **1111** (2011) 025 [arXiv:1109.2035 [hep-th]].
- [16] I. Jeon, K. Lee and J. H. Park, “Supersymmetric Double Field Theory: Stringy Reformulation of Supergravity,” *Phys. Rev. D* **85** (2012) 081501 [Erratum-ibid. *D* **86** (2012) 089903] [arXiv:1112.0069 [hep-th]].
- [17] I. Jeon, K. Lee, J. H. Park and Y. Suh, “Stringy Unification of Type IIA and IIB Supergravities under $N = 2$ $D = 10$ Supersymmetric Double Field Theory,” *Phys. Lett. B* **723** (2013) 245 [arXiv:1210.5078 [hep-th]].
- [18] O. Hohm, C. Hull and B. Zwiebach, “Generalized metric formulation of double field theory,” *JHEP* **1008** (2010) 008 [arXiv:1006.4823 [hep-th]].
- [19] W. Cho, J. J. Fernandez-Melgarejo, I. Jeon and J. H. Park, “Supersymmetric gauged double field theory: systematic derivation by virtue of twist,” *JHEP* **1508** (2015) 084 [arXiv:1505.01301 [hep-th]].

- [20] D. Geissbuhler, “Double Field Theory and N=4 Gauged Supergravity,” JHEP **1111** (2011) 116 [arXiv:1109.4280 [hep-th]].
- [21] G. Aldazabal, W. Baron, D. Marques and C. Nunez, “The effective action of Double Field Theory,” JHEP **1111** (2011) 052 Erratum: [JHEP **1111** (2011) 109] [arXiv:1109.0290 [hep-th]].
- [22] M. Grana and D. Marques, “Gauged Double Field Theory,” JHEP **1204** (2012) 020 [arXiv:1201.2924 [hep-th]].
- [23] G. Dibitetto, J. J. Fernandez-Melgarejo, D. Marques and D. Roest, “Duality orbits of non-geometric fluxes,” Fortsch. Phys. **60** (2012) 1123 [arXiv:1203.6562 [hep-th]].
- [24] K. S. Choi and J. H. Park, “Standard Model as a Double Field Theory,” Phys. Rev. Lett. **115** (2015) no.17, 171603 [arXiv:1506.05277 [hep-th]].
- [25] K. Lee and J. H. Park, “Covariant action for a string in doubled yet gauged spacetime,” Nucl. Phys. B **880** (2014) 134 [arXiv:1307.8377 [hep-th]].
- [26] S. M. Ko, C. Melby-Thompson, R. Meyer and J. H. Park, “Dynamics of Perturbations in Double Field Theory & Non-Relativistic String Theory,” JHEP **1512** (2015) 144 [arXiv:1508.01121 [hep-th]].
- [27] J. H. Park, “Green-Schwarz superstring on doubled-yet-gauged spacetime,” JHEP **1611** (2016) 005 [arXiv:1609.04265 [hep-th]].
- [28] J. Gomis and H. Ooguri, “Nonrelativistic closed string theory,” J. Math. Phys. **42** (2001) 3127 doi:10.1063/1.1372697 [hep-th/0009181].
- [29] As a matter of fact, the Gomis-Ooguri non-relativistic string theory [28] can be derived from the doubled-yet-gauged string action (30) with a non-Riemannian DFT-metric of the type (1, 1) [26].
- [30] The numerators in the quotient groups of (17) are not target-spacetime T-duality group: they act on the local Lorentz vector indices and correspond to the stabilizer of the defining relation of the DFT-vielbeins (5).
- [31] J. Maharana and J. H. Schwarz, “Noncompact symmetries in string theory,” Nucl. Phys. B **390** (1993) 3 [hep-th/9207016].
- [32] K. Peeters, “Introducing Cadabra: A Symbolic computer algebra system for field theory problems,” hep-th/0701238.
- [33] I. Jeon, K. Lee and J. H. Park, “Double field formulation of Yang-Mills theory,” Phys. Lett. B **701** (2011) 260 [arXiv:1102.0419 [hep-th]].
- [34] O. Hohm and S. K. Kwak, “Double Field Theory Formulation of Heterotic Strings,” JHEP **1106** (2011) 096 [arXiv:1103.2136 [hep-th]].
- [35] D. S. Berman and K. Lee, “Supersymmetry for Gauged Double Field Theory and Generalised Scherk-Schwarz Reductions,” Nucl. Phys. B **881** (2014) 369 [arXiv:1305.2747 [hep-th]].
- [36] O. Hohm, A. Sen and B. Zwiebach, “Heterotic Effective Action and Duality Symmetries Revisited,” JHEP **1502** (2015) 079 [arXiv:1411.5696 [hep-th]].
- [37] E. Malek, “From Exceptional Field Theory to Heterotic Double Field Theory via K3,” JHEP **1703** (2017) 057 [arXiv:1612.01990 [hep-th]].
- [38] G. Aldazabal, E. Andres, M. Mayo and V. Penas, “Double Field Theory description of Heterotic gauge symmetry enhancing-breaking,” JHEP **1710** (2017) 046 [arXiv:1708.07148 [hep-th]].
- [39] With (13), we can rewrite the Yang-Mills kinetic term, $-\frac{1}{8}\mathcal{H}^{AC}\mathcal{H}^{BD}F_{AB}{}^{\dot{A}}F_{CD\dot{A}} = \frac{1}{2}P^{AC}\bar{P}^{BD}\mathcal{F}_{AB}{}^{\dot{A}}\mathcal{F}_{CD\dot{A}}$, where $\mathcal{F}_{AB}{}^{\dot{C}} := \nabla_A W_B{}^{\dot{C}} - \nabla_B W_A{}^{\dot{C}} + f_{\dot{A}\dot{B}}{}^{\dot{C}}W_A{}^{\dot{A}}W_B{}^{\dot{B}}$ [24, 33]. Irrespective of (13) $P_A{}^C\bar{P}_B{}^D\mathcal{F}_{CD}{}^{\dot{A}}$ is diffeomorphism-covariant, but $F_{AB}{}^{\dot{A}}$ is so if (13) holds [10].
- [40] S. F. Hassan and A. Sen, “Twisting classical solutions in heterotic string theory,” Nucl. Phys. B **375** (1992) 103 [hep-th/9109038].
- [41] E. Bergshoeff, M. de Roo, B. de Wit and P. van Nieuwenhuizen, “Ten-Dimensional Maxwell-Einstein Supergravity, Its Currents, and the Issue of Its Auxiliary Fields,” Nucl. Phys. B **195** (1982) 97.
- [42] G. F. Chapline and N. S. Manton, “Unification of Yang-Mills Theory and Supergravity in Ten-Dimensions,” Phys. Lett. B **120** (1983) 105.
- [43] M. B. Green and J. H. Schwarz, “Anomaly Cancellation in Supersymmetric D=10 Gauge Theory and Superstring Theory,” Phys. Lett. **149B** (1984) 117.
- [44] O. A. Bedoya, D. Marques and C. Nunez, “Heterotic α' -corrections in Double Field Theory,” JHEP **1412** (2014) 074 [arXiv:1407.0365 [hep-th]].
- [45] A. Coimbra, R. Minasian, H. Triendl and D. Waldram, “Generalised geometry for string corrections,” JHEP **1411** (2014) 160 [arXiv:1407.7542 [hep-th]].
- [46] K. Lee, “Quadratic α' -corrections to heterotic double field theory,” Nucl. Phys. B **899** (2015) 594 [arXiv:1504.00149 [hep-th]].
- [47] C. M. Hull, “Doubled Geometry and T-Folds,” JHEP **0707** (2007) 080 [hep-th/0605149].
- [48] A. S. Arvanitakis and C. D. A. Blair, “Unifying Type-II Strings by Exceptional Groups,” Phys. Rev. Lett. **120** (2018) no.21, 211601 [arXiv:1712.07115 [hep-th]].
- [49] A. S. Arvanitakis and C. D. A. Blair, “The Exceptional Sigma Model,” JHEP **1804** (2018) 064 [arXiv:1802.00442 [hep-th]].
- [50] J. H. Park, “Comments on double field theory and diffeomorphisms,” JHEP **1306** (2013) 098 [arXiv:1304.5946 [hep-th]].
- [51] A. Giveon and M. Rocek, “Generalized duality in curved string backgrounds,” Nucl. Phys. B **380** (1992) 128 [hep-th/9112070].
- [52] For the reduction of the type II critical superstring [17, 27] to Minkowskian four-dimensions ($\hat{D} = 10 \rightarrow D = 4$), we may consider $D' = 6$, (3, 3)-type non-Riemannian internal space, as suggested previously in [14]. This will set the twofold spin groups to be equally ten-dimensional, and the corresponding coset structure to be $\frac{\mathbf{O}(10,10)}{\mathbf{O}(4,6) \times \mathbf{O}(6,4)}$, which contrasts with the asymmetric spin groups in (17).
- [53] D. S. Berman, C. D. A. Blair and R. Otsuki, “Non-Riemannian geometry of M-theory,” arXiv:1902.01867 [hep-th].
- [54] O. Hohm and H. Samtleben, “Leibniz-Chern-Simons Theory and Phases of Exceptional Field Theory,” arXiv:1805.03220 [hep-th].