ENTROPY DIMENSION FOR DETERMINISTIC WALKS IN RANDOM SCENERIES

DOU DOU AND KYEWON KOH PARK

Abstract. Entropy dimension is an entropy-type quantity which takes values in [0, 1] and classifies different levels of intermediate growth rate of complexity for dynamical systems. In this paper, we consider the complexity of skew products of irrational rotations with Bernoulli systems, which can be viewed as deterministic walks in random sceneries, and show that this class of models can have any given entropy dimension by choosing suitable rotations for the base system.

1. Introduction

Complexity for dynamical systems describes the growth rate of orbits under the iteration of the actions. We measure the complexity via the iterated partitions in a probability measure space and via the iterated open covers in a topological setting. If a system has positive entropy, then it has exponential growth rate. Positive entropy systems have been studied and much understood in recent decades and they are known to be chaotic and unpredictable. Hence the word complexity is used mostly for entropy zero systems. Many examples of polynomial growth rate, such as irrational rotations, adding machines, interval exchange maps and some billiards on polygons, are wellknown. However the systems of sub-exponential but super-polynomial growth rate (called intermediate growth rate) are not well understood yet, and hence are wide open to further development.

In the study of more general group actions, for example, \mathbb{Z}^2 -systems, the subexponential growth arises naturally and some of their properties have been investigated in [\[19,](#page-17-0) [20\]](#page-17-1) and also in [\[15\]](#page-17-2). The notions of slow entropy or entropy dimension have been introduced to distinguish different levels of intermediate growth rate of zero entropy systems ([\[16,](#page-17-3) [7,](#page-17-4) [12,](#page-17-5) [13\]](#page-17-6)). For \mathbb{Z} -actions, several definitions of entropy dimensions for topological and measurable dynamical systems are introduced and investigated in [\[7,](#page-17-4) [12,](#page-17-5) [9,](#page-17-7) [10,](#page-17-8) [3,](#page-17-9) [11\]](#page-17-10).

In [\[6\]](#page-17-11), a class of uniformly recurrent infinite words with given intermediate complexity was constructed and it was shown in [\[3\]](#page-17-9) that the systems generated by the words are uniquely ergodic. Moreover it was shown in [\[3\]](#page-17-9) that entropy dimension does not have the property of variational principle. In $[9, 10]$ $[9, 10]$, systems of uniform entropy dimension (u.d., for short) were constructed. They have the property that all finite open covers have the same topological entropy dimension which corresponds to topological property

²⁰¹⁰ Mathematics Subject Classification. Primary: 37A35, 37A05, 54H20.

Key words and phrases. entropy dimension, complexity, deterministic walk, sub-exponential growth.

u.p.e. or all finite measurable partitions have the same metric entropy dimension which corresponds to the metric property of K-mixing. Since the study of systems with intermediate complexity is still at the beginning stage, we have many interesting and basic questions. Besides the properties mentioned above, very little is known in the way of general results and the developments have been focused on examples. Some examples related to intermediate complexity can be found in [\[14,](#page-17-12) [18\]](#page-17-13), but all of theses examples are via constructive methods to have desired properties. Hence we would like to provide with examples of intermediate growth rate out of more familiar models.

Aaronson showed the relative complexity of intermediate growth rate for a class of random walks in random sceneries $([1])$ $([1])$ $([1])$. However the model itself has positive entropy (exponential growth rate) since the base is Bernoulli. We consider a class of models which we call deterministic walks in random sceneries (DWRS, for short). The deterministic walks are chosen to be the irrational rotations of the unit circle T and the random sceneries are chosen to be the orbits of Bernoulli systems (or more generally, positive entropy systems). One can refer to [\[2,](#page-17-15) [4,](#page-17-16) [8\]](#page-17-17) for DWRS's on \mathbb{Z} .

In our DWRS models, the unit circle of the base is divided into two subintervals of the same length. If after a rotation, a base point lies in the first interval, it will walk forward along its scenery and walk backward otherwise. Due to the construction, at time n, the point can only walk $o(n)$ distance away from the starting point along the orbit of the scenery system, hence the entropy should be zero. The scenery supplies the "randomness" for this skew-product system and the base controls the growth rate of the complexity. By choosing suitable irrational rotation number α of the base, we will show that our DWRS model has intermediate growth rate. More precisely, for any $\tau \in [0, 1]$, there exists α such that the skew-product system has the upper entropy dimension τ .

We should notice that there are several ways to give the definitions of entropy dimensions. In this paper, we proceed with those appeared in [\[3,](#page-17-9) [9,](#page-17-7) [10,](#page-17-8) [11\]](#page-17-10) via entropy generating sequences. Due to the choice of taking lim sup or lim inf in the definitions of entropy dimensions, we have upper entropy dimensions and lower entropy dimensions. It is shown in [\[9,](#page-17-7) [10\]](#page-17-8) that both of them are isomorphism invariants. However it is not hard to construct non-isomorphic systems of the same upper entropy dimension. In most of constructive examples we can increase upper dimension by adding more independencies and also decrease lower dimension by adding more repetitions. In fact making the upper and lower dimensions agree demands more careful choices of parameters at the successive steps $(12, 11, 10)$ $(12, 11, 10)$ $(12, 11, 10)$. We remark that this is unlike computing the entropy of a system, which can never increase in longer blocks(names). When we consider factor maps between systems, the so called localization theory in dynamics or the u.d. property, the upper entropy dimension appears to carry more meaning ([\[9,](#page-17-7) [10\]](#page-17-8)). Hence in this paper, when we talk about entropy dimension, it refers to the upper entropy dimension.

The paper is organized as follows. In section 2, we will give the definitions of entropy dimensions via entropy generating sequences in both topological settings and measure-theoretic settings. Together with a brief review on the best approximation and continued fraction expansion of irrationals, we will also introduce the construction and some preliminaries of the DWRS models. In section 3, we will discuss some recurrence properties of the irrational rotation and in section 4 and 5, we will show the existence of any given topological and metric entropy dimensions by choosing the rotations properly.

2. Preliminaries

2.1. **Entropy dimension.** Let $S = \{s_1 \leq s_2 \leq \cdots\}$ be an increasing sequence of integers. We define the upper dimension and the lower dimension of the sequence S by

$$
\overline{D}(S) = \limsup_{n \to \infty} \frac{\log n}{\log s_n}
$$

and

$$
\underline{D}(S) = \liminf_{n \to \infty} \frac{\log n}{\log s_n}
$$

respectively. Equivalently,

$$
\overline{D}(S) = \inf\{\tau \ge 0 : \limsup_{n \to \infty} \frac{n}{(s_n)^{\tau}} = 0\} = \sup\{\tau \ge 0 : \limsup_{n \to \infty} \frac{n}{(s_n)^{\tau}} = \infty\}
$$

and

$$
\underline{D}(S) = \inf \{ \tau \ge 0 : \liminf_{n \to \infty} \frac{n}{(s_n)^{\tau}} = 0 \} = \sup \{ \tau \ge 0 : \liminf_{n \to \infty} \frac{n}{(s_n)^{\tau}} = \infty \}.
$$

Clearly $0 \leq \underline{D}(S) \leq \overline{D}(S) \leq 1$. When $\overline{D}(S) = \underline{D}(S) = \tau$, we say the sequence S has dimension τ . For example, the dimension of the sequence $\{1, 4, 9, \cdots, n^2, \cdots\}$ is $\frac{1}{2}$ and more general, the dimension of the sequence $\{n^t\}$ (*t* is a positive integer) is $\frac{1}{t}$.

By a topological dynamical system (TDS, for short) we mean a pair (X, T) where X is a compact metric space and T is a continuous onto self-map on X. Let (X, T) be a TDS and U be a finite open cover of X. We say an increasing sequence of positive integers $S = \{s_1 < s_2 < \cdots\}$ is an entropy generating sequence of U if

$$
\liminf_{n \to \infty} \frac{1}{n} \log N(\bigvee_{i=1}^{n} T^{-s_i} \mathcal{U}) > 0,
$$

where $N(\mathcal{V})$ denotes the minimal cardinality of subcovers of the open cover \mathcal{V} of X.

Then the *(upper)* entropy dimension of U is defined by

 $\overline{D}(T,\mathcal{U}) = \sup \{ \overline{D}(S) : S \text{ is an entropy generating sequence of } \mathcal{U} \}.$

And the *(upper)* topological entropy dimension of the TDS (X, T) is defined by

$$
\overline{D}(X,T) = \sup_{\mathcal{U}} \overline{D}(T,\mathcal{U}),
$$

where the supremum is taken over all the finite open covers $\mathcal U$ of X.

Remark 2.1. The following will be used for the computation of topological entropy dimension which appeared in [\[9\]](#page-17-7).

4 D. Dou and K.K. Park

(1) There are alternative equivalent definitions for $\overline{D}(T,\mathcal{U})$

$$
\overline{D}(T,\mathcal{U}) = \inf \{ \tau \ge 0 : \limsup_{n \to \infty} \frac{1}{n^{\tau}} N(\bigvee_{i=0}^{n-1} T^{-i} \mathcal{U}) = 0 \}
$$

$$
= \sup \{ \tau \ge 0 : \limsup_{n \to \infty} \frac{1}{n^{\tau}} N(\bigvee_{i=0}^{n-1} T^{-i} \mathcal{U}) = \infty \}.
$$

(2) If U is a generating open cover, i.e. $\lim_{n\to\infty} \text{diam}(\bigvee_{i=0}^{n-1} T^{-i}U) = 0$, then $\overline{D}(X,T) = \overline{D}(T,\mathcal{U}).$

Similarly, we can define entropy dimension for measurable dynamical systems. By a measurable dynamical system (MDS, for short), we mean a quadruple (X, \mathcal{B}, μ, T) , where (X, \mathcal{B}, μ) is a probability space and T is a measurable transformation on X that preserves μ . Let (X, \mathcal{B}, μ, T) be a MDS and \mathcal{P} be a finite measurable partition of X. An increasing sequence of positive integers $S = \{s_1 < s_2 < \cdots\}$ is called an *entropy* generating sequence of P (w.r.t. μ) if

$$
\liminf_{n \to \infty} \frac{1}{n} H_{\mu}(\bigvee_{i=1}^{n} T^{-s_i} \mathcal{P}) > 0.
$$

Then the *(upper)* entropy dimension of P is defined by

 $\overline{D}_{\mu}(T,\mathcal{P}) = \sup \{ \overline{D}(S) : S \text{ is an entropy generating sequence of } \mathcal{P} \text{ (w.r.t } \mu \text{)} \}.$

The (upper) entropy dimension of the MDS (X, \mathcal{B}, μ, T) is defined by

$$
\overline{D}_{\mu}(X,T) = \sup_{\mathcal{P}} \overline{D}_{\mu}(T,\mathcal{P}),
$$

where the supremum is taken over all the finite measurable partitions P of X. It is shown in [\[10\]](#page-17-8), $D_{\mu}(X,T)$ equals $D_{\mu}(T,\mathcal{P})$ when $\mathcal P$ is a generating partition.

2.2. DWRS. Let $\mathbf{T} = \mathbb{R}/\mathbb{Z}$ be the unit circle and $\alpha \in (0, 1)$ be an irrational number. Let (T, α) be the rotation by α on the unit circle. For any $\theta \in T$, we define a bi-infinite $\{-1, 1\}$ sequence $z = z(\theta) = (\cdots, z_{-1}, z_0, z_1 \cdots)$ by

$$
z_n = \begin{cases} 1 & \text{if } \theta + n\alpha \mod 1 \in [0, \frac{1}{2}) \\ -1 & \text{if } \theta + n\alpha \mod 1 \in [\frac{1}{2}, 1) \end{cases}.
$$

Let $Z = cl\{\sigma^n z(0) : n \in \mathbb{Z}\}\,$, where σ is the left shift map. Then (Z, σ) forms a TDS which is called a Sturmian flow of type $(\alpha, \frac{1}{2})$ (see [\[21,](#page-17-18) (2.53)]). Since α and $\frac{1}{2}$ are rationally independent, (Z, σ) is minimal and uniquely ergodic. Moreover, (Z, σ) is an almost 1-1 extension of the irrational rotation (\mathbf{T}, α) . Let φ be the factor map between (Z,σ) and (\mathbf{T},α) . It is known that $\varphi^{-1}(\theta)$ has exact two points if $\theta \in Orb(0)\cup Orb(\frac{1}{2})$ $(\frac{1}{2})$ and $\varphi^{-1}(\theta) = z(\theta)$ otherwise.

Let $\mathcal{B}_{\mathbf{T}}$ and \mathcal{B}_{Z} be the Borel σ -algebra of **T** and Z respectively. Let **m** and μ be the unique ergodic invariant probability measures of (T, α) and (Z, σ) respectively. Then

 is the Haar-Lebesgue measure of $**T**$ **and moreover, from the measure-theoretic point** of view, the MDS's $(\mathbf{T}, \mathcal{B}_{\mathbf{T}}, \mathbf{m}, \alpha)$ and $(Z, \mathcal{B}_Z, \mu, \sigma)$ are conjugate.

Sometimes we will not distinguish \bf{T} and \bf{Z} in this paper.

Let (Y, S) be an invertible TDS. We define the map $T : Z \times Y \to Z \times Y$ by

$$
T(z, y) = (\sigma z, S^{z_0} y),
$$

for $z = (\dots, z_{-1}z_0z_1\dots) \in Z$. That is, for any point $(z, y) \in Z \times Y$, if z_0 is 1, we then push y forward to Sy , otherwise we pull y backward to $S^{-1}y$. Hence we have constructed a skew-product system $(Z \times Y, T)$. For measure-theoretic case, for an invertible MDS (Y, \mathcal{D}, ν, S) , this skew-product system can also be viewed as a deterministic walk along the orbits of (Y, \mathcal{D}, ν, S) .

For simplicity, we let 2) denote either the invertible TDS (Y, S) or the invertible MDS (Y, \mathcal{D}, ν, S) . And denote by $\mathfrak{X} = \mathfrak{X}_{\alpha, \mathfrak{Y}}$ either the TDS $(Z \times Y, T)$ or the MDS $(Z \times Y, \mathcal{B}_Z \otimes \mathcal{D}, \mu \times \nu, T)$ and call it a deterministic walk with rotation α along \mathfrak{Y} . Especially, when $\mathfrak Y$ is a Bernoulli system (or a positive entropy system), we call $\mathfrak X$ a DWRS.

Remark 2.2. Since $(T, \mathcal{B}_T, m, \alpha)$ and $(Z, \mathcal{B}_Z, \mu, \sigma)$ are conjugate, sometimes \mathfrak{X} is also viewed as $(\mathbf{T} \times Y, \mathcal{B}_{\mathbf{T}} \otimes \mathcal{D}, \mathbf{m} \times \nu, T)$. In this case, T also denotes the corresponding skew-product map on $\mathbf{T} \times Y$.

It is easy to see that the following hold and we omit the proofs.

Proposition 2.3. \mathfrak{X} is minimal (respectively, transitive or ergodic) if and only if \mathfrak{Y} is.

Proposition 2.4. For every $\alpha \in \mathbf{T}$ and any system \mathfrak{Y} , the entropy of \mathfrak{X} is zero.

Let $z = (\dots, z_{-1}z_0z_1 \dots) \in Z$. For any $m, n \in \mathbb{N}$ with $m \leq n$, denote by

$$
\omega(z,m,n) = \sum_{i=m}^{n} z_i, \ \omega(z,1,n) = \omega(z,n).
$$

For $\theta \in \mathbf{T}$, denote by

$$
\omega(\theta,m,n)=\omega(z(\theta),m,n),\quad \omega(\theta,n)=\omega(z(\theta),n)=\omega(\theta,1,n)
$$

and

$$
\omega(n) = \omega(0, n) = \omega(z(0), n).
$$

If $\varphi(z)$ does not belong to the orbits of 0 and $\frac{1}{2}$, then $\omega(z, k) = \omega(\varphi(z), k)$. Otherwise, $|\omega(z, k) - \omega(\varphi(z), k)| \leq 2.$

Denote by

$$
M(\theta, n) = \max_{1 \le \ell \le n} \omega(\theta, \ell), \quad m(\theta, n) = \min_{1 \le \ell \le n} \omega(\theta, \ell)
$$

and

$$
M(z,n)=\max_{1\leq \ell\leq n}\omega(z,\ell),\ \ m(z,n)=\min_{1\leq \ell\leq n}\omega(z,\ell).
$$

If $\varphi(z)$ does not belong to the orbits of 0 and $\frac{1}{2}$, then $M(z,n) = M(\varphi(z),n)$ and $m(z, n) = m(\varphi(z), n)$. Otherwise, $|M(z, n) - M(\varphi(z), n)| \leq 2$ and $|m(z, n) - M(\varphi(z), n)| \leq 2$ $m(\varphi(z), n)| \leq 2.$

Let $\theta \in \mathbf{T}$ and $z = z(\theta) = (\cdots, z_{-1}z_0z_1 \cdots) \in Z$. Recall that $T(z, y) = (\sigma z, S^{z_0}y)$. Hence, for $i > 0$,

(2.1)
$$
T^{i}(z, y) = (\sigma^{i} z, S^{\omega(z, 0, i-1)} y),
$$

$$
T^{-i}(z, y) = (\sigma^{-i} z, S^{-\omega(\sigma^{-i} z, 0, i-1)} y),
$$

where $\omega(z, 0, -1) := 0$.

2.3. The best approximation and continued fraction expansion of irrationals. Fix an irrational number $\alpha \in (0,1)$. The growth of $\omega(n)$ depends on the recurrent properties of α . Here we collect some basic facts on the continued fraction expansion of irrationals (see Khinchin's classic book [\[17\]](#page-17-19) for details).

For a real number β , let $\|\beta\| = \min\{\{\beta\}, 1 - \{\beta\}\}\$, where $\{\beta\}$ denotes the fractional part of β .

Let $[a_0; a_1, a_2, \dots]$ be the continued fraction expansion of α and $\frac{p_n}{q_n} = [a_0; a_1, a_2, \dots, a_n]$ be the continued fraction approximation of α . Then

$$
\frac{p_0}{q_0} < \frac{p_2}{q_2} < \dots < \alpha < \dots < \frac{p_3}{q_3} < \frac{p_1}{q_1}
$$

and

$$
(2.2) \t\t q_n = a_n q_{n-1} + q_{n-2},
$$

where $q_{-1} = 0$ and $q_{-2} = 1$. Since α is chosen from $(0, 1)$, we have that $a_0 = 0$ and $q_0 = 1$. Moreover, p_n, q_n 's have the following properties:

(2.3)
$$
||q_n\alpha|| = \begin{cases} \{q_n\alpha\} & \text{if } n = 2k, \\ 1 - \{q_n\alpha\} & \text{if } n = 2k + 1; \end{cases}
$$

(2.4)
$$
||q_{n-1}\alpha|| < ||k\alpha||, \forall k < q_n, k \neq q_{n-1};
$$

(2.5)
$$
\frac{1}{q_n(q_n + q_{n+1})} \le (-1)^n (\alpha - \frac{p_n}{q_n}) \le \frac{1}{q_n q_{n+1}}.
$$

3. SOME RECURRENCE PROPERTIES OF (T, α)

Let α be a fixed irrational in [0, 1). In this section we will discuss the growth of $\omega(\theta, n)$ for $\theta \in \mathbf{T}$, which will be related with the complexity of our DWRS models. Firstly, let's study the growth of $\omega(n)$. For convenience, we will identify **T** with the interval $[0, 1)$. When we talk about some point $\ell \alpha$, it should be understood as $\ell \alpha$ mod 1 due to the context. And when we talk about some interval, it should be understood as an arc of **T**. For each $n \in \mathbb{N}$, let $Q_n = \{k\alpha : 1 \leq k \leq q_n\} \subset \mathbf{T}$.

The following two propositions will show that the points in Q_n have "almost equidistribution" properties on **T** with respect to any translation of the partition $\{[0, \frac{1}{2}$ $(\frac{1}{2}),$ $\left[\frac{1}{2}\right]$ $\frac{1}{2}, 1) \}.$

Proposition 3.1.

$$
\omega(q_n) = \begin{cases} 0 & \text{if } q_n = 2r, \\ 1 & \text{if } q_n = 2r + 1 \text{ and } n = 2k, \\ -1 & \text{if } q_n = 2r + 1 \text{ and } n = 2k + 1. \end{cases}
$$

Proof. We will just prove the case $n = 2k$ since the parallel argument holds for $n =$ $2k + 1$.

We note that $||q_n\alpha|| = \{q_n\alpha\}.$

For $1 \leq \ell \leq q_n-1$, if $\ell \alpha \in (0, \frac{1}{2})$ $(\frac{1}{2}), \text{ then } -\ell \alpha \in (\frac{1}{2})$ $\frac{1}{2}$, 1). Hence $(q_n - \ell)$ α ∈ $(\frac{1}{2} + ||q_n \alpha||, 1 + \frac{1}{2})$ $\|q_n\alpha\|$). Since no element in \overline{Q}_n belongs to $[0, \|q_n\alpha\|)$, we know that $(q_n - \ell)\alpha \in$ $\left(\frac{1}{2} + ||q_n\alpha||, 1\right)$. Likewise, if $\ell\alpha \in \left(\frac{1}{2} + ||q_n\alpha||, 1\right)$, then $(q_n - \ell)\alpha \in (0, \frac{1}{2})$ $\frac{1}{2}$). Moreover, $\ell \alpha \in (\frac{1}{2})$ $\frac{1}{2}, \frac{1}{2} + ||q_n \alpha||$) if and only if $(q_n - \ell) \alpha \in (\frac{1}{2})$ $\frac{1}{2}, \frac{1}{2} + ||q_n \alpha||$.

If $q_n = 2r$, then $r\alpha = \frac{1}{2} + \frac{1}{2}$ $\frac{1}{2}||q_n\alpha|| \in \left(\frac{1}{2}\right)$ $\frac{1}{2}, \frac{1}{2} + ||q_n \alpha||$). Note that $q_n \alpha \in [0, \frac{1}{2}]$ $(\frac{1}{2})$, hence $\omega(q_n) = 0$. If $q_n = 2r + 1$, then no element of the set Q_n belongs to $(\frac{1}{2}, \frac{1}{2} + ||q_n \alpha||)$ since otherwise there will be two elements of Q_n belonging to this interval, which contradicts to the fact that $\frac{p_n}{q_n}$ is the best approximation of α . Hence $\omega(q_n) = 1$.

Proposition 3.2. For $\theta \in \mathbf{T}$, the following hold.

- (1) If $\omega(q_n) = 0$, then $\omega(\theta, q_n) = 0$ or ± 2 .
- (2) If $\omega(q_n) = \pm 1$, then $\omega(\hat{\theta}, q_n) = \pm 1$ or ± 3 .
- (3) If $q_n = 2r + 1$ and $kq_n < q_{n+1}$, then $|\omega(kq_n)| \leq k$.

Proof. (1). Suppose $\omega(q_n) = 0$. If $\theta = 0$ or $\frac{1}{2}$, obviously $\omega(\theta, q_n) = 0$. Now we treat the case for $0 < \theta < \frac{1}{2}$ and the case for $\frac{1}{2} < \theta < 1$ is similar. Then

$$
\omega(\theta, q_n) = #((\theta + Q_n) \cap [0, \frac{1}{2})) - #((\theta + Q_n) \cap [\frac{1}{2}, 1))
$$

\n
$$
= #\left(Q_n \cap ([-\theta, 0) \cup [0, \frac{1}{2} - \theta)\right) - #\left(Q_n \cap ([\frac{1}{2} - \theta, \frac{1}{2}) \cup [\frac{1}{2}, -\theta)\right)\right)
$$

\n
$$
= #(Q_n \cap [-\theta, 0)) + #(Q_n \cap [0, \frac{1}{2})) - #(Q_n \cap [\frac{1}{2} - \theta, \frac{1}{2}))
$$

\n
$$
- #(Q_n \cap [\frac{1}{2} - \theta, \frac{1}{2})) - #(Q_n \cap [\frac{1}{2}, 1)) + #(Q_n \cap [-\theta, 0)).
$$

Since $\omega(q_n) = \#(Q_n \cap [0, \frac{1}{2})]$ $(\frac{1}{2})$) – # $(Q_n \cap [\frac{1}{2})]$ $(\frac{1}{2}, 1)$ = 0, we have that

(3.1)
$$
\omega(\theta, q_n) = 2\left(\#(Q_n \cap [-\theta, 0)) - \#(Q_n \cap [\frac{1}{2} - \theta, \frac{1}{2}))\right).
$$

From Proposition [3.1,](#page-6-0) $\omega(q_n) = 0$ if and only if $q_n = 2r$ for some r. As before we will just discuss for the case $n = 2k$. Note that in this case, $||q_n\alpha|| = \{q_n\alpha\}$ and $r\alpha = \frac{1}{2} + \frac{1}{2}$ $\frac{1}{2}||q_n\alpha||$. Hence $Q_n \cap [\frac{1}{2} - \frac{1}{2}]$ $\frac{1}{2}||q_n\alpha||, \frac{1}{2}+\frac{1}{2}$ $\frac{1}{2}||q_n\alpha||)=\emptyset.$

For $0 < \ell \leq r$, we have that

(3.2)
$$
\ell\alpha \in [-\theta, 0] \Leftrightarrow (\ell + r)\alpha \in [\frac{1}{2} - \theta + \frac{1}{2} ||q_n \alpha||, \frac{1}{2} + \frac{1}{2} ||q_n \alpha||)
$$

$$
\Leftrightarrow (\ell + r)\alpha \in [\frac{1}{2} - \theta + \frac{1}{2} ||q_n \alpha||, \frac{1}{2})
$$

For $r < \ell \leq q_n$, we have that

(3.3)
$$
\ell\alpha \in [-\theta, 0) \Leftrightarrow (\ell - r)\alpha \in \left[\frac{1}{2} - \theta - \frac{1}{2}||q_n\alpha||, \frac{1}{2} - \frac{1}{2}||q_n\alpha||\right)
$$

$$
\Leftrightarrow (\ell - r)\alpha \in \left[\frac{1}{2} - \theta - \frac{1}{2}||q_n\alpha||, \frac{1}{2}\right)
$$

Notice that $\left[\frac{1}{2} - \theta - \frac{1}{2}\right]$ $\frac{1}{2}||q_n\alpha||, \frac{1}{2}-\theta+\frac{1}{2}$ $\frac{1}{2}||q_n\alpha||$) contains at most one point of Q_n (since the length of this interval is $||q_n\alpha||$. Together with [\(3.1\)](#page-6-1), [\(3.2\)](#page-7-0) and [\(3.3\)](#page-7-1), we conclude that

$$
\omega(\theta, q_n) = \begin{cases} 2, & \text{if } (\ell - r)\alpha \in \left[\frac{1}{2} - \theta - \frac{1}{2}\right] |q_n \alpha|, \frac{1}{2} - \theta) \text{ for some } r < \ell \le q_n; \\ -2, & \text{if } (\ell + r)\alpha \in \left[\frac{1}{2} - \theta, \frac{1}{2} - \theta + \frac{1}{2}\right] |q_n \alpha| \text{ for some } 0 < \ell \le r; \\ 0, & \text{otherwise.} \end{cases}
$$

(2). We just consider the case for $\omega(q_n) = 1$ and the case for $\omega(q_n) = -1$ is similar. By Proposition [3.1,](#page-6-0) if $\omega(q_n) = 1$ then $n = 2k$ and $q_n = 2r + 1$.

It is clear that $\omega(\theta, q_n) = 1$ if $\theta = 0$ and $\omega(\theta, q_n) = -1$ if $\theta = \frac{1}{2}$ $\frac{1}{2}$. Now assume $0 < \theta < \frac{1}{2}$. Similar with [\(3.1\)](#page-6-1), we have

(3.4)
$$
\omega(\theta, q_n) = 2\left(\#(Q_n \cap [-\theta, 0)) - \#(Q_n \cap [\frac{1}{2} - \theta, \frac{1}{2}))\right) + 1.
$$

Assume that $\frac{1}{2} \in (s_1\alpha, s_2\alpha)$, where $s_1\alpha, s_2\alpha \in Q_n$ and $(s_1\alpha, s_2\alpha) \cap Q_n = \emptyset$. If $s_2 \leq q_n-q_{n-1}$, then $(s_2+q_{n-1})\alpha \in Q_n$ and no element of Q_n can lie between $(s_2+q_{n-1})\alpha$ and $s_2\alpha$. Hence $s_1 = s_2 + q_{n-1}$ and $|s_2\alpha - s_1\alpha| = ||q_{n-1}\alpha||$. If $s_2 > q_n - q_{n-1}$, then $(s_2 + q_{n-1} - q_n)\alpha \in Q_n$ and no element of Q_n can lie between $(s_2 + q_{n-1} - q_n)\alpha$ and $s_2\alpha$. Hence $s_1 = s_2 + q_{n-1} - q_n$ and $|s_2\alpha - s_1\alpha| = ||q_{n-1}\alpha|| + ||q_n\alpha||$. So $(-\theta + s_1\alpha, -\theta + s_2\alpha)$ contains at most two elements in Q_n .

For $0 < \ell \leq q_n - s_2$, since $Q_n \cap \left[\frac{1}{2}\right]$ $(\frac{1}{2}, s_2 \alpha) = \emptyset$, we have that

(3.5)
$$
\ell \alpha \in [-\theta, 0] \Leftrightarrow (\ell + s_2) \alpha \in [-\theta + s_2 \alpha, s_2 \alpha)
$$

$$
\Leftrightarrow (\ell + s_2) \alpha \in [-\theta + s_2 \alpha, \frac{1}{2}).
$$

For $q_n - s_2 < \ell \le q_n$, since no element of Q_n lies between $s_2 \alpha - ||q_n \alpha||$ and $\frac{1}{2}$, we have that

(3.6)
$$
\ell \alpha \in [-\theta, 0) \Leftrightarrow (\ell + s_2 - q_n)\alpha \in [-\theta + s_2\alpha - ||q_n\alpha||, s_2\alpha - ||q_n\alpha||)
$$

$$
\Leftrightarrow (\ell + s_2 - q_n)\alpha \in [-\theta + s_2\alpha - ||q_n\alpha||, \frac{1}{2}).
$$

In the proof of Proposition [3.1](#page-6-0) (the last paragraph), we have shown that Q_n does not intersect with the interval $(\frac{1}{2}, \frac{1}{2} + ||q_n \alpha||)$ when $q_n = 2r + 1$. So $s_2 \alpha - ||q_n \alpha|| \ge \frac{1}{2}$. Noticing that $[-\theta + \frac{1}{2}]$ $\frac{1}{2}$, $-\theta + s_2\alpha$) contains at most two elements of Q_n and $[-\theta + \frac{1}{2}]$ $\frac{1}{2}$, $-\theta$ + $s_2\alpha - ||q_n\alpha||$ contains at most one element of Q_n , together with [\(3.4\)](#page-7-2), [\(3.5\)](#page-7-3) and [\(3.6\)](#page-7-4), we conclude that $\omega(\theta, q_n) = \pm 1$ or -3 .

If $\frac{1}{2} < \theta < 1$, then $\omega(\theta, q_n) = -\omega(\theta - \frac{1}{2})$ $(\frac{1}{2}, q_n)$, and hence $\omega(\theta, q_n) = \pm 1$ or 3.

(3). If $q_n = 2r + 1$ and $n = 2t$, i.e. $\omega(q_n) = 1$, then $||q_n\alpha|| = \{q_n\alpha\}$. We now order the points in Q_n on **T** by $0 < q_n \alpha < \cdots < s_1 \alpha < \frac{1}{2} < s_2 \alpha < \cdots < q_{n-1} \alpha < 1$. Noticing that $kq_n < q_{n+1}$, hence the points in $\{\ell \alpha\}_{\ell=1}^{kq_n}$ should be ordered as

$$
0 < q_n \alpha < 2q_n \alpha < \dots < kq_n \alpha
$$
\n
$$
\dots
$$
\n
$$
\alpha < s_1 \alpha < (s_1 + q_n) \alpha < \dots < (s_1 + (k-1)q_n) \alpha
$$
\n
$$
\alpha < s_2 \alpha < (s_2 + q_n) \alpha < \dots < (s_2 + (k-1)q_n) \alpha
$$
\n
$$
\dots
$$
\n
$$
\alpha < q_{n-1} \alpha < (q_{n-1} + q_n) \alpha < \dots < (q_{n-1} + (k-1)q_n) \alpha < 1.
$$

Due to the place where $\frac{1}{2}$ lies, obviously $|\omega(kq_n)| \leq k$.

Similarly, when $q_n = 2r + 1$ and $n = 2t + 1$, we also have $|\omega(kq_n)| \leq k$.

$$
\Box
$$

The following proposition gives estimation on $m(\theta, q_n)$ and $M(\theta, q_n)$.

Proposition 3.3. Let $\theta \in \mathbf{T}$, then we have the following estimation.

(1) For $q_n < t \leq q_{n+1}$, $M(\theta, t) \leq M(\theta, q_n) + 3\left[\frac{t}{a}\right]$ $\frac{t}{q_n}$. (2) For $q_n < t \leq q_{n+1}$, $m(\theta, t) \geq m(\theta, q_n) - 3\left[\frac{t}{q_n}\right]$ $\frac{t}{q_n}$. (3) max $\{M(0, q_{n+1}), |m(0, q_{n+1})|\} \geq |\omega(q_n)| \lfloor \frac{q_n+1}{6q_n} \rfloor$.

Proof. For any $0 < \ell \leq t$, we write $\ell = \ell_0 + kq_n$, where $0 \leq \ell_0 < q_n$ and $0 \leq k \leq \lfloor \frac{t}{q_n} \rfloor$. Then

$$
\omega(\theta,\ell) = \omega(\theta,\ell_0) + \omega(\theta,\ell_0+1,\ell_0+q_n) + \cdots + \omega(\theta,\ell_0+1+(k-1)q_n,\ell_0+kq_n)
$$

=
$$
\omega(\theta,\ell_0) + \omega(\theta+\ell_0\alpha,q_n) + \cdots + \omega(\theta+(\ell_0+(k-1)q_n)\alpha,q_n).
$$

Hence by Proposition [3.2,](#page-6-2)

$$
m(\theta, q_n) - 3k \le \omega(\theta, \ell) \le M(\theta, q_n) + 3k.
$$

This proves (1) and (2) .

Now we prove (3). Obviously (3) is true if $\omega(q_n) = 0$. Without loss of generality, we assume that $\omega(q_n) = 1$.

Suppose $s_1 \alpha < \frac{1}{2} < s_2 \alpha$, where $0 < s_1, s_2 < q_n$ and no element in Q_n lies between $s_1 \alpha$ and $s_2\alpha$. From the property of $\frac{p_n}{q_n}$ (inequality [\(2.4\)](#page-5-0)), $|s_1-s_2|\leq q_{n-1}$ and $||s_1\alpha-s_2\alpha||\geq$ $||q_{n-1}\alpha||$. If $\frac{1}{2} - \{s_1\alpha\} < ||q_n\alpha||$, then

(3.7)
$$
\omega(0, (k-1)q_n + 1, kq_n) = \begin{cases} 1, & \text{if } k = 1, \\ -1, & \text{if } k = 2, 3, \dots, \lfloor \frac{\|q_{n-1}\alpha\|}{\|q_n\alpha\|} \rfloor. \end{cases}
$$

If $\frac{1}{2} - \{s_1 \alpha\} \ge ||q_n \alpha||$, then

(3.8)
$$
\omega(0, (k-1)q_n + 1, kq_n) = \begin{cases} 1, & \text{if } k = 1, 2, \dots, \lfloor \frac{\frac{1}{2} - \{s_1 \alpha\}}{\|q_n \alpha\|} \rfloor, \\ -1, & \text{if } k = \lfloor \frac{\frac{1}{2} - \{s_1 \alpha\}}{\|q_n \alpha\|} \rfloor + 1, \dots, \lfloor \frac{\|q_{n-1} \alpha\|}{\|q_n \alpha\|} \rfloor. \end{cases}
$$

We also note that by inequality (2.5) ,

$$
\frac{||q_{n-1}\alpha||}{||q_n\alpha||} \ge \frac{1}{q_{n-1}(q_{n-1} + q_n)} q_n q_{n+1} \ge \lfloor \frac{q_{n+1}}{q_n} \rfloor \frac{q_n^2}{q_{n-1}(q_{n-1} + q_n)}
$$

$$
\ge \frac{1}{2} \lfloor \frac{q_{n+1}}{q_n} \rfloor \ge \lfloor \frac{q_{n+1}}{2q_n} \rfloor.
$$

This implies that

(3.9)
$$
\lfloor \frac{q_{n+1}}{6q_n} \rfloor \|q_n \alpha\| \le \frac{1}{3} \|q_{n-1} \alpha\|.
$$

We assume that $q_{n+1} \geq 6q_n$ (otherwise (3) already holds).

If $\frac{1}{2} - s_1 \alpha > \frac{1}{3} ||q_{n-1} \alpha|| \ge ||q_n \alpha||$, then $\left[\frac{\frac{1}{2} - \{s_1 \alpha\}}{||q_n \alpha||} \right] \ge \left[\frac{q_{n+1}}{6q_n} \right]$. By [\(3.8\)](#page-9-0), $\omega(kq_n) = k$ for any $k = 1, 2, \cdots, \lfloor \frac{\frac{1}{2} - \{s_1 \alpha\}}{\|a_n \alpha\|} \rfloor$ $\frac{-(s_1\alpha)}{\|q_n\alpha\|}$. So in this case $M(0, q_{n+1}) \geq |\omega(q_n)| \lfloor \frac{q_{n+1}}{6q_n} \rfloor$.

If $||q_n\alpha|| \leq \frac{1}{2} - s_1\alpha \leq \frac{1}{3}$ $\frac{1}{3} ||q_{n-1} \alpha||$, then by [\(3.8\)](#page-9-0), $\omega(kq_n) = 2 \lfloor \frac{\frac{1}{2} - \{s_1 \alpha\}}{||q_n \alpha||}$ $\frac{-\{s_1\alpha\}}{\|q_n\alpha\|}\big] - k$ for any $\frac{\frac{1}{2} - \{s_1 \alpha\}}{\|a - \alpha\|}$ $\frac{-\{s_1\alpha\}}{\|q_n\alpha\|}\big] < k \leq \lfloor \frac{\|q_{n-1}\alpha\|}{\|q_n\alpha\|}\big]$. So in this case

$$
m(0, q_{n+1}) \le \omega \left(\left\lfloor \frac{\|q_{n-1}\alpha\|}{\|q_n\alpha\|} \right\rfloor q_n \right) = 2 \left\lfloor \frac{\frac{1}{2} - \{s_1 \alpha\}}{\|q_n \alpha\|} \right\rfloor - \left\lfloor \frac{\|q_{n-1}\alpha\|}{\|q_n \alpha\|} \right\rfloor
$$

$$
\le - \left\lfloor \frac{q_{n+1}}{6q_n} \right\rfloor.
$$

If $\frac{1}{2} - s_1 \alpha \leq \frac{1}{3}$ $\frac{1}{3} ||q_{n-1}\alpha||$ and $\frac{1}{2} - \{s_1\alpha\} < ||q_n\alpha||$, then by [\(3.7\)](#page-9-1), it obviously holds that

$$
m(0, q_{n+1}) \le \omega \left(\left\lfloor \frac{\|q_{n-1}\alpha\|}{\|q_n\alpha\|} \right\rfloor q_n \right) = -\left\lfloor \frac{\|q_{n-1}\alpha\|}{\|q_n\alpha\|} \right\rfloor + 2
$$

$$
\le -\left\lfloor \frac{q_{n+1}}{6q_n} \right\rfloor.
$$

Denote

$$
M(k) = \max_{z \in Z} M(z, k) \text{ and } m(k) = \min_{z \in Z} m(z, k).
$$

Proposition 3.4. For $q_n < k \leq q_{n+1}$, we have

$$
M(k) \le q_1 + 3\lceil \frac{k}{q_n} \rceil + 3 \sum_{j=2}^n \lceil \frac{q_j}{q_{j-1}} \rceil + 2
$$

and

$$
m(k) \ge -q_1 - 3\lceil \frac{k}{q_n} \rceil - 3 \sum_{j=2}^n \lceil \frac{q_j}{q_{j-1}} \rceil - 2.
$$

Proof. By (1) of Proposition [3.3,](#page-8-0) for $q_n < k \leq q_{n+1}$,

$$
M(\theta, k) \le M(\theta, q_n) + 3\lceil \frac{k}{q_n} \rceil \le \cdots
$$

$$
\le M(\theta, q_1) + 3\lceil \frac{k}{q_n} \rceil + 3 \sum_{j=2}^n \lceil \frac{q_j}{q_{j-1}} \rceil
$$

$$
\le q_1 + 3\lceil \frac{k}{q_n} \rceil + 3 \sum_{j=2}^n \lceil \frac{q_j}{q_{j-1}} \rceil.
$$

Hence

$$
M(k) = \max_{z \in Z} M(z, k) \le \max_{\theta \in \mathbf{T}} M(\theta, k) + 2
$$

$$
\le q_1 + 3\lceil \frac{k}{q_n} \rceil + 3 \sum_{j=2}^n \lceil \frac{q_j}{q_{j-1}} \rceil + 2.
$$

The proof is similar for $m(k)$.

4. Topological entropy dimension

Let $\mathfrak{X} = \mathfrak{X}_{\alpha, \mathfrak{Y}}$ be the DWRS with rotation α along \mathfrak{Y} , where \mathfrak{Y} is an invertible TDS with positive entropy. In this section we will show that for any $\tau \in [0,1)$, there exists α such that $\mathfrak X$ has topological entropy dimension τ .

Let $\tau \in [0, 1)$ be given. For any positive integers $\{t_1, t_2, \dots, t_k\}$, we define an irrational number $\alpha(t_1, t_2, \dots, t_k) = [0; a_1, a_2, \dots] \in [0, 1)$ as follows:

(4.1)
$$
a_n = t_n
$$
 for $1 \le n \le k$;
$$
\begin{cases} a_{k+1} = a_{k+2} = 2, & \text{if } q_k q_{k-1} \text{ is odd,} \\ a_{k+1} = 3, a_{k+2} = 2, & \text{if } q_{k-1} \text{ is even and } q_k \text{ is odd,} \\ a_{k+1} = a_{k+2} = 3, & \text{if } q_k \text{ is even and } q_{k-1} \text{ is odd;} \end{cases}
$$

$$
a_{n+1} = 2 \lfloor q_n^{\frac{\tau}{1-\tau}} \rfloor \text{ for } n > k+2.
$$

By [\(2.2\)](#page-5-2), $q_{n+1} = a_{n+1}q_n + q_{n-1}$ is odd for every $n \geq k$. We remark here that if q_n is even then q_{n-1} must be odd (otherwise q_{n-2}, \dots, q_1, q_0 are all even, which contradicts to the fact $q_0 = 1$).

Denote by E_{τ} the collection of such irrational number $\alpha(t_1, t_2, \dots, t_k)$'s. Then E_{τ} is dense in $[0, 1)$.

Let $\alpha \in E_{\tau}$ be fixed. By [\(2.2\)](#page-5-2), a simple estimation shows that for any $c > 0$,

(4.2)
$$
\frac{q_{n+1}}{q_n} \sim a_{n+1} \sim 2q_n^{\frac{\tau}{1-\tau}} \gg n^c,
$$

when n is sufficiently large.

Let $l_n = \lfloor \frac{q_{n+1}}{12q_n} \rfloor$ $\frac{q_{n+1}}{12q_n}$. By (2.5) ,

$$
\frac{||q_{n-1}\alpha||}{||q_n\alpha||} \ge \frac{\frac{1}{q_{n-1}+q_n}}{\frac{1}{q_{n+1}}} \ge \frac{q_{n+1}}{2q_n}.
$$

Hence

(4.3)
$$
l_n \leq \lfloor \frac{||q_{n-1}\alpha||}{6||q_n\alpha||} \rfloor.
$$

 ∞

Moreover, by [\(4.2\)](#page-11-0), there exists n_0 such that $l_n > 0$ and $\sum_{i=n_0}^{n-1} l_i < l_n$ whenever $n \ge n_0$. Now we set

(4.4)
$$
F = \bigcup_{n=n_0} \{q_n, 2q_n, \cdots, l_n q_n\} = \{s_1 < s_2 < \cdots\}.
$$

Lemma 4.1.

$$
\overline{D}(F) \geq \tau.
$$

Proof. From the definition of F,

$$
\overline{D}(F) = \limsup_{n \to \infty} \frac{\log n}{\log s_n} \ge \lim_{m \to \infty} \frac{\log \sum_{i=n_0}^{m} l_i}{\log(l_m q_m)}
$$

\n
$$
\ge \lim_{m \to \infty} \frac{\log l_m}{\log(l_m q_m)} = \lim_{m \to \infty} \frac{\log \lfloor \frac{q_{m+1}}{12q_m} \rfloor}{\log \lfloor \frac{q_{m+1}}{12q_m} \rfloor q_m)}
$$

\n
$$
= \lim_{m \to \infty} \frac{\log \frac{q_{m+1}}{q_m}}{\log q_{m+1}} = \frac{\frac{\tau}{1-\tau}}{\frac{\tau}{1-\tau} + 1} \text{ (by (4.2))}
$$

\n
$$
= \tau.
$$

 $\hfill\square$

The following lemma will be used while computing the entropy dimension.

Lemma 4.2. When n is large enough,

$$
\mathbf{m}\{\theta\in\mathbf{T}:\omega(\theta,iq_n)=i,1\leq i\leq \lfloor\frac{||q_{n-1}\alpha||}{6||q_n\alpha||}\rfloor\}>\frac{1}{8}.
$$

Proof. We only consider the case $n = 2t$ and the case $n = 2t + 1$ is similar. By Proposition [3.1,](#page-6-0) $\omega(q_n) = 1$. And we note that $\{q_n \alpha\} = ||q_n \alpha||$.

Order the points in Q_n on **T** by $0 < q_n \alpha < \cdots < s_1 \alpha < \frac{1}{2} < s_2 \alpha < \cdots < s_{n-1} \alpha < 1$. Then $||s_2\alpha - s_1\alpha|| \ge ||q_{n-1}\alpha||$. We note that from [\(4.2\)](#page-11-0), when *n* is sufficiently large, $||q_{n-1}\alpha|| \gg ||q_n\alpha||.$

Case 1. $||\frac{1}{2} - s_1 \alpha|| \ge \frac{1}{2} ||q_{n-1} \alpha||.$

Claim 1. For any $\ell \alpha \in Q_n$ and $0 \le \theta < ||\frac{1}{2} - s_1\alpha|| - ||q_n\alpha||$, $\omega(\ell \alpha + \theta, q_n) = \omega(q_n) = 1$.

Proof of the Claim 1. Notice that

 $\ell\alpha+\theta+Q_n=\big(\theta+\ell\alpha+\{\alpha,2\alpha,\cdots,(q_n-\ell)\alpha\}\big)\Big|\ \big|(\theta+||q_n\alpha||+\{\alpha,2\alpha,\cdots,\ell\alpha\}\big).$

Compare elements in $\ell \alpha + \theta + Q_n$ with elements in Q_n , we see that the first part of the right-hand side of the above equality is a translation by θ of a subset of Q_n and the second part is a translation by $\theta + ||q_n\alpha||$ of a subset of Q_n . Since the distance of $s_1\alpha$ and $\frac{1}{2}$ is larger than $\theta + ||q_n\alpha||$, we have that $\omega(\ell\alpha + \theta, q_n) = \omega(q_n) = 1$.

For any $\ell \alpha \in Q_n$, $0 \leq \theta < \frac{1}{4} ||q_{n-1} \alpha||$ and $0 \leq i \leq \lfloor \frac{||q_{n-1} \alpha||}{4||q_n \alpha||} \rfloor - 1$, it is easy to see that $\|\theta+i\|q_n\alpha\|+||q_n\alpha||<||\frac{1}{2}-s_1\alpha||.$ Hence for $1\leq i\leq \lfloor \frac{||q_{n-1}\alpha||}{4||q_n\alpha||}\rfloor-1$, by the Claim, $\omega(\ell\alpha + \theta, i a_n) = \omega(\ell\alpha + \theta, a_n) + \omega(\ell\alpha + \theta + ||a_n\alpha||, a_n)$

$$
\omega(\ell\alpha + \theta, i q_n) = \omega(\ell\alpha + \theta, q_n) + \omega(\ell\alpha + \theta + ||q_n\alpha||, q_n)
$$

$$
+ \cdots + \omega(\ell\alpha + \theta + (i - 1)||q_n\alpha||, q_n)
$$

$$
= i.
$$

By [\(2.5\)](#page-5-1),

$$
||q_{n-1}\alpha|| = q_{n-1}(-1)^{n-1}(\alpha - \frac{p_{n-1}}{q_{n-1}}) \ge \frac{1}{q_n + q_{n-1}} > \frac{1}{2q_n}.
$$

Let *n* be large enough to make $\lfloor \frac{||q_{n-1}\alpha||}{6||q_n\alpha||} \rfloor \leq \lfloor \frac{||q_{n-1}\alpha||}{4||q_n\alpha||} \rfloor - 1$. Since

$$
\{\theta\in\mathbf{T}: \omega(\theta, iq_n)=i, 1\leq i\leq \lfloor\frac{||q_{n-1}\alpha||}{6||q_n\alpha||}\rfloor\}\supset \bigsqcup_{\ell=1}^{q_n}\big(\ell\alpha+[0,\frac{1}{4}||q_{n-1}\alpha||)\big),
$$

we have that

$$
\mathbf{m}\{\theta \in \mathbf{T} : \omega(\theta, iq_n) = i, 1 \le i \le \lfloor \frac{||q_{n-1}\alpha||}{6||q_n\alpha||}\rfloor\} \ge q_n \frac{1}{4} ||q_{n-1}\alpha|| > \frac{1}{8}.
$$

Case 2. $\left| \frac{1}{2} - s_1 \alpha \right| \right| < \frac{1}{2}$ $\frac{1}{2}||q_{n-1}\alpha||.$

Claim 2. For any $\ell \alpha \in Q_n$ and $\left|\left|\frac{1}{2} - s_1 \alpha\right|\right| < \theta \leq ||q_{n-1} \alpha|| - ||q_n \alpha||$, $\omega(\ell \alpha + \theta, q_n) =$ −1.

Proof of the Claim 2. As in the proof of Claim 1,

$$
\ell\alpha+\theta+Q_n=\big(\theta+\ell\alpha+\{\alpha,2\alpha,\cdots,(q_n-\ell)\alpha\}\big)\bigsqcup\big(\theta+||q_n\alpha||+\{\alpha,2\alpha,\cdots,\ell\alpha\}\big).
$$

Then elements in $\ell \alpha + \theta + Q_n$ are obtained by translating elements in Q_n by θ or $\theta + ||q_n\alpha||$. We see that only one element $s_1\alpha$ is moved from upper semi-circle to lower semi-circle. Hence $\omega(\ell \alpha + \theta, q_n) = -1$.

Similar to Case 1, when n is large enough, we also have

$$
\mathbf{m}\{\theta \in \mathbf{T} : \omega(\theta, iq_n) = i, 1 \le i \le \lfloor \frac{||q_{n-1}\alpha||}{6||q_n\alpha||}\rfloor\} \ge q_n \frac{1}{4} ||q_{n-1}\alpha|| > \frac{1}{8}.
$$

$$
13\quad
$$

 \Box

Theorem 4.3. Let $\mathfrak{X} = \mathfrak{X}_{\alpha, \mathfrak{Y}}$ be the DWRS with rotation α along \mathfrak{Y} , where α is defined by (4.1) and $\mathfrak Y$ is a TDS with positive entropy. Then the topological entropy dimension of \mathfrak{X} is τ .

Proof. Suppose $\mathfrak{Y} = (Y, S)$ and $\mathfrak{X} = (Z \times Y, T)$ as defined in section 2. Let U be a finite open cover of Z and V be a finite open cover of Y. For $q_n \leq k \leq q_{n+1}$ and $i = 1, \dots, k$, we now partition Z due to the range of $\omega(z, i)$ for $z \in Z$. Note that $m(i) \leq \omega(z, i) \leq M(i)$. Let $\widetilde{U}_{i,\ell} = \{z \in Z : \omega(z, i) = \ell\}$, where $m(i) \leq \ell \leq M(i)$. Then $\mathcal{U}_i = \{\tilde{U}_{i,m(i)}, \tilde{U}_{i,m(i)+1}, \cdots, \tilde{U}_{i,M(i)}\}$ is an open cover of Z. Note that $z \in \sigma^{i+1} U_{i,\ell}$ if and only if $\omega(\sigma^{-i}z, 0, i-1) = \omega(\sigma^{-(i+1)}z, i) = \ell$. Hence

$$
\bigvee_{i=1}^{k} T^{-i}(\mathcal{U} \times \mathcal{V}) \prec \bigvee_{i=1}^{k} T^{-i}((\mathcal{U} \vee \sigma^{i+1}\widetilde{\mathcal{U}}_i) \times \mathcal{V})
$$

$$
\prec \big(\bigvee_{i=1}^{k} \sigma^{-i}(\mathcal{U} \vee \sigma^{i+1}\widetilde{\mathcal{U}}_i)\big) \times \big(\bigvee_{j=m(k)}^{M(k)} S^{-j}\mathcal{V}\big).
$$

So

$$
\log N\left(\bigvee_{i=1}^{k} T^{-i}(\mathcal{U} \times \mathcal{V})\right)
$$
\n
$$
\leq \log N\left(\left(\bigvee_{i=1}^{k} \sigma^{-i}(\mathcal{U} \vee \sigma^{i+1}\widetilde{\mathcal{U}}_{i})\right) \times \left(\bigvee_{j=m(k)}^{M(k)} S^{-j} \mathcal{V}\right)\right)
$$
\n
$$
\leq \log N\left(\bigvee_{i=1}^{k} \sigma^{-i} \mathcal{U}\right) + \log N\left(\bigvee_{i=1}^{k} \sigma(\widetilde{\mathcal{U}}_{i})\right) + \log N\left(\bigvee_{j=m(k)}^{M(k)} S^{-j} \mathcal{V}\right)
$$
\n
$$
\leq \log N\left(\bigvee_{i=1}^{k} \sigma^{-i} \mathcal{U}\right) + \log N\left(\bigvee_{i=1}^{k} \sigma(\widetilde{\mathcal{U}}_{i})\right) + \left(M(k) - m(k) + 1\right) \log N(\mathcal{V}).
$$

For the Sturmian system (Z, σ) , its sequence entropy and entropy dimension are both zero. Noticing that $\bigvee_{i=1}^k \sigma(\widetilde{\mathcal{U}}_i) \prec \{\text{the cover formed by the blocks of length } k+1\},\$ have that for any $\epsilon > 0$,

$$
\limsup_{k\to\infty}\frac{1}{k^{\tau+\epsilon}}\bigg(\log N\big(\bigvee_{i=1}^k\sigma^{-i}\mathcal{U}\big)+\log N\big(\bigvee_{i=1}^k\sigma(\widetilde{\mathcal{U}}_i)\big)\bigg)=0.
$$

For each k, there exists a unique $n(k)$ such that $q_{n(k)} < k \leq q_{n(k)+1}$. By Proposition [3.4,](#page-10-1)

$$
\limsup_{k \to \infty} \frac{1}{k^{\tau+\epsilon}} \left(M(k) - m(k) + 1 \right) \le \limsup_{k \to \infty} \frac{1}{k^{\tau+\epsilon}} \left(6\left\lceil \frac{k}{q_{n(k)}} \right\rceil + 6 \sum_{j=2}^{n} (k) \frac{q_j}{q_{j-1}} + 2q_1 + 5 \right)
$$

$$
\le \limsup_{k \to \infty} \frac{6 \frac{k}{q_{n(k)}}}{k^{\tau+\epsilon}} + \limsup_{k \to \infty} \frac{6n(k) \frac{q_{n(k)}}{q_{n(k)-1}}}{k^{\tau+\epsilon}}
$$

$$
\leq 6\left(\limsup_{k\to\infty}\frac{q_{n(k)+1}^{1-\tau}}{q_{n(k)}k^{\epsilon}}+\limsup_{k\to\infty}\frac{q_{n(k)}^{1-\tau}n(k)}{q_{n(k)-1}k^{\epsilon}}\right)=0.
$$

Hence by (1) of Remark [2.1,](#page-2-0) $\overline{D}(X,T) \leq \tau$.

If $\tau = 0$, $\overline{D}(X,T) = 0$ has already been proved. In the following, we will show $D(X,T) \geq \tau$ for $\tau > 0$.

Since $h_{top}(Y, S) > 0$, there exist two non-empty disjoint closed sets A_1, A_2 of Y, such that $h_{top}(S, \{A_1^c, A_2^c\}) > 0$ (c.f. [\[5\]](#page-17-20)). Let $\mathcal{U} = \{Z \times A_1^c, Z \times A_2^c\}$, which is an open cover of X .

By Lemma [4.2,](#page-11-1) when n is large enough, we can take any small closed interval from $\{\theta \in \mathbf{T} : \omega(\theta, iq_n) = i, 1 \leq i \leq \lfloor \frac{||q_{n-1}\alpha||}{6||q_n\alpha||} \rfloor\}$ and denote it by *I*. Together with [\(4.3\)](#page-11-2), for any fixed $1 \leq i \leq l_n$ and any point $z \in \varphi^{-1}(I)$, $\omega(z, iq_n) = i$. By (2.1) , for $j = 1, 2$,

$$
T^{-iq_n}(\varphi^{-1}((iq_n+1)\alpha+I)\times A_j)=\varphi^{-1}(\alpha+I)\times S^{-i}A_j.
$$

Hence for any $1 \leq \ell \leq l_n$,

$$
N\Big(\bigvee_{i=1}^{\ell} T^{-iq_n} \Big\{ Z \times A_1^c, Z \times A_2^c \Big\} \Big)
$$

\n
$$
\geq N\Big(\bigvee_{i=1}^{\ell} T^{-iq_n} \Big\{ \big(\varphi^{-1}((iq_n+1)\alpha+I) \times A_1\big)^c, \big(\varphi^{-1}((iq_n+1)\alpha+I) \times A_2\big)^c \Big\} \Big)
$$

\n
$$
= N\Big(\bigvee_{i=1}^{\ell} \Big\{ \big(\varphi^{-1}(\alpha+I) \times S^{-i}A_1\big)^c, \big(\varphi^{-1}(\alpha+I) \times S^{-i}A_2\big)^c \Big\} \Big)
$$

\n
$$
= N\Big(\bigvee_{i=1}^{\ell} \Big\{ \varphi^{-1}(\alpha+I)^c \times Y \cup \varphi^{-1}(\alpha+I) \times S^{-i}A_1^c,
$$

\n
$$
\varphi^{-1}(\alpha+I)^c \times Y \cup \varphi^{-1}(\alpha+I) \times S^{-i}A_2^c \Big\} \Big)
$$

\n
$$
= N\Big(\bigvee_{i=1}^{\ell} S^{-i} \{A_1^c, A_2^c\} \Big).
$$

Now let $F = \{s_1 < s_2 < \cdots\}$ as defined in [\(4.4\)](#page-11-3). For any k, there exists $n(k)$ such that $q_{n(k)} \leq s_k < q_{n(k)+1}$. Assume that $s_k = \ell q_{n(k)}$. Then $k = \sum_{i=n_0}^{n(k)-1} l_i + \ell$ and $\max\{\sum_{i=n_0}^{n(k)-1} l_i, \ell\} \ge \frac{1}{2}k.$

Hence along the sequence F ,

$$
\liminf_{k \to \infty} \frac{1}{k} N\left(\bigvee_{i=1}^{k} T^{-s_i} \{Z \times A_1^c, Z \times A_2^c\}\right)
$$
\n
$$
\geq \liminf_{k \to \infty} \frac{1}{k} \max \left\{ N\left(\bigvee_{i=1}^{l_{n(k)-1}} T^{-iq_{n(k)-1}} \{Z \times A_1^c, Z \times A_2^c\}\right),\right\}
$$

$$
N\Big(\bigvee_{i=1}^{\ell} T^{-iq_{n(k)}}\{Z \times A_1^c, Z \times A_2^c\}\Big)\Big\}
$$

\n
$$
\geq \liminf_{k \to \infty} \frac{1}{k} \max \Big\{ N\Big(\bigvee_{i=1}^{l_{n(k)-1}} S^{-i}\{A_1^c, A_2^c\}\Big), N\Big(\bigvee_{i=1}^{\ell} S^{-i}\{A_1^c, A_2^c\}\Big)\Big\}
$$

\n
$$
= \liminf_{k \to \infty} \max \Big\{ \frac{l_{n(k)-1}}{k} \frac{N\Big(\bigvee_{i=1}^{l_{n(k)-1}} S^{-i}\{A_1^c, A_2^c\}\Big)}{l_{n(k)-1}}, \frac{\ell}{k} \frac{N\Big(\bigvee_{i=1}^{\ell} S^{-i}\{A_1^c, A_2^c\}\Big)}{\ell} \Big\}
$$

\n
$$
\geq \frac{1}{4} h_{top}(S, \{A_1^c, A_2^c\}) > 0.
$$

So F is an entropy generating sequence of \tilde{U} . Noticing that $\overline{D}(F) = \tau$, we can deduce that $\overline{D}(X,T) \geq \tau$. Hence $\overline{D}(X,T) = \tau$.

5. Metric entropy dimension

Let $\mathfrak{X} = \mathfrak{X}_{\alpha, \mathfrak{Y}}$ be the DWRS with rotation α along \mathfrak{Y} , where $\alpha \in E_{\tau}$ is defined as in section 4 and $\mathfrak V$ is a MDS with positive metric entropy. In this section, we will show that the metric entropy dimension of $\mathfrak X$ also equals τ .

By Remark [2.2,](#page-4-0) in this section we assume $\mathfrak{X} = (X, \mathcal{B}, \mu, T) = (\mathbf{T} \times Y, \mathcal{B}_{\mathbf{T}} \otimes \mathcal{D}, \mathbf{m} \times \mathcal{B}_{\mathbf{T}})$ ν, T).

Let us consider a measurable partition of X, say $\mathcal{P} = \{T \times A, T \times A^c\}$, where $\{A, A^c\}$ is a measurable partition of Y with $h_{\nu}(S, \{A, A^c\}) > 0$.

Proposition 5.1. Let $F = \{s_1 < s_2 < \cdots\}$ be defined by [\(4.4\)](#page-11-3). Then F is an entropy generating sequence of P . Hence by Lemma [4.1,](#page-11-4)

$$
\overline{D}_{\mu}(T,\mathcal{P}) \geq \tau.
$$

Proof. For $A = \{A_1, A_2, \cdots, A_k\}$, a collection of measurable subsets of X (need not to be a partition), we still denote $H_{\mu}(\mathcal{A}) = \sum_{i=1}^{k} -\mu(A_i) \log \mu(A_i)$.

Let $W = A$ or A^c and denote by $I_i = \{ \theta \in \mathbf{T} : \omega(\theta, iq_n) = i \}$ and $I = \bigcap_{i=1}^{\ell} I_i$, then by (2.1) ,

$$
T^{-iq_n}(\mathbf{T} \times W) = \bigcup_{\theta \in \mathbf{T}} (\theta - iq_n \alpha, S^{-\omega(\theta - iq_n \alpha, 0, iq_n - 1)} W)
$$

$$
\supset \{ \theta \in \mathbf{T} : \omega(\theta, 0, iq_n - 1) = i \} \times S^{-i} W
$$

$$
= (I_i + \alpha) \times S^{-i} W.
$$

So

$$
H_{\mu}(\bigvee_{i=1}^{\ell} T^{-iq_n} \mathcal{P}) \ge H_{\mu}\left(\bigvee_{i=1}^{\ell} (I_i + \alpha) \times S^{-i}\{A, A^c\}\right)
$$

$$
= H_{\mu}\left(\bigcap_{i=1}^{\ell} (I_i + \alpha) \times \bigvee_{i=1}^{\ell} S^{-i}\{A, A^c\}\right)
$$

$$
= -\mathbf{m}(I) \log \mathbf{m}(I) + \mathbf{m}(I) H_{\nu}(\bigvee_{i=1}^{\ell} S^{-i} \{A, A^{c}\})
$$

$$
\geq \frac{1}{8} H_{\nu}(\bigvee_{i=1}^{\ell} S^{-i} \{A, A^{c}\}).
$$

For any k, there exists $n(k)$ such that $q_{n(k)} \leq s_k < q_{n(k)+1}$. Assume that $s_k = \ell q_{n(k)}$. Then $k = \sum_{i=1}^{n(k)-1} l_i + \ell$ and $\max\{\sum_{i=n_0}^{n(k)-1} l_i, \ell\} \ge \frac{1}{2}k$.

Hence along the sequence F ,

$$
\liminf_{k \to \infty} \frac{1}{k} H_{\mu} \Big(\bigvee_{i=1}^{k} T^{-s_{i}} \mathcal{P} \Big)
$$
\n
$$
\geq \liminf_{k \to \infty} \frac{1}{k} \max \Big\{ H_{\mu} \Big(\bigvee_{i=1}^{l_{n(k)-1}} T^{-iq_{n(k)-1}} \mathcal{P} \Big), H_{\mu} \Big(\bigvee_{i=1}^{\ell} T^{-iq_{n(k)}} \mathcal{P} \Big) \Big\}
$$
\n
$$
\geq \liminf_{k \to \infty} \frac{1}{k} \max \Big\{ \frac{1}{8} H_{\nu} \Big(\bigvee_{i=1}^{l_{n(k)-1}} S^{-i} \{A, A^{c}\}, \frac{1}{8} H_{\nu} \Big(\bigvee_{i=1}^{\ell} S^{-i} \{A, A^{c}\} \Big) \Big\}
$$
\n
$$
= \liminf_{k \to \infty} \max \Big\{ \frac{l_{n(k)-1}}{8k} \frac{H_{\nu} \Big(\bigvee_{i=1}^{l_{n(k)-1}} S^{-i} \{A, A^{c}\}, \frac{\ell}{8k} \frac{H_{\nu} \Big(\bigvee_{i=1}^{\ell} S^{-i} \{A, A^{c}\} \Big)}{\ell} \Big\}
$$
\n
$$
\geq \frac{1}{32} h_{\nu} \big(S, \{A, A^{c}\} \big) > 0.
$$

So F is an entropy generating sequence of \mathcal{P} .

Since the metric entropy dimension is bounded from above by the topological entropy dimension, together with Theorem [4.3](#page-13-0) and Proposition [5.1,](#page-15-0) we have

Theorem 5.2. The metric entropy dimension of $\mathfrak X$ equals τ .

- **Remark 5.3.** (1) Take an increasing sequence $\{\tau_n\}$ such that $\lim_{n\to\infty}\tau_n = 1$. In the construction of the irrational number $\alpha = \alpha(t_1, t_2, \dots, t_k) = [0; a_1, a_2, \dots]$ (see [\(4.1\)](#page-10-0)), when $n > k + 2$, modify a_{n+1} by $a_{n+1} = 2\lfloor q_n^{\frac{1}{1-r_n}} \rfloor$. By similar proofs of Lemma [4.1](#page-11-4) and Proposition [5.1,](#page-15-0) we can show that the metric entropy dimension of $\mathfrak X$ equals 1 for such α 's. Hence the topological entropy dimension is also 1.
	- (2) For $\tau \in [0, 1)$, since E_{τ} is dense in $[0, 1)$, by Theorem [4.3](#page-13-0) and Theorem [5.2,](#page-16-0) the collection of irrational α 's such that the system $\mathfrak X$ has entropy dimension τ is dense.

Acknowledgements We are grateful to the referee for the careful reading and many valuable comments to improve the paper. The first author is supported by NNSF of China (Grant No. 11790274, 11401220 and 11431012). The second author is supported in part by NRF 2010-0020946.

18 D. Dou and K.K. Park

REFERENCES

- [1] J. Aaronson, Relative complexity of random walks in random sceneries, Ann. Probab., 40 (2012), no. 6, 2460-2482.
- [2] J. Aaronson and M. Keane, The visits to zero of some deterministic random walks, Proceedings of the London Mathematical Society, 44 (1982), 535-553.
- [3] Y. Ahn, D. Dou and K. K. Park, Entropy dimension and its variational principle, Studia Math., 199 (2010), no. 3 295-309.
- [4] A. Avila, D. Dolgopyat, E. Duryev, O. Sarig, The visits to zero of a random walk driven by an irrational rotation, Israel J. Math., 207 (2015), 653-717.
- [5] F. Blanchard, A disjointness theorem involving topological entropy, Bull. Soc. Math. France, 121(1993), 465-478.
- [6] Julien Cassaigne, Constructing infinite words of intermediate complexity, Developments in language theory, 173–184, Lecture Notes in Comput. Sci., 2450, Springer, Berlin, 2003.
- [7] M. De Carvalho, Entropy dimension of dynamical systems, Portugal. Math., 54 (1997), no. 1, 19-40.
- [8] J.-P. Conze and M. Keane, Ergodicité d'un flot cylindrique, in Séminaire de Probabilités, I (Univ. Rennes, Rennes, 1976), Dépt. Math. Informat., Université de Rennes, Rennes, 1976, Exp. No. 5.
- [9] D. Dou, W. Huang and K. K. Park, Entropy dimension of topological dynamics, Trans. Amer. Math. Soc., **363** (2011), 659-680.
- [10] D. Dou, W. Huang and K. K. Park, Entropy dimension of measure preserving systems, Trans. Amer. Math. Soc., **371** (2019), 7029-7065.
- [11] D. Dou and K. K. Park, Examples of entropy generating sequence, Sci. China Math., **54** (2011), no. 3, 531-538.
- [12] S. Ferenczi and K. K. Park, Entropy dimensions and a class of constructive examples, Discrete Cont. Dyn. Syst., 17 (2007), no. 1, 133-141.
- [13] M. Hochman, Slow entropy and differentiable models for infinite-measure preserving \mathbb{Z}^k actions, Ergod. Th. Dynam. Sys., 32 (2012), no. 2, 653-674.
- [14] U. Jung, J. Lee and K. K. Park, Constructions of subshifts with positive topological entropy dimension, [arXiv:1601.07259v](http://arxiv.org/abs/1601.07259)1.
- [15] U. Jung, J. Lee and K. K. Park, Topological entropy dimension and directional entropy dimension for \mathbb{Z}^2 -subshifts, *Entropy*, **19** (2017), no. 2, Paper No. 46, 13 pp.
- [16] A. Katok and J.-P. Thouvenot, Slow entropy type invariants and smooth realization of commuting measure-preserving transformations, Ann. Inst. H. Poincare Probab. Statist., 33 (1997), no. 3, 323-338.
- [17] A. Ya. Khinchin, Continued fractions, Groningen: Noordhoff, 1963.
- [18] H. Kim and S. Park, Toeplitz sequences of intermediate complexity, Journal of The Korean Mathematical Society, 48 (2011), no. 2, 383-395.
- [19] K. K. Park, On directional entropy functions, Israel J. Math., 113 (1999), 243-267.
- [20] K. K. Park, A counterexample of the entropy of a skew product, Indagationes Mathematicae, 9 (1998), no. 4, 537-548.
- [21] J. de Vries, Elements of topological dynamics, Kluwer, Dordrecht, 1993.

Department of Mathematics, Nanjing University, Nanjing, Jiangsu, 210093, P.R. China

Email address: doumath@163.com

Center for Mathematical Challenges, Korea Institute for Advanced Study, Seoul 130-722, Korea

Email address: kkpark@kias.re.kr