

## Effect of Acute and Repeated Exposure to Traumatic Conditions on Gene Expression

Akram Yazdani<sup>1\*</sup>, Fatemeh Haghighi<sup>1, 2,\*</sup>

<sup>1</sup>Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, USA

<sup>2</sup>James J. Peters VA Medical Center, New York, NY, USA

\*To whom correspondence should be addressed

### Abstract

Repeated exposure to low-level blast may initiate a range of adverse health problem such as traumatic brain injury, yet mechanisms underpinning its actions are not fully understood. In this study, we investigated underlying relationship among genes through constructing transcript network using RNA-seq data. Data for pre- and post-transcripts collected from 32 individuals in Army training sites combined with our system approach provided unique opportunity to compare the pre- and post-transcript networks. Digging into the networks, we identified four pathways that are dysregulated and disrupted due to the exposure waves produced by blast. The pathways that are related to immune system and inflammatory process are complementary to experimental studies while they hypothesize new target for prevention and intervention. Among genes with relatively high Fold-Change in their transcript expression levels, genes *PELI*, *PTGS2*, *TRIM58*, *B2M*, *BCL2AL*, *S100A8* and *ZNF654* showed major impact on dysregulation of the pathways. In addition, this study reveals how repeated exposures to traumatic conditions increase the transcripts' Fold-Changes.

### Introduction

There is a major concern that repeated exposure to low-level blast may lead to traumatic brain injury (mTBI) or subclinical exposure (Elder, Stone, & Ahlers, 2014). Although no apparent physical damage presented virtually due to low-level blast exposure, it activates multiple apoptotic and inflammatory pathways through acute effects on the brain (Carr et al., 2015). Furthermore, many studies provide evidence that the effect of exposure to blast is cumulative and long-lasting (Elder et al., 2014). All these make necessary to understand the latent pathological processes of chronic exposure and its association with mTBI.

While mTBI is a complex disease and multiple pathways are involved in the pathology and mechanism of the disease, complementary studies to experimental studies are required. Overwhelming studies based on microarrays (Redell et al., 2013; Rojo et al., 2011; Samal et al., 2015) and more recently RNA sequencing (Lipponen, Paananen, Puhakka, & Pitkänen, 2016) focus on differential gene expression in response to TBI and successfully identified biomarkers associated to TBI. However, those studies do not provide information regarding changes in the system while analyzing each gene separately regardless of their relationships. To address this limitation, approaches like network analysis are applied.

In this study, we conducted a systems approach using Bayesian network to predict essential disruption in the gene-gene interactions due to pressure waves produced by blast. We constructed the network on RNA-seq data collected from 32 individuals participating in a 2-week cycle at U.S. army explosive entry training sites. The blood samples of the individuals were collected pre- and post-training to profile transcript abundance which makes possible comparison between pre- and post-identified pathways. This systems approach combined with specific data set available in our study provided a unique opportunity to study the mechanism of disease through identifying pathways altered by blast exposure.

## Materials and methods

**Samples:** This study includes 32 samples from site 3 of a study designed by Naval Medical Research Center and Walter Reed Army Institute of Research Institutional Review Boards (Reference). In this training program, all participants are male while 19 participants reported with history of TBI and 24 reported with prior experience in explosive breaching. Figure 1, right plot, represents the overlap between subsets of individuals with history of TBI and history of breacher through a Venn diagram. All individuals who were enrolled as students or designated as instructors for the training programs were eligible to participate in the study. The average ages of participants were 31 with standard deviation 4.6, Figure 1, left plot. During the 2-week period, trainees were exposed to multiple blasts and their blood draw was taken at baseline and the end of training program.




Figure 1: Left: Histogram of age. Right: Venn diagram for history of TBI and breacher (BR).

**RNA-Seq:** Total RNA sequencing libraries were prepared using the Illumina Stranded for the first day (before exposing to blast) and last day (after exposing to blast during 2-week training program). Final libraries were evaluated using PicoGreen (Life Technologies) and Fragment Analyzer (Advanced Analytics) and were sequenced on an Illumina HiSeq2500 sequencer (v4 chemistry) using 2 x 125bp read length. RNA-seq reads were aligned to the human reference hg19 using STAR aligner (v2.4.0c) (Dobin et

al., 2013). Quantification of genes annotated in Gencode v18 was performed using feature Counts (v1.4.3). The QC metrics were collected with Picard (v1.83) and RSeQC (Wang, Wang, & Li, 2012).

**Statistical Method:** Having pre- and post-transcripts for the same individuals, we constructed the gene expression Bayesian network (Friedman, Geiger, & Goldszmit, 1997) and identified changes in gene-expression pathways that are due to the exposing to blast. In this analysis, we calculated matrix counts of transcriptome reads and then normalize for gene length and sequencing depth. To have high sequencing coverage, we selected 2578 genes from the pre- and post-data with at least 10 nonzero recorded transcripts over samples, provided in Supplementary 1. We constructed the Bayesian networks separately for the pre- and post-data with tuning parameter 0.001. RNA-seq Bayesian network is a probabilistic network and seeks to determine the dependency between genes in the system. In the RNA-seq networks, each node stands for a gene and the edge between nodes represents interaction between corresponding nodes. To compare the identified networks, we focused on subnetworks with at least 4 genes in the pre-network as identified pathways in observational study. We then selected paths that show one or two changes in their gene interactions in post-network. After that, we mapped TBI associated genes from previous studies on the constructed networks. This integration categorized the results into two classes: one class comprises pathways with TBI associated genes and the other one comprises pathways with no prior knowledge associated with TBI. Figure 2 shows the strategy for selecting the target pathways in this study.




Figure 2: Strategy of selecting target subnetwork or pathways.

## Results

To determine whether the effects of exposure waves produced by blast changed the number of genes in each pathway, we compared the size of pathways in pre- and post-transcriptomic analyses, Figure 4. The size of pathway is equivalent with the number of genes in each pathway. We observed that the largest changes occurred for the pathways of size 3, 4 and 5 which are decreased 18%, 27%, 45% respectively after exposing to blast.




Figure 3: Following blast induces perturbation is measured as the network connectivity and the size of subnetworks. Left: Distribution of the size of subnetworks in pre-network analysis. Right: Distribution of the size of subnetworks in post-network analysis.

In both networks, each gene interacted with at most 3 other genes directly. In overall, the direct interaction level of each gene slightly reduced in the post-network, Table 1.

Table 1: Degree of connectivity in the pre- and post- data analyses

Level of direct interaction	1	2	3
Pre_transcriptomic analysis	1498	303	6
Post_transcriptomic analysis	1432	242	4

To extract more instructive information, we dug the network and found 4 pathways based on the strategy in Figure 2. Calculating the Fold-Change of transcript levels in each path elucidates which gene in the path has major impact on dysregulation of the pathway. Fold-Change is computed as the average of ratio of pre vs post transcriptomic levels of each sample. Fold-Change value greater than 1 indicates upregulation and the value less than one indicates downregulation in posteriori. Genes with relatively large up/down regulation in identified pathways are visualized in Figure 4 since Supplementary 2 includes the Fold-Change of all genes in identified pathways.




Figure 4: Log Fold-Change of genes with highest impact in identified pathways. Left: over individuals with TBI history. Right: over individuals without TBI history. Vertical lines represent the threshold for significant changes.

In addition, we separated the samples into two subsets with and without TBI history and calculated the Fold-change for each subset. Interestingly, upregulated genes due to exposing to blast showed greater changes among individuals without TBI history. In contrast, downregulated genes showed greater changes among individuals with TBI history. This may be due to the differences between acute effect of blast exposure and accumulative effect of repeated blast exposure. We did not present similar comparison for subsets of individuals with and without breacher history since the size of subsamples are very different (24 to 8).

### Properties and functions of identified pathways

**Pathway 1:** This path includes 4 genes (*PELI1*, *ST8SIA4*, *ITM2B*, and *PTGS2*) in the pre-transcriptomic study. Among these genes (*PELI1*, *ITM2B* and *PTGS2*) are TBI associated genes (Huang et al., 2017; Wu et al., 2013; Graber, Costine, & Hickey, 2015). However, as Figure 5 displays, there are no interactions between (*PTGS2*, *ITM2B*) and (*ST8SIA4*, *ITM2B*) in the post-analysis. The transcript Fold-Change revealed that the effect of blast on individuals without history of TBI is larger. Figure 6 shows the violin plot of transcript genes with relatively higher Fold-Change.




Figure 5- Left: Identified pathway; black/green links represent gene interactions prior/posterior to the training and red cross represents missing link in posteriori. Right: Log Fold-Change of transcripts for all individuals, individuals with and without history of TBI.

While overexpression of *PELI1* can alter expression of *ST8SIA4* through their interaction, we conclude that induced transcript level of *PELI1* and *PTGS2* may play a key role in disruption of this pathway. These two genes previously identified as TBI associated genes. Ingenuity pathway analysis (IPA) indicates that all genes in pathway 1 have common function related to organismal injury and abnormalities. More discussion on the function and potential role in neurological disease of this pathway is presented in the discussion section.




Figure 6: Violin plots of transcripts of individuals without TBI history for the genes with relatively higher Fold-Change.

**Pathway 2:** Genes involve in this pathway are *BCL2L1*, *DCAF12*, *PSMF1* and *TRIM58* in the pre-transcriptomic analysis. However, in the post-analysis *PSMF1* did not show any interaction with other genes in the pathway, Figure 7. The analysis revealed that TBI associated gene *TRIM58* has highest Fold-Change and directly interacts with *DCAF12* while it shows a slight changes in its expression level. Furthermore, overexpression of these genes influenced the relation between *PSMF1* and *DCAF12*.




Figure 7- Left: Identified pathway; black/green links represent genes interaction prior/posterior to training and red cross represents missing link in posteriori. Right: Log Fold-Change of transcripts for all individuals with and without history of TBI.

We presented the violin plots of the *TRIM58* with relatively larger Fold-Change in the pathway in Figure 8 to show the probability density of the data at different transcript values. All genes in this pathway have common function related to organismal injury and abnormalities using IPA.




Figure 8: Violin plots of *TRIM58* transcripts of individuals without TBI history.

**Pathway 3:** This pathway comprises 5 genes (*B2M*, *ATP6V1G1*, *BCL2AL*, *LY96*, and *S100A8*) in the pre-analysis. In the post-analysis, after exposing the participants to blast, this path breaks into two shorter pathways (*B2M*, *ATP6V1G1*) and (*BCL2AL*, *LY96*, *S100A8*). As shown in Figure 9, gene *ATP6V1G1* and *BCL2AL* stop interacting due to exposure effect. In addition, *S100A8* interacts indirectly with *LY96* despite its direct interaction in pre-analysis.




Figure 9: Left: Identified pathway with 5 genes; black/green links represent gene interactions prior/posterior to training and red crosses represent missing links in the posteriori. Right: Log Fold-Change of transcripts for all individuals with and without history of TBI.

Any alteration observed in this pathway after exposing to waves produced by blast leads to downregulation of the genes. Right panel, Figure 8, shows the effect of exposure on individuals with history of TBI is higher than individuals without TBI history. In addition, Figure 10 displays probability densities of the genes with highest change through violin plots. This result may be an evidence of accumulated effect of repeated exposures to blast or traumatic conditions in biological reaction.




Figure 10: Violin plots of transcripts of individuals with TBI history for the genes with relatively higher Fold-Change.

All genes in this pathway are related to immune system; in particular, *B2M* and *ATP6V1G1* are related to innate immune system discussed in the discussion section. Similar to the aforementioned pathways, IPA indicates that all genes in this pathway have common function related to organismal injury and abnormalities.

**Pathway 4:** This path includes 5 genes (*PRDX3*, *RNF139*, *SDHD*, *SERP1*, and *TMED2*) in the pre-transcriptomic analysis. The post-transcriptomic analysis shows that *RNF139* interacts with *ZNF654*



while its interaction with *TMED2* and *SERP1* are disrupted in the pathway. The heatmap in Figure 11 displays upregulation of *ZNF654* in the group of individuals with no history of TBI while small Fold-Change can be seen for *RNF139* and *TMED2*. The violin plots in Figure 12 represent further details on changes in transcript values.




Figure 11: Left: The identified pathway with 6 genes; black/green links represent gene interactions prior/posterior to training and red crosses represent missing links in posteriori. Right: Log Fold-Change of transcripts for all the individuals with and without history of TBI.

Interestingly, IPA analysis does show that all genes in the pathway identified prior to blast exposure have common function related to organismal injury and abnormalities.




Figure 12: Violin plots of transcripts of individuals without TBI history for the genes with relatively higher Fold-Change.

## Discussion

In this study, we investigated underlying relationships among transcripts on RNA-seq data collected prior to and following the training program. Comparing the identified pathways in each set of data provided insights into mechanisms of blast exposure impact on gene interactions and potentially mTBI progress. This study generated blood-based biomarkers which are desirable with advantages of being cost- and time-effective while simultaneously being feasible at the population level compared with the collection of cerebrospinal fluid (CSF) or neuroimaging, (O'Bryant et al., 2016); (O'Bryant et al., 2017); (Lista, Faltraco, Prvulovic, & Hampel, 2013). Moreover, this study provided systematic knowledge and narrows down the search space for further studies regarding treatment and prevention. Therefore, these kinds of systems approaches are complementary to experimental studies especially in the age of big data.

In this research, we observed that the effect of traumatic conditions on people varies depending on their history of TBI. Although, limited sample prevents us to have group specific analysis for individuals with and without TBI history. That will help to find specific pathways for each group separately, and consequently have better prevention approaches. In addition, increasing the number of samples, leads to identify modules and longer pathways through network analysis.

Here, we discuss two of the identified pathways with more details to show the significant of these findings. One of these pathways includes 4 genes (*PELII*, *ST8SIA4*, *ITM2B*, and *PTGS2*) that are overexpressed among individuals without TBI history. This path that is dysregulated immediately after two weeks trainees can be a good target for preventing acute health effects of exposure to blast. In addition, all these genes are expressed in brain tissue and influence brain function through beta amyloid.

Gene *PELII* with the highest Fold-Change in the pathway is known to play a pivotal role in inflammatory and autoimmune processes (Choe et al., 2016). Park J. et al. (2017) observed *PELII* overexpression have several distinct phenotypes depending on the expression level of *PELII*. Interestingly, our analysis is consistent with their experimental observation since our study showed higher expression of *PELII* on individuals without TBI history while *ST8SIA4* expression altered as its direct interacted gene

*ST8SIA4* catalyzes the polycondensation of alpha-2,8-linked sialic acid required for the synthesis of polysialic acid. Molecular, cellular, and genetic studies implicate polysialic acid in the control of cell-cell and cell-matrix interactions, intermolecular interactions at cell surfaces, and interactions with other molecules in the cellular environment. Therefore, dysregulation in relation between *ST8SIA4* and *ITM2B* may be the cause of observing *ITM2B* immunoreactivity in extracellular deposits while glycosylation plays an important role in protein trafficking of *ITM2B*. The *ITM2B* deposition starts in early stages of the Alzheimer's disease (AD) pathology (Eliss et al., 1989) which support our result since our research showed this path is overexpressed among individuals without TBI history.

The path between *ITM2B* and *PTGS2* is also disrupted while we observed overexpression of *PTGS2*. The gene *PTGS2* (*Cox-2*) is responsible for the prostaglandin biosynthesis involved in inflammation (<https://www.genecards.org/cgi-bin/carddisp.pl?gene=PTGS2>). Furthermore, *PTGS2* is the key enzyme that controls the production of PGE which is also involved in various inflammatory process and degenerative disorder. Overexpression of *PTGS2* elevates PGE2 levels. As suggested by Ellis et al. (1989), increased levels of PGE2 after injury may be a body wide response including brain. PGE2 in brain does dependently inhibits macrophage-mediated phagocytosis of aggregated A $\beta$  (Fox et al., 2015;

Aronoff, Canetti, & Peters-Golden, 2004). Several lines of evidence indicate that PGE2 inhibits the microglial-mediated phagocytosis of aggregated A $\beta$  through the EP2 receptor (Fox et al., 2015).

The other Pathway with 5 genes (*B2M*, *ATP6V1G1*, *BCL2AL*, *LY96*, and *S100A8*) is related to the immune system. Except *LY96*, all other genes are low expressed among individuals with history of TBI. Thus, we can hypothesize that this path represents the accumulated effect of repeated exposures to blast or traumatic conditions on immune system.

Genes *B2M* and *ATP6V1G1*, which interact prior to and following the training program, are related to the innate immune system. Exposing to the blast wave resulted in a low expression level of both genes such that low expression of *B2M* causes low level of *MHC* class I antigens which influence a proper response of immune system (Safa, Saadatzadeh, Gadol, Pollok, & Vishehsaraei, 2016). Low expression of *ATP6V1G1* induces cell death since *ATP6V1G1* encodes a component of vacuolar ATPase, a multi-subunit enzyme necessary for intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation (Safa et al., 2016).

Three other genes in the Pathway (*BCL2AL*, *LY96*, *S100A8*) continued interacting after the training program although *S100A8* interacts with *LY96* indirectly through *BCL2AL* in posteriori. *S100A8* and *BCL2AL* with relatively large downregulation after exposing to blast are involved in the immune system process. Expression of *BCL2AL* with important role in immune system protects cells from various death stimuli. Down regulation of *BCL2A1* causes loss of survival of pre-T cell (Mandal et al., 2005). *S100A8* also stimulates innate immune cells and acts as an alarm in or a danger associated molecular pattern (DAMP) molecule. It is involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. Interestingly, protein of *S100A8* interacts with *LY96* (protein *MD-2*) with a role in the innate immune system (Deguchi et al., 2016) as we observed in identified pathway in this study. This interaction breaks after the training program as the effect of exposing to the blast waves. Therefore, this study suggests that preventing dysregulation of identified pathway may reduce the effect of exposing to traumatic conditions on immune system.

## References

- Aronoff, D. M., Canetti, C., & Peters-Golden, M. (2004). Prostaglandin E2 Inhibits Alveolar Macrophage Phagocytosis through an E-Prostanoid 2 Receptor-Mediated Increase in Intracellular Cyclic AMP. *The Journal of Immunology*, *173*(1), 559–565. <https://doi.org/10.4049/jimmunol.173.1.559>
- Carr, W., Yarnell, A. M., Ong, R., Walilko, T., Kamimori, G. H., Da Silva, U., ... LoPresti, M. L. (2015). Ubiquitin carboxy-terminal hydrolase-L1 as a serum neurotrauma biomarker for exposure to occupational low-level blast. *Frontiers in Neurology*, *6*(MAR), 1–11. <https://doi.org/10.3389/fneur.2015.00049>
- Choe, J.-Y., Park, M., Yun, J. Y., Na, H. Y., Go, H., Kim, H.-J., ... Kim, J. E. (2016). PELI1 expression is correlated with MYC and BCL6 expression and associated with poor prognosis in diffuse large B-cell lymphoma. *Modern Pathology*, *29*(11), 1313–1323. <https://doi.org/10.1038/modpathol.2016.128>
- Deguchi, A., Tomita, T., Ohto, U., Takemura, K., Kitao, A., Akashi-Takamura, S., ... Maru, Y. (2016). Eritoran inhibits S100A8-mediated TLR4/MD-2 activation and tumor growth by changing the

- immune microenvironment. *Oncogene*, 35(11), 1445–1456. <https://doi.org/10.1038/onc.2015.211>
- Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., ... Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, 29(1), 15–21. <https://doi.org/10.1093/bioinformatics/bts635>
- Elder, G. A., Stone, J. R., & Ahlers, S. T. (2014). Effects of low-level blast exposure on the nervous system: Is there really a controversy? *Frontiers in Neurology*, 5(DEC). <https://doi.org/10.3389/fneur.2014.00269>
- Ellis, E. F., Police, R. J., Rice, L. Y., Grabeel, M., & Holt, S. (1989). Increased Plasma PGE<sub>2</sub>, 6-Keto-PGF<sub>1α</sub>, and 12-HETE Levels Following Experimental Concussive Brain Injury. *Journal of Neurotrauma*, 6(1), 31–37. <https://doi.org/10.1089/neu.1989.6.31>
- Fox, B. M., Beck, H. P., Roveto, P. M., Kayser, F., Cheng, Q., Dou, H., ... Olson, S. H. (2015). A Selective Prostaglandin E<sub>2</sub> Receptor Subtype 2 (EP2) Antagonist Increases the Macrophage-Mediated Clearance of Amyloid-Beta Plaques. *Journal of Medicinal Chemistry*, 58(13), 5256–5273. <https://doi.org/10.1021/acs.jmedchem.5b00567>
- Friedman, N., Geiger, D., & Goldszmit, M. (1997). Bayesian Network Classifiers. *Machine Learning*, 29, 131–163. <https://doi.org/10.1023/a:1007465528199>
- Graber, D. J., Costine, B. A., & Hickey, W. F. (2015). Early inflammatory mediator gene expression in two models of traumatic brain injury: Ex vivo cortical slice in mice and in vivo cortical impact in piglets. *Journal of Neuroinflammation*, 12(1), 1–8. <https://doi.org/10.1186/s12974-015-0298-4>
- Huang, X.-P., Peng, J.-H., Pang, J.-W., Tian, X.-C., Li, X.-S., Wu, Y., ... Sun, X.-C. (2017). Peli1 Contributions in Microglial Activation, Neuroinflammatory Responses and Neurological Deficits Following Experimental Subarachnoid Hemorrhage. *Frontiers in Molecular Neuroscience*, 10(November), 1–13. <https://doi.org/10.3389/fnmol.2017.00398>
- Lipponen, A., Paananen, J., Puhakka, N., & Pitkänen, A. (2016). Analysis of Post-Traumatic Brain Injury Gene Expression Signature Reveals Tubulins, Nfe2l2, Nfkb, Cd44 and S100a4 as Treatment Targets. *Scientific Reports*, 6(1), 31570. <https://doi.org/10.1038/srep31570>
- Lista, S., Faltraco, F., Prvulovic, D., & Hampel, H. (2013). Blood and plasma-based proteomic biomarker research in Alzheimer's disease. *Progress in Neurobiology*, 101–102, 1–17. <https://doi.org/10.1016/j.pneurobio.2012.06.007>
- Mandal, M., Borowski, C., Palomero, T., Ferrando, A. A., Oberdoerffer, P., Meng, F., ... Aifantis, I. (2005). The BCL2A1 gene as a pre-T cell receptor-induced regulator of thymocyte survival. *The Journal of Experimental Medicine*, 201(4), 603–614. <https://doi.org/10.1084/jem.20041924>
- O'Bryant, S. E., Edwards, M., Johnson, L., Hall, J., Villarreal, A. E., Britton, G. B., ... Graff-Radford, N. R. (2016). A blood screening test for Alzheimer's disease. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*, 3, 83–90. <https://doi.org/10.1016/j.dadm.2016.06.004>
- O'Bryant, S. E., Mielke, M. M., Rissman, R. A., Lista, S., Vanderstichele, H., Zetterberg, H., ... Biofluid Based Biomarker Professional Interest Area, the B. B. B. P. I. (2017). Blood-based biomarkers in Alzheimer disease: Current state of the science and a novel collaborative paradigm for advancing from discovery to clinic. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association*, 13(1), 45–58. <https://doi.org/10.1016/j.jalz.2016.09.014>
- Park, J., Park, H.-Y., Kim, S., Kim, H.-S., Park, J. Y., Go, H., & Lee, C.-W. (2017). Pellino 1 inactivates

- mitotic spindle checkpoint by targeting BubR1 for ubiquitination degradation. *Oncotarget*, 8(19), 32055–32067. <https://doi.org/10.18632/oncotarget.16762>
- Redell, J. B., Moore, A. N., Grill, R. J., Johnson, D., Zhao, J., Liu, Y., & Dash, P. K. (2013). Analysis of Functional Pathways Altered after Mild Traumatic Brain Injury. *Journal of Neurotrauma*, 30(9), 752–764. <https://doi.org/10.1089/neu.2012.2437>
- Rojo, D. R., Prough, D. S., Falduto, M. T., Boone, D. R., Micci, M.-A., Kahrig, K. M., ... Hellmich, H. L. (2011). Influence of Stochastic Gene Expression on the Cell Survival Rheostat after Traumatic Brain Injury. *PLoS ONE*, 6(8), e23111. <https://doi.org/10.1371/journal.pone.0023111>
- Safa, A. R., Saadatzaheh, M. R., Gadol, A. A. C., Pollok, K. E., & Vishehsaraei, K. B. (2016). Emerging targets for glioblastoma stem cell therapy. *Journal of Biomedical Research*, 30(1), 19–31. <https://doi.org/10.7555/JBR.30.20150100>
- Samal, B. B., Waites, C. K., Almeida-Suhett, C., Li, Z., Marini, A. M., Samal, N. R., ... Eiden, L. E. (2015). Acute Response of the Hippocampal Transcriptome Following Mild Traumatic Brain Injury After Controlled Cortical Impact in the Rat. *Journal of Molecular Neuroscience*, 57(2), 282–303. <https://doi.org/10.1007/s12031-015-0626-2>
- Wang, L., Wang, S., & Li, W. (2012). RSeQC: quality control of RNA-seq experiments. *Bioinformatics (Oxford, England)*, 28(16), 2184–2185. <https://doi.org/10.1093/bioinformatics/bts356>
- Wu, P., Zhao, Y., Haidacher, S. J., Wang, E., Parsley, M. O., Gao, J., ... Denner, L. (2013). Detection of Structural and Metabolic Changes in Traumatically Injured Hippocampus by Quantitative Differential Proteomics. *Journal of Neurotrauma*, 30(9), 775–788. <https://doi.org/10.1089/neu.2012.2391>

## Supplementary 1

Table 1: Fold-Change for individuals with/out TBI history and the total in Pathway 1

	<i>ITM2B</i>	<i>PEL1</i>	<i>ST8SIA4</i>	<i>PTGS2</i>
<b>With TBI history</b>	1.163	1.393	1.233	1.221
<b>Without TBI history</b>	1.337	1.629	1.547	1.472
<b>Total</b>	<b>1.055</b>	<b>1.251</b>	<b>1.055</b>	<b>1.075</b>

Table 2: Fold-Change for individuals with/out TBI history and the total in Pathway 2

	<i>BCL2L1</i>	<i>DCAF12</i>	<i>PSMF1</i>	<i>TRIM58</i>
<b>With TBI history</b>	1.085	1.316	1.152	1.573
<b>Without TBI history</b>	1.084	1.341	1.111	1.707
<b>Total</b>	<b>1.085</b>	<b>1.299</b>	<b>1.181</b>	<b>1.487</b>

Table 3: Fold-Change for individuals with/out TBI history and the total in Pathway 3

	<i>ATP6V1G1</i>	<i>BCL2A1</i>	<i>B2M</i>	<i>LY96</i>	<i>S100A8</i>
<b>With TBI history</b>	-0.565	0.527	0.480	0.832	0.448
<b>Without TBI history</b>	0.881	1.091	0.740	1.2849	0.802
<b>Total</b>	<b>0.677</b>	<b>0.708</b>	<b>0.579</b>	<b>0.993</b>	<b>1.764</b>

Table 4: Fold-Change for individuals with/out TBI history and the total in Pathway 4

	<i>PRDX3</i>	<i>RNF139</i>	<i>TMED2</i>	<i>SDHD</i>	<i>SERP1</i>	<i>ZNF654</i>
<b>With TBI history</b>	-0.158	-0.177	-0.289	-0.140	-0.150	0.532
<b>Without TBI history</b>	0.095	0.200	-0.063	0.146	0.199	-0.053
<b>Total</b>	<b>-0.055</b>	<b>-0.024</b>	<b>-0.197</b>	<b>-0.024</b>	<b>-0.008</b>	<b>0.184</b>