# Binomial Thue equations and power integral bases in pure quartic fields

István Gaál<sup>\*</sup> and László Remete University of Debrecen, Mathematical Institute H–4010 Debrecen Pf.12., Hungary e-mail: igaal@science.unideb.hu, hagyma42@gmail.com

October 2, 2018

#### Abstract

It is a classical problem in algebraic number theory to decide if a number field admits power integral bases and further to calculate all generators of power integral bases. This problem is especially delicate to consider in an infinite parametric family of number fields. In the present paper we investigate power integral bases in the infinite parametric family of pure quartic fields  $K = \mathbb{Q}(\sqrt[4]{m})$ .

We often pointed out close connection of various types of Thue equations with calculating power integral bases (cf. [9], [7]). In this paper we determine power integral bases in pure quartic fields by reducing the corresponding index form equation to quartic binomial Thue equations. We apply recent results on the solutions of binomial

<sup>\*</sup>This work was partially supported by the European Union and the European Social Fund through project Supercomputer, the national virtual lab (grant no.: TAMOP-4.2.2.C-11/1/KONV-2012-0010) and also supported in part by K67580 and K75566 from the Hungarian National Foundation for Scientific Research,

<sup>2010</sup> Mathematics Subject Classification: Primar 11R16; Secondary 11D59, 11Y50 Key words and phrases: pure quartic fields, power integral bases, supercomputers

Thue equations of type  $a^4 - mb^4 = \pm 1$  (in  $a, b \in \mathbb{Z}$ ) as well as we perform an extensive calculation by a high performance computer to determine "small" solutions (with  $\max(|a|, |b|) < 10^{500}$ ) of binomial Thue equations for  $m < 10^7$ .

Using these results on binomial Thue equations we characterize power integral bases in infinite subfamilies of pure quartic fields. For  $1 < m < 10^7$  we give all generators of power integral bases with coefficients  $< 10^{1000}$  in the integral basis.

### 1 Introduction

Let K be an algebraic number field of degree n with ring of integers  $\mathbb{Z}_K$ . There is an extensive literature (cf. [7]) of *power integral bases* in K, that is integral bases of the form  $\{1, \gamma, \gamma^2, \ldots, \gamma^{n-1}\}$ . In general, if  $\{1, \omega_2, \ldots, \omega_n\}$  is an integral basis of K, then the discriminant of the linear form  $\omega_2 X_2 + \ldots + \omega_n X_n$  can be written as

$$D_{K/Q}(\omega_2 X_2 + \ldots + \omega_n X_n) = (I(X_2, \ldots, X_n))^2 D_K$$

where  $D_K$  denotes the discriminant of the field K, and  $I(X_2, \ldots, X_n)$  is the *index form* corresponding to the above integral basis. As is known, for any primitive element  $\gamma = x_1 + \omega_2 x_2 + \ldots + \omega_n x_n \in \mathbb{Z}_K$  (that is  $K = \mathbb{Q}(\gamma)$ ) we have

$$I(\gamma) = (\mathbb{Z}_K^+ : \mathbb{Z}[\gamma]^+) = |I(x_2, \dots, x_n)|,$$

where the index of the additive groups of the corresponding rings are taken.  $I(\gamma)$  is the *index of*  $\gamma$ , which does not depend on  $x_1$ . Therefore the element  $\gamma = x_1 + \omega_2 x_2 + \ldots + \omega_n x_n$  generates a power integral basis of K if and only if  $x_1 \in \mathbb{Z}$  and  $(x_2, \ldots, x_n) \in \mathbb{Z}^{n-1}$  is a solution of the *index form equation* 

$$I(x_2, \dots, x_n) = \pm 1$$
 with  $x_2, \dots, x_n \in \mathbb{Z}$ . (1)

This is the way the problem of power integral bases reduces to the resolution of diophantine equations. The elements  $\beta, \gamma \in \mathbb{Z}_K$  are called *equivalent*, if  $\beta \pm \gamma \in \mathbb{Z}$ . In this case  $I(\beta) = I(\gamma)$ . We only consider generators of power integral bases up to equivalence.

There is an extensive literature of index form equations and power integral bases. Until today there are efficient algorithms for calculating generators of power integral bases in number fields of degree up to 6, as well as for some higher degree fields of special structure. (The interested reader should consult I.Gaál [7] for a general overview, for fields up to degree 5 and for special higher degree fields. For sextic fields see Y.Bilu, I.Gaál and K.Győry [4].) Considering this problem not only in a single number field but in an infinite family of fields is an especially challenging task. However we have succeeded in several cases, see e.g. I.Gaál [8], I.Gaál and M.Pohst [11], I.Gaál and T.Szabó [13], [14], I.Gaál [7].

### 2 Pure quartic fields

The purpose of the present paper is to consider power integral bases in pure quartic fields. Recently pure cubic fields were studied by I.Gaál and T. Szabó [12]. Therefore it was straithforward to closely consider the problem of power integral bases in pure quartic fields that has not been investigated before. The main goal of this paper is to show that the problem of power integral bases in pure quartic fields can be reduced to quartic binomial Thue equations.

Let *m* be a positive integer and consider the pure quartic field  $K = \mathbb{Q}(\sqrt[4]{m})$ . Set  $\alpha = \sqrt[4]{m}$ . Assume that *m* is square free,  $m \neq \pm 1$  and  $m \equiv 2, 3 \pmod{4}$ . According to A.Hameed, T.Nakahara, S.M.Husnine and S.Ahmad [16]  $\{1, \alpha, \alpha^2, \alpha^3\}$  is an integer basis in *K*. We shall use this integer basis to construct the index form equation therefore in the following we continue to assume these conditions on *m*.

Using results on binomial Thue equations and making direct calculations we obtained the following statements.

As we shall see it is easy to deal with negative values of m.

**Theorem 1.** Let m < -1 be a square free integer with  $m \equiv 2, 3 \pmod{4}$ . Then up to equivalence  $\alpha = {}^4\sqrt{m}$  is the only generator of power integral bases in  $K = \mathbb{Q}({}^4\sqrt{m})$ .

Next we consider values of m composed of at most two primes less than or equal to 29.

**Theorem 2.** Let m > 1 be a square free integer with  $m \equiv 2,3 \pmod{4}$ composed of at most two primes p, q with  $2 \leq p, q \leq 29$ . Set  $\alpha = \sqrt[4]{m}$ and let  $K = \mathbb{Q}(\sqrt[4]{m})$ . Then up to equivalence the only generators of power integral bases in  $K = \mathbb{Q}(\sqrt[4]{m})$  are  $\vartheta = x\alpha + y\alpha^2 + z\alpha^3$  with (x, y, z) = (1, 0, 0) for arbitrary m,  $(x, y, z) = (4, \pm 2, 1)$  for m = 15,  $(x, y, z) = (25, \pm 10, 4)$  for m = 39.

We construct an infinite series of values of m for which the corresponding binomial Thue equation is easy to solve therefore we can characterize the power integral bases in  $K = \mathbb{Q}(\sqrt[4]{m})$ .

**Theorem 3.** Let s, t be positive integer parameters and let

$$m = \frac{(s^4t \pm 1)^4 - 1}{s^4}$$

be a square free integer with m > 1 and  $m \equiv 2,3 \pmod{4}$ . Set  $\alpha = \sqrt[4]{m}$ . Then up to equivalence the only generators of power integral bases in  $K = \mathbb{Q}(\sqrt[4]{m})$  are  $\vartheta = x\alpha + y\alpha^2 + z\alpha^3$  with  $(x, y, z) = (1, 0, 0), ((s^4t \pm 1)^2, \pm (s^4t \pm 1)s, s^2).$ 

The following theorem uses results on binomial Thue equations for small values of m.

**Theorem 4.** Let 1 < m < 2000 be a square free integer, with  $m \equiv 2, 3$  (mod 4). For 400 < m < 2000 we also assume that m is odd. Set  $\alpha = {}^{4}\sqrt{m}$  and let  $K = \mathbb{Q}({}^{4}\sqrt{m})$ . Then up to equivalence the only generators of power integral bases in  $K = \mathbb{Q}({}^{4}\sqrt{m})$  are  $\vartheta = x\alpha + y\alpha^{2} + z\alpha^{3}$  with (x, y, z) = (1, 0, 0) for arbitrary m,  $(x, y, z) = (1, \pm 1, 1)$  for m = 2,  $(x, y, z) = (4, \pm 2, 1)$  for m = 15,  $(x, y, z) = (25, \pm 10, 4)$  for m = 39,  $(x, y, z) = (9, \pm 3, 1)$  for m = 82,  $(x, y, z) = (16, \pm 4, 1)$  for m = 255,  $(x, y, z) = (121, \pm 22, 4)$  for m = 915,  $(x, y, z) = (36, \pm 6, 1)$ , for m = 1295.

Our last theorem was obtained by an extensive calculation using a supercomputer of "small" solutions of binomial Thue equations. The range  $m < 10^7$  and the bound  $10^{1000}$  for the solutions must be sufficient for practical applications.

**Theorem 5.** Let  $1 < m < 10^7$  be a square free integer with  $m \equiv 2, 3 \pmod{4}$ . 4). Set  $\alpha = {}^4\sqrt{m}$  and let  $K = \mathbb{Q}({}^4\sqrt{m})$ . Then up to equivalence the only generators  $\vartheta = x\alpha + y\alpha^2 + z\alpha^3$  of power integral bases in  $K = \mathbb{Q}({}^4\sqrt{m})$  with

 $\max(|x|, |y|, |z|) < 10^{1000}$ 

are (x, y, z) = (1, 0, 0) for arbitrary m, moreover the following (x, y, z) for

the given values of m:

m	x	$\pm y$	z
2	1	1	1
15	4	2	1
39	25	10	4
82	9	3	1
255	16	4	1
410	81	18	4
626	25	5	1
915	121	22	4
1295	36	6	1
2402	49	7	1
6562	81	9	1
12155	441	42	4
14642	121	11	1
17490	529	46	4
20735	144	12	1
24414	625	50	4
28562	169	13	1
33215	729	54	4
38415	196	14	1
50626	225	15	1
61535	3969	252	16
65535	256	16	1
83522	289	17	1
130322	361	19	1
144590	1521	78	4
159999	400	20	1
194482	441	21	1
234255	484	22	1
279842	529	23	1
390626	625	25	1
505679	6400	240	9
531442	729	27	1

m	x	$\pm y$	z
707282	841	29	1
757335	3481	118	4
809999	900	30	1
923522	961	31	1
1081730	16641	516	16
1185922	1089	33	1
1336335	1156	34	1
1416695	4761	138	4
1500626	1225	35	1
1679615	1296	36	1
1755519	33124	910	25
1874162	1369	37	1
1977539	5625	150	4
2313442	1521	39	1
2559999	1600	40	1
2825762	1681	41	1
3111695	1764	42	1
3262539	7225	170	4
3418802	1849	43	1
3580610	7569	174	4
3851367	196249	4430	100
4100626	2025	45	1
4879682	2209	47	1
5764802	2401	49	1
6765202	2601 51		1
7034430	10609	206	4
7311615	2704	52	1
7596914	11025	210	4
7890482	2809	53	1
8503055	2916	54	1
8608519	105625	1950	36
9150626	150626 3025 55		1
9834495	3136	56	1

г

## 3 Auxiliary tools I: Index form equations in quartic fields

Calculating generators of power integral bases is equivalent to solving the index form equation (1). For quartic number fields it is an equation of degree 6 in 3 variables. However by using the method of I.Gaál, A.Pethő and M.Pohst [10] (see also I.Gaál [7]) this can be reduced to cubic and quartic Thue equations that can be solved by using standard technics (cf. [7]). We cite here this result in detail since this will be a basic tool in our proofs.

Let K be a quartic number field generated by a root  $\xi$  with minimal polynomial  $f(x) = x^4 + a_1 x^3 + a_2 x^2 + a_3 x + a_4 \in \mathbb{Z}[x]$ . We represent any  $\vartheta \in \mathbb{Z}_K$  in the form

$$\vartheta = \frac{1}{d} \left( a + x\xi + y\xi^2 + z\xi^3 \right) \tag{2}$$

with coefficients  $a, x, y, z \in \mathbb{Z}$  and with a common denominator  $d \in \mathbb{Z}$ . Let  $n = I(\xi)$ ,

$$F(u,v) = u^3 - a_2 u^2 v + (a_1 a_3 - 4a_4) uv^2 + (4a_2 a_4 - a_3^2 - a_1^2 a_4) v^3$$

a binary cubic form over  $\mathbb{Z}$  and

$$Q_1(x, y, z) = x^2 - xya_1 + y^2a_2 + xz(a_1^2 - 2a_2) + yz(a_3 - a_1a_2) + z^2(-a_1a_3 + a_2^2 + a_4)$$
  

$$Q_2(x, y, z) = y^2 - xz - a_1yz + z^2a_2$$

ternary quadratic forms over  $\mathbb{Z}$ .

**Lemma 6.** (I.Gaál, A.Pethő and M.Pohst [10]) If  $\vartheta$  of (2) generates a power integral basis in K then there is a solution  $(u, v) \in \mathbb{Z}^2$  of

$$F(u,v) = \pm \frac{d^6m}{n} \tag{3}$$

such that

$$u = Q_1(x, y, z),$$
  
 $v = Q_2(x, y, z).$  (4)

### 4 Auxiliary tools II: Binomial Thue equations

Let g > 0 be a given integer. The index form equation in pure quartic fields will be reduced to a *binomial Thue equation* of type

$$a^4 - gb^4 = \pm 1 \quad (\text{in } a, b \in \mathbb{Z}).$$
 (5)

In order to solve the index form equation we shall use some corresponding results on binomial Thue equations.

If we know a positive solution of the binomial Thue equation, then the following result of M.A.Bennett is useful to exclude any other positive solutions.

**Lemma 7.** (M.A.Bennett [3]) If f, g are integers with  $fg \neq 0$  and  $n \geq 3$  then the equation

$$fa^n - gb^n = \pm 1 \quad (\text{in } a, b \in \mathbb{Z})$$

has at most one solution in positive integers (a, b).

The following theorem describes all solutions of binomial Thue equation if its coefficients are composed of at most two primes less than or equal to 29.

**Lemma 8.** (K.Győry and Á.Pintér [15]) If f < g are positive integers composed of at most two primes p, q with  $2 \le p, q \le 29$  and  $n \ge 3$  then the only solutions of the equation

$$fa^n - gb^n = \pm 1$$
 (in  $a, b \in \mathbb{Z}$ )

are those with

$$f \in \{1, 2, 3, 4, 7, 8, 16\}, a = b = 1,$$

and

$$n = 3, (f, a) = (1, 2), (1, 3), (1, 4), (1, 8), (1, 9), (1, 18), (1, 19), (1, 23),$$

$$(2, 2), (3, 2), (5, 3), (5, 11), (11, 6),$$
  

$$n = 4, (f, a) = (1, 2), (1, 3), (1, 5), (3, 2),$$
  

$$n = 5, (f, a) = (1, 2), (1, 3), (9, 2),$$
  

$$n = 6, (f, a) = 1, 2).$$

The following results of A.Bazsó, A.Bérczes, K.Győry and Á.Pintér [1] describe the solutions of binomial Thue equations with f = 1. For simplicity we quote the case n = 4 only, that is we consider equation (5).

**Lemma 9.** (A.Bazsó, A.Bérczes, K.Győry and Á.Pintér [1]) If  $1 < g \leq 400$  then all positive integer solutions (a, b), with |ab| > 1 of equation (5) are

> (g, a, b) = (5, 3, 2), (15, 2, 1), (17, 2, 1), (39, 5, 2),(80, 3, 1), (82, 3, 1), (150, 7, 2), (255, 4, 1), (257, 4, 1).

**Lemma 10.** (A.Bazsó, A.Bérczes, K.Győry and Á.Pintér [1]) If  $400 < g \leq 2000$  is odd, then all positive integer solutions (a,b), with |ab| > 1 of equation (5) are

$$(g, a, b) = (915, 11, 2), (1295, 6, 1), (1297, 6, 1), (1785, 13, 2).$$

Remark. The solutions with m = 82,915,1295 of Lemmas 9, 10 are indicated in the Corregindum [2] to [1].

Our last theorem was obtained by an extensive computation using a high performance computer. We calculate the "small" solutions up to  $10^{500}$  of binomial Thue equations (5) for g up to  $10^7$ . Note that in several cases the calculation of fundamental units of the field  $K = \mathbb{Q}(\sqrt[4]{g})$  is extremely time consuming. For example computing the fundamental unit in  $K = \mathbb{Q}(\sqrt[4]{18071})$ takes more than 10 minutes using Kash [6] and the coefficients of the fundamental unit in the integral basis have ca. 1000 digits. Therefore using the present machinery it does not seem feasible to completely solve all these Thue equations. However, according to our experience, these solutions yield all solutions with very high probability.

Theorem 11. All solutions with

$$\max(|a|, |b|) < 10^{500}$$

of the binomial Thue equation (5) for  $1 < g < 10^7$  (with irreducible left hand side) are listed in the following table.

g	a	Ь	
2	1	1	6
5	3	2	6
15	2	1	6
17	2	1	7
39	5	2	8
80	3	1	8
82	3	1	- 9
150	7	2	10
255	4	1	10
257	4	1	11
410	9	2	11
624	5	1	13
626	5	1	13
915	11	2	14
1295	6	1	15
1297	6	1	16
1785	13	2	17
2400	7	1	19
2402	7	1	19
3164	15	2	21
4095	8	1	23
4097	8	1	23
5220	17	2	25
6560	9	1	27
6562	9	1	27
7140	239	26	30
8145	19	2	-33
9999	10	1	-33
10001	10	1	36
12155	21	2	39
14640	11	1	39
14642	11	1	42
17490	23	2	45
20735	12	1	45
20737	12	1	49
24414	25	2	50
28560	13	1	51
28562	13	1	53
33215	27	2	53
38415	14	1	54
38417	14	1	55
44205	29	2	57
50624	15	1	61
50626	15	1	61
57720	31	2	65
61535	63	4	70

g	a	b	
5535	16	1	
5537	16	1	
9729	65	4	
4120	33	2	
3520	17	1	
3522	17	1	
3789	35	2	
04975	18	1	
04977	18	1	
14240	239	13	
17135	37	2	
30320	19	1	
30322	19	1	
$44\overline{590}$	39	2	
59999	20	1	
50001	20	1	
76610	41	2	
94480	21	1	
94482	21	1	
13675	43	2	
34255	22	1	
34257	22	1	
56289	45	2	
79840	23	1	
79842	23	1	
)4980	47	2	
31775	24	1	
31777	24	1	
50300	49	2	
90624	25	1	
90626	25	1	
22825	51	2	
56975	26	1	
56977	26	1	
93155	53	2	
J5679	80	3	
18440	161	6	
31440	27	1	
31442	27	1	
44685	163	6	
58175	82	3	
71914	55	2	
14655	28	1	
14657	28	1	
59750	57	2	
)7280	29	1	

*g* 

 $\begin{array}{c} 1048577 \\ 1081730 \\ 1115664 \end{array}$ 

 $\frac{1185922}{1259445}$ 

 $\frac{1336335}{1336337}$ 

 $\frac{1588230}{1679615}$ 

 $\frac{1977539}{2085135}$ 

 a

b	l ſ	g	a	b
1		3580610	87	2
2		3748095	44	1
1		3748097	44	1
1		3851367	443	10
2		3921390	89	2
1		4100624	45	1
1		4100626	45	1
2		4285935	91	2
4		4477455	46	1
1		4477457	46	1
1		4675325	93	2
4		4879680	47	1
2		4879682	47	1
1		5090664	95	2
1		5198685	191	4
2		5308415	48	1
1		5308417	48	1
1		5419875	193	4
2		5533080	97	2
1		5764800	49	1
1		5764802	49	1
2		6003725	99	2
1		6249999	50	1
1	ſ	6250001	50	1
5		6503775	101	2
2		6765200	51	1
1		6765202	51	1
1		7034430	103	2
2		7311615	52	1
1		7311617	52	1
1		7596914	105	2
2		7890480	53	1
1		7890482	53	1
1		8192475	107	2
2		8295040	161	3
1		8398565	323	6
1		8503055	54	1
2		8503057	54	1
1		8608519	325	6
1		8714960	163	3
2		8822385	109	2
1		9150624	55	1
1		9150626	55	1
2		9487940	111	2
1		9834495	56	1
1	ll	9834497	56	1

### 5 Proofs

Let  $\alpha = {}^{4}\sqrt{m}$  (*m* is square free,  $m \neq \pm 1$ , with  $m \equiv 2,3 \pmod{4}$ ). Then by the result of A.Hameed, T.Nakahara, S.M.Husnine and S.Ahmad [16]  $\{1, \alpha, \alpha^{2}, \alpha^{3}\}$  is an integer basis in the pure quartic field  $K = \mathbb{Q}(\alpha)$ . This means  $n = I(\alpha) = 1$  and any  $\vartheta \in \mathbb{Z}_{K}$  can be represented in the form (2) with d = 1. Therefore the equations (3) and (4) can be written in the following form:

$$u(u^2 + 4mv^2) = \pm 1, (6)$$

$$x^2 - mz^2 = u, (7)$$

$$y^2 - xz = v. ag{8}$$

Equation (6) implies  $u = \pm 1$  and  $u^2 + 4mv^2 = \pm 1$ , whence v = 0.

**Case I.** Let m < 0. Equation (7) implies x = 1 and z = 0. By equation (8) we get y = 0. This yields that up to equivalence  $\vartheta = \alpha$  is the only generator of power integral bases. This proves Theorem 1.

**Case II.** Let m > 1. Equation (7), that is  $x^2 - mz^2 = \pm 1$  implies that gcd(x, z) = 1. Equation (8), that is  $y^2 = xz$  implies that x and z are both squares. (Note that if (x, y, z) is a solution then so also is (-x, -y, -z), therefore we may assume that x, z are both nonnegative.)

Set  $x = a^2$  and  $z = b^2$  with  $a, b \in \mathbb{Z}$ . Then equation (7) yields the binomial Thue equation

$$a^4 - mb^4 = \pm 1 \quad (\text{in } a, b \in \mathbb{Z}).$$
 (9)

All solutions (up to sign) of (7) and (8) are obtained by

$$x = a^2, \ y = \pm ab \ z = b^2$$
 (10)

where a, b are solutions of (9).

Theorem 2 is a consequence of Lemma 8.

To prove Theorem 3 observe that  $(a, b) = (s^4t \pm 1, s)$  is a positive solution of equation (9). By Lemma 7 this and (a, b) = (1, 0) are all solutions of this equation. (x, y, z) are calculated from (a, b) by (10).

Theorem 4 is a consequence of Lemma 9 and of Lemma 10.

Theorem 11 was proved by calculating "small" solutions of binomial Thue equations (9) with  $\max(|a|, |b|) < 10^{500}$  by using the method of Pethő [17]. The procedure was implemented in Maple [5], we used 1200 digits accuracy. This extensive calculation involving almost  $10^7$  binomial Thue equations was performed on the supercomputer (high performance computer) network situated in Debrecen-Budapest-Pécs-Szeged in Hungary. The total running time was about 120 hours calculated for a single node which yields a few hours using parallel computing with a couple of nodes.

Theorem 5 is a consequence of Theorem 11 using (10).

### References

- [1] A.Bazsó, A.Bérczes, K.Győry and Á.Pintér, On the resolution of equations  $Ax^n - By^n = C$  in integers x, y and  $n \ge 3$ , II, Publ. Math. (Debrecen), **76**(2010), 227–250.
- [2] A.Bazsó, A.Bérczes, K.Győry and Á.Pintér, Corregindum to the paper "On the resolution of equations  $Ax^n - By^n = C$  in integers x, y and  $n \ge 3$ , II", Publ. Math. (Debrecen), to appear.
- [3] M.A.Bennett, Rational approximation to algebraic numbers of small height: The Diophantine equation |ax<sup>n</sup> by<sup>n</sup>| = 1, J. Reine Angew. Math. 535(2001), 1-49.
- [4] Y.Bilu, I.Gaál and K.Győry, Index form equations in sextic fields: a hard computation, Acta Arithm., 115(2004), 85-96.
- [5] B.W.Char, K.O.Geddes, G.H.Gonnet, M.B.Monagan, S.M.Watt (eds.) MAPLE, Reference Manual, Watcom Publications, Waterloo, Canada, 1988.
- [6] M. Daberkow, C.Fieker, J.Klüners, M.Pohst, K.Roegner and K.Wildanger, KANT V4, J.Symbolic Comput., 24(1997), 267–283.
- [7] I.Gaál, *Diophantine equations and power integral bases*, Boston, Birkhäuser, 2002.

- [8] I.Gaál, Power integral bases in orders of families of quartic fields, Publ. Math. (Debrecen), 42 (1993), 253–263.
- [9] I.Gaál, Application of Thue equations to computing power integral bases in algebraic number fields, Proc. Conf. ANTS II, Talence, France, 1996. Lecture Notes in Computer Science 1122, Springer 1996, pp. 151–155.
- [10] I.Gaál, A.Pethő and M.Pohst, Simultaneous representation of integers by a pair of ternary quadratic forms – with an application to index form equations in quartic number fields, J.Number Theory, 57(1996), 90–104.
- [11] I.Gaál and M.Pohst, Power integral bases in a parametric family of totally real quintics, Math. Comput., 66(1997), 1689–1696.
- [12] I.Gaál and T. Szabó, A note on the minimal indices of pure cubic fields, JP Journal of Algebra, Number Theory and Applications, 19(2010), 129–139
- [13] I.Gaál and T. Szabó, Power integral bases in parametric families of biquadratic fields, JP Journal of Algebra, Number Theory and Applications, 21(2012), 105–114.
- [14] I.Gaál and T.Szabó, Relative power integral bases in infinite families of quartic extensions of quadratic field, JP Journal of Algebra, Number Theory and Applications, 29(2013), 31–43.
- [15] K.Győry and A.Pintér, Binomial Thue equations, ternary equations, and power values of polynomials, J. Math. Sciences 180(2012), 569–580.
- [16] A.Hameed, T.Nakahara, S.M.Husnine and S.Ahmad, On the existence of canonical number system in certain classes of pure algebraic number fields, J. Prime Research in Math., 7(2011), 19–24.
- [17] A.Pethő, On the resolution of Thue inequalities, J.Symbolic Comput., 4(1987), 103–109.