# MOST VERTEX SUPERALGEBRAS ASSOCIATED TO AN ODD UNIMODULAR LATTICE OF RANK 24 HAVE AN N=4 SUPERCONFORMAL STRUCTURE

## GERALD HÖHN AND GEOFFREY MASON

ABSTRACT. Odd, positive-definite, integral, unimodular lattices N of rank 24 were classified by Borcherds. There are 273 isometry classes of such lattices. Associated to them are vertex superalgebras  $V_N$  of central charge  $c{=}24$ . We show that at least 267 of these vertex operator superalgebras contain an  $N{=}4$  superconformal subalgebra of central charge  $c'{=}6$ . This is achieved by studying embeddings  $L^{+}{\subseteq}N$  of a certain rank 6 lattice  $L^{+}$ .

### 1. Introduction

The results discussed in this paper concern positive-definite, integral, unimodular lattices of rank 24 and the vertex algebras that they define. We find it very convenient to call any such lattice a *Niemeier* lattice, adding the adjective *odd* or *even* when appropriate. Thus for us an *even Niemeier lattice* is what is now called simply a Niemeier lattice in the literature.

More often than not, the basic axiom for a vertex algebra, i.e., the *Jacobi identity*, is supplemented by additional conditions such as the existence of a Virasoro field, or by a superconformal structure. In the case of vertex superalgebras there is a hierarchy of such structures, often denoted by N=1, N=2, N=4, ... becoming more intricate with increasing N. For general background we refer the reader to [4], [9], [8] and for an early comparison of these super structures, see [6].

Vertex algebras containing an  $N{=}4$  superconformal structure of central charge 6, in particular, have come into recent prominence because of their intervention in  $Mathieu\ Moonshine\ [5]$ . Experience suggests that moonshine-like phenomena may be related to Niemeier lattices as we have defined them and thus one is lead to consider odd Niemeier lattices N of rank 24 and the vertex superalgebras  $V_N$  that they define. A natural question presents itself: does  $V_N$  have an  $N{=}4$  structure? In this paper we will show that most of the vertex operator superalgebras  $V_N\ do$  have an  $N{=}4$  superconformal structure.

This question was considered in [10] for general lattice vertex superalgebras  $V_L$ . However, the precise definition of the (small)  $N{=}4$  algebra is awkward to handle (cf. [10], Section 2) and makes the question hard to deal with. A way around this difficulty was proposed in [10]. Namely, it was shown (loc. cit., Proposition 26) that a certain integral rank 6 lattice which we denote by  $L^+$ , has the property that  $V_{L^+}$  contains an  $N{=}4$  superconformal algebra A and that  $V_{L^+}$  and A share the same Virasoro element (the canonical one defined by the Siegel-Sugawara construction).

<sup>1991</sup> Mathematics Subject Classification. MSC(2010): 17B69, 17B81.

Key words and phrases. Niemeier lattice, N=4 algebra, vertex superalgebra.

The first author was supported by the Simons Foundation Collaboration Grant #355294.

The second author was supported by the Simons Foundation #427007.

The lattice  $L^+$  is spanned by elements  $\alpha_1, \ldots, \alpha_6$  and  $h:=\frac{1}{3}(\alpha_1+\cdots+\alpha_6)$  and equipped with a bilinear form satisfying  $(\alpha_i, \alpha_j)=3$   $\delta_{ij}$ ,  $(i, j=1, \ldots, 6)$ .

In this way one can establish the existence of an N=4 superconformal subalgebra of central charge 6 in  $V_N$  by proving the existence of a sublattice  $L^+\subseteq N$ , for we then have the tower  $A\subseteq V_{L^+}\subseteq V_N$ . Our main results, obtained by a computer search, identify which odd Niemeier lattices admit such an embedding.

### 2. Results

In order to state our main results, we need to review the theory of Niemeier lattices [1, 3]. The important idea of *neighboring* lattices is due to Kneser. Suppose that A and B are a pair of equal rank lattices. We say that A and B are *neighbors* in case their intersection D (in a common Euclidean space E containing both) has index 2 in each of them. If A and B are neighboring even Niemeier lattices then there is a unique odd Niemeier lattice N that neighbors both A and B. (A, B and A are the lattices in E containing D with index 2). Furthermore, every odd Niemeier lattice N arises in this way from an unordered pair (A, B) of even Niemeier lattices that is unique up to isometry. We say that A has A has A neven Niemeier lattice is characterized by its (semisimple) root system (which by convention is empty in the case of the Leech lattice A). Thus the type of an odd Niemeier lattice A is determined by the corresponding pair (A, A, A, of root systems, which we also refer to as the A type of A.

We are also interested in the *minimum norm*  $\mu$  of a Niemeier lattice N, i.e., the least value of  $(\alpha, \alpha)$  for  $0 \neq \alpha \in N$ . If N is even then  $\mu = 2$  with the single exception of the Leech lattice  $\Lambda$  which has  $\mu = 4$ . For odd N there are three possibilities as in the following table:

TABLE 1. Minimum norms for odd Niemeier lattices

$$\begin{array}{c|ccccc} \mu & 1 & 2 & 3 \\ \hline \# \text{ of classes} & 116 & 156 & 1 \\ \end{array}$$

The unique lattice of minimum norm 3 is called the *odd Leech lattice*  $\Lambda^{\text{odd}}$ ; it has type  $(\emptyset, A_1^{24})$ .

If N is an odd Niemeier lattice of minimum norm 1, its two even Niemeier neighbors are necessarily isometric. This is because the Weyl reflection determined by a norm 1 vector *exchanges* the two even neighbors. Thus the type of such a lattice always takes the form  $(\Phi,\Phi)$  where  $\Phi$  is the root system common to the even neighbors.

This situation dos not necessarily pertain in the case of odd Niemeier lattices with  $\mu$ = 2. The *neighboring graph*, with nodes given by even Niemeier lattices and edges between A and B indexed by the odd lattices of type (A, B) and with  $\mu$ =2 was first constructed by Borcherds [1]. It is Figure 17.1 in [3].

We can now state our main results.

**Theorem 1.** Suppose that N is an odd Niemeier lattice. Then the following hold: (1) If  $\mu=1$ , then either N is of type  $(\emptyset,\emptyset)$ , or else there is a sublattice  $L^+ \subseteq N$ .

- (2) If  $\mu$ =2, exactly one of the following holds:
  - (a) N has type  $(E_8^3, D_8^3)$ ,  $(D_{16}E_8, A_{15}D_9)$ ,  $(D_{12}^2, A_{12}^2)$  or  $(D_{24}, A_{24})$ ,
  - (b) there exists a sublattice  $L^+ \subseteq N$ .
- (3) If  $\mu$ =3, the odd Leech lattice  $\Lambda$ <sup>odd</sup> does not contain L<sup>+</sup>.
- Remark 2. (i) There is a unique odd Niemeier lattice for each of the types listed in Theorem 1(2)(a). Thus there are exactly 152 isometry classes of odd Niemeier lattices of minimum norm 2 that contain the lattice  $L^+$ .
- (ii) The odd Leech lattice  $\Lambda^{\text{odd}}$  cannot possibly contain  $L^+$  because the even sublattice of  $\Lambda^{\text{odd}}$  is contained in  $\Lambda$  and hence has minimum norm  $\geq 4$ , whereas  $L^+$  contains vectors of norm 2 (namely  $\pm h$ ).
- (iii) For similar reasons, the unique lattice of type  $(\emptyset, \emptyset)$  in Theorem 1 (1) cannot contain  $L^+$ . Thus we can paraphrase Theorem 1 (1) as follows: every one of the 116 odd Niemeier lattices with  $\mu$ =1 that *could* contain  $L^+$ , *does* contain  $L^+$ .

The proof of Theorem 1 is computational. We used the neighborhood method as implemented in MAGMA [2] together with some optimizations to create the list of 273 odd Niemeier lattices together with their two even neighbors. We also computed the automorphism group and checked the result by a mass formula ([3], Chapter 16).

The lattice  $L^+$  has a basis consisting of the vector h of norm 2 and the five pairwise orthogonal norm 3 vectors  $\alpha_1, \ldots, \alpha_5$  with scalar product  $(\alpha_i, h) = 1$ . For each odd Niemeier lattice N, this permits use of the following search method for the sublattice  $L^+$ : do a backtracking search by selecting all possible norm 2 vectors h in N, then for given vector h take all norm 3 vectors  $\alpha_1$  with  $(h, \alpha_1) = 1$ , then for such pairs  $\{h, \alpha_1\}$  select all norm 3 vectors  $\alpha_2$  with  $(h, \alpha_2) = 1$  and  $(\alpha_1, \alpha_2) = 0$ , then, in similar way, all possible vectors  $\alpha_3$ ,  $\alpha_4$  and  $\alpha_5$ . If we find a tuple  $(h, \alpha_1, \ldots, \alpha_5)$  we stop our search for the lattice N. Otherwise we continue until all possibilities are exhausted.

To speed up the computation, we used the automorphism group O(N) of the lattice N. First we compute the orbits of O(N) on the roots of norm 2. Then it is enough to choose a random vector h in each orbit. Then we compute the pointwise stabilizer  $H_1$  of h in O(N) and decompose the set of norm 3 vectors  $\alpha$  with  $(h,\alpha)=1$  into  $H_1$ -orbits. Then it is enough to choose a random vector  $\alpha_1$  in each such orbit. Then we compute the pointwise stabilizer  $H_2$  of the two vectors h and  $\alpha_1$  in O(N) and select a random vector among the possible  $\alpha_2$  in each  $H_2$ -orbit. For  $\alpha_3$ ,  $\alpha_4$  and  $\alpha_5$  we went through all possible choices since this was quicker than computing additional stabilizers.

- *Remark* 3. (i) The computation took less than 24 hours on a single processor machine.
- (ii) We did not classify all orbits of sublattices  $L^+ \subseteq N$  under the action of O(N), although this can easily be achieved with our method.
- (iii) We also tried a different computational approach. We constructed a single embedding of  $L^+$  into an odd Niemeier lattice N and determined the orthogonal complement lattice  $K = (L^+ \otimes \mathbf{R})^\perp \cap N$ . Then we tried the neighborhood method to compute all lattice K' in the genus of K. By computing all gluings of all such lattices K' with  $L^+$ , we obtain odd Niemeier lattices N' having  $L^+$  as a sublattice and all such lattices can be constructed that way.

We found more than 10,000 rank 18 lattices K' covering most of the genus of K. However, the estimated run time for completing the calculation was more than a month. Apart from a few missing cases, the list of lattices N' obtained in this way was identical with the one from our theorem.

The following two questions remain open:

- (i) Do the four exceptional lattice vertex superalgebras  $V_N$  in part (2) of Theorem 1 have an  $N{=}4$  super conformal structure not coming from a lattice embedding  $L^+ \subset N$ ?
- (ii) Do most of the self-dual lattice vertex superalgebras of central charge 24 (cf. [7]), which are not of lattice type have an N=4 superconformal structure?

#### REFERENCES

- [1] R. E. Borcherds, The Leech lattice and other lattices, Ph.D. thesis, Cambridge, 1985.
- [2] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput., **24** (1997), 235–265.
- [3] J. Conway and N. Sloane, *Sphere packings, Lattices and Groups*, Grundleheren No. **290**, Springer, New York, 1988.
- [4] C. Dong and J. Lepowsky, *Generalized Vertex Algebras and Relative Vertex Operators*, Progress in Mathematics No. **112**, Birkhäuser, Boston., 1993.
- [5] T. Eguchi, Tachikawa and H. Ooguri, Notes on the K3 surface and the Mathieu group  $M_{24}$ , Exper. Math. **20**, 91-96 (2011).
- [6] M. Green, J. Schwarz and E. Witten, *Superstring Theory* Vol.1, Cambridge Monographs in Mathematical Physics, C.U.P., 1987.
- [7] G. Höhn, *Selbstduale Vertexoperatorsuperalgebren und das Babymonster*, Bonner Mathematische Schriften **286** (1996), 1–85.
- [8] V. Kac, Vertex Algebras for Beginners, Second Ed., Amer. Math. Soc., 1998.
- [9] J. Lepowsky and H. Li, *Introduction to Vertex Operator Algebras and Their Representations*, Progress in Mathematics No. **227**, Birkhäuser, Boston, 2004.
- [10] G. Mason, M. Tuite and G. Yamskulna, N=2 and N=4 subalgebras of super vertex operator algebras, J. Phys. A: Math and Theoretical Vol. **51**, No. 6. (2017).

DEPARTMENT OF MATHEMATICS, KANSAS STATE UNIVERSITY, MANHATTAN, KS 66506 *E-mail address*: gerald@monstrous-moonshine.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT SANTA CRUZ, CA 95064 *E-mail address*: gem@ucsc.edu