

**ON THE CAUCHY PROBLEM FOR NONDEGENERATE
PARABOLIC INTEGRO-DIFFERENTIAL EQUATIONS IN
THE SCALE OF GENERALIZED HÖLDER SPACES**

REMIGIJUS MIKULEVIČIUS AND FANHUI XU

ABSTRACT. Parabolic integro-differential nondegenerate Cauchy problem is considered in the scale of Hölder spaces of functions whose regularity is defined by a radially O-regularly varying Lévy measure. Existence and uniqueness and the estimates of the solution are derived.

CONTENTS

1.	Introduction	1
2.	Notation, function spaces and norm equivalence	5
2.1.	Basic notation	5
2.2.	Function spaces and norm equivalence	6
3.	Proof of main theorem	21
3.1.	Hölder estimates of the smooth solution	22
3.2.	General Hölder inputs	26
4.	Appendix	27
	References	32

1. INTRODUCTION

Let $\alpha \in (0, 2)$ and \mathfrak{A}^α be the class of all nonnegative measures ν on $\mathbf{R}_0^d = \mathbf{R}^d \setminus \{0\}$ such that $\int |y|^2 \wedge 1 d\nu < \infty$ and

$$\alpha = \inf \left\{ \sigma < 2 : \int_{|y| \leq 1} |y|^\sigma d\nu < \infty \right\}.$$

In addition, we assume that for $\nu \in \mathfrak{A}^\alpha$,

$$\begin{aligned} \int_{|y| > 1} |y| d\nu &< \infty \text{ if } \alpha \in (1, 2), \\ \int_{R < |y| \leq R'} y d\nu &= 0 \text{ if } \alpha = 1 \text{ for all } 0 < R < R' < \infty. \end{aligned}$$

Date: August 28, 2018.

1991 Mathematics Subject Classification. 35R09, 60J75, 35B65.

Key words and phrases. non-local parabolic integro-differential equations, Lévy processes.

In this paper we consider the parabolic Cauchy problem with $\lambda \geq 0$

$$(1.1) \quad \begin{aligned} \partial_t u(t, x) &= Lu(t, x) - \lambda u(t, x) + f(t, x) \text{ in } H_T = [0, T] \times \mathbf{R}^d, \\ u(0, x) &= 0, \quad x \in \mathbf{R}^d, \end{aligned}$$

and integro-differential operator

$$L\varphi(x) = L^\nu \varphi(x) = \int [\varphi(x+y) - \varphi(x) - \chi_\alpha(y) y \cdot \nabla \varphi(x)] \nu(dy), \quad \varphi \in C_0^\infty(\mathbf{R}^d),$$

where $\nu \in \mathfrak{A}^\alpha$, $\chi_\alpha(y) = 0$ if $\alpha \in (0, 1)$, $\chi_\alpha(y) = 1_{\{|y| \leq 1\}}(y)$ if $\alpha = 1$, and $\chi_\alpha(y) = 1$ if $\alpha \in (1, 2)$. Given a Lévy measure $\nu \in \mathfrak{A}^\alpha$ on $\mathbf{R}_0^d = \mathbf{R}^d \setminus \{0\}$, there exist a Poisson random measure $J(ds, dy)$ on $[0, \infty) \times \mathbf{R}_0^d$ such that

$$\mathbf{E}[J(ds, dy)] = \nu(dy) ds,$$

and a Lévy process Z_t^ν so that

$$(1.2) \quad Z_t^\nu = \int_0^t \int_{\mathbf{R}_0^d} \chi_\alpha(y) y \tilde{J}(ds, dy) + \int_0^t \int_{\mathbf{R}_0^d} (1 - \chi_\alpha(y)) y J(ds, dy), \quad t \geq 0,$$

with $\tilde{J}(ds, dy) = J(ds, dy) - \nu(dy) ds$. For $\nu \in \mathfrak{A}^\alpha$, set

$$\begin{aligned} \delta(r) &= \delta_\nu(r) = \nu(\{|y| > r\}), \quad r > 0, \\ w(r) &= w_\nu(r) = \delta_\nu(r)^{-1}, \quad r > 0. \end{aligned}$$

Our main assumption is that $w(r) = w_\nu(r) = \delta_\nu(r)^{-1}$, $r > 0$, is an O-RV function (O-regular variation function) at zero (see [2] and [3]), that is

$$r_1(\varepsilon) = \lim_{x \rightarrow 0} \frac{\delta(\varepsilon x)^{-1}}{\delta(x)^{-1}} < \infty, \quad \varepsilon > 0.$$

By Theorem 2 in [2], the following limits exist:

$$(1.3) \quad p_1 = p_1^\nu = \lim_{\varepsilon \rightarrow 0} \frac{\log r_1(\varepsilon)}{\log \varepsilon}, \quad q_1 = q_1^\nu = \lim_{\varepsilon \rightarrow \infty} \frac{\log r_1(\varepsilon)}{\log \varepsilon},$$

and $p_1 \leq q_1$. It can be shown (see Remark 5) that $p_1 \leq \alpha \leq q_1$. In this paper, we study the Cauchy problem (1.1) in the scale of spaces of generalized Hölder functions whose regularity is determined by the Lévy measure ν . We use w to define generalized Besov norms $|\cdot|_{\beta, \infty}$ and generalized spaces $\tilde{C}_{\infty, \infty}^\beta(H_T)$, $\beta > 0$ (See Section 2.2). They are Besov spaces of generalized smoothness (see [12], [13], [11]) with admissible sequence $w(N^{-j})^{-\beta}$, $j \geq 0$, and covering sequence N^j , $j \geq 0$, with $N > 1$. In particular (see Section 2), for $\beta \in (0, q_1^{-1})$, the norm $|\cdot|_{\beta, \infty}$ for the functions on \mathbf{R}^d is equivalent to

$$\|u\|_\beta = \sup_x |u(x)| + \sup_{x \neq y} \frac{|u(x) - u(y)|}{w(|x - y|)^\beta}.$$

When ν is “close” to an α -stable measure, they reduce to the classical Besov (or equiv. Hölder-Zygmund) spaces.

Let

$$\tilde{\nu}_R(dy) = w(R)\nu(Rdy), R \in [0, 1].$$

The main results of this paper is

Theorem 1. *Let $\beta \in (0, \infty)$, $\lambda \geq 0$. Let $\nu \in \mathfrak{A}^\alpha$, and $w = w_\nu$ be an O-RV function at zero with p_1, q_1 defined in (1.3). Assume*

A.

$$\begin{aligned} 0 &< p_1 \leq q_1 < 1 \text{ if } \alpha \in (0, 1), \quad 1 \leq p_1 \leq q_1 < 2 \text{ if } \alpha = 1, \\ 1 &< p_1 \leq q_1 < 2 \text{ if } \alpha \in (1, 2); \end{aligned}$$

B.

$$\inf_{R \in (0, 1], |\hat{\xi}|=1} \int_{|y| \leq 1} \left| \hat{\xi} \cdot y \right|^2 \tilde{\nu}_R(dy) > 0;$$

C. *There is $N_0 > 2$ so that*

$$\int_1^\infty w(t)^{\frac{1}{q_1}} \frac{dt}{t^{N_0}} < \infty.$$

Then for each $f \in \tilde{C}_{\infty, \infty}^\beta(H_T)$ there is a unique solution $u \in \tilde{C}_{\infty, \infty}^{1+\beta}(H_T)$ solving (1.1). Moreover,

$$(1.4) \quad |u|_{\beta, \infty} \leq C \rho_\lambda(T) |f|_{\beta, \infty},$$

$$(1.5) \quad |u|_{1+\beta, \infty} \leq C [1 + \rho_\lambda(T)] |f|_{\beta, \infty}$$

and

$$\begin{aligned} &|u(t, \cdot) - u(t', \cdot)|_{\mu+\beta, \infty} \\ &\leq C \left\{ (t-t')^{1-\mu} + [1 + \rho_\lambda(T)] |t-t'| \right\} |f|_{\beta, \infty} \end{aligned}$$

for any $\mu \in [0, 1]$ and $t' < t \leq T$, where $\rho_\lambda(T) = (\frac{1}{\lambda} \wedge T)$. The constant C does not depend on f, λ, T, μ .

More specific examples could be the following.

Example 1. *According to [8], Chapter 3, 70-74, any Lévy measure $\nu \in \mathfrak{A}^\alpha$ can be disintegrated as*

$$\nu(\Gamma) = - \int_0^\infty \int_{S_{d-1}} \chi_\Gamma(rw) \Pi(r, dw) d\delta(r), \Gamma \in \mathcal{B}(\mathbf{R}_0^d),$$

where $\delta = \delta_\nu$, and $\Pi(r, dw), r > 0$, is a measurable family of measures on the unit sphere S_{d-1} with $\Pi(r, S_{d-1}) = 1, r > 0$. If δ is an O-RV function, $|\{s \in [0, 1] : r_1(s) < 1\}| > 0$, **A**, **C** and

$$\inf_{|\hat{\xi}|=1} \int_{S_{d-1}} \left| \hat{\xi} \cdot w \right|^2 \Pi(r, dw) \geq c_0 > 0, r > 0,$$

hold, then all assumptions of Theorem 1 are satisfied (see Corollary 6).

Example 2. Consider Lévy measures in radial and angular coordinate system ($r = |y|$, $w = \frac{y}{|y|}$) in the form

$$\nu(B) = \int_0^\infty \int_{|w|=1} 1_B(rw) a(r, w) j(r) r^{d-1} S(dw) dr, B \in \mathcal{B}(\mathbf{R}_0^d),$$

where $S(dw)$ is a finite measure on the unit sphere.

Assume

(i) There is $C > 1, c > 0, 0 < \delta_1 \leq \delta_2 < 1$, such that

$$C^{-1} \phi(r^{-2}) \leq j(r) r^d \leq C \phi(r^{-2})$$

and for all $1 < r \leq R$,

$$c^{-1} \left(\frac{R}{r}\right)^{\delta_1} \leq \frac{\phi(R)}{\phi(r)} \leq c \left(\frac{R}{r}\right)^{\delta_2}.$$

(ii) There is a function $\rho_0(w)$ defined on the unit sphere such that $\rho_0(w) \leq a(r, w) \leq 1, \forall r > 0$, and for all $|\hat{\xi}| = 1$,

$$\int_{S^{d-1}} |\hat{\xi} \cdot w|^2 \rho_0(w) S(dw) \geq c > 0.$$

Under these assumptions, it can be shown that **B** and **C** hold, and δ_v is an O -RV function with $2\delta_1 \leq p_1 \leq q_1 \leq 2\delta_2$. Among the options for ϕ could be (see [15])

- (1) $\phi(r) = \sum_{i=1}^n r^{\alpha_i}, \alpha_i \in (0, 1), i = 1, \dots, n;$
- (2) $\phi(r) = (r + r^\alpha)^\beta, \alpha, \beta \in (0, 1);$
- (3) $\phi(r) = r^\alpha (\ln(1+r))^\beta, \alpha \in (0, 1), \beta \in (0, 1-\alpha);$
- (4) $\phi(r) = (r + m^{1/\alpha})^\alpha - m, \alpha \in (0, 1), m > 0;$
- (5) $\phi(r) = [\ln(\cosh \sqrt{r})]^\alpha, \alpha \in (0, 1).$

Equations in classical Hölder spaces with non-local nondegenerate operators of the form

$$\begin{aligned} \mathcal{L}u(x) &= 1_{\alpha \in (0,2)} \int [u(x+y) - u(x) - 1_{\alpha \geq 1} 1_{|y| \leq 1} y \cdot \nabla u(x)] m(x, y) \nu(dy) \\ &+ 1_{\alpha=2} a^{ij}(x) \partial_{ij}^2 u(x) + 1_{\alpha \geq 1} \tilde{b}^i(x) \partial_i u(x) + l(x) u(x), x \in \mathbf{R}^d, \end{aligned}$$

were considered in many papers. In [1], the existence and uniqueness of a solution to a parabolic equation with \mathcal{L} in Hölder spaces was proved analytically for m Hölder continuous in x and smooth in y , $\nu(dy) = dy/|y|^{d+\alpha}$. The elliptic problem $\mathcal{L}u = f$ in \mathbf{R}^d with $\nu(dy) = dy/|y|^{d+\alpha}$ was considered in [4], [6] and [9]. In [6], the interior Hölder estimates (in a non-linear case as well) were studied assuming that m is symmetric in y . In [4], with $\nu(dy) = dy/|y|^{d+\alpha}$, the a priori estimates were derived in Hölder classes assuming Hölder continuity of m in x , except the case $\alpha = 1$. Similar results, including the case $\alpha = 1$ were proved in [9]. In [7] (see references therein),

in the classical Hölder spaces the case of a nondegenerate

$$\nu(\Gamma) = \int_0^\infty \int_{S_{d-1}} \chi_\Gamma(rw) a(r, w) S(dw) \frac{dr}{r^{1+\alpha}}, \Gamma \in \mathcal{B}(\mathbf{R}_0^d),$$

with a finite measure $S(dw)$ on the unit sphere was considered. Finally, in [17], for (1.1) with x -dependent density $m(x, y)$ at ν , under different assumptions than **A-C**, existence and uniqueness in generalized smoothness classes is derived.

Our paper is organized as follows. In section 2, notation is introduced, the scale of generalized Hölder function spaces is defined, and various equivalent norms are introduced. In particular, using some probabilistic considerations, we prove the equivalence of $|u|_{\beta, \infty}$ to the norms involving fractional powers of nondegenerate L^ν . The continuity of the operator is proved as well. Study of function spaces of generalized smoothness dates back to the seventies-eighties, (see [12], [13] and references therein). Later, this interest continued in connection with the construction problems of Markov processes with jumps (see [11], [10] and references therein). In section 3, we prove the main theorem by starting with smooth input functions. Then we derive the key uniform estimates for the corresponding smooth solutions to (1.1). We handle generalized Hölder inputs by passing to the limit. Finally, Appendix contains all needed results about O-RV functions. The regular variation functions were introduced in [14] and used in tauberian theorems which were extended to O-RV functions as well (see [2], [3], and references therein). They are very convenient for the derivation of our main estimates.

2. NOTATION, FUNCTION SPACES AND NORM EQUIVALENCE

2.1. Basic notation. We denote $\mathbf{N} = \{0, 1, 2, 3, \dots\}$, $\mathbf{N}_+ = \mathbf{N} \setminus \{0\}$; $H_T = [0, T] \times \mathbf{R}^d$; S^{d-1} is the unit sphere in \mathbf{R}^d . For $B \in \mathcal{B}(\mathbf{R}^d)$, we denote by $|B|$ the Lebesgue measure of B .

For a function u on H_T , we denote its partial derivatives by $\partial_t u = \partial u / \partial t$, $\partial_i u = \partial u / \partial x_i$, $\partial_{ij}^2 u = \partial^2 u / \partial x_i \partial x_j$, and denote its gradient with respect to x by $\nabla u = (\partial_1 u, \dots, \partial_d u)$ and $D^{|\gamma|} u = \partial^{|\gamma|} u / \partial x_1^{\gamma_1} \dots \partial x_d^{\gamma_d}$, where $\gamma = (\gamma_1, \dots, \gamma_d) \in \mathbf{N}^d$ is a multi-index.

We use $C_b^\infty(\mathbf{R}^d)$ to denote the set of bounded infinitely differentiable functions on \mathbf{R}^d whose derivative of arbitrary order is bounded, and $C^k(\mathbf{R}^d)$, $k \in \mathbf{N}$ the class of k -times continuously differentiable functions.

We denote $\mathcal{S}(\mathbf{R}^d)$ the Schwartz space of rapidly decreasing functions on \mathbf{R}^d , and $\mathcal{S}'(\mathbf{R}^d)$ denotes the space of continuous functionals on $\mathcal{S}(\mathbf{R}^d)$, i.e. the space of tempered distributions.

We adopt the normalized definition for Fourier and its inverse transforms for functions in $\mathcal{S}(\mathbf{R}^d)$, i.e.,

$$\begin{aligned}\mathcal{F}\varphi(\xi) &= \hat{\varphi}(\xi) := \int e^{-i2\pi x \cdot \xi} \varphi(x) dx, \\ \mathcal{F}^{-1}\varphi(x) &= \check{\varphi}(x) := \int e^{i2\pi x \cdot \xi} \varphi(\xi) d\xi, \quad \varphi \in \mathcal{S}(\mathbf{R}^d).\end{aligned}$$

Recall that Fourier transform can be extended to a bijection on $\mathcal{S}'(\mathbf{R}^d)$.

Throughout the sequel, Z_t^ν represents the Lévy process associated to the Lévy measure $\nu \in \mathfrak{A}^\alpha$, see (1.2).

For any Lévy measure $\nu \in \mathfrak{A}^\alpha$ and $R > 0$,

$$\nu_R(B) := \int 1_B(y/R) \nu(dy), \quad B \in \mathcal{B}(\mathbf{R}_0^d), \quad \tilde{\nu}_R(dy) := w(R) \nu_R(dy).$$

For any Lévy measure $\nu \in \mathfrak{A}^\alpha$, we denote its symmetrization

$$\nu_{sym}(dy) = \frac{1}{2} [\nu(dy) + \nu(-dy)].$$

And $\mathfrak{A}_{sym}^\alpha = \{\mu \in \mathfrak{A}^\alpha : \mu = \mu_{sym}\}$.

If $A(\mathbf{R}^d)$ is a space of functions v on \mathbf{R}^d with norm $|v|_A = |v|_{A(\mathbf{R}^d)}$, then $A(H_T)$ denotes the spaces of functions u on $H_T = [0, T] \times \mathbf{R}^d$ with finite norm

$$|u|_A = |u|_{A(H_T)} = \sup_{t \in [0, T]} |u(t, \cdot)|_{A(\mathbf{R}^d)}.$$

We have specific values assigned for c_0, c_1, c_2, N_0, N_1 , but we allow C to vary from line to line. In particular, $C(\dots)$ represents a constant depending only on quantities in the parentheses.

2.2. Function spaces and norm equivalence. We fix a constant $N > 1$. For such an N , by Lemma 6.1.7 in [5] and appropriate scaling, there exists $\phi \in C_0^\infty(\mathbf{R}^d)$ such that $\text{supp}(\phi) = \{\xi : \frac{1}{N} \leq |\xi| \leq N\}$, $\phi(\xi) > 0$ in the interior of its support, and

$$\sum_{j=-\infty}^{\infty} \phi(N^{-j}\xi) = 1 \text{ if } \xi \neq 0.$$

We denote throughout this paper

$$(2.2) \quad \varphi_j = \mathcal{F}^{-1}[\phi(N^{-j}\xi)], \quad j = 1, 2, \dots, \xi \in \mathbf{R}^d,$$

$$(2.3) \quad \varphi_0 = \mathcal{F}^{-1}\left[1 - \sum_{j=1}^{\infty} \phi(N^{-j}\xi)\right].$$

Apparently, $\varphi_j \in \mathcal{S}(\mathbf{R}^d)$, $j \in \mathbf{N}$. They are convolution functions we use to define *generalized Besov spaces*. Namely, for $\beta > 0$ we write $\tilde{C}_{\infty, \infty}^\beta(\mathbf{R}^d)$ as

the set of functions in $\mathcal{S}'(\mathbf{R}^d)$ for which

$$(2.4) \quad |u|_{\beta, \infty} := \sup_j w(N^{-j})^{-\beta} |u * \varphi_j|_0 < \infty,$$

where $w = w_\nu$ with $\nu \in \mathfrak{A}^\alpha$.

Lemma 1. *Let $\nu \in \mathfrak{A}^\alpha$, $w = w_\nu$ be an O-RV function at zero and **A** holds for it. Let $\beta \in (0, \infty)$. If $u \in \tilde{C}_{\infty, \infty}^\beta(\mathbf{R}^d)$, then u is bounded and continuous,*

$$u(x) = \sum_{j=0}^{\infty} (u * \varphi_j)(x), x \in \mathbf{R}^d,$$

where the series converges uniformly. Moreover,

$$|u|_0 \leq C |u|_{\beta, \infty}, u \in \tilde{C}_{\infty, \infty}^\beta(\mathbf{R}^d).$$

Proof. Note that $u * \varphi_j$ is continuous of moderate growth and $\sum_{j=0}^{\infty} u * \varphi_j = u$ in $\mathcal{S}'(\mathbf{R}^d)$. Obviously, by Corollary 5 in Appendix,

$$\begin{aligned} \sum_{j=0}^{\infty} |u * \varphi_j|_0 &= \sum_{j=0}^{\infty} w(N^{-j})^\beta w(N^{-j})^{-\beta} |u * \varphi_j|_0 \\ &\leq \sup_{j \geq 0} w(N^{-j})^{-\beta} |u * \varphi_j|_0 \sum_{j=0}^{\infty} w(N^{-j})^\beta \\ &\leq C |u|_{\beta, \infty} \sum_{j=0}^{\infty} w(N^{-j})^\beta \leq C |u|_{\beta, \infty}. \end{aligned}$$

□

Let $\nu \in \mathfrak{A}^\alpha$, $w = w_\nu$, $\beta > 0$. For $u \in C_b^\infty(\mathbf{R}^d)$, set

$$|u|_0 = \sup_x |u(x)|, [u]_\beta = \sup_{x, h \neq 0} \frac{|u(x+h) - u(x)|}{w(|h|)^\beta},$$

and

$$|u|_\beta := |u|_0 + [u]_\beta.$$

Proposition 1. *Let $\nu \in \mathfrak{A}^\alpha$, $w = w_\nu$ be an O-RV function at zero so that **A** and **C** hold for it. Let $\beta \in (0, q_1^{-1})$. Then the norm $|u|_\beta$ and norm $|u|_{\beta, \infty}$ are equivalent on $C_b^\infty(\mathbf{R}^d)$. Namely, there is $C > 0$ depending only on d, β, N such that*

$$C^{-1} |u|_\beta \leq |u|_{\beta, \infty} \leq C |u|_\beta, u \in C_b^\infty(\mathbf{R}^d).$$

Proof. Let $u \in C_b^\infty(\mathbf{R}^d)$. Then, by Lemma 8, $|u|_\beta < \infty$. If $j = 0$, then

$$|u * \varphi_0|_0 \leq |u|_0 \int |\varphi_0(y)| dy \leq C |u|_\beta.$$

If $j \neq 0$, then by the construction of φ_j , $\int \varphi_j(y) dy = \hat{\varphi}_j(0) = 0$. Therefore, denoting $\varphi = \mathcal{F}^{-1}\phi$,

$$\begin{aligned} & |u * \varphi_j|_0 \\ &= \left| \int [u(y) - u(x)] \varphi_j(x-y) dy \right|_0 \\ &\leq [u]_\beta \int w(|y-x|)^\beta N^{jd} |\varphi(N^j(x-y))| dy \\ &= [u]_\beta \int w(N^{-j}|y|)^\beta |\varphi(y)| dy. \end{aligned}$$

Since for $N_0 > d+1$,

$$|\varphi(y)| \leq C(1+|y|)^{-N_0}, y \in \mathbf{R}^d,$$

for some $C > 0$, we have

$$\begin{aligned} & \int w(N^{-j}|y|)^\beta |\varphi(y)| dy \\ & \leq C \int_0^1 w(N^{-j}|y|)^\beta dy + C \int_1^\infty w(N^{-j}|y|)^\beta |y|^{-N_0} dy = A_1 + A_2. \end{aligned}$$

By Lemma 8

$$N^{-j(N_0-d)} \leq Cw(N^{-j})^\beta, j \geq 0,$$

and,

$$\begin{aligned} A_1 & \leq CN^{jd} \int_0^{N^{-j}} w(r)^\beta r^d \frac{dr}{r} \leq Cw(N^{-j})^\beta, \\ A_2 & = CN^{-j(N_0-d)} \int_{N^{-j}}^\infty w(r)^\beta r^{-(N_0-d)} \frac{dr}{r} \\ & = CN^{-j(N_0-d)} \int_{N^{-j}}^1 w(r)^\beta r^{-(N_0-d)} \frac{dr}{r} + N^{-j(N_0-d)} \int_1^\infty w(r)^\beta r^{-(N_0-d)} \frac{dr}{r} \\ & \leq Cw(N^{-j})^\beta, j \geq 0. \end{aligned}$$

That is to say $|u|_{\beta, \infty} \leq C|u|_\beta, u \in C_b(\mathbf{R}^d)$ for some constant $C(\beta, d) > 0$.

Let $\tilde{\phi}, \tilde{\phi}_0 \in C_0^\infty(\mathbf{R}^d)$, be such that $0 \notin \text{supp}(\tilde{\phi})$, $\tilde{\phi}\phi = \phi, \tilde{\phi}_0\phi_0 = \phi_0$, where $\phi_0 = \mathcal{F}\varphi_0$, and ϕ, φ_0 are the functions introduced in (2.3), (2.2). Let

$$(2.5) \quad \tilde{\varphi} = \mathcal{F}^{-1}\tilde{\phi}, \quad \tilde{\varphi}_j = \mathcal{F}^{-1}\tilde{\phi}(N^{-j}\cdot), j \geq 1,$$

$$(2.6) \quad \tilde{\varphi}_0 = \mathcal{F}^{-1}\tilde{\phi}_0.$$

Hence

$$\varphi_j = \varphi_j * \tilde{\varphi}_j, j \geq 0,$$

where in particular,

$$\tilde{\varphi}_j(x) = N^{jd}\tilde{\varphi}(N^jx), j \geq 1, x \in \mathbf{R}^d.$$

Obviously,

$$\begin{aligned} & |u * \varphi_0(x+y) - u * \varphi_0(x)| \\ & \leq \int |\tilde{\varphi}_0(x+y-z) - \tilde{\varphi}_0(x-z)| |u * \varphi_0(z)| dz \\ & \leq C(|y| \wedge 1) |u * \varphi_0|_0, \quad x, y \in \mathbf{R}^d, \end{aligned}$$

and

$$\begin{aligned} & |u * \varphi_j(x+y) - u * \varphi_j(x)| \\ & \leq N^{jd} \int |\tilde{\varphi}(N^j(x+y-z)) - \tilde{\varphi}(N^j(x-z))| |u * \varphi_j(z)| dz \\ & \leq C(|N^j y| \wedge 1) |u * \varphi_j|_0, \quad j \geq 1, \quad x, y \in \mathbf{R}^d. \end{aligned}$$

By Lemma 1, for $x, y \in \mathbf{R}^d$,

$$\begin{aligned} & |u(x+y) - u(x)| \\ & \leq \sum_{j=0}^{\infty} |u * \varphi_j(x+y) - u * \varphi_j(x)| \leq C \sum_{j=0}^{\infty} (N^j |y| \wedge 1) |u * \varphi_j|_0. \end{aligned}$$

Let $\beta q_1 < 1, k \in \mathbf{N}$. For $|y| \in (N^{-k-1}, N^{-k}]$,

$$\begin{aligned} |u(x+y) - u(x)| & \leq C |u|_{\beta, \infty} \sup_{|y| \leq N^{-k}} \sum_{j=0}^{\infty} (N^j |y| \wedge 1) w(N^{-j})^\beta \\ & \leq C |u|_{\beta, \infty} \left[\sum_{j=0}^k N^{j-k} w(N^{-j})^\beta + \sum_{j=k+1}^{\infty} w(N^{-j})^\beta \right]. \end{aligned}$$

Then, by Lemma 8,

$$\begin{aligned} N^{-k} \sum_{j=0}^k N^j w(N^{-j})^\beta & \leq C_2 N^{-k} \int_0^{k+1} N^x w(N^{-x})^\beta dx \\ & \leq C N^{-k} \int_{N^{-k-1}}^1 x^{-1} w(x)^\beta \frac{dx}{x} \leq C w(|y|)^\beta. \end{aligned}$$

Again, by Lemma 8,

$$\sum_{j=k+1}^{\infty} w(N^{-j})^\beta \leq C \int_{k+1}^{\infty} w(N^{-x})^\beta dx \leq C \int_0^{N^{-k-1}} w(x)^\beta \frac{dx}{x} \leq C w(|y|)^\beta.$$

The statement is proved. \square

2.2.1. *Equivalent norms on $C_b^\infty(\mathbf{R}^d)$.* Now we will introduce some other norms on $C_b^\infty(\mathbf{R}^d)$ involving the powers of the operators $L^\nu, I - L^\nu$:

$$\begin{aligned} |u|_{\nu, \kappa, \beta} & = |u|_{\kappa, \beta} = |u|_0 + |L^{\nu; \kappa} u|_{\beta, \infty}, \quad u \in C_b^\infty(\mathbf{R}^d), \\ \|u\|_{\nu, \kappa, \beta} & = \|u\|_{\kappa, \beta} = |(I - L^\nu)^\kappa u|_{\beta, \infty}, \quad u \in C_b^\infty(\mathbf{R}^d), \end{aligned}$$

with $\kappa, \beta > 0$, $L^{\nu;\kappa} = (L^\nu)^\kappa$, and ν satisfying **A** and **B**. In addition, we assume that $\nu \in \mathfrak{A}_{sym}^\alpha = \{\mu \in \mathfrak{A}^\alpha : \mu = \mu_{sym}\}$ if κ is fractional. First, we define those powers and corresponding norms on $C_b^\infty(\mathbf{R}^d)$. Then we study their relations and extend them to $\tilde{C}_{\infty\infty}^{\kappa+\beta}(\mathbf{R}^d)$.

For $\nu \in \mathfrak{A}_{sym}^\alpha$, $\kappa \in (0, 1)$, $a \geq 0$, and $f \in \mathcal{S}(\mathbf{R}^d)$, we see easily that

$$\begin{aligned} & (a - \psi^\nu(\xi))^\kappa \hat{f}(\xi) \\ &= c_\kappa \int_0^\infty t^{-\kappa} [e^{-at} \exp(\psi^\nu(\xi)t) - 1] \frac{dt}{t} \hat{f}(\xi), \xi \in \mathbf{R}^d, \end{aligned}$$

and define

$$\begin{aligned} (2.7) \quad & (aI - L^\nu)^\kappa f(x) \\ &= \mathcal{F}^{-1} \left[(a - \psi^\nu)^\kappa \hat{f} \right] (x) \\ &= c_\kappa \mathbf{E} \int_0^\infty t^{-\kappa} [e^{-at} f(x + Z_t^\nu) - f(x)] \frac{dt}{t}, x \in \mathbf{R}^d, \end{aligned}$$

where

$$c_\kappa = \left(\int_0^\infty (e^{-t} - 1) t^{-\kappa} \frac{dt}{t} \right)^{-1}.$$

We denote, with $a = 0$, $f \in \mathcal{S}(\mathbf{R}^d)$, $\kappa \in (0, 1)$,

$$L^{\nu;\kappa} f := \mathcal{F}^{-1} \left[-(-\psi^\nu)^\kappa \hat{f} \right].$$

For $f \in C_b^\infty(\mathbf{R}^d)$, $\kappa \in (0, 1)$, $a \geq 0$, we define

$$(aI - L^\nu)^\kappa f(x) = c_\kappa \mathbf{E} \int_0^\infty t^{-\kappa} [e^{-at} f(x + Z_t^\nu) - f(x)] \frac{dt}{t}, x \in \mathbf{R}^d.$$

For $\kappa = 1$, $(aI - L^\nu)^1 f = (aI - L^\nu) f = af - L^\nu f$, $f \in C_b^\infty(\mathbf{R}^d)$. Note that for $\kappa \in (0, 1)$, $a \geq 0$,

$$\begin{aligned} (2.8) \quad & (aI - L^\nu)^\kappa f(x) \\ &= c_\kappa \mathbf{E} \int_1^\infty t^{-\kappa} [e^{-at} f(x + Z_t^\nu) - f(x)] \frac{dt}{t} \\ & \quad + c_\kappa \mathbf{E} \int_0^1 t^{-\kappa} \int_0^t e^{-as} (-a + L^\nu) f(x + Z_s^\nu) ds \frac{dt}{t}, x \in \mathbf{R}^d. \end{aligned}$$

For $a > 0$, $f \in C_b^\infty(\mathbf{R}^d)$, set

$$(aI - L^\nu)^{-\kappa} f(x) = c'_\kappa \int_0^\infty t^\kappa e^{-at} \mathbf{E} f(x + Z_t^\nu) \frac{dt}{t}, x \in \mathbf{R}^d,$$

where

$$c'_\kappa = \left(\int_0^\infty t^\kappa e^{-t} \frac{dt}{t} \right)^{-1},$$

and $\nu \in \mathfrak{A}_{sym}^\alpha$, $\kappa > 0$, or $\nu \in \mathfrak{A}^\alpha$, $\kappa \in \mathbf{N}$.

Note that for $g \in \mathcal{S}(\mathbf{R}^d)$,

$$\begin{aligned}\mathcal{F}[(aI - L^\nu)^{-\kappa} g] &= (a - \psi^\nu)^{-\kappa} \hat{g}, a > 0, \kappa > 0, \\ \mathcal{F}[(aI - L^\nu)^\kappa g] &= (a - \psi^\nu)^\kappa \hat{g}, a \geq 0, \kappa \in (0, 1].\end{aligned}$$

We use the formulas above to define $(a - L^\nu)^\kappa, a \geq 0, \kappa = 1, 0, -1, \dots$, for $\nu \in \mathfrak{A}^\alpha$.

Remark 1. Assume $\kappa \in (0, 1], a \geq 0$ or $\kappa \in (-\infty, 0), a > 0$. It is easy to see that

a) for any $f \in C_b^\infty(\mathbf{R}^d)$, we have $(aI - L^\nu)^\kappa f \in C_b^\infty(\mathbf{R}^d)$ and for any multiindex γ , $D^\gamma (aI - L^\nu)^\kappa f = (aI - L^\nu)^\kappa D^\gamma f, \nu \in \mathfrak{A}_{sym}^\alpha$. The same holds for $\nu \in \mathfrak{A}^\alpha$ and $\kappa = 1, 0, -1, \dots$.

b) for any $f \in C_b^\infty(\mathbf{R}^d)$ such that for any multiindex γ , $D^\gamma f \in L^1(\mathbf{R}^d) \cap L^2(\mathbf{R}^d)$, we have

$$\begin{aligned}\mathcal{F}[(aI - L^\nu)^{-\kappa} f] &= (a - \psi^\nu)^{-\kappa} \hat{f}, a > 0, \kappa > 0, \\ \mathcal{F}[(aI - L^\nu)^\kappa f] &= (a - \psi^\nu)^\kappa \hat{f}, a \geq 0, \kappa \in (0, 1],\end{aligned}$$

for $\nu \in \mathfrak{A}_{sym}^\alpha$. The same holds for $\nu \in \mathfrak{A}^\alpha$ and $\kappa = 1, 0, -1, \dots$.

The following obvious claim holds.

Lemma 2. Let $\nu \in \mathfrak{A}_{sym}^\alpha$. Assume $\kappa \in (0, 1], a \geq 0$ or $\kappa \in (-\infty, 0), a > 0$. Let $f, f_n \in C_b^\infty(\mathbf{R}^d)$ be so that for any multiindex γ , $D^\gamma f_n \rightarrow D^\gamma f$ as $n \rightarrow \infty$ uniformly on compact subsets of \mathbf{R}^d and

$$\sup_{x,n} |D^\gamma f_n(x)| < \infty.$$

Then for any multiindex γ ,

$D^\gamma (aI - L^\nu)^\kappa f_n = (aI - L^\nu)^\kappa D^\gamma f_n \rightarrow D^\gamma (aI - L^\nu)^\kappa f = (aI - L^\nu)^\kappa D^\gamma f$ uniformly on compact subsets of \mathbf{R}^d , and

$$\sup_{x,n} |(aI - L^\nu)^\kappa D^\gamma f_n(x)| < \infty.$$

The same holds for $\nu \in \mathfrak{A}^\alpha$ and $\kappa = 1, 0, -1, \dots$.

Remark 2. Given $f \in C_b^\infty(\mathbf{R}^d)$ there is a sequence $f_n \in C_0^\infty(\mathbf{R}^d)$ so that for any multiindex γ , $D^\gamma f_n \rightarrow D^\gamma f$ as $n \rightarrow \infty$ uniformly on compact subsets of \mathbf{R}^d and

$$\sup_{x,n} |D^\gamma f_n(x)| < \infty.$$

Indeed, choose $g \in C_b^\infty(\mathbf{R}^d), 0 \leq g \leq 1, g(x) = 1$ if $|x| \leq 1$, and $g(x) = 0$ if $|x| > 2$. Given $f \in C_b^\infty(\mathbf{R}^d)$, take

$$f_n(x) = f(x) g(x/n), x \in \mathbf{R}^d, n \geq 1.$$

Lemma 3. *Let $\nu \in \mathfrak{A}_{sym}^\alpha$. Assume $a > 0, \kappa \in (0, 1]$. Then $(aI - L^\nu)^\kappa : C_b^\infty(\mathbf{R}^d) \rightarrow C_b^\infty(\mathbf{R}^d)$ is bijective whose inverse is $(aI - L^\nu)^{-\kappa}$:*

$$(aI - L^\nu)^\kappa (aI - L^\nu)^{-\kappa} f(x) = (aI - L^\nu)^{-\kappa} (aI - L^\nu)^\kappa f(x) = f(x), x \in \mathbf{R}^d,$$

for any $f \in C_b^\infty(\mathbf{R}^d)$.

Proof. It is an easy consequence of Lemma 2 and Remarks 1 and 2. \square

For an integer $k \in \mathbf{N}$, we define for $\nu \in \mathfrak{A}^\alpha$,

$$(aI - L^\nu)^k = \underbrace{(aI - L^\nu) \dots (aI - L^\nu)}_{k \text{ times}}.$$

For a non integer $\kappa > 0, \kappa = [\kappa] + s$ with $s \in (0, 1)$ and $\nu \in \mathfrak{A}_{sym}^\alpha$, we set

$$\begin{aligned} (aI - L^\nu)^\kappa f &= (aI - L^\nu)^{[\kappa]} (aI - L^\nu)^s f \\ &= (aI - L^\nu)^s (aI - L^\nu)^{[\kappa]} f, f \in C_b^\infty(\mathbf{R}^d). \end{aligned}$$

Remark 3. *Let $\nu \in \mathfrak{A}_{sym}^\alpha$, and $f \in C_b^\infty(\mathbf{R}^d)$ be such that for any multiindex γ , $D^\gamma f \in L^1(\mathbf{R}^d) \cap L^2(\mathbf{R}^d)$. Then*

$$\begin{aligned} \mathcal{F}[(aI - L^\nu)^{-s} f] &= (a - \psi^\nu)^{-s} \hat{f}, a > 0, s > 0, \\ \mathcal{F}[(aI - L^\nu)^s f] &= (a - \psi^\nu)^s \hat{f}, a \geq 0, s > 0. \end{aligned}$$

The same holds with $\nu \in \mathfrak{A}^\alpha$ if $s \in \mathbf{N}$.

Lemma 4. *Assume $a > 0$. Then*

(i) *for any $\kappa, s \geq 0$, we have $L^{\nu; \kappa} L^{\nu; s} = L^{\nu; \kappa+s}$; for any $\kappa, s \in \mathbf{R}$,*

$$\begin{aligned} (aI - L^\nu)^\kappa (aI - L^\nu)^s &= (aI - L^\nu)^{\kappa+s}, \\ (aI - L^\nu)^{-\kappa} (aI - L^\nu)^{-s} &= (aI - L^\nu)^{-(\kappa+s)}, \end{aligned}$$

for $\nu \in \mathfrak{A}_{sym}^\alpha$.

The same holds with $\nu \in \mathfrak{A}^\alpha$ if $\kappa, s \in \mathbf{N}$.

(ii) *for any $\kappa > 0$, the mapping $(aI - L^\nu)^\kappa : C_b^\infty(\mathbf{R}^d) \rightarrow C_b^\infty(\mathbf{R}^d)$ is bijective whose inverse is $(aI - L^\nu)^{-\kappa}$:*

$$(aI - L^\nu)^\kappa (aI - L^\nu)^{-\kappa} f(x) = (aI - L^\nu)^{-\kappa} (aI - L^\nu)^\kappa f(x) = f(x), x \in \mathbf{R}^d.$$

for any $f \in C_b^\infty(\mathbf{R}^d)$.

The same holds with $\nu \in \mathfrak{A}^\alpha$ if $\kappa \in \mathbf{N}$.

Proof. The statement is an easy consequence of Lemma 2 and Remarks 1, 2, and 3. \square

Lemma 5. *Let $\nu \in \mathfrak{A}_{sym}^\alpha$ satisfy **A**.*

(i) *Let $a \geq 0, \kappa > 0, m = [\kappa] + 1$. For any $f \in C_b^\infty(\mathbf{R}^d)$,*

$$\sup_{R \in (0, 1], x} \left| (a - L^{\tilde{\nu}_R})^\kappa D^\gamma f(x) \right| \leq C(1+a)^\kappa \max_{|\mu| \leq |\gamma| + 2m} |D^\mu f|_0 < \infty.$$

If, in addition, for any multiindex γ , $\int |D^\gamma f(x)| dx < \infty$, then

$$\sup_{R \in (0,1], x} \int \left| (a - L^{\tilde{\nu}_R})^\kappa D^\gamma f(x) \right| dx \leq C(1+a)^\kappa \max_{|\mu| \leq |\gamma| + 2m} \int |D^\mu f(x)| dx.$$

The same holds with $\nu \in \mathfrak{A}^\alpha$ satisfying **A** if $\kappa \in \mathbf{N}$.

(ii) Let $a > 0, \kappa > 0$. For any $f \in C_b^\infty(\mathbf{R}^d)$,

$$\sup_{R \in (0,1], x} \left| (a - L^{\tilde{\nu}_R})^{-\kappa} D^\gamma f(x) \right| \leq Ca^{-\kappa} \max_{|\mu| \leq |\gamma|} |D^\mu f|_0 < \infty.$$

If in addition, for any multiindex γ , $\int |D^\gamma f(x)| dx < \infty$, then

$$\sup_{R \in (0,1]} \int \left| (a - L^{\tilde{\nu}_R})^{-\kappa} D^\gamma f(x) \right| dx \leq Ca^{-\kappa} \max_{|\mu| \leq |\gamma|} \int |D^\mu f(x)| dx.$$

The same holds with $\nu \in \mathfrak{A}^\alpha$ satisfying **A** if $\kappa \in \mathbf{N}$.

Proof. (i) Let $\kappa \in (0, 1]$. Then

$$\begin{aligned} & (a - L^{\tilde{\nu}_R})^\kappa f(x) \\ &= c_\kappa \int_0^\infty t^{-\kappa} \mathbf{E} \left[e^{-at} f(x + Z_t^{\tilde{\nu}_R}) - f(x) \right] \frac{dt}{t} = c_\kappa \int_1^\infty \dots \\ & \quad + c_\kappa \int_0^1 t^{-\kappa} \int_0^t e^{-as} \mathbf{E} \left[(L^{\tilde{\nu}_R} - a) f(x + Z_s^{\tilde{\nu}_R}) \right] ds \frac{dt}{t}, \quad x \in \mathbf{R}^d. \end{aligned}$$

By Lemma 9, we have

$$\begin{aligned} \sup_{R \in (0,1]} \int (|y| \wedge 1) \tilde{\nu}_R(dy) &< \infty \text{ if } \alpha \in (0, 1), \\ \sup_{R \in (0,1]} \int (|y|^2 \wedge 1) \tilde{\nu}_R(dy) &< \infty \text{ if } \alpha = 1, \\ \sup_{R \in (0,1]} \int (|y|^2 \wedge |y|) \tilde{\nu}_R(dy) &< \infty \text{ if } \alpha \in (1, 2), \end{aligned}$$

and both inequalities easily follow. Applying them repeatedly we obtain the claim for an arbitrary $\kappa > 0$.

(ii) Indeed, for any $\kappa > 0, a > 0$, and any multiindex γ ,

$$D^\gamma (a - L^{\tilde{\nu}_R})^{-\kappa} f(x) = c_\kappa \int_0^\infty e^{-at} t^\kappa \mathbf{E} D^\gamma f(x + Z_t^{\tilde{\nu}_R}) \frac{dt}{t}, x \in \mathbf{R}^d,$$

and the claim obviously follows. \square

Lemma 6. Let $\nu \in \mathfrak{A}^\alpha$ satisfy **A** and **B**. Let $g \in \mathcal{S}(\mathbf{R}^d)$ be such that $\hat{g} \in C_0^\infty(\mathbf{R}^d)$, $0 \notin \text{supp}(\hat{g})$. Then there are constants C, c so that

$$\sup_{R \in (0,1]} \int \left| \mathbf{E} g(x + Z_t^{\tilde{\nu}_R}) \right| dx \leq Ce^{-ct}, t > 0.$$

Proof. Let $F(t, x) = \mathbf{E}g\left(x + Z_t^{\tilde{\nu}_R}\right)$, $x \in \mathbf{R}^d$, $t > 0$. We choose $\varepsilon > 0$ so that $\text{supp}(\hat{g}) \subseteq \{\xi : |\xi| \leq \varepsilon^{-1}\}$. Let $\tilde{\nu}_{R,\varepsilon}(dy) = \chi_{\{|y| \leq \varepsilon\}} \tilde{\nu}_R(dy)$, $R \in [0, 1]$. Then for $\xi \in \text{supp}(\hat{g})$ and $|y| \leq \varepsilon$,

$$1 - \cos(\xi \cdot y) \geq \frac{1}{\pi} |\xi \cdot y|^2 = \frac{|\xi|^2}{\pi} |\hat{\xi} \cdot y|^2$$

with $\hat{\xi} = \xi/|\xi|$. Therefore there is $c_0 > 0$ so that for any $\xi \in \text{supp}(\hat{g})$ and $R \in (0, 1]$,

$$(2.9) \quad \begin{aligned} -\text{Re} \psi^{\tilde{\nu}_{R,\varepsilon}}(\xi) &= \int_{|y| \leq \varepsilon} [1 - \cos(\xi \cdot y)] \tilde{\nu}_R(dy) \\ &\geq \frac{|\xi|^2}{\pi} \int_{|y| \leq \varepsilon} |\hat{\xi} \cdot y|^2 \tilde{\nu}_R(dy) \geq c_0 |\xi|^2. \end{aligned}$$

Then

$\hat{F}(t, \xi) = \exp\{\psi^{\tilde{\nu}_R}(\xi)t\} \hat{g}(\xi) = \exp\{\psi^{\tilde{\nu}_{R,\varepsilon}}(\xi)t\} \exp\{\psi(\xi)t\} \hat{g}(\xi)$, $\xi \in \mathbf{R}^d$, where $\exp\{\psi(\xi)t\}$ is a characteristic function of a probability distribution $P_{R,t}(dy)$ on \mathbf{R}^d . Hence

$$F(t, x) = \int H(t, x - y) P_{R,t}(dy), \quad x \in \mathbf{R}^d,$$

with

$$H(t, x) = \mathcal{F}^{-1}[\exp\{\psi^{\tilde{\nu}_{R,\varepsilon}}t\} \hat{g}] = \mathbf{E}g\left(x + Z_t^{\tilde{\nu}_{R,\varepsilon}}\right), \quad x \in \mathbf{R}^d.$$

Since

$$\int |F(t, x)| dx \leq \int |H(t, x)| dx,$$

it is enough to prove that

$$(2.10) \quad \int |H(t, x)| dx \leq C e^{-ct}, \quad t > 0.$$

Now, (2.9) implies that for any multiindex $|\gamma| \leq n = [\frac{d}{2}] + 3$,

$$\begin{aligned} \int |x^\gamma H(t, x)|^2 dx &\leq C \int |D^\gamma [\hat{g}(\xi) \exp\{\psi^{\tilde{\nu}_{R,\varepsilon}}(\xi)t\}]|^2 d\xi \\ &\leq C_1 e^{-c_2 t}, \quad t > 0. \end{aligned}$$

Hence, denoting $d_0 = [\frac{d}{2}] + 1$,

$$\begin{aligned} \int |H(t, x)| dx &= \int (1 + |x|^2)^{-d_0} |H(t, x)| (1 + |x|^2)^{d_0} dx \\ &\leq C \int |H(t, x)|^2 (1 + |x|^2)^{2d_0} dx \leq C_1 e^{-c_2 t}, \quad t > 0. \end{aligned}$$

Thus (2.10) follows, and

$$(2.11) \quad \int \left| \mathbf{E}g\left(x + Z_t^{\tilde{\nu}_R}\right) \right| dx \leq C_1 e^{-c_2 t}, \quad t > 0.$$

□

Corollary 1. *Let $\nu \in \mathfrak{A}_{sym}^\alpha$ satisfy **A** and **B**. Let $g \in \mathcal{S}(\mathbf{R}^d)$ be such that $\hat{g} \in C_0^\infty(\mathbf{R}^d)$, $0 \notin \text{supp}(\hat{g})$. Then for $a \geq 0, \kappa > 0, R \in (0, 1]$,*

$$\begin{aligned} (a - L^{\tilde{\nu}_R})^{-\kappa} g(x) &= -\mathcal{F}^{-1} \left[(a - \psi^{\tilde{\nu}_R})^{-\kappa} \hat{g} \right] (x) \\ &= c_{-\kappa} \int_0^\infty e^{-at} t^\kappa \mathbf{E}g \left(x + Z_t^{\tilde{\nu}_R} \right) \frac{dt}{t}, x \in \mathbf{R}^d, \end{aligned}$$

is C_b^∞ -function, and for every multiindex γ , we have $D^\gamma (a - L^{\tilde{\nu}_R})^{-\kappa} g = (a - L^{\tilde{\nu}_R})^{-\kappa} D^\gamma g, R > 0$, and

$$\sup_{R \in (0, 1], a \geq 0} \int \left| D^\gamma (a - L^{\tilde{\nu}_R})^{-\kappa} g(x) \right|^p dx < \infty, p \geq 1.$$

The same holds with $\nu \in \mathfrak{A}^\alpha$ satisfying **A** and **B** if $\kappa \in \mathbf{N}$.

Proof. Take $\eta \in C_0^\infty(\mathbf{R}^d)$ so that $\eta \hat{g} = \hat{g}, 0 \notin \text{supp}(\eta)$, and let $\tilde{\eta} = \mathcal{F}^{-1}\eta$. Let

$$F_R(t, x) = \mathbf{E}g \left(x + Z_t^{\tilde{\nu}_R} \right), t > 0, x \in \mathbf{R}^d.$$

Then

$$\hat{F}_R(t, \xi) = \exp \{ \psi^{\tilde{\nu}_R}(\xi) t \} \eta(\xi) \hat{g}(\xi), \xi \in \mathbf{R}^d,$$

and

$$F_R(t, x) = \int H_R(t, x - y) g(y) dy = \int g(x - y) H_R(t, y) dy, x \in \mathbf{R}^d,$$

with

$$H_R(t, x) = \mathcal{F}^{-1} \left[\exp \{ \psi^{\tilde{\nu}_R} t \} \eta \right] = \mathbf{E} \tilde{\eta} \left(x + Z_t^{\tilde{\nu}_R} \right), t > 0, x \in \mathbf{R}^d.$$

By Lemma 6,

$$\sup_{R \in (0, 1]} \int |H_R(t, y)| dy \leq C e^{-ct}, t > 0.$$

Hence $F_R(t, \cdot) \in C_b^\infty(\mathbf{R}^d), t > 0$, and for each multiindex γ and $p \geq 1$,

$$\sup_R \left(\int |D^\gamma F_R(t, x)|^p dx \right)^{1/p} \leq C \left(\int |D^\gamma g(x)|^p dx \right)^{1/p} e^{-ct}, t > 0.$$

□

Corollary 2. *Let $\nu \in \mathfrak{A}^\alpha$ satisfy **A** and **B**. Let $g \in \mathcal{S}(\mathbf{R}^d)$ be such that $\hat{g} \in C_0^\infty(\mathbf{R}^d), 0 \notin \text{supp}(\hat{g})$. Then there are constants $C, c > 0$ so that*

$$\sup_{R \in (0, 1]} \int \left| \mathbf{E} L^{\tilde{\nu}_R} g \left(x + Z_t^{\tilde{\nu}_R} \right) \right| dx \leq C e^{-ct}, t > 0.$$

Proof. Let $h \in C_0^\infty(\mathbf{R}^d)$, $0 \leq h \leq 1$, and $h(\xi) = 1$ if $\xi \in \text{supp}(g)$, $h(\xi) = 0$ in a neighborhood of zero. Let

$$G_R(t, x) = \mathbf{E}L^{\tilde{\nu}_R}g\left(x + Z_t^{\tilde{\nu}_R}\right), x \in \mathbf{R}^d.$$

Then

$$\begin{aligned}\hat{G}_R(t, \xi) &= \exp\{\psi^{\tilde{\nu}_R}(\xi)t\} \psi^{\tilde{\nu}_R}(\xi) \hat{g}(\xi) \\ &= \exp\{\psi^{\tilde{\nu}_R}(\xi)t\} h(\xi) \psi^{\tilde{\nu}_R}(\xi) \hat{g}(\xi), \xi \in \mathbf{R}^d.\end{aligned}$$

Hence

$$G_R(t, x) = \int H_R(t, x - y) B_R(y) dy, x \in \mathbf{R}^d,$$

where

$$B_R(x) = L^{\tilde{\nu}_R}g(x), H_R(t, x) = \mathbf{E}h\left(x + Z_t^{\tilde{\nu}_R}\right) x \in \mathbf{R}^d.$$

Thus, by Lemma 6

$$\begin{aligned}\sup_{R \in (0, 1]} \int |G_R(t, x)| dx &\leq \sup_{R \in (0, 1]} \int |H_R(t, x)| dx \sup_{R \in (0, 1]} \int |B_R(x)| dx \\ &\leq C e^{-ct}, t > 0.\end{aligned}$$

□

Lemma 7. Let $\nu \in \mathfrak{A}_{sym}^\alpha$ satisfy **A** and **B**. Then

(i) For each $\beta, \kappa > 0$, there is $C > 0$ so that

$$\begin{aligned}|L^{\nu; \kappa}u|_{\beta, \infty} &\leq C |u|_{\beta + \kappa, \infty}, u \in C_b^\infty(\mathbf{R}^d), \\ |(I - L^\nu)^\kappa u|_{\beta, \infty} &\leq C |u|_{\beta + \kappa, \infty}, u \in C_b^\infty(\mathbf{R}^d),\end{aligned}$$

(ii) For each $0 < \kappa < \beta$, there is $C > 0$ so that

$$\begin{aligned}|u|_{\beta, \infty} &\leq C \left[|L^{\nu; \kappa}u|_{\beta - \kappa, \infty} + |u|_0 \right], u \in C_b^\infty(\mathbf{R}^d), \\ |u|_{\beta, \infty} &\leq C |(I - L^\nu)^\kappa u|_{\beta - \kappa, \infty}, u \in C_b^\infty(\mathbf{R}^d).\end{aligned}$$

The same holds with $\nu \in \mathfrak{A}^\alpha$ satisfying **A** and **B** if $\kappa \in \mathbf{N}$.

Proof. Let $u \in C_b^\infty(\mathbf{R}^d)$, $\tilde{\phi}, \tilde{\phi}_0 \in C_0^\infty(\mathbf{R}^d)$, be such that $\tilde{\phi}\phi = \phi$, $\tilde{\phi}_0\phi_0 = \phi_0$, where $\phi_0 = \mathcal{F}^{-1}\varphi_0$, and ϕ, φ_0 are the functions in the definition of the spaces.

(i) Let $r \in [0, 1]$. Then

$$\begin{aligned}(r - L^\nu)^\kappa u * \varphi_j &= \mathcal{F}^{-1} \left[(r - \psi^\nu)^\kappa \tilde{\phi}(N^{-j}\cdot) \hat{u}\phi(N^{-j}\cdot) \right] \\ &= \int H_r^j(x - y) u * \varphi_j(y) dy, x \in \mathbf{R}^d, j \geq 1, \\ (r - L^\nu)^\kappa u * \varphi_0 &= \mathcal{F}^{-1} \left[(r - \psi^\nu)^\kappa \tilde{\phi}_0 \hat{u}\phi_0 \right] = \int H_r^0(x - y) u * \varphi_0(y) dy, x \in \mathbf{R}^d,\end{aligned}$$

where $H_r^j = \mathcal{F}^{-1} \left[(r - \psi^\nu)^\kappa \tilde{\phi} (N^{-j} \cdot) \right]$, $j \geq 1$, $H_r^0 = \mathcal{F}^{-1} \left[(r - \psi^\nu)^\kappa \tilde{\phi}_0 \right]$.
Let

$$G_j = w (N^{-j})^{-\kappa} \mathcal{F}^{-1} \left[(rw (N^{-j}) - \psi^{\tilde{\nu}_{N^{-j}}})^\kappa \tilde{\phi} \right], j \geq 1.$$

Since

$$(r - \psi^\nu (\xi))^\kappa \tilde{\phi} (N^{-j} \xi) = w (N^{-j})^{-\kappa} [rw (N^{-j}) - \psi^{\tilde{\nu}_{N^{-j}}} (N^{-j} \xi)]^\kappa \tilde{\phi} (N^{-j} \xi), \xi \in \mathbf{R}^d,$$

it follows by Lemma 5 that

$$\begin{aligned} \int |H_r^j (x)| dx &= \int |G_j (x)| dx \\ &= w (N^{-j})^{-\kappa} \int \left| (rw (N^{-j}) - L^{\tilde{\nu}_{N^{-j}}})^\kappa \tilde{\varphi} (x) \right| dx \\ &\leq Cw (N^{-j})^{-\kappa}, j \geq 0. \end{aligned}$$

(ii) Let $0 < \kappa < \beta$, $r \in [0, 1]$. Then for $j \geq 1$,

$$\begin{aligned} u * \varphi_j &= (r - L^\nu)^\kappa (r - L^\nu)^{-\kappa} u * \varphi_j \\ &= \mathcal{F}^{-1} \left[(r - \psi^\nu)^{-\kappa} \tilde{\phi} (N^{-j} \cdot) (r - \psi^\nu)^\kappa \hat{u} \phi (N^{-j} \cdot) \right] \\ &= \int H_r^j (x - y) (r - L^\nu)^\kappa u * \varphi_j (y) dy, x \in \mathbf{R}^d, j \geq 1, \end{aligned}$$

where

$$H_r^j = \mathcal{F}^{-1} \left[(r - \psi^\nu)^{-\kappa} \tilde{\phi} (N^{-j} \cdot) \right], j \geq 1, r \geq 0.$$

Let

$$G_j = w (N^{-j})^\kappa \mathcal{F}^{-1} \left[(rw (N^{-j}) - \psi^{\tilde{\nu}_{N^{-j}}})^{-\kappa} \tilde{\phi} \right], j \geq 1.$$

It follows by Corollary 1 that there is C independent of $r \geq 0, j \geq 1$, so that

$$\begin{aligned} \int |H_r^j (x)| dx &= \int |G_j (x)| dx = w (N^{-j})^\kappa \int \left| (rw (N^{-j}) - L^{\tilde{\nu}_{N^{-j}}})^{-\kappa} \tilde{\varphi} (x) \right| dx \\ &\leq Cw (N^{-j})^\kappa. \end{aligned}$$

On the other hand,

$$|u * \varphi_0|_0 \leq C |u|_0.$$

The statement follows. \square

For $\beta > 0, \kappa > 0$, we define the following norms:

$$\begin{aligned} |u|_{\nu, \kappa, \beta} &= |u|_0 + |L^{\nu; \kappa} u|_{\beta, \infty}, u \in C_b^\infty (\mathbf{R}^d), \\ ||u||_{\nu; \kappa, \beta} &= |(I - L^\nu)^\kappa u|_{\beta, \infty}, u \in C_b^\infty (\mathbf{R}^d), \end{aligned}$$

with ν satisfying **A** and **B**. An immediate consequence of Lemma 7 is the following norm equivalence.

Corollary 3. *Let $\nu \in \mathfrak{A}_{sym}^\alpha$ be a Lévy measure satisfying **A** and **B**, $\beta > 0, \kappa > 0$. Then norms $|u|_{\nu, \kappa, \beta}$, $\|u\|_{\nu, \kappa, \beta}$ and $|u|_{\beta+\kappa, \infty}$ are equivalent on $C_b^\infty(\mathbf{R}^d)$.*

*The same holds with $\nu \in \mathfrak{A}^\alpha$ satisfying **A** and **B** if $\kappa \in \mathbf{N}$.*

Proof. Let $\beta, \kappa > 0$. By Lemma 7,

$$\begin{aligned} |(I - L^\nu)^\kappa u|_{\beta, \infty} &\leq C |u|_{\beta+\kappa, \infty} \leq C \left[|L^{\nu; \kappa} u|_{\beta, \infty} + |u|_0 \right] \\ &= C |u|_{\nu; \beta, \kappa}, u \in C_b^\infty(\mathbf{R}^d). \end{aligned}$$

On the other hand, by Lemmas 1 and 7,

$$\begin{aligned} |u|_0 &\leq C |u|_{\beta, \infty} \leq C |u|_{\beta+\kappa, \infty} \leq C |(I - L^\nu)^\kappa u|_{\beta, \infty}, \\ |L^{\nu; \kappa} u|_{\beta, \infty} &\leq C |u|_{\beta+\kappa, \infty} \leq C |(I - L^\nu)^\kappa u|_{\beta, \infty}, u \in C_b^\infty(\mathbf{R}^d). \end{aligned}$$

□

Corollary 4. *Let $\nu \in \mathfrak{A}_{sym}^\alpha$ and $\pi \in \mathfrak{A}^\alpha$ be a Lévy measure satisfying **A** and **B** such that $w_\pi \sim w_\nu$. Then for any $\kappa \in \mathbf{N}, \beta > 0$, there are constants $c, C > 0$ so that*

$$|(L^\pi)^\kappa u|_{\beta, \infty} \leq C_1 |u|_{\nu; \kappa, \beta} \leq C_2 \left[|(L^\pi)^\kappa u|_{\beta, \infty} + |u|_0 \right], u \in C_b^\infty(\mathbf{R}^d).$$

Proof. Indeed, by Corollary 3,

$$\begin{aligned} |(L^\pi)^\kappa u|_{\beta, \infty} &\leq C |u|_{\kappa+\beta, \infty} \leq C |u|_{\nu; \kappa, \beta} \\ &\leq C |u|_{\kappa+\beta, \infty} \leq C_2 \left[|(L^\pi)^\kappa u|_{\beta, \infty} + |u|_0 \right], u \in C_b^\infty(\mathbf{R}^d). \end{aligned}$$

□

2.2.2. *Extension of norm equivalence to $\tilde{C}_{\infty\infty}^\beta(\mathbf{R}^d)$.* We extend the definition of $(a - L^\nu)^\kappa$ and the norm equivalence (see Corollary 3 above) from $C_b^\infty(\mathbf{R}^d)$ to $\tilde{C}_{\infty\infty}^\beta(\mathbf{R}^d)$. We start with the following observation.

Remark 4. *Let $0 < \beta' < \beta$. Then for each $\varepsilon > 0$ there is a constant $C_\varepsilon > 0$ so that*

$$|u|_{\beta', \infty} \leq \varepsilon |u|_{\beta, \infty} + C_\varepsilon |u|_0, u \in \tilde{C}_{\infty\infty}^\beta(\mathbf{R}^d).$$

Indeed, For each $\varepsilon > 0$ there is $K > 1$ so that $w(N^{-j})^{\beta-\beta'} \leq \varepsilon$ if $j \geq K$. Hence

$$\begin{aligned} w(N^{-j})^{-\beta'} |u * \varphi_j|_0 &= w(N^{-j})^{\beta-\beta'} w(N^{-j})^{-\beta} |u * \varphi_j|_0 \\ &\leq \varepsilon |u|_{\beta, \infty} + \max_{k \leq K} \left[w(N^{-k})^{-\beta'} |u * \varphi_k|_0 \right] \\ &\leq \varepsilon |u|_{\beta, \infty} + C_\varepsilon |u|_0. \end{aligned}$$

Proposition 2. *Let $\beta \in (0, \infty)$, $u \in \tilde{C}_{\infty, \infty}^{\beta}(\mathbf{R}^d)$. Then there exists a sequence $u_n \in C_b^{\infty}(\mathbf{R}^d)$, such that*

$$|u|_{\beta, \infty} \leq \liminf_n |u_n|_{\beta, \infty}, \quad |u_n|_{\beta, \infty} \leq C |u|_{\beta, \infty}$$

for some $C > 0$ that only depends on d, N .

Moreover, for any $0 < \beta' < \beta$,

$$|u_n - u|_{\beta', \infty} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Proof. Set $u_n = \sum_{j=0}^n u * \varphi_j$, $n \geq 1$. Obviously, $u_n \in C_b^{\infty}(\mathbf{R}^d)$, $n \geq 1$, and by Lemma 1, $u = \sum_{j=0}^{\infty} u * \varphi_j$ is a bounded continuous function. Since

$$\varphi_k = \sum_{l=-1}^1 \varphi_{k+l} * \varphi_k, \quad k \geq 1, \quad \varphi_0 = (\varphi_0 + \varphi_1) * \varphi_0,$$

we have for $n > 1$,

$$\begin{aligned} (u - u_n) * \varphi_k &= 0, \quad k < n, \\ (u - u_n) * \varphi_k &= (u * \varphi_{k-1} + u * \varphi_k + u * \varphi_{k+1}) * \varphi_k, \quad k > n + 1, \\ (u - u_n) * \varphi_n &= (u * \varphi_{n+1}) * \varphi_n, \\ (u - u_n) * \varphi_{n+1} &= (u * \varphi_{n+1} + u * \varphi_{n+2}) * \varphi_{n+1}. \end{aligned}$$

Hence there is a constant C so that

$$|u_n * \varphi_j|_0 \leq C |u * \varphi_j|_0, \quad j \geq 0, \quad n > 1,$$

and for $n > 1$,

$$\sup_{j < n} w(N^{-j})^{-\beta} |u * \varphi_j|_0 = \sup_{j < n} w(N^{-j})^{-\beta} |u_n * \varphi_j|_0 \leq |u_n|_{\beta, \infty}.$$

Thus

$$|u|_{\beta, \infty} \leq \underline{\lim}_n |u_n|_{\beta, \infty},$$

and

$$|u_n|_{\beta, \infty} \leq C |u|_{\beta, \infty}, \quad n > 1.$$

Now, by Remark 4, for each $\varepsilon > 0$, there is a constant C_{ε} so that

$$|u_n - u|_{\beta', \infty} \leq \varepsilon \left(|u_n|_{\beta, \infty} + |u|_{\beta, \infty} \right) + C_{\varepsilon} |u_n - u|_0.$$

Since by Lemma 1, $|u_n - u|_0 \rightarrow 0$, the statement follows. \square

Using the approximating sequence introduced in Proposition 2, we can extend $L^{\nu; \kappa} u, (I - L^{\nu})^{\kappa} u$, $0 < \kappa < \beta$, to all $u \in \tilde{C}_{\infty, \infty}^{\beta}(\mathbf{R}^d)$, $\beta > 0$.

Proposition 3. *Let ν be a Lévy measure satisfying **A** and **B**, $\beta > 0$ and $u \in \tilde{C}_{\infty, \infty}^{\beta}(\mathbf{R}^d)$. Let $u_n \in C_b^{\infty}(\mathbf{R}^d)$ be an approximating sequence of u in Proposition 2. Then for each $\kappa \in (0, \beta)$ there are bounded continuous functions, denoted $(I - L^{\nu})^{\kappa} u, L^{\nu; \kappa} u \in \tilde{C}_{\infty, \infty}^{\beta - \kappa}$, so that for any $0 < \beta' < \beta - \kappa$,*

$$|L^{\nu; \kappa} u_n - L^{\nu; \kappa} u|_{\beta', \infty} + |(I - L^{\nu})^{\kappa} u - (I - L^{\nu})^{\kappa} u_n|_{\beta', \infty} \rightarrow 0$$

as $n \rightarrow \infty$. Moreover, for each $\kappa \in (0, \beta)$ there is $C > 0$ independent of $u \in \tilde{C}_{\infty, \infty}^{\beta}(\mathbf{R}^d)$ so that

$$(2.12) \quad |L^{\nu; \kappa} u|_{\beta - \kappa, \infty} \leq C |u|_{\beta, \infty}, |(I - L^{\nu})^{\kappa} u|_{\beta - \kappa, \infty} \leq C |u|_{\beta, \infty},$$

and

$$(2.13) \quad |u|_{\beta, \infty} \leq C \left[|L^{\nu; \kappa} u|_{\beta - \kappa, \infty} + |u|_0 \right], |u|_{\beta, \infty} \leq C |(I - L^{\nu})^{\kappa} u|_{\beta - \kappa, \infty}.$$

Proof. Let $u \in \tilde{C}_{\infty, \infty}^{\beta}(\mathbf{R}^d)$. By Proposition 2, there is a sequence $u_n \in C_b^{\infty}(\mathbf{R}^d)$ such that

$$|u|_{\beta, \infty} \leq \liminf_n |u_n|_{\beta, \infty}, \quad |u_n|_{\beta, \infty} \leq C |u|_{\beta, \infty}, n \geq 1,$$

for some $C > 0$ independent of u , and for any $\kappa < \beta, \beta' \in (0, \beta - \kappa)$, see Lemma 7 as well,

$$\begin{aligned} & |L^{\nu; \kappa} u_n - L^{\nu; \kappa} u_m|_{\beta', \infty} + |(I - L^{\nu})^{\kappa} u_n - (I - L^{\nu})^{\kappa} u_m|_{\beta', \infty} \\ & \leq C |u_n - u_m|_{\beta'} \rightarrow 0 \text{ as } n, m \rightarrow \infty. \end{aligned}$$

Hence there are bounded continuous functions, denoted $L^{\mu; \kappa} u, (I - L^{\nu})^{\kappa} u$, so that

$$|L^{\nu; \kappa} u_n - L^{\nu; \kappa} u|_0 + |(I - L^{\nu})^{\kappa} u_n - (I - L^{\nu})^{\kappa} u|_0 \rightarrow 0$$

as $n \rightarrow \infty$. Thus

$$\begin{aligned} & |L^{\nu; \kappa} u_n * \varphi_j - L^{\nu; \kappa} u * \varphi_j|_0 \\ & + |(I - L^{\nu})^{\kappa} u_n * \varphi_j - (I - L^{\nu})^{\kappa} u * \varphi_j|_0 \\ & \rightarrow 0, \quad j \geq 0, \end{aligned}$$

as $n \rightarrow \infty$. Now, for each $m > 1$, and $a = 0, 1$,

$$\begin{aligned} & \sup_{j \leq m} w(N^{-j})^{-\beta - \kappa} |(a - L^{\nu})^{\kappa} u * \varphi_j|_0 \\ & = \lim_{n \rightarrow \infty} \sup_{j \leq m} w(N^{-j})^{-\beta - \kappa} |(a - L^{\nu})^{\kappa} u_n * \varphi_j|_0 \\ & \leq \sup_n |(a - L^{\nu})^{\kappa} u_n|_{\beta - \kappa} \leq \sup_n C |u_n|_{\beta, \infty} \leq C |u|_{\beta, \infty}. \end{aligned}$$

Hence $(a - L^{\nu})^{\kappa} u \in \tilde{C}_{\infty, \infty}^{\beta - \kappa}(\mathbf{R}^d)$, $a = 0, 1$, and (2.12) holds.

Now for every $j \geq 0$, we have

$$(2.14) \quad \begin{aligned} [(a - L^{\nu})^{\kappa} u] * \varphi_j &= \lim_n [(a - L^{\nu})^{\kappa} u_n] * \varphi_j \\ &= (a - L^{\nu})^{\kappa} [u * \varphi_j] \end{aligned}$$

uniformly. By the definition of the approximation sequence (see proof of Proposition 2),

$$|(a - L^{\nu})^{\kappa} [u_n * \varphi_j]|_0 \leq C |(a - L^{\nu})^{\kappa} [u * \varphi_j]|_0, j \geq 0.$$

Hence

$$\begin{aligned} |u|_{\beta, \infty} &\leq \liminf_n |u_n|_{\beta, \infty} \leq C \liminf_n |(I - L^\nu)^\kappa u_n|_{\beta - \kappa, \infty} \\ &\leq C |(I - L^\nu)^\kappa|_{\beta - \kappa, \infty}, \end{aligned}$$

and similarly,

$$\begin{aligned} |u|_{\beta, \infty} &\leq \liminf_n |u_n|_{\beta, \infty} \leq C \liminf_n \left[|L^{\nu; \kappa} u_n|_{\beta - \kappa, \infty} + |u_n|_0 \right] \\ &\leq C \left[|L^{\nu; \kappa} u|_{\beta - \kappa, \infty} + |u|_0 \right]. \end{aligned}$$

The statement is proved. \square

Proposition 4. *Let $\nu \in \mathfrak{A}_{sym}^\alpha$ be a Lévy measure satisfying **A** and **B**, $\beta > 0, \kappa > 0$. Then norms $|u|_{\nu, \kappa, \beta}$, $\|u\|_{\nu, \kappa, \beta}$ and $|u|_{\beta + \kappa, \infty}$ are equivalent on $\tilde{C}_{\infty \infty}^{\beta + \kappa}(\mathbf{R}^d)$.*

Proof. We show the equivalence by repeating proof of Corollary 3 where the equivalence of the same norms on $C_b^\infty(\mathbf{R}^d)$ was derived. Only instead of Lemma 7 we use Proposition 3. \square

3. PROOF OF MAIN THEOREM

We assume in this section that **A**, **B** and **C** hold. First we solve the equation with smooth input functions.

Proposition 5. *Let $\nu \in \mathfrak{A}^\alpha$, $\beta \in (0, 1)$, $\lambda \geq 0$. Assume that $f(t, x) \in C_b^\infty(H_T)$. Then there is a unique solution $u \in C_b^\infty(H_T)$ to*

$$(3.1) \quad \begin{aligned} \partial_t u(t, x) &= L^\nu u(t, x) - \lambda u(t, x) + f(t, x), \\ u(0, x) &= 0, \quad (t, x) \in [0, T] \times \mathbf{R}^d. \end{aligned}$$

Proof. EXISTENCE. Denote $F(r, Z_r^\nu) = e^{-\lambda(r-s)} f(s, x + Z_r^\nu - Z_s^\nu)$, $s \leq r \leq t$, and apply the Itô formula to $F(r, Z_r^\nu)$ on $[s, t]$.

$$\begin{aligned} &e^{-\lambda(t-s)} f(s, x + Z_t^\nu - Z_s^\nu) - f(s, x) \\ &= -\lambda \int_s^t F(r, Z_r^\nu) dr + \int_s^t \int \chi_\alpha(y) y \cdot \nabla F(r, Z_{r-}^\nu) \tilde{J}(dr, dy) \\ &\quad + \int_s^t \int [F(r, Z_{r-}^\nu + y) - F(r, Z_{r-}^\nu) - \chi_\alpha(y) y \cdot \nabla F(r, Z_{r-}^\nu)] J(dr, dy). \end{aligned}$$

Take expectation for both sides and use the stochastic Fubini theorem,

$$\begin{aligned} &e^{-\lambda(t-s)} \mathbf{E} f(s, x + Z_t^\nu - Z_s^\nu) - f(s, x) \\ &= -\lambda \int_s^t e^{-\lambda(r-s)} \mathbf{E} f(s, x + Z_r^\nu - Z_s^\nu) dr + \int_s^t L^\nu e^{-\lambda(r-s)} \mathbf{E} f(s, x + Z_r^\nu - Z_s^\nu) dr. \end{aligned}$$

Integrate both sides over $[0, t]$ with respect to s and obtain

$$\begin{aligned} & \int_0^t e^{-\lambda(t-s)} \mathbf{E} f(s, x + Z_t^\nu - Z_s^\nu) ds - \int_0^t f(s, x) ds \\ = & -\lambda \int_0^t \int_0^r e^{-\lambda(r-s)} \mathbf{E} f(s, x + Z_r^\nu - Z_s^\nu) ds dr \\ & + \int_0^t L^\nu \int_0^r e^{-\lambda(r-s)} \mathbf{E} f(s, x + Z_r^\nu - Z_s^\nu) ds dr, \end{aligned}$$

which shows $u(t, x) = \int_0^t e^{-\lambda(t-s)} \mathbf{E} f(s, x + Z_{t-s}^\nu) ds$ solves (3.1) in the integral sense. Obviously, as a result of the dominated convergence theorem and Fubini's theorem, $u \in C_b^\infty(H_T)$. And by the equation, u is continuously differentiable in t .

UNIQUENESS. Suppose there are two solutions u_1, u_2 solving the equation, then $u := u_1 - u_2$ solves

$$(3.2) \quad \begin{aligned} \partial_t u(t, x) &= L^\nu u(t, x) - \lambda u(t, x), \\ u(0, x) &= 0. \end{aligned}$$

Fix any $t \in [0, T]$. Apply the Itô formula to $v(t-s, Z_s^\nu) := e^{-\lambda s} u(t-s, x + Z_s^\nu)$, $0 \leq s \leq t$, over $[0, t]$ and take expectation for both sides of the resulting identity, then

$$u(t, x) = -\mathbf{E} \int_0^t e^{-\lambda s} [(-\partial_t u - \lambda u + L^\nu u)(t-s, x + Z_{s-}^\nu)] ds = 0.$$

□

3.1. Hölder estimates of the smooth solution. First we derive the estimates of the solution corresponding to a smooth input function.

Proposition 6. *Let $\nu \in \mathfrak{A}^\alpha$, $\beta > 0$ and **A-C** hold. Let $u \in C_b^\infty(H_T)$ be the unique solution u to (3.1) with $f \in C_b^\infty(H_T)$. Then*

$$(3.3) \quad |u|_{\beta, \infty} \leq C \rho_\lambda(T) |f|_{\beta, \infty},$$

$$(3.4) \quad |u|_{1+\beta, \infty} \leq C [1 + \rho_\lambda(T)] |f|_{\beta, \infty}$$

and for any $\mu \in [0, 1]$, $t' < t \leq T$,

$$(3.5) \quad \begin{aligned} & |u(t, \cdot) - u(t', \cdot)|_{\mu+\beta, \infty} \\ & \leq C \left\{ (t-t')^{1-\mu} + [1 + \rho_\lambda(T)] |t-t'| \right\} |f|_{\beta, \infty}, \end{aligned}$$

where $\rho_\lambda(T) = \frac{1}{\lambda} \wedge T$.

Proof. Since $f \in C_b^\infty(H_T)$, by Lemma 1,

$$\begin{aligned} f(t, x) &= (f(t, \cdot) * \varphi_0(\cdot))(x) + \sum_{j=1}^{\infty} (f(t, \cdot) * \varphi_j(\cdot))(x) \\ &= f_0(t, x) + \sum_{j=1}^{\infty} f_j(t, x), (t, x) \in H_T. \end{aligned}$$

Accordingly, for $j \geq 0$,

$$u_j(t, x) = u(t, x) * \varphi_j(x) = \int_0^t e^{-\lambda(t-s)} \mathbf{E} f_j(s, x + Z_{t-s}^\nu) ds, (t, x) \in H_T,$$

is the solution to (3.1) with input $f_j = f * \varphi_j$. In terms of Fourier transform,

$$\begin{aligned} &\hat{u}_j(t, \xi) \\ &= \int_0^t \exp\{-(\lambda - \psi^\nu(\xi))(t-s)\} \hat{f}(s, \xi) \phi(N^{-j}\xi) ds \\ &= \int_0^t e^{-\lambda(t-s)} \exp\{\psi^{\bar{\nu}}_{N^{-j}}(N^{-j}\xi) w(N^{-j})^{-1}(t-s)\} \tilde{\phi}(N^{-j}\xi) \hat{f}_j(s, \xi) ds, j \geq 1. \end{aligned}$$

Denote $w_j = w(N^{-j})^{-1}$. Then for $j \geq 0$,

$$\begin{aligned} &u_j(t, x) \\ &= \int_0^t e^{-\lambda(t-s)} \int H^j(t-s, x-y) f_j(s, y) dy ds, t \in [0, T], x \in \mathbf{R}^d, \end{aligned}$$

with

$$\begin{aligned} H^j(t, x) &= N^{jd} \mathbf{E} \tilde{\varphi}\left(N^j x + Z_{w_j t}^{\bar{\nu}_{N^{-j}}}\right), (t, x) \in H_T, j \geq 1, \\ H^0(t, x) &= \mathbf{E} \tilde{\varphi}_0(x + Z_t^\nu), (t, x) \in H_T. \end{aligned}$$

Hence

$$(3.6) \quad \int |H^j(t, x)| dx = \int |G^j(t, x)| dx, t > 0, j \geq 0,$$

with $G^0 = H^0$ and

$$G^j(t, x) = \mathbf{E} \tilde{\varphi}\left(x + Z_{w_j t}^{\bar{\nu}_{N^{-j}}}\right), (t, x) \in \mathbf{R}^d, j \geq 1.$$

First we estimate the solution itself. For $j \geq 1$, by Lemma 6,

$$\begin{aligned} |u_j(t, \cdot)|_0 &\leq |f_j|_0 \int_0^t e^{-\lambda(t-s)} \int |G^j(t-s, x)| dx ds \\ &\leq |f_j|_0 \int_0^t e^{-\lambda(t-s)} e^{-cw_j(t-s)} ds \leq C w_j^{-1} |f_j|_0. \end{aligned}$$

Directly,

$$|u_0(t, \cdot)|_0 \leq |f_0|_0 \int_0^t e^{-\lambda(t-s)} ds \leq \left(\frac{1}{\lambda} \wedge T\right) |f_0|_0.$$

Hence

$$|u|_{1+\beta, \infty} \leq C \left[1 + \left(\frac{1}{\lambda} \wedge T \right) \right] |f|_{\beta, \infty}.$$

Now we estimate time differences. For fixed $0 < t' < t \leq T, j \geq 0$,

$$\begin{aligned} & u_j(t, x) - u_j(t', x) \\ &= \int_{t'}^t e^{-\lambda(t-s)} \int H^j(t-s, x-y) f_j(s, y) dy ds \\ & \quad + \left(e^{-\lambda(t-t')} - 1 \right) \int_0^{t'} e^{-\lambda(t'-s)} \int H^j(t-s, x-y) f_j(s, y) dy ds \\ & \quad + \int_0^{t'} e^{-\lambda(t'-s)} \int [H^j(t-s, x-y) - H^j(t'-s, x-y)] f_j(s, y) dy ds \\ &= A_1^j(x) + A_2^j(x) + A_3^j(x), x \in \mathbf{R}^d. \end{aligned}$$

First, by Lemma 6, for $j \geq 1$,

$$\begin{aligned} |A_1^j|_0 &\leq \int_{t'}^t e^{-\lambda(t-s)} \int |G^j(t-s, y)| dy ds |f_j|_0 \\ &\leq C \int_{t'}^t e^{-\lambda(t-s)} e^{-cw_j(t-s)} ds |f_j|_0 \leq C \int_{t'}^t e^{-cw_j(t-s)} ds |f_j|_0 \\ &\leq C w_j^{-1} \left[1 - e^{-cw_j(t-t')} \right] |f_j|_0. \end{aligned}$$

And

$$|A_1^0| \leq C |f_0|_0 \int_{t'}^t e^{-\lambda(t-s)} ds \leq C |f_j|_0 |t - t'|.$$

By (3.6) and Lemma 6, for $j \geq 1$,

$$\begin{aligned} & |A_2^j|_0 \\ &\leq \left(1 - e^{-\lambda(t-t')} \right) \int_0^{t'} e^{-\lambda(t'-s)} \int |G^j(t-s, y)| dy ds |f_j|_0 \\ &\leq C \left(1 - e^{-\lambda(t-t')} \right) \int_0^{t'} e^{-\lambda(t'-s)} e^{-cw_j(t-s)} ds |f_j|_0. \end{aligned}$$

Thus for $j \geq 1$,

$$(3.7) \quad \begin{aligned} & |A_2^j|_0 \\ &\leq C \left(1 - e^{-\lambda(t-t')} \right) \int_0^{t'} e^{-\lambda(t'-s)} ds |f_j|_0 \leq C |f_j|_0 |t - t'|, \end{aligned}$$

in the mean time,

$$(3.8) \quad |A_2^j|_0 \leq C |f_j|_0 \int_0^{t'} e^{-cw_j(t'-s)} ds \leq C |f_j|_0 w_j^{-1}.$$

For $j = 0$,

$$\begin{aligned} |A_2^0|_0 &\leq C \left(1 - e^{-\lambda(t-t')}\right) \int_0^{t'} e^{-\lambda(t'-s)} ds |f_j|_0 \\ &\leq C |t - t'| \lambda \int_0^{t'} e^{-\lambda(t'-s)} ds |f_j|_0 \leq C |f_j|_0 |t - t'|. \end{aligned}$$

At last, for $j \geq 1$,

$$|A_3^j|_0 \leq |f_j|_0 \int_0^{t'} \int |G^j(t-s, y) - G^j(t'-s, y)| dy ds.$$

Note for $s \leq t'$,

$$\begin{aligned} &G^j(t-s, y) - G^j(t'-s, y) \\ &= \mathbf{E} \left[\tilde{\varphi} \left(y + Z_{w_j(t-s)}^{\tilde{\nu}_{N-j}} \right) - \tilde{\varphi} \left(y + Z_{w_j(t'-s)}^{\tilde{\nu}_{N-j}} \right) \right] \\ &= \mathbf{E} \int_{w_j(t'-s)}^{w_j(t-s)} L^{\tilde{\nu}_{N-j}} \tilde{\varphi} \left(y + Z_r^{\tilde{\nu}_{N-j}} \right) dr, \end{aligned}$$

and by Corollary 2,

$$\begin{aligned} &\int |G^j(t-s, y) - G^j(t'-s, y)| dy \\ &\leq C \int_{w_j(t'-s)}^{w_j(t-s)} e^{-cr} dr \leq C e^{-cw_j(t'-s)} \left[1 - e^{-cw_j(t-t')}\right]. \end{aligned}$$

Thus for $j \geq 1$,

$$\begin{aligned} |A_3^j|_0 &\leq C |f_j|_0 \left[1 - e^{-cw_j(t-t')}\right] \int_0^{t'} e^{-cw_j(t'-s)} ds \\ &= C w_j^{-1} |f_j|_0 \left[1 - e^{-cw_j(t-t')}\right] \left[1 - e^{-cw_j t'}\right] \\ &\leq C |f_j|_0 w_j^{-1} \left(1 - e^{-cw_j(t-t')}\right). \end{aligned}$$

In addition,

$$\begin{aligned} |A_3^0|_0 &\leq C |f_0|_0 \int_0^{t'} e^{-\lambda(t'-s)} ds |t - t'| \\ &\leq C \left(\frac{1}{\lambda} \wedge T\right) |f_0|_0 |t - t'|. \end{aligned}$$

Summarizing,

$$|u_0(t, \cdot) - u_0(t', \cdot)|_0 \leq C \left[1 + \left(\frac{1}{\lambda} \wedge T\right)\right] |f_0|_0 |t - t'|,$$

and

$$\begin{aligned} & |u_j(t, \cdot) - u_j(t', \cdot)|_0 \\ & \leq C |f_j|_0 \left[(|t - t'| \wedge w_j^{-1}) + w_j^{-1} (1 - e^{-cw_j(t-t')}) \right] \\ & = C |f_j|_0 w_j^{-1} \left[(|t - t'| w_j) \wedge 1 + (1 - e^{-cw_j(t-t')}) \right], \end{aligned}$$

which leads to

$$|u_j(t, \cdot) - u_j(t', \cdot)|_0 \leq C w_j^{-\mu} (t - t')^{1-\mu}, \quad \mu \in [0, 1], \quad j \geq 1.$$

Thus

$$\begin{aligned} & |u(t, \cdot) - u(t', \cdot)|_{\mu+\beta, \infty} \\ & \leq C |f|_{\beta, \infty} \left\{ (t - t')^{1-\mu} + \left[1 + \left(\frac{1}{\lambda} \wedge T \right) \right] |t - t'| \right\} \end{aligned}$$

for any $\mu \in [0, 1]$. The statement is proved. \square

3.2. General Hölder inputs. EXISTENCE AND ESTIMATES. Given $f \in \tilde{C}_{\infty, \infty}^{\beta}(H_T)$, by Proposition 2, we can find a sequence of functions f_n in $C_b^{\infty}(H_T)$ such that

$$|f_n|_{\beta, \infty} \leq C |f|_{\beta, \infty}, \quad |f|_{\beta, \infty} \leq \liminf_n |f_n|_{\beta, \infty},$$

and for any $0 < \beta' < \beta$,

$$|f_n - f|_0 \leq C |f_n - f|_{\beta', \infty} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

According to Theorems 5 and 6, for each $f_n \in C_b^{\infty}(\mathbf{R}^d)$, there is a corresponding solution $u_n \in C_b^{\infty}(H_T)$:

(3.9)

$$u_n(t, x) = \int_0^t [L^{\nu} u_n(r, x) - \lambda u_n(r, x) + f_n(r, x)] dr, \quad (t, x) \in [0, T] \times \mathbf{R}^d.$$

By Theorem 6,

$$\begin{aligned} & |L^{\nu} u_m - L^{\nu} u_n|_{\beta', \infty} \\ & \leq C |L^{\mu} u_m - L^{\mu} u_n|_{\beta', \infty} \leq C |u_m - u_n|_{1+\beta', \infty} \\ & \leq C |f_m - f_n|_{\beta', \infty} \rightarrow 0, \text{ as } m, n \rightarrow \infty \end{aligned}$$

for all $\beta' \in (0, \beta)$, which by Lemma 1 implies that

$$|u_n - u_m|_0 + |L^{\nu} u_m - L^{\mu} u_n|_0 \rightarrow 0 \text{ as } m, n \rightarrow \infty.$$

So, there is $u \in \tilde{C}_{\infty, \infty}^{1+\beta'}(H_T)$ for any $\beta' \in (0, \beta)$ such that $|u_n - u|_{1+\beta', \infty} \rightarrow 0$ as $n \rightarrow \infty$. Passing to the limit in (3.9) we see that (3.9) holds for u . Let $\beta' \in (0, \beta)$ and $\beta - \beta' < q_1^{-1}$. Then

$$\left| L^{1+\beta'} u_n \right|_{\beta-\beta', \infty} \leq C |u_n|_{1+\beta, \infty} \leq C |f_n|_{\beta, \infty} \leq C |f|_{\beta, \infty}$$

implies that

$$\left| L^{1+\beta'} u_n(t, x) - L^{1+\beta'} u_n(t, y) \right| \leq C |f|_{\beta, \infty} w(|x - y|^{\beta - \beta'}), \quad x, y \in \mathbf{R}^d.$$

and passing to the limit we see that

$$\left| L^{1+\beta'} u(t, x) - L^{1+\beta'} u(t, y) \right| \leq C |f|_{\beta, \infty} w(|x - y|^{\beta - \beta'}), \quad x, y \in \mathbf{R}^d.$$

Hence $L^{1+\beta'} u \in \tilde{C}_{\infty, \infty}^{\beta - \beta'}(\mathbf{R}^d)$, i.e., $u \in \tilde{C}_{\infty, \infty}^{1+\beta}(H_T)$ and

$$|u|_{1+\beta, \infty} \leq C |f|_{\beta, \infty}.$$

The convergence of u_n to u implies easily other estimates.

UNIQUENESS. Suppose there are two solutions $u_1, u_2 \in \tilde{C}_{\infty, \infty}^{1+\beta}(H_T)$ to (1.1), then $u := u_1 - u_2$ solves

$$(3.10) \quad u(t, x) = \int_0^t [L^\nu u(r, x) - \lambda u(r, x)] dr, \quad (t, x) \in [0, T] \times \mathbf{R}^d.$$

Let $g \in C_0^\infty(\mathbf{R}^d)$, $0 \leq g \leq 1$, $\int g dx = 1$. For $\varepsilon > 0$, set

$$u_\varepsilon(t, x) = \int u(t, y) g_\varepsilon(x - y) dy = \int v(t, x - y) g_\varepsilon(y) dy, \quad (t, x) \in H_T,$$

with $g_\varepsilon(x) = \varepsilon^{-d} g(x/\varepsilon)$, $x \in \mathbf{R}^d$. Then $u_\varepsilon \in \tilde{C}_b^\infty(H_T)$ solves (3.10). Hence $u_\varepsilon = 0$ for all $\varepsilon > 0$. Thus $u = 0$, the solution is unique.

4. APPENDIX

We simply state a few results that were used in this paper. Let $\nu \in \mathfrak{A}^\alpha$, and

$$\begin{aligned} \delta(r) &= \delta_\nu(r) = \nu(\{|y| > r\}) > 0, r > 0, \\ w &= w_\nu(r) = \delta(r)^{-1}, r > 0, \lim_{r \rightarrow 0} w(r) = 0. \end{aligned}$$

We assume that $w = w_\nu$ is an O-RV function at zero, i.e.,

$$r_1(\varepsilon) = \lim_{x \rightarrow 0} \frac{\delta(\varepsilon x)^{-1}}{\delta(x)^{-1}} < \infty, \varepsilon > 0.$$

By Theorem 2 in [2], the following limits exist:

$$(4.1) \quad p_1 = \lim_{\varepsilon \rightarrow 0} \frac{\log r_1(\varepsilon)}{\log \varepsilon} \leq q_1 = \lim_{\varepsilon \rightarrow \infty} \frac{\log r_1(\varepsilon)}{\log \varepsilon}.$$

Lemma 8. *Assume $w = w_\nu$ is an O-RV function at zero.*

a) *Let $\beta > 0$ and $\tau > -\beta p_1$. There is $C > 0$ so that*

$$\int_0^x t^\tau w(t)^\beta \frac{dt}{t} \leq C x^\tau w(x)^\beta, \quad x \in (0, 1],$$

and $\lim_{x \rightarrow 0} x^\tau w(x)^\beta = 0$.

b) Let $\beta > 0$ and $\tau < -\beta q_1$. There is $C > 0$ so that

$$\int_x^1 t^\tau w(t)^\beta \frac{dt}{t} \leq C x^\tau w(x)^\beta, x \in (0, 1],$$

and $\lim_{x \rightarrow 0} x^\tau w(x)^\beta = \infty$.

c) Let $\beta < 0$ and $\tau > -\beta q_1$. There is $C > 0$ so that

$$\int_0^x t^\tau w(t)^\beta \frac{dt}{t} \leq C x^\tau w(x)^\beta, x \in (0, 1],$$

and $\lim_{x \rightarrow 0} x^\tau w(x)^\beta = 0$.

d) Let $\beta < 0$ and $\tau < -\beta p_1$. There is $C > 0$ so that

$$\int_x^1 t^\tau w(t)^\beta \frac{dt}{t} = \int_1^{x^{-1}} t^{-\tau} w\left(\frac{1}{t}\right)^\beta \frac{dt}{t} \leq C x^\tau w(x)^\beta, x \in (0, 1],$$

and $\lim_{x \rightarrow 0} x^\tau w(x)^\beta = \infty$.

Proof. The claims follow easily by Theorems 3, 4 in [2]. Because of the similarities, we will prove c) only. Let $\beta < 0$ and $\tau > -\beta q_1$. Then

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{w\left(\frac{1}{\varepsilon t}\right)^\beta}{w\left(\frac{1}{t}\right)^\beta} &= \lim_{x \rightarrow 0} \frac{w(x)^{-\beta}}{w(\varepsilon^{-1}x)^{-\beta}} = \lim_{x \rightarrow 0} \frac{w(\varepsilon \varepsilon^{-1}x)^{-\beta}}{w(\varepsilon^{-1}x)^{-\beta}} \\ &= \lim_{x \rightarrow 0} \frac{w(\varepsilon x)^{-\beta}}{w(x)^{-\beta}} = r_1(\varepsilon)^{-\beta} < \infty, \varepsilon > 0. \end{aligned}$$

Hence $w\left(\frac{1}{t}\right)^\beta, t \geq 1$, is an O-RV function at infinity with

$$p = \lim_{\varepsilon \rightarrow 0} \frac{\log r_1(\varepsilon)^{-\beta}}{\log \varepsilon} = -\beta p_1 \leq -\beta q_1 = \lim_{\varepsilon \rightarrow \infty} \frac{\log r_1(\varepsilon)^{-\beta}}{\log \varepsilon} = q.$$

Then for $x \in (0, 1]$,

$$\int_0^x t^\tau w(t)^\beta \frac{dt}{t} = \int_{x^{-1}}^\infty t^{-\tau} w\left(\frac{1}{t}\right)^\beta \frac{dt}{t} \leq C x^\tau w(x)^\beta$$

by Theorem 3 in [2], and $\lim_{x \rightarrow 0} x^\tau w(x)^\beta = 0$ according to Theorem 4 in [2]. \square

Corollary 5. Assume $w = w_\nu$ is an O-RV function at zero and $p_1 > 0$. Let $N > 1, \beta > 0$. Then

$$\sum_{j=0}^{\infty} w(N^{-j})^\beta < \infty.$$

Proof. Indeed,

$$\sum_{j=0}^{\infty} w(N^{-j})^\beta \leq \int_0^\infty w(N^{-x})^\beta dx \leq C \int_0^1 w(t)^\beta \frac{dt}{t} < \infty,$$

because, by Lemma 8a),

$$\int_0^x w(t)^\beta \frac{dt}{t} \leq Cw(x)^\beta, x \in [0, 1].$$

□

We will need some Lévy measure moment estimates.

Lemma 9. *Let $\nu \in \mathfrak{A}^\alpha$, and $w = w_\nu$ be an O-RV function at zero with p_1, q_1 defined in (4.1). Assume*

$$\begin{aligned} 0 &< p_1 \leq q_1 < 1 \text{ if } \alpha \in (0, 1), \\ 1 &\leq p_1 \leq q_1 < 2 \text{ if } \alpha = 1, \\ 1 &< p_1 \leq q_1 < 2 \text{ if } \alpha \in (1, 2). \end{aligned}$$

Then

(i)

$$\begin{aligned} \sup_{R \in (0, 1]} \int (|y| \wedge 1) \tilde{\nu}_R(dy) &< \infty \text{ if } \alpha \in (0, 1), \\ \sup_{R \in (0, 1]} \int (|y|^2 \wedge 1) \tilde{\nu}_R(dy) &< \infty \text{ if } \alpha = 1, \\ \sup_{R \in (0, 1]} \int (|y|^2 \wedge |y|) \tilde{\nu}_R(dy) &< \infty \text{ if } \alpha \in (1, 2). \end{aligned}$$

(ii)

$$\inf_{R \in (0, 1]} \int_{|y| \leq 1} |y|^2 \tilde{\nu}_R(dy) \geq c_1,$$

for some $c_1 > 0$.

Proof. (i) Let $\alpha \in (0, 1)$. Then by Lemma 8,

$$\begin{aligned} \int_{|y| \leq 1} |y| \tilde{\nu}_R(dy) &= R^{-1} \int_{|y| \leq R} |y| \nu(dy) \\ &= R^{-1} \int_0^R [\delta(s) - \delta(R)] ds, \end{aligned}$$

and

$$\int_{|y| \leq 1} (|y| \wedge 1) \tilde{\nu}_R(dy) = R^{-1} \int_0^R w(s)^{-1} ds \leq C, R \in (0, 1].$$

Let $\alpha = 1$. Then, using Lemma 8 we have

$$\int_{|y| \leq 1} (|y|^2 \wedge 1) \tilde{\nu}_R(dy) = 2R^{-2} \int_0^R s^2 w(s)^{-1} \frac{ds}{s} \leq C, R \in (0, 1].$$

Let $\alpha \in (1, 2)$. Then similarly,

$$\begin{aligned} R^{-1} \int_{|y|>R} |y| \nu(dy) &= R^{-1} \int_0^\infty \delta(s \vee R) ds \\ &= \delta(R) + \int_R^\infty \delta(s) ds = \delta(R) + \int_R^\infty w(s)^{-1} ds \end{aligned}$$

and with $R \in (0, 1]$,

$$\begin{aligned} (4.2) \quad R^{-2} \int_{|y|\leq R} |y|^2 \nu(dy) &= 2R^{-2} \int_0^R s^2 [w(s)^{-1} - w(R)^{-1}] \frac{ds}{s} \\ &= 2R^{-2} \int_0^R s^2 w(s)^{-1} \frac{ds}{s} - w(R)^{-1}. \end{aligned}$$

Hence, by Lemma 8,

$$\begin{aligned} &\int \left(|y|^2 \wedge |y| \right) \tilde{\nu}_R(dy) \\ &\leq 2R^{-2} \int_0^R s^2 w(s)^{-1} \frac{ds}{s} + \int_R^1 w(s)^{-1} ds + \int_1^\infty w(s)^{-1} ds \\ &= 2R^{-2} \int_0^R s^2 w(s)^{-1} \frac{ds}{s} + \int_R^1 w(s)^{-1} ds + \int_{|y|>1} |y| \nu(dy) \\ &\leq Cw(R)^{-1}, \quad R \in (0, 1]. \end{aligned}$$

(ii) By (4.2), for $R \in (0, 1]$,

$$\begin{aligned} \int_{|y|\leq 1} |y|^2 \tilde{\nu}_R(dy) &= w(R) \int_{|y|\leq 1} |y|^2 \nu_R(dy) \\ &= 2R^{-2} \int_0^R s^2 \left[\frac{w(R)}{w(s)} - 1 \right] \frac{ds}{s} = 2 \int_0^1 s^2 \left[\frac{w(R)}{w(Rs)} - 1 \right] \frac{ds}{s}. \end{aligned}$$

Hence, by Fatou's lemma,

$$\underline{\lim}_{R \rightarrow 0} \int_{|y|\leq 1} |y|^2 \tilde{\nu}_R(dy) \geq 2 \int_0^1 s^2 \left[\frac{1}{r_1(s)} - 1 \right] \frac{ds}{s} = c_1 > 0$$

if $|\{s \in [0, 1] : r_1(s) < 1\}| > 0$, and

$$\liminf_{R \rightarrow 0} \frac{w(R)}{w(Rs)} = \frac{1}{\limsup_{R \rightarrow 0} \frac{w(Rs)}{w(R)}} = \frac{1}{r_1(s)}, \quad s \in (0, 1].$$

□

According to [8], Chapter 3, 70-74, any Lévy measure $\nu \in \mathfrak{A}^\alpha$ can be disintegrated as

$$\nu(\Gamma) = - \int_0^\infty \int_{S_{d-1}} \chi_\Gamma(rw) \Pi(r, dw) d\delta(r), \quad \Gamma \in \mathcal{B}(\mathbf{R}_0^d),$$

where $\delta = \delta_\nu$, and $\Pi(r, dw), r > 0$, is a measurable family of measures on the unit sphere S_{d-1} with $\Pi(r, S_{d-1}) = 1, r > 0$. The following is a straightforward consequence of Lemma 9(ii).

Corollary 6. *Let $\nu \in \mathfrak{A}^\alpha$,*

$$\nu(\Gamma) = - \int_0^\infty \int_{S_{d-1}} \chi_\Gamma(rw) \Pi(r, dw) d\delta(r), \Gamma \in \mathcal{B}(\mathbf{R}_0^d),$$

where $\delta = \delta_\pi, \Pi(r, dw), r > 0$, is a measurable family of measures on S_{d-1} with $\Pi(r, S_{d-1}) = 1, r > 0$. Assume $w = w_\nu = \delta_\nu^{-1}$ be an O-RV function at zero satisfying assumptions of Lemma 9, and

$$(4.3) \quad \inf_{|\hat{\xi}|=1} \int_{S_{d-1}} |\hat{\xi} \cdot w|^2 \Pi(r, dw) \geq c_0 > 0.$$

Then assumption **B** holds.

Proof. Indeed, for $|\hat{\xi}| = 1, R \in (0, 1]$, with $C > 0$,

$$\begin{aligned} & \int_{|y| \leq 1} |\hat{\xi} \cdot y|^2 \nu_R(dy) \\ &= R^{-2} \int_{|y| \leq R} |\hat{\xi} \cdot y|^2 \nu(dy) = -R^{-2} \int_0^R \int_{S_{d-1}} |\hat{\xi} \cdot w|^2 \Pi(r, dw) r^2 d\delta(r) \\ &\geq -c_0 R^{-2} \int_0^R r^2 d\delta(r) = c_0 R^{-2} \int_{|y| \leq R} |y|^2 \nu(dy) = c_0 \int_{|y| \leq 1} |y|^2 \nu_R(dy). \end{aligned}$$

Hence by Lemma 9(ii),

$$\begin{aligned} \inf_{R \in (0, 1]} \inf_{|\hat{\xi}|=1} \int_{|y| \leq 1} |\hat{\xi} \cdot y|^2 \tilde{\nu}_R(dy) &\geq c_0 \inf_{R \in (0, 1]} \int_{|y| \leq 1} |y|^2 \tilde{\nu}_R(dy) \\ &\geq c_0 c_1 > 0. \end{aligned}$$

□

Remark 5. *Let $\alpha \in (0, 2)$, $\nu \in \mathfrak{A}^\alpha$, and w_ν be an O-RV function at zero, $p_1 > 0$. By Theorems 3 and 4 in [2], for any $\sigma \in (0, p_1)$,*

$$\begin{aligned} \int_{r < |y| \leq 1} |y|^\sigma \nu(dy) &= \sigma \int_r^1 t^\sigma w(t)^{-1} \frac{dt}{t} - \delta(1) \\ &\geq cr^\sigma w(r)^{-1} - \delta(1) \rightarrow \infty \end{aligned}$$

as $r \rightarrow 0$. Hence $p_1 \leq \alpha$. On the other hand for any $\sigma > q_1$, by Lemma 9,

$$\int_{0 < |y| \leq 1} |y|^\sigma \nu(dy) \leq \sigma \int_0^1 t^\sigma w(t)^{-1} \frac{dt}{t} < \infty,$$

and $\alpha \leq q_1$.

REFERENCES

- [1] Abels, H., Kassman, M.: The Cauchy problem and the martingale problem for integrodifferential operators with non-smooth kernels. *Osaka J. Math.* 46, 661–683 (2009).
- [2] Aljančić, S. and Arandelović, D., O -regularly varying functions, *PIMB (NS)*, 22(36), 1977, pp 5-22.
- [3] Bingham, N.H., Goldie, C.M. and Teugels, J.L., *Regular Variation*, Cambridge University Press, 1987.
- [4] Bass, R.F.: Regularity results of stable-like operators. *J. Funct. Anal.* 257, 2693–2722 (2009)
- [5] Bergh, J. and Löfström, J., *Interpolation Spaces. An Introduction*, Springer Verlag, 1976.
- [6] Caffarelli, L., Silvestre, L.: Regularity theory for fully nonlinear integro-differential equations. *Comm. Pure Appl. Math.* 62, 597–638 (2009)
- [7] Chen, Z., Zhang, X. and Zhao, G., Well posedness of supercritical; SDE driven by Lévy processes with irregular drifts, arXiv:1709.04632 [math.PR], 2017.
- [8] Dellacherie, C. and Meyer, P.-A., *Probabilities and Potential -A*, North-Holland, 1978.
- [9] Dong, H., Kim, D.: Schauder estimates for a class of non-local elliptic equations. *Discrete Cont. Dyn.-A* 33(6), 2319–2347 (2013)
- [10] Farkas, W., Jacob, N. and Schilling, R.L., Function spaces related to continuous negative definite functions: ψ -Bessel potential spaces, *Dissertationes Mathematicae*, 1-60, 2001.
- [11] Farkas, W. and Leopold, H.G., Characterization of function spaces of generalized smoothness, *Annali di Matematica*, 185, 1-62, 2006.
- [12] Kalyabin, G.A., Description of functions in classes of Besov-Lizorkin-Triebel type, *Trudy Mat. Inst. Steklov*, 156, 160-173, 1980.
- [13] Kalyabin, G.A. and Lizorkin, P.I., Spaces of functions of generalized smoothness, *Math. Nachr.*, 133, 7-32, 1987.
- [14] Karamata, J., Sur un mode de croissance régulière des fonctions, *Mathematica (Cluj)*, 4, 1930, pp. 38-53.
- [15] Kim, P., Song, R. and Vondraček, Z., Heat Kernels of non-symmetric jump processes: beyond the stable case, *Potential Analysis*, 49(1), 2018, pp. 37–90.
- [16] Mikulevičius, R. and Pragarauskas, H., On the Cauchy problem for parabolic integro-differential operators in Hölder classes and the uniqueness of the Martingale problem, *Potential Analysis*, 40(4), 2014, pp 539-563.
- [17] Xu, F., On the Cauchy problem for parabolic integro-differential equations with space-dependent coefficients in generalized Hölder classes, 2018.

E-mail address: mikulvcs@usc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES

E-mail address: fanhuixu@usc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES