# On Exact and $\infty$ -Rényi Common Informations

Lei Yu and Vincent Y. F. Tan, Senior Member, IEEE

Abstract-Recently, two extensions of Wyner's common information-exact and Rényi common informations-were introduced respectively by Kumar, Li, and El Gamal (KLE), and the present authors. The class of common information problems involves determining the minimum rate of the common input to two independent processors needed to exactly or approximately generate a target joint distribution. For the exact common information problem, exact generation of the target distribution is required, while for Wyner's and  $\alpha$ -Rényi common informations, the relative entropy and Rényi divergence with order  $\alpha$  were respectively used to quantify the discrepancy between the synthesized and target distributions. The exact common information is larger than or equal to Wyner's common information. However, it was hitherto unknown whether the former is strictly larger than the latter for some joint distributions. In this paper, we first establish the equivalence between the exact and  $\infty$ -Rényi common informations, and then provide single-letter upper and lower bounds for these two quantities. For doubly symmetric binary sources, we show that the upper and lower bounds coincide, which implies that for such sources, the exact and  $\infty$ -Rényi common informations are completely characterized. Interestingly, we observe that for such sources, these two common informations are strictly larger than Wyner's. This answers an open problem posed by KLE. Furthermore, we extend Wyner's,  $\infty$ -Rényi, and exact common informations to sources with countably infinite or continuous alphabets, including Gaussian sources.

*Index Terms*—Wyner's common information, Rényi common information, Exact common information, Exact channel simulation, Exact source simulation, Communication complexity of correlation

#### I. INTRODUCTION

How much common randomness is needed to simulate two correlated sources in a distributed fashion? This problem (depicted in Fig. I), termed *distributed source simulation*, was first studied by Wyner [2], who used the normalized relative entropy (Kullback-Leibler divergence or KL divergence) to measure the discrepancy or "distance" between the simulated joint distribution and the joint distribution of the original correlated sources  $\pi_{XY}$ . He defined the minimum rate needed to ensure that the normalized relative entropy vanishes asymptotically as the *common information* (denoted as  $T_1(\pi_{XY})$ ) between the sources  $\pi_{XY}$ . He also established a single-letter

This work was supported by a Singapore Ministry of Education Tier 2 Grant (R-263-000-C83-112). This paper was presented in part at the 2019 IEEE International Symposium on Information Theory (ISIT) [1].

L. Yu is with the Department of Electrical and Computer Engineering, National University of Singapore (NUS), Singapore 117583 (e-mail: leiyu@nus.edu.sg). V. Y. F. Tan is with the Department of Electrical and Computer Engineering and the Department of Mathematics, NUS, Singapore 119076 (e-mail: vtan@nus.edu.sg).

Communicated by M. Raginsky, Associate Editor for Probability and Statistics.

Copyright (c) 2019 IEEE. Personal use of this material is permitted. However, permission to use this material for any other purposes must be obtained from the IEEE by sending a request to pubs-permissions@ieee.org. characterization for the common information, i.e., the common information between correlated sources X and Y is

$$T_1(\pi_{XY}) = C_{\mathsf{Wyner}}(\pi_{XY}) \tag{1}$$

$$:= \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} I(XY; W).$$
(2)

For Gray-Wyner's source coding problem subject to the condition that the total rate of all three messages is  $H_{\pi}(XY)$ (the joint entropy of the correlated sources  $\pi_{XY}$ ), the quantity in (2) was also used to characterize the minimum rate of the common message [2].

Recently, the present authors [3], [4] introduced the notion of  $\alpha$ -Rényi common information with  $\alpha \in [0, \infty]$ , which is defined as the minimum common rate when the KL divergence is replaced by more general divergences - the family of Rényi divergences with order  $\alpha \in [0, \infty]$ . When  $\alpha = 1$ , Rényi common information reduces to Wyner's common information. We proved that for Rényi divergences of order  $\alpha \in (0, 1]$ , the minimum rate needed to guarantee that the (normalized and unnormalized) Rényi divergences vanish asymptotically is equal to Wyner's common information. However, for Rényi divergences of order  $\alpha \in (1, 2]$ , we only provided lower and upper bounds. Numerical results show that our lower and upper bounds coincide for doubly symmetric binary sources (DSBSes), and for this case, both of them are strictly larger than Wyner's common information. Furthermore, the common information with approximation error measured by the total variation (TV) distance is also equal to Wyner's common information [3], [5], [6]; and exponential achievability and converse results for this case were established in [3], [5], [7].

Kumar, Li, and El Gamal (KLE) [6] extended Wyner's common information in a different way. They assumed variablelength codes and exact generation of the correlated sources  $(X,Y) \sim \pi_{XY}$ , instead of block codes and approximate simulation of  $\pi_{XY}$  as assumed by Wyner [2] and by us [3], [4]. For such exact generation problem, KLE [6] characterized the minimum common rate, coined *exact common information*, by

$$T_{\text{Exact}}(\pi_{XY}) := \lim_{n \to \infty} \frac{1}{n} G(\pi_{XY}^n).$$
(3)

where the common entropy

$$G(\pi_{XY}) := \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} H(W).$$
(4)

The exact common information is no smaller than Wyner's common information. However, it was previously unknown whether they are equal for all sources  $\pi_{XY}$ . Even for simple sources, e.g., DSBSes, the exact common information was still unknown. It is worth noting that the quantities  $G(\pi_{XY})$  and  $T_{\text{Exact}}(\pi_{XY})$  were first considered by Witsenhausen in 1976 [8, p. 331]. In [8], Witsenhausen studied properties of Wyner's common information. He provided an example [8, p. 331] (in

the framework of Gray-Wyner's source coding problem [2]) for which Wyner's common information can be attained by a one-shot coding scheme (i.e., block coding with  $n \ge 2$  is unnecessary), and at the same time, zero error is realized by this one-shot scheme. For this example, he showed that

$$G(\pi_{XY}) = C_{\mathsf{Wyner}}(\pi_{XY}),\tag{5}$$

which suggests that one may avoid block coding and also attain zero error. In order to better understand the relation between  $G(\pi_{XY})$  and  $C_{Wyner}(\pi_{XY})$  for an arbitrary  $\pi_{XY}$  (not specified to that example), Witsenhausen stated the following relation between  $T_{Exact}(\pi_{XY})$  and  $C_{Wyner}(\pi_{XY})$  for arbitrary  $\pi_{XY}$  with finite support:

$$T_{\text{Exact}}(\pi_{XY}) \stackrel{?}{=} C_{\text{Wyner}}(\pi_{XY}). \tag{6}$$

However, he did not provide a proof for (6). In this paper, we first completely characterize the exact common information for DSBSes, and then show that for this class of sources, the exact common information is strictly larger than Wyner's common information. This implies (6) does not always hold. Furthermore, sufficient conditions for (6) to hold (i.e., for equality of Wyner's common information and the exact common information) were investigated in [9].

The exact common information for *continuous* sources was studied by Li and El Gamal [10]. In [10], Li and El Gamal adopted dyadic decomposition schemes to construct a discrete common random variable W for continuous random variables with log-concave probability density functions (pdfs). By using such schemes, they established the first known upper bound on the exact common information for continuous sources. Specifically, for a pair of correlated sources  $(X, Y) \sim \pi_{XY}$  with a log-concave pdf, they showed that

$$I_{\pi}(X;Y) \le T_{\text{Exact}}(\pi_{XY}) \tag{7}$$

$$\leq G(\pi_{XY}) \tag{8}$$

$$\leq I_{\pi}(X;Y) + 24\log 2$$
 nats/symbol, (9)

where  $I_{\pi}(X; Y)$  denotes the mutual information between  $(X, Y) \sim \pi_{XY}$ . This result implies that the exact common information for continuous sources with log-concave pdfs is finite. Furthermore, it is worth noting that Li and El Gamal's dyadic decomposition scheme is a one-shot scheme, i.e., it is valid for the case with blocklength equal to 1. For Gaussian sources with correlation coefficient  $\rho \in [0, 1)$ , Li and El Gamal's upper bound in (9) reduces to

$$\frac{1}{2}\log\left[\frac{1}{1-\rho^2}\right] + 24\log 2 \text{ nats/symbol.}$$
(10)

It is known that  $T_{\text{Exact}}(\pi_{XY}) \geq C_{\text{Wyner}}(\pi_{XY})$  [6] and for joint Gaussian sources,  $C_{\text{Wyner}}(\pi_{XY}) = \frac{1}{2} \log \left[\frac{1+\rho}{1-\rho}\right]$  [11], [12]. Hence for joint Gaussian sources,  $T_{\text{Exact}}(\pi_{XY}) \geq \frac{1}{2} \log \left[\frac{1+\rho}{1-\rho}\right]$ . Note that there is a large gap between this lower bound and Li and El Gamal's upper bound in (10). In this paper, we prove a new upper bound on  $T_{\text{Exact}}(\pi_{XY})$ which is at most 0.72 bits/symbol larger than the lower bound  $\frac{1}{2} \log \left[\frac{1+\rho}{1-\rho}\right]$  and hence much tighter than Li and El Gamal's upper bound, albeit with the use of a block coding scheme.



Fig. 1. The distributed source simulation problem. For the exact common information problem, the discrete random variable  $W_n$  can be arbitrarily distributed, but for the Rényi common information problem, it is restricted to be uniformly distributed. Hence for the latter case, we use  $M_n$  to denote the common randomness, in place of the  $W_n$ .

#### A. Main Contributions

Our contributions include the following aspects.

- We first consider sources with finite alphabets. We establish the equivalence between the exact common information and ∞-Rényi common information. We provide a multi-letter characterization for the exact and ∞-Rényi common informations. Using this multi-letter characterization, we derive single-letter upper and lower bounds.
- When specialized to DSBSes, the upper and lower bounds coincide. This implies that the exact and ∞-Rényi common informations for DSBSes are completely solved. Interestingly, we show that they are both strictly larger than Wyner's common information. This solves an open problem posed by Kumar, Li, and El Gamal [6].
- We extend the exact and  $\infty$ -Rényi common informations, and also the relative entropy version and the TV distance version of Wyner's common information to sources with general (countable or continuous) alphabets, including Gaussian sources. We establish the equivalence between the exact and  $\infty$ -Rényi common informations for such general sources. We provide an upper bound on the exact and  $\infty$ -Rényi common informations for Gaussian sources, which is at least 22.28 bits/symbol smaller than Li and El Gamal's bound [10]. However, it is worth noting that theirs is a one-shot bound that is obtained by a scheme with blocklength 1, but ours is an asymptotic one which requires the blocklength to tend to infinity. Furthermore, we also completely characterize Wyner's common information for Gaussian sources which is equal to  $C_{\text{Wyner}}(\pi_{XY})$  that was computed in [11], [12] for Gaussian sources.
- Concerning the innovations in our proofs, they rely on the so-called *mixture decomposition* or *splitting technique*, which was previously used in [6], [9], [14]– [17]. However, in this paper, we combine it with various truncation techniques to deal with sources with countably infinite alphabets, and also combine it with truncation, discretization, and Li and El Gamal's dyadic decomposition techniques [10] to deal with sources with continuous alphabets. Besides the mixture decomposition technique, a superblock coding approach is also adopted to prove the equivalence between the exact and ∞-Rényi common informations. Furthermore, as by-products of our analyses, various lemmas are derived, e.g., the "chain

Com. Inf.	Fixed or Variable-Length	Exact or Approx.	Expressions for Various Alphabets
Wyner's CI [2]	Fixed	Approx. $(\frac{1}{n}D)$	1) Finite $(\mathcal{X}, \mathcal{Y})$ : $C_{\text{Wyner}}(\pi_{XY})$ [2] 2) Countably Infinite/Continuous* $(\mathcal{X}, \mathcal{Y})$ : $C_{\text{Wyner}}(\pi_{XY})$ [P, Cor. 2, 3]
$\alpha$ -Rényi CI, $\alpha \in [0, 2]$ [3], [4]	Fixed	Approx. $(\frac{1}{n}D_{\alpha} \text{ or } D_{\alpha})$	Finite $(\mathcal{X}, \mathcal{Y})$ : $\begin{cases} 0 & \alpha = 0 \\ C_{\text{Wyner}}(\pi_{XY}) & \alpha \in (0, 1] \ [3], [4] \\ \left[\Gamma^{\text{LB}}_{\alpha}(\pi_{XY}), \Gamma^{\text{UB}}_{\alpha}(\pi_{XY})\right] & \alpha \in (1, 2] \end{cases}$
∞-Rényi CI [P]	Fixed	Approx. $(\frac{1}{n}D_{\infty} \text{ or } D_{\infty})$	<ol> <li>Equivalent for Finite and Countably Infinite/Continuous* (X, Y) [P, Thm. 1].</li> <li>Multiletter Expressions: lim<sub>n→∞</sub> <sup>1</sup>/<sub>n</sub>G(π<sup>n</sup><sub>XY</sub>) [6] and lim<sub>n→∞</sub> <sup>1</sup>/<sub>n</sub>Γ(π<sup>n</sup><sub>XY</sub>) (for finite (X, Y)) [P, Thm. 1].</li> </ol>
Exact CI [6]	Variable	Exact $(P_{X^nY^n} = \pi^n_{XY})$	3) Singleletter Bounds: $[C_{Wyner}(\pi_{XY}), G(\pi_{XY})]$ [6]; $[\Gamma^{LB}(\pi_{XY}), \Gamma^{UB}(\pi_{XY})]$ (for finite $(\mathcal{X}, \mathcal{Y})$ ) [P, Thm. 2]. 4) Gaussian Sources: $\left[\frac{1}{2}\log\frac{1+\rho}{1-\rho}, \frac{1}{2}\log\frac{1}{1-\rho^2} + 24\log 2\right]$ [6], [10]; $\left[\frac{1}{2}\log\frac{1+\rho}{1-\rho}, \frac{1}{2}\log\frac{1+\rho}{1-\rho} + \frac{\rho}{1+\rho}\right]$ [P, Thm. 8].
Gács-Körner's CI [13]	Fixed	$\begin{array}{l} \text{Approx.} \\ (\mathbb{P}\left(f\left(X^{n}\right) \neq g\left(Y^{n}\right)\right) \leq \\ \varepsilon \text{ and } f\left(X^{n}\right), g\left(Y^{n}\right) \\ \text{almost uniform on} \\ \left[1:e^{nR}\right]\right) \end{array}$	Finite $(\mathcal{X}, \mathcal{Y})$ : For $(X^n, Y^n) \sim \pi_{XY}^n$ and any $\varepsilon \in (0, 1)$ , $\max_{f,g} R = C_{\mathrm{GK}}(\pi_{XY}) := \max_{\hat{f}, \hat{g}: \hat{f}(X) = \hat{g}(Y)} H(\hat{f}(X))$ [13]
A Variant of Gács-Körner's CI [13]	Variable	Exact $(f(X^n) = g(Y^n) \text{ a.s.})$	Finite $(\mathcal{X}, \mathcal{Y})$ : For $(X^n, Y^n) \sim \pi_{XY}^n$ , $\max_{f,g} \frac{1}{n} H(f(X^n)) = \frac{1}{n} C_{\text{GK}}(\pi_{XY}^n) = C_{\text{GK}}(\pi_{XY}) $ [13]

 TABLE I

 SUMMARY OF VARIOUS COMMON INFORMATIONS.

Here [P] refers to the present paper. "\*" means that some regular conditions are required.  $\Gamma_{\alpha}^{\text{LB}}(\pi_{XY})$  and  $\Gamma_{\alpha}^{\text{UB}}(\pi_{XY})$  were defined in [4]. In the rightmost box in the rows of " $\infty$ -Rényi CI" and "Exact CI", Points 1) and 2) hold for unnormalized version of  $\infty$ -Rényi CI (i.e., with  $D_{\infty}$  measure). Points 3) and 4) hold for both normalized and unnormalized versions of  $\infty$ -Rényi CI.

rule" for coupling (Lemma 9), the distributed Rényicovering lemmas (for sources with finite alphabets and Gaussian sources) (Lemmas 7 and 17), and a lemma on the estimation of conditional mutual information (Lemma 15).

#### **B.** Notations

We use  $P_X$  to denote the probability distribution of a random variable X. For brevity, we also use  $P_X(x)$  to denote the corresponding probability mass function (pmf) for discrete distributions, and the corresponding probability density function (pdf) for continuous distributions. This will also be denoted as P(x) (when the random variable X is clear from the context). We also use  $\pi_X, \tilde{P}_X, \tilde{P}_X$  and  $Q_X$  to denote various probability distributions on alphabet  $\mathcal{X}$ . The set of probability measures on  $\mathcal{X}$  is denoted as  $\mathcal{P}(\mathcal{X})$ , and the set of conditional probability measures on  $\mathcal{Y}$  given a variable in  $\mathcal{X}$  is denoted as  $\mathcal{P}(\mathcal{Y}|\mathcal{X}) := \{ P_{Y|X} : P_{Y|X}(\cdot|x) \in \mathcal{P}(\mathcal{Y}), x \in \mathcal{X} \}.$ Furthermore, the support of a distribution  $P \in \mathcal{P}(\mathcal{X})$  is denoted as  $\operatorname{supp}(P) = \{x \in \mathcal{X} : P(x) > 0\}$ . For two distributions P and Q defined on the same measurable space, we use  $P \ll Q$  to denote that P is absolutely continuous with respect to Q. If  $P \ll Q$ , we use  $\frac{dP}{dQ}$  to denote the Radon–Nikodym derivative of P with respect to Q.

The TV distance between two probability mass functions P and Q with a common alphabet  $\mathcal{X}$  is defined as

$$|P - Q| := \frac{1}{2} \sum_{x \in \mathcal{X}} |P(x) - Q(x)|.$$
(11)

We use  $T_{x^n}(x) := \frac{1}{n} \sum_{i=1}^n 1\{x_i = x\}$  to denote the type (empirical distribution) of a sequence  $x^n$ ,  $T_X$  and  $V_{Y|X}$  to re-

spectively denote a type of sequences in  $\mathcal{X}^n$  and a conditional type of sequences in  $\mathcal{Y}^n$  (given a sequence  $x^n \in \mathcal{X}^n$ ). For a type  $T_X$ , the type class (set of sequences having the same type  $T_X$ ) is denoted by  $\mathcal{T}_{T_X}$ . For a conditional type  $V_{Y|X}$  and a sequence  $x^n$ , the  $V_{Y|X}$ -shell of  $x^n$  (the set of  $y^n$  sequences having the same conditional type  $V_{Y|X}$  given  $x^n$ ) is denoted by  $\mathcal{T}_{V_{Y|X}}(x^n)$ . For brevity, sometimes we use T(x, y) to denote the joint distributions T(x)V(y|x) or T(y)V(x|y).

For  $X \sim P_X$ , we denote the *entropy* of X as

$$H_P(X) = H(P_X) := -\sum_{x \in \text{supp}(P_X)} P_X(x) \log P_X(x).$$
 (12)

For  $(X, Y) \sim P_{XY}$ , we denote the *conditional entropy* of X given Y as

$$H_P(X|Y) = H(P_{X|Y}|P_Y)$$
(13)

$$:= -\sum_{x,y} P_{XY}(x,y) \log P_{X|Y}(x|y).$$
(14)

For  $(X, Y) \sim P_{XY}$ , we denote the *mutual information* between X and Y as

$$I_P(X;Y) = H_P(X) - H_P(X|Y).$$
 (15)

For brevity and when entropies, conditional entropies, and mutual informations are computed respect to a distribution denoted by "P", we omit the subscript and denote them respectively as H(X), H(X|W), and I(X;Y) instead of the more verbose  $H_P(X)$ ,  $H_P(X|W)$ , and  $I_P(X;Y)$ . The  $\epsilon$ -strongly,  $\epsilon$ -weakly, and  $\epsilon$ -unified typical sets [18]– [21] of  $P_X$  are respectively denoted as

$$\mathcal{T}_{\epsilon}^{(n)}(P_X) := \left\{ x^n \in \mathcal{X}^n : |T_{x^n}(x) - P_X(x)| \le \epsilon P_X(x), \forall x \in \mathcal{X} \right\},$$
(16)

$$\mathcal{A}_{\epsilon}^{(n)}(P_X) := \left\{ x^n \in \mathcal{X}^n : \left| -\frac{1}{n} \log P_X^n(x^n) - H(P_X) \right| \le \epsilon \right\}, \quad (17)$$

$$\mathcal{U}_{\epsilon}^{(n)}\left(P_{X}\right) := \left\{ x^{n} \in \mathcal{X}^{n} : \\ D\left(T_{x^{n}} \| P_{X}\right) + \left| H\left(T_{x^{n}}\right) - H\left(P_{X}\right) \right| \leq \epsilon \right\}.$$
(18)

Note that  $\mathcal{T}_{\epsilon}^{(n)}(P_X)$  only applies to sources with finite alphabets, and  $\mathcal{U}_{\epsilon}^{(n)}(P_X)$  applies to sources with countable alphabets. For  $\mathcal{A}_{\epsilon}^{(n)}(P_X)$ , if  $P_X$  is an absolutely continuous distribution, in (17),  $P_X^n(x^n)$  and  $H(P_X)$  are respectively replaced with the corresponding pdf and differential entropy. The corresponding jointly typical sets are defined similarly. The conditionally  $\epsilon$ -strongly typical set of  $P_{XY}$  is denoted as

$$\mathcal{T}_{\epsilon}^{(n)}(P_{XY}|x^n) := \left\{ y^n \in \mathcal{Y}^n : (x^n, y^n) \in \mathcal{T}_{\epsilon}^{(n)}(P_{XY}) \right\},\tag{19}$$

and the conditionally  $\epsilon$ -weakly and  $\epsilon$ -unified typical sets are defined similarly. For brevity, sometimes we write  $\mathcal{T}_{\epsilon}^{(n)}(P_X), \mathcal{A}_{\epsilon}^{(n)}(P_X)$  and  $\mathcal{U}_{\epsilon}^{(n)}(P_X)$  as  $\mathcal{T}_{\epsilon}^{(n)}, \mathcal{A}_{\epsilon}^{(n)}$  and  $\mathcal{U}_{\epsilon}^{(n)}$ , respectively.

For distributions  $P_X, Q_X \in \mathcal{P}(\mathcal{X})$ , the *relative entropy* and the *Rényi divergence of order*  $1 + s \in (0, 1) \cup (1, \infty)$  are respectively defined as<sup>1</sup>

$$D(P_X || Q_X) := \sum_{x \in \text{supp}(P_X)} P_X(x) \log \frac{P_X(x)}{Q_X(x)}$$
(20)

$$D_{1+s}(P_X \| Q_X) := \frac{1}{s} \log \sum_{x \in \text{supp}(P_X)} P_X(x)^{1+s} Q_X(x)^{-s},$$
(21)

and the conditional versions are respectively defined as

$$D(P_{Y|X} \| Q_{Y|X} | P_X) := D(P_X P_{Y|X} \| P_X Q_{Y|X})$$
(22)

$$D_{1+s}(P_{Y|X} \| Q_{Y|X} | P_X) := D_{1+s}(P_X P_{Y|X} \| P_X Q_{Y|X}),$$
(23)

where the summations in (20) and (21) are taken over the elements in  $\operatorname{supp}(P_X)$ . The Rényi divergence of order  $1 + s \in \{0, \infty\}$  is defined by the continuous extensions of  $D_{1+s}$ . The Rényi divergence of order 1 is defined as  $D_1(P_X || Q_X) := \lim_{s \uparrow 0} D_{1+s}(P_X || Q_X)$ . Throughout, log and exp are to the natural base e and  $s \ge -1$ . It is known that  $D_1(P_X || Q_X) = D(P_X || Q_X)$  so a special case of the Rényi divergence (or the conditional version) is the usual relative entropy (or the conditional version). The Rényi divergence of order  $\infty$  satisfies

$$D_{\infty}(P_X \| Q_X) = \log \sup_{x \in \operatorname{supp}(P_X)} \frac{P_X(x)}{Q_X(x)}.$$
 (24)

If  $\mathcal{X}$  is a countable alphabet or  $\mathbb{R}$  and we replace  $Q_X$  by respectively the counting or the Lebesgue measures, then the Rényi divergence  $D_{1+s}(P_X || Q_X)$  of order  $1 + s \in [0, \infty]$ reduces to the Rényi entropy  $-H_{1+s}(P_X)$  of the same order.

Denote the coupling sets of  $(P_X, P_Y)$  and  $(P_{X|W}, P_{Y|W})$  respectively as

$$C(P_X, P_Y) := \{Q_{XY} \in \mathcal{P}(\mathcal{X} \times \mathcal{Y}) :$$

$$Q_X = P_X, Q_Y = P_Y\}$$

$$C(P_{X|W}, P_{Y|W}) := \{Q_{XY|W} \in \mathcal{P}(\mathcal{X} \times \mathcal{Y}|\mathcal{W}) :$$

$$Q_{X|W} = P_{X|W}, Q_{Y|W} = P_{Y|W}\}.$$
(26)

For  $i, j \in \mathbb{Z}$ , and  $i \leq j$ , we define  $[i : j] := \{i, i+1, \ldots, j\}$ . Given a number  $a \in [0, 1]$ , we define  $\overline{a} = 1 - a$ . Define  $[x]^+ = \max\{x, 0\}$ . Denote  $\mathcal{A}^c$  as the complement of the set  $\mathcal{A}$ . Finally, we write  $f(n) \sim g(n)$  if  $\lim_{n \to \infty} \frac{f(n)}{g(n)} = 1$ .

#### **II. PROBLEM FORMULATIONS**

#### A. Rényi Common Information

Consider the distributed source simulation setup depicted in Fig. I. Two terminals both have access to a uniformly distributed common randomness  $M_n$ . Given a target distribution  $\pi_{XY}$ , one of terminals uses  $M_n$  and his own local randomness to generate  $X^n$  and the other one uses  $M_n$  and his own local randomness to generate  $Y^n$  such that the the generated (or synthesized) distribution  $P_{X^nY^n}$  is close to the product distribution  $\pi_{XY}^n$  under Rényi divergence measures. We wish to find the limit on the least amount of common randomness satisfying such a requirement. More specifically, given a target distribution  $\pi_{XY}$ , we wish to minimize the alphabet size of a random variable  $M_n$  that is uniformly distributed over<sup>2</sup>  $\mathcal{M}_n := [1 : e^{nR}]$  (*R* is a positive number known as the *rate*), such that the generated (or synthesized) distribution

$$P_{X^{n}Y^{n}}(x^{n}, y^{n})$$
  
:=  $\frac{1}{|\mathcal{M}_{n}|} \sum_{m \in \mathcal{M}_{n}} P_{X^{n}|M_{n}}(x^{n}|m) P_{Y^{n}|M_{n}}(y^{n}|m)$  (27)

forms a good approximation to the product distribution  $\pi_{XY}^n$ .

**Definition 1.** A fixed-length (n, R)-code consists of a pair of random mappings  $P_{X^n|W_n} : \mathcal{W}_n \to \mathcal{X}^n, P_{Y^n|W_n} : \mathcal{W}_n \to \mathcal{Y}^n$ for some countable set  $\mathcal{W}_n$  such that  $\frac{1}{n} \log |\mathcal{W}_n| \leq R$ .

In the Rényi common information problem [3], the unnormalized Rényi divergence  $D_{1+s}(P_{X^nY^n} \| \pi_{XY}^n)$  and the normalized Rényi divergence  $\frac{1}{n}D_{1+s}(P_{X^nY^n} \| \pi_{XY}^n)$  are adopted to measure the discrepancy between  $P_{X^nY^n}$  and  $\pi_{XY}^n$ .

**Definition 2.** [3] The unnormalized and normalized Rényi common informations  $T_{1+s}(\pi_{XY})$  and  $\widetilde{T}_{1+s}(\pi_{XY})$  of order

<sup>&</sup>lt;sup>1</sup>When the alphabet  $\mathcal{X}$  is uncountable, it is understood that  $\frac{P_X}{Q_X}$  should be replaced by the Radon-Nikodym derivative  $\frac{dP_X}{dQ_X}$  of  $P_X$  respect to  $Q_X$ . If  $P_X$  is not absolutely continuous respect to  $Q_X$ , then as assumed conventionally, the relative entropy and the Rényi divergence of order  $1 + s \in (1, \infty)$  are defined as  $\infty$ . (Note that for this case, the Rényi divergence of order  $1 + s \in (0, 1)$  is well-defined.)

<sup>&</sup>lt;sup>2</sup>For simplicity, we assume that  $e^{nR}$  and similar expressions are integers.

 $1 + s \in [0, \infty]$  between two sources with joint distribution  $\pi_{XY}$  are defined as

$$T_{1+s}(\pi_{XY})$$
  
:= inf  $\left\{ \begin{array}{c} R : \exists \{ \text{fixed-length } (n, R) \text{ code} \}_{n=1}^{\infty} \text{ s.t.} \\ \lim_{n \to \infty} D_{1+s}(P_{X^nY^n} \| \pi_{XY}^n) = 0 \end{array} \right\}$  (28)

and

$$T_{1+s}(\pi_{XY})$$

$$:= \inf \left\{ \begin{array}{c} R: \exists \{ \text{fixed-length } (n,R) \text{ code} \}_{n=1}^{\infty} \text{ s.t.} \\ \lim_{n \to \infty} \frac{1}{n} D_{1+s}(P_{X^nY^n} \| \pi_{XY}^n) = 0 \end{array} \right\}.$$
(29)

It is clear that

$$\tilde{T}_{1+s}(\pi_{XY}) \le T_{1+s}(\pi_{XY}).$$
 (30)

If s = 0, then the unnormalized and normalized Rényi common informations respectively reduce to the unnormalized and normalized versions of Wyner's common informations [2].

#### **B.** Exact Common Information

In the formulation of the Rényi common information problem, fixed-length block codes and approximate generation of the target distribution  $\pi_{XY}^n$  are assumed. In contrast, in the exact common information problem [6], KLE considered variable-length codes and exact generation of  $\pi_{XY}^n$ . The target is also to find the limit on the least amount of common randomness satisfying such a requirement, but the amount here is quantified in term of per-letter expected codeword length, rather than the exponent of alphabet size described in the previous subsection.

Define  $\{0,1\}^* := \bigcup_{n \ge 1} \{0,1\}^n$  as the set of finite-length strings of symbols from a binary alphabet  $\{0,1\}$ . Denote the alphabet of the common random variable  $W_n$  as  $W_n$ , which can be any countable set (without loss of generality, one can assume  $W_n = \mathbb{N}$ ). Consider a prefix-free code  $f : W_n \to$  $\{0,1\}^*$ . Then for each symbol  $w \in W_n$  and the code f, let  $\ell_f(w)$  denote the length of the codeword f(w).

**Definition 3.** The expected codeword length  $L_f(W_n)$  for compressing the random variable  $W_n$  by a uniquely decodable code f is defined as  $L_f(W_n) := \mathbb{E} [\ell_f(W_n)].$ 

**Definition 4.** A variable-length (n, R)-code consists of  $(P_{W_n}, f, P_{X^n|W_n}, P_{Y^n|W_n})$ , i.e., consists of a distribution  $P_{W_n}$  on for some countable set  $\mathcal{W}_n$ , a pair of random mappings  $P_{X^n|W_n} : \mathcal{W}_n \to \mathcal{X}^n, P_{Y^n|W_n} : \mathcal{W}_n \to \mathcal{Y}^n$ , and a prefix-free code  $f : \mathcal{W}_n \to \{0, 1\}^*$  such that the expected codeword length for  $W_n$  satisfies  $L_f(W_n)/n \leq R$ .

By using variable-length codes,  $W_n$  is transmitted to two terminals with error free. The generated (or synthesized) distribution for such setting is

$$P_{X^{n}Y^{n}}(x^{n}, y^{n}) = \sum_{w \in \mathcal{W}_{n}} P_{W_{n}}(w) P_{X^{n}|W_{n}}(x^{n}|w) P_{Y^{n}|W_{n}}(y^{n}|w), \quad (31)$$

which is required to be  $\pi_{XY}^n$  exactly.

**Definition 5.** [6] The exact common information  $T_{\text{Exact}}(\pi_{XY})$  between two sources with joint distribution  $\pi_{XY}$  is defined as the minimum asymptotic rate required to ensure  $P_{X^nY^n} = \pi_{XY}^n$  for all  $n \ge 1$ , i.e.,

$$T_{\text{Exact}}(\pi_{XY}) = \inf \left\{ \begin{array}{c} R : \exists \{ \text{variable-length } (n, R^{(n)}) \text{ code} \}_{n=1}^{\infty} \text{ s.t.} \\ P_{X^nY^n} = \pi_{XY}^n, \forall n \ge 1 \\ R \ge \limsup_{n \to \infty} R^{(n)} \end{array} \right\}$$
(32)

By observing that the expected codeword length  $L_f(W_n)$  satisfies  $H(W_n) \leq L_f(W_n) < H(W_n) + 1$ , it is easy to verify that  $\frac{1}{n} (L_f(W_n) - H(W_n)) \rightarrow 0$  as  $n \rightarrow \infty$ . Based on such an argument, KLE [6] provided the following multi-letter characterization of the exact common information:

$$T_{\text{Exact}}(\pi_{XY}) = \lim_{n \to \infty} \frac{1}{n} \min_{P_W P_{X^n | W} P_{Y^n | W} : P_{X^n Y^n} = \pi_{XY}^n} H(W).$$
(33)

Hence a variable-length synthesis code can be represented by  $(P_{W_n}, P_{X^n|W_n}, P_{Y^n|W_n})$ , where the dependence on the variable-length compression code f is omitted.

#### III. MAIN RESULTS FOR SOURCES WITH FINITE Alphabets

#### A. Equivalence and Multi-letter Characterization

We first establish the equivalence between the exact and  $\infty$ -Rényi common informations, and characterize them using a multi-letter expression. The proof of Theorem 1 is given in Appendix A.

**Theorem 1** (Equivalence). For a source with distribution  $\pi_{XY}$  defined on a finite alphabet,

$$T_{\text{Exact}}(\pi_{XY}) = T_{\infty}(\pi_{XY}) = \lim_{n \to \infty} \frac{1}{n} \Gamma(\pi_{XY}^n), \qquad (34)$$

where<sup>3</sup>

$$\Gamma(\pi_{XY}^{n}) := \inf_{\substack{P_{W}P_{X^{n}|W}P_{Y^{n}|W}: \\ P_{X^{n}Y^{n}} = \pi_{XY}^{n} \\ C(P_{X^{n}Y^{n}|W}, P_{Y^{n}|W}) \\ -\sum_{w} P(w) \sum_{x^{n}, y^{n}} Q(x^{n}, y^{n}|w) \log \pi^{n} (x^{n}, y^{n}) .$$
(35)

*Remark* 2. By using a proof similar to that for the converse part of Theorem 1, one can show the following lower bound on the normalized  $\infty$ -Rényi common information.

$$\widetilde{T}_{\infty}(\pi_{XY}) \ge \lim_{\epsilon \downarrow 0} \lim_{n \to \infty} \frac{1}{n} \Gamma_{\epsilon}(\pi_{XY}^n),$$
(36)

<sup>3</sup>Note that per Subsection I-B, the conditional entropy  $H(X^nY^n|W)$  is computed with respect to  $P_WP_{X^n|W}P_{Y^n|W}$ . Hence in fact,  $H(X^nY^n|W) = H(X^n|W) + H(Y^n|W)$ .

where

$$\Gamma_{\epsilon}(\pi_{XY}^{n}) := \inf_{\substack{P_{W}P_{X^{n}|W}P_{Y^{n}|W}: \quad Q_{X^{n}Y^{n}|W} \in \\ \frac{1}{n}D_{\infty}(P_{X^{n}Y^{n}} \| \pi_{XY}^{n}) \le \epsilon \ C(P_{X^{n}|W}, P_{Y^{n}|W})}{-\sum_{w} P(w) \sum_{x^{n}, y^{n}} Q(x^{n}, y^{n}|w) \log \pi^{n} (x^{n}, y^{n}) . \quad (37)$$

It is easy to verify that given  $\epsilon > 0$ ,  $\Gamma_{\epsilon}(\pi_{XY}^n)$  is subadditive in n, i.e.,  $\Gamma_{\epsilon}(\pi_{XY}^n) \leq \Gamma_{\epsilon}(\pi_{XY}^{n_1}) + \Gamma_{\epsilon}(\pi_{XY}^{n_2})$  for all  $n_1 + n_2 = n$ . Hence the limit in (36) exists.

*Remark* 3. A similar equivalence as the first equality in (34) has been found by Kumar, Li, and El Gamal in [6, Remark on Page 164]. They showed that the exact common information is equal to a variant of the  $\infty$ -Rényi common information in which variable-length codes are allowed. Our equivalence enhances their equivalence for the direction of  $T_{\text{Exact}}(\pi_{XY}) \ge T_{\infty}(\pi_{XY})$ . Such a difference enables us to derive the converse part of the multiletter characterization given in (34).

#### B. Single-letter Bounds

Define the maximal cross-entropy over couplings  $C(P_{X|W=w}, P_{Y|W=w'})$  as<sup>4</sup>

$$\mathcal{H}(P_{X|W=w}, P_{Y|W=w'} \| \pi_{XY}) \\ := \sup_{Q_{XY} \in C(P_{X|W=w}, P_{Y|W=w'})} \sum_{x,y} Q(x,y) \log \frac{1}{\pi (x,y)}.$$
(38)

Here for the finite alphabet case, the supremum is in fact a maximum.

Define

$$\Gamma^{\text{UB}}(\pi_{XY}) := \Gamma(\pi_{XY}) \\
= \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} \left\{ -H(XY|W) + \sum_{w} P(w) \mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY}) \right\},$$
(39)

and

$$\Gamma^{\text{LB}}(\pi_{XY}) := \inf_{\substack{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY} \\ + \inf_{\substack{Q_{WW'} \in C(P_W, P_W) \\ w, w'}} \sum_{w, w'} Q(w, w')} \times \mathcal{H}(P_{X|W=w}, P_{Y|W=w'} \| \pi_{XY}) \Big\}.$$
(40)

For (39), it suffices to restrict the size of the alphabet of W such that  $|W| \leq |\mathcal{X}||\mathcal{Y}|$ . This is because

$$-H(XY|W) + \sum_{w} P(w)\mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY})$$

$$(41)$$

is a linear function of  $P_W$ . Hence by standard cardinality bounding techniques (e.g., the support lemma in [18, Appendix

C]), there exists an optimal distribution  $P_W P_{X|W} P_{Y|W}$  with  $P_{XY} = \pi_{XY}$  and  $|\operatorname{supp}(W)| \leq |\mathcal{X}||\mathcal{Y}|$  attaining the minimization in (39).

By utilizing the multi-letter expression in Theorem 1, we provide single-letter lower and upper bounds for the exact and  $\infty$ -Rényi common informations. The proof of Theorem 2 is given in Appendix B.

**Theorem 2** (Single-letter Bounds). The exact and  $\infty$ -Rényi common informations for a source with distribution  $\pi_{XY}$  defined on a finite alphabet satisfy

$$\max\left\{\Gamma^{\mathrm{LB}}(\pi_{XY}), C_{\mathsf{Wyner}}(\pi_{XY})\right\} \le \widetilde{T}_{\infty}(\pi_{XY}) \tag{42}$$

$$\leq T_{\infty}(\pi_{XY})$$
 (43)

$$=T_{\mathrm{Exact}}(\pi_{XY})$$
 (44)

$$\leq \Gamma^{\rm UB}(\pi_{XY}).$$
 (45)

Note that the only difference between the upper and lower bounds is that in the lower bound, the minimization operation is taken over all couplings of  $(P_W, P_W)$ , but in the upper bound, it is not (or equivalently, the expectation in (39) can be seen as being taken under the equality coupling of  $(P_W, P_W)$ , namely  $P_W(w) 1\{w' = w\}$ ). The upper bound  $\Gamma^{UB}(\pi_{XY})$  and lower bound  $\Gamma^{LB}(\pi_{XY})$  are consistent with the bounds for  $\alpha$ -Rényi common information for  $\alpha \in [0, \infty]$  [3], [4].

The  $\infty$ -Rényi common information code we adopt in the proof is a truncated i.i.d. code. For such a code, the codewords are independent and each codeword is drawn according to a distribution  $P_{W^n}$  which is generated by truncating a product distribution  $Q_W^n$  onto some (strongly) typical set. Truncated i.i.d. codes are rather useful (i.e., strictly better than i.i.d. codes without truncation) for  $\infty$ -Rényi-approximate synthesis (but achieve the same performance as i.i.d. codes for Wyner's synthesis, i.e., 1-Rényi-approximate synthesis). This follows from the following argument. Observe that for both  $\infty$ -Rényiapproximate synthesis and Wyner's synthesis,  $X^n \to W_n \to$  $Y^n$  forms a Markov chain. Hence given  $W_n = w$ , the support of  $P_{X^n|W_n}(\cdot|w) P_{Y^n|W_n}(\cdot|w)$  is a product set, which in turn implies that the support of  $P_{X^nY^n}$  is the union of a family of product sets. Such a requirement leads to the fact that the support of  $P_{X^nY^n}$  includes not only a jointly typical set, but also other joint type classes, which is termed by us as the type overflow phenomenon. Wyner's synthesis (under the relative entropy measure) only requires the sequences in a typical set to be well-simulated. However,  $\infty$ -Rényi-approximate synthesis requires all the sequences in the support of  $P_{X^nY^n}$  to be well-simulated. Hence the type overflow phenomenon does not affect Wyner's synthesis asymptotically, but plays a critical role in minimizing the rate of  $\infty$ -Rényi-approximate synthesis (or exact synthesis). Truncated i.i.d. coding is an efficient approach to control the possible types of the output sequence of a code (or more precisely, to mitigate the effects of type overflow). Furthermore, truncated i.i.d. codes have also been used by the present authors [3], [4], [25] to study  $\alpha$ -Rényi common informations, and by Vellambi and Kliewer [9], [26] to study sufficient conditions for equality of the exact and Wyner's common informations.

The maximal cross-entropy in (38) has the following intuitive interpretation. Consider a joint distribution  $\pi_{XY}$ , a

<sup>&</sup>lt;sup>4</sup>Note that the maximization in (38) is an optimal transport problem [22], [23]. Hence Kantorovich duality can be used to bound the maximal crossentropy if it is required. For more details about the maximal cross-entropy, please refer to [24, Section III.A].

pair of distributions  $(P_X, P_Y)$ , and a sequence of pairs of types  $\{(T_X^{(n)}, T_Y^{(n)}) \in \mathcal{P}_n(\mathcal{X}) \times \mathcal{P}_n(\mathcal{Y})\}_{n \in \mathbb{N}}$  such that  $(T_X^{(n)}, T_Y^{(n)}) \to (P_X, P_Y)$  as  $n \to \infty$ . The minimum of the exponents of probabilities  $\pi_{XY}^n(x^n, y^n)$  such that  $T_{x^n} = T_X^{(n)}, T_{y^n} = T_Y^{(n)}$  satisfy that

$$\lim_{n \to \infty} \min_{\substack{(x^n, y^n): \\ T_x = T_X^{(n)}, \\ T_y = T_Y^{(n)}, \\ x_y = T_Y^{(n)}, \\ x_y = T_Y^{(n)}, \\ T_x = T_X^{(n)}, \\ T_y = T_X^{(n)}, \\ T_y = T_Y^{(n)}, \\ T_y = T_Y^{(n)}, \\ x_y = T_Y^{(n)}, \\ (46)$$

Based on the type overflow argument and the intuitive explanation of the maximal cross-entropy given above, our bounds are easy to comprehend intuitively. The exact synthesis requires that there exists a sequence of variable-length codes with asymptotic rate R satisfying  $\frac{P_{Y^n|X^n}(y^n|x^n)}{\pi_{Y|X}^n(y^n|x^n)} = 1$  for all  $(x^n, y^n) \in \mathcal{X}^n \times \mathcal{Y}^n$ . By using the mixture decomposition technique, the exact synthesis problem can be relaxed to the  $\infty$ -Rényi-approximate synthesis problem, which requires that there exists a sequence of fixed-length codes with asymptotic rate R satisfying

$$\frac{P_{X^nY^n}(x^n, y^n)}{\pi_{XY}^n(x^n, y^n)} \le 1 + o(1) \tag{48}$$

for all  $(x^n, y^n) \in \text{supp}(P_{X^nY^n})$ ; see Lemma 4. By using truncated i.i.d. codes, to mitigate the effect of type overflow we can restrict  $(W^n, X^n) \in \mathcal{T}_{\epsilon}^{(n)}(P_{WX})$  and  $(W^n, Y^n) \in$  $\mathcal{T}_{\epsilon}^{(n)}(P_{WY})$ . Suppose that  $M_n$  is the message for  $\infty$ -Rényiapproximate synthesis. Then for sufficiently large n and sufficiently small  $\epsilon$ ,

$$P_{X^{n}Y^{n}}(x^{n}, y^{n}) \approx \sum P_{M_{n}}(m)P_{X|W}^{n}(x^{n}|w^{n}(m))P_{Y|W}^{n}(y^{n}|w^{n}(m)) \quad (49)$$

$$\approx N(x^{n}, y^{n})e^{-nR}e^{-nH(X|W)}e^{-nH(Y|W)},$$
(50)

where  $N(x^n, y^n)$  denotes the number of codewords  $w^n(m)$  that cover  $x^n$  and  $y^n$  (i.e., that are jointly typical with  $x^n$  and jointly typical with  $y^n$ ). On the other hand,

$$\min_{\substack{(x^n, y^n) \in \operatorname{supp}(P_{X^n Y^n})}} \pi^n_{XY}(x^n, y^n)$$
  
$$\approx \min_{\substack{(w^n, x^n, y^n): T_{w^n x^n} \approx P_{WX}, T_{w^n y^n} \approx P_{WY}}} \pi^n_{XY}(x^n, y^n) \quad (51)$$

$$\approx e^{-n\sum_{w} P_W(w)\mathcal{H}(P_{X|W=w}, P_{Y|W=w}\|\pi_{XY})}.$$
(52)

Substituting (50) and (52) into (48) and observing that  $N(x^n, y^n) \ge 1$  for  $(x^n, y^n) \in \operatorname{supp}(P_{X^nY^n})$ , we obtain

$$R \gtrsim -H(XY|W) + \sum_{w} P(w)\mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY}).$$
(53)

Taking the minimum over all distributions  $P_W P_{X|W} P_{Y|W}$ such that  $P_{XY} = \pi_{XY}$ , we obtain the upper bound  $\Gamma^{\text{UB}}(\pi_{XY})$ . We make this argument precise in Appendix B.

#### C. Doubly Symmetric Binary Sources

A doubly symmetric binary source (DSBS) is a source (X, Y) with distribution

$$\pi_{XY} := \begin{bmatrix} \alpha_0 & \beta_0 \\ \beta_0 & \alpha_0 \end{bmatrix}$$
(54)

where  $\alpha_0 = \frac{1-p}{2}, \beta_0 = \frac{p}{2}$  with  $p \in (0, \frac{1}{2})$ . This is equivalent to  $X \sim \text{Bern}(\frac{1}{2})$  and  $Y = X \oplus E$  with  $E \sim \text{Bern}(p)$ independent of X; or  $X = W \oplus A$  and  $Y = W \oplus B$  with  $W \sim \text{Bern}(\frac{1}{2}), A \sim \text{Bern}(a)$ , and  $B \sim \text{Bern}(a)$  mutually independent, where  $a := \frac{1-\sqrt{1-2p}}{2} \in (0, \frac{1}{2})$  or equivalently,  $\alpha_0 = \frac{1}{2} \left(a^2 + (1-a)^2\right), \beta_0 = a(1-a)$ . Here we do not lose any generality by restricting p or  $a \in (0, \frac{1}{2})$ , since otherwise, we can set  $X \oplus 1$  to X.

By utilizing the lower and upper bounds in Theorem 2, we completely characterize the exact and  $\infty$ -Rényi common informations for DSBSes. The proof of Theorem 3 is given in Appendix C.

**Theorem 3.** For a DSBS (X, Y) with distribution  $\pi_{XY}$  given in (54),

$$\widetilde{T}_{\infty}(\pi_{XY}) = T_{\infty}(\pi_{XY}) = T_{\text{Exact}}(\pi_{XY}) = -2H_2(a) - (1 - 2a) \log \left[\frac{1}{2} \left(a^2 + (1 - a)^2\right)\right] - 2a \log \left[a(1 - a)\right],$$
(55)

where

$$H_2(a) := -a \log a - (1-a) \log(1-a)$$
 (56)

denotes the binary entropy function.

**Corollary 1.** For a DSBS (X, Y) with distribution  $\pi_{XY}$  given in (54),

$$\widetilde{T}_{\infty}(\pi_{XY}) = T_{\infty}(\pi_{XY}) = T_{\text{Exact}}(\pi_{XY}) > C_{\text{Wyner}}(\pi_{XY})$$
(57)

for the parameter  $a \in (0, \frac{1}{2})$ .

*Remark* 4. For this case, the exact common information is strictly larger than Wyner's common information. This answers an open problem posed by KLE [6].

Proof: For DSBSes, Wyner [2] showed that

$$T_1(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY})$$
  
=  $-2H_2(a) - (a^2 + (1-a)^2) \log \left[\frac{1}{2}(a^2 + (1-a)^2)\right]$   
 $- 2a(1-a) \log [a(1-a)].$  (58)

Hence

=

$$T_{\infty}(\pi_{XY}) - C_{\text{Wyner}}(\pi_{XY})$$
  
=  $\left(\left(a^{2} + (1-a)^{2}\right) - (1-2a)\right)\log\left[\frac{1}{2}\left(a^{2} + (1-a)^{2}\right)\right]$   
+  $\left(2a(1-a) - 2a\right)\log\left[a(1-a)\right]$  (59)

$$2a^{2}\log\left|\frac{\frac{1}{2}\left(a^{2}+(1-a)^{2}\right)}{a(1-a)}\right| > 0.$$
(60)

We obtain the desired result.

The exact,  $\infty$ -Rényi, and Wyner's common informations for DSBSes are illustrated in Fig. 2.



Fig. 2. Illustrations of the exact and  $\infty$ -Rényi common informations (55) and Wyner's common information (58) for DSBSes (X, Y) such that  $X \sim \text{Bern}(\frac{1}{2})$  and  $Y = X \oplus E$  with  $E \sim \text{Bern}(p)$  independent of X.

### D. Sufficient Conditions for Equality of Exact and Wyner's Common Informations

In Corollary 1, we showed that for a DSBS, the exact common information is strictly larger than Wyner's common information. Now we study sufficient conditions for equality of exact and Wyner's common informations. Obviously, if  $\Gamma^{\rm UB}(\pi_{XY}) = C_{\rm Wyner}(\pi_{XY})$ , then the exact and Wyner's common informations are equal. We first introduce a condition on  $\pi_{XY}$ .

Condition (\*): There exists some optimal distribution  $P_W P_{X|W} P_{Y|W}$  attaining  $C_{Wyner}(\pi_{XY})$  such that for any  $w \in \text{supp}(P_W)$ ,  $\pi_{XY}$  when restricted to  $\mathcal{A}_w := \text{supp}(P_{X|W=w}) \times \text{supp}(P_{Y|W=w})$  is a product distribution, i.e.,  $\pi_{XY}(\cdot|\mathcal{A}_w)$  is a product distribution for each  $w \in \text{supp}(P_W)$ .

**Theorem 4.** If  $\pi_{XY}$  satisfies Condition (\*), then the exact and Wyner's common informations are equal, i.e.,

$$T_{\text{Exact}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY}). \tag{61}$$

*Remark* 5. Theorem 4 generalizes the sufficient conditions given in [6], [9], [26].

Theorem 4 follows from the following lemma.

**Lemma 1.**  $\Gamma^{\text{UB}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY})$  if and only if  $\pi_{XY}$  satisfies Condition (\*).

*Remark* 6. Lemma 1 implies that if the upper bound  $\Gamma^{\text{UB}}(\pi_{XY})$  is tight for the exact common information (i.e.,  $T_{\text{Exact}}(\pi_{XY}) = \Gamma^{\text{UB}}(\pi_{XY})$ ), then Condition (\*) is necessary and sufficient for  $T_{\text{Exact}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY})$ .

*Proof:* "If" Part: Suppose that  $\pi_{XY}$  satisfies Condition (\*). Then by [24, Proposition 2], we obtain that for any  $w \in \text{supp}(P_W)$ ,

$$\mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY})$$
  
=  $\sum_{x,y} P(x|w) P(y|w) \log \frac{1}{\pi(x,y)}.$  (62)

After taking the expectation respect to  $P_W$ , we obtain

$$\sum_{w} P(w) \mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY}) = H(XY).$$
(63)

Therefore, substituting the distribution  $P_W P_{X|W} P_{Y|W}$  into  $\Gamma^{\text{UB}}(\pi_{XY})$ , we obtain that  $\Gamma^{\text{UB}}(\pi_{XY}) \leq C_{\text{Wyner}}(\pi_{XY})$ . Since  $\Gamma^{\text{UB}}(\pi_{XY}) \geq C_{\text{Wyner}}(\pi_{XY})$ , we obtain that  $\Gamma^{\text{UB}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY})$ .

"Only If" Part: Suppose that  $\Gamma^{UB}(\pi_{XY}) = C_{Wyner}(\pi_{XY})$ . For а distribution  $\pi_{XY}$ , denote  $P_W P_{X|W} P_{Y|W}$  $\Gamma^{\rm UB}(\pi_{XY}).$ as an optimal distribution attaining Then we have that for any w $\in$  $\operatorname{supp}(P_W),$  $\operatorname{supp}\left(P_{X|W=w}\right) \times \operatorname{supp}\left(P_{Y|W=w}\right)$  $\subseteq$  $\operatorname{supp}(\pi_{XY}),$ otherwise,  $\sum_{w} P(w) \mathcal{H}(P_{X|W=w}, P_{Y|W=w} || \pi_{XY}) = \infty$ which contradicts the optimality of  $P_W P_{X|W} P_{Y|W}$ . On the other hand, we have that

$$\Gamma^{\text{UB}}(\pi_{XY})$$

$$= -H(XY|W) + \sum_{w} P(w)\mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY})$$
(64)

$$\geq -H(XY|W) + H(XY) \tag{65}$$

$$\geq C_{\mathsf{Wyner}}(\pi_{XY}) \tag{66}$$

By assumption, the inequalities in (65) and (66) are equalities. Hence  $P_W P_{X|W} P_{Y|W}$  also attains  $C_{Wyner}(\pi_{XY})$  and the following equality holds:

$$\sum_{w} P(w) \mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY}) = H(XY).$$
 (67)

Equation (67) implies

$$\mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY})$$
  
=  $\sum_{x,y} P(x|w)P(y|w)\log\frac{1}{\pi(x,y)}$  (68)

for every  $w \in \text{supp}(P_W)$ . By [24, Proposition 2], for every  $w \in \text{supp}(P_W)$ ,  $\pi_{XY}$  is product on the set  $\text{supp}(P_{X|W=w}) \times \text{supp}(P_{Y|W=w})$ . Hence  $\pi_{XY}$  satisfies Condition (\*).

The following is a special case of Condition (\*).

**Definition 6.** A joint distribution  $\pi_{XY}$  is pseudo-product if for some  $A \subseteq \mathcal{X} \times \mathcal{Y}$ ,

$$\pi_{XY}(x,y) = \begin{cases} \alpha(x)\beta(y) & (x,y) \in A\\ 0 & \text{otherwise} \end{cases}$$
(69)

where  $\alpha : \mathcal{X} \to \mathbb{R}_{>0}$  and  $\beta : \mathcal{Y} \to \mathbb{R}_{>0}$  are two positive functions such that  $\sum_{(x,y)\in A} \alpha(x)\beta(y) = 1$ .

*Remark* 7. In general, a pseudo-product distribution may not be a product distribution. For example,

$$\frac{1}{\alpha_0\beta_0 + \alpha_0\beta_1 + \alpha_1\beta_0} \begin{bmatrix} \alpha_0\beta_0 & \alpha_0\beta_1\\ \alpha_1\beta_0 & 0 \end{bmatrix}$$
(70)

is a pseudo-product distribution but not a product distribution. However, if  $supp(\pi_{XY})$  is a product set, then a pseudoproduct distribution  $\pi_{XY}$  is a product distribution.

Obviously, pseudo-product distributions satisfy Condition (\*). Hence for pseudo-product distributions, the exact and Wyner's common informations are equal.

#### IV. EXTENSION TO SOURCES WITH GENERAL ALPHABETS

In Section III, we derived exact and  $\infty$ -Rényi common informations for sources with finite alphabets. In this section, we generalize the results to sources with countably infinite alphabets and a certain class of continuous sources. Furthermore, note that for Wyner's common information, till date, only the case of sources with finite alphabets was studied by Wyner [2], and there is no characterization<sup>5</sup> for sources with countably infinite alphabets and continuous sources. Hence in this section, before generalizing the exact and  $\infty$ -Rényi common informations, we first generalize Wyner's common information to such sources. In the proofs of the converse parts, the mixturedecomposition technique is used extensively. We show that for sources with countably infinite alphabets and a certain class of continuous sources (including Gaussian sources), Wyner's common information remains  $C_{Wyner}(\pi_{XY})$ . Moreover, for a source with countably infinite alphabets, we show that Wyner's common information can be obtained by computing the common information for an alphabet-truncated version of the source and then taking limits to enlarge the domain of the truncated alphabet.

#### A. Wyner's Common Information

Wyner [2] only characterized the common information for sources with finite alphabets. Here we extend his results to sources with countably infinite alphabets and continuous sources. The proof of Theorem 5 is given in Appendix D.

**Theorem 5** (Wyner's Common Information for General Sources). Let (X, Y) be a source with distribution  $\pi_{XY}$  defined on the product of two arbitrary alphabets (i.e., on the product of two arbitrary measurable spaces). Then we have

$$\widetilde{C}_{\text{Wyner}}(\pi_{XY}) \le \widetilde{T}_1(\pi_{XY}) \le T_1(\pi_{XY}) \le \widehat{C}_{\text{Wyner}}(\pi_{XY}),$$
(71)

where<sup>6</sup>

$$\widetilde{C}_{\text{Wyner}}(\pi_{XY}) := \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_X \mid W P_Y \mid W:\\ D(P_X \lor \| \pi_{YY}) \le \epsilon}} I(XY; W)$$
(72)

and

$$C_{Wyner}(\pi_{XY}) := \inf_{\substack{P_W P_X|W} P_Y|W: \ s \downarrow 0 \\ P_{XY} = \pi_{XY}}} \lim_{\substack{P_W P_X|W} P_Y|W: \ s \downarrow 0} D_{1+s}(P_X|W P_Y|W ||P_{XY}|P_W).$$
(73)

Since  $D(P_{X|W}P_{Y|W}||P_{XY}|P_W) = I(XY; W)$  and for any fixed (P,Q),  $D_{1+s}(P||Q)$  is non-decreasing in s, we know that

$$\widetilde{C}_{Wyner}(\pi_{XY}) \le C_{Wyner}(\pi_{XY}) \le \widehat{C}_{Wyner}(\pi_{XY}).$$
 (74)

Note that for any fixed (P,Q), the Rényi divergence  $D_{1+s}(P||Q)$  is continuous in  $s \in [-1,0] \cup \{s \in (0,\infty] : D_{1+s}(P||Q) < \infty\}$ . However, van Erven and Harremoës in [27] showed that there exists a pair of distributions (P,Q) such that the Rényi divergence  $D_{1+s}(P||Q)$  is not continuous at s = 0. Hence we do not know if the inequalities in (74) are equalities in general.

**Proposition 1.** The following are sufficient conditions to ensure  $\widehat{C}_{Wyner}(\pi_{XY}) = C_{Wyner}(\pi_{XY})$ .

1) There exists a joint distribution  $P_W P_{X|W} P_{Y|W}$  that attains  $C_{Wyner}(\pi_{XY})$  and satisfies

$$D_{1+s}(P_{X|W}P_{Y|W}||P_{XY}|P_W) < \infty$$
(75)

<sup>6</sup>For this arbitrary alphabet case, the distributions  $P_W P_{X|W} P_{Y|W}$  in the infimizations in the definitions of  $\tilde{C}_{Wyner}(\pi_{XY})$  and  $C_{Wyner}(\pi_{XY})$  (for the latter, see (2)) are restricted to satisfy that the mutual information I(XY; W) exists; if there is no such distribution, then  $\tilde{C}_{Wyner}(\pi_{XY}) := \infty$  and  $C_{Wyner}(\pi_{XY}) := \infty$ . Here we say the mutual information I(U; V) of two random variables U and V exists if  $P_{UV} \ll P_U P_V$  and the integral  $\int_{\mathcal{U} \times \mathcal{V}} \left| \log \frac{\mathrm{d}P_{UV}}{\mathrm{d}(PUP_V)} \right| \mathrm{d}P_{UV} < \infty$ . The mutual information always exists for distributions with finite alphabets but does not always exist for other distributions. Hence here we need to add this constraint. Similarly, the distribution  $P_W P_{X|W} P_{Y|W}$  in the infimization in the definition of  $\hat{C}_{Wyner}(\pi_{XY})$  is restricted to satisfy that  $D_{1+s}(P_X|WP_Y|W||P_{XY}|P_W)$  exists for some s > 0; if there is no such distribution, then  $\hat{C}_{Wyner}(\pi_{XY}) := \infty$ . Here we say  $D_{1+s} \left( P_U|V| \|P_U|P_V \right)$  exists if  $P_{UV} \ll P_U P_V$  and the integral  $\frac{1}{s} \log \int_{\mathcal{U} \times \mathcal{V}} \left( \frac{\mathrm{d}P_{UV}}{\mathrm{d}(P_U P_V)} \right)^s \mathrm{d}P_{UV} < \infty$ .

<sup>&</sup>lt;sup>5</sup>More precisely, there is no converse result derived for sources with countably infinite alphabets and continuous sources. As for the achievability part, several existing results on channel resolvability (e.g., [5], [7], [25]) can be applied to obtain achievability results for the Wyner's common information problem.

for some s > 0.

2) There exists a sequence of joint distributions  $P_{WXY}^{(k)} := P_W^{(k)} P_{X|W}^{(k)} P_{Y|W}^{(k)}$  such that they attain  $C_{Wyner}(\pi_{XY})$  asymptotically, i.e.,  $P_{XY}^{(k)} = \pi_{XY}$ ,  $\lim_{k\to\infty} I_{P^{(k)}}(XY;W) = C_{Wyner}(\pi_{XY})$ , and for every k, there exists some  $s_k > 0$  satisfying

$$D_{1+s_k}(P_{X|W}^{(k)}P_{Y|W}^{(k)}\|P_{XY}^{(k)}|P_W^{(k)}) < \infty.$$
(76)

*Proof:* Here we only prove Statement 2). Statement 1) follows similarly.

Suppose that there exists a sequence of joint distributions  $P_W^{(k)}P_{X|W}^{(k)}P_{Y|W}^{(k)}$  satisfying the conditions given in Statement 2). Then (76) implies that given  $P_W^{(k)}P_{X|W}^{(k)}P_{Y|W}^{(k)}$ , the conditional Rényi divergence  $D_{1+s}(P_{X|W}^{(k)}P_{Y|W}^{(k)}||P_{XY}^{(k)}|P_W^{(k)})$  is continuous in  $s \in [-1, s_k]$ . Hence

$$\lim_{s \downarrow 0} D_{1+s}(P_{X|W}^{(k)} P_{Y|W}^{(k)} \| P_{XY}^{(k)} | P_W^{(k)}) = I_{P^{(k)}}(XY; W).$$
(77)

Therefore, by the definition of  $\widehat{C}_{Wyner}(\pi_{XY})$ , for all k,

$$\widehat{C}_{\text{Wyner}}(\pi_{XY}) \le \lim_{s \downarrow 0} D_{1+s}(P_{X|W}^{(k)} P_{Y|W}^{(k)} \| P_{XY}^{(k)} | P_{W}^{(k)})$$
(78)

$$= I_{P^{(k)}}(XY;W).$$
(79)

By assumption,  $I_{P^{(k)}}(XY;W) \to C_{Wyner}(\pi_{XY})$  as  $k \to \infty$ . Hence

$$\widehat{C}_{\text{Wyner}}(\pi_{XY}) \le C_{\text{Wyner}}(\pi_{XY}).$$
(80)

Observe that the requirements (75) and (76) are respectively equivalent to

$$\int_{\mathcal{W}\times\mathcal{X}\times\mathcal{Y}} \left(\frac{\mathrm{d}\left(P_{X|W}P_{Y|W}\right)}{\mathrm{d}P_{XY}}\right)^{s} \mathrm{d}\left(P_{W}P_{X|W}P_{Y|W}\right) < \infty$$
(81)

and

$$\int_{\mathcal{W}\times\mathcal{X}\times\mathcal{Y}} \left( \frac{\mathrm{d}\left(P_{X|W}^{(k)}P_{Y|W}^{(k)}\right)}{\mathrm{d}P_{XY}^{(k)}} \right)^{s} \mathrm{d}\left(P_{W}^{(k)}P_{X|W}^{(k)}P_{Y|W}^{(k)}\right) < \infty$$
(82)

for s > 0. Note that for s = 0, (81) and (82) are satisfied. Hence we conjecture that the conditions given in Proposition 1 hold for a large class of sources.

For the finite alphabet case, it is easy to verify that  $\widetilde{C}_{Wyner}(\pi_{XY}) = C_{Wyner}(\pi_{XY}) = \widehat{C}_{Wyner}(\pi_{XY})$ . Hence  $\widetilde{T}_1(\pi_{XY}) = T_1(\pi_{XY}) = C_{Wyner}(\pi_{XY})$  for this case. The result  $\widetilde{T}_1(\pi_{XY}) = C_{Wyner}(\pi_{XY})$  was first proven by Wyner [2]. The case concerning sources with countably infinite alphabets and the case concerning a certain class of continuous sources are considered in the following corollaries. The proofs are given in Appendices E and F.

**Corollary 2.** Let (X, Y) be a source with distribution  $\pi_{XY}$  defined on the product of two countably infinite alphabets. Assume  $H_{\alpha}(\pi_{XY})$  exists (and hence is finite) for some  $\alpha \in [0, 1)$ . Then we have

$$T_1(\pi_{XY}) = T_1(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY}).$$
 (83)

Remark 8. In our proof, we show that

$$\widetilde{C}_{\text{Wyner}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY}) \tag{84}$$

$$= \lim_{k \to \infty} C_{\text{Wyner}}(\pi_{XY}^{(k)}) \tag{85}$$

$$= \lim_{k \to \infty} C_{\mathrm{Wyner}}(\pi_{[X]_k[Y]_k})$$
(86)

where

$$\pi_{XY}^{(k)}(x,y) := \frac{\pi_{XY}(x,y) \mathbb{1}\left\{(x,y) \in [-k,k]^2\right\}}{\pi_{XY}([-k,k]^2)}$$
(87)

and  $\pi_{[X]_k[Y]_k}$  with  $[z]_k := z$ , if  $|z| \le k$ , and k + 1, otherwise, denote distributions induced by truncation operations. That is to say, we can compute Wyner's common information for countably-infinite-valued sources by computing the common information for their truncated versions and then taking limit in k.

**Corollary 3.** Assume  $\pi_{XY}$  is an absolutely continuous distribution on  $\mathbb{R}^2$  such that  $\widehat{C}_{Wyner}(\pi_{XY}) = C_{Wyner}(\pi_{XY})$  (e.g., at least one of the conditions given in Proposition 1 is satisfied) and its pdf<sup>7</sup>  $\pi_{XY}$  is log-concave<sup>8</sup> and differentiable. Assume I(X;Y) exists (and hence is finite). For d > 0, define

$$L_{d} := \sup_{(x,y)\in[-d,d]^{2}} \left| \frac{\partial}{\partial x} \log \pi_{XY}(x,y) \right| + \left| \frac{\partial}{\partial y} \log \pi_{XY}(x,y) \right|,$$
(88)

and

$$\epsilon_d := 1 - \pi_{XY} \left( [-d, d]^2 \right).$$
 (89)

Assume that  $\epsilon_d \log (dL_d) \to 0$  as  $d \to +\infty$ . Then we have

$$\widetilde{T}_1(\pi_{XY}) = T_1(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY}).$$
 (90)

*Remark* 9. If the pdf  $\pi_{XY}$  is *not* differentiable, then Corollary 3 still holds if 1) the pdf  $\pi_{XY}$  is continuous (this is also implied by the log-concavity of the pdf  $\pi_{XY}$ ) and 2) the definition of  $L_d$  in (88) is replaced with

$$L_{d} := \sup_{\Delta \ge 0} \frac{1}{\Delta} \log \sup_{\substack{(x,y), (\hat{x}, \hat{y}) \in [-d,d]^{2}: \\ |x-\hat{x}|, |y-\hat{y}| \le \Delta}} \frac{\pi_{XY}(x,y)}{\pi_{XY}(\hat{x}, \hat{y})}.$$
 (91)

This claim follows since in our proof of Corollary 3, the assumption of differentiability of the pdf  $\pi_{XY}$  is used to upper bound the RHS of (91) by using  $L_d$  (see Lemma 13); however, adopting the definition of  $L_d$  in (91) avoids this complicated derivation, since it directly relates  $L_d$  to the RHS of (91).

Now we consider bivariate Gaussian sources (X, Y). Without loss of any generality, we assume that the correlation coefficient  $\rho$  between X and Y is nonnegative; otherwise, we can set -X to X. For this case,

$$C_{\text{Wyner}}(\pi_{XY}) = \frac{1}{2} \log \left[ \frac{1+\rho}{1-\rho} \right]$$
(92)

and it is attained by the joint Gaussian distribution  $P_W P_{X|W} P_{Y|W}$  with  $P_W = \mathcal{N}(0,\rho), P_{X|W}(\cdot|w) = \mathcal{N}(w, 1-\rho), P_{Y|W}(\cdot|w) = \mathcal{N}(w, 1-\rho)$  [11], [12]. Using the formula

<sup>7</sup>For brevity, we use the same notation  $\pi_{XY}$  to denote both an absolutely continuous distribution and the corresponding pdf.

<sup>8</sup>A pdf  $\pi_{XY}$  is log-concave if  $\log \pi_{XY}$  is concave.

for Rényi divergences between Gaussian distributions derived in [28], we obtain that for  $0 < s \le \sqrt{\frac{1+\rho}{2\rho}}$ ,

$$D_{1+s}(P_{X|W}P_{Y|W} || P_{XY} | P_W) = \frac{1}{2} \log \left[ \frac{1+\rho}{1-\rho} \right] - \frac{1}{2s} \log \left( 1 - \frac{2s^2\rho}{1+\rho} \right).$$
(93)

Hence Gaussian sources satisfy the sufficient condition 1) given in Proposition 1, which in turn implies  $\widehat{C}_{Wyner}(\pi_{XY}) = C_{Wyner}(\pi_{XY})$ . Furthermore, it is easy to verify that other conditions given in Corollary 3 are also satisfied by Gaussian sources. Hence we obtain the following result.

**Corollary 4.** For a bivariate Gaussian source (X, Y) with correlation coefficient  $\rho \in [0, 1)$ , we have

$$\widetilde{T}_{1}(\pi_{XY}) = T_{1}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY}) = \frac{1}{2} \log \left[ \frac{1+\rho}{1-\rho} \right].$$
(94)

*Proof:* The last equality in (94) was proven in [11], [12]. The first two equalities in (94) are implied by Corollary 3, since it is easy to verify that the hypotheses as stated in Corollary 3 are satisfied by Gaussian sources.

If we replace the relative entropy measure with the TVdistance, we can define the TV-distance version of Wyner's common information as

$$T_{\rm TV}(\pi_{XY}) := \inf \Big\{ R : \lim_{n \to \infty} |P_{X^n Y^n} - \pi_{XY}^n| = 0 \Big\}.$$
(95)

By replacing the relative entropy with the TV-distance in our proofs, one can easily obtain the following result. The proof is similar to the ones for the relative entropy versions, and hence is omitted here.

#### Theorem 6. Redefine

$$\widetilde{C}_{\text{Wyner}}(\pi_{XY}) := \lim_{\epsilon \downarrow 0} \inf_{P_W P_{X|W} P_{Y|W} : |P_{XY} - \pi_{XY}| \le \epsilon} I(XY;W)$$
(96)

Then Theorem 5 as well as Corollaries 2, 3, and 4 hold mutatis mutandis for the TV-distance version of Wyner's common information.

*Remark* 10. For the TV-distance version, the upper bound in Theorem 5  $\widehat{C}_{Wyner}(\pi_{XY})$  can be tightened to be  $C_{Wyner}(\pi_{XY})$ , by replacing Lemma 10 in the proof with [5, Theorem VII.1]. This in turn implies that the requirement of existence of  $H_{\alpha}(\pi_{XY})$  for some  $\alpha \in [0, 1)$  in Corollary 2 can be relaxed to that of existence of  $H(\pi_{XY})$ , and the requirement that  $\widehat{C}_{Wyner}(\pi_{XY}) = C_{Wyner}(\pi_{XY})$  in Corollary 3 can be removed.

Since it is difficult to obtain closed-form expressions for the optimal joint distributions  $P_W P_{X|W} P_{Y|W}$  attaining  $C_{Wyner}(\pi_{XY})$ , the sufficient conditions for  $\hat{C}_{Wyner}(\pi_{XY}) =$  $C_{Wyner}(\pi_{XY})$  given in Proposition 1 is difficult to verify. However, for the TV-distance version, the requirement of  $\hat{C}_{Wyner}(\pi_{XY}) = C_{Wyner}(\pi_{XY})$  in Corollary 3 can be removed. Hence for this case, the conditions in Corollary 3 can be easily verified. For example, besides Gaussian sources, "Laplacian" sources<sup>9</sup>  $\pi_{XY}(x, y) \propto \exp(-|ax + by|)$  with

 $^9 \rm Note$  that here  $\pi_{XY}(x,y) \propto \exp{(-|ax+by|)}$  is not the common bivariate Laplacian distribution.

 $a, b \in \mathbb{R}$  also satisfy the required conditions, and hence for "Laplacian" sources,  $T_{\text{TV}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY})$ .

The exponential strong converse holds for the TV-distance version of Wyner's common information when the alphabet is finite; see [3], [4]. We conjecture that the exponential strong converse also holds when the alphabet is infinite (countably infinite or uncountable).

#### B. Exact and ∞-Rényi Common Informations

Now we generalize exact and  $\infty$ -Rényi common informations to sources with countably infinite alphabets and a certain class of continuous sources.

1) Equivalence: In Theorem 1, we established the equivalence between the exact and  $\infty$ -Rényi common informations for sources with finite alphabets. Now we extend it to the countably infinite alphabet case.

**Theorem 7** (Equivalence). Let (X, Y) be a source with distribution  $\pi_{XY}$  defined on the product of two countably infinite alphabets. Assume  $H(\pi_{XY})$  exists (and hence is finite). Then we have

$$T_{\text{Exact}}(\pi_{XY}) = T_{\infty}(\pi_{XY}). \tag{97}$$

For sources with discrete (finite or countably infinite) or continuous alphabets, we have shown  $T_{\text{Exact}}(\pi_{XY}) \geq T_{\infty}(\pi_{XY})$  in Lemma 6 in Appendix A. Thus it suffices to prove the reverse inequality.

**Lemma 2.** Let (X, Y) be a source with distribution  $\pi_{XY}$ defined on the product of two countably infinite alphabets. Assume  $H(\pi_{XY})$  exists (and hence is finite). Then for a source with such a distribution  $\pi_{XY}$ , if there exists a sequence of fixed-length codes with rate R that generates  $P_{X^nY^n}$  such that  $D_{\infty}(P_{X^nY^n} || \pi_{XY}^n) \to 0$ , then there must exist a sequence of variable-length codes with rate R that exactly generates  $\pi_{XY}^n$ . That is,  $T_{\text{Exact}}(\pi_{XY}) \leq T_{\infty}(\pi_{XY})$ .

The proof of Lemma 2 is given in Appendix G.

Until now, we have shown that  $T_{\text{Exact}}(\pi_{XY}) \geq T_{\infty}(\pi_{XY})$ holds for sources with discrete or continuous alphabets, and  $T_{\text{Exact}}(\pi_{XY}) \leq T_{\infty}(\pi_{XY})$  holds for sources with discrete alphabets. However, we do not know whether  $T_{\text{Exact}}(\pi_{XY}) \leq$  $T_{\infty}(\pi_{XY})$  always holds for continuous sources. Next we prove that it indeed holds if continuous sources satisfy certain regularity conditions, and the optimal (minimum)  $\infty$ -Rényi divergence  $D_{\infty}(P_{X^nY^n} || \pi_{XY}^n)$  converges to zero sufficiently fast. The proof of Lemma 3 is given in Appendix H.

**Lemma 3.** Assume  $\pi_{XY}$  is an absolutely continuous distribution on  $\mathbb{R}^2$  with  $\mathbb{E}[X^2]$ ,  $\mathbb{E}[Y^2] < \infty$ . Without loss of generality, we assume  $\mathbb{E}[X^2] = \mathbb{E}[Y^2] = 1$ . Assume the pdf of  $\pi_{XY}$  is log-concave, and continuously differentiable. Assume I(X;Y) exists (and hence is finite). For  $\epsilon > 0$  and  $n \in \mathbb{N}$ , define

$$L_{\epsilon,n} := \sup_{(x,y)\in\mathcal{L}^{2}_{\epsilon,n}} \left\{ \left| \frac{\partial}{\partial x} \log \pi_{XY}(x,y) \right| + \left| \frac{\partial}{\partial y} \log \pi_{XY}(x,y) \right| \right\},$$
(98)

where

$$\mathcal{L}_{\epsilon,n} := \left\{ x \in \mathbb{R} : |x| \le \sqrt{n\left(1+\epsilon\right)} \right\}.$$
(99)

Assume  $\log L_{\epsilon,n}$  is sub-exponential in n for fixed  $\epsilon$  (i.e.,  $\lim_{n\to\infty} \frac{1}{n} \log \log L_{\epsilon,n} = 0$  for all fixed  $\epsilon > 0$ ). Then for a source with such a distribution  $\pi_{XY}$ , if there exists a sequence of fixed-length codes with rate R that generates  $P_{X^nY^n}$  such that  $D_{\infty}(P_{X^nY^n} || \pi_{XY}^n) = o\left(\frac{1}{n + \log L_{\epsilon,n}}\right)$  for any  $\epsilon > 0$ , then there must exist a sequence of variablelength codes with rate R that exactly generates  $\pi_{XY}^n$ . That is,  $T_{\text{Exact}}(\pi_{XY}) \leq T'_{\infty}(\pi_{XY})$ , where

$$T'_{\infty}(\pi_{XY}) := \inf \left\{ R : D_{\infty}(P_{X^nY^n} \| \pi^n_{XY}) \\ = o\left(\frac{1}{n + \log L_{\epsilon,n}}\right), \forall \epsilon > 0 \right\}.$$
(100)

*Remark* 11. Similar to Remark 9, if the pdf  $\pi_{XY}$  is not differentiable, then Corollary 3 still holds if 1) the pdf  $\pi_{XY}$  is continuous and 2) the definition of  $L_{\epsilon,n}$  in (98) is replaced with

$$L_{\epsilon,n} := \sup_{\Delta \ge 0} \frac{1}{\Delta} \log \sup_{\substack{(x,y), (\hat{x}, \hat{y}) \in \mathcal{L}^2_{\epsilon,n}: \\ |x-\hat{x}|, |y-\hat{y}| \le \Delta}} \frac{\pi_{XY}(x, y)}{\pi_{XY}(\hat{x}, \hat{y})}.$$
 (101)

*Remark* 12. One important example satisfying the conditions in the lemma above is bivariate Gaussian sources. Consider a bivariate Gaussian source  $\pi_{XY} = \mathcal{N}(0, \Sigma_{XY})$  where  $\Sigma_{XY} = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}$  with  $\rho \in [0, 1)$ . For this case,

$$L_{\epsilon,n} = \sup_{(x,y)\in\mathcal{L}^2_{\epsilon,n}} \left| \frac{x-\rho y}{1-\rho^2} \right| + \left| \frac{y-\rho x}{1-\rho^2} \right|$$
(102)

$$=\frac{2\sqrt{n\left(1+\epsilon\right)}}{1-\rho}.$$
(103)

Hence  $\log L_{\epsilon,n}$  is sub-exponential in n for fixed  $\epsilon$ . Observe that  $\frac{1}{n+\log L_{\epsilon,n}} \sim \frac{1}{n}$ . Hence, by this lemma, if there exists a sequence of fixed-length codes with rate R that generates  $P_{X^nY^n}$  such that  $D_{\infty}(P_{X^nY^n} \| \pi_{XY}^n) = o(\frac{1}{n})$ , then there must exist a sequence of variable-length codes with rate R that exactly generates  $\pi_{XY}^n$ .

2) Discrete Sources with Countably Infinite Alphabets: We now generalize the exact and  $\infty$ -Rényi common informations to sources with countably infinite alphabets. In the proof of Theorem 1, a truncated i.i.d. code was adopted to prove the achievability part, in which the codewords are i.i.d. with each drawn according to a set of truncated distributions (obtained by truncating a set of product distributions into some (strongly) typical sets). For the countably infinite alphabet case, we need replace strongly typical sets with unified typical sets (defined in (18)). Then we establish the following result. **Corollary 5.** Let (X, Y) be a source with distribution  $\pi_{XY}$  defined on the product of two countably infinite alphabets. Assume  $H(\pi_{XY})$  exists (and hence is finite). We have

$$\max\left\{\widehat{\Gamma}^{\text{LB}}(\pi_{XY}), C_{\text{Wyner}}(\pi_{XY})\right\} \le \widetilde{T}_{\infty}(\pi_{XY})$$
(104)

$$\leq T_{\infty}(\pi_{XY}) \tag{105}$$

$$=T_{\mathrm{Exact}}(\pi_{XY})$$
 (106)

$$\leq \widehat{\Gamma}^{\mathrm{UB}}(\pi_{XY}),$$
 (107)

where

$$\widehat{\Gamma}^{\text{UB}}(\pi_{XY}) := \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_X | W P_Y | W: \\ P_{XY} = \pi_{XY}}} \sup_{\substack{Q_{XYW}: \\ D(Q_{WX} || P_{WX}) \le \epsilon, \\ D(Q_{WY} || P_{WY}) \le \epsilon}} \left\{ -\sum_{w,x,y} P(w) Q(x, y | w) \log \pi(x, y) - H(XY | W) \right\}$$
(108)

and

$$\widehat{\Gamma}^{\text{LB}}(\pi_{XY}) := \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_X | W P_Y | W : \\ D(P_{XY} | | \pi_{XY}) \le \epsilon}} \left\{ -H(XY|W) + \inf_{\substack{Q_{WW'} \in C(P_W, P_W) \\ w, w'}} \sum_{w, w'} Q(w, w') \right\} \\
\times \mathcal{H}(P_{X|W=w}, P_{Y|W=w'} || \pi_{XY}) \left\}.$$
(109)

For the finite alphabet case, the  $\epsilon$ 's in the optimizations in (108) and (109) can be removed by using the compactness technique or the splitting technique. For the countably infinite alphabet case, in general we cannot apply the compactness technique. However, it may be possible to apply the splitting technique to remove  $\epsilon$ 's, similarly as in the proof of Corollary 2. Nevertheless, we need carefully deal with the terms involving  $\log \pi (x, y)$  in (108) and (109), since a little difference between  $Q_{XY}$  and  $\pi_{XY}$  could lead to a large increase of  $\sum_{x,y} Q(x, y) \log \frac{1}{\pi(x,y)}$ . 3) Gaussian Sources: Next we generalize the exact and  $\infty$ -

3) Gaussian Sources: Next we generalize the exact and  $\infty$ -Rényi common informations to a certain class of continuous sources. We provide an upper bound on  $T_{\text{Exact}}(\pi_{XY})$  and  $T_{\infty}(\pi_{XY})$  for bivariate Gaussian sources  $\pi_{XY}$ . Without loss of any generality, we assume that the correlation coefficient  $\rho$  between (X, Y) is nonnegative. The proof of Theorem 8 is given in Appendix I.

**Theorem 8.** For a Gaussian source (X, Y) with correlation coefficient  $\rho \in [0, 1)$ , we have

$$\frac{1}{2}\log\left[\frac{1+\rho}{1-\rho}\right] \le \widetilde{T}_{\infty}(\pi_{XY}) \tag{110}$$

$$\leq T_{\infty}(\pi_{XY}) \tag{111}$$

$$=T_{\mathrm{Exact}}(\pi_{XY}) \tag{112}$$

$$\leq \frac{1}{2} \log \left[ \frac{1+\rho}{1-\rho} \right] + \frac{\rho}{1+\rho}.$$
 (113)

*Remark* 13. For Gaussian sources (X, Y) with correlation coefficient  $\rho \in [0, 1)$ , Li and El Gamal [10] provided the following upper bound

$$T_{\text{Exact}}(\pi_{XY}) \le \frac{1}{2} \log\left[\frac{1}{1-\rho^2}\right] + 24 \log 2.$$
 (114)

Such an upper bound is a one-shot bound, and hence it is also valid for the case with blocklength equal to 1. However, our upper bound requires blocklength to be infinity. Furthermore, for the asymptotic case, Li and El Gamal's bound is rather loose, since the difference between the upper bounds in (114) and (113) is

$$\frac{1}{2}\log\left[\frac{1}{1-\rho^2}\right] + 24\log 2 - \left(\frac{1}{2}\log\left[\frac{1+\rho}{1-\rho}\right] + \frac{\rho}{1+\rho}\right)$$
$$= 24\log 2 - 1 + \frac{1}{1+\rho} + \log\left[\frac{1}{1+\rho}\right]$$
(115)

 $\geq 15.44$  Nats/Symbol

$$= 22.28 \text{ Bits/Symbol.}$$
 (117)

Li and El Gamal's bound was proven by using a dyadic decomposition scheme which decomposes the joint distribution into a sequence of uniform distributions. For such a scheme, even if the source  $\pi_{XY}$  is comprised of two independent components and at least one of them is not uniform (i.e.,  $\pi_{XY} = \pi_X \pi_Y$ but either  $\pi_X$  or  $\pi_Y$  is not uniform), the induced common randomness rate between them is still strictly positive. This is because for this case, Li and El Gamal's dyadic decomposition scheme cannot identify the optimal decomposition  $\pi_{XY} = \pi_X \pi_Y$ . Hence the common randomness rate induced by Li and El Gamal's scheme does not cross 0 for  $\rho = 0$ . In addition, it is worth noting that our exact common information scheme is a mixture of Li and El Gamal's scheme and an  $\infty$ -Rényi common information scheme. In our scheme, Li and El Gamal's scheme is invoked with asymptotically vanishing probability, and hence the performance of our scheme is dominated by the  $\infty$ -Rényi common information scheme which requires a much lower rate.

For the DSBS case, our upper bound is tight. Hence it is natural to conjecture that for Gaussian sources, the upper bound in (113) is also tight. Similarly to the discrete source case, one can show the following lower bound on  $T_{\text{Exact}}(\pi_{XY})$  and  $T_{\infty}(\pi_{XY})$  holds for continuous sources (including Gaussian sources).

$$\widehat{\Gamma}^{\text{LB}}(\pi_{XY}) := \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_X | W P_Y | W: \\ D(P_{XY} || \pi_{XY}) \le \epsilon}} \left\{ -h(XY|W) + \inf_{\substack{Q_{WW'} \in C(P_W, P_W) \\ w, w'}} \sum_{w, w'} Q(w, w') \right\} \\
\times \mathcal{H}(P_{X|W=w}, P_{Y|W=w'} || \pi_{XY}) \right\},$$
(118)

where  $P_W$  is a discrete distribution, given w, w',  $P_{X|W=w}, P_{Y|W=w'}$  are continuous distributions, and  $\mathcal{H}(P_{X|W=w}, P_{Y|W=w'} || \pi_{XY})$  is the maximal (differential) cross-entropy defined in (38) (with  $Q_{XY}, \pi_{XY}$  denoting the pdfs rather than pmfs). However, we do not know how to prove  $\widehat{\Gamma}^{\text{LB}}(\pi_{XY}) \geq \frac{1}{2} \log \left[ \frac{1+\rho}{1-\rho} \right] + \frac{\rho}{1+\rho}$ . Furthermore, it is possible to generalize the upper bound in Theorem 8 to other continuous sources by utilizing general typicality, e.g., [29], [30].

For Gaussian sources, Li and El Gamal's upper bound in (114), our upper bound in (113), and Wyner's common information in (94) are illustrated in Fig. 3. The exact and  $\infty$ -Rényi common informations are lower bounded by Wyner's common information. Hence the exact and  $\infty$ -Rényi common informations are between Wyner's common information and our bound. The gap between them is  $\frac{\rho}{1+\rho} \leq 0.5$  nats/symbol or 0.72 bits/symbol.

# V. CONNECTION TO OTHER PROBLEMS

The exact common information problem is related to (or can be generalized to) the following problems.

#### • Distributed Channel Synthesis

(116)

In both the exact and TV-approximate senses, the common information problem is equivalent to the distributed channel simulation problem (with no shared information). The distributed channel simulation problem (or the communication complexity problem for generating correlation), illustrated in Fig. 4, was studied in [5], [31]–[34]. The distributed exact (resp. TV-approximate) channel simulation problem refers to determining the minimum communication rate needed to generate two correlated sources  $(X^n, Y^n)$  respectively at the encoder and decoder such that the induced joint distribution  $P_{X^nY^n}$  exactly equals  $\pi_{XY}^n$  (resp. the TV distance  $P_{X^nY^n}$ and  $\pi_{XY}^n$  vanishes asymptotically).

The exact common information problem (or exact correlation generation problem) is essentially equivalent to the distributed channel simulation problem with no shared information (or the communication complexity problem for generating correlation) [5], [31]–[34] (illustrated in Fig. 4 with  $R_0 = 0$ ). This can be easily obtained by observing that if there exists an exact common information code  $(P_{M_n}, P_{X^n|M_n}, P_{Y^n|M_n})$ then  $(P_{M_n|X^n}, P_{Y^n|M_n})$  forms an exact channel synthesis code; and vice versa.

In the literature, Bennett et al. [31] studied exact syntheses of a target channel when there is unlimited shared randomness, i.e.,  $R_0 = \infty$ , available at the encoder and decoder. They showed that the minimum communication rates for this case is equal to the mutual information  $I_{\pi}(X;Y)$  in which  $(X, Y) \sim \pi_{XY}$ . Harsha *et al.* [34] used a rejection sampling scheme to prove a one-shot bound for exact simulation for finitely-supported (X, Y). They showed that the number of bits of the shared randomness can be limited to  $O(\log \log |\mathcal{X}| +$  $\log |\mathcal{Y}|$  if the expected description length is increased by  $O(\log (I_{\pi}(X;Y) + 1) + \log \log |\mathcal{Y}|)$  bits from the lower bound  $I_{\pi}(X;Y)$ . Li and El Gamal [35] used functional representation lemma to prove that if the expected description length is increased by  $\log(I_{\pi}(X;Y)+1)+5$  bits from  $I_{\pi}(X;Y)$ , then the number of bits of the shared randomness can be upper bounded by  $\log(|\mathcal{X}|(|\mathcal{Y}|-1)+2)$ . The tradeoff between the communication rate and the shared randomness rate for exact synthesis of the symmetric binary erasure source (SBES) was characterized by Kumar, Li, and El Gamal [6]. Recently, we extend the results and the proof techniques in this paper to study the tradeoff between the communication rate and the shared randomness rate for exact synthesis of discrete and continuous memoryless channels. In particular, we completely characterized the tradeoff for DSBSes. Furthermore, there are also multiple works, e.g., [5], [31], [32], studying approximate syntheses of a target channel, in which the distance between the generated channel and the target channel is required to converge to zero asymptotically.



Fig. 3. Illustrations of Wyner's common information (94), as well as Li and El Gamal's upper bound (114) and our upper bound (113) on the exact and  $\infty$ -Rényi common informations for Gaussian sources with correlation coefficient  $\rho \in [0, 1)$ . For ease of comparison, here we plot Li and El Gamal's upper bound minus 20 (bits/symbol), rather than their bound itself, since their bound is much larger than our bound and Wyner's common information.



Fig. 4. The exact channel synthesis problem. We would like to design the code  $(P_{W_n|X^nK_n}, P_{Y^n|W_nK_n})$  such that the induced conditional distribution  $P_{Y^n|X^n}$  satisfies  $P_{Y^n|X^n} = \pi_{Y|X}^n$ .

#### • Exact α-Rényi Common Informations

As shown in [6] (or (33)), the exact common information for  $\pi_{XY}$  is equal to

$$\lim_{n \to \infty} \frac{1}{n} \min_{P_W P_{X^n | W} P_{Y^n | W} : P_{X^n Y^n} = \pi_{XY}^n} H(W).$$
(119)

Note that the  $\alpha$ -Rényi entropy with  $\alpha \in [-\infty, \infty]$  is defined as

$$H_{\alpha}(W) := \frac{1}{1-\alpha} \log \sum_{w \in \operatorname{supp}(P_W)} P_W(w)^{\alpha}$$
(120)

for  $\alpha \notin \{-\infty, 1, \infty\}$  and where  $H_{-\infty} = \lim_{\alpha \downarrow -\infty} H_{\alpha}, H_{\infty} = \lim_{\alpha \uparrow \infty} H_{\alpha}$ , and  $H_1 = \lim_{\alpha \uparrow 1} H_{\alpha}$ . The  $\alpha$ -Rényi entropy is a natural generalization of the Shannon entropy. For  $\pi_{XY}$ , we define the *common*  $\alpha$ -*Rényi entropy* with  $\alpha \in [-\infty, \infty]$  as

$$G_{\alpha}(\pi_{XY}) := \min_{P_{W}P_{X|W}P_{Y|W}: P_{XY} = \pi_{XY}} H_{\alpha}(W).$$
(121)

(The common  $\alpha$ -Rényi entropy is a generalization of the common entropy [6], [8]; see the definition of the common entropy in (4)). The exact common information can be generalized to

the exact  $\alpha$ -Rényi common information with  $\alpha \in [-\infty, \infty]$ , which is defined as

$$T_{\text{Exact}}^{(\alpha)}(\pi_{XY}) := \lim_{n \to \infty} \frac{1}{n} G_{\alpha}(\pi_{XY}^n).$$
(122)

Here the existence of the limit in (122) follows by the subadditivity of the sequence of  $\{G_{\alpha}(\pi_{XY}^n)\}_{n\in\mathbb{N}}$ .

Since  $H_{\alpha}$  is non-increasing in  $\alpha \in [-\infty, \infty]$ , we have that  $G_{\alpha}(\pi_{XY})$  and  $T_{\text{Exact}}^{(\alpha)}(\pi_{XY})$  are also non-increasing in  $\alpha \in [-\infty, \infty]$ . Furthermore, for  $\alpha \in \{0, 1, \infty\}$ , we have the following characterization of  $G_{\alpha}(\pi_{XY})$  and  $T_{\text{Exact}}^{(\alpha)}(\pi_{XY})$ . The proof is provided in Appendix J.

Proposition 2. We have

$$G_{\alpha}(\pi_{XY}) = \begin{cases} \log \operatorname{rank}^{+}(\pi_{XY}), & \alpha = 0\\ \min_{P_{W}P_{X|W}P_{Y|W}:P_{XY} = \pi_{XY}} H(W), & \alpha = 1\\ \min_{Q_{X},Q_{Y}} D_{\infty}(Q_{X}Q_{Y} \| \pi_{XY}), & \alpha = \infty \end{cases}$$
(123)

where rank<sup>+</sup>(**A**) denotes the nonnegative rank of a matrix **A**, *i.e.*, the minimum  $k \in \mathbb{N}$  such that there exist nonnegative

matrices  $\mathbf{U} \in \mathbb{R}_{\geq 0}^{k \times |\mathcal{X}|}, \mathbf{V} \in \mathbb{R}_{\geq 0}^{k \times |\mathcal{Y}|}$  satisfying  $\mathbf{U}\mathbf{V}^{\top} = \mathbf{A}$ . Furthermore, for  $\alpha = \infty$ ,

$$T_{\text{Exact}}^{(\infty)}(\pi_{XY}) = \min_{Q_X, Q_Y} D_{\infty}(Q_X Q_Y \| \pi_{XY}).$$

By definition, the exact 0-Rényi common information corresponds to the minimum common randomness rate for exact generation of the target distribution in which the common randomness is only allowed to be compressed by *fixed-length* codes. By (123), the exact 0-Rényi common information can be expressed as

$$T_{\text{Exact}}^{(0)}(\pi_{XY}) = \lim_{n \to \infty} \frac{1}{n} \log \operatorname{rank}^+(\pi_{XY}^{\otimes n}), \qquad (124)$$

where  $\pi_{XY}^{\otimes n}$  denotes the Kronecker product of *n* copies of the matrix  $\pi_{XY}$ . That is,  $T_{\text{Exact}}^{(0)}(\pi_{XY})$  is the exponent of  $\operatorname{rank}^+(\pi_{XY}^{\otimes n})$  as  $n \to \infty$ . By definition, we can easily obtain

$$\log \operatorname{rank}(\pi_{XY}) \le T_{\operatorname{Exact}}^{(0)}(\pi_{XY}) \le \log \operatorname{rank}^+(\pi_{XY}).$$
(125)

Yannakakis [36] first related the nonnegative rank of a matrix to the communication complexity (the minimum number of communication bits) of distributively computing a matrix (or a bivariate function). The equivalence between the nonnegative rank and the exact common information (when only fixedlength codes allowed), as shown in (124), was previously obtained in [37], [38].

#### • Nonnegative $\alpha$ -Rank

The class of common information problems can also be cast in the light of approximate or exact decomposition of a joint distribution. Let  $\mathbf{P}_W$  be the diagonal matrix with the probability values of  $P_W$  as its diagonal elements and let  $\top$  denote the transposition operation. The exact common information problem is equivalent to decomposing a joint distribution as a mixture of product conditional distributions

$$\mathbf{P}_{XY} = \mathbf{P}_{X|W}^{\top} \mathbf{P}_{W} \mathbf{P}_{Y|W}$$
(126)

such that the entropy  $H(\mathbf{P}_W)$  is minimized. Such a decomposition is closely related to nonnegative matrix factorization (NMF) and the nonnegative rank [39]. The nonnegative rank and NMF play a crucial role in many subdisciplines of theoretical computer science and discrete mathematics, including signal processing, machine learning, communication complexity, and combinatorial optimization, e.g., [40].

Recall the common  $\alpha$ -Rényi entropy defined in (121). When  $\alpha = 0$ , this quantity is equal to the logarithm of nonnegative rank of the joint distribution matrix. Inspired by this relationship, we can generalize the nonnegative rank to the nonnegative  $\alpha$ -rank as follows. For a nonnegative matrix (but not zero matrix) **A** and  $\alpha \in [-\infty, \infty]$ , we define the *nonnegative*  $\alpha$ -rank of **A** as

$$\operatorname{rank}_{\alpha}^{+}(\mathbf{A}) := \exp\left\{G_{\alpha}\left(\frac{\mathbf{A}}{\|\mathbf{A}\|_{1}}\right)\right\}.$$
 (127)

Here, note that the argument of  $G_{\alpha}$  is the normalized version of the matrix **A** because the argument of  $G_{\alpha}$  needs to be a joint probability distribution. This makes sense since any reasonable definition of "rank" should satisfy invariance under scaling operations (with non-zero scale factors). When  $\alpha = 0$ , the nonnegative 0-rank defined in (127) reduces to the traditional nonnegative rank, i.e.,  $\operatorname{rank}_0^+(\mathbf{A}) = \operatorname{rank}^+(\mathbf{A})$ . Equivalently, the nonnegative  $\alpha$ -rank  $\operatorname{rank}_{\alpha}^+(\mathbf{A})$  can be alternatively expressed as<sup>10</sup>

$$\operatorname{rank}_{\alpha}^{+}(\mathbf{A}) = \min_{\mathbf{U}, \mathbf{D}, \mathbf{V}} \|\mathbf{D}\|_{\alpha}^{\frac{1}{1-\alpha}}$$
(128)

where the minimization in (128) is taken over all nonnegative matrices  $\mathbf{U} \in \mathbb{R}_{\geq 0}^{k \times |\mathcal{X}|}, \mathbf{D} \in \mathbb{R}_{\geq 0}^{k \times k}, \mathbf{V} \in \mathbb{R}_{\geq 0}^{k \times |\mathcal{Y}|}$  for some  $k \in \mathbb{N}$  such that  $\mathbf{D}$  is diagonal and  $\mathbf{U}^{\top} \mathbf{D} \mathbf{V} = \frac{\mathbf{A}}{\|\mathbf{A}\|_{1}}$ .

Here we only provide the definition of nonnegative  $\alpha$ -rank. Investigation on applications of nonnegative  $\alpha$ -rank is outside the scope of this paper, which remains to be done in the future.

#### VI. CONCLUDING REMARKS

In this paper, we established the equivalence between the exact and  $\infty$ -Rényi common informations; provided single-letter upper and lower bounds on these two quantities; completely characterized them for DSBSes; and extended the exact and  $\infty$ -Rényi common informations, and also Wyner's common information to sources with general (countable or continuous) alphabets, including Gaussian sources.

For DSBSes, we observed that the exact and  $\infty$ -Rényi common informations are both strictly larger than Wyner's common information. This resolves an open problem posed by Kumar, Li, and El Gamal [6]. For Gaussian sources with correlation coefficient  $\rho \in [0, 1)$ , we provided an upper bound on the exact and  $\infty$ -Rényi common informations, which is at most 0.72 (exactly,  $\frac{\rho}{1+\rho} \log_2 e$ ) bits/symbol larger than Wyner's common information, and at least 22.28 bits/symbol smaller than Li and El Gamal's one-shot bound [10]. We conjectured our upper bound is tight.

Due to the equivalence between the exact common information and exact channel simulation, we apply our results on the former problem to the latter problem. In [5], [31]– [33], it was shown that when there exists unlimited shared randomness, the minimum communication rates are the same for TV-approximate and exact channel simulation problems, and this rate is equal to the mutual information. However, this is not the case when there is no shared randomness. Our results imply that with no shared randomness, the minimum communication rate for TV-approximate channel simulation is Wyner's common information; however the minimum rate for exact channel simulation is the exact common information which is larger than Wyner's common information.

We also connected the common information problem to the distributed channel synthesis problem. Our results imply that with no shared randomness, the minimum rate for exact channel simulation is the exact common information which is larger than Wyner's common information. When there is randomness shared by the encoder and decoder, the best tradeoff between the shared randomness rate and the communication

<sup>&</sup>lt;sup>10</sup>One can also define a variant  $\overline{\operatorname{rank}}^+_{\alpha}(\mathbf{A})$  of the nonnegative  $\alpha$ -rank by replacing  $\|\mathbf{D}\|^{\frac{\alpha}{1-\alpha}}_{\alpha}$  with  $\|\mathbf{D}\|^{\alpha}_{\alpha}$ . This variant can be written as  $\overline{\operatorname{rank}}^+_{\alpha}(\mathbf{A}) = (\operatorname{rank}^+_{\alpha}(\mathbf{A}))^{1-\alpha}$  with  $\operatorname{rank}^+_{\alpha}(\mathbf{A})$  denoting the nonnegative  $\alpha$ -rank defined in (127) or (128). Hence  $\overline{\operatorname{rank}}^+_{\alpha}(\mathbf{A})$  and  $\operatorname{rank}^+_{\alpha}(\mathbf{A})$  are uniquely determined by each other except for  $\alpha \in \{-\infty, 1, \infty\}$ .

rate were studied in our paper [24]. In the future, we are planning to work on various closely-related problems, e.g., the exact versions of various coordination problems [41].

# APPENDIX A Proof of Theorem 1

A. Proof of  $T_{\text{Exact}}(\pi_{XY}) = T_{\infty}(\pi_{XY})$ 

One direction of the equivalence  $T_{\text{Exact}}(\pi_{XY}) = T_{\infty}(\pi_{XY})$  follows from the following lemmas.

**Lemma 4.** [6] If there exists a sequence of fixed-length synthesis codes with rate R that generates  $P_{X^nY^n}$  such that  $D_{\infty}(P_{X^nY^n} || \pi_{XY}^n) \to 0$ , then there must exist a sequence of variable-length synthesis codes with asymptotic rate R that exactly generates  $\pi_{XY}^n$ . That is,  $T_{\text{Exact}}(\pi_{XY}) \leq T_{\infty}(\pi_{XY})$ .

This lemma was proven by Kumar, Li, and El Gamal in [6, Remark on Page 164] using the following mixture decomposition technique<sup>11</sup> (also termed "splitting technique"). According to the definition of  $D_{\infty}$ ,  $D_{\infty}(P_{X^nY^n} || \pi_{XY}^n) \leq \epsilon$ with  $\epsilon > 0$  implies that  $P_{X^nY^n}(x^n, y^n) \leq e^{\epsilon} \pi_{XY}^n(x^n, y^n)$ for all  $x^n, y^n$ . Define

$$\widehat{P}_{X^{n}Y^{n}}(x^{n}, y^{n}) := \frac{e^{\epsilon} \pi_{XY}^{n}(x^{n}, y^{n}) - P_{X^{n}Y^{n}}(x^{n}, y^{n})}{e^{\epsilon} - 1},$$
(129)

then obviously,  $\hat{P}_{X^nY^n}(x^n, y^n)$  is a distribution. Hence  $\pi_{XY}^n$  can be written as a mixture distribution

$$\pi_{XY}^{n}(x^{n}, y^{n}) = e^{-\epsilon} P_{X^{n}Y^{n}}(x^{n}, y^{n}) + (1 - e^{-\epsilon}) \widehat{P}_{X^{n}Y^{n}}(x^{n}, y^{n}).$$
(130)

The encoder first generates a Bernoulli random variable U with  $P_U(1) = e^{-\epsilon}$ , compresses it by using 1 bit, and transmits it to the two generators. If U = 1, then the encoder generates a uniform random variable  $M \sim \text{Unif}[1 : e^{nR}]$ , and the encoder and two generators use the fixed-length synthesis codes with rate R to generate  $P_{X^nY^n}$ . If U = 0, then the encoder generates  $(X^n, Y^n) \sim \widehat{P}_{X^nY^n}$ , and uses a variable-length compression code with rate  $\leq \log |\mathcal{X}||\mathcal{Y}|$  to generate  $\widehat{P}_{X^nY^n}$ . The distribution generated by such a mixed code is  $e^{-\epsilon}P_{X^nY^n}(x^n, y^n) + (1 - e^{-\epsilon})\widehat{P}_{X^nY^n}(x^n, y^n)$ , i.e.,  $\pi_{XY}^n(x^n, y^n)$ . The total code rate is no larger than  $\frac{1}{n} + e^{-\epsilon}R + (1 - e^{-\epsilon})\log |\mathcal{X}||\mathcal{Y}|$ , which converges to R upon taking the limit in  $n \to \infty$  and the limit in  $\epsilon \to 0$ .

The mixture decomposition (or split) of a distribution in (130) can be generalized to general distributions.

**Lemma 5** (Mixture Decomposition of General Distributions). Assume P, Q are two distributions defined on the same Borelmeasurable space. Assume<sup>12</sup>  $D_{\infty}(P||Q) \leq \epsilon$  for some  $\epsilon \in [0, \infty]$ . Then

$$Q = e^{-\epsilon}P + \left(1 - e^{-\epsilon}\right)\widehat{P},\tag{131}$$

where

$$\widehat{P} := \begin{cases} any \ distribution \quad \epsilon = 0\\ \frac{e^{\epsilon}Q - P}{e^{\epsilon} - 1} \qquad \epsilon \in (0, \infty) \\ Q \qquad \epsilon = \infty \end{cases}$$
(132)

Moreover, if we define

$$\Lambda(Q, P) := \sup \Big\{ \alpha : \exists \ a \ distribution \ \widehat{P} \ s.t. \\ Q = \alpha P + (1 - \alpha) \ \widehat{P}, \alpha \in [0, 1] \Big\},$$
(133)

then

$$\Lambda(Q,P) = e^{-D_{\infty}(P \parallel Q)} = \begin{cases} \frac{1}{\operatorname{ess\,sup}_{P} \frac{\mathrm{d}P}{\mathrm{d}Q}} & P \ll Q\\ 0 & P \not\ll Q \end{cases}.$$
 (134)

*Remark* 14. Given a set of distributions  $\{P_i : i \in [1:n]\}$  and a target distribution Q defined on the same space  $(\mathcal{Y}, \mathbb{B})$ , a natural question is to determine the minimum value  $\alpha_0 \ge 0$  such that

$$Q = \sum_{i=1}^{n} \alpha_i P_i + \alpha_0 \widehat{P} \tag{135}$$

for some distribution  $\widehat{P}$  and some values  $\alpha_i \ge 0, i \in [1:n]$ and  $\sum_{i=0}^{n} \alpha_i = 1$ . By Lemma 5 such a mixture decomposition problem is equivalent to

$$\min_{\{\hat{\alpha}_i\}:\hat{\alpha}_i \ge 0, \sum_{i=1}^n \hat{\alpha}_i = 1} D_{\infty}(\sum_{i=1}^n \hat{\alpha}_i P_i \| Q).$$
(136)

If we consider  $\{P_i\}$  as a channel  $P_{Y|X}$  with  $P_{Y|X=i} = P_i$ and denote  $Q_Y := Q$ , then (136) can be rewritten as

$$\min_{P_X} D_{\infty}(P_Y \| Q_Y) \tag{137}$$

where  $P_X$  is a distribution on [1:n] and  $P_Y$  is the output distribution of  $P_{Y|X}$  when the input distribution is  $P_X$ . The problem in (137) is just the so-called  $\infty$ -*Rényi resolvability* problem (or channel resolvability problem under  $\infty$ -*Rényi* divergence measure). In [25], the present authors studied the  $\infty$ -Rényi resolvability problem in which the channels and target distributions are of product forms and  $P_X$  restricted to be a function of a given uniform random variable.

Now we consider the other direction of the equivalence  $T_{\text{Exact}}(\pi_{XY}) = T_{\infty}(\pi_{XY}).$ 

**Lemma 6.** If there exists a sequence of variable-length synthesis codes with asymptotic rate R that exactly generates  $\pi_{XY}^n$ , then there must exist a sequence of fixed-length synthesis codes with rate R that generates  $P_{X^nY^n}$  such that  $D_{\infty}(P_{X^nY^n} \| \pi_{XY}^n) \to 0$ . That is,  $T_{\text{Exact}}(\pi_{XY}) \ge T_{\infty}(\pi_{XY})$ .

<sup>12</sup>For general distributions P, Q such that  $P \ll Q$ ,  $D_{\infty}(P||Q) := \log \operatorname{ess\,sup}_P \frac{\mathrm{d}P}{\mathrm{d}Q}$ , where  $\frac{\mathrm{d}P}{\mathrm{d}Q}$  denotes Radon–Nikodym derivative of P respect to Q, and  $\operatorname{ess\,sup}_P \frac{\mathrm{d}P}{\mathrm{d}Q}$  denotes the essential supremum of  $\frac{\mathrm{d}P}{\mathrm{d}Q}$  with respect to P. Moreover, if  $P \ll Q$ , then  $D_{\infty}(P||Q) := +\infty$ .

<sup>&</sup>lt;sup>11</sup>The decomposition of a distribution into a mixture of several distributions, as in (130) and (131), is termed the mixture decomposition (or split) of a distribution. This mixture decomposition is rather useful to construct a desired distribution from a given one. Such an idea originated from Nummelin' work [14] and Athreya and Ney's work [15]. In both of [14] and [15], the authors used this splitting technique to study limiting theorems of recurrent Markov processes. Furthermore, such a technique was also used to study the mixing rate of Markov Chain Monte Carlo (MCMC) methods [16], by constructing a coupling of an original Markov chain and an target Markov chain. Besides as a tool, the mixture decomposition is also an important topic in probability and statistics theories that has independent interest; see [42] (or more general decomposition theories [43]). The mixture decomposition is also related to other information-theoretic problems. For example, such a technique was used in the proof of [17, Theorem 16]. Furthermore as mentioned in Remark 14, finding an optimal mixture decomposition (with the minimum coefficient for the residual part) is equivalent to the  $\infty$ -Rényi resolvability problem [25].

Remark 15. Note that by checking our proof, one can find that this lemma holds not only for sources with finite alphabets, but also for those with countably infinite or continuous/uncountable alphabets.

*Proof:* Let  $\{c_k\}_{k=1}^{\infty}$  be a sequence of variable-length codes with rate R that exactly generates  $\pi_{XY}^k$ . Let  $W_k$  be the common random variable, and  $P_{X^k|W_k}$ and  $P_{Y^k|W_k}$  the two generators that define  $c_k$ . Hence  $\sum_{w} P_{W_k}(w) P_{X^k|W_k}(\cdot|w) P_{Y^k|W_k}(\cdot|w) = \pi_{XY}^k, \text{ and } \frac{1}{k} H(W_k) \to R \text{ as } k \to \infty. \text{ Now we consider a superblock}$ =  $\pi^k_{XY}$ , and code that consists of n independent k-length codes as defined above. That is,  $W_k^n \sim P_{W_k}^n$  is the common random variable and  $P_{X^k|W_k}^n$  and  $P_{Y^k|W_k}^n$  are the two generators. Observe that  $W_k^n$  is an *n*-length i.i.d. random sequence with each  $W_{k,i} \sim P_{W_k}$ . Hence we have

$$\mathbb{P}\left(W_k^n \in \mathcal{A}_{\epsilon}^{(n)}(P_{W_k})\right) \to 1$$
(138)

as  $n \to \infty$  for fixed k. Furthermore,  $|\mathcal{A}_{\epsilon}^{(n)}| \leq e^{n(H(W_k) + \epsilon)}$ . Define a truncated distribution

$$Q_{W_k^n}\left(w_k^n\right) := \frac{P_{W_k}^n\left(w_k^n\right) \mathbf{1}\left\{w_k^n \in \mathcal{A}_{\epsilon}^{(n)}\right\}}{P_{W_k}^n\left(\mathcal{A}_{\epsilon}^{(n)}\right)}.$$
(139)

Now we adopt a simulation scheme  $f_n$  as used in [44, Theorem 7] to simulate the truncated distribution  $Q_{W_{k}^{n}}$  from a uniform random variable  $M \sim \text{Unif}[1 : e^{nkR^{\delta}}]$ . For each  $w_k^n \in \mathcal{A}_{\epsilon}^{(n)}$ , we map either  $|e^{nkR'}Q_{W_k^n}(w_k^n)|$  or  $\left[e^{nkR'}Q_{W_{k}^{n}}\left(w_{k}^{n}\right)\right]$  number of elements  $m \in \left[1 : e^{nkR'}\right]$ to it. Hence the output distribution  $\widetilde{P}_{W_k^n}$  induced by such a mapping satisfies  $\widetilde{P}_{W_k^n}(w_k^n) = e^{-nkR'} [e^{nkR'}Q_{W_k^n}(w_k^n)]$ or  $\widetilde{P}_{W_k^n}(w_k^n) = e^{-nkR'} [e^{nkR'}Q_{W_k^n}(w_k^n)]$  for  $w_k^n \in \mathcal{A}_{\epsilon}^{(n)}$ . Hence

$$D_{\infty}(\tilde{P}_{W_{k}^{n}} \| Q_{W_{k}^{n}})$$

$$= \log \max_{w_{k}^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{\tilde{P}_{W_{k}^{n}}(w_{k}^{n})}{Q_{W_{k}^{n}}(w_{k}^{n})}$$
(140)

$$\leq \log \max_{w_{k}^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{Q_{W_{k}^{n}}(w_{k}^{n}) + e^{-nkR'}}{Q_{W_{k}^{n}}(w_{k}^{n})}$$
(141)

$$\leq \log\left(1 + \frac{e^{-nkR'}}{e^{-n(H(W_k)+\epsilon)}}\right) \tag{142}$$

$$= \log\left(1 + e^{-nk\left(R' - \frac{1}{k}\left(H(W_k) + \epsilon\right)\right)}\right).$$
(143)

Therefore, if  $R' > \frac{1}{k} (H(W_k) + \epsilon)$ , then

$$D_{\infty}(\widetilde{P}_{W_{k}^{n}} \| Q_{W_{k}^{n}})$$

$$= \log \max_{w_{k}^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{\widetilde{P}_{W_{k}^{n}}(w_{k}^{n})}{Q_{W_{k}^{n}}(w_{k}^{n})} \to 0, \qquad (144)$$

as  $n \to \infty$  for fixed k. Such a simulation code  $f_n$  is also valid for simulating  $P_{W_{h}}^{n}$ . This is because

$$D_{\infty}(\widetilde{P}_{W_{k}^{n}} \| P_{W_{k}}^{n})$$

$$= \log \max_{w_{k}^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{\widetilde{P}_{W_{k}^{n}}(w_{k}^{n})}{P_{W_{k}}^{n}(w_{k}^{n})}$$
(145)

$$\leq \log \max_{w_k^n \in \mathcal{A}_{\epsilon}^{(n)}} \frac{\widetilde{P}_{W_k^n}(w_k^n)}{Q_{W_k^n}(w_k^n)} + \log \max_{w_k^n \in \mathcal{A}_{\epsilon}^{(n)}} \frac{Q_{W_k^n}(w_k^n)}{P_{W_k}^n(w_k^n)}$$
(146)

$$= D_{\infty}(\widetilde{P}_{W_{k}^{n}} \| Q_{W_{k}^{n}}) - \log P_{W_{k}}^{n} \left( \mathcal{A}_{\epsilon}^{(n)} \right)$$
(147)

$$\rightarrow 0,$$
 (148)

as  $n \to \infty$  for fixed k.

Now we consider a cascaded synthesis code by concatenating the simulation code  $f_n$  above with the two generators  $P_{X^k|W_k}^n$  and  $P_{Y^k|W_k}^n$  of the variable-length synthesis code. Observe that  $P_{X^{kn}Y^{kn}}$  and  $\pi_{XY}^{kn}$  are respectively the outputs of the channel  $P_{X^k|W_k}^n P_{Y^k|W_k}^n$  respectively induced by the channel inputs  $P_{W_k^n}$  and  $P_{W_k}^n$ . Hence by the data processing inequality [27], for such a cascaded code, we have

$$D_{\infty}(P_{X^{kn}Y^{kn}} \| \pi_{XY}^{kn})$$

$$\leq D_{\infty}(\widetilde{P}_{W_k^n} \| P_{W_k}^n) \qquad (149)$$

$$\to 0 \qquad (150)$$

as  $n \to \infty$  for fixed k, as long as the code rate R' > $\frac{1}{k}(H(W_k)+\epsilon).$ 

As for the case where the blocklength n' is not a multiple of k, i.e., n' = kn + l with  $l \in [1 : k - 1]$ , we need to construct a code with blocklength k(n+1) and then truncate the outputs  $(X^{k(n+1)}, Y^{k(n+1)})$  to  $(X^{n'}, Y^{n'})$ . Obviously,  $D_{\infty}(P_{X^{n'}Y^{n'}} \| \pi_{XY}^{n'}) \leq D_{\infty}(P_{X^{k(n+1)}Y^{k(n+1)}} \| \pi_{XY}^{k(n+1)}) \to 0$ as  $n \to \infty$ . Furthermore, the code rate for such a code is  $\frac{k(n+1)R'}{n'} \leq (1+\frac{1}{n})R' \to R' \text{ as } n \to \infty.$  On the other hand,  $\frac{1}{k}H(W_k) \to R$  as  $k \to \infty$ . Therefore, there exists a sequence of fixed-length synthesis codes with asymptotic rate R that generates  $P_{X^{n'}Y^{n'}}$  such that  $D_{\infty}(P_{X^{n'}Y^{n'}} \| \pi_{XY}^{n'}) \to 0$ as  $n' \to \infty$ .

# B. Proof of $T_{\infty}(\pi_{XY}) \leq \liminf_{n \to \infty} \frac{1}{n} \Gamma(\pi_{XY}^n)$

Here we prove the achievability result from the perspective of  $\infty$ -Rényi common information problem. We borrow an idea from [9]. The corresponding coding scheme was also independently used by the present authors in [3], [25].

To show the achievability part, we only need to show that the single-letter expression  $\Gamma(\pi_{XY})$  satisfies  $T_{\infty}(\pi_{XY}) \leq$  $\Gamma(\pi_{XY})$ . This is because we can obtain the upper bound  $\Gamma(\pi_{XY}^k)$  by substituting  $\pi_{XY}$  with  $\pi_{XY}^k$  into the single-letter expression<sup>13</sup>. For fixed  $\epsilon > 0$  and a fixed joint distribution

<sup>&</sup>lt;sup>13</sup>Note that by definition,  $T_{\infty}(\pi_{XY}^k)$  is additive in k, i.e.,  $T_{\infty}(\pi_{XY}^k) = kT_{\infty}(\pi_{XY})$ . This is because, on one hand, the superblock code that consists of k independent copies of a (n, R) code for  $\pi_{XY}$  forms a (n, kR) code for  $\pi_{XY}^k$ . On the other hand, a (n, kR) code for  $\pi_{XY}^k$  forms a (nk, R) code for  $\pi_{XY}$ .

 $Q_{WXY} = Q_W Q_{X|W} Q_{Y|W}$ , define the distributions

$$P_{W^{n}}\left(w^{n}\right) \propto Q_{W}^{n}\left(w^{n}\right) \mathbb{1}\left\{w^{n} \in \mathcal{T}_{\frac{\epsilon}{2}}^{\left(n\right)}\left(Q_{W}\right)\right\},$$
(151)

$$P_{X^{n}|W^{n}}(x^{n}|w^{n}) \propto Q_{X|W}^{n}(x^{n}|w^{n})$$

$$\times 1\left\{x^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WX}|w^{n})\right\}, (152)$$

$$P_{Y^{n}|W^{n}}(y^{n}|w^{n}) \propto Q_{Y|W}^{n}(y^{n}|w^{n})$$

$$\times 1\left\{y^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WY}|w^{n})\right\}. (153)$$

We set  $C_n = \{W^n(m)\}_{m \in \mathcal{M}_n}$  with  $W^n(m), m \in \mathcal{M}_n$ drawn independently for different *m*'s and according to the same distribution  $P_{W^n}$ . Upon receiving  $W^n(M_n)$ , the generators respectively use random mappings  $P_{X^n|W^n}$  and  $P_{Y^n|W^n}$ to generate  $X^n$  and  $Y^n$ . Define  $P_{\mathcal{M}_n} := \text{Unif}[1 : e^{nR}]$ . For random mappings  $(P_{X^n|W^n}, P_{Y^n|W^n})$ , we define

$$P_{X^{n}Y^{n}|\mathcal{C}_{n}}(x^{n}, y^{n}|\{W^{n}(m)\})$$
  
:=  $\sum_{m} P_{M_{n}}(m)P_{X^{n}|W^{n}}(x^{n}|W^{n}(m))P_{Y^{n}|W^{n}}(y^{n}|W^{n}(m))$   
(154)

which is the output distribution induced by the codebook  $C_n$  in a distributed source simulation system with simulators  $(P_{X^n|W^n}, P_{Y^n|W^n})$ . For such a code, we have the following distributed Rényi-covering lemma.

**Lemma 7** (Distributed Rényi-Covering). *For the random code described above, if* 

$$R > \mathcal{I}(Q) := -H_Q(XY|W) + \sum_{w} Q(w) \mathcal{H}(Q_{X|W=w}, Q_{Y|W=w} ||Q_{XY}), \quad (155)$$

then there exists some  $\alpha, \epsilon > 0$  such that

$$\mathbb{P}_{\mathcal{C}_n}\left(D_{\infty}(P_{X^nY^n|\mathcal{C}_n} \| Q_{XY}^n) \le e^{-n\alpha}\right) \to 1$$
(156)

#### doubly exponentially fast.

*Remark* 16. The soft-covering problem under the  $\infty$ -Rényi divergence measure was also studied in [9] as a key step of proving sufficient conditions for equality of Wyner's common information and the exact common information. However, no explicit rate bound (e.g.,  $\mathcal{I}(Q)$  as defined in (155)) for an arbitrary  $\pi_{XY}$  was provided in [9].

Setting  $Q_{WXY}$  as an optimal distribution attaining  $\Gamma(\pi_{XY})$ , we obtain  $\mathcal{I}(Q) = \Gamma(\pi_{XY})$ . Hence this lemma implies that there exists a sequence of codebooks  $\{c_n\}$  with rate R such that  $D_{\infty}(P_{X^nY^n|\mathcal{C}_n=c_n} || Q_{XY}^n) \leq e^{-n\alpha}$  as long as R > $\Gamma(\pi_{XY})$ . This completes the proof of  $T_{\infty}(\pi_{XY}) \leq \Gamma(\pi_{XY})$ . Hence what we need to do is to prove Lemma 7. The proof is provided in the following. *Proof of Lemma 7:* For the fixed  $\epsilon > 0$  and the fixed joint distribution  $Q_{WXY} = Q_W Q_{X|W} Q_{Y|W}$ , define

$$\mathcal{B}_{\epsilon} := \left\{ P_{WXY} \in \mathcal{P}(\mathcal{W} \times \mathcal{X} \times \mathcal{Y}) : \\ \forall w, |P_{W}(w) - Q_{W}(w)| \leq \frac{\epsilon}{2} Q_{W}(w), \\ \forall (w, x), |P_{WX}(w, x) - Q_{WX}(w, x)| \leq \epsilon Q_{WX}(w, x), \\ \forall (w, y), |P_{WY}(w, y) - Q_{WY}(w, y)| \leq \epsilon Q_{WY}(w, y) \right\},$$

$$(157)$$

and

$$\mathcal{I}_{\epsilon}\left(Q\right) := \max_{\widetilde{P}_{WXY} \in \mathcal{B}_{\epsilon}} \sum_{w,x} \widetilde{P}\left(w,x\right) \log Q\left(x|w\right) \\ + \sum_{w,y} \widetilde{P}\left(w,y\right) \log Q\left(y|w\right) \\ - \sum_{x,y} \widetilde{P}\left(x,y\right) \log Q\left(x,y\right).$$
(158)

Obviously,  $\mathcal{I}_{\epsilon}(Q) \geq \mathcal{I}(Q)$ , hence  $\lim_{\epsilon \downarrow 0} \mathcal{I}_{\epsilon}(Q) \geq \mathcal{I}(Q)$ , where  $\mathcal{I}(Q)$  is defined in (155). Now we prove  $\lim_{\epsilon \downarrow 0} \mathcal{I}_{\epsilon}(Q) \leq \mathcal{I}(Q)$ . Let  $\{\epsilon_k\}_{k=1}^{\infty}$  be a sequence of decreasing positive numbers with  $\lim_{k \to \infty} \epsilon_k = 0$ . Let  $\{P_{WXY}^{(k)}\}_{k=1}^{\infty}$  be a sequence of optimal distributions attaining  $\widetilde{\Gamma}_{\epsilon_k}(\pi_{XY})$ . Since  $\mathcal{P}(\mathcal{W} \times \mathcal{X} \times \mathcal{Y})$  is compact, there exists some subsequence  $\{\epsilon_{k_i}\}_{i=1}^{\infty}$  such that  $P_{WXY}^{(k_i)}$  converges to some distribution  $\widehat{P}_{WXY}$  as  $i \to \infty$ . Since  $\lim_{i \to \infty} \epsilon_{k_i} = 0$ , we must have

$$\widehat{P}_{WX} = Q_{WX} \tag{159}$$

$$\widehat{P}_{WY} = Q_{WY}.\tag{160}$$

Since the objective function in the right hand side of (158) is continuous in  $\tilde{P}_{WXY}$ , we have

$$\lim_{\epsilon \downarrow 0} \mathcal{I}_{\epsilon} (Q)$$

$$= \sum_{w,x} \widehat{P}(w,x) \log Q(x|w) + \sum_{w,y} \widehat{P}(w,y) \log Q(y|w)$$

$$- \sum_{x,y} \widehat{P}(x,y) \log Q(x,y)$$
(161)

$$\leq \max_{\widetilde{P}_{WXY}:\widetilde{P}_{WX}=Q_{WX},\widetilde{P}_{WY}=Q_{WY}} \sum_{w,x} \widetilde{P}(w,x) \log Q(x|w) + \sum_{w,y} \widetilde{P}(w,y) \log Q(y|w) - \sum_{x,y} \widetilde{P}(x,y) \log Q(x,y)$$
(162)

$$=\mathcal{I}\left(Q\right).\tag{163}$$

Therefore,

$$\lim_{\epsilon \downarrow 0} \mathcal{I}_{\epsilon} \left( Q \right) = \mathcal{I} \left( Q \right). \tag{164}$$

By the continuity of  $\epsilon \mapsto \mathcal{I}_{\epsilon}(Q)$  shown in (164), we can choose  $\epsilon > 0$ , used in definitions (151)-(153), so small such that

$$R > \mathcal{I}_{\epsilon}\left(Q\right) + \epsilon. \tag{165}$$

The reason for this choice of  $\epsilon$  is to ensure (189) (at the end of this proof) to hold.

For brevity, in the following we denote  $M = e^{nR}$ . According to the definition of the Rényi divergence, we first have<sup>14</sup>

$$e^{D_{\infty}(P_{X^{n_{Y^n}}} \| \pi_{XY}^n)} = \max_{x^n, y^n} \frac{P_{X^n Y^n}(x^n, y^n)}{Q_{XY}^n(x^n, y^n)}$$
(166)

$$= \max_{x^n, y^n} \widetilde{g}(x^n, y^n | \mathcal{C}_n), \tag{167}$$

where  $\widetilde{g}(x^n, y^n | \mathcal{C}_n) := \sum_{m \in \mathcal{M}_n} \frac{1}{\mathsf{M}} g(x^n, y^n | W^n(m))$  with  $g(x^n, y^n | w^n) := \frac{1}{Q_{XY}^n(x^n, y^n)} P_{X^n | W^n}(x^n | w^n) P_{Y^n | W^n}(y^n | w^n).$ Then for  $w^n \in \mathcal{T}^n_{\frac{\epsilon}{2}}(Q_W)$ ,

$$g(x^{n}, y^{n}|w^{n}) = \frac{1}{Q_{XY}^{n}(x^{n}, y^{n})} \frac{Q_{X|W}^{n}(x^{n}|w^{n}) 1\left\{x^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WX}|w^{n})\right\}}{Q_{X|W}^{n}\left(\mathcal{T}_{\epsilon}^{(n)}(Q_{WX}|w^{n})|w^{n}\right)} \\ \times \frac{Q_{Y|W}^{n}(y^{n}|w^{n}) 1\left\{y^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WY}|w^{n})\right\}}{Q_{Y|W}^{n}\left(\mathcal{T}_{\epsilon}^{(n)}(Q_{WY}|w^{n})|w^{n}\right)}$$
(168)  
$$\leq \frac{1\left\{x^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WX}|w^{n}), y^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WY}|w^{n})\right\}}{(1-\delta_{1,n})(1-\delta_{2,n})} \\ \times e^{n\sum_{w,y} T_{w^{n}x^{n}}(w,y) \log Q(y|w)} \\ \times e^{-n\sum_{x,y} T_{x^{n}y^{n}}(x,y) \log Q(x,y)}$$
(169)

$$\leq \frac{1}{(1-\delta_{1,n})(1-\delta_{2,n})}e^{n\mathcal{I}_{\epsilon}(Q)}$$
(170)

$$=:\beta_n,\tag{171}$$

where by [3, Lemma 4], both  $\delta_{1,n} := 1 - Q_{X|W}^n \left( \mathcal{T}_{\epsilon}^{(n)} \left( Q_{WX} | w^n \right) | w^n \right)$  and  $\delta_{2,n} := 1 - Q_{Y|W}^n \left( \mathcal{T}_{\epsilon}^{(n)} \left( Q_{WY} | w^n \right) | w^n \right)$  converge to zero exponentially fast as  $n \to \infty$ , and  $\mathcal{I}_{\epsilon} \left( Q \right)$  is defined in (158).

Continuing (167), we get for any  $\delta > 0$ ,

$$\mathbb{P}_{\mathcal{C}_n}\left(D_{\infty}(P_{X^nY^n} \| \pi_{XY}^n) \ge \delta\right)$$
  
$$\leq \mathbb{P}_{\mathcal{C}_n}\left(e^{D_{\infty}(P_{X^nY^n} \| \pi_{XY}^n)} - 1 \ge \delta\right)$$
(172)

$$= \mathbb{P}_{\mathcal{C}_n}\left(\max_{x^n, y^n} \widetilde{g}(x^n, y^n | \mathcal{C}_n) \ge 1 + \delta\right)$$
(173)

$$\leq \left|\mathcal{X}\right|^{n} \left|\mathcal{Y}\right|^{n} \max_{x^{n}, y^{n}} \mathbb{P}_{\mathcal{C}_{n}}\left(\widetilde{g}(x^{n}, y^{n} | \mathcal{C}_{n}) \geq 1 + \delta\right), \quad (174)$$

where (174) follows from the union bound. Obviously,  $|\mathcal{X}|^n |\mathcal{Y}|^n$  is only exponentially growing. Therefore, if the probability vanishes doubly exponentially fast, then  $\max_{x^n,y^n} \tilde{g}(x^n,y^n|\mathcal{C}_n) < 1 + \delta$  with probability of failure decaying to zero doubly exponentially fast as  $n \to \infty$ . To this end, we use the Bernstein inequality to bound the probability.

<sup>14</sup>For brevity, we denote  $P_{X^nY^n|\mathcal{C}_n}$  as  $P_{X^nY^n}$ .

Observe that  $g(x^n, y^n | W^n(m)), m \in \mathcal{M}_n$  are i.i.d. random variables with mean

$$\mu_{n} := \mathbb{E}_{W^{n}} \left[ g(x^{n}, y^{n} | W^{n}) \right]$$

$$= \frac{1}{Q_{XY}^{n}(x^{n}, y^{n})} \sum_{w^{n}} \frac{Q_{W}^{n}(w^{n}) 1\left\{ w^{n} \in \mathcal{T}_{\frac{\epsilon}{2}}^{(n)}(Q_{W}) \right\}}{Q_{W}^{n}\left(\mathcal{T}_{\frac{\epsilon}{2}}^{(n)}(Q_{W})\right)}$$

$$\times \frac{Q_{X|W}^{n}(x^{n} | w^{n}) 1\left\{ x^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WX} | w^{n}) \right\}}{Q_{X|W}^{n}\left(\mathcal{T}_{\epsilon}^{(n)}(Q_{WX} | w^{n}) | w^{n}\right)}$$

$$\times \frac{Q_{Y|W}^{n}(y^{n} | w^{n}) 1\left\{ y^{n} \in \mathcal{T}_{\epsilon}^{(n)}(Q_{WY} | w^{n}) \right\}}{Q_{Y|W}^{n}\left(\mathcal{T}_{\epsilon}^{(n)}(Q_{WY} | w^{n}) | w^{n}\right)}$$

$$(176)$$

$$\leq \frac{1}{(1-\delta_{0,n})(1-\delta_{1,n})(1-\delta_{2,n})}$$
(177)

$$\rightarrow 1$$
 exponentially fast as  $n \rightarrow \infty$ , (178)

and variance

$$\operatorname{Var}_{W^{n}}\left[g(x^{n}, y^{n}|W^{n})\right] \leq \mathbb{E}_{W^{n}}\left[g(x^{n}, y^{n}|W^{n})^{2}\right] \quad (179)$$
$$\leq \beta_{n}\mu_{n}. \quad (180)$$

Here (178) follows since  $\delta_{0,n} := 1 - Q_W^n\left(\mathcal{T}_{\frac{\epsilon}{2}}^{(n)}(Q_W)\right)$  converges to zero exponentially fast as  $n \to \infty$ . Then we bound the probability in (174) as follows:

$$\mathbb{P}_{\mathcal{C}_{n}}\left(\widetilde{g}(x^{n}, y^{n} | \mathcal{C}_{n}) \geq 1 + \delta\right)$$

$$= \mathbb{P}_{\mathcal{C}_{n}}\left(\sum_{m \in \mathcal{M}_{n}} g(x^{n}, y^{n} | W^{n}(m)) - \mu_{n} \mathsf{M}\right)$$

$$\geq (1 + \delta - \mu_{n}) \mathsf{M}\right)$$
(181)

$$\leq \exp\left(-\frac{\frac{1}{2}\left(1+\delta-\mu_{n}\right)^{2}\mathsf{M}^{2}}{\mathsf{M}\beta_{n}\mu_{n}+\frac{1}{3}\left(1+\delta-\mu_{n}\right)\mathsf{M}\beta_{n}}\right)$$
(182)

$$= \exp\left(-\frac{3\left(1+\delta-\mu_n\right)^2 \mathsf{M}}{2\left(1+\delta+2\mu_n\right)\beta_n}\right),\tag{183}$$

where (182) follows from Bernstein's inequality, stated here for the readers' convenience.

**Lemma 8.** [45] Let  $X_1, \ldots, X_n$  be independent zero-mean random variables such that  $|X_i| \leq M$  almost surely, for all *i*. Then, for any t > 0,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_i > t\right) \le \exp\left(-\frac{\frac{1}{2}t^2}{\sum_{i=1}^{n} \mathbb{E}\left[X_i^2\right] + \frac{1}{3}Mt}\right). \quad (184)$$

Observe that

$$\frac{\mathsf{M}}{\beta_n} = (1 - \delta_{1,n}) (1 - \delta_{2,n}) e^{n(R - \mathcal{I}_{\epsilon}(Q))}.$$
 (185)

Denote  $\alpha_0$  as the exponent of  $\frac{1}{(1-\delta_{0,n})(1-\delta_{1,n})(1-\delta_{2,n})} - 1$ . By [3, Lemma 4],

$$\alpha_0 \ge \min\left\{\frac{1}{3}\epsilon^2 Q_W^{(\min)}, \frac{1}{3}\left(\frac{\epsilon}{2+\epsilon}\right)^2 \min\left\{Q_{X|W}^{(\min)}, Q_{Y|W}^{(\min)}\right\}\right\},\tag{186}$$

where  $Q_W^{(\min)} := \min_{w:Q_W(w)>0} Q_W(w), \quad Q_{X|W}^{(\min)} := \min_{(x,w):Q_{X|W}(x|w)>0} Q_{X|W}(x|w)$ , and similarly for  $Q_{Y|W}^{(\min)}$ . By (177),  $\mu_n - 1 \le e^{-n\frac{\alpha_0}{2}}$  for all sufficiently large n.

Set  $\delta = e^{-n\alpha_1}$  with  $\alpha_1 := \min\left\{\frac{\alpha_0}{4}, \frac{\epsilon}{4}\right\} > 0$ , then

$$\begin{split} \liminf_{n \to \infty} \frac{1}{n} \log \frac{3\left(1 + \delta - \mu_n\right)^2 \mathsf{M}}{2\left(1 + \delta + 2\mu_n\right)\beta_n} \\ \geq R - \mathcal{I}_{\epsilon}\left(Q\right) \\ + \liminf_{n \to \infty} \frac{1}{n} \log \frac{3\left(e^{-n\alpha_1} - e^{-n\frac{\alpha_0}{2}}\right)^2}{2\left(1 + e^{-n\alpha_1} + 2\left(1 + e^{-n\frac{\alpha_0}{2}}\right)\right)} \\ = R - \mathcal{I}_{\epsilon}\left(Q\right) - 2\alpha_1, \end{split}$$
(188)

where (188) follows since  $\alpha_1 < \frac{\alpha_0}{2}$ . Hence the exponent of  $\frac{3(1+\delta-\mu_n)^2M}{2(1+\delta+2\mu_n)\beta_n}$  is lower bounded by

$$R - \mathcal{I}_{\epsilon}(Q) - 2\alpha_1 \ge \frac{\epsilon}{2}, \tag{189}$$

where (189) holds due to (165) and the choice of  $\alpha_1$ . Hence (183) converges to zero doubly exponentially fast in *n*. Combined this with (174) yields

$$\mathbb{P}_{\mathcal{C}_n}\left(D_{\infty}(P_{X^nY^n|\mathcal{C}_n}\|\pi_{XY}^n) \ge e^{-n\alpha_1}\right) \to 0$$
(190)

doubly exponentially fast as  $n \to \infty$ .

# C. Proof of $T_{\text{Exact}}(\pi_{XY}) \geq \limsup_{n \to \infty} \frac{1}{n} \Gamma(\pi_{XY}^n)$

We prove the converse result from the perspective of exact common information, i.e.,

$$T_{\text{Exact}}(\pi_{XY}) \ge \limsup_{n \to \infty} \frac{1}{n} \Gamma(\pi_{XY}^n).$$
(191)

Similar to the idea used in Appendix A-A, we first independently replicate a k-length optimal exact common information code  $(P_{W_k}, P_{X^k|W_k}, P_{Y^k|W_k})$  n times. Then the resulting superblock code is also an exact common information code, i.e.,  $\sum_{w^n} P_{W_k}^n(w^n) P_{X^k|W_k}^n(\cdot|w^n) P_{Y^k|W_k}^n(\cdot|w^n) = \pi_{XY}^{kn}$ . Observe that  $W_k^n = (W_{k,1}, W_{k,2}, ..., W_{k,n})$  is an n-length i.i.d. random sequence with each  $W_{k,i} \sim P_{W_k}$ . Hence we have for  $\epsilon > 0$ ,

$$\mathbb{P}\left(W_{k}^{n} \in \mathcal{A}_{\epsilon}^{(n)}\left(P_{W_{k}}\right)\right) \to 1$$
(192)

as  $n \to \infty$  for fixed k. Furthermore,  $|\mathcal{A}_{\epsilon}^{(n)}| \leq e^{n(H(W_k)+\epsilon)}$ .

Consider

$$D_{\infty}(P_{X^{kn}Y^{kn}} \| \pi_{XY}^{kn}) = \log \left( \max_{x^{kn}, y^{kn}} \frac{\sum_{w^n} P_{W_k}^n(w^n) P_{X^k|W_k}^n(x^{kn}|w^n) P_{Y^k|W_k}^n(y^{kn}|w^n)}{\pi_{XY}^{kn}(x^{kn}, y^{kn})} \right)$$
(193)

$$\geq \log \left( \max_{x^{kn}, y^{kn}} \max_{w^n \in \mathcal{A}_{\epsilon}^{(n)}} \frac{P_{W_k}^n(w^n) P_{X^k|W_k}^n(x^{kn}|w^n) P_{Y^k|W_k}^n(y^{kn}|w^n)}{\pi_{XY}^{kn}(x^{kn}, y^{kn})} \right)$$
(194)

$$\geq \log \left( \max_{x^{kn}, y^{kn}} \max_{w^n \in \mathcal{A}_{\epsilon}^{(n)}} \frac{e^{-n(H(W_k)+\epsilon)} P_{X^k|W_k}^n(x^{kn}|w^n) P_{Y^k|W_k}^n(y^{kn}|w^n)}{\pi_{XY}^{kn}(x^{kn}, y^{kn})} \right)$$
(195)

$$= -n \left( H(W_{k}) + \epsilon \right) + \log \left( \max_{x^{kn}, y^{kn}} \max_{w^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{P_{X^{k}|W_{k}}^{n}(x^{kn}|w^{n})P_{Y^{k}|W_{k}}^{n}(y^{kn}|w^{n})}{\pi_{XY}^{kn}(x^{kn}, y^{kn})} \right).$$
(196)

Since for the exact common information superblock code,  $D_{\infty}(P_{X^{kn}Y^{kn}} || \pi_{XY}^{kn}) = 0$ , we have

$$\frac{\frac{1}{k}(H(W_{k}) + \epsilon)}{\sum_{n \to \infty} \sum_{n \to \infty} \frac{1}{kn} \log} \left( \max_{x^{kn}, y^{kn}} \max_{w^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{P_{X^{k}|W_{k}}^{n}(x^{kn}|w^{n})P_{Y^{k}|W_{k}}^{n}(y^{kn}|w^{n})}{\pi_{XY}^{kn}(x^{kn}, y^{kn})} \right)$$
(197)

$$= \limsup_{n \to \infty} \frac{1}{kn} \max_{w^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \sum_{i=1}^{n} \max_{x^{k}, y^{k}} \log \frac{P_{X^{k}|W_{k}}(x^{k}|w_{i})P_{Y^{k}|W_{k}}(y^{k}|w_{i})}{\pi_{XY}^{k}(x^{k}, y^{k})}.$$
 (198)

Continuing (198), we obtain

$$\max_{w^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \sum_{i=1}^{n} \max_{x^{k}, y^{k}} \log \frac{P_{X^{k}|W_{k}}(x^{k}|w_{i})P_{Y^{k}|W_{k}}(y^{k}|w_{i})}{\pi_{XY}^{k}(x^{k}, y^{k})} \\
\geq \sum_{w^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{P_{W_{k}}^{n}(w^{n})}{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})} \sum_{i=1}^{n} \max_{\substack{Q_{X^{k}Y^{k}|W_{k}} \in \\ C(P_{X^{k}|W_{k}}, P_{Y^{k}|W_{k}})}} \\
\sum_{x^{k}, y^{k}} Q_{X^{k}Y^{k}|W_{k}}(x^{k}, y^{k}|w_{i}) \\
\times \log \frac{P_{X^{k}|W_{k}}(x^{k}|w_{i})P_{Y^{k}|W_{k}}(y^{k}|w_{i})}{\pi_{XY}^{k}(x^{k}, y^{k})} \qquad (199) \\
= \sum_{w^{n} \in \mathcal{A}_{\epsilon}^{(n)}} \frac{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})}{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})} \sum_{i=1}^{n} g(w_{i}, P_{X^{k}|W_{k}}, P_{Y^{k}|W_{k}}) \qquad (200)$$

$$= \sum_{w^{n}} \frac{P_{W_{k}}^{n}(w^{n})}{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})} \sum_{i=1}^{n} g\left(w_{i}, P_{X^{k}|W_{k}}, P_{Y^{k}|W_{k}}\right) \\ - \sum_{w^{n} \notin \mathcal{A}_{\epsilon}^{(n)}} \frac{P_{W_{k}}^{n}(w^{n})}{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})} \sum_{i=1}^{n} g\left(w_{i}, P_{X^{k}|W_{k}}, P_{Y^{k}|W_{k}}\right)$$
(201)

$$\geq \frac{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})}{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})} \mathbb{E}_{W \sim P_{W_{k}}}g\left(W, P_{X^{k}|W_{k}}, P_{Y^{k}|W_{k}}\right) - \frac{1 - P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})}{P_{W_{k}}^{n}(\mathcal{A}_{\epsilon}^{(n)})} nk \log \min_{x,y:\pi_{XY}(x,y)>0} \pi_{XY}\left(x,y\right),$$
(202)

where (199) follows since the maximum is no smaller than the average; in (200),

$$g\left(w, P_{X^{k}|W_{k}}, P_{Y^{k}|W_{k}}\right)$$
  
:=  $-H(X^{k}|W_{k} = w) - H(Y^{k}|W_{k} = w)$  (203)

$$+ \mathcal{H}(P_{X^{k}|W_{k}=w}, P_{Y^{k}|W_{k}=w} \| \pi_{XY}^{k});$$
(204)

and (202) follows since

$$g\left(w, P_{X^{k}|W_{k}}, P_{Y^{k}|W_{k}}\right) \leq \mathcal{H}(P_{X^{k}|W_{k}=w}, P_{Y^{k}|W_{k}=w} \| \pi_{XY}^{k})$$
(205)

$$\leq -k \log \min_{x,y:\pi_{XY}(x,y)>0} \pi_{XY}(x,y) \,. \tag{206}$$

Since  $P_{W_k}^n(\mathcal{A}_{\epsilon}^{(n)}) \to 1$ , combining this fact with (198) and (200), we have

$$\frac{1}{k} (H(W_k) + \epsilon)$$

$$\geq \frac{1}{k} \left( -H(X^k | W_k) - H(Y^k | W_k) + \sum_{w} P_{W_k}(w) \mathcal{H}(P_{X^k | W_k = w}, P_{Y^k | W_k = w} \| \pi_{XY}^k) \right)$$
(207)

$$\geq \frac{1}{k} \left( \inf_{\substack{P_{W_{k}}P_{X^{k}|W_{k}}P_{Y^{k}|W_{k}}:\\P_{X^{k}Y^{k}}=\pi_{XY}^{k}}} -H(X^{k}|W_{k}) - H(Y^{k}|W_{k}) + \sum_{w} P_{W_{k}}(w) \mathcal{H}(P_{X^{k}|W_{k}=w}, P_{Y^{k}|W_{k}=w} \| \pi_{XY}^{k}) \right)$$

$$(208)$$

$$=\frac{1}{k}\Gamma(\pi_{XY}^k).$$
(209)

Furthermore, since  $\frac{1}{k}H(W_k) \to R$  as  $k \to \infty$ , we have

$$R \ge \limsup_{k \to \infty} \frac{1}{k} \Gamma(\pi_{XY}^k).$$
(210)

#### APPENDIX B Proof of Theorem 2

The inequality  $\widetilde{T}_{\infty}(\pi_{XY}) \geq C_{\text{Wyner}}(\pi_{XY})$  follows since  $\widetilde{T}_{\infty}(\pi_{XY}) \geq \widetilde{T}_{1}(\pi_{XY}) \geq C_{\text{Wyner}}(\pi_{XY})$ , where the last inequality is the converse result for Wyner's common information [2]. On the other hand, the upper bound  $\Gamma^{\text{UB}}(\pi_{XY})$  (i.e.,  $\Gamma(\pi_{XY})$ ) has been proved in Appendix A-B. Hence we only

need to prove the lower bound, i.e.,  $\widetilde{T}_{\infty}(\pi_{XY}) \geq \Gamma^{\text{LB}}(\pi_{XY})$ . The proof for this inequality is divided into two parts: singleletterization and simplifying constraints.

#### A. Single-letterization

Observe by Remark 2, since  $\widetilde{T}_{\infty}(\pi_{XY}) \geq \lim_{\epsilon \downarrow 0} \lim_{n \to \infty} \frac{1}{n} \Gamma_{\epsilon}(\pi_{XY}^{n})$ , in order to lower bound  $\widetilde{T}_{\infty}(\pi_{XY})$ , it suffices to lower bound  $\frac{1}{n} \Gamma_{\epsilon}(\pi_{XY}^{n})$ . According to the definition of  $\Gamma_{\epsilon}(\pi_{XY}^{n})$  in (37), we have (211) (given on page 22). Denote  $J \sim P_{J} := \text{Unif}[1:n]$  as a time index which is independent of  $(W, X^{n}, Y^{n}) \sim P_{W} P_{X^{n} \mid W} P_{Y^{n} \mid W}$ . Then

$$-\frac{1}{n}\sum_{i=1}^{n}H(X_{i}|X^{i-1}W) - \frac{1}{n}\sum_{i=1}^{n}H(Y_{i}|Y^{i-1}W)$$
  
=  $-H(X_{J}|X^{J-1}WJ) - H(Y_{J}|Y^{J-1}WJ).$  (212)

Next we single-letterize the last term in (211). On one hand,

$$\sum_{x^{i-1}, y^{i-1}} Q(x^{i-1}, y^{i-1} | w) \\ \times \sum_{x_i, y_i} Q(x_i, y_i | x^{i-1}, y^{i-1}, w) \log \frac{1}{\pi(x_i, y_i)} \\ \geq \min_{\substack{\widetilde{Q}_{X^{i-1}Y^{i-1} | W} \in \\ C(P_{X^{i-1} | W}, P_{Y^{i-1} | W})}} \sum_{x^{i-1}, y^{i-1}} \widetilde{Q}(x^{i-1}, y^{i-1} | w) \\ \times \sum_{x_i, y_i} Q(x_i, y_i | x^{i-1}, y^{i-1}, w) \log \frac{1}{\pi(x_i, y_i)}.$$
(213)

On the other hand, in order to get a further lower bound on (213), we need the following "chain rule" for coupling sets.

**Lemma 9** ("Chain Rule" for Coupling Sets). For a pair of conditional distributions  $(P_{X^n|W}, P_{Y^n|W})$ , we have

$$\prod_{i=1}^{n} C(P_{X_{i}|X^{i-1}W}, P_{Y_{i}|Y^{i-1}W}) \subseteq C(P_{X^{n}|W}, P_{Y^{n}|W}),$$
(214)

where

$$C(P_{X_{i}|X^{i-1}W}, P_{Y_{i}|Y^{i-1}W})$$
  
:=  $\left\{ Q_{X_{i}Y_{i}|X^{i-1}Y^{i-1}W} : Q_{X_{i}|X^{i-1}Y^{i-1}W} = P_{X_{i}|X^{i-1}W}, Q_{Y_{i}|X^{i-1}Y^{i-1}W} = P_{Y_{i}|Y^{i-1}W} \right\}, i \in [1:n]$  (215)

and

$$\prod_{i=1}^{n} C(P_{X_{i}|X^{i-1}W}, P_{Y_{i}|Y^{i-1}W})$$
  
$$:= \left\{ \prod_{i=1}^{n} Q_{X_{i}Y_{i}|X^{i-1}Y^{i-1}W} : Q_{X_{i}Y_{i}|X^{i-1}Y^{i-1}W} \in C(P_{X_{i}|X^{i-1}W}, P_{Y_{i}|Y^{i-1}W}), i \in [1:n] \right\}.$$
(216)

 $\begin{array}{rll} \textit{Proof:} & \mathrm{If} & \left\{Q_{X_iY_i|X^{i-1}Y^{i-1}W}\right\}_{i\in[1:n]} \text{ is a set} \\ \mathrm{of} & \mathrm{distributions} & \mathrm{such} & \mathrm{that} & Q_{X_iY_i|X^{i-1}Y^{i-1}W} & \in \end{array}$ 

$$\frac{1}{n}\Gamma_{\epsilon}(\pi_{XY}^{n}) \geq \frac{1}{n} \inf_{\substack{P_{W}P_{X^{n}|W}P_{Y^{n}|W}: \\ \frac{1}{n}D_{\infty}(P_{X^{n}Y^{n}}||\pi_{XY}^{n}) \leq \epsilon \ C(P_{X^{n}|W},P_{Y^{n}|W})}} -\frac{1}{n}\sum_{i=1}^{n}H(X_{i}|X^{i-1}W) - \frac{1}{n}\sum_{i=1}^{n}H(Y_{i}|Y^{i-1}W) - \sum_{i=1}^{n}H(Y_{i}|Y^{i-1}W) - \sum_{i=1}^{n}P(w)\frac{1}{n}\sum_{i=1}^{n}\int_{i=1}^{n}\left(\sum_{x^{i-1},y^{i-1}}Q(x^{i-1},y^{i-1}|w)\sum_{x_{i},y_{i}}Q(x_{i},y_{i}|x^{i-1},y^{i-1},w)\log\pi(x_{i},y_{i})\right). \quad (211)$$

 $C(P_{X_i|X^{i-1}W},P_{Y_i|Y^{i-1}W})$  for all  $i\in[1:n],$  then we have that for any  $(w,x^n),$ 

$$\sum_{y^{n}} \prod_{i=1}^{n} Q\left(x_{i}, y_{i} | x^{i-1}, y^{i-1}, w\right)$$

$$= \sum_{y^{n-1}} \prod_{i=1}^{n-1} Q\left(x_{i}, y_{i} | x^{i-1}, y^{i-1}, w\right)$$

$$\times \sum_{y_{n}} Q\left(x_{n}, y_{n} | x^{n-1}, y^{n-1}, w\right)$$
(217)

$$= \sum_{y^{n-1}} \prod_{i=1}^{n-1} Q\left(x_i, y_i | x^{i-1}, y^{i-1}, w\right) Q\left(x_n | x^{n-1}, y^{n-1}, w\right)$$
(218)

$$= P\left(x_n | x^{n-1}, w\right) \sum_{y^{n-1}} \prod_{i=1}^{n-1} Q\left(x_i, y_i | x^{i-1}, y^{i-1}, w\right)$$
(219)

$$= P\left(x_{n}|x^{n-1},w\right) P\left(x_{n-1}|x^{n-2},w\right)$$
$$\times \sum_{y^{n-2}} \prod_{i=1}^{n-2} Q\left(x_{i},y_{i}|x^{i-1},y^{i-1},w\right)$$
(220)

$$=\prod_{i=1}^{n} P\left(x_{i}|x^{i-1},w\right)$$
(221)

$$=P\left(x^{n}|w\right),\tag{222}$$

where (219) follows since  $Q_{X_iY_i|X^{i-1}Y^{i-1}W} \in C(P_{X_i|X^{i-1}W}, P_{Y_i|Y^{i-1}W}).$ 

Hence  $\prod_{i=1}^{n} Q_{X_i Y_i | X^{i-1} Y^{i-1} W}$  has marginal conditional distributions  $P_{X^n | W}$  and  $P_{Y^n | W}$ , i.e.,  $\prod_{i=1}^{n} Q_{X_i Y_i | X^{i-1} Y^{i-1} W} \in C(P_{X^n | W}, P_{Y^n | W})$ . Since for any  $i \in [1:n]$ ,  $Q_{X_i Y_i | X^{i-1} Y^{i-1} W}$  is an arbitrary distribution in  $C(P_{X_i | X^{i-1} W}, P_{Y_i | Y^{i-1} W})$ , we have that  $\prod_{i=1}^{n} C(P_{X_i | X^{i-1} W}, P_{Y_i | Y^{i-1} W}) \subseteq C(P_{X^n | W}, P_{Y^n | W})$ . By Lemma 9, we have that for any function  $f: \mathcal{P}(\mathcal{X}^n \times \mathcal{Y}^n) \to \mathbb{R}$ ,

$$\max_{\substack{Q_{X^{n}Y^{n}|W} \in \\ C(P_{X^{n}|W}, P_{Y^{n}|W})}} f\left(Q_{X^{n}Y^{n}|W}\right) \\
\geq \max_{\substack{Q_{X^{n}Y^{n}|W} \in \\ \prod_{i=1}^{n} C(P_{X_{i}|X^{i-1}W}, P_{Y_{i}|Y^{i-1}W})}} f\left(\prod_{i=1}^{n} Q_{X_{i}Y_{i}|X^{i-1}Y^{i-1}W}\right).$$
(223)

Therefore, substituting (213) into the last term in (211) and utilizing (223), we obtain (227)-(229) (given on page 23), where the swapping of min and max in (228) follows since

on one hand, maximin is no larger than minimax, and on the other hand,

$$(227) \geq \sum_{w} P(w) \frac{1}{n} \sum_{i=1}^{n} \min_{\substack{\tilde{Q}_{X^{i-1}Y^{i-1}|W} \in \\ C(P_{X^{i-1}|W}, P_{Y^{i-1}|W})}} \sum_{\substack{x^{i-1}, y^{i-1}} \tilde{Q}(x^{i-1}, y^{i-1}|w) \\ \times \sum_{x_{i}, y_{i}} Q^{*}(x_{i}, y_{i}|x^{i-1}, y^{i-1}, w) \log \frac{1}{\pi(x_{i}, y_{i})}$$

$$(224)$$

$$(225)$$

$$= (228)$$
 (225)

with

$$Q_{X_{i}Y_{i}|X^{i-1}Y^{i-1}W}^{*} := \arg \max_{\substack{Q_{X_{i}Y_{i}|X^{i-1}Y^{i-1}W \in \\ C(P_{X_{i}|X^{i-1}W}, P_{Y_{i}|Y^{i-1}W})}} \sum_{x_{i}, y_{i}} Q(x_{i}, y_{i}|x^{i-1}, y^{i-1}, w) \times \log \frac{1}{\pi(x_{i}, y_{i})}.$$
(226)

For brevity, we set

$$W \leftarrow WJ, U \leftarrow X^{J-1}, V \leftarrow Y^{J-1}, X \leftarrow X_J, Y \leftarrow Y_J.$$
(230)

Then  $\frac{1}{n}D_{\infty}(P_{X^nY^n} \| \pi_{XY}^n) \leq \epsilon$  implies that  $D(P_{XY} \| \pi_{XY}) \leq \epsilon$ . Since  $\pi_{XY}$  has finite support,  $D(P_{XY} \| \pi_{XY}) \rightarrow 0$  if and only if  $D_{\infty}(P_{XY} \| \pi_{XY}) \rightarrow 0$ . Therefore, substituting (212) and (229) into (211) and utilizing the identification of the random variables in (230), we obtain (231) (given on page 23). For  $\tilde{Q}_{UV|W} \in C(P_{U|W}, P_{V|W})$ , define a joint distribution induced by  $\tilde{Q}_{UV|W}$  as

$$\widehat{Q}_{(U,V',W),(U',V,W')}(u,v',w,u',v,w') 
:= P_W(w)\widetilde{Q}_{UV|W}(u,v|w)1\{w'=w\} 
\times P_{V|W}(v'|w)P_{U|W}(u'|w').$$
(232)

Then this joint distribution satisfies the following marginal constraints:

$$\widehat{Q}_{UVW}(u, v, w) = P_W(w)\widetilde{Q}_{UV|W}(u, v|w)$$
(233)

$$\widehat{Q}_{UV'W}(u, v', w) = P_{UVW}(u, v', w)$$
(234)

$$\widehat{Q}_{U'VW'}(u', v, w') = P_{UVW}(u', v, w').$$
(235)

$$\begin{split} & \max_{\substack{Q_{X^n Y^n | W} \in \\ C(P_{X^n | W}, P_{Y^n | W}) \\ W}} \sum_{w} P(w) \frac{1}{n} \sum_{i=1}^{n} \sum_{x^{i-1}, y^{i-1}} Q(x^{i-1}, y^{i-1} | w) \sum_{x_i, y_i} Q(x_i, y_i | x^{i-1}, y^{i-1}, w) \log \frac{1}{\pi(x_i, y_i)} \\ & \geq \sum_{w} P(w) \frac{1}{n} \sum_{i=1}^{n} \max_{\substack{Q_{X_i Y_i | X^{i-1} W \in \\ C(P_{X_i | X^{i-1} W}, P_{Y_i | Y^{i-1} W}) \\ C(P_{X_i | Y^{i-1} W}) \sum_{x_i, y_i} Q(x_i, y_i | x^{i-1}, w^{i-1}, w) \log \frac{1}{\pi(x_i, y_i)}} \\ & \sum_{x^{i-1}, y^{i-1}} \widetilde{Q}(x^{i-1}, y^{i-1} | w) \sum_{x_i, y_i} Q(x_i, y_i | x^{i-1}, y^{i-1}, w) \log \frac{1}{\pi(x_i, y_i)} \\ & = \sum_{w} P(w) \frac{1}{n} \sum_{i=1}^{n} \min_{\substack{Q_{X^{i-1} Y^{i-1} | W \in \\ C(P_{X^{i-1} | W}, P_{Y^{i-1} | W)}}} \sum_{x_i, y_i} Q(x_i, y_i | x^{i-1}, y^{i-1}, w) \log \frac{1}{\pi(x_i, y_i)} \\ & \times \max_{\substack{Q_{X_i Y_i | X^{i-1} W \in \\ C(P_{X^{i-1} | W}, P_{Y^{i-1} | W)}}} \sum_{x_i, y_i} Q(x_i, y_i | x^{i-1}, y^{i-1}, w) \log \frac{1}{\pi(x_i, y_i)} \\ & = \sum_{w} P(w) \sum_{j=1}^{n} P_J(j) \max_{\substack{Q_{X^{J-1} Y^{J-1} | W \in \\ C(P_{X^{J-1} | W, P_{Y^{J-1} | W, J}) \\ C(P_{X^{J-1} | W, P_{Y^{J-1} | W, J})} \sum_{x_j, y_j} Q(x_j, y_j | x^{j-1}, y^{j-1}, w, j) \log \frac{1}{\pi(x_j, y_j)}, \end{split}$$
(228)  

$$& = \sum_{w} P(w) \sum_{j=1}^{n} P_J(j) \max_{\substack{Q_{X^{J-1} Y^{J-1} | W \in \\ C(P_{X^{J-1} | W, J, P_{Y^{J-1} | W, J})}} \sum_{x_j, y_j} Q(x_j, y_j | x^{j-1}, y^{j-1}, w, j) \log \frac{1}{\pi(x_j, y_j)}, \end{cases}$$
(229)

$$\widetilde{T}_{\infty}(\pi_{XY}) \geq \lim_{\epsilon \downarrow 0} \inf_{\substack{P_{W}P_{U|W}P_{Y|W}P_{X|U}P_{Y|V}:\\ D_{\infty}(P_{XY}||\pi_{XY}) \leq \epsilon}} -H(X|UW) - H(Y|VW) + \sum_{w} P(w) \inf_{\substack{\widetilde{Q}_{UV|W} \in \\ C(P_{U|W},P_{V|W})}} \sum_{u,v} \widetilde{Q}(u,v|w) \max_{\substack{Q_{XY}|UVW \in \\ C(P_{X|UW},P_{Y|VW})}} \sum_{x,y} Q(x,y|u,v,w) \log \frac{1}{\pi(x,y)}.$$
(231)

Utilizing this induced distribution, its properties in (233)-(235), and the lower bound in (231), we obtain

$$\widetilde{T}_{1+s}(\pi_{XY}) \geq \lim_{\epsilon \downarrow 0} \inf_{\substack{P_{W}P_{U|W}P_{Y|W}P_{X|U}P_{Y|V}:\\D_{\infty}(P_{XY}||\pi_{XY}) \leq \epsilon}} - (H(X|UW) + H(Y|VW)) + \frac{1}{Q_{(U,V',W),(U',V,W')} \leq \epsilon} \sum_{\substack{u,u',v,v',w,w'\\C(P_{UVW},P_{UVW})}} \widehat{Q}(u,v',w,u',v,w') + \frac{1}{Q_{XY} \leq \epsilon} \sum_{\substack{Q_{XY} \in C(P_{X|UW=u,w},P_{Y}|VW=v,w')}} \sum_{\substack{x,y}} Q(x,y) \log \frac{1}{\pi(x,y)}.$$
(236)

Observe that  $P_{X|UW=u,w} = P_{X|(U,V,W)=(u,v',w)}$  and  $P_{Y|VW=v,w'} = P_{Y|(U,V,W)=(u',v,w')}$  (since  $X \to UW \to V$  and  $Y \to VW \to U$  form Markov chains under P). Hence the coupling set  $C(P_{X|UW=u,w}, P_{Y|VW=v,w'})$  in the last term in (236) can be replaced by  $C(P_{X|(U,V,W)=(u,v',w)}, P_{Y|(U,V,W)=(u',v,w')})$ .

Substituting  $W \leftarrow (U, V, W)$ , we can simplify (236) as follows:

 $\geq \lim_{\epsilon \downarrow 0} \inf_{\substack{P_{W}P_{X|W}P_{Y|W}:\\D_{\infty}(P_{XY} \| \pi_{XY}) \leq \epsilon}} - (H(X|W) + H(Y|W)) + \lim_{\substack{Q_{WW'} \in\\C(P_{W}, P_{W})}} \sum_{w,w'} Q(w,w') \mathcal{H}(P_{X|W=w}, P_{Y|W=w'} \| \pi_{XY}).$ (237)

#### B. Simplifying Constraints

Next we prove that the constraint  $D_{\infty}(P_{XY} || \pi_{XY}) \leq \epsilon$  in (237) can be replaced with  $P_{XY} = \pi_{XY}$ . For  $D_{\infty}(P_{XY} || \pi_{XY}) \leq \epsilon$ , using the splitting technique, we can write

$$\pi_{XY}(x,y) = e^{-\epsilon} P_{XY}(x,y) + (1 - e^{-\epsilon}) \hat{P}_{XY}(x,y)$$
(238)

where

$$\widehat{P}_{XY}(x,y) := \frac{e^{\epsilon} \pi_{XY}(x,y) - P_{XY}(x,y)}{e^{\epsilon} - 1}.$$
(239)

Define

$$\widetilde{P}_{XYWU}(x, y, w, u) = \begin{cases} e^{-\epsilon} P_W P_{X|W} P_{Y|W} & \text{if } u = 1\\ (1 - e^{-\epsilon}) \widehat{P}_{XY}(x, y) \mathbf{1} \{ w = (x, y) \} & \text{if } u = 0 \end{cases}. (240)$$

consider (241)-(243) (given on page 25), Then (241) follows since  $\overline{P}_U(u)\overline{P}_U(u')Q_{WW'}(w,w')$ where with  $\in$  $C(P_{W|U=u}, P_{W|U=u'})$  forms a  $Q_{WW'}$ coupling of  $(\tilde{P}_{WU}, \tilde{P}_{WU})$ , and (242) follows since  $\mathcal{H}(Q_X, Q_Y \| \pi_{XY}) \leq \max_{(x,y) \in \operatorname{supp}(\pi_{XY})} \log \frac{1}{\pi(x,y)}$  for any  $(Q_X, Q_Y).$ 

Hence substituting (243) into (237), we obtain (244)-(247) (given on page 25).

#### APPENDIX C **PROOF OF THEOREM 3**

Upper Bound: Set  $X = W \oplus A$  and  $Y = W \oplus B$  with  $W \sim \text{Bern}(\frac{1}{2}), A \sim \text{Bern}(a), \text{ and } B \sim \text{Bern}(a)$  mutually independent, where  $a := \frac{1-\sqrt{1-2p}}{2} \in (0, \frac{1}{2}).$ 

$$\mathcal{H}(P_{X|W=w}, P_{Y|W=w} \| \pi_{XY}) = \max_{Q_{XY} \in C(P_{X|W=w}, P_{Y|W=w})} \sum_{x,y} Q_{XY}(x, y) \log \frac{1}{\pi(x, y)}$$
(248)

$$= \log \frac{1}{\alpha_0} + 2\min\{a, \overline{a}\} \log \frac{\alpha_0}{\beta_0}$$
(249)

$$= \log \frac{1}{\alpha_0} + 2a \log \frac{\alpha_0}{\beta_0} \tag{250}$$

Hence we have

$$\Gamma^{\rm UB}(\pi_{XY}) \le -H_2(a) - H_2(a) + \log \frac{1}{\alpha_0} + 2a \log \frac{\alpha_0}{\beta_0}$$
(251)

$$= -2H_2(a) + \log\frac{1}{\alpha_0} + 2a\log\frac{\alpha_0}{\beta_0}.$$
 (252)

Substituting  $\alpha_0, \beta_0$  into (252), we get the right hand side of (55).

Lower Bound: We adopt similar techniques as ones used by Wyner [2]. Denote

$$\alpha(w) := \mathbb{P}\left(X = 0 | W = w\right) \tag{253}$$

$$\beta(w) := \mathbb{P}\left(Y = 0 | W = w\right). \tag{254}$$

Hence  $P_{XY} = \pi_{XY}$  implies

$$\mathbb{E}\alpha(W) = \mathbb{P}\left(X = 0\right) = \frac{1}{2}$$
(255)

$$\mathbb{E}\beta(W) = \mathbb{P}\left(Y=0\right) = \frac{1}{2}$$
(256)  
$$\alpha(W)\beta(W) = \mathbb{P}\left(X=0, Y=0\right) = \alpha_0.$$
(257)

$$\mathbb{E}\alpha(W)\beta(W) = \mathbb{P}\left(X = 0, Y = 0\right) = \alpha_0.$$
 (257)

Observe that

$$\mathcal{H}(P_{X|W=w}, P_{Y|W=w'} \| \pi_{XY}) = \max_{Q_{XY} \in C(P_{X|W=w}, P_{Y|W=w'})} \sum_{x,y} Q_{XY}(x, y) \log \frac{1}{\pi(x, y)}$$
(258)

$$= \log \frac{1}{\alpha_0} + \left(\min\{\alpha(w), \overline{\beta(w')}\} + \min\{\overline{\alpha(w)}, \beta(w')\right) \log \frac{\alpha_0}{\beta_0}$$
(259)
$$= \log \frac{1}{\alpha_0} + \min\{\alpha(w) + \beta(w'), \overline{\alpha(w)} + \overline{\beta(w')}\} \log \frac{\alpha_0}{\beta_0}$$
(260)

$$\geq \log \frac{1}{\alpha_0} + \left(\min\{\alpha(w), \overline{\alpha(w)}\} + \min\{\beta(w'), \overline{\beta(w')}\right) \log \frac{\alpha_0}{\beta_0}.$$
(261)

Here  $\overline{a} = 1 - a$ .

Define  $\alpha'(W) := |\alpha(W) - \frac{1}{2}|, \beta'(W) := |\beta(W) - \frac{1}{2}|,$  $\gamma(W) := \frac{\alpha'(W) + \beta'(W)}{2}, \quad \delta(W) := \gamma^2(W), \text{ and } \theta := \sqrt{\mathbb{E}\delta(W)}.$  Then we can lower bound  $\Gamma^{\text{LB}}(\pi_{XY})$  as (262)-(269) (given on page 26), where (264) follows from [2, Prop. 3.2]; (265) follows since  $-H_2(t)$  is convex in t; (267) follows from [2, Prop. 3.3] and the fact  $x \mapsto \sqrt{x}$  is a concave function; (269) follows since the objective function in (268) is nondecreasing in  $\theta$  (this can be seen from the facts that the stationary point  $\theta^* = \frac{1}{2} \left( \frac{\alpha_0}{\beta_0} - 1 \right) / \frac{\alpha_0}{\beta_0} + 1$  of the objective function is not larger than  $\sqrt{\alpha_0 - \frac{1}{4}}$ , the objective function is convex, and the derivative of the objective function is continuous).

Substituting  $a = \frac{1}{2} + \sqrt{\alpha_0 - \frac{1}{4}}$  into (269) , we obtain the desired result.

#### APPENDIX D **PROOF OF THEOREM 5**

Achievability Part: The achievability part is obtained by the following lemma.

**Lemma 10** (One-Shot Soft-Covering). [25] Assume  $P_W$  and  $P_{X|W}$  are unconditional and conditional distributions respectively (which can be defined on any countable or uncountable alphabets). Consider a random codebook  $\mathcal{C} = \{W(i)\}_{i \in \mathcal{M}}$ with  $W(i) \sim P_W, i \in \mathcal{M}$ , where  $\mathcal{M} = \{1, \ldots, e^R\}$ . We define

$$P_{X|\mathcal{C}}(\cdot|\{w(i)\}_{i\in\mathcal{M}}) := \frac{1}{|\mathcal{M}|} \sum_{m\in\mathcal{M}} P_{X|W}(\cdot|w(m)) \quad (270)$$

Assume  $\pi_X$  is a distribution such that for some  $s \in (0, 1]$ ,  $D_{1+s}\left(P_{X|W} \| \pi_X | P_W\right)$  and  $D_{1+s}(P_X \| \pi_X)$  exist (and hence are finite). Then we have

$$e^{sD_{1+s}(P_{X|C} \|\pi_X|P_C)} \le e^{sD_{1+s}(P_{X|W} \|\pi_X|P_W) - sR} + e^{sD_{1+s}(P_X \|\pi_X)}.$$
 (271)

Now we set 
$$\pi_X, P_{X|W}, P_W, R$$
 to  $\pi_{XY}^n, P_{X|W}^n, P_{Y|W}^n, P_W^n, nR$  respectively<sup>15</sup> for some

<sup>15</sup>The pair  $(X^n, Y^n)$  plays the role of X in Lemma 10.

$$-H_{\widetilde{P}}(XY|WU) + \inf_{\substack{Q_{WUW'U'} \in \\ C(\widetilde{P}_{WU},\widetilde{P}_{WU})}} \sum_{w,u,w',u'} Q(w,u,w',u') \mathcal{H}(\widetilde{P}_{X|(W,U)=(w,u)},\widetilde{P}_{Y|(W,U)=(w',u')} \| \pi_{XY})$$

$$\leq -e^{-\epsilon}H(XY|W) + \sum_{u,u'} \widetilde{P}_{U}(u)\widetilde{P}_{U}(u')$$

$$\times \inf_{\substack{Q_{WW'} \in \\ C(\widetilde{P}_{W|U=u},\widetilde{P}_{W|U=u'})}} \sum_{w,w'} Q(w,w') \mathcal{H}(\widetilde{P}_{X|(W,U)=(w,u)},\widetilde{P}_{Y|(W,U)=(w',u')} \| \pi_{XY})$$
(241)

$$\leq -e^{-\epsilon}H(XY|W) + e^{-2\epsilon} \inf_{Q_{WW'} \in C(P_W, P_W)} \sum_{w,w'} Q(w, w') \mathcal{H}(P_{X|W=w}, P_{Y|W=w'} \| \pi_{XY}) + (1 - e^{-2\epsilon}) \max_{w,w'} \log \frac{1}{2\epsilon}$$
(242)

$$+ (1 - e^{-\epsilon}) \max_{(x,y)\in \operatorname{supp}(\pi_{XY})} \log \frac{1}{\pi(x,y)}$$

$$\leq e^{-\epsilon} \left( -H(XY|W) + \inf_{\substack{Q_{WW'}\in C(P_W,P_W)}} \sum_{w,w'} Q(w,w') \mathcal{H}(P_{X|W=w}, P_{Y|W=w'} \| \pi_{XY}) \right) + O(\epsilon).$$
(242)
$$(242)$$

$$\begin{split} \widetilde{T}_{\infty}(\pi_{XY}) &\geq \lim_{\epsilon \downarrow 0} \inf_{\substack{P_{W}P_{X|W}P_{Y|W}:\\ D_{\infty}(P_{XY} || \pi_{XY}) \leq \epsilon}} \epsilon^{\epsilon} \left\{ -H_{\tilde{P}}(XY|WU) \\ &+ \inf_{\substack{Q_{WUW'U'} \in \\ C(\tilde{P}_{WU,\tilde{P}_{WU}}) w, u, w', u'}} \sum_{\substack{Q(w, u, w', u') \mathcal{H}(\tilde{P}_{X|(W,U)=(w,u)}, \tilde{P}_{Y|(W,U)=(w',u')} || \pi_{XY}) + O(\epsilon)} \right\} \quad (244) \\ &= \lim_{\epsilon \downarrow 0} \inf_{\substack{P_{W}P_{X|W}P_{Y|W}: \\ D_{\infty}(P_{XY} || \pi_{XY}) \leq \epsilon}} -H_{\tilde{P}}(XY|WU) \\ &+ \inf_{\substack{Q_{WUW'U'} \in \\ C(\tilde{P}_{WU,\tilde{P}_{WU}}) w, u, w', u'}} \sum_{\substack{Q(w, u, w', u') \mathcal{H}(\tilde{P}_{X|(W,U)=(w,u)}, \tilde{P}_{Y|(W,U)=(w',u')} || \pi_{XY})} \\ &\geq \inf_{\substack{\tilde{P}_{WU} \tilde{P}_{X|W} \tilde{P}_{Y|WU}} \sum_{\substack{P_{XY} = \pi_{XY} \\ P_{XY} = \pi_{XY}}} -H_{\tilde{P}}(XY|WU) \\ &+ \inf_{\substack{Q_{WUW'U'} \in \\ C(\tilde{P}_{WU,\tilde{P}_{WU})} w, u, w', u'}} Q(w, u, w', u') \mathcal{H}(\tilde{P}_{X|(W,U)=(w,u)}, \tilde{P}_{Y|(W,U)=(w',u')} || \pi_{XY}) \\ &= \Gamma^{\text{LB}}(\pi_{XY}). \quad (245) \end{split}$$

distribution  $P_W P_{X|W} P_{Y|W}$  such that the marginal distribution of  $P_W P_{X|W} P_{Y|W}$  on (X, Y) is equal to  $\pi_{XY}$ . Then Lemma 10 implies that if

$$R > D_{1+s}(P_{X|W}P_{Y|W} \| \pi_{XY} | P_W), \qquad (272)$$

then  $D_{1+s}(P_{X^nY^n|\mathcal{C}_n} \| \pi_{XY}^n | \mathcal{P}_{\mathcal{C}_n}) \to 0$ . That is, there exists at least one sequence of codebooks indexed by  $\{c_n\}_{n=1}^{\infty}$  such that  $D(P_{X^nY^n|\mathcal{C}_n=c_n} \| \pi_{XY}^n) \le D_{1+s}(P_{X^nY^n|\mathcal{C}_n=c_n} \| \pi_{XY}^n) \to 0.$ This completes the achievability proof.

Converse Part: Observe that

ſ

$$R = \frac{1}{n} H\left(M\right) \tag{273}$$

$$\geq \frac{1}{n} I\left(X^n Y^n; M\right) \tag{274}$$

$$= \frac{1}{n} D\left( P_{X^n Y^n M} \| P_{X^n Y^n} P_M \right)$$
(275)

$$= \frac{1}{n} D\left( P_{X^n Y^n M} \| \pi_{XY}^n P_M \right) - \frac{1}{n} D\left( P_{X^n Y^n} \| \pi_{XY}^n \right).$$
(276)

We lower bound the first term in (276) as follows:

$$\frac{1}{n}D\left(P_{X^{n}Y^{n}M}\|\pi_{XY}^{n}P_{M}\right) = \frac{1}{n}\sum_{i=1}^{n}D\left(P_{X_{i}Y_{i}|MX^{i-1}Y^{i-1}}\|\pi_{XY}|P_{MX^{i-1}Y^{i-1}}\right) \quad (277)$$

Ι

$$\geq \frac{1}{n} \sum_{i=1}^{n} D\left( P_{X_i Y_i | M} \| \pi_{XY} | P_M \right)$$
(278)

$$= D\left(P_{X_J Y_J | MJ} \| \pi_{XY} | P_{MJ}\right) \tag{279}$$

$$= D\left(P_{XY|W} \| \pi_{XY} | P_W\right), \tag{280}$$

where (277) follows by chain rule, (278) follows by the convexity of relative entropy [19, Theorem 2.7.2], in (279),  $J \sim P_J := \text{Unif}[1 : n]$  is a time index independent of  $(M, X^n, Y^n)$ , and in (280),  $X := X_J, Y := Y_J, W := MJ$ .

On the other hand, by assumption, the second term in (276) satisfies -1

$$\frac{1}{n}D\left(P_{X^{n}Y^{n}}\|\pi_{XY}^{n}\right) \to 0$$
(281)

$$\begin{split} \Gamma^{\text{LB}}(\pi_{XY}) \\ &\geq \inf_{\substack{B_{V},\alpha(1),\beta(1)\\B_{Z}(W)=\frac{1}{2}\\B_{Z}(W)=$$

as  $n \to \infty$ . Moreover, by similar derivation as (277)-(278), we can lower bound it as follows:

$$\frac{1}{n}D\left(P_{X^{n}Y^{n}}\|\pi_{XY}^{n}\right) \ge D\left(P_{XY}\|\pi_{XY}\right).$$
(282)

Hence combining (276), (280), (281), and (282) yields that

$$R \ge \lim_{\epsilon \downarrow 0} \inf_{P_W P_X|_W P_Y|_W : D(P_{XY} \| \pi_{XY}) \le \epsilon} D\left(P_{XY|W} \| \pi_{XY} | P_W\right)$$
(283)

$$= \lim_{\epsilon \downarrow 0} \inf_{P_W P_{X|W} P_{Y|W}: D(P_{XY} \parallel \pi_{XY}) \le \epsilon} I(XY; W).$$
(284)

### APPENDIX E Proof of Corollary 2

First we introduce the following lemma, which upper bounds Rényi divergences in terms of Rényi entropies.

**Lemma 11.** For a distribution  $P_{UV}$  with  $\mathcal{U}$  countable, we have for  $s \in [-1, \infty]$ ,

$$D_{1+s}(P_{UV} \| P_U P_V) \le H_{1-s}(P_U)$$
(285)

Proof: Consider,

$$D_{1+s}(P_{UV} \| P_U P_V) = \frac{1}{s} \log \mathbb{E}_{P_{UV}} \left( \frac{P_{U|V}(U|V)}{P_U(U)} \right)^s \quad (286)$$
  
$$\leq \frac{1}{s} \log \mathbb{E}_{P_{UV}} \left( \frac{1}{P_U(U)} \right)^s \quad (287)$$

$$= \frac{1}{s} \circ P_{U} \left( P_{U}(U) \right)$$
$$= H_{1} \left( P_{U} \right)$$
(288)

$$H_{1-s}(P_U).$$
 (288)

By the lemma above, we obtain  $D_{1+s}(P_{X|W}P_{Y|W}||P_{XY}|P_W) \leq H_{1-s}(\pi_{XY}) < \infty$  for all  $P_W P_{X|W} P_{Y|W}$  such that  $P_{XY} = \pi_{XY}$ . Then by Proposition 1, we have that

$$\widehat{C}_{\text{Wyner}}(\pi_{XY}) = C_{\text{Wyner}}(\pi_{XY}).$$
(289)

Furthermore, by Theorem 5, to prove Corollary 2, we only need prove

$$\widetilde{C}_{\mathrm{Wyner}}(\pi_{XY}) \ge \widehat{C}_{\mathrm{Wyner}}(\pi_{XY}).$$
 (290)

Hence we only need prove  $C_{Wyner}(\pi_{XY}) \ge C_{Wyner}(\pi_{XY})$ . In this appendix, we combine the distribution truncation technique and the mixture decomposition to prove this. Without loss of generality, we assume X, Y are integervalued. Define the *n*-truncation operator  $[\cdot]_n$  as follows:  $[z]_n := z$  if  $|z| \le n$ , and  $[z]_n := n + 1$  if |z| > n. We introduce a random variable (in fact, a function of (X, Y) or  $([X]_n, [Y]_n)$ )

$$V := 1\left\{ (X, Y) \in [-n:n]^2 \right\} = 1\left\{ ([X]_n, [Y]_n) \in [-n:n]^2 \right\}$$
(291)

Hence  $P_{V|W[X]_n[Y]_n}(v|w, x, y) = 1 \{(x, y) \in [-n : n]^2\}$ , and  $q_n := P_V(1) = P_{[X]_n[Y]_n} ([-n : n]^2)$ . Then

$$\widetilde{C}_{Wyner}(\pi_{XY}) = \lim_{\substack{\epsilon \downarrow 0 \\ PWP_{X|W}P_{Y|W}: \\ D(P_{XY} \parallel \pi_{XY}) \le \epsilon}} I(XY; W)$$
(292)

 $\geq \limsup_{n \to \infty} \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_X | W P_Y | W:\\ D(P_{XY} \| \pi_{XY}) \leq \epsilon}} I\left( \left[ X \right]_n \left[ Y \right]_n; W \right)$ (293)

$$\geq \limsup_{n \to \infty} \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_{X|W} P_{Y|W}:\\ D\left(P_{[X]_n[Y]_n} \| \pi_{[X]_n[Y]_n}\right) \leq \epsilon}} I\left(\left[X\right]_n \left[Y\right]_n; W\right)$$
(294)

$$= \limsup_{n \to \infty} \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_{[X]_n \mid W} P_{[Y]_n \mid W}:\\D\left(P_{[X]_n [Y]_n} \parallel \pi_{[X]_n [Y]_n}\right) \le \epsilon}} I\left([X]_n [Y]_n; W\right)$$
(295)

$$= \limsup_{n \to \infty} \min_{\substack{P_W P_{[X]_n | W} P_{[Y]_n | W}:\\P_{[X]_n [Y]_n} = \pi_{[X]_n [Y]_n}}} I\left([X]_n [Y]_n; W\right), \quad (296)$$

where (293) follows by the data processing inequality  $I([X]_n[Y]_n; W) \leq I(XY; W)$ ; (294) follows by the data processing inequality  $D(P_{[X]_n[Y]_n} || \pi_{[X]_n[Y]_n}) \leq$  $D(P_{XY} || \pi_{XY})$ ; (295) follows since the objective function and the constraint depend on (W, X, Y) through their truncated version  $(W, [X]_n, [Y]_n)$ ; and (296) follows since the alphabet size of W can be restricted to be no larger than  $(2n+1)^2$  (by standard cardinality bounding techniques) and hence for such discrete W, the probability simplex defined on the alphabet of (W, X, Y) is compact.

By basic information-theoretic inequalities, we obtain that

$$I([X]_{n}[Y]_{n};W) = I([X]_{n}[Y]_{n}V;W)$$
(297)  
=  $I([X]_{n}[Y]_{n}W|V) + I(V;W)$ (298)

$$\geq I\left([X]_{n}[Y]_{n};W|V\right)$$
(299)

$$\geq q_n I([X]_n [Y]_n; W | V = 1).$$
 (300)

Observe that under the condition  $P_{[X]_n[Y]_n} = \pi_{[X]_n[Y]_n}$ , it holds that  $q_n = P_{[X]_n[Y]_n}([-n:n]^2) = \pi_{[X]_n[Y]_n}([-n:n]^2) \to 1 \text{ as } n \to \infty$ . Hence

$$I([X]_{n}[Y]_{n};W) \ge I([X]_{n}[Y]_{n};W|V=1).$$
(301)

Combining (296) and (301), we obtain

$$\widehat{C}_{\text{Wyner}}(\pi_{XY}) \\
\geq \limsup_{n \to \infty} \sup_{\substack{P_{W}P_{[X]_{n}|W}P_{[Y]_{n}|W}:\\P_{[X]_{n}[Y]_{n}}=\pi_{[X]_{n}[Y]_{n}}} I\left([X]_{n}[Y]_{n};W|V=1\right).$$
(302)

To simplify the RHS of (302), we need the following lemma.

**Lemma 12** (Conditional Markov Chain). If  $X \to W \to Y$ form a Markov chain, then  $X \to W \to Y$  also form a Markov chain conditioned on  $\{X \in A, Y \in B\}$  for any  $A \subseteq \mathcal{X}, B \subseteq \mathcal{Y}$  such that  $\mathbb{P}(X \in A), \mathbb{P}(Y \in B) > 0$ .

Proof: Consider,

$$\mathbb{P}((W, X, Y) = (w, x, y) | X \in A, Y \in B) 
= \frac{P_W(w) P_{X|W}(x|w) P_{Y|W}(y|w) 1\{(x, y) \in A \times B\}}{P_{XY}(A \times B)}$$
(303)

$$= \frac{P_{W}(w)P_{X|W}(A|w)P_{Y|W}(B|w)}{P_{XY}(A \times B)} \times \frac{P_{X|W}(x|w)1\{x \in A\}}{P_{X|W}(A|w)} \frac{P_{Y|W}(y|w)1\{y \in B\}}{P_{Y|W}(B|w)}$$
(304)

$$=: \widetilde{P}_{W}(w)\widetilde{P}_{X|W}(x|w)\widetilde{P}_{Y|W}(y|w), \tag{305}$$

i.e.,  $X \to W \to Y$  forms a Markov chain under  $\tilde{P}$ . By Lemma 12, for  $(x, y) \in [-n : n]^2$ ,  $P_{W[X]_n[Y]_n|V}(w, x, y|1)$  can be factorized as

$$P_{W[X]_n[Y]_n|V}(w, x, y|1) = \widetilde{P}_W(w)\widetilde{P}_{X|W}(x|w)\widetilde{P}_{Y|W}(y|w)$$
(306)

i.e.,  $X \to W \to Y$  forms a Markov chain under  $\widetilde{P}$ . Hence

$$I([X]_{n}[Y]_{n};W|V=1) = I_{\widetilde{P}}(XY;W).$$
(307)

On the other hand,  $P_{[X]_n[Y]_n} = \pi_{[X]_n[Y]_n}$  implies

$$\sum_{w} \widetilde{P}_{W}(w) \widetilde{P}_{X|W}(x|w) \widetilde{P}_{Y|W}(y|w)$$
$$= P_{[X]_{n}[Y]_{n}|V}(x,y|1)$$
(308)

$$=\frac{\pi_{[X]_n[Y]_n}(x,y)P_{V|[X]_n[Y]_n}(1|x,y)}{P_V(1)}$$
(309)

$$=\frac{\pi_{[X]_n[Y]_n}(x,y)1\left\{(x,y)\in[-n,n]^2\right\}}{\pi_{[X]_n[Y]_n}\left([-n,n]^2\right)}$$
(310)

$$=:\pi_{XY}^{(n)}(x,y).$$
(311)

(299) Hence (302) implies that

$$C_{\text{Wyner}}(\pi_{XY}) \geq \limsup_{n \to \infty} \min_{\widetilde{P}_{W} \widetilde{P}_{X|W} \widetilde{P}_{Y|W}: \widetilde{P}_{XY} = \pi_{XY}^{(n)}} I_{\widetilde{P}}(XY;W) \quad (312)$$

$$= \limsup_{n \to \infty} C_{\text{Wyner}}(\pi_{XY}^{(n)}).$$
(313)

Next we prove  $C_{\text{Wyner}}(\pi_{XY}) \leq \lim_{n \to \infty} \inf_{0 \to \infty} C_{\text{Wyner}}(\pi_{XY}^{(n)})$ . Obviously,  $p_n := \pi_{XY}([-n,n]^2) \to 1$  as  $n \to \infty$ . Then for  $(x,y) \in \operatorname{supp}(\pi_{XY})$ ,

$$\frac{\pi_{XY}^{(n)}(x,y)}{\pi_{XY}(x,y)} = \frac{1\left\{(x,y)\in[-n,n]^2\right\}}{p_n}$$
(314)

$$\leq \frac{1}{p_n},\tag{315}$$

and

$$H\left(\pi_{XY}^{(n)}\right) = -\sum_{(x,y)\in[-n,n]^2} \frac{\pi_{XY}(x,y)}{p_n} \log \frac{\pi_{XY}(x,y)}{p_n}$$
(316)  
$$= \log p_n - \frac{1}{p_n} \sum_{(x,y)\in[-n,n]^2} \pi_{XY}(x,y) \log \pi_{XY}(x,y).$$
(317)

According to the definition of entropy,

$$-\sum_{(x,y)\in[-n,n]^2}\pi_{XY}(x,y)\log\pi_{XY}(x,y)\to H(\pi_{XY})$$
(318)

as  $n \to \infty$ . Hence

$$\lim_{n \to \infty} H\left(\pi_{XY}^{(n)}\right) \to H(\pi_{XY}). \tag{319}$$

We construct a new distribution

$$\widehat{\pi}_{XY}^{(n)}(x,y) := \frac{\frac{1}{p_n} \pi_{XY}(x,y) - \pi_{XY}^{(n)}(x,y)}{\frac{1}{p_n} - 1}$$
(320)

$$=\frac{\pi_{XY}(x,y)\,\mathbf{1}\,\{(x,y)\notin[-n:n]^2\}}{1-p_n}.$$
 (321)

Hence  $\pi_{XY}$  can be written as a mixture distribution  $\pi_{XY}(x,y) = p_n \pi_{XY}^{(n)}(x,y) + (1-p_n) \hat{\pi}_{XY}^{(n)}(x,y)$ . Define U as a Bernoulli random variable U with  $P_U(1) = p_n$ . Define

$$Q_{XYWU}^{(n)}(x, y, w, u) = \begin{cases} p_n \pi_{XY}^{(n)}(x, y) P_{W|XY}^{(n)}(w|x, y) & \text{if } u = 1\\ (1 - p_n) \,\widehat{\pi}_{XY}^{(n)}(x, y) \, 1 \, \{w = (x, y)\} & \text{if } u = 0 \end{cases},$$
(322)

where  $P_{W|XY}^{(n)}$  is induced by an optimal joint distribution  $P_W^{(n)}P_{X|W}^{(n)}P_{Y|W}^{(n)}$  (with  $P_{XY}^{(n)} = \pi_{XY}^{(n)}$  and W having a finite support) attaining  $C_{Wyner}(\pi_{XY}^{(n)})$ . Obviously,  $Q_{XY}^{(n)} = \pi_{XY}$ , and  $X \to (W, U) \to Y$  under  $Q^{(n)}$ . Therefore, we have

$$C_{\text{Wyner}}(\pi_{XY}) = \inf_{\substack{P_W P_{X|W} : P_{XY} = \pi_{XY}}} I(XY;W)$$
(323)

$$\leq I_{Q^{(n)}}\left(XY;WU\right) \tag{324}$$

$$= H(\pi_{XY}) - H_{Q^{(n)}}(XY|WU)$$
(325)

$$= H(\pi_{XY}) - p_n H_{Q^{(n)}}(XY|W, U = 1)$$
(1 m)  $H_{Q^{(n)}}(YY|W, U = 0)$ 
(326)

$$= H(\pi_{XY}) - p_n H_{P^{(n)}}(XY|W)$$
(327)

$$= H(\pi_{XY}) - p_n H\left(\pi_{XY}^{(n)}\right) + p_n I_{P^{(n)}}(XY;W).$$
(328)

Taking limits and using (319) and the fact that  $p_n \to 1$  as  $n \to \infty$ , we have

$$C_{\text{Wyner}}(\pi_{XY}) \le \liminf_{n \to \infty} I_{P^{(n)}}(XY;W)$$
(329)

$$= \liminf_{n \to \infty} C_{\text{Wyner}}(\pi_{XY}^{(n)}). \tag{330}$$

Combining (313) and (330) gives us the desired result.

=

# Appendix F

#### PROOF OF COROLLARY 3

In this section, we extend the proof in Appendix E to the continuous distribution case by combining it with the discretization technique and dyadic decomposition results in [10].

By assumption,  $\widehat{C}_{Wyner}(\pi_{XY}) = C_{Wyner}(\pi_{XY})$ . Hence to prove Corollary 3, we only need to prove  $\widetilde{C}_{Wyner}(\pi_{XY}) \ge C_{Wyner}(\pi_{XY})$ . To this end, similar to (291), we introduce a random variable

$$V_d := 1\left\{ (X, Y) \in [-d, d)^2 \right\}.$$
(331)

Denote  $P_{WXY} = P_W P_{X|W} P_{Y|W}$ . Similarly to (306), we define  $\tilde{P}_{WXY}(\cdot) := P_{WXY|V_d}(\cdot|1)$ . Then  $\tilde{P}_{WXY} = \tilde{P}_W \tilde{P}_{X|W} \tilde{P}_{Y|W}$ , i.e.,  $X \to W \to Y$  forms a Markov chain under  $\tilde{P}$ . Define  $q_d := P_{XY}([-d, d)^2)$ . The conclusions similar to (307) and (311) hold.

A. Proof of 
$$C_{Wyner}(\pi_{XY}) \geq \lim_{d\to\infty} \widetilde{C}_{Wyner}(\pi_{XY}|_{V_d=1})$$

Consider that

$$\widetilde{C}_{\text{Wyner}}(\pi_{XY}) = \lim_{\substack{\epsilon \downarrow 0 \\ D(P_{XY} \mid ||\pi_{XY}) \leq \epsilon}} \prod_{I (XY;W)} I(XY;W)$$
(332)

$$\geq \limsup_{d \to \infty} \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_X | W P_Y | W:\\ D(P_{\mathbf{Y}\mathbf{Y}} || \pi_{\mathbf{Y}\mathbf{Y}}) \leq \epsilon}} q_d I\left(XY; W | V_d = 1\right) \quad (333)$$

$$= \limsup_{d \to \infty} \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_{X|W} P_{Y|W}:\\D(P_{XY} \| \pi_{XY}) \le \epsilon}} \pi_{XY} \left( [-d, d)^2 \right)$$
$$\times I \left( XY; W | V_d = 1 \right)$$
(334)

$$d \to \infty^{-\epsilon} \epsilon \downarrow 0 \qquad P_W P_{X|W} P_{Y|W}: \\ (\pi_{XY} ([-d,d)^2) - \sqrt{2\epsilon}) D(P_{XY|V_d=1} \| \pi_{XY|V_d=1}) \leq \epsilon \\ \pi_{XY} ([-d,d)^2) I(XY; W|V_d=1)$$
(335)

$$= \limsup_{d \to \infty} \lim_{\epsilon \downarrow 0} \inf_{\substack{P_W P_X | W P_Y | W:\\ D\left(P_{XY} | V_d = 1 || \pi_{XY} | V_d = 1\right) \le \epsilon}} I\left(XY; W | V_d = 1\right)$$

$$\geq \limsup_{d \to \infty} \lim_{\epsilon \downarrow 0} \inf_{\substack{\tilde{P}_{W} \tilde{P}_{X|W} \tilde{P}_{Y|W}:\\D(\tilde{P}_{XY} \| \pi_{XY|V,-1}) \le \epsilon}} I_{\tilde{P}}(XY;W)$$
(337)

$$= \limsup_{d \to \infty} \widetilde{C}_{\text{Wyner}}(\pi_{XY|V_d=1}), \tag{338}$$

where (333) follows similarly as (297)-(300); (334) follows from that by Pinsker's inequality  $|P_{XY} - \pi_{XY}| \le \sqrt{2D(P_{XY}||\pi_{XY})} \le \sqrt{2\epsilon}$ , we have

$$q_d \in \pi_{XY}\left([-d,d)^2\right) + \left[-\sqrt{2\epsilon},\sqrt{2\epsilon}\right];\tag{339}$$

(335) follows from (339) and the fact that

$$q_{d}D\left(P_{XY|V_{d}=1} \| \pi_{XY|V_{d}=1}\right) \\ \leq D\left(P_{XY|V_{d}} \| \pi_{XY|V_{d}} | P_{V_{d}}\right)$$
(340)

$$\leq D\left(P_{XY|V_d} \| \pi_{XY|V_d} | P_{V_d}\right) + D\left(P_{V_d} \| \pi_{V_d}\right)$$
(341)

$$= D\left(P_{XYV_d} \| \pi_{XYV_d}\right) \tag{342}$$

$$= D\left(P_{XY} \| \pi_{XY}\right),\tag{343}$$

((343) follows since  $V_d$  is a function of (X, Y)); and (336) follows since  $\pi_{XY}([-d,d)^2) \to 1$  as  $d \to \infty$ .

B. Proof of 
$$C_{\text{Wyner}}(\pi_{XY}) \leq \liminf_{d \to \infty} \widetilde{C}_{\text{Wyner}}(\pi_{XY|V_d=1})$$

Next prove  $C_{\text{Wyner}}(\pi_{XY})$ we  $\liminf_{d\to\infty} C_{\text{Wyner}}(\pi_{XY|V_d=1})$ . Since in definition of  $C_{\text{Wyner}}(\pi_{XY|V_d=1})$ , a joint distribution  $\widetilde{P}_W \widetilde{P}_{X|W} \widetilde{P}_{Y|W}$ generates a distribution  $P_{XY}$ , which is an approximate version of  $\pi_{XY|V_d=1}$  and hence is also an approximation of  $\pi_{XY}$ . In this subsection, we combine mixture decomposition technique with dyadic decomposition schemes [10] to make the joint distribution  $P_{XY}$  exactly equal to  $\pi_{XY}$  (by constructing a modified version of  $P_W P_{X|W} P_{Y|W}$ ).

Define

$$\pi_{XY}^{(d)}(x,y) := \frac{1}{p_d} \pi_{XY}(x,y) \mathbb{1}\left\{ (x,y) \in [-d,d)^2 \right\}$$
(344)  
=  $\pi_{XY}(y,y) \mathbb{1}\left\{ (x,y) \in [-d,d)^2 \right\}$ (345)

$$=\pi_{XY|V_d=1},\tag{345}$$

where  $p_d := \pi_{XY} \left( [-d, d)^2 \right) \rightarrow 1$  as  $d \rightarrow \infty$ . Then given an integer n > 0, we define  $\Delta := \frac{d}{n}$ and we quantize X, Y as  $A := \lfloor \frac{X}{\Delta} \rfloor, B_{(-)} := \lfloor \frac{Y}{\Delta} \rfloor$ . The induced distribution of (A, B) is  $\pi_{AB}^{(n)}(a, b)$  $\begin{array}{l} \frac{1}{p_d} \int_{\Delta(a,b)+[0,\Delta)^2} \pi_{XY}(x,y) \mathrm{d}x \mathrm{d}y \mathbbm{1}\left\{(a,b) \in [-n,n-1]^2\right\}. \\ \text{By adding an independent uniform vector } (U,V) \sim \end{array}$ Unif $([0,\Delta)^2)$  to  $\Delta(A,B)$  with  $(A,B) \sim \pi_{AB}^{(n)}$ , we get a continuous distribution

$$\Delta(A,B) + (U,V) \sim \pi_{XY}^{(n)}(x,y) \coloneqq \frac{1}{\Delta^2} \pi_{AB}^{(n)}\left(\left\lfloor \frac{x}{\Delta} \right\rfloor, \left\lfloor \frac{y}{\Delta} \right\rfloor\right).$$
(346)

Then for  $(x, y) \in \text{supp}(\pi_{XY})$ ,

$$\frac{\pi_{XY}^{(n)}(x,y)}{\pi_{XY}(x,y)} = \frac{\frac{1}{\Delta^2} \pi_{AB}^{(n)}(\lfloor \frac{x}{\Delta} \rfloor, \lfloor \frac{y}{\Delta} \rfloor)}{\pi_{XY}(x,y)} \tag{347}$$

$$= \frac{\frac{1}{\Delta^2} \int_{\Delta}(\lfloor \frac{x}{\Delta} \rfloor, \lfloor \frac{y}{\Delta} \rfloor) + [0,\Delta)^2 \pi_{XY}(x,y) \mathrm{d}x \mathrm{d}y}{\pi_{XY}(x,y)p_d}$$

$$\times 1\left\{ (x,y) \in [-d,d)^2 \right\}$$

$$\pi \dots \left\{ \widehat{x} \ \widehat{y} \right\} \left\{ (x,y) \in [-d,d)^2 \right\}$$
(348)

$$=\frac{\pi_{XY}(x,y)1\{(x,y)\in [-d,d)^2\}}{\pi_{XY}(x,y)p_d}$$
(349)

$$\leq \sup_{(x,y)\in [-d,d)^2} \frac{\pi_{XY}(\hat{x},\hat{y})}{\pi_{XY}(x,y)p_d}$$
(350)

where (349) follows by the mean value theorem, and it holds for some  $(\widehat{x}, \widehat{y}) \in \Delta\left(\left\lfloor \frac{x}{\Delta} \right\rfloor, \left\lfloor \frac{y}{\Delta} \right\rfloor\right) + [0, \Delta)^2$ .

**Lemma 13.** Assume  $\pi_{XY}$  is differentiable. Then for any  $(x, y), (\hat{x}, \hat{y}) \in [-d, d]^2$  satisfying  $|x - \hat{x}|, |y - \hat{y}| \leq \Delta$ , we have

$$\exp\left(-\Delta L_d\right) \le \frac{\pi_{XY}\left(x,y\right)}{\pi_{XY}\left(\hat{x},\hat{y}\right)} \le \exp\left(\Delta L_d\right),\tag{351}$$

where  $L_d$  is defined in (88).

Proof of Lemma 13: By Taylor's theorem,

$$\log \pi_{XY}(x,y) = \log \pi_{XY}(\hat{x},\hat{y}) + \frac{\partial}{\partial x} \log \pi_{XY}(\tilde{x},\tilde{y})(x-\hat{x}) + \frac{\partial}{\partial y} \log \pi_{XY}(\tilde{x},\tilde{y})(y-\hat{y})$$
(352)  
$$\leq \log \pi_{XY}(\hat{x},\hat{y}) + \left( \left| \frac{\partial \pi_{XY}}{\partial x}(\tilde{x},\tilde{y}) \right| + \left| \frac{\partial \pi_{XY}}{\partial y}(\tilde{x},\tilde{y}) \right| \right) \Delta$$

$$\leq \log \pi_{XY}(x, y) + \frac{\pi_{XY}(\widetilde{x}, \widetilde{y})}{\pi_{XY}(\widetilde{x}, \widetilde{y})}$$
(353)

$$\leq \log \pi_{XY}\left(\hat{x},\hat{y}\right) + \Delta L_d,\tag{354}$$

where (352) holds for some  $(\tilde{x}, \tilde{y})$  on the line segment joining  $(\hat{x}, \hat{y})$  and (x, y). By symmetry,  $\log \pi_{XY}(\hat{x}, \hat{y}) \leq$  $\log \pi_{XY}(x,y) + \Delta L_d$  also holds.

Using Lemma 13, we obtain

$$\frac{\pi_{XY}^{(n)}(x,y)}{\pi_{XY}(x,y)} \le \frac{1}{p_d} \exp\left(\Delta L_d\right) = \exp\left(\Delta L_d - \log p_d\right).$$
(355)

Define

$$\epsilon'_n := \Delta L_d - \log p_d + \delta_n \tag{356}$$

for some positive sequence  $\delta_n \to 0$  as  $n \to \infty$ , which will be specified later. Then (355) implies

$$e^{D_{\infty}\left(\pi_{XY}^{(n)} \| \pi_{XY}\right)} = \sup_{x,y} \frac{\pi_{XY}^{(n)}(x,y)}{\pi_{XY}(x,y)} \le e^{\epsilon'_n - \delta_n}, \qquad (357)$$

i.e.,

$$\frac{e^{\epsilon'_n}\pi_{XY}(x,y)}{\pi_{XY}^{(n)}(x,y)} \ge e^{\delta_n} \tag{358}$$

for all  $(x, y) \in [-d, d)^2$ .

 $\langle \rangle$ 

We construct a new distribution

$$\widehat{\pi}_{XY}^{(n)}(x,y) := \frac{e^{\epsilon'_n} \pi_{XY}(x,y) - \pi_{XY}^{(n)}(x,y)}{e^{\epsilon'_n} - 1}.$$
(359)

Hence  $\pi_{XY}$  can be written as a mixture distribution  $\pi_{XY}(x,y) = e^{-\epsilon'_n} \pi_{XY}^{(n)}(x,y) + \left(1 - e^{-\epsilon'_n}\right) \widehat{\pi}_{XY}^{(n)}(x,y).$  Furthermore, by (358), we have

$$\widehat{\pi}_{XY}^{(n)}(x,y) \ge \frac{e^{\delta_n} - 1}{e^{\epsilon'_n} - 1} \pi_{XY}^{(n)}(x,y)$$
(360)

$$= \frac{e^{\epsilon_n} - 1}{e^{\epsilon'_n} - 1} \frac{1}{\Delta^2} \pi^{(n)}_{AB} \left( \left\lfloor \frac{x}{\Delta} \right\rfloor, \left\lfloor \frac{y}{\Delta} \right\rfloor \right).$$
(361)

Define U as a Bernoulli random variable U with  $P_U(1) =$  $e^{-\epsilon'_n}$ . Let  $[z]_n := z$ , if  $z \in [-n, n-1]$ ; n, if  $z \ge n$ ; and -(n+1), otherwise, denote the truncation operation on integers. Define

$$\widehat{\pi}_{XY|W_{1}}^{(n)}(x',y'|w_{1}) = \frac{\widehat{\pi}_{XY}^{(n)}(x',y') \mathbf{1}\left\{\left(\left[\left\lfloor\frac{x'}{\Delta}\right\rfloor\right]_{n},\left[\left\lfloor\frac{y'}{\Delta}\right\rfloor\right]_{n}\right) = w_{1}\right\}}{\widehat{\pi}_{XY}^{(n)}\left\{(x',y'):\left(\left[\left\lfloor\frac{x'}{\Delta}\right\rfloor\right]_{n},\left[\left\lfloor\frac{y'}{\Delta}\right\rfloor\right]_{n}\right) = w_{1}\right\}}$$
(362)

for  $w_1 \in [-(n+1), n]^2$ . Define

$$\begin{aligned} Q_{XYWU}^{(n)}(x,y,w,u) \\ &:= \begin{cases} e^{-\epsilon'_n} \pi_{XY}^{(n)}(x,y) P_{W|AB}^{(n)}\left(w \mid \left\lfloor \frac{x}{\Delta} \right\rfloor, \left\lfloor \frac{y}{\Delta} \right\rfloor\right) & \text{if } u = 1\\ \left(1 - e^{-\epsilon'_n}\right) \widehat{\pi}_{XY}^{(n)}(x,y) \, \widehat{P}_{W|XY}^{(n)}(w|x,y) & \text{if } u = 0 \end{cases}, \end{aligned}$$

$$(363)$$

where  $P_{W|AB}^{(n)}$  is induced by an optimal joint distribution  $P_{W}^{(n)}P_{A|W}^{(n)}P_{B|W}^{(n)}$  (with  $P_{AB}^{(n)} = \pi_{AB}^{(n)}$  and W having a finite support) attaining  $C_{Wyner}(\pi_{AB}^{(n)})$ ;

$$\widehat{P}_{W|XY}^{(n)}((w_1, w_2)|x, y) \\
:= \widehat{P}_{W_1|XY}^{(n)}(w_1|x, y) \widehat{P}_{W_2|XYW_1}^{(n)}(w_2|x, y, w_1)$$
(364)

with  $\hat{P}_{W_1|XY}^{(n)}(w_1|x,y) = 1 \left\{ w_1 = \left( \left[ \left\lfloor \frac{x}{\Delta} \right\rfloor \right]_n, \left[ \left\lfloor \frac{y}{\Delta} \right\rfloor \right]_n \right) \right\}$  (i.e.,  $W = (W_1, W_2)$  and  $W_1 = \left( \left[ \left\lfloor \frac{x}{\Delta} \right\rfloor \right]_n, \left[ \left\lfloor \frac{y}{\Delta} \right\rfloor \right]_n \right)$  under  $\hat{P}^{(n)}$ ); and  $\hat{P}_{W_2|XYW_1}^{(n)}$  is induced by an optimal joint distribution  $\hat{P}_{W_1W_2}^{(n)}\hat{P}_{X|W_1W_2}^{(n)}\hat{P}_{Y|W_1W_2}^{(n)}$  (with  $\hat{P}_{XY|W_1}^{(n)} = \hat{\pi}_{XY|W_1}^{(n)}$ ) such that  $\hat{P}_{W_2|W_1=w_1}^{(n)}\hat{P}_{X|W_2,W_1=w_1}^{(n)}\hat{P}_{Y|W_2,W_1=w_1}^{(n)}$ attains the common entropy  $G(\hat{\pi}_{XY|W_1=w_1}^{(n)})$  defined in (4) (or  $G(\hat{\pi}_{XY|W_1=w_1}^{(n)}) + \delta'_n$  for a sequence  $\delta'_n > 0$  satisfying  $\delta'_n \to 0$  as  $n \to \infty$  if the infimization in  $G(\hat{\pi}_{XY|W_1=w_1}^{(n)})$  is not attained) for  $w_1 \in [-(n+1), n]^2$ .

Partition  $\mathbb{R}^2$  into 9 subregions by the lines  $x = \pm d$  and  $y = \pm d$ . Denote them as  $\mathbb{R}_0, \mathbb{R}_1, ..., \mathbb{R}_8$ , where  $\mathbb{R}_0 := [-d, d)^2$  and  $\mathbb{R}_1, \mathbb{R}_2, ..., \mathbb{R}_8$  denote others. Obviously,  $\mathbb{R}_k, 0 \le k \le 8$  can be expressed as  $\mathbb{R}_k = I_1^{(k)} \times I_2^{(k)}$  with  $I_i^{(k)} \in \{\mathcal{L}_d^-, \mathcal{L}_d, \mathcal{L}_d^+\}$ , where  $\mathcal{L}_d^- := (-\infty, -d), \mathcal{L}_d := [-d, d)$ , and  $\mathcal{L}_d^+ := [d, +\infty)$ . Note that  $(X, Y) \in \mathbb{R}_0$  corresponds to  $W_1 \in [-n, n-1]^2$ ; and  $(X, Y) \in \bigcup_{k=1}^8 \mathbb{R}_k$  corresponds to the case that the first or the second component of  $W_1$  is -(n+1) or n. According to the definition of  $\widehat{\pi}_{XY|W_1}^{(n)}$ , for the subregion  $\mathbb{R}_0$ , we have  $\widehat{\pi}_{XY|W_1}^{(n)}(\cdot|(a,b)) = \widehat{\pi}_{XY}^{(n)}(\cdot|I_{\Delta}^2)$  with  $I_{\Delta}^2 := \Delta(a,b) + [0,\Delta)^2$  for  $(a,b) \in [-n,n-1]^2$ ; and for the subregion  $\mathbb{R}_k, 1 \le k \le 8$ , we have  $\widehat{\pi}_{XY|W_1}^{(n)}(\cdot|w_1) = \widehat{\pi}_{XY}^{(n)}(\cdot|\mathbb{R}_k) = \pi_{XY}^{(n)}(\cdot|\mathbb{R}_k)$  for some  $1 \le k \le 8$ , where the first or the second component of  $w_1$  is -(n+1) or n.

By the following lemma, we know that  $\widehat{\pi}_{XY}^{(n)}$  is log-concave.

**Lemma 14** (Invariance of Log-Concavity). [46, Exercise 3.48] If a pdf  $P_{Z^n}$  is log-concave, then for any  $0 \le a < \inf_{z^n} P_{Z^n}(z^n)$ ,  $P_{Z^n} - a$  is also log-concave.

Since  $\widehat{\pi}_{XY}^{(n)}$  is log-concave and so is  $\widehat{\pi}_{XY|W_1}^{(n)}(\cdot|w_1)$  for each  $w_1$ , the dyadic decomposition scheme in [10] can be applied to  $\widehat{\pi}_{XY|W_1}^{(n)}(\cdot|w_1)$ . Hence [10] implies that the common entropy  $G(\widehat{\pi}_{XY|W_1}^{(n)}(\cdot|w_1))$  defined in (4) satisfies

$$H_{\widehat{P}^{(n)}}(W_2|W_1 = w_1) = G(\widehat{\pi}_{XY|W_1}^{(n)}(\cdot|w_1))$$
(365)

$$\leq I_{\hat{\pi}_{XY|W_1}^{(n)}(\cdot|w_1)}(X;Y) + 24\log 2 \tag{366}$$

nats/symbol for  $w_1 \in [-(n+1), n]^2$ . We first consider the case of  $w_1 \in [-n, n-1]^2$ . For any square  $I_{\Delta}^2 = \Delta(a, b) + [0, \Delta)^2$ in  $\mathbb{R}_0$  with  $(a, b) \in [-n, n-1]^2$ , we have that

$$I_{\widehat{\pi}_{XY}^{(n)}}\left(X;Y|\left(X,Y\right)\in I_{\Delta}^{2}\right)$$

$$=\int_{I_{\Delta}^{2}}\widehat{\pi}^{(n)}(x,y|I_{\Delta}^{2})\log\frac{\widehat{\pi}^{(n)}(x,y|I_{\Delta}^{2})}{\widehat{\pi}^{(n)}(x|I_{\Delta})\widehat{\pi}^{(n)}(y|I_{\Delta})}\mathrm{d}x\mathrm{d}y \quad (367)$$

$$\widehat{\pi}^{(n)}(x,y)$$

$$= \int_{I_{\Delta}^{2}} \widehat{\pi}^{(n)}(x, y | I_{\Delta}^{2}) \log \frac{\frac{\pi}{\widehat{\pi}^{(n)}(I_{\Delta}^{2})}}{\frac{\widehat{\pi}^{(n)}(I_{\Delta})}{\widehat{\pi}^{(n)}(I_{\Delta}^{2})} \frac{\widehat{\pi}^{(n)}(I_{\Delta}, y)}{\widehat{\pi}^{(n)}(I_{\Delta}^{2})}} \mathrm{d}x \mathrm{d}y \qquad (368)$$
$$\widehat{\pi}^{(n)}(x, y)$$

$$\leq \sup_{(x,y)\in I_{\Delta}^{2}} \log \frac{\frac{\overline{\widehat{\pi}^{(n)}(I_{\Delta}^{(n)})}(I_{\Delta}^{(n)})}{\frac{\widehat{\pi}^{(n)}(I_{\Delta})}{\widehat{\pi}^{(n)}(I_{\Delta}^{(n)})} \frac{\widehat{\pi}^{(n)}(I_{\Delta},y)}{\widehat{\pi}^{(n)}(I_{\Delta}^{(n)})}}$$
(369)

$$= \sup_{(x,y)\in I_{\Delta}^2} \log \frac{\widehat{\pi}^{(n)}(x,y)\widehat{\pi}^{(n)}(I_{\Delta}^2)}{\widehat{\pi}^{(n)}(x,I_{\Delta})\widehat{\pi}^{(n)}(I_{\Delta},y)}$$
(370)

$$= \sup_{(x,y)\in I_{\Delta}^{2}} \log \frac{\widehat{\pi}^{(n)}(x,y)\widehat{\pi}^{(n)}(x',y')}{\widehat{\pi}^{(n)}(x,\widehat{y})\widehat{\pi}^{(n)}(\widehat{x},y)}$$
(371)

$$\leq \sup_{(x,y)\in I_{\Delta}^{2}} \log \left\{ \frac{1}{\left(\frac{e^{\delta_{n}}-1}{e^{\epsilon_{n}'}-1}\frac{1}{\Delta^{2}}\pi_{AB}^{(n)}(a,b)\right)^{2}} \times \left(\frac{e^{\epsilon_{n}'}\pi_{XY}(x,y) - \frac{1}{\Delta^{2}}\pi_{AB}^{(n)}(a,b)}{e^{\epsilon_{n}'}-1}\right) \times \left(\frac{e^{\epsilon_{n}'}\pi_{XY}(x',y') - \frac{1}{\Delta^{2}}\pi_{AB}^{(n)}(a,b)}{e^{\epsilon_{n}'}-1}\right) \right\}$$
(372)

$$\leq 2 \sup_{(x,y)\in I_{\Delta}^{2}} \log \frac{e^{\epsilon'_{n}} \pi_{XY}(x,y) - \frac{1}{\Delta^{2}} \pi_{AB}^{(n)}(a,b)}{(e^{\delta_{n}} - 1) \frac{1}{\Delta^{2}} \pi_{AB}^{(n)}(a,b)}$$
(373)

$$\leq 2\log\frac{e^{\epsilon'_{n}+\Delta L_{d}}\frac{1}{\Delta^{2}}\pi_{AB}^{(n)}(a,b)-\frac{1}{\Delta^{2}}\pi_{AB}^{(n)}(a,b)}{(e^{\delta_{n}}-1)\frac{1}{\Delta^{2}}\pi_{AB}^{(n)}(a,b)}$$
(374)

$$= 2\log\frac{e^{\epsilon'_{n} + \Delta L_{d}} - 1}{e^{\delta_{n}} - 1}$$
(375)

$$= 2\log\frac{(\epsilon'_{n} + \Delta L_{d})(1 + o(1))}{\delta_{n}(1 + o(1))}$$
(376)

$$= 2\log\left(\frac{\epsilon'_n + \Delta L_d}{\delta_n}\right) + o(1) \tag{377}$$

$$\leq \frac{4\Delta L_d - 2\log p_d}{\delta_n} + o(1), \qquad (378)$$

where (369) follows since the average is no greater than the supremum; (371) holds for some  $(x',y'), (\hat{x}, \hat{y}) \in I_{\Delta}^2$ , since by the mean value theorem,  $\hat{\pi}^{(n)}(I_{\Delta}^2) = \Delta^2 \hat{\pi}^{(n)}(x',y'), \hat{\pi}^{(n)}(x,I_{\Delta}) = \Delta \hat{\pi}^{(n)}(x,\hat{y}), \hat{\pi}^{(n)}(I_{\Delta},y) = \Delta \hat{\pi}^{(n)}(\hat{x},y)$  for some  $(x',y'), (\hat{x},\hat{y}) \in I_{\Delta}^2$ ; (372) follows from (359) and (361); (374) follows from (439); (375) follows from (445); in (376), o(1) denotes a term tending to zero as  $\epsilon'_n, \Delta L_d, \delta_n \to 0$ ; and (378) follows from (356). By introducing the positive sequence  $\delta_n$ , the denominators in equations after (372) are ensured to be positive. This is the reason why we introduce  $\delta_n$  in (356).

On the other hand, for the case of  $w_1 \notin [-n, n-1]^2$ , i.e., for the subregions  $R_k = I_1^{(k)} \times I_2^{(k)}, 1 \le k \le 8$ , we have

$$\widehat{\pi}_{XY}^{(n)}(\cdot|\mathbf{R}_{k}) = \pi_{XY}(\cdot|I_{1}^{(k)} \times I_{2}^{(k)}). \text{ Hence} 
I_{\widehat{\pi}_{XY}^{(n)}}(X;Y|(X,Y) \in \mathbf{R}_{k}) 
= I_{\pi}\left(X;Y|(X,Y) \in I_{1}^{(k)} \times I_{2}^{(k)}\right).$$
(379)

Now we bound the RHS of (379) by using the following lemma.

**Lemma 15** (Estimation of Conditional Mutual Information). Assume  $\pi_{XY}$  is an absolutely continuous distribution such that  $\lim_{x\to+\infty} \pi_X(x) = \lim_{x\to-\infty} \pi_X(x) = \lim_{y\to+\infty} \pi_Y(y) =$  $\lim_{y\to-\infty} \pi_Y(y) = 0$ . For  $A, B \in \{\mathcal{L}_d^-, \mathcal{L}_d, \mathcal{L}_d^+\}$ , we have

$$\pi \left( X;Y \right| \left( X,Y \right) \in A \times B \right) \le \Upsilon_{\pi} \left( A,B \right), \tag{380}$$

where

I

$$\Upsilon_{\pi} (A, B) = \begin{cases} \frac{1}{\pi_{XY}(A \times B)} \left( I_{\pi} \left( X; Y \right) + o(1) \right) & A = B = \mathcal{L}_{d} \\ \frac{1}{\pi_{XY}(A \times B)} o(1) & otherwise \end{cases}$$
(381)

and o(1) denotes a term tending to zero as  $d \to \infty$ .

The proof of Lemma 15 is deferred to Appendix F-B1.

It is easy to verify that a absolutely continuous log-concave pdf satisfies the conditions prescribed in Lemma 15. Hence by Lemma 15, we have

$$I_{\hat{\pi}_{XY}^{(n)}}(X;Y|(X,Y) \in \mathbf{R}_k) \le \Upsilon_{\pi}\left(I_1^{(k)}, I_2^{(k)}\right).$$
(382)

Substituting this into (366), we have

(---)

$$G(\widehat{\pi}_{XY|(X,Y)\in\mathbf{R}_{k}}^{(n)})$$

$$\leq I_{\widehat{\pi}_{XY}^{(n)}}(X;Y|(X,Y)\in\mathbf{R}_{k}) + 24\log 2$$
(383)

$$\leq \Upsilon_{\pi} \left( I_1^{(k)}, I_2^{(k)} \right) + 24 \log 2.$$
 (384)

According to the definition of  $Q_{XY}^{(n)}$ , we have  $Q_{XY}^{(n)} = \pi_{XY}$ , and  $X \to (W, U) \to Y$  under  $Q^{(n)}$ . Similarly to the countable case, we obtain that

$$C_{\text{Wyner}}(\pi_{XY}) = \inf_{\substack{P_W P_X|_W : P_{YY} = \pi_{XY}}} I(XY;W)$$
(385)

$$\leq I_{Q^{(n)}}\left(XY;WU\right) \tag{386}$$

$$= I_{Q^{(n)}} (XY; U) + e^{-\epsilon'_n} I_{Q^{(n)}} (XY; W|U = 1) + \left(1 - e^{-\epsilon'_n}\right) I_{Q^{(n)}} (XY; W|U = 0)$$
(387)

$$\leq H(U) + e^{-\epsilon'_{n}} I_{P^{(n)}}(XY;W) + \left(1 - e^{-\epsilon'_{n}}\right) H_{Q^{(n)}}(W_{1}W_{2}|U=0).$$
(388)

Since  $\epsilon'_n \to 0$  as  $n \to \infty$ , the first term in (388) is bounded as  $H(U) = H\left(e^{-\epsilon'_n}\right) \to 0$  as  $n \to \infty$ . For the second term in (388),

$$e^{-\epsilon'_{n}}I_{P^{(n)}}(XY;W) = e^{-\epsilon'_{n}}I_{P^{(n)}}(AB;W)$$
(389)

$$\leq C_{\rm Wyner}(\pi_{AB}^{(n)}) \tag{390}$$

$$= \widetilde{C}_{\mathrm{Wyner}}(\pi_{AB}^{(n)}) \tag{391}$$

$$\leq \widetilde{C}_{\mathrm{Wyner}}(\pi_{XY}^{(d)}),$$
 (392)

where (391) follows by Corollary 2 since  $\pi_{AB}^{(n)}$  is supported on a finite alphabet, and (392) follows by the data processing inequality.

We bound the last term in (388) as

$$H_{Q^{(n)}}(W_1 W_2 | U = 0)$$
  
=  $H_{\widehat{P}^{(n)}}(W_1 W_2)$  (393)

$$= H_{\widehat{P}(n)}(W_1) + H_{\widehat{P}(n)}(W_2|W_1)$$
(394)

$$\leq H_{\widehat{P}^{(n)}}(W_1) + \sum_{w_1} P^{(n)}(w_1) \\ \times \left( I_{\widehat{\pi}_{XY|W_1}^{(n)}}(\cdot|w_1)(X;Y) + 24\log 2 \right)$$
(395)

$$\leq 2\log(2n+2) + I_{\widehat{P}^{(n)}}(X;Y|W_1) + 24\log 2, \quad (396)$$

where (396) follows from  $H_{\widehat{P}^{(n)}}(W_1) \leq 2\log(2n+2)$  since  $W_1$  is defined on  $[-(n+1), n]^2$ .

On the other hand, by applying (378) and (384), we obtain that

$$\left(1 - e^{-\epsilon'_n}\right) I_{\widehat{p}(n)}(X; Y|W_1)$$

$$\leq \left(1 - e^{-\epsilon'_n}\right) \widehat{\pi}_{XY}^{(n)} \left(\left[-d, d\right)^2\right) \left(\frac{4\Delta L_d - 2\log p_d}{\delta_n} + o\left(1\right)\right)$$

$$+ \left(1 - e^{-\epsilon'_n}\right) \sum_{k=1}^8 \widehat{\pi}_{XY}^{(n)}(\mathbf{R}_k) \Upsilon_{\pi} \left(I_1^{(k)}, I_2^{(k)}\right)$$

$$\leq \left(1 - e^{-\epsilon'_n}\right) \left(\frac{4\Delta L_d - 2\log p_d}{\delta_n} + o\left(1\right)\right)$$

$$(397)$$

$$+\sum_{k=1}^{8}\pi_{XY}(\mathbf{R}_{k})\Upsilon_{\pi}\left(I_{1}^{(k)},I_{2}^{(k)}\right)$$
(398)

$$= \left(1 - e^{-\epsilon'_n}\right) \left(\frac{4\Delta L_d - 2\log p_d}{\delta_n} + o\left(1\right)\right) + o\left(1\right) \quad (399)$$

where (398) follows since  $\widehat{\pi}_{XY}^{(n)}([-d,d)^2) \leq 1$  and  $\widehat{\pi}_{XY}^{(n)}(\mathbf{R}_k) = \frac{\pi_{XY}(\mathbf{R}_k)}{1-e^{-\epsilon'_n}}$  (the latter follows by (359) and the fact that  $\pi_{XY}^{(n)}(x,y)$  is defined on  $[-d,d)^2$ ); and (399) follow by Lemma 15.

Combining (388), (392), (396), and (399) yields (400)-(401) (given on page 32).

Choose  $\delta_n = 2\Delta L_d - \log p_d$ , then to ensure that the RHS of (401) is no larger than  $\liminf_{d\to\infty} \widetilde{C}_{\text{Wyner}}(\pi_{XY}^{(d)})$ , we only require

$$\left(1 - e^{-(3\Delta L_d - 2\log p_d)}\right)\log n \to 0,\tag{402}$$

i.e.,

$$(3\Delta L_d - 2\log p_d)\log n \to 0. \tag{403}$$

Set  $\Delta$  to  $\Delta_d = (dL_d)^{-\alpha}$  for  $\alpha > 1$ . Recall  $n = \frac{d}{\Delta}$ . Then we have

$$\Delta L_d \log n \to 0. \tag{404}$$

Recall  $p_d = 1 - \epsilon_d$ . By the hypothesis that  $\epsilon_d \log (dL_d) \to 0$ as  $d \to +\infty$ , we have

$$(\log p_d) (\log n) \to 0. \tag{405}$$

$$C_{\text{Wyner}}(\pi_{XY}) \leq \liminf_{n \to \infty} \left\{ \widetilde{C}_{\text{Wyner}}(\pi_{XY}^{(d)}) + \left(1 - e^{-\epsilon_n'}\right) \left( 2\log(2n+2) + 24\log 2 + \frac{4\Delta L_d - 2\log p_d}{\delta_n} + o\left(1\right) \right) + o\left(1\right) \right\}$$
(400)  
$$= \liminf_{n \to \infty} \left\{ \widetilde{C}_{\text{Wyner}}(\pi_{XY}^{(d)}) + \left(1 - e^{-(\Delta L_d - \log p_d + \delta_n)}\right) \left( 2\log(2n+2) + \frac{4\Delta L_d - 2\log p_d}{\delta_n} \right) \right\}.$$
(401)

Hence for such a choice of  $\Delta,$  (403) is satisfied, which implies that

$$C_{\text{Wyner}}(\pi_{XY}) \le \liminf_{d \to \infty} \widetilde{C}_{\text{Wyner}}(\pi_{XY}^{(d)}).$$
(406)

1) Proof of Lemma 15 : Consider that

$$I_{\pi} (X; Y| (X, Y) \in A \times B)$$

$$= \int_{A \times B} \frac{\pi_{XY}(x, y)}{\pi_{XY}(A \times B)}$$

$$\times \log \frac{\pi_{XY}(x, y)\pi_{XY}(A \times B)}{\pi_{X}(x)\pi_{Y|X}(B|x)\pi_{Y}(y)\pi_{X|Y}(A|y)} dxdy \quad (407)$$

$$= \frac{1}{\pi_{XY}(A \times B)} \left\{ \int_{A \times B} \pi_{XY}(x, y) \log \frac{\pi_{XY}(x, y)}{\pi_{X}(x)\pi_{Y}(y)} dxdy + \log \pi_{XY}(A \times B) - \int_{A} \pi_{X}(x)\pi_{Y|X}(B|x) \log \pi_{Y|X}(B|x) dx - \int_{B} \pi_{Y}(y)\pi_{X|Y}(A|y) \log \pi_{X|Y}(A|y) dy \right\} \quad (408)$$

$$\leq \Upsilon_{\pi} \left( A, B \right) \tag{409}$$

where (409) follows from the facts that  $\log \pi_{XY}(A \times B) \leq 0$ and

$$\lim_{n \to \infty} \int_{A \times B} \pi_{XY}(x, y) \log \frac{\pi_{XY}(x, y)}{\pi_X(x)\pi_Y(y)} dx dy$$
$$= \begin{cases} I_{\pi}(X; Y) & A = B = \mathcal{L}_d \\ 0 & \text{otherwise} \end{cases},$$
(410)

as well as the following arguments. For all  $B \in \{\mathcal{L}_d^-, \mathcal{L}_d, \mathcal{L}_d^+\},\$ 

$$-\pi_X(x)\pi_{Y|X}(B|x)\log\pi_{Y|X}(B|x) \to 0$$
 (411)

pointwise,

$$\left|-\pi_X(x)\pi_{Y|X}(B|x)\log\pi_{Y|X}(B|x)\right| \le e^{-1}\pi_X(x)$$
 (412)

and  $e^{-1}\pi_X(x)$  is integrable. Hence by Lebesgue's dominated convergence theorem, we have

$$\lim_{n \to \infty} -\int_A \pi_X(x) \pi_{Y|X}(B|x) \log \pi_{Y|X}(B|x) dx = 0.$$
(413)

Similarly,

$$\lim_{n \to \infty} -\int_B \pi_Y(y) \pi_{X|Y}(A|y) \log \pi_{X|Y}(A|y) \mathrm{d}y = 0.$$
(414)

#### APPENDIX G Proof of Lemma 2

The proof techniques used in this section are similar to those used in Appendix E.

Assume  $(P_M, P_{X^n|M}, P_{Y^n|M})$  is a sequence of fixedlength codes with rate R that generates  $P_{X^nY^n}$  such that  $D_{\infty}(P_{X^nY^n} || \pi_{XY}^n) \to 0$ , where  $P_M$  is the uniform distribution on  $[1 : e^{nR}]$ . Similarly to (291), we introduce a random variable

$$V := 1\left\{ \left(X^{n}, Y^{n}\right) \in \mathcal{A}_{\epsilon}^{(n)}\left(\pi_{X}\right) \times \mathcal{A}_{\epsilon}^{(n)}\left(\pi_{Y}\right) \right\}.$$
(415)

Similarly to (306), we define  $\widetilde{P}_{MX^nY^n}$  :=  $P_{MX^nY^n|V}(m, x^n, y^n|1)$ . Then  $\widetilde{P}_{MX^nY^n} = \widetilde{P}_M \widetilde{P}_{X^n|M} \widetilde{P}_{Y^n|M}$ , i.e.,  $X^n \to M \to Y^n$  forms a Markov chain under  $\widetilde{P}$ . On the other hand,

$$H(\widetilde{P}_M) \le R \tag{416}$$

(since  $\widetilde{P}_M$  is defined on an alphabet with size  $e^{nR}$ ) and

$$D_{\infty}(\tilde{P}_{X^{n}Y^{n}} \| \pi_{XY}^{n}) = D_{\infty}(P_{X^{n}Y^{n}} | V=1 \| \pi_{XY}^{n})$$

$$= \log \sup_{\substack{(x^{n}, y^{n}) \in \\ \mathcal{A}_{\epsilon}^{(n)}(\pi_{X}) \times \mathcal{A}_{\epsilon}^{(n)}(\pi_{Y})}} \frac{P_{X^{n}Y^{n}}(x^{n}, y^{n})}{\pi_{XY}^{n}(x^{n}, y^{n})} - \log P_{V}(1)$$
(418)

$$\leq D_{\infty}(P_{X^{n}Y^{n}} \| \pi_{XY}^{n}) - \log P_{V}(1).$$
(419)

We now prove Lemma 2 by a argument similar as that in Appendix A. According to the definition of  $D_{\infty}$ ,  $D_{\infty}(P_{X^nY^n} \| \pi_{XY}^n) \leq \epsilon_n$  implies  $D_{\infty}(\widetilde{P}_{X^nY^n} \| \pi_{XY}^n) \leq \epsilon_n - \log P_V(1)$ , i.e.,

$$\sup_{x^{n},y^{n}} \frac{\tilde{P}_{X^{n}Y^{n}}(x^{n},y^{n})}{\pi_{XY}^{n}(x^{n},y^{n})} \le e^{\epsilon_{n} - \log P_{V}(1)} =: e^{\epsilon'_{n}}.$$
 (420)

Define  $\widehat{P}_{X^nY^n}(x^n, y^n) := \frac{e^{\epsilon'_n \pi^n_{XY}(x^n, y^n) - \widetilde{P}_{X^nY^n}(x^n, y^n)}{e^{\epsilon'_n - 1}}$ , then obviously  $\widehat{P}_{X^nY^n}(x^n, y^n)$  is a distribution. Hence  $\pi^n_{XY}$  can be written as a mixture distribution  $\pi^n_{XY}(x^n, y^n) = e^{-\epsilon'_n} \widetilde{P}_{X^nY^n}(x^n, y^n) + (1 - e^{-\epsilon'_n}) \widehat{P}_{X^nY^n}(x^n, y^n)$ . The encoder first generates a Bernoulli random variable U with  $P_U(1) = e^{-\epsilon'_n}$ , compresses it with 1 bit, and transmits it to the two generators. If U = 1, then the encoder and two generators use the synthesis codes  $(\widetilde{P}_M, \widetilde{P}_{X^n|M}, \widetilde{P}_{Y^n|M})$  with rate R (by fixed-length codes) to generate  $\widetilde{P}_{X^nY^n}$ . If U = 0, then the encoder generates  $(X^n, Y^n) \sim \widehat{P}_{X^nY^n}$ , and uses a variablelength compression code with rate

$$\frac{1}{n} \left( H\left(\widehat{P}_{X^{n}Y^{n}}\right) + 1 \right) \\
\leq \frac{1}{n} \left( H_{\widehat{P}}(V) + \widehat{P}_{V}(1) \log \left| \mathcal{A}_{\epsilon}^{(n)}\left(\pi_{X}\right) \times \mathcal{A}_{\epsilon}^{(n)}\left(\pi_{Y}\right) \right| \\
+ \widehat{P}_{V}(0) H\left(\widehat{P}_{X^{n}Y^{n}|V=0}\right) + 1 \right)$$
(421)

to generate  $\widehat{P}_{X^nY^n}$ . The distribution generated by such a mixed code is  $e^{-\epsilon'_n}\widetilde{P}_{X^nY^n}(x^n,y^n) + (1-e^{-\epsilon'_n})\widehat{P}_{X^nY^n}(x^n,y^n)$ , i.e.,  $\pi^n_{XY}(x^n,y^n)$ . The total code rate is no larger than

$$\frac{1}{n} + e^{-\epsilon'_n} R + \left(1 - e^{-\epsilon'_n}\right) \frac{1}{n} \left(H_{\widehat{P}}(V) + \widehat{P}_V(1) \log \left|\mathcal{A}_{\epsilon}^{(n)}(\pi_X) \times \mathcal{A}_{\epsilon}^{(n)}(\pi_Y)\right| + \widehat{P}_V(0) H\left(\widehat{P}_{X^n Y^n | V = 0}\right) + 1\right).$$
(422)

Observe that  $\pi_V(0) \to 0$ , and by the data processing inequality,  $P_V(0) \leq \pi_V(0)e^{\epsilon_n} \to 0$ . Hence  $\epsilon'_n = \epsilon_n - \log P_V(1) \to 0$  as  $n \to \infty$ . On the other hand, we have

$$H_{\widehat{P}}(V) \le \log 2 \tag{423}$$

$$\frac{1}{n}\log\left|\mathcal{A}_{\epsilon}^{(n)}\left(\pi_{X}\right)\times\mathcal{A}_{\epsilon}^{(n)}\left(\pi_{Y}\right)\right|\to H\left(\pi_{X}\right)+H\left(\pi_{Y}\right),\tag{424}$$

and

$$H\left(\widehat{P}_{X^{n}Y^{n}|V=0}\right) = H\left(\pi_{X^{n}Y^{n}|V=0}\right)$$

$$= \log \pi_{V}(0) - \frac{1}{\pi_{V}(0)}$$

$$\times \sum_{\substack{(x^{n},y^{n})\notin\\\mathcal{A}_{\epsilon}^{(n)}(\pi_{X})\times\mathcal{A}_{\epsilon}^{(n)}(\pi_{Y})}} \pi_{XY}^{n}(x^{n},y^{n}) \log \pi_{XY}^{n}(x^{n},y^{n})$$
(426)

$$= \log \pi_{V}(0) + \frac{1}{\pi_{V}(0)} \left( nH(\pi_{XY}) + \sum_{\substack{(x^{n}, y^{n}) \in \\ \mathcal{A}_{\epsilon}^{(n)}(\pi_{X}) \times \mathcal{A}_{\epsilon}^{(n)}(\pi_{Y})}} \pi_{XY}^{n}(x^{n}, y^{n}) \log \pi_{XY}^{n}(x^{n}, y^{n}) \right)$$
(427)

$$\leq \log \pi_V(0) + \frac{1}{\pi_V(0)} \left( nH(\pi_{XY}) - n \left( 1 - \epsilon \right) \left( H(\pi_{XY}) - \epsilon \right) \right)$$
(428)

$$= \frac{n}{\pi_V(0)} \left( \epsilon \left( H(\pi_{XY}) + 1 - \epsilon \right) + \frac{\pi_V(0) \log \pi_V(0)}{n} \right)$$
(429)

$$=\frac{n\left(\epsilon\left(H(\pi_{XY})+1-\epsilon\right)+o(1)\right)}{\pi_V(0)},\tag{430}$$

where (428) follows since  $\mathcal{A}_{\epsilon}^{(n)}(\pi_{XY}) \subseteq \mathcal{A}_{\epsilon}^{(n)}(\pi_X) \times \mathcal{A}_{\epsilon}^{(n)}(\pi_Y)$  and

$$\sum_{\substack{(x^n, y^n) \in \mathcal{A}_{\epsilon}^{(n)}(\pi_{XY}) \\ \leq -n \sum_{(x^n, y^n) \in \mathcal{A}_{\epsilon}^{(n)}(\pi_{XY})}} \pi_{XY}^n (x^n, y^n) (H(\pi_{XY}) - \epsilon)$$
(431)

$$n\pi_{XY}^{n}\left(\mathcal{A}_{\epsilon}^{(n)}\left(\pi_{XY}\right)\right)\left(H(\pi_{XY})-\epsilon\right)$$
(432)

$$\leq -n\left(1-\epsilon\right)\left(H(\pi_{XY})-\epsilon\right).\tag{433}$$

Here (431) follows by the definition of the  $\epsilon$ -weakly jointly typical set  $\mathcal{A}_{\epsilon}^{(n)}(\pi_{XY})$ , and (433) follows by [19, Theorem 3.1.2].

Hence to ensure (422) converges to R, we only require

$$\left(1 - e^{-\epsilon'_n}\right)\widehat{P}_V(0)\frac{\epsilon\left(H(\pi_{XY}) + 1 - \epsilon\right) + o(1)}{\pi_V(0)} \to 0.$$
(434)

According to the definitions of  $\hat{P}_{X^nY^n}$  and V, we know  $\hat{P}_V(0) = \frac{e^{\epsilon'_n}\pi_V(0)}{e^{\epsilon'_n}-1}$ . Hence

$$\left(1 - e^{-\epsilon'_n}\right) \widehat{P}_V(0) \frac{\epsilon \left(H(\pi_{XY}) + 1 - \epsilon\right) + o(1)}{\pi_V(0)}$$
  
=  $\pi_V(0) \frac{\epsilon \left(H(\pi_{XY}) + 1 - \epsilon\right) + o(1)}{\pi_V(0)}$  (435)

$$= \epsilon \left( H(\pi_{XY}) + 1 - \epsilon \right) + o(1) \to 0 \tag{436}$$

by letting  $n \to \infty$  first and letting  $\epsilon \to \infty$  then. This completes the proof.

#### APPENDIX H Proof of Lemma 3

# Some proof techniques used in this section are similar to

#### A. A Modified Version of $\infty$ -Rényi Code

those used in Appendix F.

By assumption, there exists a sequence of fixed-length  $\infty$ -Rényi codes with rate R such that  $D_{\infty}(P_{X^nY^n} || \pi_{XY}^n) \to 0$ . In this subsection, we construct another sequence of fixedlength  $\infty$ -Rényi codes by cascading the original  $\infty$ -Rényi codes with truncation, discretization, and adding noise. These new  $\infty$ -Rényi codes will be used to construct the final exact synthesis scheme in Appendix H-C. The original  $\infty$ -Rényi codes cannot be applied directly, since in the final exact synthesis scheme, we mix the  $\infty$ -Rényi codes and dyadic decomposition schemes [10]. The  $\infty$ -Rényi codes are used to generate an approximate distribution  $P_{X^nY^n}$  of  $\pi_{XY}^n$ . The dyadic decomposition schemes are used to generate the residual distribution after subtracting (a scaled version of)  $P_{X^nY^n}$ from  $\pi_{XY}^n$ . The dyadic decomposition schemes require the residual distribution to be log-concave. The original  $\infty$ -Rényi codes cannot generate a log-concave residual distribution. Hence it is necessary to construct new  $\infty$ -Rényi codes to ensure the residual distribution to be log-concave.

By respectively scaling X, Y, we can obtain a bivariate source with  $\mathbb{E}[X^2] = \mathbb{E}[Y^2] = 1$ . Hence without loss of

generality, we assume  $\pi_{XY}$  satisfying  $\mathbb{E}[X^2] = \mathbb{E}[Y^2] = 1$ . Define an *n*-ball with radius  $\sqrt{n(1+\epsilon)}$  as

$$\mathcal{B}_{\epsilon}^{(n)} := \left\{ x^n \in \mathbb{R}^n : \|x^n\| \le \sqrt{n\left(1+\epsilon\right)} \right\}.$$
(437)

Note that  $\mathcal{B}_{\epsilon}^{(n)}$  is a high probability set for any memoryless source with unit second moment, i.e.,  $\pi_X^n(\mathcal{B}_{\epsilon}^{(n)}), \pi_Y^n(\mathcal{B}_{\epsilon}^{(n)}) \rightarrow 1$ . Hence  $\pi_{XY}^n(\mathcal{B}_{\epsilon}^{(n)} \times \mathcal{B}_{\epsilon}^{(n)}) \rightarrow 1$ . Obviously,  $\mathcal{B}_{\epsilon}^{(n)}$  is contained in the *n*-cube  $\mathcal{L}_{\epsilon,n}^n$  with  $\mathcal{L}_{\epsilon,n}$  defined in (99). Hence  $\pi_{XY}^n(\mathcal{L}_{\epsilon,n}^{2n}) \rightarrow 1$ .

Assume  $\Delta_n$  is a decreasing positive sequence such that  $\Delta_n \to 0$  and  $n\Delta_n L_{\epsilon,n} \to 0$ . By Lemma 13, we have that for any  $(x, y), (\hat{x}, \hat{y}) \in \mathcal{L}^2_{\epsilon,n}$  satisfying  $|x - \hat{x}|, |y - \hat{y}| \leq \Delta_n$ ,

$$\frac{\pi_{XY}(x,y)}{\pi_{XY}(\hat{x},\hat{y})} \le \exp\left(\Delta_n L_{\epsilon,n}\right). \tag{438}$$

Hence for  $(x^n, y^n), (\hat{x}^n, \hat{y}^n) \in \mathcal{L}^n_{\epsilon,n} \times \mathcal{L}^n_{\epsilon,n}$ , satisfying  $|x_i - \hat{x}_i|, |y_i - \hat{y}_i| \leq \Delta_n, \forall i$ , we have

$$\frac{\pi_{XY}^n\left(x^n, y^n\right)}{\pi_{XY}^n\left(\hat{x}^n, \hat{y}^n\right)} \le \exp\left(n\Delta_n L_{\epsilon, n}\right). \tag{439}$$

Assume  $(P_M, P_{X^n|M}, P_{Y^n|M})$  is a sequence of fixedlength  $\infty$ -Rényi codes with rate R. That is,  $P_M$  is the uniform distribution on  $[1 : e^{nR}]$ , and this sequence of codes generates distributions  $P_{X^nY^n}$  such that  $\epsilon_n := D_{\infty}(P_{X^nY^n} || \pi_{XY}^n) \to 0$ . Similar to (415), we introduce a random variable

$$V := 1\left\{ (X^n, Y^n) \in \mathcal{L}^{2n}_{\epsilon, n} \right\}.$$

$$(440)$$

We define  $\widetilde{P}_{MX^nY^n} := P_{MX^nY^n|V}(m, x^n, y^n|1)$ . Then by Lemma 12,  $\widetilde{P}_{MX^nY^n} = \widetilde{P}_M \widetilde{P}_{X^n|M} \widetilde{P}_{Y^n|M}$ , i.e.,  $X^n \to M \to Y^n$  forms a Markov chain under  $\widetilde{P}$ . (416) and (419) still hold. Define  $[z]^n := \Delta_n \left\lfloor \frac{z^n}{\Delta_n} \right\rfloor$  as componentwise quantization operation of a vector  $z^n$  with step  $\Delta_n$  (for simplicity, we choose  $\Delta_n$  such that  $\sqrt{n(1+\epsilon)}$  is a multiple of  $\Delta_n$ ). Define  $U^n, V^n \sim \text{Unif}([0, \Delta_n]^n)$  are mutually independent, and also independent of  $[X]^n, [Y]^n$ . Then

$$\sup_{x^{n},y^{n}} \frac{\tilde{P}_{[X]^{n}+U^{n},[Y]^{n}+V^{n}}(x^{n},y^{n})}{\pi^{n}_{XY}(x^{n},y^{n})} \\
\leq \exp\left(n\Delta_{n}L_{\epsilon,n}\right) \sup_{x^{n},y^{n}} \frac{\tilde{P}_{[X]^{n}[Y]^{n}}\left(\left[x\right]^{n},\left[y\right]^{n}\right)/\Delta_{n}^{n}}{\pi^{n}_{XY}(\hat{x}^{n},\hat{y}^{n})} \quad (441)$$

$$= \exp\left(n\Delta_n L_{\epsilon,n}\right) \sup_{[x]^n, [y]^n} \frac{\widetilde{P}_{[X]^n[Y]^n}\left([x]^n, [y]^n\right)}{\pi^n_{[X][Y]}\left([x]^n, [y]^n\right)}$$
(442)

$$\leq \exp\left(n\Delta_n L_{\epsilon,n}\right) \sup_{x^n,y^n} \frac{\widetilde{P}_{X^n Y^n}\left(x^n,y^n\right)}{\pi_{XY}^n\left(x^n,y^n\right)} \tag{443}$$

$$\leq \exp\left(n\Delta_n L_{\epsilon,n} + D_{\infty}(P_{X^nY^n} \| \pi_{XY}^{\circ}) - \log P_V(1)\right)$$

$$= \exp\left(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n\right).$$
(445)

$$= \exp\left(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n\right), \tag{445}$$

where  $(\hat{x}^n, \hat{y}^n)$  in (441) is a point in  $([x]^n, [y]^n) + [0, \Delta_n]^{2n}$ such that  $\pi_{XY}^n(\hat{x}^n, \hat{y}^n) = \pi_{[X][Y]}^n([x]^n, [y]^n) / \Delta_n^n$  (the existence of such a point follows from the mean value theorem), (441) follows from (439), (443) follows from the data processing inequality, and (444) follows from (419). Define

$$\epsilon'_{n} := n\Delta_{n}L_{\epsilon,n} - \log P_{V}(1) + \epsilon_{n} + \delta_{n} \qquad (446)$$

for some positive sequence  $\delta_n \to 0$  as  $n \to \infty$ , which will be specified later. Then (445) implies for all  $(x^n, y^n) \in \mathcal{L}^{2n}_{\epsilon,n}$ ,

$$\frac{e^{\epsilon_n'}\pi_{XY}^n(x^n, y^n)}{\widetilde{P}_{[X]^n + U^n, [Y]^n + V^n}(x^n, y^n)} \ge e^{\delta_n}.$$
 (447)

Define

Obviously  $\hat{P}_{X^nY^n}(x^n, y^n)$  is a distribution. Then  $\pi_{XY}^n$  can be written as a mixture distribution

$$\pi_{XY}^{n}(x^{n}, y^{n}) = e^{-\epsilon'_{n}} \widetilde{P}_{[X]^{n} + U^{n}, [Y]^{n} + V^{n}}(x^{n}, y^{n}) + \left(1 - e^{-\epsilon'_{n}}\right) \widehat{P}_{X^{n}Y^{n}}(x^{n}, y^{n}).$$
(449)

Furthermore, by (447), we have

$$\widehat{P}_{X^{n}Y^{n}}(x^{n}, y^{n}) \geq \frac{e^{\delta_{n}} - 1}{e^{\epsilon_{n}'} - 1} \widetilde{P}_{[X]^{n} + U^{n}, [Y]^{n} + V^{n}}(x^{n}, y^{n})$$

$$e^{\delta_{n}} - 1 \widetilde{P}_{[X]^{n}[Y]^{n}}([x]^{n}, [y]^{n})$$
(450)
(451)

$$=\frac{e^{\delta_n}-1}{e^{\epsilon'_n}-1}\frac{P_{[X]^n[Y]^n}\left([x]^n,[y]^n\right)}{\Delta_n^2}.$$
 (451)

Now we partition the space  $\mathbb{R}^{2n}$  into a finite number of subregions so that we can apply dyadic decomposition schemes to each subregion. Specifically, partition the whole space  $\mathbb{R}^{2n}$  into  $3^{2n}$  subregions by 2n hyperplanes  $x_i = \pm \sqrt{n(1+\epsilon)}$  and  $y_i = \pm \sqrt{n(1+\epsilon)}, 1 \leq i \leq n$ . These subregions can be expressed as  $I_1 \times I_2 \times \ldots \times I_{2n}$ , where  $I_i \in \{\mathcal{L}_{\epsilon,n}^-, \mathcal{L}_{\epsilon,n}^-, \mathcal{L}_{\epsilon,n}^+\}, 1 \leq i \leq 2n \text{ with } \mathcal{L}_{\epsilon,n}^- :=$  $(-\infty, -\sqrt{n(1+\epsilon)})$  and  $\mathcal{L}_{\epsilon,n}^+ := (\sqrt{n(1+\epsilon)}, +\infty)$ . For brevity, we denote these subregions by  $\mathbb{R}_0, \mathbb{R}_1, \ldots, \mathbb{R}_{3^{2n}-1}$ , where  $\mathbb{R}_0 := \mathcal{L}_{\epsilon,n}^{2n}$  and  $\mathbb{R}_1, \mathbb{R}_2, \ldots, \mathbb{R}_{3^{2n}-1}$  denote the remaining subregions. For  $\mathbb{R}_k, 0 \leq k \leq 3^{2n} - 1$ , we use  $I_i^{(k)} \in \{\mathcal{L}_{\epsilon,n}^-, \mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}^+\}, 1 \leq i \leq 2n$  to denote the *i*th component of  $\mathbb{R}_k$ . That is,  $\mathbb{R}_k = I_1^{(k)} \times I_2^{(k)} \times \ldots \times I_{2n}^{(k)}$ . Furthermore, observe that  $\widetilde{P}_{[X]^n + U^n, [Y]^n + V^n}$  is supported on  $\mathbb{R}_0$ . Hence for  $1 \leq k \leq 3^{2n} - 1$ ,  $\widehat{P}(\cdot|\mathbb{R}_k) = \pi_{XY}^n(\cdot|\mathbb{R}_k)$ . This implies that  $\widehat{P}(x^n, y^n|\mathbb{R}_k) = \prod_{i=1}^n \pi_{XY}(x_i, y_i|I_i^{(k)} \times I_{n+i}^{(k)}),$ i.e.,  $(X_i, Y_i), 1 \leq i \leq n$  are i.i.d. under the distribution  $\widehat{P}(\cdot|\mathbb{R}_k)$  for  $1 \leq k \leq 3^{2n} - 1$ .

Next we derive upper bounds on  $T_{\text{Exact}}(\widehat{P}_{X^nY^n|(X^n,Y^n)\in\mathbb{R}_k})$  for  $0 \leq k \leq 3^{2n} - 1$ , by using dyadic decomposition schemes proposed in [10].

B. Dyadic Decomposition Schemes for  $\widehat{P}(\cdot|\mathbf{R}_k)$ ,  $0 \leq k \leq 3^{2n}-1$ 

We first consider k = 0. Denote  $I_{\Delta_n}^{2n}$  as a 2*n*-cube in  $\mathbb{R}_0$ 

$$I_{\Delta_n}^{2n} := ([x]^n, [y]^n) + [0, \Delta_n]^{2n} \subseteq \mathbf{R}_0$$
(452)

for  $\frac{[x]_i}{\Delta_n}, \frac{[y]_i}{\Delta_n} \in \left[-\frac{\sqrt{n(1+\epsilon)}}{\Delta_n}: \frac{\sqrt{n(1+\epsilon)}}{\Delta_n} - 1\right], 1 \le i \le n$ . By derivations similar to (367)-(378), we obtain that for a 2n-cube  $I_{\Delta_n}^{2n} \subseteq \mathbb{R}_0$  and for the distribution  $\widehat{P}_{X^nY^n}$ ,

$$I_{\widehat{P}}\left(X^{i}; X_{i+1}^{n}Y^{n} | (X^{n}, Y^{n}) \in I_{\Delta_{n}}^{2n}\right)$$

$$\leq \frac{4\left(n\Delta_{n}L_{\epsilon,n} - \log P_{V}(1) + \epsilon_{n}\right)}{\delta_{n}} + o\left(1\right).$$
(453)

for every  $1 \le i \le n$ , where o(1) denotes a term tending to zero as  $\epsilon'_n, \epsilon_n, n\Delta_n L_{\epsilon,n}, \delta_n \to 0$ .

By Lemma 14,  $\widehat{P}_{X^nY^n}\left(\cdot |I_{\Delta_n}^{2n}\right)$  is log-concave. On the other hand, for a log-concave distribution  $\pi_{Z^m}$ , the dyadic decomposition scheme in [10] realizes exactly generating  $Z^m$  in a distributed way (with  $Z_i$  realized at the *i*th terminal,  $1 \leq i \leq m$ ) as long as the rate of common randomness  $R \geq I^{(D)}(Z^m) + m^2 + 9 (\log 2) m \log m$  bits/symbol, where the dual total correlation

$$I^{(D)}(Z^{m}) := h(Z^{m}) - \sum_{i=1}^{m} h\left(Z_{i}|Z^{i-1}Z^{m}_{i+1}\right)$$
(454)
$$= \sum_{i=1}^{m} h\left(Z_{i}|Z^{i-1}\right) - \sum_{i=1}^{n} h\left(Z_{i}|Z^{i-1}Z^{m}_{i+1}\right)$$

<sup>*i*=1</sup> (455)

$$=\sum_{i=1}^{m} I\left(Z_{i}; Z_{i+1}^{m} | Z^{i-1}\right)$$
(456)

$$\leq \sum_{i=1}^{m} I\left(Z^{i}; Z^{m}_{i+1}\right).$$
(457)

That is, the exact common information  $T_{\text{Exact}}(\pi_{Z^m}) \leq I^{(D)}(Z^m) + m^2 + 9(\log 2) m \log m.$ 

Substituting  $\widehat{P}_{X^nY^n}\left(\cdot|I^{2n}_{\Delta_n}\right)$  into the dual total correlation, we have

$$I_{\widehat{P}}^{(D)}\left(X^{n}Y^{n}|\left(X^{n},Y^{n}\right)\in I_{\Delta_{n}}^{2n}\right) \leq \sum_{i=1}^{n} I_{\widehat{P}}\left(X^{i};X_{i+1}^{n}Y^{n}|\left(X^{n},Y^{n}\right)\in I_{\Delta_{n}}^{2n}\right) + \sum_{i=1}^{n} I_{\widehat{P}}\left(Y^{i};Y_{i+1}^{n}X^{n}|\left(X^{n},Y^{n}\right)\in I_{\Delta_{n}}^{2n}\right)$$
(458)

$$\leq 2n \left( \frac{4 \left( n \Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n \right)}{\delta_n} + o\left( 1 \right) \right).$$
 (459)

Now we consider the subregions  $R_k, 1 \le k \le 3^{2n}-1$ . Since  $(X_i, Y_i), 1 \le i \le n$  are i.i.d. under the distribution  $\widehat{P}(\cdot | R_k)$ , we have

$$I_{\widehat{P}}(X_{i};Y_{i}|(X^{n},Y^{n}) \in \mathbf{R}_{k}) = I_{\pi}\left(X_{i};Y_{i}|(X_{i},Y_{i}) \in I_{i}^{(k)} \times I_{n+i}^{(k)}\right).$$
(460)

By Lemma 15, we further have

$$I_{\widehat{P}}\left(X_{i};Y_{i}|\left(X^{n},Y^{n}\right)\in\mathbb{R}_{k}\right)\leq\Upsilon_{\pi}\left(I_{i}^{\left(k\right)},I_{n+i}^{\left(k\right)}\right).$$
(461)

For m = 2, the dyadic decomposition scheme in [10] realizes exactly generating  $Z^2$  in a distributed way as long as the rate  $R \ge I(Z_1; Z_2) + 24 \log 2$  nats/symbol. Applying this to the distribution  $\widehat{P}_{X_iY_i|(X^n,Y^n)\in\mathbb{R}_k}$ , we have that the exact common information

$$T_{\text{Exact}}(P_{X_i Y_i | (X^n, Y^n) \in \mathbf{R}_k}) \leq I_{\widehat{P}}(X_i; Y_i | (X^n, Y^n) \in \mathbf{R}_k) + 24 \log 2$$

$$(462)$$

$$\leq \Upsilon_{\pi} \left( I_{i}^{(k)}, I_{n+i}^{(k)} \right) + 24 \log 2.$$
(463)

Since  $\widehat{P}_{X^nY^n|(X^n,Y^n)\in\mathbb{R}_k}$  is a product distribution, we have for  $1 \le k \le 3^{2n} - 1$ ,

$$T_{\text{Exact}}(P_{X^{n}Y^{n}|(X^{n},Y^{n})\in\mathbb{R}_{k}}) \leq \frac{1}{n} \sum_{i=1}^{n} I_{\widehat{P}}(X_{i};Y_{i}|(X^{n},Y^{n})\in\mathbb{R}_{k}) + 24\log 2 \quad (464)$$

$$\leq \frac{1}{n} \sum_{i=1}^{n} \Upsilon_{\pi} \left( I_{i}^{(k)}, I_{n+i}^{(k)} \right) + 24 \log 2.$$
(465)

#### C. Exact Synthesis Scheme for $\pi_{XY}$

Now we construct an exact synthesis scheme for the distribution  $\pi_{XY}$ , which is similar as that in Appendix A. Our scheme is a mixture of the dyadic decomposition schemes above and the modified fixed-length  $\infty$ -Rényi code  $\left(\widetilde{P}_{M}, \widetilde{P}_{X^{n}|M}, \widetilde{P}_{Y^{n}|M}\right)$  constructed in Subsection H-A. The encoder first generates a Bernoulli random variable U with  $P_U(1) = e^{-\epsilon'_n}$ , compresses it with 1 bit, and transmits it to the two generators. If U = 1, then the encoder and two generators use the modified  $\infty$ -Rényi code  $\left(\widetilde{P}_{M}, \widetilde{P}_{X^{n}|M}, \widetilde{P}_{Y^{n}|M}\right)$  with rate R to generate  $\widetilde{P}_{X^{n}Y^{n}}$ . Then by quantizing  $(X^{n}, Y^{n})$  and adding uniform random variables to them, the generators obtain  $P_{[X]^n+U^n,[Y]^n+V^n}$ . If U = 0, then the encoder generates  $(X^n, Y^n) \sim \hat{P}_{X^n Y^n}$ , uses  $\frac{1}{n} \log \left(3^{2n}\right) + \frac{1}{n} \log \left(\frac{\sqrt{n(1+\epsilon)}}{\Delta_n}\right)^n$  rate to encode the index of the subregion  $R_k$  and the 2*n*-cube (if  $(X^n, Y^n) \in R_0$ ) that  $(X^n, Y^n)$  belongs to, and uses the dyadic decomposition scheme in [10] to generate  $\hat{P}_{X^nY^n|(X^n,Y^n)\in I_{\Delta_n}^{2n}}$  with rate  $I_{\widehat{P}}^{(D)}(X^nY^n|(X^n,Y^n)\in I_{\Delta_n}^{2n}) + 4n^2 + 18(\log 2) n \log(2n)$ if  $(X^n,Y^n)$  belongs to some 2n-cube  $I_{\Delta_n}^{2n} \subseteq \mathbf{R}_0$ ; to generate  $\widehat{P}_{X^nY^n|(X^n,Y^n)\in\mathbb{R}_k}$  with rate  $T_{\mathrm{Exact}}(\widehat{P}_{X^nY^n|(X^n,Y^n)\in\mathbb{R}_k})$ if  $(X^n,Y^n)$  belongs to some subregion  $\mathbb{R}_k$  for  $1 \leq k \leq 3^{2n} - 1$ . The distribution generated by such a mixed code is  $e^{-\epsilon'_n} \widetilde{P}_{[X]^n+U^n,[Y]^n+V^n}(x^n,y^n)$  +  $\left(1-e^{-\epsilon'_n}\right)\widehat{P}_{X^nY^n}\left(x^n,y^n\right)$ , i.e.,  $\pi^n_{XY}\left(x^n,y^n\right)$ . The total code rate is no larger than (475)-(476) (given on page 37), where the sum  $\sum_{I_{\Delta n}} \subseteq \mathbb{R}_0$  is taken over all 2*n*-cubes  $I_{\Delta_n}^{2n} \subseteq \mathbb{R}_0$  (see (452)); (475) follows from (459), (465), and the fact that  $\pi_{XY}^{n}(\mathbf{R}_{k}) = (1 - e^{-\epsilon_{n}'}) \widehat{P}_{X^{n}Y^{n}}(\mathbf{R}_{k})$ (since  $\pi_{XY}^{n}(x^{n}, y^{n}) = (1 - e^{-\epsilon_{n}'}) \widehat{P}_{X^{n}Y^{n}}(x^{n}, y^{n})$  for  $(x^n, y^n) \notin \mathcal{L}^{2n}_{\epsilon,n}$ ; see (448)); and (476) follows since on one hand,

$$\sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n}(\mathbf{R}_{k}) \left\{ \frac{1}{n} \sum_{i=1}^{n} \Upsilon_{\pi} \left( I_{i}^{(k)}, I_{n+i}^{(k)} \right) + 24 \log 2 \right\}$$
  
= 
$$\sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n}(\mathbf{R}_{k}) \left\{ \Upsilon_{\pi} \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) + 24 \log 2 \right\}$$
(466)  
= 
$$\sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n}(\mathbf{R}_{k})$$

$$\times \left(\Upsilon_{\pi}\left(I_{1}^{(k)}, I_{n+1}^{(k)}\right) \mathbb{1}\left\{\left(I_{1}^{(k)}, I_{n+1}^{(k)}\right) \neq \left(\mathcal{L}_{\epsilon, n}, \mathcal{L}_{\epsilon, n}\right)\right\}$$

+ 
$$\Upsilon_{\pi} \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) 1 \left\{ \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) = (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}) \right\}$$
  
+ 24 log 2 (467)

$$\leq \sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n} (\mathbf{R}_{k}) \\ \times \left( \Upsilon_{\pi} \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) \mathbf{1} \left\{ \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) \neq (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}) \right\} \\ + \frac{I_{\pi} (X; Y) + o(1)}{\pi_{XY} (\mathcal{L}_{\epsilon,n} \times \mathcal{L}_{\epsilon,n})} + 24 \log 2 \right)$$
(468)
$$= \sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n} (\mathbf{R}_{k}) \Upsilon_{\pi} \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right)$$

$$\times 1\left\{ \left(I_1^{(k)}, I_{n+1}^{(k)}\right) \neq \left(\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}\right) \right\} + o(1)$$
(469)

and on the other hand,

$$\sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n} (\mathbf{R}_{k}) \Upsilon_{\pi} \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) \times 1 \left\{ \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) \neq (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}) \right\}$$

$$= \sum_{k=0}^{3^{2n}-1} \pi_{XY}^{n} (\mathbf{R}_{k}) \Upsilon_{\pi} \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) \times 1 \left\{ \left( I_{1}^{(k)}, I_{n+1}^{(k)} \right) \neq (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}) \right\}$$

$$= \sum_{x^{n}, y^{n}} \pi_{XY}^{n} (x^{n}, y^{n}) \sum_{I^{2n} \in \left\{ \mathcal{L}_{\epsilon,n}^{-}, \mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}^{+} \right\}^{2n}} 1 \left\{ (x^{n}, y^{n}) \in I^{2n} \right\} \times \Upsilon_{\pi} (I_{1}, I_{n+1}) 1 \left\{ (I_{1}, I_{n+1}) \neq (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}) \right\}$$

$$= \sum_{x_{1}, y_{1}} \pi_{XY} (x_{1}, y_{1}) \sum_{I_{1}, I_{n+1} \in \left\{ \mathcal{L}_{\epsilon,n}^{-}, \mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}^{+} \right\}, \\ (I_{1}, I_{n+1}) \neq (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n})$$

$$= \sum_{x_{1}, y_{1}} \pi_{XY} (x_{1}, y_{1}) (472)$$

$$= \sum_{\substack{I_1, I_{n+1} \in \left\{\mathcal{L}_{\epsilon,n}^-, \mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}^+\right\}, \\ (I_1, I_{n+1}) \neq (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n})}} \pi_{XY} \left(I_1 \times I_{n+1}\right) \Upsilon_{\pi} \left(I_1, I_{n+1}\right)$$
(473)

$$= o(1).$$
 (474)

Here (466) follows by symmetry:  $\Upsilon_{\pi}\left(I_{1}^{(k)}, I_{n+1}^{(k)}\right) =$  $\Upsilon_{\pi}\left(I_{i}^{(k)}, I_{n+i}^{(k)}\right)$  for all *i*; (469) follows since  $\sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n}(\mathbf{R}_{k}) = 1 - \pi_{XY}^{n}(\mathbf{R}_{0}) = o(1)$ ; and (470) follows since for  $\mathbf{R}_{0}, \left(I_{1}^{(k)}, I_{n+1}^{(k)}\right) = (\mathcal{L}_{\epsilon,n}, \mathcal{L}_{\epsilon,n}).$ 

Observe that  $\pi_V(0) \to 0$  exponentially fast, and by the data processing inequality,  $P_V(0) \leq \pi_V(0)e^{\epsilon_n} \to 0$  exponentially fast. Hence if  $n\Delta_n L_{\epsilon,n}, \delta_n \to 0$ , then  $\epsilon'_n$  (defined in (446)) satisfies  $\epsilon'_n \to 0$  as  $n \to \infty$ . On the other hand,

$$H_{\widehat{P}}(V) \le \log 2. \tag{477}$$

Hence to ensure (476) converges to R, we only require

$$n\Delta_n L_{\epsilon,n}, \delta_n \to 0 \tag{478}$$

and

$$\left(1 - e^{-\epsilon'_n}\right) \left(\log\left(\frac{\sqrt{n\left(1+\epsilon\right)}}{\Delta_n}\right) + \frac{8\left(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n\right)}{\delta_n} + 4n\right) \to 0.$$
(479)

Note that (479) is equivalent to (480) (given on page 37).

Choose  $\delta_n = n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n$ , then we only require

$$(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n) (4n - \log \Delta_n) \to 0.$$
 (481)

Observe that  $\pi_V(0) \to 0$  exponentially fast, and by the data processing inequality,  $-\log P_V(1) = -\log (1 - P_V(0)) \sim P_V(0) \to 0$  exponentially fast. Choose  $\Delta_n = \frac{1}{(nL_{\epsilon,n})^3}$ , then  $n\Delta_n L_{\epsilon,n} \to 0$  and

$$\left(\frac{1}{\left(nL_{\epsilon,n}\right)^{2}} - \log P_{V}(1) + \epsilon_{n}\right) \left(4n + 3\log n + 3\log L_{\epsilon,n}\right)$$

$$\sim \left(\frac{1}{\left(nL_{\epsilon,n}\right)^{2}} + P_{V}(0) + \epsilon_{n}\right) \left(4n + \log L_{\epsilon,n}\right)$$
(482)

$$\sim \left(\frac{1}{\left(nL_{\epsilon,n}\right)^2} + P_V(0) + \epsilon_n\right) \left(4n + \log L_{\epsilon,n}\right) \tag{482}$$

$$= (P_V(0) + \epsilon_n) (4n + \log L_{\epsilon,n}) + o(1)$$
(483)

$$= P_V(0) \log L_{\epsilon,n} + \epsilon_n \left(4n + \log L_{\epsilon,n}\right) + o(1).$$
(484)

Hence we only require

$$P_V(0)\log L_{\epsilon,n} \to 0 \tag{485}$$

$$\epsilon_n \left( n + \log L_{\epsilon,n} \right) \to 0. \tag{486}$$

That is,  $\epsilon_n = o\left(\frac{1}{n + \log L_{\epsilon,n}}\right)$  and  $\log L_{\epsilon,n}$  is sub-exponentially growing in n. These are the assumptions given in the lemma. Hence the proof is complete.

#### APPENDIX I Proof of Theorem 8

In this section, we extend the proof in Appendix A-B to the Gaussian case by combining it with discretization techniques.

Define  $Q_W = \mathcal{N}(0,\rho), Q_{X|W}(\cdot|w) = \mathcal{N}(w, 1-\rho), Q_{Y|W}(\cdot|w) = \mathcal{N}(w, 1-\rho)$ . Then  $Q_{XY} = \pi_{XY}$ . For  $\epsilon > 0$ , we define the distributions

$$P_{W^{n}}\left(w^{n}\right) \propto Q_{W}^{n}\left(w^{n}\right) \mathbb{1}\left\{w^{n} \in \mathcal{A}_{\frac{\epsilon}{2}}^{\left(n\right)}\left(Q_{W}\right)\right\},$$
(487)

$$P_{X^{n}|W^{n}}(x^{n}|w^{n}) \propto Q_{X|W}^{n}(x^{n}|w^{n}) \\ \times 1\left\{x^{n} \in \mathcal{A}_{\epsilon}^{(n)}(Q_{WX}|w^{n})\right\}, (488) \\ P_{Y^{n}|W^{n}}(y^{n}|w^{n}) \propto Q_{Y|W}^{n}(y^{n}|w^{n}) \\ \times 1\left\{y^{n} \in \mathcal{A}_{\epsilon}^{(n)}(Q_{WY}|w^{n})\right\}. (489)$$

According to the definition of weakly typical sets,

$$\mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_W) = \left\{ w^n \in \mathbb{R}^n : \left| \frac{\|w^n\|^2}{n\rho} - 1 \right| \le \epsilon \right\}$$
(490)

$$\frac{1}{n} + e^{-\epsilon'_{n}}R + (1 - e^{-\epsilon'_{n}}) \left\{ \frac{1}{n} \log \left( 3^{2n} \right) + \frac{1}{n} \log \left( \frac{\sqrt{n(1+\epsilon)}}{\Delta_{n}} \right)^{n} + \sum_{\substack{I_{\Delta_{n}}^{2n} \subseteq \mathbf{R}_{0}}} \widehat{P}_{X^{n}Y^{n}} \left( I_{\Delta_{n}}^{2n} \right) \left( I_{\widehat{P}}^{(D)} \left( X^{n}Y^{n} \right) \left( X^{n}, Y^{n} \right) \in I_{\Delta_{n}}^{2n} \right) + 4n + 18 \left( \log 2 \right) \log \left( 2n \right) \right) + \sum_{k=1}^{3^{2n}-1} \widehat{P}_{X^{n}Y^{n}} \left( \mathbf{R}_{k} \right) T_{\text{Exact}} \left( \widehat{P}_{X^{n}Y^{n} \mid (X^{n}, Y^{n}) \in \mathbf{R}_{k}} \right) \right\}$$

$$\leq \frac{1}{n} + e^{-\epsilon'_{n}}R + (1 - e^{-\epsilon'_{n}}) \left\{ \log \left( \frac{9\sqrt{n(1+\epsilon)}}{\Delta_{n}} \right) + \frac{8 \left( n\Delta_{n}L_{\epsilon,n} - \log P_{V}(1) + \epsilon_{n} \right)}{\delta_{n}} + o\left( 1 \right) + 4n + 18 \left( \log 2 \right) \log \left( 2n \right) \right\}$$

$$+ \sum_{k=1}^{3^{2n}-1} \pi_{XY}^{n} \left( \mathbf{R}_{k} \right) \left\{ \frac{1}{n} \sum_{i=1}^{n} \Upsilon_{\pi} \left( I_{i}^{(k)}, I_{n+i}^{(k)} \right) + 24 \log 2 \right\}$$

$$\sim \frac{1}{n} + e^{-\epsilon'_{n}}R + (1 - e^{-\epsilon'_{n}}) \left\{ \log \left( \frac{\sqrt{n(1+\epsilon)}}{\Delta_{n}} \right) + \frac{8 \left( n\Delta_{n}L_{\epsilon,n} - \log P_{V}(1) + \epsilon_{n} \right)}{\delta_{n}} + 4n \right\} + o\left( 1 \right).$$

$$(476)$$

$$\left(1 - e^{-(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n + \delta_n)}\right) \left(\log\left(\frac{\sqrt{n(1+\epsilon)}}{\Delta_n}\right) + \frac{8\left(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n\right)}{\delta_n} + 4n\right) \\ \sim \left(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n + \delta_n\right) \left(4n - \log\Delta_n + \frac{8\left(n\Delta_n L_{\epsilon,n} - \log P_V(1) + \epsilon_n\right)}{\delta_n}\right) \to 0.$$
(480)

and

$$\mathcal{A}_{\epsilon}^{(n)}(Q_{WX}) = \mathcal{A}_{\epsilon}^{(n)}(Q_{WY})$$

$$= \left\{ (w^{n}, x^{n}) \in \mathbb{R}^{2n} : \left| \frac{\|w^{n}\|^{2}}{n\rho} - 1 \right| \leq 2\epsilon \\ \left| \frac{\|w^{n}\|^{2}}{n\rho} + \frac{\|x^{n} - w^{n}\|^{2}}{n(1-\rho)} - 2 \right| \leq 2\epsilon \right\}.$$
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(491)
(4

Hence for  $(w^n, x^n) \in \mathcal{A}_{\epsilon}^{(n)}(Q_{WX}),$ 

$$\left|\frac{\|x^{n} - w^{n}\|^{2}}{n(1-\rho)} - 1\right| \le 4\epsilon$$
(493)

and

$$\left| \frac{1}{n} (x^n - w^n)^\top w^n \right|$$
  
=  $\left| \frac{1}{2n} \left( \|w^n\|^2 + \|x^n - w^n\|^2 - \|x^n\|^2 \right) \right|$  (494)

$$\leq \frac{1}{2} \left| \rho \left( 1 + 2\epsilon \right) + (1 - \rho) \left( 1 + 4\epsilon \right) - (1 - 2\epsilon) \right|$$
(495)

$$= (3 - \rho) \epsilon. \tag{496}$$

Define

$$\delta_{0,n} := 1 - Q_W^n \left( \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_W) \right)$$

$$\delta_{1,n} := 1 - \inf_{\substack{w^n \in \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_W)}} Q_{X|W}^n \left( \mathcal{A}_{\epsilon}^{(n)}(Q_{WX}|w^n) \, | w^n \right)$$
(497)
(497)
(497)
(497)

$$\delta_{2,n} := 1 - \inf_{w^n \in \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_W)} Q_{Y|W}^n \left( \mathcal{A}_{\epsilon}^{(n)} \left( Q_{WY} | w^n \right) | w^n \right).$$
(499)

Then  $\delta_{0,n}, \delta_{1,n}, \delta_{2,n} \to 0$  exponentially fast, as shown in the following lemma.

**Lemma 16** (Gaussian Typicality Lemma).  $\delta_{0,n}, \delta_{1,n}, \delta_{2,n} \to 0$ exponentially fast.

Proof of Lemma 16: By large deviation theory, we know that  $\delta_{0,n} \rightarrow 0$  exponentially fast. Next we prove  $\delta_{1,n} \rightarrow 0$ exponentially fast. (That  $\delta_{2,n} \rightarrow 0$  exponentially fast follows by symmetry.)

Under the condition  $w^n \in \mathcal{A}^{(n)}_{\frac{\epsilon}{2}}(Q_W)$ ,

$$\left|\frac{\left\|w^{n}\right\|^{2}}{n\rho} - 1\right| \le \epsilon \tag{500}$$

is satisfied automatically. Denote  $Z^n = X^n - w^n$ . Then  $Z_i$ 's are i.i.d., and  $Z_i \sim Q_Z = \mathcal{N}(0, 1 - \rho)$ . By large deviation theory,

$$\mathbb{P}_{Z^n \sim Q_Z^n} \left( \left| \frac{\|w^n\|^2}{n\rho} + \frac{\|Z^n\|^2}{n(1-\rho)} - 2 \right| \le 2\epsilon \right) \to 1 \quad (501)$$

exponentially fast. Now we consider the condition  $\left|\frac{\|X^n\|^2}{n} - 1\right| \le 2\epsilon$ , which is equivalent to  $\left|\frac{\|w^n + Z^n\|^2}{n} - 1\right| \le 2\epsilon$ . Observe that

$$\|w^{n} + Z^{n}\|^{2} = \|w^{n}\|^{2} + \|Z^{n}\|^{2} + 2\sum_{i=1}^{n} w_{i}Z_{i}.$$
 (502)

By the large deviation theory, for  $\epsilon' > 0$ ,

$$\mathbb{P}_{Z^n \sim Q_Z^n} \left( \left| \frac{\|Z^n\|^2}{n \left( 1 - \rho \right)} - 1 \right| \le \epsilon' \right) \to 1$$
 (503)

exponentially fast. On the other hand, observe that  $\frac{1}{n} \sum_{i=1}^{n} w_i Z_i \sim \mathcal{N}(0, \frac{1}{n^2} \|w^n\|^2 (1-\rho))$ . Hence

$$\mathbb{P}_{Z^n \sim Q_Z^n} \left( \left| \frac{1}{n} \sum_{i=1}^n w_i Z_i \right| \le \epsilon' \right) = 1 - 2 \mathsf{Q} \left( \frac{n\epsilon'}{\|w^n\| \sqrt{1-\rho}} \right),\tag{504}$$

where Q is the Q-function for the standard normal distribution. Since  $Q(x) \le e^{-\frac{x^2}{2}}$ , x > 0, we have

$$\mathbb{P}_{Z^n \sim Q_Z^n} \left( \left| \frac{1}{n} \sum_{i=1}^n w_i Z_i \right| \le \epsilon' \right) \\\ge 1 - 2 \exp\left( -\frac{1}{2} \left( \frac{n\epsilon'}{\|w^n\| \sqrt{1-\rho}} \right)^2 \right)$$
(505)

$$\geq 1 - 2\exp\left(-\frac{1}{2}\frac{n\epsilon'^2}{\rho\left(1-\epsilon\right)\left(1-\rho\right)}\right) \tag{506}$$

$$\rightarrow 1$$
 (507)

exponentially fast. Hence  $\inf_{w^n \in \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_W)} Q_Z^n(\mathcal{B}_{\epsilon'}(w^n)) \to 1$ exponentially fast, where

$$\mathcal{B}_{\epsilon'}(w^n) := \left\{ z^n \in \mathbb{R}^n : \begin{array}{c} \left| \frac{\|z^n\|^2}{n(1-\rho)} - 1 \right| \le \epsilon' \\ \left| \frac{1}{n} \sum_{i=1}^n w_i z_i \right| \le \epsilon' \end{array} \right\}.$$
(508)

Now we claim that for sufficiently small  $\epsilon'$ , if  $w^n \in \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_W)$  and  $z^n = x^n - w^n \in \mathcal{B}_{\epsilon'}(w^n)$ , then  $(w^n, x^n) \in \mathcal{A}_{\epsilon}^{(n)}(Q_{WX})$ . Since  $\inf_{w^n \in \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_W)} Q_Z^n(\mathcal{B}_{\epsilon'}(w^n)) \to 1$  exponentially fast, this claim implies that  $\delta_{1,n} \to 0$  exponentially fast as well. Hence the rest is to prove this claim.

Observe that for  $w^n \in \mathcal{A}^{(n)}_{\frac{\epsilon}{2}}(Q_W)$  and  $z^n = x^n - w^n \in \mathcal{B}_{\epsilon'}(w^n)$ , we have

$$\frac{\left\|\frac{\|x^{n}\|^{2}}{n} - 1\right| }{\left|\frac{\|w^{n} + z^{n}\|^{2}}{n} - 1\right|$$
(509)

$$= \left| \frac{\|w^n\|^2 + \|z^n\|^2 + 2\sum_{i=1}^n w_i z_i}{n} - 1 \right|$$
(510)

$$\leq \max \left\{ \left| \rho \left( 1 + \epsilon \right) + \left( 1 - \rho \right) \left( 1 + \epsilon' \right) + 2\epsilon' - 1 \right|, \\ \left| \rho \left( 1 - \epsilon \right) + \left( 1 - \rho \right) \left( 1 - \epsilon' \right) - 2\epsilon' - 1 \right| \right\}$$
(511)

$$= \rho \epsilon + (1 - \rho) \epsilon' + 2\epsilon', \tag{512}$$

and

$$\left| \frac{\|w^n\|^2}{n\rho} + \frac{\|x^n - w^n\|^2}{n(1-\rho)} - 2 \right| \\
= \left| \frac{\|w^n\|^2}{n\rho} + \frac{\|z^n\|^2}{n(1-\rho)} - 2 \right| \quad (513) \\
\leq \epsilon + \epsilon'. \quad (514)$$

Now we choose

$$\epsilon' \le \min\left\{\epsilon, \frac{(2-\rho)\epsilon}{3-\rho}\right\} = \frac{(2-\rho)\epsilon}{3-\rho}$$

then  $\left|\frac{\|x^n\|^2}{n} - 1\right| \leq 2\epsilon$  and  $\left|\frac{\|w^n\|^2}{n\rho} + \frac{\|x^n - w^n\|^2}{n(1-\rho)} - 2\right| \leq 2\epsilon$ . Hence we complete the proof of the claim above.

We set  $C_n = \{W^n(m)\}_{m \in \mathcal{M}_n}$  with  $W^n(m), m \in \mathcal{M}_n$ drawn independently for different *m*'s and according to the same distribution  $P_{W^n}$  such that  $P_{W^n}$ . Upon receiving  $W^n(M)$ , the two generators respectively use random mappings  $P_{X^n|W^n}$  and  $P_{Y^n|W^n}$  to generate  $X^n$  and  $Y^n$ . For a sequence of positive numbers  $\{\Delta_n\}$ , we quantize  $X^n$  and  $Y^n$ as  $[X]^n = \Delta_n \lfloor \frac{X^n}{\Delta_n} \rfloor$  and  $[Y]^n = \Delta_n \lfloor \frac{Y^n}{\Delta_n} \rfloor$ . Define  $\left[\mathcal{A}_{\epsilon}^{(n)}\right] \times \left[\mathcal{A}_{\epsilon}^{(n)}\right] := \left(\Delta \mathbb{Z}^n \cap \mathcal{A}_{\epsilon}^{(n)}(\pi_X)\right) \times \left(\Delta \mathbb{Z}^n \cap \mathcal{A}_{\epsilon}^{(n)}(\pi_Y)\right)$ . Define  $U^n, V^n \sim \text{Unif } (I^n_{\Delta_n})$  with  $I^n_{\Delta_n} = [0, \Delta_n]^n$  are mutually independent, and also independent of  $[X]^n, [Y]^n$ . For such a code, we have the following Gaussian version of distributed Rényi-covering lemma.

Lemma 17 (Distributed Gaussian Rényi-Covering). For the random code described above, if

$$R > \frac{1}{2} \log\left[\frac{1+\rho}{1-\rho}\right] + \frac{\rho}{1+\rho},\tag{515}$$

then there exists some  $\alpha, \epsilon > 0$  and some positive sequence  $\{\Delta_n\}$  such that

$$\mathbb{P}_{\mathcal{C}_n}\left(D_{\infty}(P_{[X]^n+U^n,[Y]^n+V^n|\mathcal{C}_n}\|\pi_{XY}^n) \le e^{-n\alpha}\right) \to 1$$
(516)

doubly exponentially fast.

This lemma implies that there exists a sequence of codebooks  $\{c_n\}$  with rate R such that  $D_{\infty}(P_{[X]^n+U^n,[Y]^n+V^n|\mathcal{C}_n=c_n} \| \pi_{XY}^n) \leq e^{-n\alpha}$  as long as  $R > \frac{1}{2} \log \left[ \frac{1+\rho}{1-\rho} \right] + \frac{\rho}{1+\rho}$ . This completes the proof of  $T_{\infty}(\pi_{XY}) \leq \frac{1}{2} \log \left[ \frac{1+\rho}{1-\rho} \right] + \frac{\rho}{1+\rho}$ . Hence what we need to do is to prove Lemma 17. The proof is provided in the following. *Proof of Lemma 17:* Assume  $\epsilon > 0$  is a number such

that

$$R > (1+\epsilon) \left(\frac{1}{2}\log\left[\frac{1+\rho}{1-\rho}\right] + \frac{\rho}{1+\rho}\right) + 3\epsilon.$$
 (517)

For brevity, in the following we denote  $M = e^{nR}$ . According to the definition of the Rényi divergence, we have

$$e^{D_{\infty}(T[X]^{n}[Y]^{n}\|[n_{[X]}[Y])} = \sup_{(x^{n},y^{n})\in [\mathcal{A}_{\epsilon}^{(n)}] \times [\mathcal{A}_{\epsilon}^{(n)}]} \frac{P_{[X]^{n}[Y]^{n}}(x^{n},y^{n})}{\pi^{n}_{[X][Y]}(x^{n},y^{n})}$$
(518)

$$= \sup_{(x^n, y^n) \in \left[\mathcal{A}_{\epsilon}^{(n)}\right] \times \left[\mathcal{A}_{\epsilon}^{(n)}\right]} \widetilde{g}_{[X]^n [Y]^n | \mathcal{C}_n}(x^n, y^n | \mathcal{C}_n), \quad (519)$$

where

מ) ת

 $|| \_ n$ 

$$\widetilde{g}_{[X]^{n}[Y]^{n}|\mathcal{C}_{n}}(x^{n}, y^{n}|\mathcal{C}_{n})$$

$$:= \sum_{m \in \mathcal{M}_{n}} \frac{g_{[X]^{n}[Y]^{n}|W^{n}}(x^{n}, y^{n}|W^{n}(m))}{\mathsf{M}}$$
(520)

with

$$g_{[X]^{n}[Y]^{n}|W^{n}}(x^{n}, y^{n}|w^{n})$$
  
:= 
$$\frac{P_{[X]^{n}|W^{n}}(x^{n}|w^{n})P_{[Y]^{n}|W^{n}}(y^{n}|w^{n})}{\pi^{n}_{[X][Y]}(x^{n}, y^{n})}.$$
(521)

By the data processing inequality,

$$\sup_{\substack{(x^n,y^n)\in\left[\mathcal{A}_{\epsilon}^{(n)}\right]\times\left[\mathcal{A}_{\epsilon}^{(n)}\right]\\\leq \sup_{(x^n,y^n)\in\mathcal{A}_{\epsilon}^{(n)}\times\mathcal{A}_{\epsilon}^{(n)}}g_{X^nY^n|W^n}(x^n,y^n|w^n).}$$
(522)

On the other hand, define

$$\mathcal{A} := \left\{ (x^n, y^n) : \exists w^n \text{ s.t. } (w^n, x^n) \in \mathcal{A}_{\epsilon}^{(n)}(Q_{WX}), \\ (w^n, y^n) \in \mathcal{A}_{\epsilon}^{(n)}(Q_{WY}) \right\}$$
(523)

and

$$\delta_{12,n} := \frac{1}{(1 - \delta_{1,n}) (1 - \delta_{2,n})}.$$
(524)

Since Lemma 16 shows that  $\delta_{1,n}, \delta_{2,n} \to 0$  exponentially fast, we know that  $\delta_{12,n} \rightarrow 1$  exponentially fast. Then similar to (171), we can show that for  $w^n \in \mathcal{A}^{(n)}_{\frac{\epsilon}{2}}(Q_W)$ , (526)-(532) (given on page 40) hold, where (528) follows from (496). Combining (522) and (532), we obtain

$$\sup_{(x^n,y^n)\in \left[\mathcal{A}_{\epsilon}^{(n)}\right]\times \left[\mathcal{A}_{\epsilon}^{(n)}\right]}g_{[X]^n[Y]^n|W^n}(x^n,y^n|w^n) \leq \beta_n.$$
 (525)

Continuing (519), we get for any  $\epsilon' > 0$ ,

$$\mathbb{P}_{\mathcal{C}_{n}}\left(e^{D_{\infty}(P_{[X]^{n}[Y]^{n}} \|\pi_{[X][Y]}^{n})} \geq 1 + \epsilon'\right) \\
= \mathbb{P}_{\mathcal{C}_{n}}\left(\sup_{(x^{n}, y^{n}) \in \left[\mathcal{A}_{\epsilon}^{(n)}\right] \times \left[\mathcal{A}_{\epsilon}^{(n)}\right]} \widetilde{g}_{[X]^{n}[Y]^{n} | \mathcal{C}_{n}}(x^{n}, y^{n} | \mathcal{C}_{n})\right) \\
\geq 1 + \epsilon'\right) \qquad (533) \\
\leq \left|\left[\mathcal{A}_{\epsilon}^{(n)}\right] \times \left[\mathcal{A}_{\epsilon}^{(n)}\right]\right| \sup_{(x^{n}, y^{n}) \in \left[\mathcal{A}_{\epsilon}^{(n)}\right] \times \left[\mathcal{A}_{\epsilon}^{(n)}\right]} \\
\mathbb{P}_{\mathcal{C}_{n}}\left(\widetilde{g}_{[X]^{n}[Y]^{n} | \mathcal{C}_{n}}(x^{n}, y^{n} | \mathcal{C}_{n}) \geq 1 + \epsilon'\right), \qquad (534)$$

where (534) follows from the union bound. If the probability in (534) vanishes doubly exponentially fast and  $\left| \begin{bmatrix} \mathcal{A}_{\epsilon}^{(n)} \end{bmatrix} \times \begin{bmatrix} \mathcal{A}_{\epsilon}^{(n)} \end{bmatrix} \right|$  is growing much slower, then  $\max_{x^n,y^n} \widetilde{g}(x^n,y^n|\mathcal{C}_n) < 1 + \epsilon'$  with high probability as  $n \rightarrow \infty$ . To this end, we use similar techniques used in Appendix A-B to bound the probability. Define  $I_{\Delta}^n := [0, \Delta]^n$ . Observe that  $g(x^n, y^n | W^n(m)), m \in \mathcal{M}_n$  are i.i.d. random variables with mean  $\mu_{\epsilon,n}$  given in (537)-(541) (given on page 40) and variance

$$\begin{aligned} &\operatorname{Var}_{W^{n}}\left[g_{[X]^{n}[Y]^{n}|W^{n}}(x^{n},y^{n}|W^{n})\right] \\ &\leq \mathbb{E}_{W^{n}}\left[g_{[X]^{n}[Y]^{n}|W^{n}}(x^{n},y^{n}|W^{n})^{2}\right] \\ &\leq \beta_{n}\mu_{\epsilon,n}. \end{aligned}$$
(536)

Here (539) follows by the following inequality. For two

functions  $f(x) \ge 0, g(x) > 0$ ,  $\int f(x) dx \qquad f(x)$ 

$$\frac{\int f(x) \,\mathrm{d}x}{\int g(x) \,\mathrm{d}x} \le \sup_{x} \frac{f(x)}{g(x)}.$$
(542)

Following steps similar to (181)-(186) (but with a lower bound on the exponent of  $\frac{1}{(1-\delta_{0,n})(1-\delta_{1,n})(1-\delta_{2,n})} - 1$  can be obtained in the proof of Lemma 16, which was derived by the large deviation theory, instead of the method of types), we get that there exists  $\epsilon'_n \to 0$  exponentially fast such that (534) with  $\epsilon'$  replaced by  $\epsilon'_n$  converges to zero doubly exponentially fast, 2) as long as  $\left| \begin{bmatrix} \mathcal{A}_{\epsilon}^{(n)} \end{bmatrix} \times \begin{bmatrix} \mathcal{A}_{\epsilon}^{(n)} \end{bmatrix} \right|$  is growing slower than doubly exponentially fast. Hence

$$\mathbb{P}_{\mathcal{C}_n}\left(e^{D_{\infty}(P_{[X]^n[Y]^n}\|\pi_{[X][Y]}^n)} \ge 1 + \epsilon'_n\right) \to 0 \tag{543}$$

doubly exponentially fast, as long as  $\left| \left[ \mathcal{A}_{\epsilon}^{(n)} \right] \times \left[ \mathcal{A}_{\epsilon}^{(n)} \right] \right|$  is growing slower than doubly exponentially fast. Obviously, (543) implies there exists a codebook such that  $D_{\infty}(P_{[X]^n[Y]^n} \| \pi^n_{[X][Y]}) \to 0$  exponentially fast. On the other hand, as shown in Remark 12, for the Gaussian

source.

$$L_{\epsilon,n} = \frac{\sqrt{n\left(1+\epsilon\right)}}{1-\rho}.$$
(544)

Similarly to (439), for  $(x^n, y^n) \in \left[\mathcal{A}_{\epsilon}^{(n)}\right] \times \left[\mathcal{A}_{\epsilon}^{(n)}\right] \subseteq \mathcal{L}_{\epsilon,n}^n \times \mathcal{L}_{\epsilon,n}^n$ , and  $|x_i - \hat{x}_i|, |y_i - \hat{y}_i| \leq \Delta_n, \forall i$ , we have

$$\frac{\pi_{XY}^n(x^n, y^n)}{\pi_{XY}^n(\hat{x}^n, \hat{y}^n)} \le \exp\left(n\Delta_n L_{\epsilon, n}\right).$$
(545)

Choose  $\Delta_n = \frac{e^{-n\delta}}{n\frac{\sqrt{n(1+\epsilon)}}{1-\rho}}$  for some  $\delta > 0$ , then

$$\left| \left[ \mathcal{A}_{\epsilon}^{(n)} \right] \times \left[ \mathcal{A}_{\epsilon}^{(n)} \right] \right| = \left( \frac{\sqrt{n\left(1+\epsilon\right)}}{\Delta_n} \right)^{2n} \tag{546}$$

$$= \left(\frac{n^2 \left(1+\epsilon\right)}{\left(1-\rho\right)e^{-n\delta}}\right)^{2n^2} \tag{547}$$

$$=e^{2n^2\delta+2n\log\frac{n}{1-\rho}},\qquad(548)$$

which grows much slower than doubly exponentially fast. Hence the doubly exponential convergence of (543) is guaranteed.

Define  $U^n, V^n \sim \text{Unif}(I^n_{\Delta_n})$  with  $I^n_{\Delta_n} = [0, \Delta_n]^n$  are mutually independent, and also independent of  $[X]^n, [Y]^n$ . Then

$$e^{D_{\infty}(P_{[X]^{n}+U^{n},[Y]^{n}+V^{n}} \| \pi_{XY}^{n})}$$

$$= \sup_{x^{n},y^{n}} \frac{P_{[X]^{n}+U^{n},[Y]^{n}+V^{n}}(x^{n},y^{n})}{\pi_{[X]^{n}+U^{n},[Y]^{n}+V^{n}}(x^{n},y^{n})}$$

$$\times \frac{\pi_{[X]^{n}+U^{n},[Y]^{n}+V^{n}}(x^{n},y^{n})}{\pi_{XY}^{n}(x^{n},y^{n})}$$

$$\leq \sup_{x^{n},y^{n}} \frac{P_{[X]^{n}+U^{n},[Y]^{n}+V^{n}}(x^{n},y^{n})}{\pi_{[X]^{n}+U^{n},[Y]^{n}+V^{n}}(x^{n},y^{n})}$$

$$\times \sup_{(x'^{n},y'^{n})\in([x]^{n}+I_{\Delta_{n}}^{n})\times([y]^{n}+I_{\Delta_{n}}^{n})} \frac{\pi_{XY}^{n}(x'^{n},y'^{n})}{\pi_{XY}^{n}(x'^{n},y'^{n})}$$

$$\leq \exp(n\Delta_{n}L_{\epsilon,n}) e^{D_{\infty}(P_{[X]^{n}[Y]^{n}} \| \pi_{[X][Y]}^{n})}$$
(549)
(551)

$$g_{X^{n}Y^{n}|W^{n}}(x^{n}, y^{n}|w^{n})$$

$$:= \frac{P_{X^{n}|W^{n}}(x^{n}|w^{n})P_{Y^{n}|W^{n}}(y^{n}|w^{n})}{\pi_{XY}^{n}(x^{n}, y^{n})}$$
(526)

$$\leq \delta_{12,n} \sup_{e^{4n\epsilon - nh(X|W) - nh(Y|W) + n\log 2\pi\sqrt{1 - \rho^2} + \frac{n(1 + 2\epsilon) - \rho\left(x^{n \top}y^n\right)}{1 - \rho^2}}$$
(527)

$$\leq \delta_{12,n} \sup_{(x^n,y^n)\in\mathcal{A}} e^{4n\epsilon - nh(X|W) - nh(Y|W) + n\log 2\pi\sqrt{1-\rho^2} + \frac{1}{1-\rho^2} \left[ n(1+2\epsilon) - \rho \left( \|w^n\|^2 + (x^n - w^n)^\top (y^n - w^n) - 2n(3-\rho)\epsilon \right) \right]}$$
(528)

$$\leq \delta_{12,n} e^{4n\epsilon - nh(X|W) - nh(Y|W) + n\log 2\pi\sqrt{1-\rho^2} + \frac{1}{1-\rho^2} [n(1+2\epsilon) - \rho(n\rho(1-2\epsilon) - n(1-\rho)(1+4\epsilon) - 2n(3-\rho)\epsilon)]}$$
(529)  
=  $\delta_{12,n} e^{\frac{6-10\rho}{1-\rho^2}n\epsilon - nh(X|W) - nh(Y|W) + n\left(\log 2\pi\sqrt{1-\rho^2} + \frac{(1-\rho)(1+2\rho)}{1-\rho^2}\right)}$ (530)  
=  $\delta_{12,n} e^{n\left(\frac{1}{2}\log\frac{1+\rho}{1-\rho} + \frac{\rho}{1+\rho} + \frac{6-10\rho}{1-\rho^2}\epsilon\right)}$ (531)

$$= \delta_{12,n} e^{n\left(\frac{1}{2}\log\frac{1+\rho}{1-\rho} + \frac{\rho}{1-\rho^2}\epsilon\right)}$$

$$= \beta_n,$$
(531)
(532)

$$=:\beta_n,$$
 (

$$\mu_{\epsilon,n} := \mathbb{E}_{W^{n}} \left[ g_{[X]^{n}[Y]^{n}|W^{n}}(x^{n}, y^{n}|w^{n}) \right]$$

$$= \int \frac{Q_{W}^{n}(w^{n}) \mathbf{1} \left\{ w^{n} \in \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_{W}) \right\}}{Q_{W}^{n} \left( \mathcal{A}_{\frac{\epsilon}{2}}^{(n)}(Q_{W}) \right)}$$
(537)

and hence

$$D_{\infty}(P_{[X]^{n}+U^{n},[Y]^{n}+V^{n}} \| \pi_{XY}^{n})$$

$$\leq n\Delta_{n}L_{\epsilon,n} + D_{\infty}(P_{[X]^{n}[Y]^{n}} \| \pi_{[X][Y]}^{n}) \qquad (552)$$

$$= e^{-n\delta} + D_{\infty}(P_{[X]^{n}[Y]^{n}} \| \pi_{[X][Y]}^{n}) \to 0 \qquad (553)$$

exponentially fast.

# APPENDIX J **PROOF OF PROPOSITION 2**

For  $\alpha = 0$  and 1, by definition, one can easily obtain that  $G_0(\pi_{XY}) = \log \operatorname{rank}^+(\pi_{XY})$  and  $G_1(\pi_{XY}) = G(\pi_{XY})$ . Next we consider the case of  $\alpha = \infty$ . For this case,

$$H_{\infty}(W) = -\log\max_{w} P_W(w).$$
(554)

# Hence

$$G_{\infty}(\pi_{XY}) \tag{555}$$

$$= \min_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} -\log\max_{w} P_W(w)$$
(556)

$$= -\log \max_{P_W P_{X|W} P_{Y|W}: P_{XY} = \pi_{XY}} \max_{w} P_W(w)$$
(557)

$$= -\log \max_{w} \max_{P_{W}P_{X|W}P_{Y|W}: P_{XY} = \pi_{XY}} P_{W}(w)$$
(558)

$$\geq -\log \max_{w} \max_{\substack{P_{X|W}P_{Y|W}:\\P_{W}(w)P_{X|W}(x|w)P_{Y|W}(y|w)\\\leq \pi_{XY}(x,y), \forall (x,y)}} P_{W}(w)$$
(559)

$$\geq \min_{w} \min_{P_{X|W=w}, P_{Y|W=w}} D_{\infty}(P_{X|W=w}P_{Y|W=w} \| \pi_{XY})$$
(560)

$$\geq \min_{Q_X, Q_Y} D_{\infty}(Q_X Q_Y \| \pi_{XY}).$$
(561)

On the other hand, denote  $(Q_X^*, Q_Y^*)$  as an optimal pair of distributions attaining the minimum in the optimization problem  $\min_{Q_X, Q_Y} D_{\infty}(Q_X Q_Y || \pi_{XY})$ . Let  $\epsilon := D_{\infty}(Q_X^* Q_Y^* || \pi_{XY})$ . By Lemma 5, we can decompose  $\pi_{XY}$  as

$$\pi_{XY} = e^{-\epsilon} Q_X^* Q_Y^* + \left(1 - e^{-\epsilon}\right) \widehat{P}, \qquad (562)$$

where

$$\widehat{P} := \begin{cases} \text{any distribution} & \epsilon = 0\\ \frac{e^{\epsilon} Q_X^* Q_Y^* - \pi_{XY}}{e^{\epsilon} - 1} & \epsilon \in (0, \infty) \\ Q_X^* Q_Y^* & \epsilon = \infty \end{cases}$$
(563)

Then set  $\mathcal{W} := (\mathcal{X} \times \mathcal{Y}) \cup \{w_0\}$  with some  $w_0 \notin \mathcal{X} \times \mathcal{Y}$ , and choose

$$P_W(w) := \begin{cases} e^{-\epsilon} & w = w_0\\ (1 - e^{-\epsilon}) \,\widehat{P}(x', y') & w = (x', y') \in \mathcal{X} \times \mathcal{Y} , \end{cases}$$
(564)

and

$$P_{X|W}(x|w) := \begin{cases} Q_X^* & w = w_0 \\ 1 \{x = x'\} & w = (x', y') \in \mathcal{X} \times \mathcal{Y} \end{cases}$$

$$P_{Y|W}(y|w) := \begin{cases} Q_Y^* & w = w_0 \\ 1 \{y = y'\} & w = (x', y') \in \mathcal{X} \times \mathcal{Y} \end{cases}$$
(566)

It is easy to verify that such a distribution  $P_W P_{X|W} P_{Y|W}$  satisfies

$$P_{XY} = \pi_{XY},\tag{567}$$

$$H_{\infty}(W) \le \epsilon. \tag{568}$$

Hence

$$G_{\infty}(\pi_{XY}) \le \min_{Q_X, Q_Y} D_{\infty}(Q_X Q_Y \| \pi_{XY}).$$
(569)

Combining (561) and (569) we conclude that

$$G_{\infty}(\pi_{XY}) = \min_{Q_X, Q_Y} D_{\infty}(Q_X Q_Y \| \pi_{XY}).$$
 (570)

Now we claim

$$G_{\infty}(\pi_{XY}^n) = nG_{\infty}(\pi_{XY}). \tag{571}$$

This follows since on one hand, by choosing  $Q_{X^n}, Q_{Y^n}$  as product distributions, we have

$$G_{\infty}(\pi_{XY}^n) \le nG_{\infty}(\pi_{XY}). \tag{572}$$

On the other hand,

=

$$D_{\infty}(Q_{X^{n}}Q_{Y^{n}} \| \pi_{XY}^{n}) = \max_{x^{n},y^{n}} \log \frac{Q_{X^{n}}(x^{n}) Q_{Y^{n}}(y^{n})}{\pi_{XY}^{n}(x^{n},y^{n})}$$
(573)
$$= \max_{x^{n},y^{n}} \sum_{i=1}^{n} \log \frac{Q_{X_{i}|X^{i-1}}(x_{i}|x^{i-1}) Q_{Y_{i}|Y^{i-1}}(y_{i}|y^{i-1})}{\pi_{XY}(x_{i},y_{i})}$$

$$\binom{n-1}{i} \qquad (574)$$

$$= \max_{x^{n-1}, y^{n-1}} \left( \sum_{i=1}^{N-1} \log \frac{Q_{X_i|X^{i-1}}(x_i|x^{i-1}) Q_{Y_i|Y^{i-1}}(y_i|y^{i-1})}{\pi_{XY}(x_i, y_i)} + \max_{x_n, y_n} \log \frac{Q_{X_n|X^{n-1}}(x_i|x^{i-1}) Q_{Y_i|Y^{i-1}}(y_i|y^{i-1})}{\pi_{XY}(x_i, y_i)} \right)$$
(575)

$$\geq \max_{x^{n-1}, y^{n-1}} \left( \sum_{i=1}^{n-1} \log \frac{Q_{X_i|X^{i-1}}\left(x_i|x^{i-1}\right) Q_{Y_i|Y^{i-1}}\left(y_i|y^{i-1}\right)}{\pi_{XY}\left(x_i, y_i\right)} + G_{\infty}(\pi_{XY}) \right)$$
(576)

$$\geq \max_{x^{n-2}, y^{n-2}} \left( \sum_{x^{n-2}, y^{n-2}}^{n-2} \log \frac{Q_{X_i|X^{i-1}}(x_i|x^{i-1}) Q_{Y_i|Y^{i-1}}(y_i|y^{i-1})}{\pi_{YY}(x_i, y_i)} \right)$$

$$+ 2G_{\infty}(\pi_{XY})$$

$$(577)$$

$$\geq nG_{\infty}(\pi_{XY}). \tag{578}$$

Therefore,

$$T_{\text{Exact}}^{(\infty)}(\pi_{XY}) = \lim_{n \to \infty} \frac{1}{n} \min_{Q_{X^n}, Q_{Y^n}} G_{\infty}(\pi_{XY}^n)$$
(579)

$$=G_{\infty}(\pi_{XY}). \tag{580}$$

# Acknowledgements

The authors would like to thank the Associate Editor Prof. Maxim Raginsky and reviewers for their extensive, constructive and helpful feedback to improve the manuscript.

#### REFERENCES

- L. Yu and V. Y. F. Tan. On exact and ∞-Rényi common informations. In 2019 IEEE International Symposium on Information Theory (ISIT), pages 2229–2233. IEEE, 2019.
- [2] A. Wyner. The common information of two dependent random variables. *IEEE Trans. on Inform. Theory*, 21(2):163–179, 1975.
- [3] L. Yu and V. Y. F. Tan. Wyner's common information under Rényi divergence measures. *IEEE Trans. on Inform. Theory*, 64(5):3616–3632, 2018.
- [4] L. Yu and V. Y. F. Tan. Corrections to "Wyner's common information under Rényi divergence measures". arXiv preprint arXiv:1810.02534, 2018.
- [5] P. Cuff. Distributed channel synthesis. *IEEE Trans. on Inform. Theory*, 59(11):7071–7096, 2013.
- [6] G. R. Kumar, C. T. Li, and A. El Gamal. Exact common information. In *IEEE International Symposium on Information Theory (ISIT)*, pages 161–165. IEEE, 2014.
- [7] M. Hayashi. General nonasymptotic and asymptotic formulas in channel resolvability and identification capacity and their application to the wiretap channel. *IEEE Trans. on Inform. Theory*, 52(4):1562–1575, 2006.

- [8] H. S. Witsenhausen. Values and bounds for the common information of two discrete random variables. *SIAM Journal on Applied Mathematics*, 31(2):313–333, 1976.
- [9] B. N. Vellambi and J. Kliewer. Sufficient conditions for the equality of exact and Wyner common information. In *Communication, Control,* and Computing (Allerton), 2016 54th Annual Allerton Conference on, pages 370–377. IEEE, 2016.
- [10] C. T. Li and A. El Gamal. Distributed simulation of continuous random variables. *IEEE Trans. on Inform. Theory*, 63(10):6329–6343, 2017.
- [11] G. Xu, W. Liu, and B. Chen. Wyner's common information: Generalizations and a new lossy source coding interpretation. arXiv preprint arXiv:1301.2237, 2013.
- [12] L. Yu, H. Li, and C. W. Chen. Generalized common informations: Measuring commonness by the conditional maximal correlation. arXiv preprint arXiv:1610.09289, 2016.
- [13] P. Gács and J. Körner. Common information is far less than mutual information. *Problems of Control and Information Theory*, 2(2):149– 162, 1973.
- [14] E. Nummelin. Uniform and ratio limit theorems for Markov renewal and semi-regenerative processes on a general state space. Annales de l'IHP Probabilités et Statistiques, 14(2):119–143, 1978.
- [15] K. B. Athreya and P. Ney. A new approach to the limit theory of recurrent Markov chains. *Transactions of the American Mathematical Society*, 245:493–501, 1978.
- [16] G. O. Roberts and J. S Rosenthal. General state space Markov chains and mcmc algorithms. *Probability Surveys*, 1:20–71, 2004.
- [17] S.-W. Ho and S. Verdú. On the interplay between conditional entropy and error probability. *IEEE Trans. on Inform. Theory*, 56(12):5930– 5942, 2010.
- [18] A. El Gamal and Y.-H. Kim. Network Information Theory. Cambridge university press, 2011.
- [19] T. M. Cover and J. A. Thomas. *Elements of Information Theory*. Wiley-Interscience, 2nd edition, 2006.
- [20] Siu-Wai S. W. Ho and R. W. Yeung. On information divergence measures and a unified typicality. *IEEE Trans. on Inform. Theory*, 56(12):5893–5905, 2010.
- [21] S. W. Ho. Markov lemma for countable alphabets. In *Information Theory Proceedings (ISIT), 2010 IEEE International Symposium on*, pages 1448–1452. IEEE, 2010.
- [22] S. T. Rachev and L. Rüschendorf. Mass Transportation Problems: Volume I: Theory, volume 1. Springer Science & Business Media, 1998.
- [23] C. Villani. Topics in optimal transportation. Number 58. American Mathematical Soc., 2003.
- [24] L. Yu and V. Y. F. Tan. Exact channel synthesis. *arXiv preprint arXiv:1810.13246*, 2018.
- [25] L. Yu and V. Y. F. Tan. Rényi resolvability and its applications to the wiretap channel. *IEEE Trans. on Inform. Theory*, 65(3):1862–1897, 2019.
- [26] B. N. Vellambi and J. Kliewer. New results on the equality of exact and Wyner common information rates. In 2018 IEEE International Symposium on Information Theory (ISIT), pages 151–155. IEEE, 2018.
- [27] T. Van Erven and P. Harremoës. Rényi divergence and Kullback-Leibler divergence. *IEEE Trans. on Inform. Theory*, 60(7):3797–3820, 2014.
- [28] M. Gil, F. Alajaji, and T. Linder. Rényi divergence measures for commonly used univariate continuous distributions. *Information Sciences*, 249:124–131, 2013.
- [29] M. Raginsky. Empirical processes, typical sequences, and coordinated actions in standard borel spaces. *IEEE Trans. on Inform. Theory*, 59(3):1288–1301, 2013.
- [30] J. Jeon. A generalized typicality for abstract alphabets. In *Information Theory (ISIT), 2014 IEEE International Symposium on*, pages 2649–2653. IEEE, 2014.
- [31] C. H. Bennett, P. W. Shor, J. A. Smolin, and A. V. Thapliyal. Entanglement-assisted capacity of a quantum channel and the reverse Shannon theorem. *IEEE Trans. on Inform. Theory*, 48(10):2637–2655, 2002.
- [32] A. Winter. Compression of sources of probability distributions and density operators. arXiv preprint quant-ph/0208131, 2002.
- [33] C. H. Bennett, I. Devetak, A. W. Harrow, P. W. Shor, and A. Winter. The quantum reverse Shannon theorem and resource tradeoffs for simulating quantum channels. *IEEE Trans. on Inform. Theory*, 60(5):2926–2959, 2014.
- [34] P. Harsha, R. Jain, D. McAllester, and J. Radhakrishnan. The communication complexity of correlation. *IEEE Trans. on Inform. Theory*, 56(1):438–449, 2010.

- [35] C. T. Li and A. El Gamal. Strong functional representation lemma and applications to coding theorems. *IEEE Trans. on Inform. Theory*, 64(11):6967–6978, 2018.
- [36] M. Yannakakis. Expressing combinatorial optimization problems by linear programs. *Journal of Computer and System Sciences*, 43(3):441– 466, 1991.
- [37] R. Jain, Y. Shi, Z. Wei, and S. Zhang. Efficient protocols for generating bipartite classical distributions and quantum states. *IEEE Trans. on Inform. Theory*, 59(8):5171–5178, 2013.
- [38] G. Braun and S. Pokutta. Common information and unique disjointness. *Algorithmica*, 76(3):597–629, 2016.
- [39] A. Vandaele, N. Gillis, F. Glineur, and D. Tuyttens. Heuristics for exact nonnegative matrix factorization. *Journal of Global Optimization*, 65(2):369–400, 2016.
- [40] L. Lovász. Communication complexity: A survey. Springer-Verlag, 1990.
- [41] P. Cuff, H. Permuter, and T. Cover. Coordination capacity. IEEE Trans. on Inform. Theory, 56(9):4181–4206, 2010.
- [42] D. M. Titterington, A. F. M. Smith, and U. E. Makov. *Statistical analysis of finite mixture distributions*. J. Willey & Sons, 1985.
- [43] R. Estrada and R. P. Kanwal. A distributional approach to Asymptotics: Theory and Applications. Springer Science & Business Media, 2012.
- [44] L. Yu and V. Y. F. Tan. Simulation of random variables under rényi divergence measures of all orders. *IEEE Trans. on Inform. Theory*, 65(6):3349–3383, 2019.
- [45] S. Boucheron, G. Lugosi, and P. Massart. Concentration inequalities: A nonasymptotic theory of independence. Oxford university press, 2013.
- [46] S. Boyd and L. Vandenberghe. Convex optimization. Cambridge university press, 2004.

Lei Yu received the B.E. and Ph.D. degrees, both in electronic engineering, from University of Science and Technology of China (USTC) in 2010 and 2015, respectively. From 2015 to 2017, he was a postdoctoral researcher at the Department of Electronic Engineering and Information Science (EEIS), USTC. From 2017 to 2019, he was a research fellow at the Department of Electrical and Computer Engineering, National University of Singapore. Currently, he is a postdoc at the Department of Electrical Engineering and Computer Sciences, University of California, Berkeley. His research interests lie in the intersection of information theory, probability theory, and combinatorics.

Vincent Y. F. Tan (S'07-M'11-SM'15) was born in Singapore in 1981. He is currently a Dean's Chair Associate Professor in the Department of Electrical and Computer Engineering and the Department of Mathematics at the National University of Singapore (NUS). He received the B.A. and M.Eng. degrees in Electrical and Information Sciences from Cambridge University in 2005 and the Ph.D. degree in Electrical Engineering and Computer Science (EECS) from the Massachusetts Institute of Technology (MIT) in 2011. His research interests include information theory, machine learning, and statistical signal processing. Dr. Tan received the MIT EECS Jin-Au Kong outstanding doctoral thesis prize in 2011, the NUS Young Investigator Award in 2014, the NUS Engineering Young Researcher Award in 2018, and the Singapore National Research Foundation (NRF) Fellowship (Class of 2018). He is also an IEEE Information Theory Society Distinguished Lecturer for 2018/9. He has authored a research monograph on "Asymptotic Estimates in Information Theory with Non-Vanishing Error Probabilities" in the Foundations and Trends in Communications and Information Theory Series (NOW Publishers). He is currently serving as an Associate Editor of the IEEE Transactions on Signal Processing.