

# Weak limits of the measures of maximal entropy for Orthogonal polynomials.

Carsten Lunde Petersen and Eva Uhre<sup>1</sup>

<sup>1</sup>Department of Science and Environment, Roskilde University,  
4000 Roskilde, Denmark \*

January 29, 2020

## Abstract

In this paper we study the sequence of orthonormal polynomials  $\{P_n(\mu; z)\}$  defined by a Borel probability measure  $\mu$  with non-polar compact support  $S(\mu) \subset \mathbb{C}$ . For each  $n \geq 2$  let  $\omega_n$  denote the unique measure of maximal entropy for  $P_n(\mu; z)$ . We prove that the sequence  $\{\omega_n\}_n$  is pre-compact for the weak-\* topology and that for any weak-\* limit  $\nu$  of a convergent sub-sequence  $\{\omega_{n_k}\}$ , the support  $S(\nu)$  is contained in the filled-in or polynomial-convex hull of the support  $S(\mu)$  for  $\mu$ . And for  $n$ -th root regular measures  $\mu$  the full sequence  $\{\omega_n\}_n$  converge weak-\* to the equilibrium measure  $\omega$  on  $S(\mu)$ .

## 1 Introduction and general results

In the classical study [5] by Stahl and Totik of general orthogonal polynomials they relate the potential and measure theoretic properties of the asymptotic zero distribution for the sequence of orthonormal polynomials defined by a Borel probability measure  $\mu$  on  $\mathbb{C}$  of infinite, but compact support  $S(\mu)$ , to the potential and measure theoretic properties of  $\mu$  and its support. In the paper [4] Christiansen, Henriksen, Pedersen and one of the authors of this paper initiated a study of the relation between the potential theoretic properties of  $\mu$  and the asymptotic (as  $n$  tend to  $\infty$ ) potential and measure theoretic properties of the Julia sets and filled Julia sets of the orthonormal polynomials  $P_n$ . In this paper we extend this with a study of the weak convergence properties of the measures of maximal entropy for the orthonormal polynomials.

For  $\mu$  a Borel probability measure on  $\mathbb{C}$  with infinite and compact support  $S(\mu)$  we denote by  $\{P_n(z)\} := \{P_n(\mu; z)\}$  the unique sequence

$$P_n(z) = \gamma_n z^n + \text{lower order terms.} \quad (1)$$

---

\*The first author would like to thank the Danish Council for Independent Research | Natural Sciences for support via the grant DFF – 4181-00502.

of orthonormal polynomials wrt.  $\mu$ . Then  $P_n$  is also characterized as the unique normalized polynomial of the form (1) which is orthogonal to all lower degree polynomials or equivalently for which the monic polynomial  $p_n(z) = P_n(z)/\gamma_n$  is the unique monic degree  $n$  polynomial of minimal norm in  $L^2(\mu)$ .

For  $S \subset \mathbb{C}$  a compact non-polar subset such as  $S(\mu)$  above we denote by  $\Omega = \Omega(S)$  the unbounded connected component of  $\mathbb{C} \setminus S$ , by  $K = K(S) := \mathbb{C} \setminus \Omega$  the filled-in or just filled  $S$  (denoted by the polynomial convex hull in e.g. [5]), by  $J = J(S) = \partial K = \partial \Omega \subset S$  the outer boundary of  $S$ , by  $g_\Omega$  the Green's function with pole at infinity for  $\Omega$  and finally by  $\omega_S$  the equilibrium measure for  $S$  (with  $\text{Supp}(\omega_S) = J$ ). For a measure  $\mu \in \mathcal{B}$  we shall write  $K(\mu)$  for  $K(S(\mu))$ ,  $J(\mu)$  for  $J(S(\mu))$  and  $\Omega(\mu)$  for  $\Omega(S(\mu))$  or simply  $K, J$  and  $\Omega$ , when the measure  $\mu$  is understood from the context.

**Definition 1.1.** We denote by  $\mathcal{B}$  the set of Borel probability measure on  $\mathbb{C}$  with compact non-polar support.

We denote by  $\mathcal{R}eg$  the set of  $n$ -th root regular measures in  $\mathcal{B}$ , that is

$$\mathcal{R}eg := \left\{ \mu \in \mathcal{B} \mid \lim_{n \rightarrow \infty} (\gamma_n)^{1/n} = \frac{1}{\text{Cap}(S(\mu))} \right\}.$$

For  $\mu \in \mathcal{B}$  and  $P_n$  the associated sequence of orthonormal polynomials, we denote by  $\Omega_n$  the attracted basin of  $\infty$  for  $P_n$ , by  $K_n = \mathbb{C} \setminus \Omega_n$  and  $J_n = \partial K_n = \partial \Omega_n$  respectively the filled Julia set and the Julia set of  $P_n$ , see Section 2. for definitions of Julia set and filled Julia set of a polynomial, by  $g_n$  the Green's function with pole at  $\infty$  for  $\Omega_n$  and by  $\omega_n$  the equilibrium measure for  $J_n$  or equivalently the measure of maximal entropy for  $P_n$  (see [1]) and see also below for definitions of these terms).

Our main result is

**Theorem A.** Let  $\mu \in \mathcal{B}$ , then the sequence  $\{\omega_n\}_{n \geq 2}$  is pre-compact for the weak-\* topology and for any limit measure  $\nu$  for a weakly convergent sub-sequence  $\{\omega_{n_k}\}_k$

$$S(\nu) \subseteq K(\mu).$$

Moreover if  $\mu \in \mathcal{R}eg$  then

$$\omega_n \xrightarrow{\text{weak}^*} \omega_{S(\mu)}.$$

The first part of the theorem is an analogue of the last statement in the following Theorem from [5], recast in the above notation.

**Theorem 1.2** ([5, Thm. 2.1.1, the first part of which was first proven by Fejér [2])). Suppose  $\mu \in \mathcal{B}$ . All zeros of the orthonormal polynomials  $P_n(z)$ ,  $n \in \mathbb{N}$ , are contained in the convex hull  $\text{Co}(K)$ , and for any compact subset  $V \subseteq \Omega$  the number of zeros of  $P_n(z)$ ,  $n \in \mathbb{N}$ , on  $V$  is bounded as  $n \rightarrow \infty$ . Consequently any weak\* limit point of (the normalized counting measures for) the zeros of  $P_n$  is supported on  $K(\mu)$ .

The second part of the theorem should be compared to [5, Thm. 3.1.4] and [5, Thm. 3.6.1]. Such a comparison reveals that the equilibrium measures  $\omega_n$  on

the Julia sets of  $P_n$  have much stronger convergence properties than the counting measure on the roots of  $P_n$ , at least in the case of regular measures  $\mu$ .

Also the main Theorem is a measure theoretic version of the following theorem from [4], which says that under a mild extra condition (see  $\mathcal{B}_+$  below) in capacity any limit of a convergent subsequence  $K_{n_k}$  (convergent in the Hausdorff topology on the space of compact subsets of  $\mathbb{C}$ ) is contained in  $K(\mu)$ .

**Theorem 1.3** ([4, Thm.1.3]). *For  $\mu \in \mathcal{B}_+ := \{\mu \in \mathcal{B} \mid \limsup_{n \rightarrow \infty} |\gamma|_n^{1/n} < \infty\}$  we have*

$$\limsup_{n \rightarrow \infty} K_n \subseteq \text{Co}(K).$$

Moreover, for any  $\epsilon > 0$  and  $V_\epsilon := \{z \in \mathbb{C} \mid g_\Omega(z) \geq \epsilon\}$ ,

$$\lim_{n \rightarrow \infty} \text{Cap}(V_\epsilon \cap K_n) = 0.$$

Our proof follows the approach of Stahl and Totik for Theorem 1.2.

## 2 Background.

### 2.1 Potential theory

We use the notation of Ransford [6]. For  $\mu$  a Borel probability measure on  $\mathbb{C}$  with compact support we define its potential as the sub-harmonic function

$$p_\mu(z) := \int_{\mathbb{C}} \log |z - w| d\mu(w) = \log |z| + o(1),$$

which is harmonic on the complement of  $S(\mu)$ . And we define its energy as

$$I(\mu) := \int_{\mathbb{C} \times \mathbb{C}} \log |z - w| d\mu(w) d\mu(z) = \int_{\mathbb{C}} p_\mu(z) d\mu(z),$$

it satisfies  $-\infty \leq I(\mu) \leq \log \text{diam}(S(\mu))$ . For a Borel set  $B \subset \mathbb{C}$  we denote by  $\mathcal{P}(B)$  the set of probability measures  $\mu$  with compact support  $\text{Supp}(\mu) \subset B$ , the energy of  $B$  as

$$I(B) := \sup\{I(\mu) \mid \mu \in \mathcal{P}(B)\}$$

and its (logarithmic) capacity as

$$\text{Cap}(B) := e^{I(B)}.$$

The set  $B$  is called polar if  $\text{Cap}(B) = 0$  or equivalently  $I(B) = -\infty$  and the set  $B$  is non-polar otherwise. For  $K \subset \mathbb{C}$  a non-polar compact subset there is a unique measure denoted by  $\omega_K$ , which realises the supremum above. It is called the equilibrium measure for  $K$  and its support is contained in the outer boundary of  $K$ . And according to Frostman's Theorem, [6, Thm. 3.3.4] its potential  $p_\mu$  is bounded from below by  $I(K) = I(\omega_K)$  and equals  $I(K)$  on  $K$

except for an  $F_\sigma$  polar subset  $E$  of  $K$ . The Green's function for  $\Omega := \mathbb{C} \setminus K$  is the non-negative sub-harmonic function

$$g_\Omega(z) = p_{\omega_K}(z) - \log(\text{Cap}(K)) = p_{\omega_K}(z) - I(K).$$

The set  $K \setminus E$  is precisely the set of Dirichlet regular boundary points of  $K$  [6, 4.4.9].

## 2.2 Polynomial dynamics

For  $P(z) = \gamma z^d + \dots$  a polynomial of degree  $d > 1$ , an easy computation shows there exists  $R = R_P > 0$  such that for any  $z$  with  $|z| > R : |P(z)| \geq 2|z|$ . Thus the orbit of such  $z$  under iteration converge to  $\infty$ . We denote by  $\Omega_P$  the basin of attraction for  $\infty$  for  $P$ , that is,

$$\Omega_P := \{z \in \mathbb{C} \mid P^k(z) \xrightarrow[k \rightarrow \infty]{} \infty\} = \bigcup_{k \geq 0} P^{-k}(\mathbb{C} \setminus \overline{D(R)}), \quad (2)$$

where  $P^k = \overbrace{P \circ P \circ \dots \circ P}^{k \text{ times}}$ . It follows immediately that  $\Omega_P$  is open and completely invariant, i.e.  $P^{-1}(\Omega_P) = \Omega_P = P(\Omega_P)$ . Denote by  $K_P := \mathbb{C} \setminus \Omega_P \subseteq \overline{D(R)}$  the filled Julia set for  $P$  and by  $J_P := \partial\Omega_P = \partial K_P$  the Julia set for  $P$ . Then  $K_P$  and  $J_P$  are compact and also completely invariant. Clearly any periodic point, i.e. solution of an equation  $P^k(z) = z$ ,  $k \in \mathbb{N}$  belongs to  $K_P$ , so that  $K_P$  is non-empty. It follows from (2) that the filled Julia set  $K_P$  can also be described as the nested intersection

$$K_P = \bigcap_{k \geq 0} P^{-k}(\overline{D(R)}). \quad (3)$$

We denote by  $g_P: \mathbb{C} \rightarrow [0, \infty)$  the Green's function for  $\Omega_P$  with pole at infinity. It follows from (3) that the Green's function  $g_P$  satisfies

$$g_P(z) = \lim_{k \rightarrow \infty} \frac{1}{d^k} \log^+(|P^k(z)|/R) = \lim_{k \rightarrow \infty} \frac{1}{d^k} \log^+ |P^k(z)|.$$

Thus  $g_P$  vanishes precisely on  $K_P$ , i.e. the exceptional set  $E$  for  $g_P$  is empty and hence every point of  $J_P$  is a Dirichlet regular boundary point of  $\Omega_P$ . Moreover

$$g_P(P(z)) = d \cdot g_P(z) \quad \text{and} \quad \text{Cap}(K_P) = \frac{1}{|\gamma|^{\frac{1}{d-1}}}.$$

According to Brolin, [1], the equilibrium measure  $\omega_P$  for the (filled) Julia set  $J_P$  equals the unique measure of maximal entropy for  $P$ .

## 3 Proof of Theorem A.

Recall that for  $\mu \in \mathcal{B}$  and  $P_n$  the associated orthonormal polynomials, we denote by  $\Omega_n$ ,  $J_n$ , and  $K_n$ , respectively, the basin of attraction for  $\infty$ , the Julia set, and

the filled Julia set of  $P_n$ . And moreover by  $g_n$  we denote the Green's function for  $\Omega_n$  and by  $\omega_n$  the equilibrium measure for  $K_n$  with support  $J_n$ .

The following Lemma gives an elementary relation between the filled support  $K(\mu)$  and the filled Julia sets  $K_n$

**Lemma 3.1** ([4, Lemma 2.2]). *Let  $\mu \in \mathcal{B}$  and choose  $R > 0$  such that  $K(\mu) \subset \mathbb{D}(0, R)$ . Then there exists  $N$  such that for all  $n \geq N$ :*

$$K_n \subset P_n^{-1}(\overline{\mathbb{D}(0, R)}) \subset \mathbb{D}(0, R).$$

**Lemma 3.2.** *For a probability measure  $\mu \in \mathcal{B}$  the following are equivalent*

1.

$$\lim_{n \rightarrow \infty} (\gamma_n)^{1/n} = \frac{1}{\text{Cap}(S(\mu))}.$$

2.

$$\text{Cap}(K_n) \xrightarrow{n \rightarrow \infty} \text{Cap}(S(\mu)).$$

3.

$$I(\omega_n) \xrightarrow{n \rightarrow \infty} I(\omega_{S(\mu)})$$

*Proof.* Since  $\text{Cap}(K_n) = \frac{1}{|\gamma|^{1/(n-1)}} = e^{I(\omega_n)}$  the lemma follows from the observation that

$$\lim_{n \rightarrow \infty} (\gamma_n)^{1/n} = \frac{1}{\text{Cap}(S(\mu))} \iff \lim_{n \rightarrow \infty} (\gamma_n)^{1/(n-1)} = \frac{1}{\text{Cap}(S(\mu))}.$$

□

We shall use the following Lemma which is a refinement of [5, Lemma 1.3.2]

**Lemma 3.3.** *Let  $V, S \subset \mathbb{C}$  be two compact subsets with  $V \cap K(S) = \emptyset$  and let  $b, 0 < b < 1$  be arbitrary. Then there exists  $M = M(b, V, S) \in \mathbb{N}$  such that for  $M$  arbitrary points  $x_1, x_2, \dots, x_M \in V$  there exists  $M$  points  $y_1, y_2, \dots, y_M \in \mathbb{C}$  for which the rational function*

$$r(z) = \prod_{j=1}^M \frac{z - y_j}{z - x_j}$$

*has supremum norm on  $S$  bounded by  $b$ :*

$$\|r\|_S \leq b.$$

*Proof.* The above cited Lemma 1.3.2 in [5] shows that the statement holds not for arbitrary  $b$ , but for some value of  $b$  call it  $a$ ,  $0 < a < 1$  and some corresponding value  $m = m(V, S) \in \mathbb{N}$ .

To pass from this to the general statement of the lemma let  $a$  and  $m$  be given as in Stahl and Totiks Lemma, and let  $b, 0 < b < 1$  be arbitrary. Choose  $n \in \mathbb{N}$  such that  $a^n < b$  and set  $M = nm$ . Given  $x_1, \dots, x_M \in V$  apply [5, Lemma 1.3.2] to each of the  $n$  sets of points  $\{x_{j+mk} | 1 \leq j \leq m\}$ ,  $0 \leq k < n$  and multiply. □

**Proposition 3.4.** *For any  $R > 0$ , for any  $\mu \in \mathcal{B}$  and any compact set  $V \subset \mathbb{C}$  with  $V \cap K(S(\mu)) = \emptyset$  there exists  $M = M(R, \mu, V) \in \mathbb{N}$  such that for any  $w, |w| \leq R$  and for any  $n$  the number of pre-images of  $w$  in  $V$  under the orthonormal polynomial  $P_n$  is less than  $M$ . That is*

$$\#(P_n^{-1}(w) \cap V) < M.$$

*Proof.* Fix  $R > 0$  and any compact set  $V \subset \mathbb{C}$  with  $V \cap K(S(\mu)) = \emptyset$ . Let  $b = 1/(1 + R)$  and let  $M = M(b, V, S(\mu))$  be as in Lemma 3.3 above. For each  $n$  let  $p_n(z) = P_n(z)/\gamma_n$  denote the monic orthogonal polynomial of degree  $n$  for  $\mu$ . Then  $p_n$  is the unique degree  $n$  monic polynomial of minimal norm  $1/\gamma_n$  in  $L^2(\mu)$ . Suppose towards a contradiction that for some  $w, |w| \leq R$  and some  $n$  the equation  $P_n(z) - w$  has at least  $M$  solutions  $x_1, \dots, x_M \in V$ . Let  $r$  be the rational function given by Lemma 3.3 such that  $\|r\|_{S(\mu)} \leq b$  and set  $q(z) := r(z) \cdot (P_n(z) - w)/\gamma_n$ . Then  $q$  is a monic polynomial of degree  $n$ . Since  $P_n$  is orthogonal to all polynomials of lower degree, and in particular to the constant polynomial  $w$  we compute

$$\begin{aligned} \|q\|_{L^2(\mu)} &\leq \frac{\|r\|_{S(\mu)}}{\gamma_n} \|P_n(z) - w\|_{L^2(\mu)} \leq \frac{b}{\gamma_n} \sqrt{1 + |w|^2} \\ &\leq \frac{\sqrt{1 + R^2}}{1 + R} \cdot \|p_n\|_{L^2(\mu)} < \|p_n\|_{L^2(\mu)}. \end{aligned}$$

This contradicts the fact that  $p_n$  is the degree  $n$  monic polynomial with minimal  $L^2(\mu)$  norm.  $\square$

**Corollary 3.5.** *For any  $\mu \in \mathcal{B}$  and any compact set  $V \subset \mathbb{C}$  with  $V \cap K(\mu) = \emptyset$  there exists  $M \in \mathbb{N}$  such that any of the orthonormal polynomials  $P_n$ ,  $n \geq 2$  and any  $z \in K_n$  the number of pre-images of  $z$  in  $V$  under  $P_n$  is less than  $M$ . That is*

$$\#(P_n^{-1}(z) \cap V) < M.$$

*Proof.* It follows from Lemma 3.1 that there exists  $R > 0$  such that

$$K_n \subset P_n^{-1}(\overline{\mathbb{D}(R)}) \subset \mathbb{D}(R),$$

for all  $n \geq 2$ , where  $K_n$  is the filled Julia set of  $P_n$ . As  $K_n$  is invariant,  $P_n^{-1}(K_n) = K_n$ . The Corollary follows immediately from this and Proposition 3.4 above.  $\square$

*Proof.* (of Theorem A) According to Lemma 3.1 the sequence of equilibrium measures  $\omega_n$  for  $K_n$  have uniformly bounded support and so the sequence of such measures is pre-compact for the weak-\* topology. Furthermore according to Brodin, [1] the equilibrium measure  $\omega_n$  is also the unique invariant balanced measure for  $P_n$ , i.e. it is the unique probability measure  $\omega$  on  $\mathbb{C}$  such that for any measurable function  $f: \mathbb{C} \rightarrow \mathbb{C}$

$$\int_{\mathbb{C}} f(z) d\omega(z) = \frac{1}{n} \int_{\mathbb{C}} \left( \sum_{w, P_n(w)=z} f(w) \right) d\omega(z). \quad (4)$$

Let  $V \subset \mathbb{C}$  be a compact subset with  $V \cap K(\mu) = \emptyset$  and let  $M \in \mathbb{N}$  be as in Corollary 3.5. Then for the measurable function  $1_V$ , the indicator function for  $V$  we have

$$\omega_n(V) = \int_{\mathbb{C}} 1_V(z) d\omega_n(z) = \frac{1}{n} \int_{\mathbb{C}} \left( \sum_{w, P_n(w)=z} 1_V(w) \right) d\omega_n(z) \leq \frac{M}{n} \xrightarrow{n \rightarrow \infty} 0.$$

This proves the first statement, that for any weak limit  $\nu$  of a convergent subsequence  $\{\omega_{n_k}\}_k$  :  $S(\nu) \subseteq K(\mu)$ .

Next suppose that  $\mu \in \mathcal{R}eg$ . Then in particular  $\mu \in \mathcal{B}_+$ , i.e.  $\limsup_{n \rightarrow \infty} |\gamma|_n^{1/n} < \infty$  and so according to [4, Cor. 4.1 *iii*]

$$\limsup_{n \rightarrow \infty} g_n(z) \leq g_\Omega(z).$$

where  $g_n$  is the Green's function with pole at infinity for  $\Omega_n = \mathbb{C} \setminus K_n$  and  $g_\Omega$  is the Green's function with pole at infinity for  $\Omega = \mathbb{C} \setminus K(\mu)$ . Since  $g_\Omega \equiv 0$  on the interior of  $K(\mu)$  we also have  $\lim_{n \rightarrow \infty} g_n \equiv 0$  uniformly (in the usual sense) on any compact subset  $L$  of the interior of  $K(\mu)$ .

Since  $\omega_n$  can be retrieved from the LaPlacian of  $g_n$  in distributions, it follows that  $\text{Supp}(\nu) \subset J = \partial\Omega$ . In more detail. For any at least  $C^2$  test-function  $\phi$  with compact support  $L$  contained in the interior of  $K(\mu)$  we have

$$\int_{\mathbb{C}} \phi(z) d\omega_n(z) = \frac{1}{2\pi} \int_{\mathbb{C}} \Delta\phi(z) g_n(z) dA(z) \xrightarrow{n \rightarrow \infty} 0$$

where  $dA(z)$  is the standard Euclidean area element. Thus

$$\int_{\mathbb{C}} \phi(z) d\nu(z) = \lim_{n \rightarrow \infty} \int_{\mathbb{C}} \phi(z) d\omega_n(z) = 0.$$

It follows that  $\nu(L) = 0$  for any compact subset  $L$  of the interior  $\overset{\circ}{K}(\mu)$  of  $K(\mu)$ . Take e.g. a compact neighbourhood  $M$  of  $L$  contained in the interior of  $K(\mu)$  and let  $\phi$  be an at least  $C^2$  function with support in  $M$ , with  $0 \leq \phi \leq 1$  and with  $\phi \equiv 1$  on  $L$ . Then  $\nu(L) \leq \int_{\mathbb{C}} \phi(z) d\nu(z) = 0$ . Finally

$$\nu(\overset{\circ}{K}(\mu)) = \sup\{\nu(L) | L \subset \overset{\circ}{K}(\mu), L \text{ compact}\} = 0$$

by regularity of  $\nu$  as a Borel, in fact Radon measure.

Finally we have, by the definition of  $n$ -th-root regularity of  $\mu$ , Lemma 3.2 and [6, Lemma 3.3.3])

$$I(\nu) \geq \limsup_{n \rightarrow \infty} I(\omega_n) = I(\omega_{S(\mu)})$$

Hence  $\nu = \omega_{S(\mu)}$ , since  $\omega_{S(\mu)}$  is the unique measure of maximal energy  $I(S(\mu))$ . Since there is a unique possible limit we have in fact

$$\omega_n \xrightarrow{\text{weak}^*} \omega_{S(\mu)}.$$

□

## References

- [1] H. Brolin, *Invariant sets under iteration of rational functions*, Arkiv f. Math. Band 6 nr 6 (1965) pp. 103-144.
- [2] L. Fejér, *Über die Lage der Nullstellen von Polynomen, die aus Minimumforderungen gewisser Art entspringen* Math. Ann. 85 (1922), 41-48.
- [3] M. Lyubich, *Entropy properties of rational endomorphisms of the Riemann sphere*. Ergod. Th. & Dynam. Sys. 3 (1983), pp.351-385.
- [4] J.S. Christiansen, C. Henriksen, H.L. Pedersen and C.L. Petersen, *Julia Sets of Orthogonal Polynomials*, Potential Analysis (2019) 50: 401. <https://doi.org/10.1007/s11118-018-9687-5>
- [5] H. Stahl and V. Totik, *General Orthogonal Polynomials*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 1992.
- [6] T. Ransford, *Potential Theory in the Complex Plane* London Mathematical Society Student Texts 28, Cambridge University Press 1995.