

THE BASS-QUILLEN CONJECTURE AND SWAN'S QUESTION

DORIN POPESCU

ABSTRACT. We present a question which implies a complete positive answer for the Bass-Quillen Conjecture.

Key words : Regular Rings, Smooth Morphisms, Projective Modules

2010 Mathematics Subject Classification: Primary 13C10, Secondary 19A13, 13H05, 13B40.

INTRODUCTION

The theory of projective modules over polynomial algebras over a regular ring R was an important subject in Commutative Algebra starting with Serre's Conjecture (see [2]) and its extension considered by Bass and Quillen.

Conjecture 1. (*Bass-Quillen Conjecture*, [1, Problem IX], [9]) *Let R be a regular ring. Every finitely generated projective module P over a polynomial R -algebra, $R[T]$, $T = (T_1, \dots, T_n)$ is extended from R , i.e. $P \cong R[T] \otimes_R (P/(T)P)$.*

Important positive answers were given by Quillen (see [9]) and Suslin (see [13]) in dimension ≤ 1 and Murthy [5] in dimension 2. Later Lindel [3] and Swan [6] gave, in particular, positive answers for many regular local rings, essentially of finite type \mathbf{Z} -algebras.

Theorem 2. (*Lindel, Swan*) *Let (R, \mathfrak{m}, k) be a regular local ring, essentially of finite type over \mathbf{Z} and $p = \text{char } k$. The following statements hold.*

- (1) *If $p = 0$ then R is essentially smooth over its prime field.*
- (2) *If $p \notin \mathfrak{m}^2$ then R is essentially smooth over \mathbf{Z} .*
- (3) *If $p = 0$ or $p \notin \mathfrak{m}^2$ then the BQ Conjecture holds for R .*

Then Swan noticed that it is useful to have a positive answer to the following question.

Question 3. (*Swan [6]*) *Is it a regular local ring a filtered inductive limit of regular local rings, essentially of finite type over \mathbf{Z} ?*

The Bass-Quillen Conjecture (shortly BQ Conjecture) is connected with the following one.

Conjecture 4. (*Bass-Quillen-Suslin Conjecture*) *Let R be a local ring, Assume either that R is regular local ring or that $1/r! \in R$. Let v be a unimodular vector over $R[T]$ of length $(r + 1)$. Then v can be completed to an invertible matrix over $R[T]$.*

This conjecture holds in many cases given for example by Rao [10], [11] who also proved the BQ Conjecture for regular local rings of dimension 3 with residue characteristic $\neq 2, 3$.

For a regular local ring (R, \mathfrak{m}, k) containing a field, or with $p := \text{char } k \notin \mathfrak{m}^2$, Swan's question has a positive answer in [7]. Using this partial positive answer and (iii) from Theorem 2 we got the following corollary (see [7, Theorem 4.1] and also [14, Theorems 2.1, 2.2]).

Corollary 5. *The Bass-Quillen Conjecture holds for R if $p = 0$ or $p \notin \mathfrak{m}^2$.*

Recently, we gave a complete positive answer to Swan's Question in [8, Theorem 17] (see here Theorem 7).

The purpose of this paper is to show that a positive answer to the following question gives a complete positive answer to the BQ Conjecture (see Theorem 12).

Question 6. *Let (R, \mathfrak{m}) be a regular local ring, which is essentially smooth over $\mathbf{Z}_{(p)}$ and $b \in \mathfrak{m}^2$. Is it true the BQ Conjecture for the regular local ring $R/(p - b)$?*

We owe thanks to Ravi Rao for some useful comments.

1. THE BASS-QUILLEN CONJECTURE

We start reminding some definitions concerning smooth morphisms after [4], or [14]. A ring morphism $R \rightarrow R'$ of Noetherian rings has *regular fibers* if for all prime ideals $\mathfrak{p} \in \text{Spec } R$ the ring $R'/\mathfrak{p}R'$ is a regular ring. It has *geometrically regular fibers* if for all prime ideals $\mathfrak{p} \in \text{Spec } R$ and all finite field extensions K of the fraction field of R/\mathfrak{p} the ring $K \otimes_{R/\mathfrak{p}} R'/\mathfrak{p}R'$ is regular. A flat morphism of Noetherian rings is *regular* if its fibers are geometrically regular. If it is regular of finite type, or essentially of finite type then it is called *smooth*, resp. *essentially smooth*.

The proof of our positive answer to Swan's Question says actually a little more (see [8, Theorem 17]).

Theorem 7. *Every regular local ring (R, \mathfrak{m}, k) with $0 \neq p = \text{char } k \in \mathfrak{m}^2$ is a filtered inductive limit of regular local rings R_i , essentially smooth over a regular local \mathbf{Z} -algebra $A_i/(p - b_i)$, where (A_i, \mathfrak{a}_i) is a regular local ring, essentially smooth over $\mathbf{Z}_{(p)}$, and $b_i \in \mathfrak{a}_i^2$.*

In fact, this theorem gives also the structure of regular local rings essentially of finite type over \mathbf{Z} in the case $0 \neq \text{char } k \in \mathfrak{m}^2$, which is not covered by (i),(ii) from Theorem 2.

Corollary 8. *Let B be a \mathbf{Z} -algebra regular local, essentially of finite type. Then B has the form $B = A/(p - b)$, where (A, \mathfrak{a}) is a regular local ring essentially smooth over \mathbf{Z} and $b \in \mathfrak{a}^2$.*

Proof. Applying Theorem 7 to B we see that there exists a regular local ring D , essentially smooth over a regular local \mathbf{Z} -algebra $A/(p - b)$, where (A, \mathfrak{a}) is a regular local ring, essentially smooth over $\mathbf{Z}_{(p)}$, and $b \in \mathfrak{a}^2$ such that the identity of B factors through D . Then $B \cong D/q$ for some prime ideal $q \subset D$. Unfortunately, we cannot conclude that B is among these D (see the next remark).

Note that D is a factor of an essentially smooth $\mathbf{Z}_{(p)}$ -algebra D' by $(p - b')$, where b' is a lifting of b to D' . Let q' be the prime ideal of D' containing $p - b'$ and such

that $q'/(p-b') = q$. Changing D by D' and q by q' we may assume that D is essentially smooth over $\mathbf{Z}_{(p)}$. Then D is an etale neighborhood of a localization of a polynomial algebra in t variables Y over $\mathbf{Z}_{(p)}$ (see e.g. [14, Theorem 2.5]) and $(p-b, Y)$ generates the maximal ideal of D . Since D, B are regular local we see that q is generated by a part of a regular system of parameters of D , let us say $p-b, z$, where $z = (z_1, \dots, z_r)$, $r \leq t$. After some linear transformations on Y we may assume that $z_i = Y_i$, $1 \leq i \leq r$. Then $B = D/q$ is an etale neighborhood of a localization of $\mathbf{Z}_{(p)}[Y_{r+1}, \dots, Y_t]/(p-b'')$, b'' being induced by b . \square

Remark 9. Let B be a regular local ring essentially of finite type over \mathbf{Z} , $D_n = B[X_n]$, $n \in \mathbf{N}$ and $\varphi_{n,n+1} : D_n \rightarrow D_{n+1}$ be the B -morphism given by $X_n \mapsto 0$. Then B is the limit of $(D_n, \varphi_{n,n+1})$, the inclusion $B \subset D_n$ has a retraction and $B \not\cong D_n$ for any $n \in \mathbf{N}$.

Next we will need the following two lemmas, the first one is elementary (see e. g. [7, Theorem 4.2]).

Lemma 10. *Let R be a regular local ring, which is a filtered inductive limit of some regular local rings $(R_i)_{i \in I}$. If the BQ Conjecture holds for all R_i , $i \in I$, then it holds for R too.*

Proof. Let M be a finitely generated projective module over $R[T]$, $T = (T_1, \dots, T_n)$. Then $M \cong R \otimes_{R_i} M_i$ for some finitely generated projective $R_i[T]$ -module M_i . Indeed, if M is defined by an idempotent φ from $\text{End}(L)$ for some $L = R^t$ then we may find i such that φ is extended from an endomorphism φ_i of R_i^t . Also we may find i such that φ_i is idempotent, and defines the wanted M_i . As BQ Conjecture holds for R_i we get M_i free and so M is free too. \square

The following lemma follows easily from [3]. However, we give here a proof in sketch.

Lemma 11. *Let $R \rightarrow R'$ be an essentially smooth morphism between regular local rings. If the BQ Conjecture holds for R then it holds for R' too.*

Proof. R' is an etale neighborhood of a localization of a polynomial algebra A over R (see e.g. [14, Theorem 2.5]). If the BQ Conjecture holds for R then it holds for A too by [12]. Now it is enough to apply the Corollary from [3]. \square

Theorem 12. *If Question 6 has a positive answer then the BQ Conjecture holds for all regular rings.*

Proof. By Quillen's Patching Theorem [9, Theorem 1] we may prove the conjecture only for regular local rings. Let (R, \mathfrak{m}, k) be a regular local ring. Using [7, Theorem 3.1] we may suppose that $0 \neq p := \text{char } k \in \mathfrak{m}^2$.

By Theorem 7, R is a filtered inductive limit of some regular local rings D , essentially smooth over a regular local ring of the form $A/(p-b)$, where (A, \mathfrak{a}) is a regular local ring, essentially smooth over \mathbf{Z} and $b \in \mathfrak{a}^2$.

Then the BQ Conjecture holds for A by Lemma 11 and for $A/(p-b)$ by Question 6. Applying Lemma 11 it follows that BQ holds for D . The final result is a consequence of Lemma 10. \square

We end the section with a special form of Theorem 7 in the frame of the discrete valuation rings (DVRs for short) in the idea of [8, Theorem 8]. Actually, [8, Theorem 8] has a complicated proof given to illustrate [8, Theorem 17] in the DVR case. The proof below is easier and does not use Néron Desingularization.

Theorem 13. *Let (A, \mathfrak{m}, k) be a DVR with $0 \neq p = \text{char } k \in \mathfrak{m}^2$. Suppose that k is separably generated over \mathbf{F}_p . Then A is a filtered inductive union of DVRs essentially of finite type over \mathbf{Z} .*

Proof. As in [8, Theorem 8], let $y = (y_i)_{i \in I}$ be a system of elements of A inducing a separable transcendence base (\bar{y}) of k over \mathbf{F}_p . Then $C_0 = \mathbf{Z}[Y]_{p\mathbf{Z}[Y]}$ for some variables $Y = (Y_i)_{i \in I}$ is a DVR and the map $C_0 \rightarrow A$, $Y \rightarrow y$ defines a ramified extension inducing an algebraic separable residue field extension $\mathbf{F}_p(\bar{y}) \subset k$.

Note that A is a filtered inductive union of DVRs $A_L = A \cap L$ with $L \subset \text{Fr}(A)$ a finite type field extension of $\mathbf{Q}(y)$. We claim that L must be finite over $\mathbf{Q}(y)$. Indeed, assume that $z \in L$ is transcendental over $\mathbf{Q}(y)$ and $\mathfrak{m}^s \subset A$ for some $s \in \mathbf{N}$. Choose r such that $p^r > s$ and consider the DVR extension $A_{L''} \subset A_{L'}$ for $L' = \mathbf{Q}(y, z)$ and $L'' = \mathbf{Q}(y, z^{p^r})$. The residue field extension induced by $A_{L''} \subset A_{L'}$ is pure inseparable and also separable by assumption. Then it is trivial and so the ramification index of $A_{L''} \subset A_{L'}$ is $p^r > s$. Contradiction!

Since L is finite over $\mathbf{Q}(y)$ it follows that A_L is a localization of the integral closure of C_0 in L and so essentially finite over C_0 , which is enough. \square

2. SWAN'S QUESTION IN THE NON-REDUCED CASE

We start this section reminding the first part of [8, Theorem 17].

Theorem 14. *Let (A, \mathfrak{m}, k) be a Noetherian local ring, $s = (s_1, \dots, s_m)$ some positive integers and $\gamma = (\gamma_1, \dots, \gamma_m)$ a system of nilpotents of A . Suppose that $0 \neq p \in \mathfrak{m}^2$, $R = A/(\gamma)$ is a regular local ring and A is a flat $\mathfrak{N} = \mathbf{Z}_{(p)}[\Gamma]/(\Gamma^s)$ -algebra, $\Gamma \mapsto \gamma$ with $\Gamma = (\Gamma_1, \dots, \Gamma_m)$ some variables, and (Γ^s) denotes the ideal $(\Gamma_1^{s_1} \cdots \Gamma_m^{s_m})$. Then A is a filtered inductive limit of some Noetherian local \mathfrak{N} -algebras $(F_i)_i$ essentially of finite type with $F_i/\Gamma F_i$ regular local rings.*

Remark 15. The above theorem holds also when $p = 0$ or $p \notin \mathfrak{m}^2$ as in [7, Theorem 4.1] because the map $\mathfrak{N} \rightarrow A$ is regular in this case.

Lemma 16. *In the notation and hypothesis of the above theorem, let (B, \mathfrak{b}) be a Noetherian local ring, and $z = (z_1, \dots, z_m)$ a system of elements of B . Suppose that*

- (1) for all $i \in [m]$ $s_i > 1$, $z_i^{s_i} = 0$,
- (2) for all $i \in [m]$ and $j \in [s_i - 1]$, $(z_1, \dots, z_{i-1}) : z_i^j = (z_1, \dots, z_{i-1}, z_i^{s_i-j})$,
- (3) $B/(z)$ is a regular local ring.

Then B is a flat \mathfrak{N} -algebra by the map $\varphi : \mathfrak{N} \rightarrow B$, $\Gamma \mapsto z$.

Proof. Suppose $m = 1$. Note that the minimal free resolution of $\mathfrak{N}/(\Gamma)$,

$$\dots \rightarrow \mathfrak{N}/(\Gamma^s) \xrightarrow{\Gamma} \mathfrak{N}/(\Gamma^s) \xrightarrow{\Gamma^{s-1}} \mathfrak{N}/(\Gamma^s) \xrightarrow{\Gamma} \mathfrak{N}/(\Gamma^s) \rightarrow \mathfrak{N}/(\Gamma) \rightarrow 0$$

gives after tensorizing with B a minimal free resolution of $B/(z)$. Thus $\mathrm{Tor}_1^{\mathfrak{A}}(\mathfrak{A}/(\Gamma), B) = 0$. As $B/(z)$ is a flat \mathbf{Z} -algebra we get φ flat using the Local Criteria of Flatness [4, Theorem 49, (20.C)].

Induct on m and assume $m > 1$. Then $B/(z_m)$ is a flat $\mathfrak{A}/(\Gamma_m)$ -algebra by induction hypothesis. As in the case $m = 1$ we see that $\mathrm{Tor}_1^{\mathfrak{A}}(\mathfrak{A}/(\Gamma_m), B) = 0$ and so φ is flat using the Local Criteria of Flatness. \square

Remark 17. Let R be a regular local ring, $s = (s_1, \dots, s_m) \in \mathbf{N}^m$ and $z = (z_1, \dots, z_m)$ be a part of a regular system of parameters of R . Then $B = R/(z_1^{s_1}, \dots, z_m^{s_m})$ satisfies the above lemma.

Using the above lemma and Theorem 14 we get the following result.

Theorem 18. *In the notation and hypothesis of the above lemma, B is a filtered inductive limit of some Noetherian local \mathbf{Z} -algebras $(F_i)_i$ essentially of finite type with $F_i/\sqrt{(0)}$ regular local rings.*

As in Corollary 8 we get the following result.

Corollary 19. *In the notation and hypothesis of Lemma 16, suppose that $B/(z)$ is essentially of finite type over \mathbf{Z} . Then B is essentially smooth over a local \mathfrak{A} -algebra D of type $C/(p-b)$, where (C, \mathfrak{q}) is a local ring, essentially smooth over \mathfrak{A} with $b \in \mathfrak{q}^2$ (so $D/\Gamma D$ and $C/\Gamma C$ are regular).*

Theorem 20. *Let (A, \mathfrak{m}, k) be a Noetherian local ring, $T = (T_1, \dots, T_n)$ some variables and \mathfrak{b} a nilpotent ideal of A . Suppose that $R = A/\mathfrak{b}$ is a regular local ring and the BQ Conjecture holds for R (for example $p := \mathrm{char} k = 0$, or $p \notin \mathfrak{m}^2$). Then any finitely generated projective $A[T]$ -module is free.*

Proof. Let M be a finitely generated projective $A[T]$ -module. Then $\bar{M} = M/\mathfrak{b}M$ is finitely generated projective over $R[T]$ and so it is free. Let $x = (x_1, \dots, x_r)$ be a system of elements from M inducing a basis in \bar{M} . Then $M = \langle x \rangle + \mathfrak{b}M = \langle x \rangle + \mathfrak{b}^t M = \langle x \rangle$ for $t \in \mathbf{N}$ with $\mathfrak{b}^t = 0$. Thus the map $\varphi : A[T]^n \rightarrow M$ given by $(a_1, \dots, a_n) \mapsto \sum_i a_i x_i$ is surjective. Set $N = \mathrm{Ker} \varphi$. Tensorizing with $A[T]/(\mathfrak{b})$ the exact sequence

$$0 \rightarrow N \rightarrow A[T]^n \rightarrow M \rightarrow 0$$

we get the following exact sequence

$$\mathrm{Tor}_1^{A[T]}(M, A[T]/(\mathfrak{b})) \rightarrow N/\mathfrak{b}N \rightarrow R[T]^n \rightarrow \bar{M} \rightarrow 0,$$

where the first module is zero because M is a flat $A[T]$ -module. As the last map is injective we see that $N = \mathfrak{b}N = \mathfrak{b}^t N = 0$, that is φ is an isomorphism. \square

REFERENCES

- [1] H. Bass, *Some problems in "classical" algebraic K-theory* in Algebraic K-theory II, Springer Lect. Notes **342**, (1973), 1–70.
- [2] T. Y. Lam, *Serre's Conjecture*, Springer Lect. Notes in Math., **635**, Berlin, 1978.
- [3] H. Lindel, *On the Bass-Quillen Conjecture concerning projective modules over polynomial rings*, Invent. Math., **65**, (1981), 319–323.

- [4] H. Matsumura, *Commutative Algebra*, Benjamin, New-York, 1980.
- [5] M. P. Murty, *Projective $A[X]$ -modules*, J. London. Math. Soc., **41**, (1966), 453–456.
- [6] M. P. Murty, A letter containing Swan’s notes on Lindel’s results, 1987.
- [7] D. Popescu, *Polynomial rings and their projective module*, Nagoya Math. J., **113**, (1989), 121–128.
- [8] D. Popescu, *On a question of Swan. With an appendix by Kestutis Cesnavicius*, Algebraic Geometry, 6(6), (2019), 716–729, arXiv/AC:1803.06956.
- [9] D. Quillen, *Projective modules over polynomial rings*, Invent. Math., **36**, (1976), 167–171.
- [10] R. Rao, *Two Examples of the Bass-Quillen-Suslin Conjectures*, Math. Ann., **279**, (1987), 227–238.
- [11] R. Rao, *The Bass-Quillen conjecture in dimension three but characteristic $\neq 2, 3$ via a question of A. Suslin*, Invent. Math., **93**, (1988), 609–618.
- [12] M. Roitman, *On projective modules over polynomial rings*, J. Algebra, **58**, (1979), 51–63.
- [13] A. A. Suslin, *Projective modules over a polynomial ring are free*, Soviet. Math. Dokl., **17**, 1976, 1160–1164.
- [14] R. Swan, *Neron-Popescu desingularization*, in “Algebra and Geometry”, Ed. M. Kang, International Press, Cambridge, (1998), 135–192.

DORIN POPESCU, SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, RESEARCH UNIT 5, UNIVERSITY OF BUCHAREST, P.O.Box 1-764, BUCHAREST 014700, ROMANIA

E-mail address: `dorin.popescu@imar.ro`