

# AUTOMATING THE CALCULATION OF HILBERT-KUNZ MULTIPLICITIES AND F-SIGNATURES

GABRIEL JOHNSON AND SANDRA SPIROFF

ABSTRACT. The Hilbert-Kunz multiplicity and  $F$ -signature are important invariants for researchers in commutative algebra and algebraic geometry. We provide software, and describe the automation of a calculation, for the two invariants in the case of intersection algebras over polynomial rings.

## Code Metadata

C1	Current code version	v1
C2	Permanent link to code/repository used for this code version	<a href="https://gitlab.com/gsjohnso/hilbert-kunz-and-f-sig">https://gitlab.com/gsjohnso/hilbert-kunz-and-f-sig</a>
C3	Legal Code License	BSD 3-Clause
C4	Code versioning system used	Git
C5	Software code languages, tools, and services used	Common Lisp
C6	Compilation requirements, operating environments & dependencies	Clozure Common Lisp, Bash/Cygwin, Sed, Mathematica (optional); Linux, Mac, Windows
C7	If available Link to developer documentation/manual	
C8	Support email for questions	<a href="mailto:gsjohnso@go.olemiss.edu">gsjohnso@go.olemiss.edu</a>

## 1. MOTIVATION AND SIGNIFICANCE

The Hilbert-Kunz multiplicity, along with the  $F$ -signature, has much importance in the related fields of commutative algebra and algebraic geometry, specifically, in characteristic  $p > 0$  methods, but is notoriously difficult to compute in practice. (The article [4] by C. Huneke provides a comprehensive overview of results on these two invariants.) In particular, very few examples exist in the literature where both values are known simultaneously. This state of affairs motivated the work of F. Enescu and S. Spiroff, who calculated the two invariants, as well as the Hilbert-Samuel multiplicity and divisor class group, for certain classes of intersection algebras [3]. Because, in their (toric) setting, the Hilbert-Kunz multiplicity and  $F$ -signature can be realized as volumes of polytopes [5, Theorem 2.2], [3, Proposition 4.2] and [6, Theorem 3.2.3], the calculation, being combinatorial in nature, lends itself perfectly to computer automation. One might expect to be able to obtain general formulæ for the invariants, however, a significant hurdle is the lack of a usable description of the unique Hilbert basis elements in terms of the parameters.

Let  $R$  be a commutative ring and  $I, J$  ideals of  $R$ . The intersection algebra of  $I$  and  $J$  is  $\mathcal{B} = \mathcal{B}_R(I, J) = \bigoplus_{r,s \in \mathbb{N}} (I^r \cap J^s)$ . In particular, when  $R = \mathbb{k}[x_1, \dots, x_n]$ , for  $\mathbb{k}$  a field, and  $I = (x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n})$  and  $J = (x_1^{b_1} x_2^{b_2} \cdots x_n^{b_n})$ , with  $a_i, b_i \in \mathbb{N}$ , the ordered pairs  $(b_i, a_i)$  partition the first quadrant of the plane into a fan made

---

*2010 Mathematics Subject Classification.* 68-04; 13-04; 15A39;  
S. Spiroff was supported by Simons Foundation Collaboration Grant #245926.

up of pointed rational cones, from which monoids, and Hilbert bases, are obtained. While for general  $n$  and  $\mathbf{a} = (a_1, \dots, a_n)$ ,  $\mathbf{b} = (b_1, \dots, b_n)$  there is no known formula for  $e_{HK}(\mathcal{B})$  or  $s(\mathcal{B})$ , for any specific numerical entries, Enescu and Spiroff have an algorithm to compute  $e_{HK}(\mathcal{B})$  and  $s(\mathcal{B})$ . This algorithm has been automated by G. Johnson. See Section 4 for obtaining and using the program.

Intersection algebras overlap with other classes of rings. For example, when  $R = \mathbb{k}[x]$ ,  $\mathcal{B}_R((x), (x))$  is a Segre product [8, §3], i.e., the homogeneous coordinate ring for the Segre embedding  $\mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^3$ , and  $\mathcal{B}_R((x^{a_1}), (x))$  is isomorphic to the rational normal scroll  $S = \mathbb{k}[T, xT, xyT, yT, x^{-1}yT, x^{-2}yT, \dots, x^{-(a_1-1)}yT]$  [9, Example 3.5]. For details on intersection algebras, especially background material, see [7, §2] and [2], and for the complete set of results obtained by Enescu and Spiroff, see [3]. References for geometric results are [4], [8], [9]. Throughout the paper,  $R$  will be a polynomial ring over a field and  $I, J$  principal monomial ideals. Since the numerical invariants in our case are given by volumes, they are independent of the characteristic of the field.

## 2. DESCRIPTION OF SOFTWARE

We extract the necessarily positive<sup>1</sup> exponents into two strings of positive integers,  $\mathbf{a} = a_1, \dots, a_n$  and  $\mathbf{b} = b_1, \dots, b_n$  since they may be treated without any reference to the ideals themselves. The command line interface can be run on a Unix shell, i.e. on Linux, Mac, or on Windows using Cygwin. For the parameters, the user need only enter the two lists of exponents delimited by spaces one after the other, with no delimiter marking the end of one list and the beginning of the next; the list of numbers entered is automatically cut in half and the first half assumed to be the exponents of one ideal and the second half the exponents of the other.

```
user directory$ ./calculate-integral a1 a2 ⋯ an b1 b2 ⋯ bn
```

The program displays the values of the Hilbert-Kunz multiplicity and the  $F$ -signature on two lines and terminates, as shown below.

```
Hilbert-Kunz Multiplicity      = p1/q1
F-Signature                    = p2/q2
```

When the `./inequalities` command is used, as shown below, the program displays the *Mathematica* code used to calculate the integrals.

```
user directory$ ./inequalities a1 a2 ⋯ an b1 b2 ⋯ bn
```

**2.1. Software architecture and functionality.** Three main technologies are used.

1. The Bash shell. This collects the arguments from the user and invokes the program itself.
2. Clozure Common Lisp. This is the programming language we used to calculate the set of inequalities bounding the solid in question. It provides a large number of

---

<sup>1</sup>The case of non-negative integers can be addressed within the scope of positive integers. See [3, Proposition 1.6].

list-processing operators as well as highly flexible looping constructs, making it well suited for dealing with the points of indefinite dimension involved in this project.

3. *Mathematica* [10]. This is used to calculate the final values of the integrals that give the Hilbert-Kunz multiplicity and  $F$ -signature, in any number of dimensions. The integration features of *Mathematica* are particularly well suited to this project, where the region of integration is defined by a set of inequalities in any number of dimensions. Other tools capable of solving integrals tend to require a function rather than a set of inequalities to define the area of integration, and limit the number of dimensions to two or three.

The terminal interface consists of a directory, called `ccl`, containing Clozure Common Lisp itself, a directory, called `src`, containing the Lisp source code as well as the Sed script used to format the output readably, and two bash scripts, `setup` and `calculate-integral`. The first, `setup`, must be run before attempting to run the other. It has Clozure compile the source code into an image file (effectively a DLL; in Lisp, all programs are essentially implemented as a DLL loaded by the language kernel with a specified top-level function to run instead of a REPL) called `hkm.image`. Once `hkm.image` exists in the same directory as the other files, `calculate-integral` can be run as specified above.

All the files and directories must occupy the same directory in order for the program to work. Further, the program also assumes that the Mathematica kernel, `MathKernel`, exists in a specified location depending on the operating system.

**2.2. Implementation details.** Below is a brief overview of the algorithm used in our program. Concurrently, we include a specific example to demonstrate the mathematics at each (hidden) step.

Recall that, to implement the program,  $a_1, a_2, \dots, a_n$  is entered as the exponents of one of the ideals, followed by  $b_1, b_2, \dots, b_n$  for the other, *with the lists, given or necessarily permuted, in fan order*. Once the arguments are collected from the command line, all steps but the last are done entirely in Lisp.

**Example 2.1.** For  $R = k[x, y]$  and  $I = (x^5y^2), J = (x^2y^3)$ , enter the command `./calculate-integral 5 2 2 3` in the Bash shell.

**Step 1.** Our program first finds the Hilbert set. As in [3, §1], [7], there are cones  $C_i$  and  $C$ , and monoids  $Q_i$  associated to the pairs  $(b_i, a_i)$ , and the unique Hilbert basis  $\mathcal{H}_i$  for each  $Q_i$  may be found via the algorithm given in [1, Algorithm 2.4].

**Definition 2.2.** [3, Definition 1.2] The set  $\mathcal{H} = \cup_{i=0, \dots, n} \mathcal{H}_i$  is called the **Hilbert set** for  $\mathcal{B}(I, J)$ . For  $v = (r, s) \in \mathcal{H}$ , let  $\mathbf{t}(v) = (\max(a_i r, b_i s))_{i=1, \dots, n}$ . Set  $\mathcal{G} = \{(v, \mathbf{t}(v)) : v \in \mathcal{H}\}$ .

e.g.,  $\mathcal{H}_0 = \{(0, 1), (1, 3), (2, 5)\}$ ,  $\mathcal{H}_1 = \{(1, 1), (1, 2), (2, 5), (3, 2)\}$ , and  $\mathcal{H}_2 = \{(1, 0), (2, 1), (3, 2)\}$ , hence  $|\mathcal{H}| = |\mathcal{H}_0 \cup \mathcal{H}_1 \cup \mathcal{H}_2| = 8$ .

**Step 2.** It finds the set  $\mathcal{G}$ . Each pair in the Hilbert set will be rewritten as an  $(n+2)$ -tuple. The extra  $n$  coordinates  $z_i$  are determined according to the formula  $z_i(x, y) = \max(a_i x, b_i y)$ , corresponding to the intersection of monomial ideals.

e.g.,  $\mathcal{G} = \{(1, 0, 5, 2), (0, 1, 2, 3), (2, 5, 10, 15), (3, 2, 15, 6), (2, 1, 10, 4), (1, 2, 5, 6), (1, 1, 5, 3), (1, 3, 6, 9)\}$

**Step 3.** It determines the set of inequalities bounding a solid in  $n+2$  dimensions. If the individual coordinates of each element of  $\mathcal{G}$  are labelled as  $(x, y, z_1, z_2, \dots, z_n)$ , then an initial set of inequalities is derived by a predetermined template:

$$\bigwedge_{i=1}^n 0 \leq a_i x \leq z_i \wedge 0 \leq b_i y \leq z_i.$$

From these initial inequalities and  $\mathcal{G}$ , further inequalities must be derived, and the conjunction of all these inequalities will produce the polytope we seek: an element  $(p, q, r_1, r_2, \dots, r_n) \in \mathcal{G}$  generates a set of further inequalities by the following function:

$$\neg \text{initialInequalities}(x - p, y - q, z_1 - r_1, z_2 - r_2, \dots, z_n - r_n),$$

where `initialInequalities` is a function that accepts an element of  $\mathcal{G}$  and returns true if and only if the given point satisfies all the initial inequalities. The program repeats this process for every element of  $\mathcal{G}$ . Then, it joins all the sets of inequalities by conjunction, producing our polytope. Its volume is the Hilbert-Kunz multiplicity [5, Theorem 2.2], [3, Proposition 4.2], and will appear in the first line of output.

```
e.g., Integrate[Boole[(x >= 0 && y >= 0 && z >= 5*x && z >= 2*y &&
w >= 2*x && w >= 3*y))
&& ((z < 5*x + 1 || z < 2*y + 1))
&& ((w < 2*x + 1 || w < 3*y + 1))
&& ((x < 1 || z < 2*y + 5 || w < 3*y + 2))
&& ((y < 1 || z < 5*x + 2 || w < 2*x + 3))
&& ((x < 2 || y < 5 || w < 2*x + 11))
&& ((x < 3 || y < 2 || z < 2*y + 11))
&& ((x < 2 || y < 1 || z < 2*y + 8 || w < 3*y + 1))
&& ((x < 1 || y < 2 || z < 2*y + 1 || w < 2*x + 4))
&& ((x < 1 || y < 1 || z < 2*y + 3 || w < 2*x + 1))
&& ((x < 1 || y < 3 || z < 5*x + 1 || w < 2*x + 7))],
{x, 0, 100}, {y, 0, 100}, {z, 0, 500}, {w, 0, 500}] }
```

**Step 4.** It determines the inequalities for the  $F$ -signature, which is much simpler, by definition, as they follow the formula

$$\bigwedge_{i=1}^n a_i x \leq z_i \leq 1 + a_i x \wedge b_i y \leq z_i \leq 1 + b_i y.$$

This can be found with a simple for loop. The  $F$ -signature is the volume of this second polytope [6, Theorem 3.2.3], and will appear in the second line of output.

```
e.g., Integrate[Boole[5*x <= z < 1 + 5*x && 2*y <= z < 1 + 2*y && 2*x <= w < 1
+ 2*x && 3*y <= w < 1 + 3*y ], {x, 0, 1}, {y, 0, 1}, {z, 0, 25}, {w, 0, 15}]
```

If the `./inequalities` command was used, then the program terminates after displaying the inequalities. Otherwise, the program passes the complete sets of inequalities to *Mathematica*, which calculates the volumes of the corresponding solids. Sed is used to format the output readably.

```
Hilbert-Kunz Multiplicity    = 37283/9900
F-Signature                  = 1087/29700
```

Implementation of the `./inequalities` command allows the user to bypass the use of *Mathematica*. This allows one to obtain the inequalities in Steps 3 and 4. Not only is this useful for 3D graphing and printing, but the user may integrate or analyze via another program, especially if he/she does not have a local installation of *Mathematica*, a proprietary software, or only wants to study the  $F$ -signature. As evinced in the examples below, the running time in higher dimensions can significantly increase depending upon the complexity of the geometry involved. However, the bottleneck in these cases is the integration related to the Hilbert-Kunz multiplicity; the steps calculating the regions determined by the inequalities, and the integration related to the  $F$ -signature, finished almost instantaneously even in the most complex cases tested.

### 3. SOME EXAMPLES-POLYTOPES AND RUNNING TIMES

**Example 3.1.** (Volumes.) The 3D polytopes associated to the invariants for  $\mathcal{B}_R((x^3), (x^2))$ , where  $R = \mathbb{k}[x]$ , show the volumes calculated by the program.




FIGURE 1. **Hilbert Kunz Multiplicity:** view from the front and from the side,  $e_{HK}(\mathcal{B}) = \frac{41}{18}$ .




FIGURE 2.  **$F$ -Signature:** view from the front and from the side,  $s(\mathcal{B}) = \frac{11}{36}$ .

**Example 3.2.** (Fan order and Running times.) When  $R = \mathbb{k}[x, y, z]$  and  $I = (x^2y^4z^7)$ ,  $J = (xy^6z^5)$ , the calculation had a running time of approximately 3.5 hours on a laptop, although the inequalities command outputs almost immediately.

```
./calculate-integral 2 7 4 1 5 6
Hilbert-Kunz Multiplicity    = 1874881259711/391184640000
F-Signature                  = 27251293/1564738560
```

On the other hand, the example `./calculate-integral 1 6 5 1 6 5` executes in a matter of seconds; i.e.,  $e_{HK}(\mathcal{B}) = \frac{1633}{864}$ ,  $s(\mathcal{B}) = \frac{95}{864}$ , for  $\mathcal{B} = \mathcal{B}_R(J, J)$ .

This last calculation illustrates the fact that if  $a_i = b_i$  for all  $i$ , then  $e_{HK}(\mathcal{B}) + s(\mathcal{B}) = 2$ . (See [3, Proposition 4.3].)

#### 4. SOFTWARE AVAILABILITY AND DIRECTIONS FOR USE

The software may be obtained at either of the websites:

`http://home.olemiss.edu/~spiroff/`

or

`https://gitlab.com/gsjohnso/hilbert-kunz-and-f-sig`

##### One time steps

1. Download the .zip file and extract it into your chosen directory.
2. In a Bash shell, go to the chosen directory using the `cd` command(s) and run `./setup`.

Commands for regular use-in a Bash shell, in the chosen directory (in fan order):

`./calculate-integral a1 a2 ⋯ an b1 b2 ⋯ bn`

`./inequalities a1 a2 ⋯ an b1 b2 ⋯ bn`

*Remark 4.1.* The details behind this automation, and the mathematical theory regarding the computations, are in the paper *Computing the invariants of intersection algebras of principal monomial ideals*, by F. Enescu and S. Spiroff, International Journal of Algebra and Computation, to appear.

#### ACKNOWLEDGEMENTS

The authors thank Florian Enescu at Georgia State University, Dawn Wilkins and Conrad Cunningham in the Department of Computer Science at the University of Mississippi, and the various specialists in the Mississippi Center for Supercomputing Research on the Oxford campus for useful conversations about this project.

#### REFERENCES

- [1] T. Cortadellas Benítez, C. D’Andrea, F. Enescu, *On the resolution of fan algebras of principal ideals over Noetherian rings*, 2016, preprint.
- [2] F. Enescu and S. Malec, *Intersection algebras for principal monomial ideals in polynomial rings*, J. Alg. Appl. **14**, no. 7 (2015), 23 pp.
- [3] Enescu, F., Spiroff, S., *Computing The Invariants of Intersection Algebras of Principal Monomial Ideals*, International Journal of Algebra and Computation, to appear.
- [4] C. Huneke, *Hilbert-Kunz Multiplicity and the F-Signature*, Commutative Algebra: Expository Papers Dedicated to David Eisenbud on the Occasion of His 65th Birthday, 485-525, Springer, New York, 2013.
- [5] E. Kazufumi, *Multiplicity and Hilbert-Kunz Multiplicity of Monoid Rings*, Tokyo J. Math, **25** no. 2 (2002).

- [6] M. von Korff, *The F-signature of toric varieties*, University of Michigan, Ph.D. thesis, 2012, ISBN: 978-1267-71896-9.
- [7] S. Malec, *On the intersection algebra of principal ideals*, *Comm. Alg.*, **43** no. 2 (2015), 623-635.
- [8] K. Watanabe, *Hilbert-Kunz Multiplicity of Toric Rings*, *Inst. Nat. Sciences Nihon Univ.*, **35** (2000), 173-177.
- [9] K. Watanabe and K. Yoshida, *Hilbert-Kunz multiplicity of three dimensional local rings*, *Nagoya Math. J.*, **177** (2005) 47-75.
- [10] Wolfram Research, Inc., *Mathematica*, version 10.0.2, Champaign, Illinois (2014).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSISSIPPI, OXFORD, MS 38677

*E-mail address:* `gsjohnson@go.olemiss.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSISSIPPI, OXFORD, MS 38677

*E-mail address:* `spiroff@olemiss.edu`