

# ON THE VARIETY OF THE INFLECTION POINTS OF PLANE CUBIC CURVES

VIK.S. KULIKOV

ABSTRACT. In the paper, we investigate properties of the nine-dimensional variety of the inflection points of the plane cubic curves. The description of local monodromy groups of the set of inflection points near singular cubic curves is given. Also, it is given a detailed description of the normalizations of the surfaces of the inflection points of plane cubic curves belonging to general two-dimensional linear systems of cubic curves, The vanishing of the irregularity a smooth manifold birationally isomorphic to the variety of the inflection points of the plane cubic curves is proved.

## 0. INTRODUCTION

The study of the properties of inflection points of plane nonsingular cubic curves has a rich and long history. Already in the middle of the 19th century, O. Hesse ([8], [9]) proved that nine inflection points of a nonsingular plane cubic is a projectively rigid configuration of points, i.e. 9-tuples of inflection points of the nonsingular plane cubic curves constitute an orbit with respect to the action of the group  $PGL(3, \mathbb{C})$  on the set of 9-tuples of the points of the projective plane. A subgroup  $Hes$  in  $PGL(3, \mathbb{C})$  of order 216 leaving invariant the set of the inflection points of Fermat cubic, given in  $\mathbb{P}^2$  by  $z_1^3 + z_2^3 + z_3^3 = 0$ , was defined by K. Jordan ([11]) and named by him the *Hesse group*. The invariants of the group  $Hes$  were described by G. Mashke in [18]. A brief historical overview of the results relating to the inflection points of plane cubic curves can be found in [1].

Let  $F(\bar{a}, \bar{z}) = \sum_{0 \leq i+j \leq 3} a_{i,j} z_1^i z_2^j z_3^{3-i-j}$  be the homogeneous polynomial of degree three in variables  $z_1, z_2, z_3$  and of degree one in variables  $a_{i,j}$ ,  $0 \leq i + j \leq 3$ . Denote by  $\mathcal{C} \subset \mathbb{P}^9 \times \mathbb{P}^2$  the complete family of plane cubic curves given by equation  $F(\bar{a}, \bar{z}) = 0$ . Let  $\kappa : \mathcal{C} \rightarrow \mathbb{P}^9$  and  $h : \mathcal{I} = \mathcal{C} \cap \mathcal{H} \rightarrow \mathbb{P}^9$  be the restrictions of the projection  $\kappa_1 : \mathbb{P}^9 \times \mathbb{P}^2 \rightarrow \mathbb{P}^9$  to  $\mathcal{C}$  and  $\mathcal{I}$ , where

$$\mathcal{H} = \{(\bar{a}, \bar{z}) \in \mathbb{P}^9 \times \mathbb{P}^2 \mid \det\left(\frac{\partial^2 F(\bar{a}, \bar{z})}{\partial z_i \partial z_j}\right) = 0\}.$$

It is well known (see, for example, [2]), that for a generic point  $\bar{a}_0 \in \mathbb{P}^9$  the intersection of the curve  $C_{\bar{a}_0} = \kappa^{-1}(\bar{a}_0)$  and its Hesse curve  $H_{C_{\bar{a}_0}}$ , given by  $\det\left(\frac{\partial^2 F(\bar{a}_0, \bar{z})}{\partial z_i \partial z_j}\right) = 0$ , is the 9 inflection points of  $C_{\bar{a}_0}$ . Therefore,  $\deg h = 9$ .

Let  $\mathcal{B} \subset \mathbb{P}^9$  be the discriminant hypersurface consisting of points  $\bar{a}$  such that the curves  $C_{\bar{a}}$  are singular. Then  $h : \mathcal{I} \setminus h^{-1}(\mathcal{B}) \rightarrow \mathbb{P}^9 \setminus \mathcal{B}$  is an unbranched covering of degree nine and, therefore, this covering determines a homomorphism  $h_* : \pi_1(\mathbb{P}^9 \setminus \mathcal{B}, \bar{a}_0) \rightarrow \mathbb{S}_9$  (here  $\mathbb{S}_9$  is a symmetric group acting on the set  $I_{\bar{a}} = C_{\bar{a}_0} \cap \mathcal{I}$ ). The group  $\mathcal{G} = \text{Im } h_*$  is called *the monodromy group of the inflection points of plane cubic curves*. In [7] has been proved

**Theorem 1.** *The group  $\mathcal{G}$  is a group of order 216. It is isomorphic to the group  $Hes$  of projective transformations of the plane leaving invariant the set of inflection points of the Fermat curve of degree three.*

In this paper, we continue the study of the properties of the varieties of inflection points of plane curves begun in [17]. In §1 we recall known facts about fundamental groups of complements to subvarieties of codimension one and on monodromy groups of dominant morphisms. In §2 we give a description of the local monodromy groups of the set of inflection points of plane cubic curves near the points  $\bar{a} \in \mathbb{P}^9$  parametrizing singular cubic curves (Propositions 3 – 5). In §3 and §4, we give a detailed description of the normalizations of the surfaces of the inflection points of plane cubic curves belonging to generic two-dimensional linear systems of cubic curves, and also prove the main result of this paper (Theorem 4), namely, we prove the vanishing of the irregularity of smooth varieties birationally isomorphic to the variety  $\mathcal{I}$ .

## 1. THE COVERING MONODROMY

**1.1. On the fundamental groups of complements to subvarieties of codimension one.** Let  $B \subset Z$  be a reduced closed codimension one subvariety of a simply connected smooth manifold  $Z$ ,  $\dim Z = K$ . It is well known that the fundamental group  $\pi_1(Z \setminus B, p)$  is generated by so-called *geometric generators* (or *bypasses* around  $B$ )  $\gamma_q$ , i.e., elements represented by loops  $\Gamma_q$ ,  $q \in B \setminus \text{Sing } B$ , of the following kind. Let  $L \subset Z$  be a germ of a smooth curve crossing transversely the variety  $B$  at  $q \in B \setminus \text{Sing } B$ , and let  $S_1 \subset L$  be a small radius circle centered at  $q$ . Right orientation of  $Z$ , defined by a complex structure, determines the orientation on  $S_1$  and in this case,  $\Gamma_q$  is a loop consisting of the path  $l$  lying in  $Z \setminus B$  and connecting the point  $p$  with some point  $q_1 \in S_1$ , the loop  $S_1$  (with the right orientation) starting and ending at the point  $q_1$ , and return to the point  $p$  along the path  $l$ . (Of course, the geometric generators  $\gamma_q$  depend not only on the choice of the point  $q$ , but also on the choice of the path  $l$ ; if we choose different paths and different points, the geometric generators  $\gamma_q$  are conjugated to each other in  $\pi_1(Z \setminus B, p)$  for points  $q$  belonging to the same irreducible component of the variety  $B$ .)

In the case when  $B$  is an hypersurface in  $\mathbb{P}^k$  and  $\mathbb{P}^n$  is a  $n$ -plane in  $\mathbb{P}^k$ , which is in general position with respect to  $B$ , then Zariski – van Kampen Theorem states that the homomorphism  $i_* : \pi_1(\mathbb{P}^n \setminus B) \rightarrow \pi_1(\mathbb{P}^k \setminus B)$ , induced by inclusion  $i : \mathbb{P}^n \hookrightarrow \mathbb{P}^k$  is an isomorphism when  $n \geq 2$  and epimorphism when  $n = 1$ . In the case  $n = 1$ , the group  $\pi_1(\mathbb{P}^1 \setminus B, p)$  is generated by  $d = \deg B$  bypasses  $\gamma_1, \dots, \gamma_d$  in  $\mathbb{P}^1 \setminus B$  around the

points belonging to the set  $\{q_1, \dots, q_d\} = B \cap \mathbb{P}^1$ . An ordered set  $\{i_*(\gamma_1), \dots, i_*(\gamma_d)\}$  is called a *good geometric base* of the group  $\pi_1(\mathbb{P}^k \setminus B, p)$  if the product  $\gamma_1 \cdot \dots \cdot \gamma_d$  is equal to the neutral element in  $\pi_1(\mathbb{P}^1 \setminus b, p)$ .

Let  $m = m_o(B)$  be the *multiplicity of singularity* of the hypersurface  $B$  at the point  $o$  and let  $z_1, \dots, z_k$  be local analytic coordinates in a certain analytic neighborhood  $V \subset Z$  of the point  $o = (0, \dots, 0) \in B$  chosen so that the intersection number  $(L_0, B)_o$  at the point  $o$  of the "line"  $L_0 \subset V$ , given by  $z_1 = \dots = z_{k-1} = 0$ , and  $B$  is equal to  $m$ .

Let

$$\Delta_{\bar{\delta}}^k = \{(z_1, \dots, z_k) \in V \mid |z_i| < \delta_i \text{ for } i = 1, \dots, k\} \subset Z$$

be a polydisk of multiradius  $\bar{\delta} = (\delta_1, \dots, \delta_k) > 0$  centered at the point  $o$ . Each time, we assume that multiradius  $\bar{\delta} = (\delta_1, \dots, \delta_k)$  is chosen so that the restriction of the projection  $\text{pr} : (z_1, \dots, z_{k-1}, z_k) \mapsto (z_1, \dots, z_{k-1})$  to  $\Delta_{\bar{\delta}}^k \cap B$  is a proper finite map of degree  $m$ . Let a "line"  $L \subset \Delta_{\bar{\delta}}^k$ , defined by equations  $z_1 = c_1 < \delta_1, \dots, z_{k-1} = c_{k-1} < \delta_{k-1}$  intersect  $B$  in  $m$  different points  $q_1, \dots, q_m$  and let  $p = (c_1, \dots, c_{k-1}, c_k) \in L \setminus B$ . The fundamental group  $\pi_1(L \setminus B, p)$  is the free group of rank  $m$  and it is generated by  $m$  bypasses  $\gamma_1, \dots, \gamma_m$  around the points  $q_1, \dots, q_m$ . An ordered set  $\{\gamma_1, \dots, \gamma_m\}$  is called a *good geometric base* of  $\pi_1(L \setminus B, p)$  if the product  $\gamma_1 \cdot \dots \cdot \gamma_m$  is equal to the element in  $\pi_1(L \setminus B, p)$  represented by the circuit along the circle  $\Delta = \{|z_k| = \delta_k\}$ . The ordered set  $\{i_*(\gamma_1), \dots, i_*(\gamma_m)\}$  is also called a *good geometric base* of the group  $\pi_1(\Delta_{\bar{\delta}}^k \setminus B, p)$ . It is easy to show that the elements of a good geometric base generate the group  $\pi_1(\Delta_{\bar{\delta}}^k \setminus B, p)$ .

For multiradii  $\bar{\delta}_1$  and  $\bar{\delta}_2$ ,  $\bar{\delta}_1 \leq \bar{\delta}_2$ , the imbedding  $\Delta_{\bar{\delta}_1}^k \subset \Delta_{\bar{\delta}_2}^k$  induces a homomorphism  $i_* : \pi_1(\Delta_{\bar{\delta}_1}^k \setminus B) \rightarrow \pi_1(\Delta_{\bar{\delta}_2}^k \setminus B)$  of the fundamental groups. The following theorem is well known (see, for example, [5]).

**Theorem 2.** *There is such a multiradius  $\bar{r} > 0$  that for any multiradius  $\bar{\delta} \leq \bar{r}$  the homomorphism  $i_* : \pi_1(\Delta_{\bar{\delta}}^k \setminus B) \rightarrow \pi_1(\Delta_{\bar{r}}^k \setminus B)$ , induced by the polydisk embedding is an isomorphism.*

The group  $\pi_1^{\text{loc}}(B, o) := \pi_1(\Delta_{\bar{r}}^k \setminus b)$  is called *local fundamental group* of  $B$  at  $o$ .

Let  $\Pi$  be a linear subspace of  $\Delta_{\bar{\delta}}^k$ ,  $\dim \Pi = n$ , passing through  $o$  and being at that point in general position with respect to  $b$ . The following theorem is a direct consequence of the results of [6].

**Theorem 3.** *The homomorphism  $i_* : \pi_1^{\text{loc}}(B \cap \Pi, o) \rightarrow \pi_1^{\text{loc}}(B, o)$ , induced by an embedding  $i : \Pi \hookrightarrow \Delta_{\bar{\delta}}^k$ , is an isomorphism for  $n \geq 3$  and an epimorphism for  $n = 2$ .*

The following claims are well known.

**Claim 1.** *Let the curve  $B \subset \Delta_{\bar{r}}^2$  is given by  $z_1^3 + z_2^2 = 0$ , i.e.,  $B$  has a singularity of type  $A_2$  at the point  $o$ . Then the group  $\pi_1^{\text{loc}}(B, o)$  has the following presentation:*

$$\pi_1^{\text{loc}}(B, o) = \langle \gamma_1, \gamma_2 : \gamma_1 \gamma_2 \gamma_1 = \gamma_1 \gamma_2 \gamma_1 \rangle, \quad (1)$$

where  $\{\gamma_1, \gamma_2\}$  is a good geometric base of  $\pi_1(\Delta_r^2 \setminus B)$ .

**Claim 2.** Let  $B \subset \Delta_r^k$  be given by equation  $z_1 \dots z_n = 0$ ,  $n \leq k$ , i.e.,  $B$  is a divisor with normal crossings at  $o$ . Then the group  $\pi_1^{loc}(B, o) \simeq \mathbb{Z}^m$  is a free abelian group and it is generated by bypasses  $\gamma_i$ ,  $i = 1, \dots, m$ , around hypersurfaces  $\{z_i = 0\}$ .

**Lemma 1.** ([16]) Let  $(B, o)$  be a germ of  $B$  in  $\Delta_r^2$  and let  $\sigma : \widetilde{\Delta}_r^2 \rightarrow \Delta_r^2$  be  $\sigma$ -process with center at  $o$ , and  $E = \sigma^{-1}(o)$  its exceptional divisor. Then the conjugacy class in the group  $\pi_1(\widetilde{\Delta}_r^2 \setminus \sigma^{-1}(B), \sigma^{-1}(p)) = \pi_1(\Delta_r^2 \setminus B, p)$  of the bypass  $\gamma$  around  $E$  contains the element  $\Delta = \gamma_1 \cdot \dots \cdot \gamma_m$ , where  $\{\gamma_1, \dots, \gamma_m\}$  is a good geometric base of the group  $\pi_1(\Delta_r^2 \setminus B, p)$ .

**1.2. On the monodromy of dominant morphisms.** Let  $f : X \rightarrow Z$  be a dominant proper holomorphic map of a complex-analytic variety  $X$  on smooth manifold  $Z$ ,  $\dim X = \dim Z = k$ . In this case, there is a subvariety  $B \subset Z$  of codimension one (which we call the *discriminant* of  $f$ ) such that  $f : Y = X \setminus f^{-1}(B) \rightarrow Z \setminus B$  is a finite unramified covering. Note that  $Y$  is a smooth manifold.

Let the degree of  $f : Y \rightarrow Z \setminus B$  be  $n$ . The map  $f$  defines a homomorphism  $f_* : \pi_1(Z \setminus B, p) \rightarrow \mathbb{S}_n$  (called the *monodromy* of  $f$ ) whose image  $G_f := f_*(\pi_1(Z \setminus B, p))$  is called the *monodromy group* of  $f$  and it is a subgroup of the symmetric group  $\mathbb{S}_n$  acting on the fibre  $f^{-1}(p) = \{p_1, \dots, p_n\}$  in the following way. A loop  $\Gamma \subset Z \setminus B$  representing an element  $\gamma \in \pi_1(Z \setminus B, p)$  can be lifted to  $Y$  and as a result, we get  $n$  paths  $\Gamma_1, \dots, \Gamma_n \subset Y$  starting and ending at the points of the fibre  $f^{-1}(p)$ . Therefore, this lift defines the action of  $f_*(\gamma)$  on  $f^{-1}(p)$ , which for each  $i = 1, \dots, n$  maps the starting point  $p_i$  of the path  $\Gamma_i$  to the end point of that path.

By Grauert - Remmert - Riemann - Stein Theorem ([21]) each epimorphism  $\varphi_* : \pi_1(Z \setminus B) \rightarrow G \subset \mathbb{S}_n$  uniquely determines a finite covering  $\varphi : X \rightarrow Z$  (which we call the *Stein covering* associated to  $\varphi_*$ ) of degree  $\deg \varphi = n$ , unramified over  $Z \setminus B$ , where  $X$  is a normal variety, and the group  $G \subset \mathbb{S}_n$  is the monodromy of  $\varphi$ , and if  $\varphi_* = f_*$  and  $Y$  is a normal variety, then there is a dominant map  $\psi : Y \rightarrow X$  (the factor map contracting the connected components of the fibres of  $f$  at points of  $X$ ) such that  $f = \varphi \circ \psi$ . In the case when  $n = |G|$  and the embedding  $G \subset \mathbb{S}_{|G|}$  is the Cayley embedding, i.e.,  $G$  acts on itself by multiplication on the right, the group  $G$  acts on  $X$  such that  $X/G = Z$  and in this case  $f$  is the *Galois covering*. The following proposition is true.

**Proposition 1.** If  $G \subset \mathbb{S}_n$  is the monodromy group of Stein covering  $\varphi : X \rightarrow Z$  and  $\mathbb{S}_n$  acts on the set  $\{q_1, \dots, q_n\}$ , then for the Galois covering  $\tilde{\varphi} : Y \rightarrow Z$  with the Galois group  $G$  there is a Galois covering  $\bar{\varphi} : Y \rightarrow X$  with Galois group  $G^1 = \{g \in G \mid g(q_1) = q_1\}$  such that  $\tilde{\varphi} = \varphi \circ \bar{\varphi}$ .

In the notation used in subsection 1.1, let  $o$  be a point in a hypersurface  $B$ . The choice of the path connecting the base points of the fundamental groups determines homomorphisms  $i_* : \pi_1^{loc}(B, o) \rightarrow \pi_1(Z \setminus B)$  and  $f_{*,loc} = f_* \circ i_* : \pi_1^{loc}(B, o) \rightarrow G$  defined uniquely up to conjugation in the group  $G$ . The group  $G_o := \text{Im } f_{*,loc} \subset G \subset \mathbb{S}_n$  is called the *local monodromy group* of  $f$  at  $o$ .

The following claim directly follows from the proof of Grauert - Remmert - Riemann - Stein Theorem.

**Claim 3.** *Let  $J_1, \dots, J_m$  be orbits of the action of the local monodromy group  $G_o \subset \mathbb{S}_n$  of Stein covering  $\varphi$  on the set  $\{1, \dots, n\}$ , where  $n_j$  is the cardinality of the orbit  $J_j$ ,  $n_1 + \dots + n_m = n$ . Then the preimage  $\varphi^{-1}(V)$  of a sufficiently small neighborhood  $V \subset Z$  of the point  $o$  splits into a disjoint union of  $m$  irreducible normal neighborhoods  $U_j$ . The degree of restriction of covering  $\varphi$  to  $U_j$  at  $q_j = U_j \cap \varphi^{-1}(o)$  is equal to  $n_j$ . In particular, if  $n_j = 1$ , then  $U_j$  is nonsingular at  $q_j$ .*

*If  $\varphi$  is a Galois cover and if the neighborhood of  $V$  is small enough, then  $\varphi^{-1}(V) = \bigsqcup_{j=1}^{i_o} U_j$  is a disjoint union of  $i_o$  connected normal pairwise biholomorphic surfaces (possibly having a singular point  $q_j \in U_j$ ), where  $i_o = (G : G_o)$  is the index of the subgroups  $G_o$  in  $G$ . The restriction of covering  $\varphi$  for each neighborhood  $U_j$  is a Galois covering with Galois group conjugated with the group  $G_o$ .*

Below we will need the following

**Claim 4.** *Let  $f : U \rightarrow \Delta_r^2$  be a proper dominant holomorphic map of degree three of a smooth surface  $U$  to the bidisk  $\Delta_r^2$  ramified in coordinate axes  $B = \{z_1 z_2 = 0\}$  and whose fibre  $L = f^{-1}(o) \simeq \mathbb{P}^1$  is a nonsingular rational curve. Assume that the images  $f_*(\gamma_1)$  and  $f_*(\gamma_2)$  of geometric generators  $\gamma_1$  and  $\gamma_2$  of the group  $\pi_1(\Delta_r^2 \setminus B)$  in the monodromy group  $G \subset \mathbb{S}_3$  of  $f$  are mutually inverse elements in  $G \simeq \mathbb{Z}_3$ . Then the self-intersection number of the curve  $L$  in  $U$  is  $(L^2)_U = -3$ .*

*Proof.* Let  $\varphi : V \rightarrow \Delta_r^2$  be the Stein covering associated with epimorphism  $f_*$ ,  $\deg \varphi = 3$ . Then the factorization map  $\psi : U \rightarrow V$  is the resolution of the singular point  $o' = \varphi^{-1}(o)$  of  $V$ , where  $o = (0, 0)$  is the origin in  $\Delta_r^2$ . On the other hand, let  $\sigma : \tilde{\Delta}_r^2 \rightarrow \Delta_r^2$  be the  $\sigma$ -process centered at  $o$ ,  $E = \sigma^{-1}(o)$ . Let  $\phi : W \rightarrow \tilde{\Delta}_r^2$  be the Stein covering associated with epimorphism  $f_* : \pi_1(\Delta_r^2 \setminus B) = \pi_1(\tilde{\Delta}_r^2 \setminus (\sigma^{-1}(B) \cup E)) \rightarrow \mathbb{S}_3$ . The covering  $\phi$  is ramified with multiplicity three only along the smooth curve  $\sigma^{-1}(B)$ , the proper preimage of the curve  $B$ , since  $f_*(\gamma) = Id$  according to Lemma 1, where  $\gamma$  is a bypass around the curve  $E$ . Hence,  $W$  is a nonsingular manifold and it is easy to see that  $L_1 = \phi^{-1}(E)$  is a smooth irreducible curve,  $(L_1^2)_W = \deg \phi \cdot (E^2)_{\tilde{\Delta}_r^2} = -3$ . According to Grauert - Remmert - Riemann - Stein Theorem, for  $\sigma \cdot \phi : W \rightarrow \Delta_r^2$  there is a canonical factorization  $\psi_1 : W \rightarrow V$ , also being the resolution of the singular point  $o' \in V$ . Therefore Claim 4 follows from the uniqueness of the minimal resolution of singular points of two-dimensional complex varieties.  $\square$

2. LOCAL MONODROMY GROUPS OF MORTHISM  $h$ 

**2.1. On the inflection points of plane cubic curves.** Consider a cubic curve  $C_{\bar{a}_0} \subset \mathbb{P}^2$  given by equation  $F(\bar{a}_0, \bar{z}) = 0$  and its Hessian curve  $H_{C_{\bar{a}_0}} \subset \mathbb{P}^2$  given by equation  $\det\left(\frac{\partial^2 F(\bar{a}_0, \bar{z})}{\partial z_i \partial z_j}\right) = 0$ . By definition, the *inflection points* of  $C_{\bar{a}_0}$  are the points belonging to the intersection  $C_{\bar{a}_0} \cap H_{C_{\bar{a}_0}}$ .

**Claim 5.** *The singular points of a cubic curve  $C_{\bar{a}_0}$  are its inflection points.*

*Proof.* If  $s = (0, 0, 1)$  is a singular point of a cubic curve  $C_{\bar{a}_0}$ , then  $C_{\bar{a}_0}$  is given by equation of the form  $z_3 \sum_{i+j=2} c_{i,j} z_1^i z_2^j + \sum_{i+j=3} c_{i,j} z_1^i z_2^j = 0$ . It is easy to see that the elements of the third column of the Hesse matrix on the left side of this equation vanish at  $s$ .  $\square$

**Claim 6.** *The points of a line  $L$  being a component of a cubic curve  $C_{\bar{a}_0}$  are inflection points of  $C_{\bar{a}_0}$ .*

*Proof.* Let the line  $L$  be given by equation  $z_1 = 0$ . In this case, the cubic curve  $C_{\bar{a}_0}$  is given by equation of the form  $z_1 \sum_{0 \leq i+j \leq 2} c_{i,j} z_1^i z_2^j z_3^{2-i-j} = 0$ . It is easy to see that only the elements of the first column and the first row of the Hesse matrix of the left side of this equation can have non-zero values at the points of  $L$ . Therefore, Hessian vanishes at the points of  $L$ .  $\square$

**2.2. Equisingular stratification of the variety of singular cubic curves.** Denote  $\mathcal{S} := \{(\bar{a}, \bar{z}) \in \mathbb{P}^9 \times \mathbb{P}^2 \mid \bar{z} \in \text{Sing } C_{\bar{a}}\}$ . We have  $\text{pr}_1(\mathcal{S}) = \mathcal{B}$ . The discriminant  $\mathcal{B} \subset \mathbb{P}^9$  is irreducible [10] and it has the natural stratification:

$$\begin{array}{ccccccc} & & & & \mathcal{B}_{3,1} & \longrightarrow & \mathcal{B}_{2,1} \\ & & & \nearrow & & & \searrow \\ \mathcal{B}_7 & \longrightarrow & \mathcal{B}_5 & \longrightarrow & \mathcal{B}_4 & & \mathcal{B}_1 \subset \mathcal{B} \\ & & & \searrow & \nearrow & & \nearrow \\ & & & & \mathcal{B}_{3,2} & \longrightarrow & \mathcal{B}_{2,2} \end{array}$$

where  $\mathcal{B}_7$  is the variety parametrizing the cubic curves consisting of triple lines;  $\mathcal{B}_5$  is the variety parametrizing the cubic curves consisting of two lines, one of which is included with multiplicity two;  $\mathcal{B}_4$  is the variety parametrizing the cubic curves consisting of three lines with a common point;  $\mathcal{B}_{3,1}$  is the variety parametrizing the cubic curves consisting of three lines in general position;  $\mathcal{B}_{3,2}$  is the variety parametrizing the cubic curves consisting of smooth conics and lines touching each other;  $\mathcal{B}_{2,1}$  is the variety parametrizing the cubic curves consisting of transversely intersecting conics and lines;  $\mathcal{B}_{2,2}$  is the variety parametrizing the cuspidal rational cubic curves;  $\mathcal{B}_1$  is the variety parametrizing the nodal rational cubic curves; arrows denote the adjoint of the strata and for the strata  $\mathcal{B}_i$  and  $\mathcal{B}_{i,j}$  the index  $i$  is equal to the codimension of these strata in  $\mathbb{P}^9$ . Note that all strata are smooth manifolds and, moreover, these strata are orbits of the action of the group  $\text{PGL}(3, \mathbb{C})$  on the space of plane cubic curves. Therefore,  $\mathcal{B}$  is an irreducible variety and therefore all geometric generators

of the groups  $\pi_1(\mathbb{P}^9 \setminus \mathcal{B})$  belong to the same conjugacy class of elements of this group and generate it. In addition, we have the following

- Claim 7.** (i) For any two points  $\bar{a}_1$  and  $\bar{a}_2$  belonging to the same stratum  $\mathcal{B}'$  of the equisingular stratification of variety  $\mathcal{B}$ , the local monodromy groups  $\mathcal{G}_{\bar{a}_1}$  and  $\mathcal{G}_{\bar{a}_2}$  are conjugated in  $\mathcal{G}$ .
- (ii) If a stratum  $\mathcal{B}'$  adjoins to a stratum  $\mathcal{B}''$ , then the local monodromy group  $\mathcal{G}_{\bar{a}_1}$  at  $\bar{a}_1$  contains a subgroup conjugated in  $\mathcal{G}$  with the local monodromy groups  $\mathcal{G}_{\bar{a}_2}$  at the points  $\bar{a}_2 \in \mathcal{B}''$ .

*Proof.* The group  $\text{PGL}(3, \mathbb{C})$  is a connected complex manifold. Therefore claim (i) follows from that each stratum  $\mathcal{B}'$  is an orbit under the action of  $\text{PGL}(3, \mathbb{C})$  on the variety of plane cubic curves.

Claim (ii) follows from that for any sufficiently small neighbourhood  $V_1 \subset \mathbb{P}^9$  of a point  $\bar{a}_1 \in \mathcal{B}'$  there is a point  $\bar{a} \in \mathcal{B}''$  lying in  $V_1$  together with some its neighbourhood  $V$ .  $\square$

The space  $\mathbb{P}^9$  has a natural covering by ten affine spaces  $\mathbb{C}_{i,j}^9 = \{\bar{a} \in \mathbb{P}^9 \mid a_{i,j} \neq 0\}$ ,  $0 \leq i + j \leq 3$ . The coordinates of the points  $\bar{a}$  in each space  $\mathbb{C}_{i_0, j_0}^9$  are the coefficients in equations

$$\sum_{0 \leq i+j \leq 3} a_{i,j} z_1^i z_2^j z_3^{3-i-j} = 0$$

of plane cubic curves normalized by condition  $a_{i_0, j_0} = 1$ .

Properties of the discriminant  $\mathcal{B}$  at the points corresponding the nodal cubic curves were investigated in [19].

**Proposition 2.** ([19]) *Let  $C_{\bar{a}_0}$  be a nodal cubic curve. Then  $\mathcal{B}$  is a divisor with normal crossings in some analytic neighbourhood of the point  $\bar{a}_0$ .*

Below, to calculate the local monodromy groups of morphism  $h$  at the points corresponding to the nodal cubic curves, we need more detail description of the discriminant  $\mathcal{B}$  at these points. For this, consider a nodal cubic curve  $C_{\bar{a}_0}$  given in non-homogeneous coordinates  $x = \frac{z_1}{z_3}$ ,  $y = \frac{z_2}{z_3}$  in  $\mathbb{C}^2 \subset \mathbb{P}^2$  by equation

$$xy + \sum_{i+j=3} c_{i,j} x^i y^j = 0.$$

The point  $\bar{a}_0$  lies in the affine space  $\mathbb{C}_{1,1}^9$  and its coordinates  $a_{0,0}$ ,  $a_{1,0}$ , and  $a_{0,1}$  vanish. Obviously, the variety  $\mathcal{S} \cap (\mathbb{C}_{1,1}^9 \times \mathbb{C}^2)$  is given by equations

$$\begin{aligned} a_{0,0} + a_{1,0}x + a_{0,1}y + xy + a_{2,0}x^2 + a_{0,2}y^2 + \sum_{i+j=3} a_{i,j}x^i y^j &= 0 \\ a_{1,0} + y + 2a_{2,0}x + \sum_{i+j=3} ia_{i,j}x^{i-1}y^j &= 0 \\ a_{0,1} + x + 2a_{0,2}y + \sum_{i+j=3} ja_{i,j}x^i y^{j-1} &= 0 \end{aligned} \tag{2}$$

It follows from the type of equations (2) that the eight-dimensional variety  $\mathcal{S}$  is smooth at the point  $(\bar{a}_0, \bar{z}_0)$ , where  $\bar{z}_0 = (0, 0, 1)$  and for  $2 \leq i + j \leq 3$ ,  $(i, j) \neq (1, 1)$  the

functions  $a_{1,0}$ ,  $a_{0,1}$ , and  $a_{i,j} - c_{i,j}$  are local parameters in  $\mathcal{S}$  at  $(\bar{a}_0, \bar{z}_0)$ . Therefore the image  $\text{pr}_1(S \cap V) \subset \mathbb{C}_{1,1}^9$  is non-singular at  $\bar{a}_0$ , where  $V$  is some sufficiently small analytic neighbourhood in  $\mathbb{C}_{1,1}^9 \times \mathbb{C}^2$  of the point  $(\bar{a}_0, \bar{z}_0)$ . In addition, it is easy to see that  $a_{0,0} = 0$  is an equation of the tangent space of  $\text{pr}_1(S \cap V)$  at  $\bar{a}_0$ .

The strata  $\mathcal{B}_1$ ,  $\mathcal{B}_{2,1}$ , and  $\mathcal{B}_{3,1}$  of the discriminant hypersurface  $\mathcal{B}$  parametrize the nodal cubic curves. These strata are orbits in  $\mathbb{P}^9$  under the action of the group  $PGL(3, \mathbb{C})$ . Therefore, without loss of generality, we can assume that a nodal cubic  $C_{\bar{a}_0}$  is given by one of the following equations:

- 1)  $z_1 z_2 z_3 + z_1^3 + z_2^3 = 0$  if  $\bar{a}_0 \in \mathcal{B}_1$ ,
- 2)  $z_1 z_2 z_3 + z_1^3 = 0$  if  $\bar{a}_0 \in \mathcal{B}_{2,1}$ ,
- 3)  $z_1 z_2 z_3 = 0$  if  $\bar{a}_0 \in \mathcal{B}_{3,1}$ .

In all three cases, the point  $\bar{a}_0$  belongs to the affine space  $\mathbb{C}_{1,1}^9$  and it follows from the above that in case 1)  $\mathcal{B}$  is a nonsingular hypersurface at  $\bar{a}_0$ ; in case 2)  $\mathcal{B}$  locally is a union of two nonsingular at point  $\bar{a}_0$  hypersurfaces intersecting transversally at that point (tangent spaces to these hypersurfaces are given by equations  $a_{0,0} = 0$  and  $a_{0,3} = 0$ ); in case 3)  $\mathcal{B}$  is locally the union of three non-singular hypersurfaces at  $\bar{a}_0$  intersecting transversally at this point (the tangent space to these hypersurfaces is defined by the equations  $a_{0,0} = 0$ ,  $a_{0,3} = 0$ , and  $a_{3,0} = 0$ ).

**2.3. Properties of the monodromy group  $\mathcal{G} = Hes$  of morphism  $h$ .** The group *Hes* was originally defined as a subgroup of the group  $PGL(3, \mathbb{C})$  of linear transformations of  $\mathbb{P}^2$  leaving invariant the *Hesse pencil*, i.e., a one-dimensional linear system of plane cubic curves defined by equation

$$C_{(t_1, t_2)} : t_1(z_1^3 + z_2^3 + z_3^3) + t_2 z_1 z_2 z_3 = 0, \quad (t_1, t_2) \in \mathbb{P}^1, \quad (3)$$

The Hesse pencil has nine fixed points

$$\begin{aligned} q_1 &= (0, 1, -1), & q_4 &= (0, 1, -\omega), & q_7 &= (0, 1, -\omega^2), \\ q_2 &= (1, 0, -1), & q_5 &= (1, 0, -\omega^2), & q_8 &= (1, 0, -\omega), \\ q_3 &= (1, -1, 0), & q_6 &= (1, -\omega, 0), & q_9 &= (1, -\omega^2, 0), \end{aligned}$$

where  $\omega = e^{2\pi i/3}$  is a primitive cubic root of unity, and a direct checking shows that these nine points are inflection points of each of the nonsingular curves entering the Hesse pencil. Therefore, the group  $Hes \subset PGL(3, \mathbb{C})$  can be defined as a subgroup of projective transformations leaving invariant the set of inflection points of the Fermat cubic  $F = C_{(1,0)}$ .

The properties of the group *Hes* are well known (see, for example, [4]). Its order is equal to 216 and the action of the group *Hes* on the nine inflection points of the curve  $F$  defines an embedding  $Hes \subset \mathbb{S}_9$  such that *Hes* is a 2-transitive subgroup of the symmetric group  $\mathbb{S}_9$  generated by permutations  $g_0 = (1, 2, 4)(5, 6, 8)(3, 9, 7)$  and  $g_1 = (4, 5, 6)(7, 9, 8)$ . The subgroups  $Hes^i = \{g \in Hes \mid g(q_i) = q_i\}$ ,  $i = 1, \dots, 9$ , consisting of elements of *Hes* leaving fixed the point  $q_i$ , are conjugated to each other in the group *Hes* and they are isomorphic to the group  $SL(2, \mathbb{Z}_3)$ . Their order is

24. As a subgroup in  $\mathbb{S}_9$ , the group  $Hes^1$  is generated by permutations  $g_1$  and  $g_2 = g_0 g_1 g_0^{-1} = (2, 8, 5)(3, 6, 9)$  and as an abstract group, it has the following presentation

$$Hes^1 = \{g_1, g_2 \mid g_1 g_2 g_1 = g_2 g_1 g_2, g_1^3 = 1\}.$$

For further we note that the class in  $Hes^1$  of elements conjugated to  $g_1$  consists of four elements:  $g_1, g_2, g_1 g_2, g_1^{-1}, g_2 g_1 g_2^{-1}$ , and any pair of these four elements generates the group  $Hes^1$  and satisfies the relation  $xyx = yxy$ . Also note (see [20]) that the group  $Hes^1$  has an exact linear presentation in  $GL(2, \mathbb{C})$  (group no. 4 in [20], see also [15]), in which images of  $g_1$  and  $g_2$  are reflections.

The group  $Hes^1$  is naturally embedded in the symmetric group  $\mathbb{S}_8$  acting on  $\{q_2, \dots, q_8\}$ . We number (left) co-sets in  $Hes^1$  of its subgroup  $\langle g_1 \rangle$  generated by  $g_1$  as follows:

$$\begin{aligned} c_2 = \langle g_1 \rangle, & \quad c_3 = g_2^2 g_1 g_2^2 \langle g_1 \rangle, & \quad c_4 = g_1^2 g_2^2 \langle g_1 \rangle, & \quad c_5 = g_2^2 \langle g_1 \rangle, \\ c_6 = g_1 g_2^2 \langle g_1 \rangle, & \quad c_7 = g_1 g_2 \langle g_1 \rangle, & \quad c_8 = g_2 \langle g_1 \rangle, & \quad c_9 = g_1^2 g_2 \langle g_1 \rangle. \end{aligned}$$

The group  $Hes^1$  acts on the set  $\{c_2, \dots, c_9\}$ .

Denote  $Hes^{1,2} = \{g \in Hes^1 \subset \mathbb{S}_8 \mid g(q_2) = q_2\}$ .

**Claim 8.** *We have  $Hes^{1,2} = \langle g_1 \rangle$  and the action of the group*

$$Hes^1 = \langle g_1 = (4, 5, 6)(7, 9, 8), g_2 = (2, 8, 5)(3, 6, 9) \rangle \subset \mathbb{S}_8$$

*on the set  $\{c_2, \dots, c_9\}$  coincides with the action on the set  $\{q_2, \dots, q_9\}$ .*

*Proof.* Writing all 24 permutations belonging to the group  $Hes^1 \subset \mathbb{S}_8$ , it is easy to verify the validity of this statement.  $\square$

#### 2.4. On the local monodromy groups of the morphism $h$ at points corresponding to nodal cubic curves.

**Proposition 3.** (i) *The local monodromy groups of morphism  $h$  at the points  $\bar{a}_1 \in \mathcal{B}_1$  are abelian groups isomorphic to the group  $\mathbb{Z}_3$  and they are conjugated in  $Hes$  to the subgroup  $\mathcal{G}_1 \subset \mathbb{S}_9$  generated by permutation  $g_2 = (2, 8, 5)(3, 6, 9)$ .*

(ii) *The local monodromy groups of morphism  $h$  at the points  $\bar{a}_{2,1} \in \mathcal{B}_{2,1}$  are abelian groups isomorphic to the group  $\mathbb{Z}_3^2$  and they are conjugated in  $Hes$  to the subgroup  $\mathcal{G}_{2,1} \subset \mathbb{S}_9$  generated by permutations  $g_2 = (2, 8, 5)(3, 6, 9)$  and  $g_3 = (1, 4, 7)(3, 9, 6)$ .*

(iii) *The local monodromy groups of morphism  $h$  at points  $\bar{a}_{3,1} \in \mathcal{B}_{3,1}$  are abelian groups isomorphic to the group  $\mathbb{Z}_3^2$  and they are conjugated in  $Hes$  to the subgroup  $\mathcal{G}_{3,1} \subset \mathbb{S}_9$  generated by permutations  $g_2 = (2, 8, 5)(3, 6, 9)$  and  $g_3 = (1, 4, 7)(3, 9, 6)$ , i.e., coincide with the group  $\mathcal{G}_{2,1}$ .*

*Proof.* We start the proof with statement (iii). Without loss of generality, we can assume that a nodal curve  $C_{\bar{a}_{3,1}}$  is given by equation  $z_1 z_2 z_3 = 0$ .

Consider a three-dimensional family  $\mathcal{C}_{\Pi_1}$  of cubic curves defined by equation

$$z_1 z_2 z_3 + a_{3,0} z_1^3 + a_{0,3} z_2^3 + a_{0,0} z_3^3 = 0 \tag{4}$$

and its projection on the affine space  $\text{pr}_1(\mathcal{C}_{\Pi_1}) = \Pi_1 \simeq \mathbb{C}^3$  in  $\mathbb{C}_{1,1}^9 \subset \mathbb{P}^9$ . Note that  $(a_{3,0}, a_{0,3}, a_{0,0})$  are affine coordinates in  $\Pi_1$  and everyone can easily check that Hessian  $H_{\mathcal{C}_{\Pi_1}}$  of this family is given by equation

$$(6^3 a_{3,0} a_{0,3} a_{0,0} + 2) z_1 z_2 z_3 + 6(a_{3,0} z_1^3 + a_{0,3} z_2^3 + a_{0,0} z_3^3) = 0. \quad (5)$$

Therefore  $h^{-1}(\Pi_1)$  in  $\Pi_1 \times \mathbb{P}^2$  is given by

$$z_1 z_2 z_3 = a_{3,0} z_1^3 + a_{0,3} z_2^3 + a_{0,0} z_3^3 = 0.$$

Denote  $p = (\delta, \delta, \delta) \in \Pi_1$ , where  $0 < \delta \ll 1$ , and let  $C_p = \kappa^{-1}(p)$ . Note that the cubic curve  $C_p$  has the same inflection points as Fermat cubic has (see subsection 2.3).

From the results set forth in subsection 2.2, it follows that locally at the point  $\bar{a}_{3,1}$  with coordinates  $a_{3,0} = a_{0,3} = a_{0,0} = 0$  the space  $\Pi_1$  intersects transversally with irreducible branches of the hypersurface  $\mathcal{B}$  in some neighborhood of this point and the surface  $\mathcal{B} \cap \Pi_1$  is defined in  $\Pi_1$  by equation  $a_{3,0} a_{0,3} a_{0,0} = 0$ . According to Theorem 3 the group  $\pi_1^{\text{loc}}(\mathcal{B}, \bar{a}_{3,1})$  is generated by elements  $\gamma_j \in \pi_1(\Pi_1 \setminus B, p)$ ,  $j = 1, 2, 3$ , represented (see Claim 2) by loops

$$\begin{aligned} c_1(t) &= \{(\delta e^{2\pi i t}, \delta, \delta) \in \Pi_1 \mid t \in [0, 1]\}, \\ c_2(t) &= \{(\delta, \delta e^{2\pi i t}, \delta) \in \Pi_1 \mid t \in [0, 1]\}, \\ c_3(t) &= \{(\delta, \delta, \delta e^{2\pi i t}) \in \Pi_1 \mid t \in [0, 1]\}. \end{aligned}$$

The preimage  $h^{-1}(c_1)$  consists of nine paths

$$\begin{aligned} q_1(t) &= (c_1(t), (0, 1, -1)), & q_2(t) &= (c_1(t), (1, 0, -\lambda(t))), & q_3(t) &= (c_1(t), (1, -\lambda(t), 0)), \\ q_4(t) &= (c_1(t), (0, 1, -\omega)), & q_5(t) &= (c_1(t), (1, 0, -\lambda(t)\omega^2)), & q_6(t) &= (c_1(t), (1, -\lambda(t)\omega, 0)), \\ q_7(t) &= (c_1(t), (0, 1, -\omega^2)), & q_8(t) &= (c_1(t), (1, 0, -\lambda(t)\omega)), & q_9(t) &= (c_1(t), (1, -\lambda(t)\omega^2, 0)), \end{aligned}$$

where where  $\omega = e^{2\pi i/3}$  and  $\lambda(t) = e^{2\pi i t/3}$ , and therefore,  $h_*(\gamma_1) = (2, 8, 5)(3, 6, 9) = g_2 \in \mathcal{G}_{3,1} \subset \mathcal{S}_9$ .

The preimage  $h^{-1}(c_2)$  consists of nine paths

$$\begin{aligned} q_1(t) &= (c_2(t), (0, 1, -\lambda(t))), & q_2(t) &= (c_2(t), (1, 0, -1)), & q_3(t) &= (c_2(t), (1, -\lambda^2(t), 0)), \\ q_4(t) &= (c_2(t), (0, 1, -\lambda(t)\omega)), & q_5(t) &= (c_2(t), (1, 0, -\omega^2)), & q_6(t) &= (c_2(t), (1, -\lambda^2(t)\omega, 0)), \\ q_7(t) &= (c_2(t), (0, 1, -\lambda(t)\omega^2)), & q_8(t) &= (c_2(t), (1, 0, -\omega)), & q_9(t) &= (c_2(t), (1, -\lambda^2(t)\omega^2, 0)) \end{aligned}$$

and therefore,  $h_*(\gamma_2) = (1, 4, 7)(3, 9, 6) = g_3$ .

Similarly, the preimage  $h^{-1}(c_3)$  consists of nine paths

$$\begin{aligned} q_1(t) &= (c_3(t), (0, 1, -\lambda^2(t))), & q_2(t) &= (c_3(t), (1, 0, -\lambda^2(t))), & q_3(t) &= (c_3(t), (1, -1, 0)), \\ q_4(t) &= (c_3(t), (0, 1, -\lambda^2(t)\omega)), & q_5(t) &= (c_3(t), (1, 0, -\lambda^2(t)\omega^2)), & q_6(t) &= (c_2(t), (1, -\omega, 0)), \\ q_7(t) &= (c_3(t), (0, 1, -\lambda^2(t)\omega^2)), & q_8(t) &= (c_3(t), (1, 0, -\lambda^2(t)\omega)), & q_9(t) &= (c_2(t), (1, -\omega^2, 0)) \end{aligned}$$

and therefore,  $h_*(\gamma_3) = (1, 7, 4)(2, 5, 8) := g_4$ . It is easy to see that  $g_2 g_3 = g_4^{-1}$ . Therefore the local monodromy group of morphism  $h$  at  $\bar{a}_{3,1}$  is  $\mathcal{G}_{3,1} \simeq \mathbb{Z}_3^2$  generated in  $\mathcal{G} = \text{Hes}$  by permutations  $g_2$  and  $g_3$ .

To prove statements (i) and (ii), we can assume that  $\bar{a}_{2,1} \in \Pi_1$  has coordinates  $a_{3,0} = a_{0,3} = 0$  and  $a_{0,0} = \varepsilon$ , where  $0 < \varepsilon \ll \delta$ , i.e., the nodal cubic  $C_{\bar{a}_{2,1}}$  is given by equation  $z_1 z_2 z_3 + \varepsilon z_3^3 = 0$ . Then it is easy to see that the local fundamental group of morphism  $h$  at  $\bar{a}_{2,1}$  is generated by elements  $\gamma_2$  and  $\gamma_3$ . And in the case (i) we can assume that the point  $\bar{a}_1 \in \Pi_1$  has coordinates  $a_{3,0} = 0$  and  $a_{0,0} = a_{0,3} = \varepsilon$ , where

$0 < \epsilon \ll \delta$ , i.e. the nodal cubic  $C_{\bar{a}_{2,1}}$  is given by the equation  $z_1 z_2 z_3 + \epsilon z_2^3 + \epsilon z_3^3 = 0$ . Then it is easy to see that the local fundamental group of morphism  $h$  at  $\bar{a}_1$  is generated by  $\gamma_2$ .  $\square$

## 2.5. On the local monodromy groups of the morphism $h$ at points corresponding to cuspidal cubic curves.

**Proposition 4.** *The local monodromy groups  $\mathcal{G}_{\bar{a}}$  of morphism  $h$  at points  $\bar{a} \in \mathcal{B}_{2,2}$  are conjugated to the group  $Hes^1$ .*

*Proof.* Without loss of generality, we can assume that the cuspidal cubic curve  $C_{\bar{a}_0} \subset \mathbb{P}^2$  is given by equation  $z_1^3 + z_2^2 z_3 = 0$ . Its singular point  $s$  has coordinates  $(0, 0, 1)$  and the point  $p = (0, 1, 0)$  is its unique nonsingular inflection point.

The fibre  $h^{-1}(\bar{a}_0)$  of morphism  $h$  is the intersection of the fibre  $C_{\bar{a}_0}$  of morphism  $\kappa : \mathcal{C} \rightarrow \mathbb{P}^9$  and Hessian variety  $\mathcal{H}$ . Calculating Hessian of the polynomial  $z_1^3 + z_2^2 z_3$ , we find that its Hessian curve  $H_{C_{\bar{a}_0}} \subset \mathbb{P}^2$  given by equation  $z_1 z_2^2 = 0$ , and it is easy to see that the intersection numbers of the curves  $C_{\bar{a}_0}$  and  $H_{C_{\bar{a}_0}}$  at points  $p$  and  $s$  are respectively  $(C_{\bar{a}_0}, H_{C_{\bar{a}_0}})_p = 1$  and  $(C_{\bar{a}_0}, H_{C_{\bar{a}_0}})_s = 8$ . It follows that in some analytic neighborhood  $V \subset \mathcal{I}$  of the point  $(\bar{a}_0, p)$  the variety  $\mathcal{I}$  is nonsingular and the restriction of the morphism  $h$  to  $V$  is a biholomorphic isomorphism of the neighborhood  $V$  and its image  $h(V)$ . Therefore, at least one of the orbits of the action of the local monodromy group  $\mathcal{G}_{\bar{a}_0}$  of morphism  $h$  on the set  $\{q_1, \dots, q_9\}$  consists of one point and, hence,  $\mathcal{G}_{\bar{a}_0}$  is contained in one of the groups conjugated to the group  $Hes^1$ .

In subsection 3.2 it will be shown that the point  $\bar{a}_0 \in \mathcal{B} \cap \Pi$  of the intersection of the variety  $\mathcal{B}$  with a generic projective plane  $\Pi \subset \mathbb{P}^9$  passing through the point  $\bar{a}_0$  is an ordinary cusp of the curve  $\mathcal{B} \cap \Pi$ . The group  $\pi_1^{loc}(\mathcal{B} \cap \Pi, \bar{a}_0)$  (see Claim 1) is generated by the geometric generators  $\gamma_1$  and  $\gamma_2$  satisfying the relation  $\gamma_1 \gamma_2 \gamma_1 = \gamma_2 \gamma_1 \gamma_2$ , and the elements  $h_*(\gamma_1)$  and  $h_*(\gamma_2)$  generate the group  $\mathcal{G}_{\bar{a}_0}$ . From the properties of the group  $Hes^1$  (see subsection 2.3), it follows that if  $h_*(\gamma_1) \neq h_*(\gamma_2)$ , then the group  $\mathcal{G}_{\bar{a}_0}$  is conjugated in  $\mathcal{G}$  to the group  $Hes^1$ , or if  $h_*(\gamma_1) = h_*(\gamma_2)$  then  $\mathcal{G}_{\bar{a}_0}$  is the cyclic group of third order.

Let us show that the second alternative has no place. To do this, consider the family of cubic curves  $C_{\bar{a}_\tau}$  parameterized by the points of the circle  $S_1 = \{\bar{a}_\tau \in \mathbb{P}^9\}$ ,  $\tau = \delta e^{2\pi i t}$ , and given by equation

$$C_{\bar{a}_\tau} : z_1^3 + z_2^2 z_3 + \tau z_3^3 = 0, \quad (6)$$

where  $t \in [0, 1]$  and  $\delta$  is some positive real number. It is easy to see that Hessian variety  $H_{C_{\bar{a}_\tau}}$  of this family is given by equation

$$z_1(3\tau z_3^2 - z_2^2) = 0. \quad (7)$$

It follows from (6) and (7) that the preimage  $h^{-1}(S_1) \subset \mathcal{I}$  of the circle  $S_1$  consists of nine paths three of which are

$$\begin{aligned} l_1(t) &= \{(\bar{a}(t), \bar{z}(t)) \in \mathcal{I} \mid \bar{a}(t) = \bar{a}_{\tau(t)}, \bar{z}(t) = (0, 1, 0)\}, \\ l_j(t) &= \{(\bar{a}(t), \bar{z}(t)) \in \mathcal{I} \mid \bar{a}(t) = \bar{a}_{\tau(t)}, \bar{z}(t) = (0, \sqrt[2]{-\delta}e^{\pi i(t+j)}, 1)\}, \quad j = 2, 3, \end{aligned}$$

and the other six are

$$l_{j,k}(t) = \{(\bar{a}(t), \bar{z}(t)) \in \mathcal{I} \mid \bar{a}(t) = \bar{a}_{\tau(t)}, \bar{z}(t) = (-\sqrt[3]{4\delta}e^{2\pi i(t+j)/3}, \sqrt[2]{3\delta}e^{\pi i(t+k)}, 1)\},$$

$j = 0, 1, 2$   $k = 0, 1$ . If  $\delta \ll 1$ , then  $S_1$  (up to conjugation) represents some element  $\gamma \in \pi_1^{loc}(\mathcal{B}, \bar{a}_0)$  and it is easy to see that the cyclic permutation type of  $h_*(\gamma) \in \mathcal{G}_{\bar{a}_0}$  is  $(6, 2, 1)$ .  $\square$

**2.6. On local monodromy groups of morphism  $h$  at the points belonging to strata of variety  $\mathcal{B}$  of codimension  $\geq 3$ .** Consider the point  $\bar{a} \in \mathcal{B}_{3,2}$ . In any neighborhood of this point there are points  $\bar{a}_1 \in \mathcal{B}_{2,1}$  and  $\bar{a}_2 \in \mathcal{B}_{2,2}$ . Therefore, according to Claim 7, the groups  $\mathcal{G}_{\bar{a}_1}$  and  $\mathcal{G}_{\bar{a}_2} \simeq Hes^1$  are subgroups of the local monodromy group  $\mathcal{G}_{\bar{a}} \subset \mathbb{S}_9$ . The action of the group  $\mathcal{G}_{\bar{a}_1}$  on the set  $\{1, \dots, 9\}$  has no fixed elements, and the group  $Hes^1$  acts transitively on the set  $\{2, \dots, 9\}$ . Hence,  $\mathcal{G}_{\bar{a}}$  contains all groups  $Hes^i$ ,  $i = 1, \dots, 9$ .

Since  $\mathcal{B}$  is an irreducible variety, it follows from Proposition 3 that the monodromy group  $\mathcal{G}$  of the morphism  $h$  is generated by permutations of cyclic type  $(3, 3, 1, 1, 1)$ . Each of these generators is contained in some group  $Hes^i$ . From this it follows

**Proposition 5.** *The local monodromy groups  $\mathcal{G}_{\bar{a}}$  of the morphism  $h$  at the points  $\bar{a} \in \mathcal{B}_{3,2}$  and the local monodromy groups  $\mathcal{G}_{\bar{a}}$  at the points  $\bar{a}$ , belonging to the strata of  $\mathcal{B}'$  of the variety  $\mathcal{B}$  of codimension  $\geq 4$ , coincide with the whole group  $\mathcal{G} = Hes$ .*

### 3. THE SURFACES OF INFLECTION POINTS OF GENERIC TWO-DIMENSIONAL LINEAR SYSTEMS OF PLANE CUBIC CURVES

**3.1. Two-dimensional linear systems of plane cubic curves.** Let  $\bar{z} = (z_1, z_2, z_3)$  be homogeneous coordinates in  $\mathbb{P}^2$  and  $\bar{t} = (t_1, t_2, t_3)$  homogeneous coordinates in a projective plane  $\Pi \subset \mathbb{P}^9$ . Consider (projectively) two-dimensional linear system of plane cubic curves  $C_{\Pi}$  given by equation  $F(\bar{t}, \bar{z}) = 0$ , where

$$F(\bar{t}, \bar{z}) = t_1 F(\bar{a}_1, \bar{z}) + t_2 F(\bar{a}_2, \bar{z}) + t_3 F(\bar{a}_3, \bar{z}).$$

Let  $H(\bar{t}, \bar{z}) = \det\left(\frac{\partial^2 F(\bar{t}, \bar{z})}{\partial z_i \partial z_j}\right)$  be the Hessian of the polynomial  $F(\bar{t}, \bar{z})$ . The equations

$$F(\bar{t}, \bar{z}) = 0, \quad H(\bar{t}, \bar{z}) = 0 \tag{8}$$

define in  $\Pi \times \mathbb{P}^2$  a surface  $\mathcal{I}_{\Pi} = \mathcal{I} \cap \text{pr}^{-1}(\Pi)$  of the inflection points of the linear system of cubic curves  $C_{\Pi}$ .

Set  $x = \frac{z_1}{z_3}$ ,  $y = \frac{z_2}{z_3}$  and  $\alpha = \frac{t_2}{t_1}$ ,  $\beta = \frac{t_3}{t_1}$ . Below, each time, without specifying, we will choose the base elements  $F_i(\bar{z}) := F(\bar{a}_i, \bar{z})$  of the linear system  $C_{\Pi}$  and homogeneous coordinates  $(z_1, z_2, z_3)$  in  $\mathbb{P}^2$  so that the form of equations (8) in non-homogeneous

coordinates  $(\alpha, \beta, x, y)$  in  $\mathbb{C}^2 \times \mathbb{C}^2 \subset \Pi \times \mathbb{P}^2$  would be most convenient to study the properties of the variety  $\mathcal{L}_\Pi$ .

**3.2. Generic projections to the plane.** The complete linear system of cubic curves, parametrized by the points of the space  $\mathbb{P}^9$ , defines, so called, Veronese-3 imbedding  $\varphi_3 : \mathbb{P}^2 \hookrightarrow \hat{\mathbb{P}}^9$ ,  $\varphi_3(\mathbb{P}^2) = \mathcal{V}_3$ , of the projective plane  $\mathbb{P}^2$  into the dual projective space of the parameter space  $\mathbb{P}^9$ . Two-dimensional linear subspaces  $\Pi \subset \mathbb{P}^9$  define linear projections  $\text{pr}_\Pi : \hat{\mathbb{P}}^9 \rightarrow \hat{\Pi}$  into the projective planes  $\hat{\Pi}$  dual to the projective planes  $\Pi$ . It is well known (see, for example, [3]) that for planes  $\Pi$  belonging to a Zariski open everywhere dense set  $\mathcal{W}$  in Grassmannian  $Gr(3, 10)$  of two-dimensional planes of  $\mathbb{P}^9$ , the compositions  $\xi := \text{pr}_\Pi \circ \varphi_3 : \mathbb{P}^2 \rightarrow \hat{\Pi}$  are generic coverings of the projective plane, i.e., they have the following properties:

- (i)  $\xi$  is a finite morphism of degree nine;
- (ii) the ramification curve  $\tilde{R} \subset \mathbb{P}^2$  of  $\xi$  is non-singular and ramification index along  $\tilde{R}$  is equal to two;
- (iii) the discriminant  $\hat{B} = \xi(\tilde{R})$  has only ordinary nodes and cusps as its singular points and the restriction of  $\xi$  to  $\tilde{R}$  is a birational morphism.

It follows from the proof of Theorem 4 in [12] that the degree of the curve  $\hat{B}$  is equal to 18, and the curve  $\hat{B}$  has 42 ordinary cusps, 84 ordinary nodes, and the geometric genus  $g(\hat{B})$  of  $\hat{B}$  is equal to 10. The curve  $\tilde{R} \subset \mathbb{P}^2$  is a smooth plane curve of degree six.

The cubic curves  $C_{\bar{\alpha}} = \xi^{-1}(L_{\bar{\alpha}})$ , parametrized by the points of  $\Pi$ , are inverse images of lines  $L_{\bar{\alpha}}$  in  $\hat{\Pi}$  dual to the points  $\bar{\alpha} \in \Pi$ . A cubic curve  $C_{\bar{\alpha}}$  is singular if and only if  $L_{\bar{\alpha}}$  is a tangent line of the curve  $\hat{B}$ . Therefore the curve  $\hat{B}$  is dual to the curve  $B := \mathcal{B} \cap \Pi$  and if a cubic curve  $C_{\bar{\alpha}}$  is singular, then its singular points belong to  $C_{\bar{\alpha}} \cap \tilde{R}$ . For a point  $q \in L_{\bar{\alpha}} \cap \hat{B}$ , the only one point from  $\xi^{-1}(q)$  can be a singular point of the curve  $C_{\bar{\alpha}}$ . In addition, the restriction of projection  $\text{pr}_2 : \Pi \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$  to  $S = \mathcal{S} \cap (\Pi \times \mathbb{P}^2)$  is a one-to-one map onto the curve  $\tilde{R}$ .

It follows from equisingular stratification of singular cubic curves (see subsection 2.2) that  $\hat{B}$  can have only bitangent and three-tangent lines  $L_{\bar{\alpha}_i}$  (in this case cubic curves  $C_{\bar{\alpha}_i}$  are unions of transversally intersecting lines and quadrics, i.e.,  $\bar{\alpha}_i \in \mathcal{B}_{2,1}$ , and, respectively, they are unions of three lines, i.e.,  $\bar{\alpha}_i \in \mathcal{B}_{3,1}$ ), and if  $L_{\bar{\alpha}_i}$  is the tangent line of the curve  $\hat{B}$  at its inflection point, then  $C_{\bar{\alpha}_i}$  is a cuspidal cubic, i.e.,  $\bar{\alpha}_i \in \mathcal{B}_{2,2}$ . Therefore, by Plücker formulae (see, for example, [13]), the degree of the curve  $B$  is equal to 12 and  $B$  has 24 ordinary cusps,  $n_1$  ordinary nodes, and  $n_2$  ordinary triple points (the singularities of type  $D_4$ ) as its singular points, where  $n_1 + 3n_2 = 21$ .

**Remark 1.** *Note that if the plane  $\Pi$  is sufficiently general, i.e.  $\Pi \cap \mathcal{B}_{3,1} = \emptyset$ , then the curves  $B \subset \Pi$  and  $\hat{B} \subset \hat{\Pi}$  give an example of two dual cuspidal curves for which the fundamental groups  $\pi_1(\Pi \setminus B)$  and  $\pi_1(\hat{\Pi} \setminus \hat{B})$  are not abelian groups.*

Denote

$$\mathfrak{B}_{\tilde{R},1} = \{\bar{z} \in \tilde{R} \mid \xi(\bar{z}) \text{ is an inflection point of } \hat{B}\},$$

$$\mathfrak{B}_{\tilde{R},2} = \{\bar{z} \in \tilde{R} \mid \xi(\bar{z}) \text{ is a cusp of } \hat{B}\}.$$

The set  $\mathfrak{B}_{\tilde{R},1}$  consists of 24 points and  $\mathfrak{B}_{\tilde{R},2}$  consists of 42 points.

Further we assume that  $\Pi \in \mathcal{W}$ .

**3.3. Properties of Stein coverings associated with morphisms  $h_\Pi$ .** A two-dimensional linear system of cubic curves  $\mathcal{C}_\Pi$  defines a generic covering of the plane  $\xi : \mathbb{P}^2 \rightarrow \hat{\Pi}$  and it follows from van Kampen – Zariski Theorem that the monodromy group of the covering  $h_\Pi : \mathcal{I}_\Pi \rightarrow \Pi$  is the group  $\mathcal{G} = Hes \subset \mathbb{S}_9$ .

Let us consider the Stein covering  $\varphi : Y \rightarrow \Pi$ ,  $\deg \varphi = 9$  associated with  $h_\Pi$ . The covering  $\varphi$  is unramified over  $\Pi \setminus B$ . The following description of properties of the covering  $\varphi : \varphi^{-1}(V) \rightarrow V$ , where  $V \subset \Pi$  is a sufficiently small complex-analytic neighbourhood of a point  $o \in B$ , based on results of section 2 and subsection 1.2.

If  $o \in \mathcal{B}_1$ , then the local monodromy group of the covering  $\varphi$  at  $o$  is the cyclic group generated by permutation whose cyclic type is  $(1, 1, 1, 3, 3)$ . Therefore  $\varphi^{-1}(V)$  is a disjoint union of five irreducible open in  $Y$  sets,  $\varphi^{-1}(V) = \cup_{i=1}^5 U_i$ , each of which is non-singular and, up to numbering, the coverings  $\varphi : U_i \rightarrow V$ ,  $i = 1, 2, 3$ , are bi-holomorphic maps and  $\varphi : U_i \rightarrow V$ ,  $i = 4, 5$ , are three-sheeted coverings branched along  $B \cap V$ .

If  $o \in \mathcal{B}_{2,1}$  and  $B \cap V = B'_1 \cup B'_2$ , where  $B'_1$  and  $B'_2$  are two branches of the curve  $B$  intersecting transversally at  $o$ , then the local monodromy group  $\mathcal{G}_{2,1} \subset \mathbb{S}_9$  of the covering  $\varphi$  at  $o$  is the abelian group generated by permutations  $g_3 = (2, 8, 5)(3, 6, 9)$  (the image of bypass around  $B'_1$ ) and  $g_4 = (1, 4, 7)(3, 9, 6)$  (the image of bypass around  $B'_2$ ). The set  $\{1, \dots, 9\}$  splits into three orbits  $\{1, 4, 7\}$ ,  $\{2, 8, 5\}$ , and  $\{3, 9, 6\}$  of the action of the group  $\mathcal{G}_{2,1}$ . Therefore  $\varphi^{-1}(V)$  is a disjoint union of three irreducible open in  $Y$  sets,  $\varphi^{-1}(V) = \cup_{i=1}^3 U_i$ . Up to numbering, the sets  $U_1$  and  $U_2$  are non-singular and  $U_3$  has a singularity at the point  $o' = U_3 \cap \varphi^{-1}(o)$ . The coverings  $\varphi : U_i \rightarrow V$ ,  $i = 1, 2, 3$ , are three-sheeted, the coverings  $\varphi : U_1 \rightarrow V$  is ramified over  $B'_1$ , the covering  $\varphi : U_2 \rightarrow V$  is ramified over  $B'_2$ , and the covering  $\varphi : U_3 \rightarrow V$  is ramified over  $B'_1 \cup B'_2$ . Note (see the proof of Claim 4) that if  $\psi : W \rightarrow U_3$  is the minimal desingularisation of the singular point  $o'$ ,  $E = \psi^{-1}(o')$ , then  $E$  is the smooth rational curve and  $(E^2)_W = -3$ .

If  $o \in \mathcal{B}_{3,1}$  and  $B \cap V = B'_1 \cup B'_2 \cup B'_3$ , where  $B'_1$ ,  $B'_2$ , and  $B'_3$  are three branches of the curve  $B$  pairwise meeting transversally at  $o$ , then the local monodromy group  $\mathcal{G}_{3,1} \subset \mathbb{S}_9$  of the covering  $\varphi$  at  $o$  is the abelian group generated by three permutations  $g_3 = (2, 8, 5)(3, 6, 9)$  (the image of bypass around  $B'_1$ ),  $g_4 = (1, 4, 7)(3, 9, 6)$  (the image of bypass around  $B'_2$ ), and  $g_5 = (2, 5, 8)(1, 7, 4)$  (the image of bypass around  $B'_3$ ). The set  $\{1, \dots, 9\}$  splits into three orbits  $\{1, 4, 7\}$ ,  $\{2, 8, 5\}$ , and  $\{3, 9, 6\}$  of the action of the group  $\mathcal{G}_{3,1}$ . Therefore  $\varphi^{-1}(V)$  is a disjoint union of three irreducible open in  $Y$  sets,  $\varphi^{-1}(V) = \cup_{i=1}^3 U_i$ . Up to numbering, the sets  $U_1$ ,  $U_2$ , and  $U_3$  have a singularity at the points  $o'_i = U_i \cap \varphi^{-1}(o)$ . The coverings  $\varphi : U_i \rightarrow V$ ,  $i = 1, 2, 3$ , are three-sheeted,

the covering  $\varphi : U_1 \rightarrow V$  is ramified over  $B'_1 \cup B'_2$ , the covering  $\varphi : U_2 \rightarrow V$  is ramified over  $B'_2 \cup B'_3$ , and the covering  $\varphi : U_3 \rightarrow V$  is ramified over  $B'_1 \cup B'_3$ . As above, note that if  $\psi : W_i \rightarrow U_i$  are the minimal desingularisations of the singular points  $o'_i$ ,  $E_i = \psi^{-1}(o'_i)$ , then  $E_i$  are smooth rational curves and  $(E_i^2)_{W_i} = -3$ .

If  $o \in \mathcal{B}_{2,2}$ , then the group of local monodromy  $\mathcal{G}_{2,2} \subset \mathbb{S}_9$  of the covering  $\varphi$  at  $o$  is the group  $\mathcal{G}_{2,2} \simeq Hes^1$  generated by two permutations  $g_1 = (4, 5, 6)(7, 9, 8)$  and  $g_2 = (2, 8, 5)(3, 6, 9)$ . The set  $\{1, \dots, 9\}$  splits into two orbits  $\{1\}$  and  $\{2, \dots, 9\}$  of the action of the group  $\mathcal{G}_{2,2}$ . Therefore  $\varphi^{-1}(V)$  is a disjoint union of two irreducible open in  $Y$  sets,  $\varphi^{-1}(V) = U_1 \cup U_2$ . The covering  $\varphi : U_1 \rightarrow V$  is a bi-holomorphic map and therefore  $U_1$  is non-singular. The degree of covering  $\varphi : U_2 \rightarrow V$  is equal to eight and this covering is branched over the curve  $B \cap V = B'$  having the singularity of type  $A_2$  at  $o$ . The group  $Hes^1 \subset \mathbb{S}_8$  is the monodromy group of the covering  $\varphi : U_2 \rightarrow V$  ( $\mathbb{S}_8$  acts on  $\{2, \dots, 9\}$ ).

Let us show that  $U_2$  is nonsingular, the ramification curve  $R' \subset U_2$  of the covering  $\varphi$  is also nonsingular, and the map  $\varphi : R' \rightarrow B'$  is two-sheeted and it is ramified at the point  $o$ . For this, let us remind that the group  $Hes^1 = \langle g_1, g_2 \rangle$  can be imbedded into  $GL(2, \mathbb{C})$  (the group no. 4 in [20], see also [15]) and under this imbedding the elements  $g_1$  and  $g_2$  are 3-reflections:

$$g_1 = \frac{-\omega}{\sqrt{2}} \begin{pmatrix} \varepsilon & \varepsilon^3 \\ \varepsilon & \varepsilon^7 \end{pmatrix}, \quad g_2 = \frac{\omega}{\sqrt{2}} \begin{pmatrix} \varepsilon^3 & \varepsilon^5 \\ \varepsilon^7 & \varepsilon^5 \end{pmatrix},$$

where  $\omega = e^{2\pi i/3}$  and  $\varepsilon = e^{2\pi i/8}$ . Since the group  $Hes^1 \subset GL(2, \mathbb{C})$  is generated by reflections, the factor-space  $\mathbb{C}^2/Hes^1$  is  $\mathbb{C}^2$ . Denote by  $\tilde{\psi} : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  the factorization morphism. We have  $\deg \tilde{\psi} = 24$  and, as it is known (see [15]),  $\tilde{\psi}$  is defined by functions  $\Psi = u^4 - 2\sqrt{-3}u^2v^2 + v^4$ ,  $\Theta = uv(u^4 - v^4)$  and it is ramified with multiplicity three along four lines  $\cup_{i=1}^4 L_i$  given by equation  $\Phi = 0$ , where  $\Phi = u^4 + 2\sqrt{-3}u^2v^2 + v^4$ . The branch curve  $\tilde{B}$  is given by  $\Psi^3 + 12\sqrt{-3}\Theta^2 = 0$ . The degree of the restriction of  $\tilde{\psi}$  to each line  $L_i$ ,  $\tilde{\psi} : L_i \rightarrow \tilde{B}$ , equals two. The group  $\langle g_1 \rangle$  transpose the lines  $L_2$ ,  $L_3$ , and  $L_4$  and leaves fixed the line  $L_1$  given by equation  $(\sqrt{2} + \omega\varepsilon)u + \omega\varepsilon^3v = 0$ . The covering  $\tilde{\psi}$  is the composition  $\tilde{\psi} = \psi \circ \bar{\psi}$ , where  $\bar{\psi} : \mathbb{C}^2 \rightarrow \mathbb{C}^2/\langle g_1 \rangle \simeq \mathbb{C}^2$  is the three-sheeted factorization map defined by the action of the group  $\langle g_1 \rangle$ . It is ramified along  $L_1$  and  $\psi : \mathbb{C}^2/\langle g_1 \rangle \rightarrow \mathbb{C}^2$  is a eight-sheeted covering ramified with multiplicity three along nonsingular curve  $\bar{R} = \bar{\psi}(L_2 \cup L_3 \cup L_4)$  and whose monodromy group is  $Hes^1 \subset \mathbb{S}_8$  acting on eight left cosets of the group  $\langle g_1 \rangle$ . The covering  $\psi : \bar{R} \rightarrow \tilde{B}$  is a two-sheeted map ramified at the origin. Now, to show that  $U_2$  is nonsingular, the ramification curve  $R' \subset U_2$  of  $\varphi$  is also nonsingular, and  $\varphi : R' \rightarrow B'$  is a two-sheeted covering branched at  $o$ , it suffices to imbed the pair  $(V, B')$  into  $(\mathbb{C}^2, \tilde{B})$  and apply Grauert - Remmert - Riemann - Stein Theorem, Proposition 1, and Claim 8 to the coverings  $\psi : \psi^{-1}(V) \rightarrow V$  and  $\varphi : U_2 \rightarrow V$ .

Denote by  $\mathcal{R} \subset Y$  the ramification curve of  $\varphi$ . It follows from above that  $\varphi : \mathcal{R} \rightarrow B$  is a two-sheeted covering branched at 24 points (the cusps of  $B$ ). Therefore, by Hurwitz formula, the geometric genus  $g(\mathcal{R})$  of  $\mathcal{R}$  is equal to

$$g(\mathcal{R}) = \frac{1}{2}[2(2g(B) - 2) + 24] + 1 = 31.$$

The number of singular points of the surface  $Y$  equals  $n_1 + 3n_2 = 21$ .

**Proposition 6.** *Let  $\sigma : X \rightarrow Y$  be the minimal resolutions of singularities of  $Y$ . The square of the canonical class  $K_X$  of  $X$  equals  $K_X^2 = 18$  and the Euler characteristic  $\chi(\mathcal{O}_X) = \sum_{i=0}^2 (-1)^i \dim H^i(X, \mathcal{O}_X)$  of the structure sheaf  $\mathcal{O}_X$  of  $X$  equals  $\chi(\mathcal{O}_X) = 1 - q + p_g = 9$ .*

*Proof.* Denote  $\psi = \varphi \circ \sigma$  and denote by  $R \subset X$  the proper preimage of the curve  $\mathcal{R}$ . By Claim 4,  $\sigma^{-1}(\text{Sing } Y)$  is the disjoint union of 21 rational curves,  $E := \sigma^{-1}(\text{Sing } Y) = \sqcup_{i=1}^{21} E_i$ . In addition, by the same Claim, we have  $(E_i^2)_X = -3$  and  $(R, E_i)_X = 2$ .

We have  $K_X = \psi^*(K_\Pi) + 2R + E$ , since  $\psi$  is ramified along  $R$  with multiplicity three,  $(E_i, \psi^*(K_\Pi))_X = 0$  and  $(K_X + E_i, E_i)_X = -2$ . In addition, we have  $(\psi^*(K_\Pi), \psi^*(K_\Pi))_X = 81$ , since  $\deg \psi = K_\Pi^2 = 9$ , and  $(R, \psi^*(K_\Pi))_X = -72$ , since the degree of the covering  $\psi : R \rightarrow B$  equals two and  $\deg B = 12$ . Therefore the equalities

$$(R^2)_X = (R, K_X)_X = 30 \tag{9}$$

follow from equalities  $(K_X + R, R)_X = (\psi^*(K_\Pi) + 3R + E, R)_X = 60$ . As a result, we have  $K_X^2 = (\psi^*(K_\Pi) + 2R + E, \psi^*(K_\Pi) + 2R + E)_X = 18$ .

Denote by  $e(M)$  Euler topological characteristic of a space  $M$ . We have  $e(X) = 9e(\Pi \setminus B) + 5e(B \setminus \text{Sing } B) + 4n_1 + 6n_2 + 2c$ , since  $\deg \psi = 9$ , the number of preimages of each nonsingular point of the curve  $B$  equals five, the preimage of each node of  $B$  is a disjoint union of projective line ( $e(\mathbb{P}^1) = 2$ ) and two points, the preimage of each triple point of  $B$  is a disjoint union of three projective lines, and the preimage of each cusp of  $B$  consists of two points. And since  $g(B) = 10$ ,  $n_1 + 3n_2 = 21$ ,  $c = 24$ , and  $e(\Pi) = 3$ , then  $e(B) = -2(g(B) - 1) - n_1 - 2n_2 = n_2 - 39$ ,  $e(\Pi \setminus B) = 42 - n_2$ , and  $e(B \setminus \text{Sing } B) = -2(g(B) - 1) - 2n_1 - 3n_2 - c = 3n_2 - 84$ . Therefore,  $e(X) = 90$  and applying Noether formula, we obtain that  $\chi(\mathcal{O}_X) = \frac{K_X^2 + e(X)}{12} = 9$ .  $\square$

**3.4. Resolution of singularities of a surface of inflection points of generic two-dimensional linear system of plane cubic curves.** Denote by  $\nu : I := I_\Pi \rightarrow \mathcal{I}_\Pi$  the normalization of variety  $\mathcal{I}_\Pi$  and let  $p_1 = h_\Pi \circ \nu : I \rightarrow \Pi$ . Denote also  $S = \mathcal{S} \cap \mathcal{C}_\Pi$ . By Claim 5, the curve  $S$  is contained in  $\mathcal{I}_\Pi$ .

**Proposition 7.** *The surface  $I$  is nonsingular and the morphism  $p_1 : I \rightarrow \Pi$  coincides with  $\psi = \varphi \circ \sigma : X \rightarrow \Pi$ , where  $\varphi : Y \rightarrow \Pi$  is the Stein covering associated with  $h_\Pi$  and  $\sigma : X \rightarrow Y$  is the minimal resolution of singularities of the surface  $Y$ .*

*Proof.* Obviously,  $\mathcal{I}_\Pi$  is nonsingular at the points belonging to  $h_\Pi^{-1}(\Pi \setminus B)$ .

**Claim 9.** *The surface  $\mathcal{I}_\Pi$  is nonsingular at the points belonging to  $\mathcal{I}_\Pi \setminus S$ .*

*Proof.* An irreducible nodal cubic  $C_{\bar{a}_1}$ ,  $\bar{a}_1 \in \mathcal{B}_1 \cap \Pi$ , has three smooth inflection points (respectively, a cuspidal cubic  $C_{\bar{a}_1}$ ,  $\bar{a}_1 \in \mathcal{B}_{2,2} \cap \Pi$ , has a single smooth inflection point). It was shown in [13] that the intersection number of the cubic curve  $C_{\bar{a}_1}$  and its Hesse curve  $H_{\bar{a}_1}$  at its singular point  $s \in C_{\bar{a}_1}$  equals  $(C_{\bar{a}_1}, H_{\bar{a}_1})_s = 6$  (respectively,  $(C_{\bar{a}_1}, H_{\bar{a}_1})_s = 8$ ). Therefore, by Claim 3, the surface  $\mathcal{I}_\Pi$  is nonsingular at the points belonging to  $\mathcal{I}_\Pi \setminus (S \cup h^{-1}(\mathcal{B}_{2,1} \cup \mathcal{B}_{3,1}))$ . Therefore, to complete the proof of Proposition 9, it suffices to show that  $\mathcal{I}_\Pi$  is non-singular at the points of  $h^{-1}(\bar{a}_1) \setminus \mathcal{S}$  for  $\bar{a}_1 \in (\mathcal{B}_{2,1} \cup \mathcal{B}_{3,1}) \cap \Pi$ .

Without loss of generality, we can assume that  $z_1 z_2 z_3 + a_{3,0} z_1^3 = 0$  is an equation of the cubic curve  $C_{\bar{a}_1}$  (if  $\bar{a}_1 \in \mathcal{B}_{3,1}$ , then  $a_{3,0} = 0$  and  $h_\Pi^{-1}(\bar{a}_1)$  are three lines in  $\{\bar{a}_1\} \times \mathbb{P}^2$  given by equations  $z_i = 0$ ,  $i = 1, 2, 3$ , and if  $\bar{a}_1 \in \mathcal{B}_{2,1}$ , then  $a_{3,0} \neq 0$  and  $h_\Pi^{-1}(\bar{a}_1)$  is the line in  $\{\bar{a}_1\} \times \mathbb{P}^2$  given by equation  $z_1 = 0$ ).

Let us show that the points of line  $E = \{\bar{a}_1\} \times \{z_1 = 0\}$  are non-singular points of  $\mathcal{I}_\Pi$  if  $z_2 z_3 \neq 0$  (in the case when  $\bar{a}_1 \in \mathcal{B}_{3,1}$  and the points belong to the lines  $\{z_2 = 0\}$  and  $\{z_3 = 0\}$  not being singular points of the curve  $C_{\bar{a}_1}$ , the proof that the surface  $\mathcal{I}_\Pi$  is non-singular at these points is similar and therefore it will be omitted). For this, let us choose base elements  $F_1(\bar{z})$ ,  $F_2(\bar{z})$ ,  $F_3(\bar{z})$  of linear system  $\mathcal{C}_\Pi$  so that in non-homogeneous coordinates  $(\alpha, \beta, x, y)$  the linear system  $\mathcal{C}_\Pi$  is given by equation

$$(xy + a_{3,0}x^3) + \alpha \left( \sum_{1 \leq i+j \leq 3} b_{i,j} x^i y^j \right) + \beta \left( 1 + \sum_{1 \leq i+j \leq 3} c_{i,j} x^i y^j \right) = 0. \quad (10)$$

Denote by  $\mathfrak{m}$  the ideal in the polynomial ring  $\mathbb{C}[\alpha, \beta, x, y]$  generated by monomials  $\alpha$ ,  $\beta$ , and  $x$  and write equation (10) in the following form:

$$xy + \alpha \left( \sum_{1 \leq j \leq 3} b_{0,j} y^j \right) + \beta \left( 1 + \sum_{1 \leq j \leq 3} c_{0,j} y^j \right) + R_1(\alpha, \beta, x) = 0, \quad (11)$$

where  $R_1(\alpha, \beta, x) \in \mathfrak{m}^2$ . Simple calculations (which we omit) show that in non-homogeneous coordinates  $(\alpha, \beta, x, y)$  the Hessian of the linear system  $\mathcal{C}_\Pi$  has the following form:

$$2[xy + \alpha \left( \sum_{1 \leq j \leq 2} b_{0,j} y^j - 3b_{0,3} y^3 \right) + \beta(-3 + \sum_{1 \leq j \leq 2} c_{0,j} y^j - 3c_{0,3} y^3)] + R_2(\alpha, \beta, x) = 0, \quad (12)$$

where  $R_2(\alpha, \beta, x) \in \mathfrak{m}^2$ .

The surface  $\mathcal{I}_\Pi \cap (\mathbb{C}^2 \times \mathbb{C}^2)$  in  $\mathbb{C}^2 \times \mathbb{C}^2$  is given by equations (11) and (12), and the line  $E$  is given by equations  $\alpha = \beta = x = 0$ . Therefore necessary and sufficient condition for a point  $p = (0, 0, 0, y_0)$  with  $y_0 \neq 0$  to be singular point of  $\mathcal{I}_\Pi$  is the proportionality of (linear in variables  $\alpha$ ,  $\beta$ , and  $x$ ) forms entering into left parts of equations (11) and (12), and in which  $y$  takes the concrete value, namely,  $y_0$ . It is easy to see that one of necessary conditions is vanishing the coefficient  $b_{0,3}$ . In this case the cubic curve  $C_{\bar{a}_2}$  given by equation  $F_2(\bar{z}) = 0$  contains not only the singular point  $q_1 = (0, 0, 1)$  of the curve  $C_{\bar{a}_1}$ , but also it contains the second singular point

$q_2 = (0, 1, 0)$  of  $C_{\bar{a}_1}$ . In our case it is impossible, since the linear system  $\mathcal{C}_\Pi$  defines the generic covering  $\xi : \mathbb{P}^2 \rightarrow \hat{\Pi}$  of the projective plane.  $\square$

**Claim 10.** *The curve  $S$  is non-singular at  $s \in S \subset \mathcal{I}_\Pi$  if  $h_\Pi(s) \in \mathcal{B}_1 \cup \mathcal{B}_{2,1} \cup \mathcal{B}_{3,1}$ .*

*Proof.* As above, we can assume that the point  $s$  lies in neighbourhood  $\mathbb{C}^2 \times \mathbb{C}^2 \subset \Pi \times \mathbb{P}^2$  with coordinates  $(\alpha, \beta, x, y)$  and has coordinates  $(0, 0, 0, 0)$ . Choose the base polynomials  $F_1(x, y)$ ,  $F_2(x, y)$ , and  $F_3(x, y)$  of linear system  $\mathcal{C}_\Pi$  as follows:

$$F_1(x, y) = xy + a_{3,0}x^3 + a_{0,3}y^3, \quad F_2(x, y) = \sum_{1 \leq i+j \leq 3} b_{i,j}x^i y^j, \quad F_3(x, y) = 1 + \sum_{1 \leq i+j \leq 3} c_{i,j}x^i y^j,$$

i.e., the linear system  $\mathcal{C}_\Pi$  is given by

$$(xy + a_{3,0}x^3 + a_{0,3}y^3) + \alpha \left( \sum_{1 \leq i+j \leq 3} b_{i,j}x^i y^j \right) + \beta \left( 1 + \sum_{1 \leq i+j \leq 3} c_{i,j}x^i y^j \right) = 0. \quad (13)$$

We can assume that  $b_{1,1} = c_{1,1} = 0$  (correcting  $F_2(x, y)$  and  $F_3(x, y)$  with help of the polynomial  $F_1(x, y)$  if it is necessary) and  $b_{1,0} = 1$ ,  $c_{1,0} = 0$ . Denote by  $\mathfrak{m}$  the ideal in the power series ring  $\mathbb{C}[[\alpha, \beta, x, y]]$  generated by monomials  $\alpha$ ,  $\beta$ ,  $x$ , and  $y$ , and denote by  $\mathfrak{m}_{\hat{\beta}}$  the ideal in the power series ring  $\mathbb{C}[[\alpha, x, y]]$  generated by monomials  $\alpha$ ,  $x$ , and  $y$ . Denote also by  $w$  the left side of equation (13) and let  $w_1 = \frac{\partial w}{\partial x}$  and  $w_2 = \frac{\partial w}{\partial y}$ . We have  $w = \beta + R_0(\alpha, \beta, x, y)$  and

$$w_1 = y + \alpha + R_1(\alpha, \beta, x, y), \quad w_2 = x + b_{0,1}\alpha + c_{0,1}\beta + R_2(\alpha, \beta, x, y), \quad (14)$$

where  $R_i(\alpha, x, y) \in \mathfrak{m}^2$  for  $i = 0, 1, 2$ . It is easy to see from (14) that the functions  $w$ ,  $w_1$ ,  $w_2$ , and  $\alpha$  are local parameters in some neighbourhood  $V$  of the point  $s$ , the variety  $U = \mathcal{C}_\Pi \cap V$  is given in  $V$  by equation  $w = 0$ , and the curve  $S \cap V$  is given in  $U$  by equations  $w_1 = w_2 = 0$ .  $\square$

**Claim 11.** *If  $h_\Pi(s) \in \mathcal{B}_1 \cup \mathcal{B}_{2,1} \cup \mathcal{B}_{3,1}$  for  $s \in S \subset \mathcal{I}_\Pi$ , then in some analytic neighbourhood of the point  $s$  the surface  $\mathcal{I}_\Pi$  is the union of two non-singular surfaces meeting transversally at  $s$ .*

*Proof.* We use notations introduced in the proof of Claim 10. Simple calculations (which we omit) show that in non-homogeneous coordinates  $(\alpha, \beta, x, y)$  the equation of Hesse surface  $H_{\mathcal{C}_\Pi}$  has the following form

$$2[-3\beta + xy + \alpha(x + b_{0,1}y) + \beta(-3 + c_{0,1}y) + 4b_{01}\alpha^2 + 4c_{0,1}\alpha\beta] + R_3(\alpha, \beta, x, y) = 0, \quad (15)$$

where  $R_3(\alpha, \beta, x, y) \in \mathfrak{m}^3$ .

It follows from (13) that the restriction of function  $\beta$  to  $V$  has the following form:

$$\beta = -[xy + \alpha(x + b_{0,1}y)] + R_4(\alpha, x, y), \quad (16)$$

where  $R_4(\alpha, x, y) \in \mathfrak{m}_{\hat{\beta}}^3$ . Therefore the surface  $\mathcal{I}_\Pi \cap U$  is given by equation (15) in which the right side of equation (16) is substituted instead of  $\beta$ . After this substitution and simple transformations, equation (15) has the following form:

$$4(y + \alpha)(x + b_{0,1}\alpha) + R_5(\alpha, x, y) = 0, \quad (17)$$

where  $R_5(\alpha, x, y) \in \mathfrak{m}_\beta^3$ . Therefore,  $s$  is a singular point of  $\mathcal{I}_\Pi$  of multiplicity two. Also, it follows from this that  $S = \text{Sing} \mathcal{I}_\Pi$ , since the point  $s$  is a generic point belonging to  $S$ .

It follows from (17) that  $s$ , as a singular point of the curve  $D = \{\alpha = 0\} \cap \mathcal{I}_\Pi$ , is an ordinary node. Let  $\sigma : \tilde{U} \rightarrow U$  be the monoidal transformation with center in the curve  $S \cap U$ ,  $\sigma^{-1}(S \cap U) = E$ . It is easy to see that the proper preimage  $\sigma^{-1}(D)$  of  $D$  intersects transversally with  $E$  at two points, and since  $s$  is a generic point of the curve  $S \cap U$ , then the intersection  $\sigma^{-1}(U \cap \mathcal{I}_\Pi) \cap E$  of the proper preimage of the surface  $U \cap \mathcal{I}_\Pi$  with  $E$  is a disjoint union of two nonsingular curves if the neighbourhood  $V$  is chosen small enough. It follows from this that the surface  $\mathcal{I}_\Pi \cap U$  is the union of two nonsingular surfaces meeting transversally, since the multiplicity of the singularity of the surface  $\mathcal{I}_\Pi \cap U$  at each point  $s \in S \cap U$  equals two.  $\square$

**Claim 12.** *If  $h_\Pi(s) \in \mathcal{B}_{2,2}$  for  $s \in S \subset \mathcal{I}_\Pi$ , then the surface  $I$  is nonsingular at the points of  $\nu^{-1}(s)$ .*

*Proof.* Let us consider a sufficiently small complex-analytic neighbourhood  $V \subset \Pi$  of the point  $p_1(s)$ . The morphism  $p_1 : p_1^{-1}(V) \rightarrow V$  is a finite covering and  $p_1^{-1}(V)$  is a normal surface (consisting of two components). Therefore Claim 12 follows from Grauert - Remmert - Riemann - Stein Theorem and results of subsection 3.3.  $\square$

To complete the proof of Proposition 7, it suffices to apply Claims 9 – 12, 4 and Grauert - Remmert - Riemann - Stein Theorem.  $\square$

#### 4. CALCULATION OF THE IRREGULARITY

**4.1. The additional condition of generality.** Further we use notations introduced in section 3. To compute the irregularity  $q = \dim H^1(\mathcal{O}_I, I)$  of a surface  $I$ , it is useful to introduce an *additional generality condition* of two-dimensional linear systems  $\mathcal{C}_\Pi$  of plane cubic curves. We say that a linear system  $\mathcal{C}_\Pi$  of cubic curves, parameterized by points of the projective plane  $\Pi \in \mathcal{W} \subset Gr(3, 10)$ , satisfies the *additional condition of generality* if for any point  $\bar{z} \in \mathbb{P}^2$  there is only a finite number of cubic curves of  $\mathcal{C}_\Pi$ , passing through  $\bar{z}$  for which  $\bar{z}$  is their inflection point.

Denote by  $J_{1,o}$  a subset of planes  $\Pi$  in  $\mathcal{W} \subset Gr(3, 10)$  such that the point  $o \in \mathbb{P}^2 \setminus \tilde{R}$  is an inflection point of each cubic from  $\mathcal{C}_\Pi$  containing the point  $o$ .

**Claim 13.** *We have  $\dim J_{1,o} \leq 17$ .*

*Proof.* Since  $o \in \mathbb{P}^2 \setminus \tilde{R}$ , then in some neighborhood of  $o$  the generic covering  $\xi : \mathbb{P}^2 \rightarrow \hat{\Pi}$  is a biholomorphic isomorphism. Therefore the base elements  $F_1(x, y)$ ,  $F_2(x, y)$ , and  $F_3(x, y)$  of the linear system  $\mathcal{C}_\Pi$  of the following form:

$$F_1(x, y) = x + \sum_{2 \leq i+j \leq 3} b_{i,j} x^i y^j, \quad F_2(x, y) = y + \sum_{2 \leq i+j \leq 3} c_{i,j} x^i y^j,$$

$$F_3(x, y) = 1 + \sum_{2 \leq i+j \leq 3} d_{i,j} x^i y^j$$

are defined uniquely by the point  $o \in \mathbb{P}^2 \setminus \widetilde{R}$  for any plane  $\Pi \in \mathcal{W}$ . Obviously, the necessary and sufficient condition for  $\Pi$  to belong to  $J_{1,o}$  is the divisibility of the quadratic form

$$(t_1 b_{2,0} + t_2 c_{2,0})x^2 + (t_1 b_{1,1} + t_2 c_{1,1})xy + (t_1 b_{0,2} + t_2 c_{0,2})y^2$$

by linear form  $t_1 x + t_2 y$  for each point  $(t_1, t_2) \in \mathbb{P}^1$ . It is easy to see (putting  $x = t_2 = 1$ ) that this condition is equivalent to the following equalities:

$$b_{2,0} = c_{1,1}, \quad b_{1,1} = c_{0,2}, \quad b_{0,2} = c_{2,0} = 0.$$

Consequently, the dimension of the variety  $J_{1,o}$  does not exceed the total (equals to 17) number of undefined coefficients  $b_{I,j}$ ,  $c_{I,j}$  and  $d_{I,j}$  for monomials in polynomials  $F_1(x, y)$ ,  $F_2(x, y)$ , and  $F_3(x, y)$ .  $\square$

Denote by  $J_{2,o}$  a subset of planes  $\Pi$  in  $W \subset Gr(3, 10)$  such that the point  $o \in \mathfrak{B}_{\widetilde{R},2}$  and it is an inflection point of each cubic curve from  $\mathcal{C}_\Pi$  passing through  $o$ .

**Claim 14.** *We have  $\dim J_{2,o} \leq 17$ .*

*Proof.* In some neighborhood of  $o$ , the covering  $\xi$  is equivalent to the restriction of the projection  $\text{pr} : (z, a, b) \mapsto (a, b)$  onto the surface in  $\mathbb{C}^3$  (with coordinates  $(z, a, b)$  in  $\mathbb{C}^3$ ) given by the equation  $z^3 + az + b = 0$ . The properties of this projection are well studied (see, for example, [12]). In particular, it is easy to show that, in our case, the pencil of cubic curves having the point  $o$  as its base point has one singular element at the point  $o$  (let us denote by  $C_1$  this cubic curve given by equation  $F_1(x, y) = 0$ ; the point  $o$  is an ordinary node of  $C_1$ ), and all other elements of the pencil are nonsingular at  $o$  and touch one of the branches of the curve  $C_1$  at that point. So we can choose in the  $\mathcal{C}_\Pi$  the base elements  $F_1(x, y)$ ,  $F_2(x, y)$ , and  $F_3(x, y)$  of the following form:

$$F_1(x, y) = xy + \sum_{i+j=3} b_{i,j} x^i y^j, \quad F_2(x, y) = x + \sum_{2 \leq i+j \leq 3} c_{i,j} x^i y^j,$$

$$F_3(x, y) = 1 + d_{0,1} y + \sum_{2 \leq i+j \leq 3} d_{i,j} x^i y^j, \quad c_{1,1} = d_{1,1} = 0.$$

It is easy to see that the necessary and sufficient condition for the plane  $\Pi$  to belong to  $J_{2,o}$  is the vanishing of the coefficient  $c_{0,2}$ . Obviously,  $\dim J_{2,o}$  does not exceed the total number (equal to 17) of undefined coefficients  $b_{i,j}$ ,  $c_{i,j}$ , and  $d_{i,j}$  for monomials in polynomials  $F_1(x, y)$ ,  $F_2(x, y)$ , and  $F_3(x, y)$ .  $\square$

**Remark 2.** From the proof of Claim 14 it is easy to see that if  $\Pi \notin J_{2,o}$  (i.e.,  $c_{0,2} \neq 0$ ), then for all nonsingular cubic curves belonging to the linear system  $\mathcal{C}_\Pi$  and passing through the point  $o \in \mathfrak{B}_{\widetilde{R},2}$ , the point  $o$  is not their inflection point.

Denote by  $J_{3,o}$  a subset of planes  $\Pi$  in  $W \subset Gr(3, 10)$  such that the point  $o \in \tilde{R} \setminus \mathfrak{B}_{\tilde{R},2}$  and it is the inflection point of each cubic curves from  $\mathcal{C}_\Pi$  passing through  $o$ .

**Claim 15.**  $J_{3,o}$  is the empty set.

*Proof.* In some neighborhood of  $o \in \tilde{R}$  the covering  $\xi$  is ramified along  $\tilde{R}$  with multiplicity two. Therefore, for any two cubic curves  $C_1$  and  $C_2$  belonging to the linear system  $\mathcal{C}_\Pi$  and passing through the point  $o$ , the intersection number  $(C_1, C_2)_o$  of the curves  $C_1$  and  $C_2$  at  $o$  is equal to two. In addition, as noted above, the pencil of cubic curves having the point  $o$  as its base point has only one cubic curve having the singularity at  $o$ , we denote this cubic by  $C_1$  (the preimage of the tangent line to one of the branches of the curve  $\hat{B}$  at point  $o'$ ), and all other elements of the pencil are nonsingular at  $o$ . In all cases, the point  $o$  is an ordinary node of the curve  $C_1$ , except when  $o'$  is the inflection point of  $\hat{B}$ . In the latter case, the point  $o$  is an ordinary cusp of the curve  $C_1$ .

Let  $F_1(x, y) = 0$  be an equation of  $C_1$  and let  $t_1F_1(x, y) + t_2F_2(x, y) = 0$  be the equation of the pencil of cubic curves passing through  $o$ . Without loss of generality, we can assume that  $F_2(x, y) = x + \sum_{2 \leq i+j \leq 3} b_{i,j}x^i y^j$ , and if  $o$  is an ordinary node of  $C_1$ , then  $F_1(x, y) = \sum_{2 \leq i+j \leq 3} c_{i,j}x^i y^j$ , where  $c_{0,2} = 1$ , since  $(C_1, C_2)_o = 2$  for the curve  $C_2$  given by equation  $F_2(x, y) = 0$ . And in the case when the point  $o$  is an ordinary cusp of  $C_1$ , we can assume that  $F_1(x, y) = y^2 + \sum_{i+j=3} c_{i,j}x^i y^j$ . In each of these cases, it is easy to see that for all elements of this pencil the point  $o$  would be the inflection point, it is necessary that for all  $(t_1, t_2) \in \mathbb{P}^1$  the equality  $t_1 + t_2 b_{0,2} = 0$  would be fulfilled, which is impossible.  $\square$

**Remark 3.** From the proof of Claim 15 it follows that if  $o$  is an ordinary cusp of  $C_1$ , then in the pencil of cubic curves passing through  $o$ , there is exactly one more cubic curve (with  $t_1 = -b_{0,2}$  and  $t_2 = 1$ ), which has inflection at this point.

**Claim 16.** *There is a non-empty Zariski open subset  $\mathcal{W}_1$  of  $\mathcal{W}$  such that two-dimensional linear systems of plane cubic curves  $\mathcal{C}_\Pi$ ,  $\Pi \in \mathcal{W}_1$ , satisfy the additional generality condition.*

*Proof.* Denote  $J = J_1 \cup J_2$ , where  $J_i = \{(\Pi, \bar{z}) \in \mathcal{W} \times \mathbb{P}^2 \mid \Pi \in J_{i,\bar{z}}\}$ ,  $i = 1, 2$ , and put  $\mathcal{W}_1 := \mathcal{W} \setminus \overline{\text{pr}_1(J)}$ , where the overline means the closure of the set. It follows from Claims 13 and 14 that  $\dim J \leq 19$  and so  $\mathcal{W}_1$  is a nonempty Zariski open set, since  $\dim \mathcal{W} = \dim Gr(3, 10) = 21$ . Furthermore, it also follows from these Claims and Claim 15 that linear systems  $\mathcal{C}_\Pi$  of cubic curves,  $\Pi \in \mathcal{W}_1$ , satisfy the additional generality condition.  $\square$

**4.2. Investigation of properties of the second projection.** Denote  $p_2 = \text{pr}_2 \circ \nu : I \rightarrow \mathbb{P}^2$ , where  $\nu : I \rightarrow \mathcal{I}_\Pi$  is the normalization morphism and  $\text{pr}_2 : \Pi \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$  is the projection to  $\mathbb{P}^2$ .

According to Proposition 7, we can identify  $I$  with the minimal resolution of singularities of Stein covering  $X$  associated with morphism  $h_\Pi$ . So, below we will use the notations introduced in the proof of Proposition 6. It follows from Proposition 6 that the surface  $I$  has the following invariants:  $K_I^2 = 18$  and  $\chi(\mathcal{O}_I) = \sum_{i=0}^2 (-1)^i \dim H^i(I, \mathcal{O}_I) = 1 - q + p_g = 9$ . The curve  $R \subset X = I$  is nonsingular and according to (9) we have the following equalities:  $(R^2)_I = (R, K_I)_I = 30$ . In addition,  $I$  contains 21 rational curves  $E_i$ ,  $i = 1, \dots, 21$ , such that  $(E_i^2)_I = -3$  and  $(E_i, K_I)_I = 1$ ,  $p_1(E_i)$  are points in  $(\mathcal{B}_{2,1} \cup \mathcal{B}_{3,1}) \cap \Pi$ , and  $p_2(E_i)$  are lines in  $\mathbb{P}^2$ .

Further, unless otherwise specified, we assume that  $\Pi \in \mathcal{W}_1$ .

**Claim 17.** *The morphism  $p_2 : I \rightarrow \mathbb{P}^2$  is a finite three-sheeted covering.*

*Proof.* Let  $L$  and  $M$  be lines respectively in  $\mathbb{P}^2$  and in  $\Pi$ . The Picard group  $\text{Pic}(\Pi \times \mathbb{P}^2)$  is a free abelian group generated by divisors  $\Pi \times L$  and  $M \times \mathbb{P}^2$ . The surface  $\mathcal{I}_\Pi$  is the complete intersection of two subvarieties  $\mathcal{C}_\Pi$  and  $H_{\mathcal{C}_\Pi}$  of codimension one. We have  $\mathcal{C}_\Pi = \Pi \times L + M \times \mathbb{P}^2$  and  $H_{\mathcal{C}_\Pi} = 3(\Pi \times L + M \times \mathbb{P}^2)$  as elements in  $\text{Pic}(\Pi \times \mathbb{P}^2)$ . The fibre of the projection  $\text{pr}_2 : \mathcal{I}_\Pi \rightarrow \mathbb{P}^2$  is also the intersection of two submanifolds  $\Pi \times L_1$  and  $\Pi \times L_2$  (where  $L_1$  and  $L_2$  are lines in  $\mathbb{P}^2$ ). It follows from this that the degree of the restriction of projection  $\text{pr}_2$  to  $\mathcal{I}_\Pi$  is equal to the intersection number  $(3(\Pi \times L + M \times \mathbb{P}^2), \Pi \times L + M \times \mathbb{P}^2, \Pi \times L, \Pi \times L)_{\Pi \times \mathbb{P}^2} = 3$ .  $\square$

To study the properties of the discriminant curve  $\mathfrak{B} \subset \mathbb{P}^2$  and the ramification curve of  $\mathfrak{R} \subset I$  of the covering  $p_2 : I \rightarrow \mathbb{P}^2$  we need the following

**Lemma 2.** *Let  $(U, o')$  and  $(V, o)$  be two germs of nonsingular surfaces at points  $o'$  and  $o$  and let  $\varphi : (U, o') \rightarrow (V, o)$  be a two-sheeted finite covering branched along a curve germ  $(B, o)$ ,  $\varphi^*((B, o)) = 2(R, o')$ . Then the germs  $(B, o)$  and  $(R, o')$  are nonsingular at  $o$  and  $o'$ , and if  $(C, o') \subset (U, o')$  and  $(\varphi(C), o) \subset (V, o)$  are two nonsingular germs of curves, then either  $\varphi : (C, o') \rightarrow (\varphi(C), o)$  is a two-sheeted covering branched at  $o'$  and the germs  $(\varphi(C), o)$  and  $(B, o)$  intersect transversally at  $o$ , or  $\varphi : (C, o') \rightarrow (\varphi(C), o)$  is biholomorphic map and the intersection numbers  $(\varphi(C), B)_o$  and  $(C, R)_{o'}$  of germs at  $o$  and  $o'$  are connected by equality  $(\varphi(C), B)_o = 2(C, R)_{o'}$ .*

*Proof.* Let  $\sum_{i+j=1}^{\infty} c_{i,j} x^i y^j = 0$  be an equation of  $(B, o)$  in  $(V, o)$ . Then the germ  $(U, o')$  is biholomorphic to the germ in  $(\mathbb{C}^3, o')$  given by  $z^2 = \sum_{i+j=1}^{\infty} c_{i,j} x^i y^j$ . Therefore, the linear part  $c_{1,0}x + c_{0,1}y$  of the equation is non-degenerate, since  $(U, o')$  is a germ of a nonsingular surface. Hence (by making a coordinate change) we can assume that the germ  $(B, o)$  is given by equation  $x = 0$  and the covering  $\varphi$  is given by equation  $z^2 = x$ .

If the germs  $(\varphi(C), o)$  and  $(B, o)$  intersect transversally at  $o$ , we can assume that  $y = 0$  is the equation of the germ  $(\varphi(C), o)$  and then, obviously,  $\varphi : (C, o') \rightarrow (\varphi(C), o)$

is a two-sheeted covering branched at  $o'$ , and germs  $(R, o')$  and  $(C, o')$  defined by equations  $z = 0$  and  $y = 0$ .

If the germs  $(\varphi(C), o)$  and  $(B, o)$  are touch each other at  $o$  with multiplicity  $r > 1$ , then by making an analytic change of coordinates, can assume that  $x = 0$  is the equation of the germ  $(B, o)$  and  $x - y^r = 0$  is the equation of the germ  $(\varphi(C), o)$ . Hence, the total preimage of  $(\varphi(C), o)$  is given by equation  $z^2 - y^r = 0$  and if  $r$  is an odd number, then this preimage is irreducible and it is a germ of a singular curve. If  $r = 2k$  is an even number, then the preimage of  $(\varphi(C), o)$  splits into two germs given by equations  $z - y^k = 0$  and  $z + y^k = 0$ .  $\square$

**Claim 18.** *The covering  $p_2$  is ramified along  $\mathfrak{R}$  with multiplicity two. The degree of the curve  $\mathfrak{B}$  is  $\deg \mathfrak{B} = 18$ . The curve  $\mathfrak{B}$  intersects transversally with  $\tilde{R}$  in 24 points belonging to the set  $\mathfrak{B}_{\tilde{R},1}$ , and, in addition,  $\tilde{R}$  and  $\mathfrak{B}$  have 42 common points of simple tangency belonging to the set  $\mathfrak{B}_{\tilde{R},2}$ .*

*Proof.* We have  $p_2(R) = \tilde{R}$ , and from the results of subsection 3.3, it follows that  $p_2 : R \rightarrow \tilde{R}$  is a double covering branched in 24 points belonging to  $\mathfrak{B}_{\tilde{R},1}$ , and it follows from Remarks 3 that for any point  $\bar{z} \in \tilde{R}$  the preimage  $p_2^{-1}(\bar{z})$  consists of not less than two points. Therefore,  $\mathfrak{B}$  does not contain irreducible components along which the covering  $p_2$  is ramified with multiplicity three, since  $\deg p_2 = 3$  and the plane curve  $\tilde{R}$  has nonempty intersection with each irreducible component of the curve  $\mathfrak{B}$ . Moreover, locally at each point  $q \in \mathfrak{R} \cap R$ , the morphism  $p_2$  is a two-sheeted covering. Hence, by Lemma 2, curves  $\mathfrak{R}$  and  $\mathfrak{B}$  are nonsingular in their intersection points, respectively, with  $R$  and  $\tilde{R}$ .

According to the adjunction formula we have  $K_I = p_2^*(K_{\mathbb{P}^2}) + \mathfrak{R}$ . Therefore, since  $p_2 : R \rightarrow \tilde{R}$  is a two-sheeted covering,  $\deg \tilde{R} = 6$  and  $(K_I, R)_I = 30$ , then

$$(\mathfrak{R}, R)_I = 30 - 2 \deg K_{\mathbb{P}^2} \cdot \deg \tilde{R} = 66. \quad (18)$$

Since  $p_2 : R \rightarrow \tilde{R}$  is ramified at 24 points belonging to the set  $\mathfrak{B}_{\tilde{R},1}$ , then  $\mathfrak{B}_{\tilde{R},1} \subset \mathfrak{B}$ . And from Remark 2 it follows that curves  $\tilde{R}$  and  $\mathfrak{B}$  also intersect at 42 points from  $\mathfrak{B}_{\tilde{R},2}$ , in which, according to Lemma 2, they touch each other. Hence (applying Lemma 2 again), we obtain that  $(\mathfrak{R}, R)_I \geq |\mathfrak{B}_{\tilde{R},1}| + |\mathfrak{B}_{\tilde{R},2}| = 66$ . Therefore, to complete the proof of Claim 18, it is enough to compare this inequality with equality (18), apply Lemma 2 again, and get equality:  $6 \deg \mathfrak{B} = (\tilde{R}, \mathfrak{B})_{\mathbb{P}^2} = |\mathfrak{B}_{\tilde{R},1}| + 2|\mathfrak{B}_{\tilde{R},2}| = 108$ .  $\square$

**Proposition 8.** *The irregularity  $q = \dim H^1(I, \mathcal{O}_I)$  of the surface  $I$  vanishes.*

*Proof.* Denote  $D = p_2^*(L)$ , where  $L$  is a line in  $\mathbb{P}^2$ . The divisor  $D$  is ample, the complete linear system  $|D|$  is free from fixed components and base points, and generic element of this system is a nonsingular curve, since  $p_2$  is a finite morphism on  $\mathbb{P}^2$ . Hence,  $\dim H^0(I, \mathcal{O}_I(D)) = 3$ , since  $(D^2)_I = 3$  and if  $\dim H^0(I, \mathcal{O}_I(D)) = r > 3$ , then the complete linear system  $|D|$  defines a birational map to a surface of degree

three in  $\mathbb{P}^{r-1}$ , which is birationally isomorphic to a ruled surface that is impossible, since  $\chi(\mathcal{O}_I) = 9$  and the Euler characteristic of structure sheafs of ruled surfaces is less than two. In addition, we have

$$(D, K_I)_I = (D, p_2^*(K_{\mathbb{P}^2}) + \mathfrak{R})_I = 3(L, K_{\mathbb{P}^2})_{\mathbb{P}^2} + (L, \mathfrak{B})_{\mathbb{P}^2} = 9.$$

Hence, a generic curve in the linear system  $|D|$  has the geometric genus  $g(D) = 7$ . Note also that from the definitions of the divisor  $D$  and the exceptional curves  $E_i$ , it follows that  $(D, E_i)_I = (K_I, E_i)_I = 1$  for all  $i = 1, \dots, 21$ .

Denote  $\hat{D} = K_I - D$  and let  $\dim H^1(I, \mathcal{O}_I(D)) = q(D)$ . We have  $(\hat{D}^2)_I = 3$  and  $(D, \hat{D})_I = 6$ . It follows from the Riemann – Roch Theorem and Serre duality that

$$\dim H^0(I, \mathcal{O}_I(D)) + \dim H^0(I, \mathcal{O}_I(\hat{D})) = \frac{(D - K_I, D)_I}{2} + p_a(I) + q(D).$$

Hence,  $\dim H^0(I, \mathcal{O}_I(\hat{D})) = 3 + q(D)$ . Note also that the linear system  $|\hat{D} - D|$  is empty, since  $(\hat{D} - D, E_i)_I = -1$  for all  $i = 1, \dots, 21$  and if the linear system  $|\hat{D} - D|$  is not empty, then all curves  $E_i$  are its fixed components. On the other hand,  $(\hat{D} - D, D)_I = 3$  and  $D$  have a positive intersection with any effective divisor. We get a contradiction.

Let us show that  $q(D) = 0$ . Assume the opposite and consider a (possibly rational) map  $\varphi_{|\hat{D}|} : I \rightarrow \mathbb{P}^{2+q(D)}$  defined by the complete linear system  $|\hat{D}|$ . Let  $N + M \in |\hat{D}|$ , where  $N$  is the (nonempty) fixed part of the linear system  $|\hat{D}|$ . Then  $\varphi_{|\hat{D}|}$  maps curves of genus  $g(D) = 7$  from the linear system  $|D|$  either birationally onto curves in  $\mathbb{P}^{2+q(D)}$  of degree  $m = (D, M)_I \leq 5$ , or two-sheeted (for  $m = 4$ ) onto curves of degree two, or  $m$ -sheeted onto lines. In all cases, images do not lie in a proper linear subspaces of the space  $\mathbb{P}^{2+q(D)}$ , since the linear system  $|\hat{D} - D|$  is empty. Therefore, the last two cases are impossible, and the first case is also impossible, since the geometric genus of curves of degree  $m \leq 5$  does not exceed six (to see this, it is enough to project birationally the image of  $D$  to the plane). Hence, the linear system  $|\hat{D}|$  has no fixed components.

Note that if  $p \in I$  is a base point of the linear system  $|\hat{D}|$ , then all (except perhaps one) curves of this linear system are nonsingular at  $q$ , since  $(\hat{D}^2)_I = 3$  and the linear system  $|\hat{D}|$  has no fixed components. Therefore, the linear system  $|\hat{D}|$  cannot be composed from a pencil and, therefore,  $\varphi_{|\hat{D}|}$  maps the surface  $I$  to a surface not lying in a proper linear subspace of the space  $\mathbb{P}^{2+q(D)}$ . It follows that  $2 \leq \deg \varphi_{|\hat{D}|}(I) \leq 3$  and  $\varphi_{|\hat{D}|} : I \rightarrow \varphi_{|\hat{D}|}(I)$  is a birational map, which is impossible, since  $\chi(\mathcal{O}_I) = 9$ . Hence,  $q(D) = 0$ .

Let us show that  $\dim H^0(D_0, \mathcal{O}_{D_0}(D)) = 2$  for a non-singular curve  $D_0 \in |D|$ . Indeed, if  $\dim H^0(D_0, \mathcal{O}_{D_0}(D)) > 2$ , then the regular sections of the sheaf  $\mathcal{O}_{D_0}(D)$

determine a birational map of the curve  $D_0$  to a curve of degree three, which is impossible, since  $g(D_0) = 7$ . Therefore, Proposition 8 follows from the exact sequence

$$0 \rightarrow H^0(I, \mathcal{O}_I) \rightarrow H^0(I, \mathcal{O}_I(D)) \rightarrow H^0(D_0, \mathcal{O}_{D_0}(D)) \rightarrow H^1(I, \mathcal{O}_I) \rightarrow 0$$

and equalities  $\dim H^0(I, \mathcal{O}_I) = 1$ ,  $\dim H^0(I, \mathcal{O}_I(D)) = 3$ ,  $\dim H^0(D_0, \mathcal{O}_{D_0}(D)) = 2$  proven above.  $\square$

**4.3. Calculation of the irregularity of the variety  $\mathcal{I}$ .** Let  $\sigma : \mathfrak{J} \rightarrow \mathcal{I}$  be a resolution of singular points of the variety  $\mathcal{I}$  such that  $\sigma : \mathfrak{J} \setminus \sigma^{-1}(\mathcal{S}) \rightarrow \mathcal{I} \setminus \mathcal{S}$  is an isomorphism.

**Theorem 4.** *The irregularity  $q(\mathfrak{J}) = \dim H^1(\mathfrak{J}, \mathcal{O}_{\mathfrak{J}})$  equals zero.*

*Proof* follows from Proposition 8 and the following theorem applied to morphism  $h \circ \sigma : \mathfrak{J} \rightarrow \mathbb{P}^9$ .  $\square$

**Theorem 5.** *Let  $f : M \rightarrow \mathbb{P}^k$  be a dominant morphism of a smooth projective variety  $M$ ,  $\dim M = k \geq 3$ . If the irregularity of the surfaces  $f^{-1}(\Pi)$  vanishes for the projective planes  $\Pi \subset \mathbb{P}^k$  belonging to an open everywhere dense in  $Gr(3, k+1)$  set  $W$ , then the irregularity  $q(M) = \dim H^1(M, \mathcal{O}_M)$  equals zero.*

*Proof.* Consider a projective variety  $N \subset \mathbb{P}^n$  and its some Zariski-open subset  $N_0$  contained in  $N \setminus \text{Sing } N$ . Let us introduce the following notations:

$Gr_2(V)$  is the Grassmannian of two-dimensional vector subspaces of a vector space  $V$ ;

$\mathfrak{Gr}_2(N_0) = \{(p, V) \mid p \in N_0, V \in Gr_2(T_{N_0, p})\}$ , where  $T_{N_0, p}$  is the the tangent space to  $N_0$  in the point  $p$ ;

$\mathfrak{P}_2(N_0) = \{(p, \Pi) \in N_0 \times Gr(3, n+1) \mid p \in \Pi \subset N\}$ , where  $\Pi$  are projective planes in  $\mathbb{P}^n$ .

Obviously, the map  $(p, \Pi) \mapsto (p, T_{\Pi, p})$  defines an imbedding  $i_N : \mathfrak{P}_2(N) \hookrightarrow \mathfrak{Gr}_2(N)$  and in the case when  $N = \mathbb{P}^k$ , the map  $i_{\mathbb{P}^k} : \mathfrak{P}_2(\mathbb{P}^k) \hookrightarrow \mathfrak{Gr}_2(\mathbb{P}^k)$  is an isomorphism.

The projection  $(p, V) \mapsto p$  defines a morphism  $\text{pr}_{N,1} : \mathfrak{Gr}_2(N_0) \rightarrow N_0$ , fibres of which over points  $p \in N_0$  have dimension  $\dim \text{pr}_{N,1}^{-1}(p) = 2(\dim N - 2)$ . Therefore,

$$\dim \mathfrak{Gr}_2(N) = 3 \dim N - 4. \quad (19)$$

Assume that  $q(M) > 0$ . Then on  $M$  there exists a nonzero holomorphic 1-form  $v$ . Denote  $K = \{p \in M \mid \ker v(p) = T_{M, p}\}$ , where  $v(p)$  is a linear mapping  $v(p) : T_{M, p} \rightarrow \mathbb{C}$  defined by the pairing  $T_{M, p} \times T_{M, p}^* \rightarrow \mathbb{C}$ . Let  $B$  be a hypersurface in  $\mathbb{P}^k$  such that  $f : M \setminus f^{-1}(B) \rightarrow \mathbb{P}^k \setminus B$  is a finite unramified covering, and denote by  $M_0 = M \setminus (f^{-1}(D))$  the open everywhere dense in  $M$  set, where  $D = f(K) \cup B \subset \mathbb{P}^k$ . Obviously, the morphism  $f : M_0 \rightarrow \mathbb{P}^k \setminus D$  induces a morphism  $f_* : \mathfrak{Gr}_2(M_0) \rightarrow \mathfrak{Gr}_2(\mathbb{P}^k \setminus D) \simeq \mathfrak{P}_2(\mathbb{P}^k \setminus D)$ .

Consider in  $\mathfrak{Gr}_2(M_0)$  the subset  $\Upsilon = \{(p, V) \in \mathfrak{Gr}_2(M_0) \mid V \subset \ker v(p)\}$  and its image  $f_*(\Upsilon) \subset \mathfrak{Gr}_2(\mathbb{P}^k \setminus D)$ . We have  $\dim \Upsilon = \dim M + \dim Gr(2, \dim M - 1) = 3k - 6$ . Therefore,

$$\dim f_*(\Upsilon) \leq 3k - 6.$$

On the other hand, let  $\text{pr}_2 : \mathfrak{P}_2(\mathbb{P}^k) \rightarrow Gr(3, k + 1)$  be the projection defined by the map  $(p, \Pi) \mapsto \Pi$ . The variety  $\mathfrak{Gr}_2(\mathbb{P}^k \setminus D) \cap \text{pr}_2^{-1}(W)$  is open and everywhere dense in  $\mathfrak{Gr}_2(\mathbb{P}^k) \simeq \mathfrak{P}_2(\mathbb{P}^k)$ . By (19), we have

$$\dim \mathfrak{Gr}_2(\mathbb{P}^k \setminus D) \cap \text{pr}_2^{-1}(W) = 3k - 4.$$

Therefore, there is a point  $(p, \Pi) \in \text{pr}_2^{-1}(W)$  that does not belong to the set  $f_*(\Upsilon)$  and, hence, the restriction of the form  $v$  to the surface  $f^{-1}(\Pi)$  does not vanish at points of  $f^{-1}(p)$ , i.e., the irregularity of the surface  $f^{-1}(\Pi)$  must be positive, that contradicts assumption that  $q(f^{-1}(\Pi)) = 0$ .  $\square$

## REFERENCES

- [1] M. Artebani, I. Dolgachev: *The Hesse Pencil of plane cubic curve*, L'Enseign. Math., **55** (2009), 235 – 273.
- [2] E. Brieskorn and H. Knörrer: *Plane algebraic curves*. Birkhäuser Verlag (1986), Basel. Boston
- [3] C. Ciliberto, F. Flamini: *On the branch curve of a general projection of a surface to a plane*, arXiv:0811.0467
- [4] H.S.M. Coxeter and W.O.J. Moser: *Generators and relations for discrete groups*, Springer-Verlag (1972), Berlin, Heidelberg, New York.
- [5] H. Hamm, Lê Dũng Tráng, *Un théorème de Zariski du type de Lefschetz*, Ann. Sci. École Norm. Sup. **6** (1973), 317 – 366.
- [6] H. Hamm, Lê Dũng Tráng, *Rectified homotopical depth and Grotendieck conjectures*, The Grotendieck Festschrift. A collection of articles written in honor of the 60th birthday of Alexander Grotendieck, Birkhauser, Boston – Basel – Berlin, vol. **II**(1991), 311 – 351.
- [7] J. Harris: *Galois groups of enumerative problems*, Duke Math. J., **46:4** (1979), 685 – 724.
- [8] O. Hesse: *Über die Elimination der Variablen aus drei algebraischen Gleichungen vom zweiten Grade mit zwei Variablen*. J. Reine Angew. Math., **28** (1844), 68 – 96.
- [9] O. Hesse: *Über die Wendepuncte der Curven dritte Ordnung*. J. Reine Angew. Math., **28** (1844), 97 – 102.
- [10] W.V.D. Hodge and D. Pedoe: (1994) [1947], *Methods of Algebraic Geometry, Volume I*, Cambridge University Press, (1947).
- [11] C. Jordan: *Mémoire sur les équations différentielles linéaires à intégrale algébrique*, J. für Reine und Angew. Math., **84** (1977), 89 – 215.
- [12] Vik.S. Kulikov: *On Chisini's conjecture*, Izv. Math., **63:6** (1999), 1139 – 1170.
- [13] Vik. S. Kulikov: *A Remark on Classical Plücker's formulae*. Ann. Fac. Sci. Toulouse. Math., **25:5** (2016), 959 – 967.
- [14] Vik.S. Kulikov: *Plane rational quartics and K3 surfaces*, Proc. Steklov Inst. Math., **294** (2016), 95 – 128.
- [15] Vik.S. Kulikov: *Dualizing coverings of the plane*, Izv. Math., **79:5** (2015), 1013 – 1042.
- [16] Vik.S. Kulikov, E.I. Shustin: *On G-Rigid Surfaces*, Proc. Steklov Inst. Math., **298** (2017), 133 – 151.

- [17] Vik.S. Kulikov: *The Hesse curve of a Lefschetz pencil of plane curves*, Russian Math. Surveys, **72:3** (2017), 574 – 576.
- [18] H. Maschke: *Aufstellung des vollen Formensystems einer quaternären Gruppe von 51840 linearen Substitutionen*, Math. Ann., **33** (1889), 317 – 344.
- [19] F. Severi: *Verlesungen über Algebraische (Anhang F)*. Leipzig, Teubner, 1921.
- [20] G.C. Shephard and J.A. Todd: *Finite unitary reflection groups*, Canadian J. Math., **6:2** (1954), 274–304.
- [21] K. Stein: *Analytische Zerlegungen komplexer Räume*, Math. Ann. **132** (1956), 63–93.

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA  
E-mail address: kulikov@mi.ras.ru