Entanglement spectroscopy of non-Abelian anyons: Reading off quantum dimensions of individual anyons

Eyal Cornfeld,^{1,2} L. Aviad Landau,¹ Kirill Shtengel,³ and Eran Sela^{1,4}

¹Raymond and Beverly Sackler School of Physics and Astronomy, Tel-Aviv University, Tel Aviv 6997801, Israel

²Department of Condensed Matter Physics, Weizmann Institute of Science, Rehovot 7610001, Israel

³Department of Physics and Astronomy, University of California, Riverside CA 92511, USA

⁴Department of Physics and Astronomy and Quantum Materials Institute,

University of British Columbia, Vancouver, B.C., V6T 1Z1, Canada

We study the entanglement spectrum of topological systems hosting non-Abelian anyons. Akin to energy levels of a Hamiltonian, the entanglement spectrum is composed of symmetry multiplets. We find that the ratio between different eigenvalues within one multiplet is universal and is determined by the anyonic quantum dimensions. This result is a consequence of the conservation of the total topological charge. For systems with non-Abelian topological order, this generalizes known degeneracies of the entanglement spectrum, which are hallmarks of topological states. Experimental detection of these entanglement spectrum signatures may become possible in Majorana wires using multicopy schemes, allowing the measurement of quantum entanglement and its symmetry resolution.

I. INTRODUCTION AND MAIN RESULT

Quantum entanglement gained a pivotal role in our understanding of many-body systems and specifically topological order¹⁻³. Its simplest measure is the entropy of entanglement between region A and region B of a bipartitioned system. In terms of the reduced density matrix of a subsystem A, $\rho_A = \text{tr}_{\text{B}}\rho$, the entanglement entropy is given by the associated entropy $S = -\text{Tr}(\rho_A \log \rho_A)$. Much more information however is encoded in the entanglement spectrum (ES), consisting of the spectrum $\{\lambda\}$ of eigenvalues of ρ_A . In topological phases, the ES provides valuable information on the physics of the edge. One may relate the ES to energy levels of an "entanglement Hamiltonian"⁴ living on the edge of region A, defined via $\rho_A = e^{-H_E}$, or $\lambda_i = e^{-\varepsilon_i}$.

The significance of the ES in topological phases was introduced in the seminal paper of Li and Haldane in 2008 which studied 2D fractional quantum Hall (FQH) ground states⁴. The low lying part of the ES was found to exhibit universal features related with the conformal field theory spectra of edge states living on the entanglement cut. This bulk-boundary correspondence was then extensively explored in a large body of studies⁵⁻¹¹ testing a variety of models such as fractional Chern insulators¹², resonating valence bond wave functions⁷, or topological spin liquids 13 ; for a recent review see Ref. 14. Along with these findings it became clear that the ES is a natural quantity for powerful numerical algorithms, based on, e.g., matrix product states or projected entangled pair states (PEPS). This general understanding of the role of the ES in topological systems included both Abelian states (e.g. the $\nu = 1/3$ FQH state) and non-Abelian states (e.g. the $\nu = 5/2$ Moore-Read state or a p + ip superconductor). Here, we take a closer look at the ES in non-Abelian states. We make progress using a topological quantum field theory language, specifically building on a 1D perspective.

The perspective of entanglement developed into a central approach towards classification of interacting topological phases, with powerful results specifically in 1D systems^{15–19}, including symmetry protected topological (SPT) states and generalizations to higher dimensions^{20,21}. Direct manifestations of topological phases have been identified in 1D as symmetry-protected degeneracies in the ES^{17} which are fingerprints of topological states in many different models $^{22-29}$. These degeneracies are a consequence of the action of symmetry on the eigenvectors of the reduced density matrix ρ_A (the "Schmidt states"). A beautiful argument¹⁷ asserts that in gapped 1D systems with a finite correlation length \mathcal{E} . every symmetry operation acts as a unitary operator on the "low energy" Schmidt states, and can be factorized into two operators $\mathcal{O}_L \mathcal{O}_R$, acting *locally* within distance ξ from the left or right ends of segment A, respectively. Thus the eigenstates of H_E provide a representation of the symmetry group. To illustrate this Ref. [17] considered the Haldane integer-spin chain. Any SU(2) spin rotation can be represented on the low lying eigenstates of H_E in terms of generators S_L and S_R acting near the two ends of region A, where $S_{tot} = S_L + S_R$. Since S_{tot} is integer, we can have two possibilities: a "trivial" phase in which both S_L and S_R are integer, or a "topological" phase with both S_L and S_R half integer. In the latter case there must be an ES degeneracy of at least 4, due to a the two-fold degeneracy associated with a half-integer spin at each end.

This should be contrasted with systems hosting non-Abelian anyons on their edges, like the Kitaev chain³⁰ with \mathbb{Z}_2 parity-number conservation, or parafermion chains³¹, where the edge operators do not support a local Hilbert space. For example in the Kitaev chain the symmetry operator measuring the total parity decomposes as $\mathcal{P}_{tot} = i\gamma_L\gamma_R$, where $\gamma_{L/R}$ are Majorana fermion modes, composed of degrees of freedom near either left or right end, which however do not support a local Hilbert space. This is a common feature of non-Abelian anyons. Generally, in the presence of symmetry, the reduced density matrix and hence H_E can be blockdiagonalized according to symmetry quantum numbers^{32–34}. Such a charge-resolution was recently applied in two-dimensional topological phases³⁵ allowing to generalize the topological entanglement entropy^{1,2} for systems protected by a symmetry.

In this paper we rely on this general symmetrydecomposition scheme, applied for the case of topological charge, defined below, in order to uncover internal structure in the ES in topological gapped phases supporting non-Abelian anyons. We demonstrate our predictions in explicit 1D lattice models of interacting anyons, and show that one can test them in experiment within available experimental setups in the context of Majorana wires.

A. Non-Abelian anyons

The idea to use the multi-dimensional, non-local space spanned by non-Abelian anyons to encode quantum information was put forward by Kitaev³⁶. Non-Abelian anyons are physically realized as guasiparticles in systems with topological order. Operations such as braiding and fusion form the basis for topological quantum computation; for a review see Ref. [37]. The universal statistical properties of anyons are encoded in their topological quantum field theory $(TQFT)^{38-40}$. Our subsequent symmetry decomposition of entanglement relies on the concept of "topological charge" or "total fusion channel", a conserved quantity in anyonic systems. Consider a collection of anyons taken from a set $\{I, a, b, c \dots\}$ satisfying certain fusion rules $a \times b = \sum_{c} N_{ab}^{c} c$, meaning that two anyons a and b can "fuse into" an anyon of type c as long as the (integer valued) fusion coefficient N_{ab}^c is finite. If $N_{aa}^c \neq 0$ for more than one fusion outcome c, then a collection of a anyons can encode information non-locally, and anyon a is said to be non-Abelian. The total topological charge of the collection of anyons can be obtained by sequentially fusing pairs of anyons ending with a single anyon a_{tot} . For N anyons of type a, the number of intermediate fusion channels grows exponentially as d_a^N , where d_a is the quantum dimension. Different choices of a sequence of fusion are related by a basis transformation. On the other hand the total fusion outcome is a conserved "topological charge".

To study the entanglement between subsystems A and B in some topological ground state $|\Psi\rangle$, it is natural to work in a basis where one fuses all anyons in each region into their total fusion channel a_A and a_B respectively. Then the total topological charge a_{tot} is determined by $a_A \times a_B = \sum_{a_{tot}} N_{a_A a_B}^{a_{tot}} a_{tot}$. While the full wave function $|\Psi\rangle$ has a well defined topological charge, importantly entangled states are a superposition of different sub-system charges a_A, a_B . We restrict our attention to the natural case of a closed system with a total topological charge corresponding to the vacuum state, *i.e.*, $a_{tot} = I$. In what follows we reserve the letter c to denote the total



FIG. 1. (a) Bipartition of a 1D system on an open segment. The wave function admits a Schmidt decomposition in terms of the boundary anyon pair a_L, \bar{a}_L and a pair a_R, \bar{a}_R at the cut between A and B. The total charge in region $A, c = a_A$, is determined by the fusion of the anyons in A. (b) Universal structure of the ES composed of many multiplets described by Eq. (1) with internal multiplicities and spacing determined by the quantum dimension of c. (c) ES of the Kitaev chain $H = \sum_{j=1}^{64-1} (c_j^{\dagger}c_{j+1} + c_j^{\dagger}c_{j+1}^{\dagger} + h.c.) + \mu \sum_{j=1}^{64} c_j^{\dagger}c_j$ for chemical potential $\mu = -3, -2.5, -2, \ldots, 3$. Here $c = 1, \psi$ takes two values with the same quantum dimension. The resulting topological degeneracy is removed at the transition, $|\mu| \geq 2$. At the sweet spot $\mu = 0$ there is a single multiplet.

fusion channel in region A, $c = a_A$, and assume $a_B = \bar{c}$. The conservation of the topological charge implies⁴¹ that ρ_A forms blocks labeled by c. Hence the ES is a union of the ES of each block with fixed topological charge. One may ask: what is the contribution to the entanglement from each charge sector³³? is there a general relation between these contributions?

For concreteness we consider 1D gapped models on an open segment. Suppose that our 1D topological phase carries an anyon of type a as edge states. One may then postulate that the wave function $|\Psi\rangle$ is described in terms of pair of anyons of type a pulled out of the vacuum across each entanglement cut between A and B. For our 1D system on an open segment we have two such pairs of anyons, as depicted in Fig. 1(a). From this picture, using the rules of TQFT we find that the ES decomposes according to the subsystem topological charge c as

$$\{\lambda\} = \bigcup_c \{\lambda\}_c, \quad \#_c = N_{aa}^c, \quad \lambda_c \propto d_c. \tag{1}$$

The ES for each charge sector c consists of a single eigenvalue λ_c with multiplicity $\#_c$. The multiplicity is just the number of possibilities for fusing the boundary anyons into the total topological charge c, $\#_c = N_{aa}^c$. Our main result is that λ_c probes the quantum dimension of the type-a anyon, $\lambda_a \propto d_a$. We can see that in general anyon systems the ES is characterized by universal ratios of its eigenvalues determined by the quantum dimensions d_c , instead of degeneracies. Equivalently, multiplets in the entanglement Hamiltonian have the same decomposition $\{\varepsilon\} = \bigcup_c \{\varepsilon\}_c$ with multiplicities $\#_c$, and with an additive entropy term

$$\delta \varepsilon_c = -\log d_c, \tag{2}$$

see Fig. 1(b). This central result demonstrates that the charge-resolved entanglement contains TQFT data, which can be interpreted as the entropy associated with a single anyon c.

A similar term appears^{1,2} as the subleading term γ , referred to as topological entanglement entropy, in 2D topologically ordered systems with a finite correlation length $S(L_A) = \alpha L_A - \gamma + \dots$ Here α is a non-universal constant, while $\gamma = \log \mathcal{D}$ is a universal number known as the total quantum dimension $\mathcal{D} = \sqrt{\sum_a d_a^2}$ characterizing the topological medium.

In generic models hosting non-Abelian anyons there are additional local degrees of freedom creating entanglement on the scale of the correlation length ξ across the entanglement cut. These are reflected in the ES by additional non-universal levels. When these local degrees of freedom are independent from the anyons, each such level becomes a multiplet with exactly the same structure as the ground state described by Eq. (1), see dashed circle in Fig. 1(b).

The Kitaev chain realizes a familiar but unfortunately non-exhaustive example of Eq. (1). It is described in terms of Ising anyons $\{I, \psi, \sigma\}$, where for open boundary conditions the edges of the chain host σ -anyons (corresponding to γ_L and γ_R above), with fusion rules $\sigma_L \times \sigma_R = I + \psi$. We thus have two topological charges Iand ψ corresponding to even and odd electron parity in region A, respectively. Accordingly the ES is decomposed as $\{\lambda\} = \{\lambda\}_I \cup \{\lambda\}_{\psi}$. However, both of anyons I and ψ are Abelian, $d_I = d_{\psi} = 1$ implying $\lambda_I = \lambda_{\psi} = 1/2$. This yields the familiar *degeneracy* in the ES of the topological phase in the Kitaev model, seen in Fig. 1(c). Thus it would be interesting to explore the ES and the prediction in Eq. (1) in models hosting more exotic anyons.

The rest of the paper is organized as follows. After deriving the main result Eq. (1) in Sec. II, in Sec. III we demonstrate its validity for specific lattice models. In contrast to the Kitaev model, in order to observe nontrivial ratios in the ES, one needs models whose edge states can fuse into a non-Abelian anyons having quantum dimension $d_a > 1$. \mathbb{Z}_n parafermions which have received considerable attention recently³¹ are richer than Majorana fermion modes since their edge states have $n \geq 2$ fusion outcomes, which themselves, however, are all Abelian anyons. Fibonnaci anyons satisfying the fusion rule $\tau \times \tau = 1 + \tau$ are rich enough to display the nontrivial structure in the ES found here. Indeed, 1D models with exotic states have been built out of these anyons^{42–44}. Here, we demonstrate our results for $SU(2)_k$ anyonic generalizations of the AKLT chain⁴⁵.

In Sec. IV we discuss an experimental protocol to detect the topological charge resolution of the entanglement. While the ES can be measured in small systems, as was recently demonstrated for a small system realizing the AKLT chain⁴⁶, here we develop methods for general systems. We focus on topological-charge resolution of entanglement measures. As a first step towards this goal, we focus on Majorana systems, specifically on their implementations in quantum wires controllable by charging energy effects⁴⁷. Various protocols have been proposed for designing entangled states $^{48-52}$ (please note that this is not a complete list). In accordance with our results, the ES in particular and other entanglement measures in general can be decomposed into degenerate even/odd sectors. Using a general protocol for measurement of charge-resolved entanglement 33,53,54 , we discuss a concrete platform for measuring the second Rényi entropy. In the topological phase hosting Majorana edge states, the separately measurable even (c = I) and odd $(c = \psi)$ contributions to the Rényi entropy become degenerate. This allows to measure experimentally the predicted topological degeneracies in the ES and assign it to the charge resolution.

II. BOUNDARY-ANYONS ENTANGLEMENT

In this section we derive Eq. (1). Bonderson *et al.*³⁹ have recently applied TQFT methods to study entanglement of anyonic systems. Here we use these methods and focus on the ES. The impatient reader may skip this TQFT-based derivation directly to Eq. 6 at the end of this technical section, and then to the next sections where its significance is exemplified.

We consider the setting depicted in Fig. 1(a), where anyons $\{a\}$ appear on the entanglement cut between regions A and B as well as on the physical boundaries. We start with a system with boundary anyons a, \bar{a} pulled out of the vacuum at the entanglement cut between regions A and B, whose joint state is represented by $|a, \bar{a}; I\rangle^R$; a second pair of anyons at the real boundary of the system is represented by $|a, \bar{a}; I\rangle^L$. The total system is in state $|\Psi\rangle = |a, \bar{a}; I\rangle^L |a, \bar{a}; I\rangle^R$ where L, R correspond to the left and right edges of region A; this state is depicted in Fig. 2(a) (here, we utilize the diagrammatic notations by Kitaev³⁸ as presented in Ref. [39]; see Appendix A). Consequently, the density matrix of the full system is

$$\rho = |a, \bar{a}; I\rangle^L |a, \bar{a}; I\rangle^R \langle a, \bar{a}; I|^L \langle a, \bar{a}; I|^R, \qquad (3)$$

see Fig. 2(b). We then partition the anyons into two sub-



FIG. 2. (a) Graphical representation of the normalized wave function (*cf.* Ref. [39]). Two pairs of anyons are fused out of the vacuum creating entanglement between A and B. (b) Normalized density matrix. (c) Reduced density matrix, separated into blocks with fixed total anyonic charge c in region A. The diagram is equivalent to two parallel lines $[||]_c$ representing the identity operator in the Hilbert subspace of two anyons a_L, \bar{a}_R , fused into c; see Eq. (4).

systems $A = a_L, \bar{a}_R, B = a_R, \bar{a}_L$, and proceed by tracing out subsystem B. This leads to a diagram representing the identity in the Hilbert space spanned by two anyons a_L, \bar{a}_R . Our main step here is facilitated by the decomposition of this space according to the total fusion channel c. This is diagrammatically represented in Fig. 2(c), and results in

$$\rho_A = \operatorname{Tr}_B \rho = \sum_c \frac{d_c}{d_{a_L} d_{\bar{a}_R}} \left[\mathbb{I}_{a_L, \bar{a}_R} \right]_c, \qquad (4)$$

where the square brackets describe the identity operator in the two-anyon Hilbert subspace with total fusion channel c^{39} . As discussed in Appendix B, this is given by

$$[\mathbb{I}_{a_L,\bar{a}_R}]_c = \sum_{\mu=1}^{N_{a_L\bar{a}_R}^c} |a_L\bar{a}_R;c,\mu\rangle\langle a_L\bar{a}_R;c,\mu|, \qquad (5)$$

where generally a_L and \bar{a}_R can fuse into c in multiple ways labelled by μ . This pair of equations yields the spectrum and degeneracies in Eq. (1),

$$\lambda_c = \frac{d_c}{d_{a_L} d_{\bar{a}_R}}, \qquad \#_c = N^c_{a_L, \bar{a}_R}.$$
 (6)

The normalization of the density matrix follows from the fusion algebra

$$\operatorname{Tr}\rho_A = \sum_c \frac{N_{a_L\bar{a}_R}^c d_c}{d_{a_L} d_{\bar{a}_R}} = 1.$$
 (7)

Below we demonstrate the result Eq. (6), illustrated in Fig. 1(b), in concrete models of interacting anyons.

III. ANYONIC CHAINS

To test the topological field theory results, in this section we study a lattice model of interacting anyons. Before focusing on a specific model in Sec. III A, we first



FIG. 3. An open chain of L anyons. We choose the total topological charge $a_{tot} = 0$. The total topological charge of region A, $c = a_A$ is given by the link connecting regions A and B.

discuss generalities about the construction of the ES in anyonic models.

Consider a system of anyons, some of which are located in region A, and the rest in B. The structure of a factorizable Hilbert space $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$, which is usually taken for granted, is actually not present, since each individual anyon does not posses a local Hilbert space. However, for every value of the total fusion channel a_A and a_B in each region, there is a local Hilbert space $\mathcal{H}_A^{a_A}$ and $\mathcal{H}_B^{a_B}$, respectively. Thus, for a specific value of the total charge a_{tot} , the Hilbert space is a *direct sum* of factorizable spaces, consistent with the fusion rules $\mathcal{H} = \bigoplus_{a_A, a_B, N_{a_A a_B}^{a_{tot}} = 1} \mathcal{H}_A^{a_A} \otimes \mathcal{H}_B^{a_B}$ (here for simplicity $N_{ab}^c = 0, 1$ for all a, b, c).

Following Singh *et al.*⁴¹ consider a chain of *L* anyons denoted $\{X_i\}$ (i = 1, ..., L), whose state can be represented by the fusion diagram in Fig. 3. A family of such 1D models was introduced in Refs. [42–45]. This basis of states is denoted as $|x_1, ..., x_{L-1}\rangle_{x_0x_L}^{X_1,...,X_L} = |\{x\}\rangle_{x_0x_L}^{\{X\}}$, with external legs $\{X\}$ corresponding to the *L* anyons, and with fixed boundary anyons x_0 and x_L . The fusion rule constraints $N_{x_i,X_{i+1}}^{x_{i+1}} \neq 0$ have to be satisfied at each vertex. Setting $x_0 = I$ to the vacuum state, we can identify x_L with the total fusion channel a_{tot} . We set it to the vacuum state $a_L = a_{tot} = I$.

Consider dividing the chain into region A containing

anyons $X_1
dots X_i$ and region B containing $X_{i+1}
dots X_L$. It is evident that the link x_i connecting A and B carries the topological charge of region A, hence we identify $x_i = a_A = c$. For each value of c, each basis vector may be partitioned into the two regions as $|\{a\}, c, \{b\}\rangle^{\{A\}, \{B\}} = |\{a\}\rangle_{0c}^{\{A\}} \otimes |\{b\}\rangle_{c0}^{\{B\}}$. This set of states with fixed boundary condition specified by c defines the Hilbert space for A and B, $\mathcal{H} = \bigoplus_c \mathcal{H}_A^c \otimes \mathcal{H}_B^{\overline{c}}$ and allows us to write any state $|\Psi\rangle = \sum_{\{x\}} \Psi_{\{x\}} |\{x\}\rangle$ as

$$|\Psi\rangle = \sum_{c} \sum_{\{a\},\{b\}} [\Psi_c]_{\{a\},\{b\}} |\{a\}\rangle_{0c}^{\{A\}} \otimes |\{b\}\rangle_{c0}^{\{B\}}.$$
 (8)

To obtain the ES one may treat the wavefunction $\Psi_{\{\{a\},c,\{b\}\}}$ as a matrix $[\Psi_c]_{\{a\},\{b\}}$ of (super-)indices $\{a\},\{b\}$, representing the *c* component of the wavefunction, and perform a singular value decomposition (SVD) on that matrix,

$$|\Psi\rangle = \sum_{c} \sum_{j} [\Lambda_c]_j |\Phi_j^A\rangle_{0c}^{\{A\}} \otimes |\Phi_j^B\rangle_{c0}^{\{B\}}.$$
 (9)

Here, the vectors $|\Phi_j^A\rangle_{0c}^{\{A\}}$ and $|\Phi_j^B\rangle_{c0}^{\{B\}}$ labeled by j belong to \mathcal{H}_A^c and \mathcal{H}_B^c , respectively. We now turn to an explicit example.

A. Anyonic-AKLT chain

We consider the anyonic- $SU(2)_k$ AKLT chain⁴⁵ as a concrete example. The model contains a family of anyons labeled by their integer "spin" quantum number j = 0, 1, ..., |k/2|. These are generally non-Abelian anyons, with fusion rules $j_1 \times j_2 = |j_1 - j_2|, \dots, \min\{j_1 + j_2\}$ $j_2, k - j_1 - j_2$, extending the usual SU(2) rules. The j = 0 anyon corresponds to the vacuum state I with zero topological charge. The chain consists of L anyons of type j = 1, *i.e.*, $X_i = 1$ $(i = 1, \dots L)$ and the $\{x_i\}$ should be consistent with the fusion rules $x_{i-1} \times X_i =$ x_i . The model contains an "AKLT sweet spot" analogous to the familiar AKLT chain⁵⁵ with Hamiltonian By the formula $H_{\text{sweet spot}} = \sum_{i=1}^{L-1} P^{(2)}(X_i, X_{i+1})$, consisting of projectors of neighbouring j = 1 anyons X_i, X_{i+1} onto the j = 2 fusion channel. A parameter θ_{12} describes deviations from the AKLT sweet spot whereby an additional term $\propto \sum_{i} P^{(1)}(X_i, X_{i+1})$ appears in the Hamiltonian

$$H = \sum_{i=1}^{L-1} [P^{(2)}(X_i, X_{i+1}) \cos \theta_{12} - P^{(1)}(X_i, X_{i+1}) \sin \theta_{12}].$$
(10)

Using F-moves one can write the Hamiltonian in the basis of Fig. 3^{45} ; see Fig. 5(b).

We focus on the $SU(2)_k$ model with odd $k \ge 5^{45}$ and discuss explicitly the case of k = 5, with 3 anyons types



FIG. 4. ES for the $SU(2)_5$ anyonic-AKLT ground states at $\theta_{12} = 0$. Purple and red correspond to fusion channel c = 0, 1, respectively. Closed expression given at Eq. (14).



FIG. 5. The F-move anyonic change of basis.

j = 0, 1, 2 of quantum dimensions d_j

$$d_0 = 1,$$
 $d_1 = \frac{\sin(3\pi/7)}{\sin(\pi/7)},$ $d_2 = \frac{\sin(5\pi/7)}{\sin(\pi/7)}.$ (11)

These anyons satisfy the fusion rules

$$1 \times 1 = 0 + 1 + 2, \qquad 1 \times 2 = 1 + 2, \qquad 2 \times 2 = 0 + 1,$$
 (12)

along with $0 \times x = x$ (x = 0, 1, 2). The following results are obtained by exact diagonalization of this Hamiltonian, supplemented by analytic treatment at the AKLT sweet spot.

1. Entanglement of the AKLT sweet spot

The ES for an equipartition of the ground state of the open chain at the AKLT sweet spot is shown in Fig. 4 for various values of L. There are only two eigenvalues with a total fusion channel of region A given by c = 0 and c = 1 (purple and red respectively). At sufficiently



FIG. 6. (a) Exact ground state for the AKLT anyonic chain with PBC^{45} . (b,c,d): Through a sequence of F-moves, the loop can be brought into contractible form. After its contraction one obtains the diagram with OBC Fig. 3. This gives the exact ground state Eq. (13).

large L their ratio approaches the ratio of the quantum dimensions d_0/d_1 . This result may seem surprising in the absence of the underlying boundary anyons assumed in Fig. 1. Through an exact solution, however, we shall see that boundary anyons do indeed emerge and, accordingly, the resulting ES is in line with our main result, Eq. (6).

2. Exact solution

To find the exact ground state for the AKLT chain with open boundary conditions (OBC), we start from the exact ground state that can be readily $guessed^{45}$ for periodic boundary conditions (PBC), see Fig. 6(a). The state depicted in Fig. 6(a) is a ground state since each pair of adjacent j = 2 anyons can never fuse into j = 2according to the fusion rules Eq. (12), yielding a zero energy state which is a ground state of the positive definite Hamiltonian $H_{\theta_{12}=0}$. Below we discuss its uniqueness. In order to obtain the corresponding ground state for OBC in the basis $|\{x\}\rangle_{x_0=0,x_L=0}$ of Fig. 3, we sequentially shrink and finally annihilate the loop. This is done by applying F-moves according to the rule in Fig. 5(c), and is implemented in Fig. 6(b,c,d). This gives back the OBC diagram of Fig. 3. The resulting state is explicitly given in terms of the F-matrices⁴⁵ by

$$\Psi_{\{x_i\}} \propto \prod_{i=0}^{L-1} (F_2^{x_i,1,2})_2^{x_{i+1}}.$$
(13)

This is the unique ground state of the total topological sector $a_{tot} = 0$. In comparison, the usual spin-1 AKLT Hamiltonian⁵⁵ has 4 ground states, obtained by the sin-

glet and triplets that the two boundary spin-1/2-s can form. In our anyonic model in Fig. 3 we have imposed a total anyonic charge 0, analogous to restricting ourselves to the the AKLT singlet state which is also unique.

As shown in Appendix C, one can analytically obtain the ES from this state. Specifically, for a bipartition of a chain with L anyons to regions $A \cup B$ consisting of $L_A + L_B = L$ anyons, we find

$$\lambda_{0} = \frac{1}{d_{2}^{2}} \cdot \frac{(1 - (-d_{1})^{1 - L_{A}})(1 - (-d_{1})^{1 - L_{B}})}{(1 - (-d_{1})^{1 - L_{A} - L_{B}})},$$

$$\lambda_{1} = \frac{d_{1}}{d_{2}^{2}} \cdot \frac{(1 - (-d_{1})^{-L_{A}})(1 - (-d_{1})^{-L_{B}})}{(1 - (-d_{1})^{1 - L_{A} - L_{B}})},$$

$$\lambda_{2} = 0.$$
(14)

One clearly sees that at the thermodynamic limit $L \to \infty$ the spectrum corresponds with our main TQFT result, Eq. (6), and $\lambda_{0,1} \to d_{0,1}/d_2^2$. This is plotted in Fig. 4 and matches the exact diagonalization numerics.

3. Boundary anyons

In Fig. 6(a), which is the PBC representation of our OBC state, we can see two j = 2 anyon lines connecting regions A and B, see Fig. 7. We associate them with the boundary anyons $a_L = a_R = \bar{a}_L = \bar{a}_R = 2$. Since the state is gapped, in the thermodynamic limit these boundary anyons are decoupled. The fusion rule $2 \times 2 = 0 + 1$ then dictates that the total charge of region A can only take on values c = 1, 2, which explains the structure of the ES.



FIG. 7. Visualisation of the "hidden" boundary anyons of the OBC model depicted in Fig. 3 using the PBC representation of Fig. 6(a).

4. ES beyond the AKLT sweet spot

Universal multiplet structure of the ES described by the $SU(2)_k$ TQFT persists beyond the AKLT sweet spot, everywhere within the topological phase of the model which extends approximately over the range of $-0.19 \leq \theta_{12}/\pi \leq 0.06^{45}$. Our results for the ES, obtained via exact diagonalization of the Hamiltonian followed by an SVD, are shown in Fig. 8 as a function of θ_{12} for L = 21. The large L convergence of the various multiplets is demonstrated in Fig. 9. One can see that in addition to the dominant c = 0, 1 doublet at the AKLT sweet spot $\theta_{12} = 0$, additional multiplets appear at finite θ_{12} . This is similar to the behaviour of the ES in the Kitaev chain away from the sweet spot, *cf.* Fig. 1(c).

The finite size results are in good agreement with the statement that the ratio between eigenvalues of states with charge c within each multiplet is proportional to the quantum dimension d_c .

The results can be understood as a generalization of Eq. (6), which itself assumes well defined boundary anyons a_L , a_R . Starting from the OBC chain at finite θ_{12} at the ground state, and going in reverse through the steps in Fig. 6, one would obtain the diagram in Fig. 6(a)with two anyons lines connecting A and B, cf. Fig. 7. Only at the AKLT sweet spot they carry a well defined charge j = 2, however away from the AKLT sweet spot the state $|\Psi\rangle$ is a linear superposition over the possible values of a_L , a_R . Even in this case, we show in Appendix B that a multiplet structure emerges. Each multiplet is characterized by a pair of boundary anyons a_L, \bar{a}_R , with possible fusion outcome c determined by their fusion rules. While the ground state multiplet corresponds to $(a_L, \bar{a}_R) = (2, 2)$, the next multiplet and its multiplicity follows from $(a_L, \bar{a}_R) = (2, 1)$ fusing into c = 1, 2 with unit multiplicity.

IV. EXPERIMENTAL TESTS

Testing the anyonic-charge-resolved entanglement in an experiment requires (i) physical systems realizing non-Abelian anyons and (ii) protocols for practical measurement of quantum entanglement. In this section we discuss controllable experimental setups based on semicon-



FIG. 8. ES for the model Eq. (10) as a function of θ_{12} parametrizing deviations from the AKLT sweet spot. Purple, red, and blue correspond to fusion channel c = 0, 1, 2, respectively. Results are displayed in both linear (top) and logarithmic (bottom) scales. The multiplet structure of Eq. (1) is evident within the topological phase, *cf.* Fig 1.

ductor quantum wires, which could realize Ising anyons. We propose to employ the schemes of Refs. [56,57] to measure the n-th Rényi entropy (RE), $s_n = \text{Tr}\rho_A^n$, which were demonstrated in a cold-atom experiments^{58,59}; for a recent work detailing this protocol, including the case of symmetry resolution, see Ref. [53]. Our suggestion is that testing the universal internal structure of the ES can be achieved by extending these schemes to measure the





FIG. 9. Convergence analysis of the results in Fig. 8.

charge-resolved entanglement $^{32-34}$.

The predicted universal structure in the ES of topological anyonic systems is reflected in the charge-resolved RE: The separation of the ES into symmetry sectors $\{\lambda\} = \bigcup_c \{\lambda\}_c$ implies the additive structure of the RE, $s_n = \sum_i \lambda_i^n = \sum_c s_n(c)$. We refer to $s_n(c)$ as the chargeresolved RE. Assuming that it is dominated by the largest eigenvalues in $\{\lambda\}_c$, for which the low-energy TQFT result Eq. (1) holds, we obtain the approximate relationship $s_n(c) \propto \#_c d_c^n$. Thus measuring the ratio between various charge-resolved contributions to the RE gives access to universal topological data.

In this section we use this simple observation to demonstrate, for systems realizing Ising anyons, that the known degeneracies in the ES can actually be measured. In this case $\#_I = \#_{\psi} = 1$ and $d_I = d_{\psi}$, yielding the familiar degeneracy in the ES, thus implying that the even and odd charged-resolved REs are equal. For Kitaev's Majorana chain model the degeneracy is symmetry protected thus this relation between the charged-resolved REs is exact.

This is explicitly shown in Fig. 10, where the same model parameters as in Fig. 1(c) are used to compute the parity resolved^{33,53} second REs, $s_2(c = I) = s_2(\text{even})$ and $s_2(c = \psi) = s_2(\text{odd})$, on which we concentrate in this section. One can see that as the system enters the topological phase by tuning the chemical potential to the regime $|\mu| < 2$, the difference between the even- and odd-REs vanishes up to corrections exponentially small in the system size.

Thus, the symmetry resolved measurement of the RE can serve as an order parameter for the topological phase:

FIG. 10. Parity resolved second Rényi entropy for the Kitaev chain. Dependence of the difference $[s_2(I) - s_2(\psi)]$ on the chemical potential for an equipartition of an open chain with total length L = 32, 64, 128, cf. Fig. 1(c).

While the total entanglement entropy is just a nonuniversal number, when symmetry resolved, this number is exactly equipartitioned in the topological phase.

We note, that a similar entanglement-based order parameter appears in the Schmidt gap⁶⁰ (*i.e.*, the difference between the two largest eigenvalues of the reduced density matrix) as was found for the an equivalent transverse field Ising model, without explicitly decomposing the ES into symmetry sectors.

A. Measurement of entanglement after fusion of Ising anyons in a quantum wire

Consider a quantum wire with spin-orbit coupling on top of which a floating superconductor is deposited. Upon application of a magnetic field a transition to topological superconductivity hosting Majorana end modes is expected⁶¹, as suggested, *e.g.*, in a recent experiment by Albrecht *et al.* by the field evolution of the Coulomb blockade peaks⁶². This putative topological superconducting state supports Majorana fermion modes below the gap allowing either even or odd fermion parity.

Aasen et al.⁵⁰ envisioned interesting manipulations of this system allowing to entangle Majorana fermion modes. Consider the system in Fig. 11(a) of a quantum wire Josephson-coupled to bulk superconducting leads. In the Coulomb blockade regime the even and odd ground states are split by Coulomb interactions, thus one may initialize an even ground state. Upon Josephson-coupling the wire to the bulk superconductors, charging effects become negligible, yielding an approximate even-odd degeneracy associated with the Majorana fermions⁵⁰. The switching-on of the Josephson coupling can be thought of as production of a pair of Majorana fermions denoted γ_1, γ_4 at the end of the wire, in the vacuum fusion channel. Now, a subsequent division of the wire into two segments denoted A and B by a central barrier, can be described as pulling the second pair of Majorana fermions γ_2 , γ_3 from the vacuum. Identifying Majorana fermion modes with Ising anyons, the resulting anyonic wave function, depicted at the bottom of Fig. 11(a), realizes the same state depicted in Fig. 1(a) with two pairs of Ising anyons produced from the vacuum. They create entanglement between two subsystems A and B, which we wish to explicitly measure.

The resulting state can be written as $|0_{14}0_{23}\rangle$, where $0_{ij}(1_{ij})$ is the even (odd) fusion channel of Majorana fermion modes γ_i and γ_j . This corresponds to the fusion of Ising anyons $\sigma \times \sigma$ with outcome $I(\psi)$. Expressing this state in the occupancy basis of two complex fermions $\gamma_1 + i\gamma_2$ and $\gamma_3 + i\gamma_4$ in regions A and B, respectively, the same state takes the form

$$|0_{14}0_{23}\rangle = \frac{1}{\sqrt{2}} \Big(|0_{12}0_{34}\rangle + |1_{12}1_{34}\rangle \Big).$$
(15)

This realizes a maximally entangled Bell state between the left and right segments of the wire.

To see the degeneracy of the ES explicitly, let us focus on the subsystem A - the left wire segment in Fig. 11(a), containing Majorana fermion modes γ_1 and γ_2 . Tracing out region B results in the reduced density matrix

$$\rho_A = \frac{1}{2} \Big(|0_{12}\rangle \langle 0_{12}| + |1_{12}\rangle \langle 1_{12}| \Big), \tag{16}$$

implying $\lambda_I = \lambda_{\psi} = \frac{1}{2}$.

In contrast to this entangled state, one may prepare a non-entangled state $\rho'_A = |0_{12}\rangle\langle 0_{12}|$ by starting with two segments individually dominated by Coulomb interactions, hence having fixed charge, and only then opening the Josephson couplings to the bulk superconductors⁵⁰. Below we show how one can distinguish these states by explicitly measuring their parity-resolved second Rényi entropy.

B. Two-wire measurement setup

Following the protocol to measure the Rényi entropy⁵⁷, we add an identical wire hosting 4 Majorana modes, and prepare the same quantum state in these wires. In the final state of the system, the Josephson couplings dominate over the charging energies so that charge may fluctuate. We use this to transfer charge between the wires via the Majorana modes. We label the two identical subsystem by i = a, b, and refer to them as copies. Subsystem A in each copy has two Majorana operators $\gamma_1^{a/b}$ and $\gamma_2^{a/b}$. Thus for the Bell state Eq. (16) the 2-copy reduced

FIG. 11. (a) Creation of an entangled state of Ising anyons in a quantum wire following Ref. [50]. Upon raising the central barrier, starting from a well define total charge state, one obtains the Bell state Eq. (15) in which the parity of the left and right segments, comprising regions A and B, respectively, are maximally entangled. (b) Two-copy system realizing a protocol to measure the parity-resolved second RE. The two copies are coupled by single particle tunnelling, J_{ab} , via the Majorana fermion modes, and then the parity of region A is measured, *e.g.*, by tunnel coupling (t_1) the Majorana fermion modes to two leads $\Psi_{1,2}$, as well as an interference arm t_0 . A similar parity measurement device has to be connected to the top wire.

density matrix is a product state

$$\rho_A^a \otimes \rho_A^b = \frac{1}{4} \Big(|0_a 0_b\rangle \langle 0_a 0_b| + |0_a 1_b\rangle \langle 0_a 1_b| + |1_a 0_b\rangle \langle 1_a 0_b| + |1_a 1_b\rangle \langle 1_a 1_b| \Big), \quad (17)$$

where we dropped the Majorana indices for brevity. The next step of the measuring protocol is to operate the following tunnelling Hamiltonian

$$H_{ab} = iJ_{ab}(t)\gamma_1^a\gamma_1^b + iJ_{ab}(t)\gamma_2^a\gamma_2^b, \qquad (18)$$

see Fig. 6(b). If we turn on this coupling for time $t = \frac{\pi\hbar}{4J_{ab}}$ then the states of the system evolve into

$$\begin{aligned} |0_a 0_b\rangle &\mapsto |0_a 0_b\rangle, \tag{19} \\ |1_a 0_b\rangle &\mapsto \frac{1}{\sqrt{2}} \Big(|1_a 0_b\rangle + |0_a 1_b\rangle \Big), \\ |0_a 1_b\rangle &\mapsto \frac{1}{\sqrt{2}} \Big(|1_a 0_b\rangle - |0_a 1_b\rangle \Big), \\ |1_a 1_b\rangle &\mapsto |1_a 1_b\rangle. \end{aligned}$$

Defining fermion a as $\gamma_1^a = \frac{a^{\dagger} + a}{\sqrt{2}}$, $\gamma_2^a = \frac{a^{\dagger} - a}{\sqrt{2}i}$, and the same for copy b with $a \leftrightarrow b$, this transformation is equivalent to⁵⁷

$$a \mapsto \frac{a+b}{\sqrt{2}}, \qquad b \mapsto \frac{b-a}{\sqrt{2}}.$$
 (20)

The final stage of the protocol is measuring the occupancy of subsystems a, b. Following Refs. [63,54] we find that the second RE, $s_2 = \text{Tr}\rho_A^2$, is given by: (i) preparing the 2-wire setup, (ii) performing the (Fourier) transformation of Eqs. (19) and (20), (iii) measuring the

N_a	N_b	f(N)	$f_{\rm even}(N)$	$f_{\rm odd}(N)$
0	0	1	1	0
0	1	0	0	0
1	0	0	0	0
1	1	1	0	1

TABLE I. Measuring protocol for the second RE in the even and odd parity fusion sector.

occupancies in each copy N_a, N_b , and (iv) repeating steps (i-iii) and averaging over a function f(N) where

$$f(N) = (-1)^{\frac{1}{2}(N_a + N_b) + N_a} \delta_{\frac{1}{2}(N_a + N_b) \in \mathbb{N}} = \delta_{\frac{1}{2}(N_a + N_b) \in \mathbb{N}},$$
(21)

see Table I. The last equality is a consequence of having only two Majorana wires present in the system, hence the occupancies $N_{a,b}$ are either 0 or 1. Moreover, since the total occupancy is invariant under the transformation we may immediately read off the second RE for each fusion sector, $s_2(\text{even/odd}) = \text{Tr}[\rho_A^2 \delta_{P,\text{even/odd}}]$, where $(-1)^P \equiv (-1)^N$ is the parity operator (P = 0, 1). We thus find

$$f_P(N) = \delta_{\frac{1}{2}(N_a + N_b), P},$$
 (22)

see Table I.

We now describe the occupancy measurement protocol for copy b; copy a should be handled analogously. Each one of the Majorana fermion modes should be coupled to externally linked normal leads described by the Hamiltonian^{51,64}

$$H_{\text{parity}} = t_0 \Psi_1^{\dagger} \Psi_2 + t_1 \Psi_1^{\dagger} \gamma_1^b + t_2 \Psi_2^{\dagger} \gamma_2^b + \text{H.c.}, \qquad (23)$$

see Fig. 11(b). This results in the effective tunnelling Hamiltonian

$$H_T = (t_0 + it_{12}\gamma_1^b\gamma_2^b)\Psi_1^{\dagger}\Psi_2 + \text{H.c.}, \qquad (24)$$

where $t_{12} \sim \frac{t_1 t_2}{E_c}$ with E_c the charging energy. Measuring the conductance, it depends on the interference term $\sim t_0 t_{12} \langle \gamma_1^b \gamma_2^b \rangle$, allowing to measure the occupancy in b. Using Eq. (22), one may average the measurement results $\{0, 1\}$ to obtain the second Rényi entropy.

V. SUMMARY AND OUTLOOK

Using a topological quantum field theory approach, the entanglement spectrum of topological phases hosting non-Abelian anyons is found to have a universal structure: each eigenvalue can be labeled by the sub-system's anyonic charge, and its weight is dictated by the corresponding quantum dimension; likewise, its degeneracy is dictated by the fusion rules of the low energy topological theory. We tested our results for solvable anyon-chain models, specifically for the $SU(2)_k$ AKLT chains. Such gapped models elucidate the fact that as the subsystem size increases beyond the correlation length, the eigenvalues of the entanglement spectrum reach the predicted universal ratio dictated by quantum dimensions. Our result generalizes the well known degeneracies in the entanglement spectrum, *e.g.*, in the Haldane chain, which serve as fingerprints of topological order, into universal ratios dictated by the quantum dimension of the possible fusion channels.

We have established a connection between our results on the entanglement spectrum, with a quantity that can be experimentally tested more easily. That is, measurements of the anyonic-charge-resolved Rényi entropy provide information on the multiplet structure in the entanglement spectrum. We demonstrated this explicitly for Ising anyons realized in the Kitaev chain. Unfortunately this illustrates our results only in a partial way, since the quantum dimensions of the total topological charge correspond to Abelian anyons, hence the entanglement spectrum is characterized by degeneracies, and not by non-trivial ratios. Yet we have also shown that these degeneracies can be measured via the parity-resolved Rényi entropy in Majorana wires. In principle it should be possible to generalize the symmetry resolved Rényi entropy measurement protocol to arbitrary anyons by projecting anyon pairs onto specific fusion channels.

It would be interesting to connect our results with the ES of non-Abelian 2D FQH phases⁴, which displays the structure of conformal towers describing 1D edge states (see Sec. I for further discussion and references). The derivation of our results assumes a simple wave function of few anyons fused from the vacuum (see Fig. 1); this is a natural description of 1D states, e.g., in the Kitaev chain or the various anyon-chain generalizations that we considered. However, the application of topological quantum field theory is not restricted to 1D. It also describes the non-Abelian sectors in 2D FQH phases, where the ES is quasi-continuous due to the gapless 1D edge states. Nevertheless, one can still label each state of the entanglement Hamiltonian by the total fusion channel of the subsystem. We expect the prediction that the ES contains universal multiplets dictated by the scaling dimensions and fusion rules to apply even in the presence of gapless edge states and leave this for future work.

Using our results, entanglement spectroscopy can be used to identify non-Abelian anyons as emergent particles in condensed matter systems of interacting bosons or fermions, both using computational methods, and with sufficient motivation, using experimental multi-copy entanglement-measurement methods.

ACKNOWLEDGEMENTS

We thank Eddy Ardonne, Reinhold Egger, Masaki Oshikawa, Stephan Plugge, Simon Trebst, and Ari Turner for discussions and remarks. E.C. acknowledges Chetan Nayak and the Boulder school for providing pedagogical background. K.S. and E.S. were supported by the US-Israel Binational Science Foundation (Grant No. 2016255).

Appendix A: Notations and normalizations of anyonic fusion algebras

In this appendix we specify the notations and conventions of the anyonic fusion algebra in the paper.

The splitting space of two anyons, a and b, with total charge, c, is a vector space V_c^{ab} of dimension N_{ab}^c ; it is indexed by $\mu = 1, \ldots, N_{ab}^c$. We label the states in this space either using bra-ket notation $|a, b; c, \mu\rangle \in V_c^{ab}$ or using the diagramatic presentation

$$|a,b;c,\mu\rangle = \left(\frac{d_c}{d_a d_b}\right)^{\frac{1}{4}} \stackrel{a}{\sim} \underbrace{c}_{c} \stackrel{b}{\mu}, \qquad (A1)$$

where the dual fusion space, V_{ab}^c , is spanned by

$$\langle a, b; c, \mu | = \left(\frac{d_c}{d_a d_b}\right)^{\frac{1}{4}} a^{c} \mu b.$$
 (A2)

We omit the μ -index for $N_{ab}^c = 1$ where it is redundant. The benefits of the above diagram normalization is the orthonormality of the states:

$$\langle a', b'; c', \mu' | a, b; c, \mu \rangle = \delta_{a,a'} \delta_{b,b'} \delta_{c,c'} \delta_{\mu,\mu'} \mathbb{I}_c, \qquad (A3)$$

$$\operatorname{Tr}\left\{|a,b;c,\mu\rangle\langle a',b';c',\mu'|\right\} = \delta_{a,a'}\delta_{b,b'}\delta_{c,c'}\delta_{\mu,\mu'}.$$
 (A4)

The partial traces are similarly nicely normalized

$$\operatorname{Tr}_{B}\left\{|a,b;c,\mu\rangle\langle a',b';c',\mu'|\right\} = \delta_{a,a'}\delta_{b,b'}\delta_{c,c'}\delta_{\mu,\mu'}\mathbb{I}_{a}.$$
(A5)

For example, we may use these normalized states to resolve the identity operator

$$a \downarrow b = \mathbb{I}_{ab} = \sum_{c,\mu} |a,b;c,\mu\rangle \langle a,b;c,\mu|$$
$$= \sum_{c,\mu} \sqrt{\frac{d_c}{d_a d_b}} a \downarrow b \downarrow b.$$
(A6)

Note, that the relation between this "bra-ket" normalization and Kitaev's ψ -notation³⁸ is $|a, b; c, \mu\rangle \Leftrightarrow (\frac{d_c}{d_a d_b})^{1/4} \psi_{c,\mu}^{ab}$. With this identification all diagrammatic representations are identical. A complete description of anyonic fusion algebras in our normalized notation is found in Appendix A of Bonderson, Knapp, and Patel³⁹; see also Ref. [65].

Appendix B: TQFT results for general edge content

The ES of anyonic models is complicated by two facts:

1. Multiple boundary anyons

First, one may have cases with multiple boundary anyons. In this case, the effective wave function is described in terms of anyons $\{a_i\}_{i=1}^n, \{\bar{a}_i\}_{i=1}^n$ fused from the vacuum and is given by

$$\rho = \prod_{i} |a_i, \bar{a}_i; I\rangle^L |a_i, \bar{a}_i; I\rangle^R \langle a_i, \bar{a}_i; I|^L \langle a_i, \bar{a}_i; I|^R.$$
(B1)

In this case, after tracing out region A, one obtains (note that taking the partial trace is not equivalent to the partial anyonic trace^{38,39}

$$\rho_A = \operatorname{Tr}_B \rho = \sum_c \frac{d_c}{1} [\widetilde{\operatorname{Tr}}_B[\rho]_1]_c = \sum_c \frac{d_c}{\prod_i d_{a_i}^2} [\mathbb{I}_{\{a_i^L, \bar{a}_i^R\}}]_c,$$
(B2)

where in general the identity operator of the collection of anyons $\{a_i^L\} \cup \{\bar{a}_i^R\}$ to fuse into total channel c contains $N_{\{a_i^L\} \cup \{\bar{a}_i^R\}}^c \geq 1$ many possibilities,

$$[\mathbb{I}_{\{a_i^L,\bar{a}_i^R\}}]_c = \sum_{\mu=1}^{N_{\{a_i^L,\bar{a}_i^R\}}^c} |\{a_i^L,\bar{a}_i^R\};c,\mu\rangle\langle\{a_i^L,\bar{a}_i^R\};c,\mu|, (B3)$$

$$\mathbb{I}_{\{a_i^L, \bar{a}_i^R\}} = \sum_c [\mathbb{I}_{\{a_i^L, \bar{a}_i^R\}}]_c,$$
(B4)

$$N^{c}_{\{a^{L}_{i},\bar{a}^{R}_{i}\}} = \sum_{d^{L},d^{R}} N^{d^{L}}_{\{a^{L}_{i}\}} N^{d^{R}}_{\{\bar{a}^{R}_{i}\}} N^{c}_{d^{L}d^{R}}, \qquad (B5)$$

$$N_{\{a_i\}}^d = \sum_{\{x_i\}_{i=2}^{n-1}} N_{a_1 a_2}^{x_2} N_{x_2 a_3}^{x_3} \cdots N_{x_{n-2} a_{n-1}}^{x_{n-1}} N_{x_{n-1} a_n}^d.$$
(B6)

This immediately yields the spectrum and degeneracies generalizing Eq. 6,

$$\lambda_c = \frac{d_c}{\prod_i d_{a_i}^2}, \qquad \#_c = N^c_{\{a_i^L, \bar{a}_i^R\}}. \tag{B7}$$

2. Superposition of anyons

Second, the wave function may take the form of a linear superposition over the boundary anyons,

$$|\Psi\rangle = \sum_{a_L,\bar{a}_R} \alpha_{a_L,\bar{a}_R} |a,\bar{a};I\rangle^L |a,\bar{a};I\rangle^R.$$
(B8)

Consequently, the density matrix of the full system is

$$\rho = \sum_{a_L,\bar{a}_R} \alpha_{a_L,\bar{a}_R} \sum_{a_L',\bar{a}'_R} \alpha^*_{a'_L,\bar{a}'_R} \\
\times |a,\bar{a};I\rangle^L |a,\bar{a};I\rangle^R \langle a',\bar{a}';I|^L \langle a',\bar{a}';I|^R.$$
(B9)

Tracing out subsystem B, as in Fig. 2(c), brings a deltafunction imposing $a_L = a'_L$, $a_R = a'_R$, and similarly for \bar{a}_L and \bar{a}_R . Thus,

$$\rho_A = \sum_{a_L, \bar{a}_R} |\alpha_{a_L, \bar{a}_R}|^2 \rho_A(a_L, \bar{a}_R),$$
(B10)

where $\rho_A(a_L, \bar{a}_R)$ denotes the reduced density matrix for fixed boundary anyons. In our main result, Eq. (6), various fusion trees associated with the possible internal fusion processes of the fixed boundary anyons lead to $\#_c$ distinct Schmidt states; see Eq. (B7). Thus, distinct values of the boundary anyons allow for a further distinction of the Schmidt states. We conclude that each block of fixed c admits a further block decomposition according to distinct values of a_L, \bar{a}_R , which together fuse into c,

$$\rho_A = \bigoplus_{a_L, \bar{a}_R} |\alpha_{a_L, \bar{a}_R}|^2 \bigoplus_{c \text{ s.t. } N^c_{a_L, \bar{a}_R} \neq 0} \rho_A^{(c)}(a_L, \bar{a}_R). \quad (B11)$$

We conclude that there are different multiplets labeled by two boundary anyons a_L, \bar{a}_R ; Each multiplet decomposes according to the total anyon charge of region A, c, with a multiplicity given by $N_{\{a_i^L\}\cup\{\bar{a}_i^R\}}^c$ which indeed depends on the edge content. This allows to understand the results in Sec. III A 4.

Appendix C: Exact ES at the AKLT sweet spot

Here, we derive Eq. (14). The open BC anyon-chain wave-function takes the form

$$\psi = \sum_{\{x_i\}} f(\{x_i\}) | x_1, \cdots, x_L \rangle, \qquad (C1)$$

where⁴⁵ $f({x_i}) = (-1)^{\#1}(d_1)^{(\#1-L)/2}(d_2)^{L/2-\#1}$, *L* is the number of *inner* anyonic legs, #j is the number of appearance of *j*-type anyons, and we impose $x_i \in \{0, 1\}$ and $x_1 = x_L = 0$. This wave function is not normalized; this is later taken into account. The form of this wave function dictates that $\lambda_2 = 0$ for all *L*, since none of the anyons can be a 2-type anyon.

Let us define $\chi_{ij}(n) = \sum_{\{x_i\}\neq x_1,x_n} f^2(\{x_i\})_{x_1=i,x_n=j}$ to be the probability to find a string with n inner legs in which the first and nth anyons are $x_1 = i$, $x_n = j$. We also define $\chi_i(1) = f^2(i)$ to be squared coefficient of a single leg anyon string of length 1. The system under consideration is a bi-partitioned string with n_1, n_2 outer anyons in each sector, see Fig. (3). Since the singular values are norm preserving, we can represent the two unnormalized values of the entanglement spectrum as a product of two strings

$$(\bar{\Lambda}_0)^2 = \frac{\chi_{00}(n_1+1)\chi_{00}(n_2+1)}{\chi_0(1)},$$

$$(\bar{\Lambda}_1)^2 = \frac{\chi_{01}(n_1+1)\chi_{10}(n_2+1)}{\chi_1(1)},$$
(C2)

where the factors $\chi_0(1)$ ($\chi_1(1)$) in the denominator are

due to double counting of a single 0 (1) anyon. The bar sign indicates that these entanglement spectrum values are not normalized. However, we can take care of normalization by observing that the string $\chi_{00}(n_1 + n_2 + 1)$ includes both $\bar{\Lambda}_0$ and $\bar{\Lambda}_1$. Therefore, the normalized values are

$$(\Lambda_0)^2 = \frac{\chi_{00}(n_1+1)\chi_{00}(n_2+1)}{\chi_0(1)\chi_{00}(n_1+n_2+1)},$$

$$(\Lambda_1)^2 = \frac{\chi_{01}(n_1+1)\chi_{10}(n_2+1)}{\chi_1(1)\chi_{00}(n_1+n_2+1)}.$$
(C3)

Using the string identities $\chi_{01}(n) = \chi_0(1)\chi_{11}(n-1)$ and $\chi_{00}(n) = \chi_{00}(2)\chi_{11}(n-2)$ we rewrite the above equations as

$$(\Lambda_0)^2 = \frac{\chi_0(1)\chi_{11}(n_1-1)\chi_{11}(n_2-1)}{\chi_{11}(n_1+n_2-1)}, \qquad (C4)$$
$$(\Lambda_1)^2 = \frac{\chi_{11}(n_1)\chi_{11}(n_2)}{\chi_1(1)\chi_{11}(n_1+n_2-1)}.$$

Consequently, in order to obtain a closed form of the entanglement spectrum for all n we need to explicitly calculate only the value of $\chi_{11}(n)$. To do so, we observe that each $\chi_{11}(n)$, which is nothing but an open string starting and ending with 1 anyons, can be recursively expressed as

$$\chi_{11}(n) = \chi_1(1)\chi_{11}(n-1) + \chi_{10}(2)\chi_{11}(n-2),$$

= $\frac{1}{d_1}\chi_{11}(n-1) + \frac{1}{d_2}\chi_{11}(n-2).$ (C5)

This equation has a simple closed form solution:

$$\chi_{11}(n) = A + \frac{B}{(-d_1)^n},$$
 (C6)

where A and B are some constants. Using $\chi_{11}(2) = \frac{1}{d_2^2}$ and the value of $\chi_1(1) = \frac{1}{d_2}$, we find

$$\chi_{11}(n) = \frac{d_1}{d_2^3} \left[1 - \frac{1}{\left(-d_1\right)^n} \right].$$
 (C7)

Plugging this at Eq. (C4) we obtain the general form of the entanglement spectrum

$$\Lambda_{0} = \frac{1}{d_{2}} \sqrt{\frac{\left[1 - (-d_{1})^{-n_{1}+1}\right] \left[1 - (-d_{1})^{-n_{2}+1}\right]}{\left[1 - (-d_{1})^{-n_{1}-n_{2}+1}\right]}},$$
$$\Lambda_{1} = \frac{\sqrt{d_{1}}}{d_{2}} \sqrt{\frac{\left[1 - (-d_{1})^{-n_{1}}\right] \left[1 - (-d_{1})^{-n_{2}}\right]}{\left[1 - (-d_{1})^{-n_{1}-n_{2}+1}\right]}}.$$
 (C8)

For equally partitioned system $n_1 = n_2 = n$ one obtains

$$\Lambda_0 = \frac{1}{d_2} \cdot \frac{1 - (-d_1)^{-n+1}}{\sqrt{1 + d_1^{-2n+1}}},$$
(C9)

$$\Lambda_1 = \frac{\sqrt{d_1}}{d_2} \cdot \frac{1 - (-d_1)^{-n}}{\sqrt{1 + d_1^{-2n+1}}}.$$

This completes the derivation of the ES $\{\lambda\} = \{\Lambda^2\}$ in Eq. (14).

¹ A. Kitaev and J. Preskill, Phys. Rev. Lett. **96**, 110404 (2006).

- ² M. Levin and X.-G. Wen, Phys. Rev. Lett. 96, 110405 (2006).
- ³ H.-C. Jiang, Z. Wang, and L. Balents, Nature Physics 8, 902 (2012).
- ⁴ H. Li and F. D. M. Haldane, Phys. Rev. Lett. **101**, 010504 (2008).
- ⁵ A. Chandran, M. Hermanns, N. Regnault, and B. A. Bernevig, Physical Review B 84, 205136 (2011).
- ⁶ X.-L. Qi, H. Katsura, and A. W. Ludwig, Physical review letters **108**, 196402 (2012).
- ⁷ D. Poilblanc, N. Schuch, D. Pérez-García, and J. I. Cirac, Physical Review B 86, 014404 (2012).
- ⁸ J. Dubail, N. Read, and E. Rezayi, Physical Review B 85, 115321 (2012).
- ⁹ A. Sterdyniak, A. Chandran, N. Regnault, B. A. Bernevig, and P. Bonderson, Physical Review B 85, 125308 (2012).
- ¹⁰ B. Estienne, Z. Papić, N. Regnault, and B. A. Bernevig, Physical Review B 87, 161112 (2013).
- ¹¹ B. Yan, R. R. Biswas, and C. H. Greene, Physical Review B **99**, 035153 (2019).
- ¹² N. Regnault and B. A. Bernevig, Physical Review X 1, 021014 (2011).
- ¹³ D. Poilblanc, J. I. Cirac, and N. Schuch, Physical Review B **91**, 224431 (2015).
- ¹⁴ N. Laflorencie, Physics Reports **646**, 1 (2016).
- ¹⁵ L. Fidkowski and A. Kitaev, Physical Review B **81**, 134509 (2010).
- ¹⁶ F. Pollmann, A. M. Turner, E. Berg, and M. Oshikawa, Physical review b 81, 064439 (2010).
- ¹⁷ A. M. Turner, F. Pollmann, and E. Berg, Physical review b 83, 075102 (2011).
- ¹⁸ L. Fidkowski and A. Kitaev, Physical review b 83, 075103 (2011).
- ¹⁹ X. Chen, Z.-C. Gu, and X.-G. Wen, Physical review b 83, 035107 (2011).
- ²⁰ X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Physical Review B 87, 155114 (2013).
- ²¹ N. Schuch, D. Pérez-García, and I. Cirac, Phys. Rev. B 84, 165139 (2011).
- 22 L. Fidkowski, Physical review letters $\mathbf{104},\,130502$ (2010).
- ²³ D. Poilblanc, Physical review letters **105**, 077202 (2010).
- ²⁴ C.-Y. Huang and F.-L. Lin, Physical Review B 84, 125110 (2011).
- ²⁵ X. Deng and L. Santos, Physical Review B 84, 085138 (2011).
- ²⁶ W. Li, A. Weichselbaum, and J. von Delft, Physical Re-

view B 88, 245121 (2013).

- ²⁷ K. Duivenvoorden and T. Quella, Physical Review B 87, 125145 (2013).
- ²⁸ H. Nonne, M. Moliner, S. Capponi, P. Lecheminant, and K. Totsuka, EPL (Europhysics Letters) **102**, 37008 (2013).
- ²⁹ T. Morimoto, H. Ueda, T. Momoi, and A. Furusaki, Physical Review B **90**, 235111 (2014).
- ³⁰ A. Y. Kitaev, Physics-Uspekhi **44**, 131 (2001).
- ³¹ J. Alicea and P. Fendley, Annual Review of Condensed Matter Physics 7, 119 (2016).
- ³² N. Laflorencie and S. Rachel, J. Stat. Mech., P11013 (2014).
- ³³ M. Goldstein and E. Sela, Physical review letters 120, 200602 (2018).
- ³⁴ J. C. Xavier, F. C. Alcaraz, and G. Sierra, Phys. Rev. B 98, 041106 (2018).
- ³⁵ S. Matsuura, X. Wen, L.-Y. Hung, and S. Ryu, Physical Review B **93**, 195113 (2016).
- ³⁶ A. Y. Kitaev, Annals of Physics **303**, 2 (2003).
- ³⁷ C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).
- ³⁸ A. Kitaev, Annals of Physics **321**, 2 (2006), january Special Issue.
- ³⁹ P. Bonderson, C. Knapp, and K. Patel, Annals of Physics 385, 399 (2017).
- ⁴⁰ J. K. Pachos, *Introduction to topological quantum compu*tation (Cambridge University Press, 2012).
- ⁴¹ S. Singh, R. N. Pfeifer, G. Vidal, and G. K. Brennen, Physical Review B 89, 075112 (2014).
- ⁴² A. Feiguin, S. Trebst, A. W. Ludwig, M. Troyer, A. Kitaev, Z. Wang, and M. H. Freedman, Physical review letters **98**, 160409 (2007).
- ⁴³ S. Trebst, E. Ardonne, A. Feiguin, D. A. Huse, A. W. Ludwig, and M. Troyer, Physical review letters **101**, 050401 (2008).
- ⁴⁴ S. Trebst, M. Troyer, Z. Wang, and A. W. Ludwig, Progress of Theoretical Physics Supplement **176**, 384 (2008).
- ⁴⁵ C. Gils, E. Ardonne, S. Trebst, D. A. Huse, A. W. Ludwig, M. Troyer, and Z. Wang, Physical Review B 87, 235120 (2013).
- ⁴⁶ K. Choo, C. W. von Keyserlingk, N. Regnault, and T. Neupert, Phys. Rev. Lett. **121**, 086808 (2018).
- ⁴⁷ R. Barends, J. Kelly, A. Megrant, A. Veitia, D. Sank, E. Jeffrey, T. C. White, J. Mutus, A. G. Fowler, B. Campbell, *et al.*, Nature **508**, 500 (2014).
- ⁴⁸ B. Van Heck, A. Akhmerov, F. Hassler, M. Burrello, and C. Beenakker, New Journal of Physics **14**, 035019 (2012).

- ⁴⁹ F. Hassler and D. Schuricht, New Journal of Physics 14, 125018 (2012).
- ⁵⁰ D. Aasen, M. Hell, R. V. Mishmash, A. Higginbotham, J. Danon, M. Leijnse, T. S. Jespersen, J. A. Folk, C. M. Marcus, K. Flensberg, and J. Alicea, Phys. Rev. X 6, 031016 (2016).
- ⁵¹ L. Landau, S. Plugge, E. Sela, A. Altland, S. Albrecht, and R. Egger, Physical review letters **116**, 050501 (2016).
- ⁵² D. Litinski and F. von Oppen, Physical Review B 97, 205404 (2018).
- ⁵³ E. Cornfeld, M. Goldstein, and E. Sela, Phys. Rev. A 98, 032302 (2018).
- ⁵⁴ E. Cornfeld, E. Sela, and M. Goldstein, arXiv preprint arXiv:1808.04471 (2018).
- ⁵⁵ I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Physical review letters **59**, 799 (1987).
- ⁵⁶ D. A. Abanin and E. Demler, Physical review letters **109**, 020504 (2012).
- ⁵⁷ A. Daley, H. Pichler, J. Schachenmayer, and P. Zoller,

Phys. Rev. Lett. 109, 020505 (2012).

- ⁵⁸ R. Islam, R. Ma, P. M. Preiss, M. Eric Tai, A. Lukin, M. Rispoli, and M. Greiner, Nature **528**, 15750 (2015).
- ⁵⁹ A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Science **353**, 794 (2016).
- ⁶⁰ G. De Chiara, L. Lepori, M. Lewenstein, and A. Sanpera, Physical review letters **109**, 237208 (2012).
- 61 J. Alicea, Reports on progress in physics $\mathbf{75},076501$ (2012).
- ⁶² S. M. Albrecht, A. Higginbotham, M. Madsen, F. Kuemmeth, T. S. Jespersen, J. Nygård, P. Krogstrup, and C. Marcus, Nature **531**, 206 (2016).
- ⁶³ H. Pichler, L. Bonnes, A. J. Daley, A. M. Läuchli, and P. Zoller, New Journal of Physics **15**, 063003 (2013).
- ⁶⁴ S. Plugge, L. Landau, E. Sela, A. Altland, K. Flensberg, and R. Egger, Physical Review B **94**, 174514 (2016).
- ⁶⁵ P. H. Bonderson, Non-Abelian anyons and interferometry, Ph.D. thesis, California Institute of Technology (2007).