On Sets of Words of Rank Two

Giuseppa Castiglione, Gabriele Fici, and Antonio Restivo

Dipartimento di Matematica e Informatica, Università di Palermo Via Archirafi 34, Palermo, Italy

{giuseppa.castiglione,gabriele.fici,antonio.restivo}@unipa.it

Abstract. Given a (finite or infinite) subset X of the free monoid A^* over a finite alphabet A, the rank of X is the minimal cardinality of a set F such that $X \subseteq F^*$. A submonoid M generated by k elements of A^* is k-maximal if there does not exist another submonoid generated by at most k words containing M. We call a set $X \subseteq A^*$ primitive if it is the basis of a |X|-maximal submonoid. This extends the notion of primitive word: indeed, $\{w\}$ is a primitive set if and only if w is a primitive word. By definition, for any set X, there exists a primitive set Y such that $X \subseteq Y^*$. The set Y is therefore called a primitive root of X. As a main result, we prove that if a set has rank 2, then it has a unique primitive root. This result cannot be extended to sets of rank larger than 2. For a single word w, we say that the set $\{x, y\}$ is a binary root of w if w can be written as a concatenation of copies of x and y and $\{x, y\}$ is a primitive set. We prove that every primitive word w has at most one binary root $\{x,y\}$ such that $|x| + |y| < \sqrt{|w|}$. That is, the binary root of a word is unique provided the length of the word is sufficiently large with respect to the size of the root. Our results are also compared to previous approaches that investigate pseudo-repetitions, where a morphic involutive function θ is defined on A^* . In this setting, the notions of θ -power, θ -primitive and θ -root are defined, and it is shown that any word has a unique θ -primitive root. This result can be obtained with our approach by showing that a word w is θ -primitive if and only if $\{w, \theta(w)\}$ is a primitive set.

Keywords: Repetition, pseudo-repetition, hidden repetition, primitive set, binary root, *k*-maximal monoid.

1 Introduction

The notion of rank plays an important role in combinatorics on words. Given a subset X of the free monoid A^* over a finite alphabet A, the rank of X, in symbols r(X), is defined as the smallest number of words needed to express all words of X, i.e., as the minimal cardinality of a set F such that $X \subseteq F^*$. Notice that this minimal set F may not be unique. For instance, the set $X = \{aabca, aa, bcaaa\}$ has rank 2 and there exist two distinct sets $F_1 = \{aa, bca\}$ and $F_2 = \{a, bc\}$ such that $X \subseteq F_1^*$ and $X \subseteq F_2^*$. It is worth noticing that since $r(X) \leq min\{|X|, |A|\}$, r(X) is always finite even if X is an infinite set.

A set X is said to be *elementary* if r(X) = |X|. The notion of rank – and the related notion of elementary set – have been investigated in several papers

 $\mathbf{2}$

(cf. [16,18,17]). In particular, in [16] it is shown that the problem to decide whether a finite set is elementary is co-NP-complete.

In this paper, we introduce the notion of primitiveness for a *set* of words, which is closely related to that of rank. We first define the notion of *k*-maximal submomoid. A submonoid M of A^* , generated by k elements, is *k*-maximal if there does not exist another submonoid generated by at most k words containing M. We then call a set $X \subseteq A^*$ primitive if it is the basis of a |X|-maximal submonoid. Notice that if X is primitive, then r(X) = |X|, i.e., X is elementary. The converse is not in general true: there exist elementary sets that are not primitive. For instance, the set $F_1 = \{aa, bca\}$ is elementary, but it is not primitive since $F_1^* \subseteq F_2^* = \{a, bc\}^*$. The set F_2 , instead, is primitive.

The notion of primitive set can be seen as an extension of the classical notion of primitive word. Indeed, given a word $w \in A^*$, the set $\{w\}$ is primitive if and only if the word w is primitive. For instance, the set $\{abab, abababab\}$ is not elementary, the set $\{abab\}$ is elementary but not primitive, and the set $\{ab\}$ is primitive.

We have from that definition that for every set X, there exists a primitive set Y such that $X \subseteq Y^*$. The set Y is therefore called a *primitive root* of X. However, the primitive root of a set is not, in general, unique. Consider for instance the set $X = \{abcbab, abcdcbab, abcdccbab\}$. It has rank 3, hence it is elementary, yet it is not primitive. Indeed, $X \subseteq \{ab, cb, cd\}^*$. The set $\{ab, cb, cd\}$ is primitive, and it is a primitive root of X. However, it is not the only primitive root of X: the set $\{abc, dc, bab\}$ is primitive and $X \subseteq \{abc, dc, bab\}^*$, hence $\{abc, dc, bab\}$ is another primitive root of X. In the special case of sets of rank 1, clearly these always have a unique primitive root. For instance, the primitive root of the set $\{abab, abababab\}$ is the set $\{ab\}$.

As a main result, we prove that if a set has rank 2, then it has a unique primitive root. This is equivalent to say that for every pair of nonempty words $\{x, y\}$ such that $xy \neq yx$ there exists a unique primitive set $\{u, v\}$ such that x and y can be written as concatenations of copies of u and v. The proof is based on the algebraic properties of k-maximal submonoids of a free monoid.

In this investigation, we also take into account another notion of rank, that of free rank (in the literature, in order to avoid ambiguity, the notion of rank we gave above is often referred to as the combinatorial rank). The free rank of a set X is the cardinality of the basis of the minimal free submonoid containing X. Closely related to the notion of free rank is the defect theorem, which states that if X is not a code (i.e., X^* is not a free submonoid), then the free rank of X is strictly smaller than its cardinality. We are specially interested in the case k = 2 (that is, the case of 2-maximal submonoids) and we use the fact that, in this special case, the notions of free rank and (combinatorial) rank coincide. A fundamental step in our argument is Theorem 14, which states that the intersection of two 2-maximal submonoids is either the empty word or a submonoid generated by one primitive word. As a consequence, for every submonoid M generated by two words that do not commute, there exists a unique 2-maximal submonoid containing M. This is equivalent to the fact that every set of rank 2 has a

unique primitive root. One of the examples we gave above shows that this result is no longer true for sets of rank 3 or larger — this highlights the very special role of sets of rank 1 or 2.

From these results we derive some consequences on the combinatorics of a single word. Given a word w, we say that $\{x, y\}$ is a binary root of w if w can be written as a concatenation of copies of x and y and $\{x, y\}$ is a primitive set. We prove that every primitive word w has at most one binary root $\{x, y\}$ such that $|x| + |y| < \sqrt{|w|}$. That is, the binary root of a word is unique provided the length of the word is sufficiently large with respect to the size of the root. The notion of binary root of a single word may be seen as a way to capture a hidden "repetitive structure", which encompasses the classical notion of integer repetition (non-primitive word). Indeed, the existence in a word w of a "short" (with respect to |w|) binary root reveals some hidden repetition in the word.

As described in the last section, our results can also be compared to previous approaches that investigate *pseudo-repetitions*, where an involutive morphism (or antimorphism) θ is defined on the set of words A^* . This idea stems from the seminal paper of Czeizler, Kari and Seki [4], where originally θ was the Watson-Crick complementarity function and the motivation was the discovery of hidden repetitive structures in biological sequences. A word w is called a θ power if there exists a word v such that w can be factored using copies of v and $\theta(v)$ — otherwise the word w is called θ -primitive. If v is a θ -primitive word, then it is called the θ -primitive root of w. Of course, since the same applies to the word $\theta(w)$, these definitions can be given in terms of the pair $\{w, \theta(w)\}$ and considering as the root the pair $\{v, \theta(v)\}$. With our results, we generalize this setting by considering as a root any pair of words $\{x, y\}$, i.e., dropping the relation between the components of the pair.

2 Preliminaries

Given a finite nonempty set A, called the *alphabet*, with A^* (resp. $A^+ = A^* \setminus \{\varepsilon\}$) we denote the *free monoid* (resp. *free semigroup*) generated by A, i.e., the set of all finite words (resp. all finite nonempty words) over A.

The length |w| of a word $w \in A^*$ is the number of its symbols. The length of the empty word ε is 0. For a word w = uvz, with $u, v, z \in A^*$, we say that v is a factor of w. Such a factor is called *internal* if $u, z \neq \varepsilon$, a *prefix* if $u = \varepsilon$, or a suffix if $z = \varepsilon$. A word w is primitive if $w = v^n$ implies n = 1, otherwise it is called a power. Equivalently, w is primitive if and only if it is not an internal factor of w^2 .

It is well known in combinatorics on words (see, e.g., [14]) that given two words x and y we have xy = yx if and only if x and y are powers of the same word.

Given a subset X of A^* , we let X^* denote the submonoid of A^* generated by X (under concatenation). Conversely, given a submonoid M of A^* , there exists a unique set X that generates M and is minimal for set inclusion. In fact, X is

G. Castiglione, G. Fici and A. Restivo

the set

$$X = (M \setminus \{\varepsilon\}) \setminus (M \setminus \{\varepsilon\})^2, \tag{1}$$

i.e., X is the set of nonempty words of M that cannot be written as a product of two nonempty words of M. The set X will be referred to as the *minimal* generating set of M, or the set of generators of M.

Let M be a submonoid of A^* and X its minimal generating set. M is said to be *free* if any word of M can be *uniquely* expressed as a product of elements of X. The minimal generating set of a free submonoid M of A^* is called a *code*; it is referred to as *the basis* of M. It is easy to see that a set X is a code if and only if, for every $x, y \in X, x \neq y$, one has $xX^* \cap yX^* = \emptyset$. We say that X is a *prefix code* (resp. a *suffix code*) if for all $x, y \in X$, one has $x \cap yA^* = \emptyset$ (resp. $x \cap A^*y = \emptyset$). A code is a *bifix code* if it is both a prefix and a suffix code. It follows from elementary automata theory that if X is a prefix code, then there exists a DFA \mathcal{A}_X recognizing X^* whose set of states Q_X verifies (cf. [2]):

$$|Q_X| \le \sum_{x \in X} |x| - |X| + 1.$$

A submonoid M of A^* is called *pure* (cf. [19]) if for all $w \in A^*$ and $n \ge 1$,

$$w^n \in M \Rightarrow w \in M.$$

By a result of Tilson [20], any nonempty intersection of free submonoids of A^* is free. As a consequence, for any subset $X \subseteq A^*$, there exists the smallest free submonoid containing X.

Here we mention the well-known Defect Theorem (cf. [1], [14, Chap. 1], [15, Chap. 6]), a fundamental result in the theory of codes that provides a relation between a given subset X of A^* and the basis of the minimal free submonoid containing X (called the *free hull* of X).

Theorem 1 (Defect Theorem). Let X be a finite nonempty subset of A^* . Let Y be the basis of the free hull of X. Then either X is a code, and Y = X, or

$$|Y| \le |X| - 1.$$

As in [9], given a set $X \subseteq A^*$, we let $r_f(X)$ denote the cardinality of the basis of the free hull of X, called the *free rank* of X. Notice that for any subset $X \subseteq A^*$, X and X^* have the same free rank. Furthermore, by r(X) we denote the *combinatorial rank* (or simply rank) of X, defined by:

$$r(X) = \min\{|Y| \mid Y \subseteq A^*, X \subseteq Y^*\}.$$

With this notation, the Defect Theorem can be stated as follows.

Theorem 2. Let X be a finite nonempty subset of A^* . Then $r_f(X) \leq |X|$, and the equality holds if and only if X is a code.

4

5

Note that, for any $X \subseteq A^+$, one has

$$r(X) \le r_f(X) \le |X|.$$

Example 3. Let $X = \{aa, ba, baa\}$. One can prove that X is a code, hence we have $r_f(X) = 3$, while r(X) = 2 since $X \subset \{a, b\}^*$. For $X = \{aa, aaa\}$, we have $r(X) = r_f(X) = 1$.

Remark 4. If |X| = 2 then $r_f(X) = r(X)$. So for sets of cardinality 2 we will not specify if we refer to the free rank or to the (combinatorial) rank.

Moreover, from the complexity point of view, Néraud proved that deciding if a set has rank 2 can be done in polynomial time [17], whereas for general rank k it is an NP-hard problem [16].

The dependency graph (cf. [9]) of a finite set $X \subset A^+$ is the graph $G_X = (X, E_X)$ where $E_X = \{(u, v) \in X \times X \mid uX^* \cap vX^* \neq \emptyset\}$. Notice that if X is a code, then G_X has no edge. Furthermore, if (u, v) is an edge, then u is a prefix of v or vice versa. In [8] and [9], the following useful lemma is proved.

Lemma 5 (Graph Lemma). Let $X \subseteq A^+$ be a finite set that is not a code. Then

$$r_f(X) \le c(X) < |X|$$

where c(X) is the number of connected components of G_X .

Example 6. Let $X = \{a, ab, abc, bca, acb, cba\}$. We have $acba = a \cdot cba = acb \cdot a$ and $abca = a \cdot bca = abc \cdot a$. The basis of the free hull of X is $Y = \{a, ab, bc, cb\}$, hence $r_f(X) = 4$. Furthermore, r(X) = 3 and c(X) = 4, as shown in Figure 1.




Fig. 1. The dependency graph of $X = \{a, ab, abc, bca, acb, cba\}$.

3 k-Maximal Monoids

With \mathcal{M}_k we denote the family of submonoids of A^* having at most k generators in A^+ . The following definition is fundamental for the theory developed in this paper.

6 G. Castiglione, G. Fici and A. Restivo

Definition 7. A submonoid $M \in \mathcal{M}_k$ is k-maximal if for every $M' \in \mathcal{M}_k$, $M \subseteq M'$ implies M = M'.

In other words, M is k-maximal if it is not possible to find another submonoid generated by at most k words containing M.

Example 8. Let $A = \{a, b, c\}$. The submonoid $M = \{a, abca\}^*$ is not 2-maximal since *abca* can be factored with *a* and *bc*, hence *M* is contained in $\{a, bc\}^*$. On the contrary, $\{a, bc\}^*$ is 2-maximal since, obviously, *a* and *bc* cannot be factored using two common factors.

Example 9. Let $A = \{a, b, c, d\}$. The submonoid $\{a, cbd, dbd\}^*$ is 3-maximal, whereas $\{a, cbd, dcbd\}^*$ is not 3-maximal since it is contained in $\{a, d, cb\}^*$.

Proposition 10. Let M be a k-maximal submonoid and X its minimal generating set. Then, X is a bifix code.

Proof. By contradiction, if X is not prefix (resp. not suffix) then there exist $u, v \in X$ and $t \in A^+$ such that v = ut (resp v = tu). It follows that $X^* \subseteq (X \setminus \{v\} \cup \{t\})^*$, whence $X^* = M$ is not k-maximal. \Box

Remark 11. By Proposition 10, it follows that if X^* is k-maximal, then $r(X) = r_f(X) = k$. The inverse implication does not hold in general. For example, the submonoid $X^* = \{a, cbd, dcbd\}^*$ of Example 9 has both rank and free rank equal to 3 and is bifix, but it is not 3-maximal.

Proposition 12. Let M be a k-maximal submonoid. Then M is a pure submonoid.

Proof. We have to show that, for every $z \in A^*$, if $z^n \in M$, for some $n \ge 1$, then $z \in M$. Let X be the minimal generating set of M. If $z^n \in M$, for some n > 1, then $z \in X$ or the set $X \cup \{z\}$ is not a code. By the Defect Theorem (Theorem 1), there exist $u_1, u_2, \ldots, u_k \in A^+$ such that $(X \cup \{z\})^* \subseteq \{u_1, u_2, \ldots, u_k\}^*$. Since $X^* \subseteq \{u_1, u_2, \ldots, u_k\}^*$ and X^* is k-maximal, we have that $X = \{u_1, u_2, \ldots, u_k\}$. Therefore, $X \cup \{z\} \subseteq X^*$, hence $z \in X^*$.

As a direct consequence of Proposition 12, we have that a k-maximal submonoid is generated by primitive words. However, not any set of k primitive words generates a k-maximal monoid (e.g., $X = \{ab, ba\}^*$ is not 2-maximal since it is contained in $\{a, b\}^*$).

Submonoids generated by two words, i.e., the elements of \mathcal{M}_2 , are of special interest for our purposes. They have been extensively studied in literature (cf. [13,10,17,12]) and play an important role in some fundamental aspects of combinatorics on words.

The reader may observe that, as a consequence of some well-known results in combinatorics on words, the submonoids in \mathcal{M}_1 have the following important property: If x^* and u^* are 1-maximal submonoids (i.e., x and u are primitive words) then $x^* \cap u^* = \{\varepsilon\}$. Next Theorem 14, which represents the main result of this section, can be seen as a generalization of this result to the case of 2-maximal submonoids.

It is known (see [10]) that if X and U both have rank 2, then the intersection $X^* \cap U^*$ is a free monoid generated either by at most two words or by an infinite set of words.

Example 13. Let $X_1 = \{abca, bc\}$ and $U_1 = \{a, bcabc\}$. One can verify that $X_1^* \cap U_1^* = \{abcabc, bcabca\}^*$. Let $X_2 = \{aab, aba\}$ and $U_2 = \{a, baaba\}$. Then $X_2^* \cap U_2^* = (a(abaaba)^*baaba)^*$.

In the previous example, we have two submonoids that are not 2-maximal. Indeed, $X_1^*, U_1^* \subseteq \{a, bc\}^*$ and $X_2^*, U_2^* \subseteq \{a, b\}^*$. We now address the question of finding the generators of the intersection of two 2-maximal submonoids.

Theorem 14. Let $X^* = \{x, y\}^*$ and $U^* = \{u, v\}^*$ be two 2-maximal submonoids. If $X^* \cap U^* \neq \{\varepsilon\}$, then there exists a word $z \in A^+$ such that $X^* \cap U^* = z^*$. Moreover, z is primitive, that is, $X^* \cap U^*$ is 1-maximal.


Proof. If $X \cap U = \{z\}$ then $X^* \cap U^* = z^*$. Indeed, if y = v = z and $X^* \cap U^* \neq z^*$ we have the following graph G_Z for $Z = \{x, u, z\}$:



since $\{x, z\}$ and $\{u, z\}$ are bifix sets. Hence, by the Graph Lemma, $r_f(Z) \leq c(Z) = 2$, contradicting the 2-maximality of X and U.

If $X \cap U = \emptyset$, let us consider the set $Z = X \cup U$. We have that $r_f(Z) > 2$ since X^* and U^* are 2-maximal, and, by the Defect Theorem (Theorem 1), $r_f(Z) < 4$ since Z^* is not free (as $X^* \cap U^*$ contains a nonempty word). Hence, the free rank of Z is equal to 3.

Let z be a generator of $X^* \cap U^*$. So, $z = x_1 x_2 \cdots x_m = u_1 u_2 \cdots u_n$, with $m, n \ge 1, x_i \in X$ and $u_j \in U$. Clearly, since z is a generator, for every p < m and q < n one has $x_1 x_2 \cdots x_p \neq u_1 u_2 \cdots u_q$. Moreover, we can suppose, without loss of generality, that $x_1 = x$ and $u_1 = u$. We want to prove that z is the unique generator of $X^* \cap U^*$. By contradiction, suppose that there exists another generator $z' \neq z$ of $X^* \cap U^*$, and let $z' = x'_1 x'_2 \cdots x'_r = u'_1 u'_2 \cdots u'_s$. If $x'_1 \neq x_1 = x$, then $x'_1 = y$ and we have $xZ^* \cap uZ^* \neq \emptyset$ and $yZ^* \cap u'_1Z^* \neq \emptyset$. In both cases $(u'_1 = u \text{ or } u'_1 = v)$, we have that the graph G_Z has two edges, i.e., c(Z) = 2, which is impossible by the Graph Lemma. So $x_1 = x'_1 = x$. In the same way we prove that $u_1 = u'_1 = u$, and therefore in the graph G_Z there is only one edge, namely the one joining x and u.



8 G. Castiglione, G. Fici and A. Restivo

Let $h = \max\{i \mid x_j = x'_j \; \forall j \leq i\}$ and $k = \max\{i \mid u_j = u'_j \; \forall j \leq i\}$. The hypothesis that $z \neq z'$ implies that h < m and k < n. We show that this leads to a contradiction, and then we conclude that z = z' is the unique generator of $X^* \cap U^*$.


Without loss of generality, we can suppose that $x_1x_2\cdots x_h$ is a prefix of $u_1u_2\cdots u_k$. Hence, there exists a nonempty word t such that $x_1x_2\cdots x_ht = u_1u_2\cdots u_k$. By definition of h, $x_{h+1} \neq x'_{h+1}$, and we can suppose that $x_{h+1} = x$ and $x'_{h+1} = y$. Then,

$$tu_{k+1}\cdots u_n = x_{h+1}\cdots x_m = x\cdots x_m$$
$$tu'_{k+1}\cdots u'_s = x'_{h+1}\cdots x'_r = y\cdots x'_r.$$

Set $Z_t = X \cup U \cup \{t\}$. We have

$$tZ_t^* \cap xZ_t^* \neq \emptyset$$
$$tZ_t^* \cap yZ_t^* \neq \emptyset.$$

Thus, the graph G_{Z_t} contains the edges depicted in figure:



By the Graph Lemma, then, the free rank of Z_t is at most 2, and this contradicts the 2-maximality of X^* and U^* .

Finally, let us prove that z is primitive. Since X^* and Y^* are 2-maximal, by Proposition 12 they are both pure, hence also their intersection z^* is pure. But it is immediate that z^* is pure if and only if z is primitive.

Example 15. Consider the two 2-maximal monoids $\{abcab, cb\}^*$ and $\{abc, bcb\}^*$. Their intersection is $\{abcabcbcb\}^*$. The intersection of $\{a, bc\}^*$ and $\{a, cb\}^*$ is a^* .

We have shown that the intersection of two 2-maximal submonoids is generated by at most one element. Moreover, we know that the intersection of two 1-maximal submonoids is the empty word, i.e., it is generated by zero elements. Thus, it is natural to ask if in general, for every $k \ge 1$, the intersection of two k-maximal submonoids is generated by at most k - 1 elements. The following examples, suggested to us by Štěpán Holub, provide a negative answer to this question.

Example 16. The intersection of the two 3-maximal monoids $\{abc, dc, bab\}^*$ and $\{ab, cb, cd\}^*$ is infinitely generated by $abc(dc)^*bab$. The intersection of the two 4-maximal monoids $\{a, b, cd, ce\}^*$ and $\{ac, bc, da, ea\}^*$ is $\{acea, bcea, acda, bcda\}^*$.

Thus, our Theorem 14 is specific for rank 2 and cannot be generalized to larger k.

For an upper bound on the length of the word that generates the intersection of two 2-maximal submonoids, we have the following proposition.

Proposition 17. With the hypotheses of Theorem 14,

$$|z| < (|x| + |y|)(|u| + |v|).$$

Proof. Let \mathcal{A}_X (resp. \mathcal{A}_U) be the minimal DFA recognizing X^* (resp. U^*) and Q_X (resp. Q_U) its set of states. Since X and U are bifix codes, we have $|Q_X| < |x| + |y|$ and $|Q_U| < |u| + |v|$. Then the automaton \mathcal{A} recognizing $X^* \cap U^*$ has a set of states Q such that |Q| < (|x| + |y|)(|u| + |v|). By Theorem 14, \mathcal{A} is composed by only one cycle, labeled by z. Thus, |z| < (|x| + |y|)(|u| + |v|). \Box

For all our examples, the bound is much smaller than the previous one, hence we pose the following

Problem 18. Find a tight bound on the length of z in terms of the lengths of x and y and u and v.

4 Primitive Sets

We now show how the previous results can be interpreted in the terminology of combinatorics on words.

Let us start with the remark that a word $x \in A^+$ is primitive if and only if

$$x \in u^*, u \in A^+ \Rightarrow x = u.$$

With our definition of maximality, we have that a word $x \in A^+$ is primitive if and only if the monoid x^* is 1-maximal. Inspired by this observation, we give the following definition.

Definition 19. A finite set $X \subseteq A^*$ is primitive if it is the basis of a |X|-maximal submonoid.

Remark 20. The definition of primitive set does not coincide with that of elementary set. A set X is said to be elementary if r(X) = |X|. If X is primitive, then r(X) = |X|, i.e., it is elementary. But there exist elementary sets that are not primitive. For instance, the set $\{aa, bca\}$ is elementary, but it is not primitive since $\{aa, bca\}^* \subseteq \{a, bc\}^*$. The set $\{a, bc\}$, instead, is primitive.

From the definition of primitive set, we have that for every set X there exists a primitive set Y such that $X \subseteq Y^*$. The set Y is therefore called a *primitive root* of X. However, the primitive root of a set is not, in general, unique. Consider for instance the set $X = \{abcbab, abcdcbab, abcdccbab\}$. It has rank 3, hence it is elementary, yet it is not primitive. Indeed, $X \subseteq \{ab, cb, cd\}^*$. The set $\{ab, cb, cd\}$ is primitive, and it is a primitive root of X. However, it is not the only primitive root of X: the set $\{abc, dc, bab\}$ is primitive and $X \subseteq \{abc, dc, bab\}^*$, hence $\{abc, dc, bab\}$ is another primitive root of X. In the special case of sets of rank 1, clearly these always have a unique primitive root. For instance, the primitive root of the set $\{abab, abababab\}$ is the set $\{ab\}$.

However, as a consequence of Theorem 14 we have the following result.

Theorem 21. A set X of rank 2 has a unique primitive root.

Proof. If $\{u_1, u_2\}$ and $\{v_1, v_2\}$ are two primitive roots of X then $X^* \subseteq \{u_1, u_2\}^* \cap \{v_1, v_2\}^*$. Hence, by Theorem 14, $X \subseteq \{z\}^*$, for some primitive word z, i.e. r(X) = 1, a contradiction.

In what follows, we find convenient call a primitive set of cardinality 2 a *primitive pair*.

Example 22. The words abca and bc are primitive words, yet the pair $\{abca, bc\}$ is not a primitive pair, since $\{abca, bc\}^* \subseteq \{a, bc\}^*$, hence $\{abca, bc\}^*$ is not 2maximal. The pair $\{abcabc, bcabca\}$ can be written as concatenations of copies of both $\{abca, bc\}$ and $\{a, bcabc\}$. However, there is a unique way to decompose each word of the pair $\{abcabc, bcabca\}$ as a concatenation of words of a primitive pair, and this pair is $\{a, bc\}$. Indeed, the primitive root of $\{abcabc, bcabca\}$ is $\{a, bc\}$.

As it is well known, a primitive word x does not have internal occurrences in xx. The next result, whose proof is omitted for brevity, provides a similar property in the case of a primitive set of two words.

Theorem 23. Let $\{x, y\}$ be a primitive pair. Then neither xy nor yx occurs internally in a word of $\{x, y\}^3$.

Example 24. Let x = abcabca, y = bcaabcabc. Then xy has an internal occurrence in yxx, yet $\{x, y\} \subset \{a, bc\}^*$. This example shows that the hypothesis that $\{x, y\}$ is primitive cannot be replaced by simply requiring that x and y are primitive words.

Differently to the case of a single primitive word, the converse of Theorem 23 does not hold. For example, $\{abcaa, bc\}$ is not primitive $(\{abcaa, bc\}^* \subseteq \{a, bc\}^*)$, yet neither *abcaabc* nor *bcabcaa* occurs internally in a word of $\{x, y\}^3$.

5 Binary Root of a Single Primitive Word

In this section, we derive some consequences on the combinatorics of a *single* word. In particular, we introduce the notion of binary root of a primitive word, and we show how this notion may be useful to reveal some hidden repetitive structure in the word.

Let w be a nonempty word. If w is not primitive, then it can be written in a unique way as a concatenation of copies of a primitive word r, called the root of w. However, if w is primitive, one can ask whether it can be written as a concatenation of copies of two words x and y. If we further require that $\{x, y\}$ is a primitive set, then we call $\{x, y\}$ a *binary root* of the word w. Note that the binary root of a single word is not, in general, unique. For instance, for w = abcbac we have $w = ab \cdot cbac = abcb \cdot ac$ and $\{ab, cbac\}$ and $\{abcb, ac\}$ are both primitive pairs, i.e., they are both binary roots of w. However, if we additionally require that the size |x| + |y| of the binary root $\{x, y\}$ is "short" with respect to the length of w, then we obtain again the uniqueness. This is shown in the next theorem.

Theorem 25. Let w be a primitive word. Then w has at most one binary root $\{x, y\}$ such that $|x| + |y| < \sqrt{|w|}$.

Proof. Suppose by contradiction there exists another binary root $\{u, v\}$ of w with $|u| + |v| < \sqrt{|w|}$. Take $X = \{x, y\}$ and $U = \{u, v\}$. By Theorem 14, there exists a primitive word z and an integer n such that $w = z^n$. As w is primitive, w = z and n = 1. By Proposition 17, we have that $|w| < (|x| + |y|)(|u| + |v|) < \sqrt{|w|} \cdot \sqrt{|w|} = |w|$, a contradiction.

The following example shows a word w that has binary roots of different sizes, but only one of size less than $\sqrt{|w|}$.

Example 26. Consider the primitive word w = abcaabcabc of length 10. The pair $\{a, bc\}$ is the only binary root of w of size smaller than $\sqrt{|w|}$.

Asking for a tight bound in the statement of Theorem 25 is of course a problem intimately related to Open Problem 18.

We observe that both the classical notion of root and that of binary root are related to some repetitive structure inside the word. If w is not primitive, the length of its root reveals its repetitive structure in the sense that, if such a length is much smaller than the length of w, then the word w can be considered highly repetitive. If w is primitive, the size of its binary root plays an analogous role. This could be illustrated by the following (negative) example. Consider a word wover the alphabet A such that all the letters of w are distinct, so that |w| = |A|. This word is not repetitive at all, and it has |w| - 1 different binary roots $\{x, y\}$, all of size |w|, corresponding to the trivial factorizations w = xy. Thus, the absence of repetitions in a word is related to the large size of its binary roots. On the contrary, the existence in a word w of a "short" (with respect to |w|) binary root corresponds to the existence of some hidden repetitive structure in the word. This approach generalizes some already-considered notions of hidden repetitions (cf. [7,6,5]).

We think that the notion of a binary root can be further explored and may have applications, e.g., in the area of string algorithms.

Notice that the minimal length of a binary root (intended as the sum of the lengths of the two components of the pair) is affected by the combinatorial properties of the word. For example, if w is a square-free word, then w cannot have a binary root $\{x, y\}$ such that |x| + |y| < |w|/4, since otherwise w would

contain a square (xx, yy, xyxy or yxyx). The previous remark suggests a possible link between the notion of a binary root and the classical notion of *binary pattern*, which has been deeply investigated in combinatorics on words and fully classified by J. Cassaigne [3] (see also [15, Chap. 3] for a survey).

6 Connections with Pseudo-Primitive Words

We now show how the notion of a primitive pair can be seen as a generalization of the notion of a pseudo-primitive word, with respect to an involutive (anti-)morphism θ , as introduced in [4].

A map $\theta : A^* \to A^*$ is a morphism (resp. antimorphism) if for each $u, v \in A^*$, $\theta(uv) = \theta(u)\theta(v)$ (resp. $\theta(uv) = \theta(v)\theta(u)$) — θ is an involution if $\theta(\theta(a)) = a$ for every $a \in A$.

Let θ be an involutive morphism or antimorphism other than the identity function. We say that a word $w \in A^*$ is a θ -power of t if $w \in t\{t, \theta(t)\}^*$. A word w is θ -primitive if there exists no nonempty word t such that w is a θ -power of t and |w| > |t|.

Theorem 27 ([4]). Given a word $w \in A^*$ and an involutive (anti-)morphism θ , there exists a unique θ -primitive word $u \in A^*$ such hat w is a θ -power of u. The word u is called the θ -root of w.

Example 28. Let $\theta : \{a, b, c\}^* \to \{a, b, c\}^*$ the involutive morphism defined by $\theta(a) = b, \ \theta(b) = a \ \text{and} \ \theta(c) = c$. The θ -root of the word *abcabcbac* is *abc*.

If θ is an involutive morphism, we show that Theorem 27 can be obtained as a consequence of Theorem 21. If θ is an involutive antimorphism, we obtain a slightly different formulation, from which we derive a new property of θ -primitive words.

Given a morphism θ and a set $X \subseteq A^*$, $\theta(X)$ denotes the set $\{\theta(u) \mid u \in X\}$. We say that X is θ -invariant if $\theta(X) \subseteq X$.

We have the following propositions.

Proposition 29. Let θ be involutive. If $\{x, y\}$ is θ -invariant, then so is its root.

Example 30. Let θ be as in Example 28. The pair {*abcabcbac*, *abcbacabc*} is θ -invariant. However, it is not a primitive pair. Its binary root is the pair {*abc*, *bac*}, which is θ -invariant since $\theta(abc) = bac$.

Remark 31. Let θ be an involutive morphism. Then $\{x, y\}$ is θ -invariant if and only if $y = \theta(x)$. If θ is an involutive antimorphism, then $\{x, y\}$ is θ -invariant if and only if either $y = \theta(x)$ or $x = \theta(x)$ and $y = \theta(y)$. In the last case, x and y are called θ -palindromes.

Example 32. Let $\theta : \{a, b, c\}^* \mapsto \{a, b, c\}^*$ be the involutive antimorphism defined by $\theta(a) = a, \theta(b) = b, \theta(c) = c$. The pair $\{abcbbcba, abcba\}$ is θ -invariant. Its binary root is $\{a, bcb\}$, which is θ -invariant since composed by θ -palindromes. With the same θ , the pair $\{abbbbabba, abbabbba\}$ is θ -invariant and its binary root is $\{abb, bba\}$, which is θ -invariant since $\theta(abb) = bba$.

Proposition 33. Let $w \in A^*$ and θ be an involutive morphism of A^* . Then, w is θ -primitive if and only if the pair $\{w, \theta(w)\}$ is a primitive pair.

Proof. Let us suppose, by contradiction, that $\{w, \theta(w)\}$ is a primitive pair and w is not θ -primitive. Then there exists t such that $w \in \{t, \theta(t)\}^*$. Hence, $\theta(w) \in \{t, \theta(t)\}^*$, so the pair $\{w, \theta(w)\}$ is not primitive. Conversely, let us suppose that w is θ -primitive and $\{w, \theta(w)\}$ is not a primitive pair. Denote by $\{u, v\}$ its binary root. Since $\{w, \theta(w)\}$ is θ -invariant, then $\{u, v\}$ is θ -invariant, i.e., $v = \theta(u)$. Hence, $w \in \{u, \theta(u)\}^*$, i.e., w is not θ -primitive.

From Theorem 21 and Proposition 33 we derive Theorem 27 when θ is an involutive morphism.

Now, let us consider the case of antimorphisms. Reasoning analogously as we did in the proof of Proposition 33, we can prove the following result.

Proposition 34. Let $w \in A^*$ and θ an involutive antimorphism of A^* . If the pair $\{w, \theta(w)\}$ is a primitive pair, then w is θ -primitive.

The converse does not hold in general, as the following example shows.

Example 35. Let θ be the antimorphic involution of Example 32. The word w = abbaabbacbc is θ -primitive, whereas the pair $\{w, \theta(w)\} = \{abbaabbacbc, cbcabbaabba\}$ is not a primitive pair, since its binary root is the pair $\{abba, cbc\}$.

Finally, we can state the following proposition, which provides a factorization property of θ -primitive words.

Proposition 36. Let $w \in A^*$ and θ an involutive antimorphism. If w is θ -primitive and $\{w, \theta(w)\}$ is not a primitive pair, then there exist two θ -palindromes p and q such that $w \in \{p, q\}^*$.

Proof. Suppose that $\{w, \theta(w)\}$ is not a primitive pair and denote by $\{u, v\}$ its binary root. Since $\{w, \theta(w)\}$ is θ -invariant, then so is $\{u, v\}$ by Proposition 29, and $v \neq \theta(u)$ since w is θ -primitive. Then, $u = \theta(u)$ and $v = \theta(v)$ are θ -palindromes.

Finally, we point out that our Theorem 23 can be viewed as a generalization of the following result of Kari, Masson and Seki [11]:

Theorem 37 (Theorem 12 of [11]). Let x be a nonempty θ -primitive word. Then neither $x\theta(x)$ nor $\theta(x)x$ occurs internally in a word of $\{x, \theta(x)\}^3$.

7 Acknowledgments

We thank Štěpán Holub for useful discussions and in particular for suggesting us the important Example 16.

References

- 1. J. Berstel, D. Perrin, J.F. Perrot, and A. Restivo. Sur le théorème du défaut. Journal of Algebra, 60(1):169–180, 1979.
- J. Berstel, D. Perrin, and C. Reutenauer. Codes and Automata (Encyclopedia of Mathematics and Its Applications). Cambridge University Press, New York, NY, USA, 1st edition, 2009.
- J. Cassaigne. Motifs évitables et régularités dans les mots. PhD thesis, Université Paris VI, 1994.
- E. Czeizler, L. Kari, and S. Seki. On a special class of primitive words. *Theoret. Comput. Sci.*, 411(3):617–630, 2010.
- 5. P. Gawrychowski, F. Manea, R. Mercas, and D. Nowotka. Hide and seek with repetitions. J. Comput. Syst. Sci., 101:42–67, 2019.
- P. Gawrychowski, F. Manea, R. Mercas, D. Nowotka, and C. Tiseanu. Finding pseudo-repetitions. In Natacha Portier and Thomas Wilke, editors, *STACS* 2013, Proceedings, volume 20 of *LIPIcs*, pages 257–268. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2013.
- P. Gawrychowski, F. Manea, and D. Nowotka. Discovering hidden repetitions in words. In Paola Bonizzoni, Vasco Brattka, and Benedikt Löwe, editors, *CiE 2013, Proceedings*, volume 7921 of *Lecture Notes in Computer Science*, pages 210–219. Springer, 2013.
- T. Harju and J. Karhumäki. On the defect theorem and simplifiability. Semigroup Forum, 33(1):199–217, 1986.
- T. Harju and J. Karhumäki. Many aspects of defect theorems. *Theoret. Comput. Sci.*, 324(1):35–54, 2004.
- J. Karhumäki. A note on intersections of free submonoids of a free monoid. Semigroup Forum, 29(1):183–205, 1984.
- L. Kari, B. Masson, and S. Seki. Properties of pseudo-primitive words and their applications. Int. J. Found. Comput. Sci., 22(2):447–471, 2011.
- E.B. Le Rest and M. Le Rest. Sur la combinatoire des codes à deux mots. *Theoret. Comput. Sci.*, 41(C):61–80, 1985.
- A. Lentin and M. Schützenberger. A combinatorial problem in the theory of free monoids. Proc. University of North Carolina, pages 128–144, 1967.
- 14. M. Lothaire. Combinatorics on Words. Addison-Wesley, 1983.
- 15. M. Lothaire. *Algebraic Combinatorics on Words*. Cambridge University Press, 2002.
- J. Néraud. Elementariness of a finite set of words is co-NP-complete. ITA, 24:459– 470, 1990.
- J. Néraud. Deciding whether a finite set of words has rank at most two. Theoret. Comput. Sci., 112(2):311–337, 1993.
- Jean Néraud. On the rank of the subsets of a free monoid. Theoret. Comput. Sci., 99(2):231–241, 1992.
- A. Restivo. On a question of McNaughton and Papert. Information and Control, 25(1):93–101, 1974.
- 20. B. Tilson. The intersection of free submonoids of a free monoid is free. Semigroup Forum, 4(1):345–350, 1972.