

Interpretation of the newly observed $\Sigma_b(6097)^\pm$ and $\Xi_b(6227)^-$ states as the P -wave bottom baryons

Kai-Lei Wang¹, Qi-Fang Lü^{1,2}*, Xian-Hui Zhong^{1,2}†

1) Department of Physics, Hunan Normal University, and Key Laboratory of Low-Dimensional Quantum Structures and Quantum Control of Ministry of Education, Changsha 410081, China and

2) Synergetic Innovation Center for Quantum Effects and Applications (SICQEA), Hunan Normal University, Changsha 410081, China

The strong decays of the P -wave singly bottom heavy baryons belonging to $\mathbf{6}_F$ are investigated with a constituent quark model in the j - j coupling scheme. The results show that the newly observed $\Sigma_b(6097)$ and $\Xi_b(6227)$ states by the LHCb collaboration can be assigned as the λ -mode P -wave bottom baryons belonging to $\mathbf{6}_F$. Given the heavy quark symmetry, both the $\Sigma_b(6097)$ and $\Xi_b(6227)$ states favor the light spin $j = 2$ states with spin-parity numbers $J^P = 3/2^-$ or $J^P = 5/2^-$. More P -wave baryons in the Σ_b , Ξ'_b , and Ω_b families are most likely to be observed in future experiments for their relatively narrow width.

PACS numbers:

I. INTRODUCTION

Recently, the LHCb collaboration announced the observation of a new bottom baryon $\Xi_b(6227)^-$ in both $\Lambda_b^0 K^-$ and $\Xi_b^0 \pi^-$ decay modes [1]. Just several weeks ago, the LHCb collaboration found two new resonances $\Sigma_b(6097)^\pm$ in the $\Lambda_b^0 \pi^\pm$ channels [2]. The measured masses and widths of the $\Xi_b(6227)^-$ and $\Sigma_b(6097)^\pm$ are presented as follows,

$$m[\Xi_b(6227)^-] = 6226.9 \pm 2.0 \pm 0.3 \pm 0.2 \text{ MeV}, \quad (1)$$

$$\Gamma[\Xi_b(6227)^-] = 18.1 \pm 5.4 \pm 1.8 \text{ MeV}, \quad (2)$$

$$m[\Sigma_b(6097)^-] = 6098.0 \pm 1.7 \pm 0.5 \text{ MeV}, \quad (3)$$

$$\Gamma[\Sigma_b(6097)^-] = 28.9 \pm 4.2 \pm 0.9 \text{ MeV}, \quad (4)$$

$$m[\Sigma_b(6097)^+] = 6095.8 \pm 1.7 \pm 0.4 \text{ MeV}, \quad (5)$$

$$\Gamma[\Sigma_b(6097)^+] = 31.0 \pm 5.5 \pm 0.7 \text{ MeV}. \quad (6)$$

Compared with the mass spectrum predicted in various models [3–13], it is found that the $\Xi_b(6227)^-$ and $\Sigma_b(6097)^\pm$ are good candidates of the λ -mode P -wave bottom baryons belonging to $\mathbf{6}_F$.

To explain the nature of $\Xi_b(6227)^-$ and $\Sigma_b(6097)^\pm$, recently the strong decays of the excited singly bottom states were studied with the 3P_0 model in Ref. [14, 15]. The results indicate that $\Xi_b(6227)^-$ and $\Sigma_b(6097)^\pm$ may be good candidates of the P -wave states with $J^P = 3/2^-$ or $J^P = 5/2^-$ in the j - j coupling scheme. In Ref. [16], we studied the strong decays of the singly bottom baryons within the chiral quark model in the L - S coupling scheme. Comparing the predictions with the observations, one can find that both $\Xi_b(6227)$ and $\Sigma_b(6097)$ could be assigned as the P -wave states with $J^P = 3/2^-$ or

$J^P = 5/2^-$ as well. However, the partial width ratios and decay modes of some P -wave Ξ'_b states predicted within the j - j coupling scheme are different from those predicted within the L - S coupling scheme.

We have noted that for the singly heavy baryons there may exist some effects of heavy quark symmetry, which was not carefully considered in our previous work [16]. To approximately preserve the heavy quark symmetry with the finite heavy quark masses, the physical states may be closer to the j - j coupling basis. In this coupling scheme, the states are mixed states between the states of the L - S coupling scheme with the same spin-parity (J^P) numbers. It is interesting to find that the newly observed state $\Omega_c(3000)$ at LHCb [17] may be explained as a $J^P = 1/2^-$ resonance via a $|^2P_{1/2}\rangle$ - $|^4P_{1/2}\rangle$ mixing [18]. This may be evidence that the physical states of singly heavy baryons more favor the j - j coupling scheme. In another word, the configuration mixing effects between the singly heavy baryon states with the same J^P should be considered if one adopts the L - S coupling scheme.

In this work, considering the requirements of the heavy quark symmetry we study the strong decay behaviors of the λ -mode P -wave bottom baryons Σ_b , Ξ'_b and Ω_b in the j - j coupling scheme with the chiral quark model. In this coupling scheme, the states with light spin $j = 0$ and $j = 1$ cannot decay into some special final states, which shows selection rules. The decay modes together with the predicted total widths suggest that both the $\Xi_b(6227)^-$ and $\Sigma_b(6097)^\pm$ are good candidates of light spin $j = 2$ states, while the $J^P = 3/2^-$ and $J^P = 5/2^-$ cannot be distinguished due to lack of data. The predicted narrow widths of other flavor symmetric partners can provide helpful information for future experiments.

This paper is organized as follows. The spectrum and notations are presented in Sec. II. The chiral quark model is briefly introduced in Sec. III. The strong decays of the λ -mode P -wave Σ_b , Ξ'_b and Ω_b are estimated in Sec. IV. A short summary is presented in the last section.

II. SPECTROSCOPY

The heavy baryon containing a heavy quark violates the $SU(4)$ symmetry. However, the $SU(3)$ symmetry between the other two light quarks (u , d , or s) is approximately kept. Ac-

*E-mail: lvqifang@hunnu.edu.cn

†E-mail: zhongxh@hunnu.edu.cn

cording to the symmetry, the heavy baryons containing a single heavy quark belong to two different SU(3) flavor representations: the symmetric sextet $\mathbf{6}_F$ and antisymmetric antitriplet $\bar{\mathbf{3}}_F$ [16]. For example, the $\Sigma_b^{-,0,+}$, $\Xi_b^{0,-}$ and Ω_b form a $\mathbf{6}_F$ representation, while Λ_b and $\Xi_b^{0,-}$ form a $\bar{\mathbf{3}}_F$ representation.

In the L - S coupling scheme, the quark model states are constructed by [8]

$$|^{2S+1}L_J\rangle = \left| \left[\left[(\ell_\rho \ell_\lambda)_L (s_\rho s_Q)_S \right]_{J^P} \right] \right\rangle, \quad (7)$$

where ℓ_ρ and ℓ_λ correspond to the eigenvalues of the orbital angular momenta ℓ_ρ and ℓ_λ for the ρ and λ oscillators, respectively, and L stands for the eigenvalue of the total orbital angular momentum $\mathbf{L} = \ell_\rho + \ell_\lambda$ [18, 25]. The parity of a state is determined by $P = (-1)^{\ell_\rho + \ell_\lambda}$. Quantum numbers s_ρ and s_Q correspond to the eigenvalues of the total spin of the two light quarks s_ρ and the spin of the heavy quark s_Q , respectively, while S stands for the eigenvalue of the total spin angular momentum $\mathbf{S} = \mathbf{s}_\rho + \mathbf{s}_Q$. According to the quark model classification, there are five λ -mode $1P$ -wave states belonging to $\mathbf{6}_F$ [16]. These states and their corresponding quantum numbers have been collected in Table I.

TABLE I: The classifications of the low-lying P -wave states belonging to $\mathbf{6}_F$ in the L - S coupling scheme. The states in the L - S coupling scheme are denoted by $|^{2S+1}L_J\rangle$.

$ ^{2S+1}L_J\rangle$	J^P	ℓ_ρ	ℓ_λ	L	s_ρ	s_Q	S
$ ^2P_{1/2}\rangle$	$\frac{1}{2}^-$	0	1	1	1	$\frac{1}{2}$	$\frac{1}{2}$
$ ^2P_{3/2}\rangle$	$\frac{3}{2}^-$	0	1	1	1	$\frac{1}{2}$	$\frac{1}{2}$
$ ^4P_{1/2}\rangle$	$\frac{1}{2}^-$	0	1	1	1	$\frac{1}{2}$	$\frac{3}{2}$
$ ^4P_{3/2}\rangle$	$\frac{3}{2}^-$	0	1	1	1	$\frac{1}{2}$	$\frac{3}{2}$
$ ^4P_{5/2}\rangle$	$\frac{5}{2}^-$	0	1	1	1	$\frac{1}{2}$	$\frac{5}{2}$

TABLE II: The classifications of the low-lying P -wave states belonging to $\mathbf{6}_F$ in the j - j coupling scheme. The states in the j - j coupling scheme are denoted by $|J^P, j\rangle$.

$ J^P, j\rangle$	J^P	j	ℓ_ρ	ℓ_λ	L	s_ρ	s_Q
$ J^P = \frac{1}{2}^-, 0\rangle$	$\frac{1}{2}^-$	0	0	1	1	1	$\frac{1}{2}$
$ J^P = \frac{1}{2}^-, 1\rangle$	$\frac{1}{2}^-$	1	0	1	1	1	$\frac{1}{2}$
$ J^P = \frac{3}{2}^-, 1\rangle$	$\frac{3}{2}^-$	1	0	1	1	1	$\frac{1}{2}$
$ J^P = \frac{3}{2}^-, 2\rangle$	$\frac{3}{2}^-$	2	0	1	1	1	$\frac{1}{2}$
$ J^P = \frac{5}{2}^-, 2\rangle$	$\frac{5}{2}^-$	2	0	1	1	1	$\frac{1}{2}$

It should be pointed out that the heavy-quark symmetry as an approximation is commonly adopted for the study of the singly heavy baryons. In the heavy quark effective theory description, the spin of the heavy quark s_Q and the total angular momentum of the two light quarks $\mathbf{j} = \mathbf{L} + \mathbf{s}_\rho$ are separately conserved [19]. The total angular momentum is given by $\mathbf{J} = \mathbf{j} + \mathbf{s}_Q$. In the heavy-quark symmetry limit, the quark model states may more favor the j - j coupling scheme

$$|J^P, j\rangle = \left| \left[\left[(\ell_\rho \ell_\lambda)_L (s_\rho)_j (s_Q) \right]_{J^P} \right] \right\rangle. \quad (8)$$

The five $1P$ -wave states belonging to $\mathbf{6}_F$ in the j - j coupling scheme and their corresponding quantum numbers have been collected in Table II.

The states of the j - j coupling scheme are linear combinations of the states of the L - S coupling scheme. The precise relationship is [8]

$$\left\{ \left[\left[(\ell_\rho \ell_\lambda)_L (s_\rho)_j (s_Q) \right]_{J^P} \right] \right\} = (-1)^{L+s_\rho+J+\frac{1}{2}} \sqrt{2j+1} \sum_S \sqrt{2S+1} \begin{Bmatrix} L & s_\rho & j \\ s_Q & J & S \end{Bmatrix} \left| \left[(\ell_\rho \ell_\lambda)_L (s_\rho s_Q)_S \right]_{J^P} \right\rangle. \quad (9)$$

The heavy-quark symmetry may tell us that there are configuration mixing between the singly heavy baryon states with the same J^P numbers in the L - S coupling scheme. In the heavy quark limit, the mixing angles are determined by Eq.(9).

III. STRONG DECAY

To study the strong decays of the low-lying P -wave bottom baryons, we apply the chiral quark model [20]. In this model, the light pseudoscalar mesons, i.e. π , K and η , are treated as point-like Goldstone boson. This method has been successfully applied to the strong decays of heavy-light mesons, charmed and strange baryons [18, 21–30]. The nonrelativistic form of the effective quark-pseudoscalar-meson interactions might be described by [18, 21–27]:

$$H_m^{nr} = \sum_j \left[\mathcal{G} \boldsymbol{\sigma}_j \cdot \mathbf{q} + h \boldsymbol{\sigma}_j \cdot \mathbf{p}_j \right] I_j e^{-i\mathbf{q} \cdot \mathbf{r}_j}, \quad (10)$$

with $\mathcal{G} \equiv -\left(1 + \frac{\omega_m}{E_f + M_f}\right)$, $h \equiv \frac{\omega_m}{2\mu_q}$. In the above equation, $\boldsymbol{\sigma}_j$ and \mathbf{p}_j stand for the Pauli spin vector and internal momentum operator for the j th quark of the initial hadron; ω_m and \mathbf{q} stand for the energy and three momenta of the emitted light meson, respectively; E_f and M_f are the energy and mass of the final heavy baryon; μ_q is a reduced mass given by $1/\mu_q = 1/m_j + 1/m'_j$ with m_j and m'_j for the masses of the j th quark in the initial and final hadrons, respectively; and I_j is the isospin operator associated with the pseudoscalar mesons, which have been defined in Refs. [22, 25]. It should be mentioned that the nonrelativistic form of quark-pseudoscalar-meson interactions expressed in Eq. (10) is similar to that in Refs. [3, 31, 32], except that the factors \mathcal{G} and h in this work have explicit dependence on the energies of final hadrons.

For a light pseudoscalar meson emission in a strong decay process, the partial decay width can be calculated with [22, 25]

$$\Gamma_m = \left(\frac{\delta}{f_m} \right)^2 \frac{(E_f + M_f)|\mathbf{q}|}{4\pi M_i(2J_i + 1)} \sum_{J_{fz}, J_{iz}} |\mathcal{M}_{J_{fz}, J_{iz}}|^2, \quad (11)$$

where $\mathcal{M}_{J_{fz}, J_{iz}}$ corresponds to the strong amplitudes. The quantum numbers J_{iz} and J_{fz} stand for the third components of the total angular momenta of the initial and final heavy baryons, respectively. M_i is the mass of the initial heavy

baryon. δ as a global parameter accounts for the strength of the quark-meson couplings. It has been determined in our previous study of the strong decays of the charmed baryons and heavy-light mesons [22, 25]. Here, we fix its value the same as that in Refs. [22, 25], i.e. $\delta = 0.557$.

In the calculation, we adopt the same quark model parameter set as that in Refs. [16, 30]. The masses of the well-established hadrons used in the calculations are taken from the Particle Data Group (PDG) [33], and the masses of the undiscovered initial states adopted from the predictions in Ref. [4].




FIG. 1: Partial and total strong decay widths of the the $J^P = 1/2^-$ and $J^P = 3/2^-$ states with $M = 6097$ MeV in the Σ_b families as a function of mixing angle ϕ . The solid curves stand for the total widths.

IV. RESULTS AND DISCUSSIONS

A. $\Sigma_b(6097)^\pm$ as the P -wave Σ_b states

The measured mass of the $\Sigma_b(6097)$ indicates that it is a good candidate of the λ -mode P -wave excitations. In the L - S coupling scheme, the strong decay properties of the P wave Σ_b baryons have been studied in Ref. [16] (see Table. III). From the decay behaviors, it is found that $\Sigma_b(6097)$ favors the $J^P = 3/2^-$ $|\Sigma_b {}^2P_{3/2}\rangle$ state or the $J^P = 5/2^-$ $|\Sigma_b {}^4P_{5/2}\rangle$ state. Moreover, the possibilities of $\Sigma_b(6097)$ as a candidate of the other $1P$ wave Σ_b states cannot be excluded, because their predicted total widths are close to that of $\Sigma_b(6097)$, and the $\Lambda_b\pi$ decay mode is allowed.

1. $J^P = 1/2^-$ states

In the j - j coupling scheme, according to Eq. (9) we can obtain two $J^P = 1/2^-$ states as mixed states between $|{}^2P_{1/2}\rangle$ and $|{}^4P_{1/2}\rangle$:

$$\left|J^P = \frac{1}{2}^-, 0\right\rangle = -\sqrt{\frac{1}{3}}|{}^2P_{1/2}\rangle + \sqrt{\frac{2}{3}}|{}^4P_{1/2}\rangle, \quad (12)$$

$$\left|J^P = \frac{1}{2}^-, 1\right\rangle = \sqrt{\frac{2}{3}}|{}^2P_{1/2}\rangle + \sqrt{\frac{1}{3}}|{}^4P_{1/2}\rangle. \quad (13)$$

If let $\cos\phi = \sqrt{\frac{2}{3}}$ and $\sin\phi = \sqrt{\frac{1}{3}}$, we get the mixing angle $\phi \approx 35^\circ$. The strong decay properties of $|J^P = \frac{1}{2}^-, 0\rangle$ and $|J^P = \frac{1}{2}^-, 1\rangle$ are calculated within the chiral quark model, our results are presented in Table. III. It is seen that the $|J^P = \frac{1}{2}^-, 0\rangle$ is a narrow state with a width of a few MeV, and mainly decays into $\Lambda_b\pi$; while the $|J^P = \frac{1}{2}^-, 1\rangle$ state has a width of ~ 30 MeV, and dominantly decays into $\Sigma_b\pi$. For a comparison, in Table. III we also list the strong decay properties of the $J^P = 1/2^-$ states $|{}^2P_{1/2}\rangle$ and $|{}^4P_{1/2}\rangle$ obtained from the L - S coupling scheme in Ref. [16]. It is found that the decay properties of $|J^P = \frac{1}{2}^-, 1\rangle$ in the j - j coupling scheme are similar to those of $|{}^2P_{1/2}\rangle$ in the L - S coupling scheme, for the strong decays of $|J^P = \frac{1}{2}^-, 1\rangle$ are governed by the component of $|{}^2P_{1/2}\rangle$. However, the strong decay properties of $|J^P = \frac{1}{2}^-, 0\rangle$ are very different from those in the L - S coupling scheme. The decay properties of the $J^P = 1/2^-$ states are inconsistent with the observations of $\Sigma_b(6097)$, thus, the $\Sigma_b(6097)$ as the $J^P = 1/2^-$ states should be excluded.

It is interesting to note that the $\Omega_c(3000)$ resonance may be explained as a mixed state $|P_\lambda \frac{1}{2}^-\rangle_1 = \cos\phi|{}^2P_{1/2}\rangle + \sin\phi|{}^4P_{1/2}\rangle$ with the mixing angle $\phi \approx 24^\circ$ [18], which is close to $\phi \approx 35^\circ$ determined by the heavy-quark symmetry in the j - j coupling scheme, but not actually equal to that determined by the heavy-quark symmetry. This indicates that on the one hand the heavy-quark symmetry requires the physical states might be mixed states between the states with the same J^P in the L - S coupling scheme, on the other hand because the heavy-quark symmetry is only an approximation, the physical mixing angle should be slightly different from that determined within the j - j coupling scheme. Thus, the physical P -wave singly-heavy baryon states might be mixed states between $|{}^2P_J\rangle$ and $|{}^4P_J\rangle$ in the L - S coupling scheme, i.e.,

$$\begin{pmatrix} |P J^P\rangle_1 \\ |P J^P\rangle_2 \end{pmatrix} = \begin{pmatrix} \cos\phi & \sin\phi \\ -\sin\phi & \cos\phi \end{pmatrix} \begin{pmatrix} |{}^2P_J\rangle \\ |{}^4P_J\rangle \end{pmatrix}, \quad (14)$$

and the mixing angle ϕ may range from the value of the L - S coupling scheme ($\phi = 0^\circ$) to that of the j - j coupling scheme. Specially, for the P -wave states with $J^P = 1/2^-$ the mixing angle $\phi \in (0, 35^\circ)$, while for the P -wave states with $J^P = 3/2^-$ the mixing angle $\phi \in (0, 24^\circ)$.

According to the mixing scheme defined in Eq. (14), the strong decay widths of the $J^P = 1/2^-$ states in the Σ_b family

TABLE III: Partial widths (MeV) and branching fractions for the strong decays of the $1P$ -wave states in the j - j coupling scheme compared with that in the L - S coupling scheme taken from Ref. [16] in the Σ_b family. The masses of the P -wave bottom baryons are adopted from the quark model predictions in Ref. [4].

State	$ J, j\rangle$	Channel	Γ_i (MeV)	\mathcal{B}_i	$ ^{2S+1}L_J\rangle$	Channel	Γ_i (MeV) [16]	\mathcal{B}_i [16]
$\Sigma_b(6101)$	$ J = \frac{1}{2}^-, 1\rangle$	$\Lambda_b\pi$	\dots	\dots	$ ^2P_{1/2}\rangle$	$\Lambda_b\pi$	1.74	7.68%
		$\Sigma_b\pi$	28.89	92.04%		$\Sigma_b\pi$	19.26	85.00%
		$\Sigma_b^*\pi$	2.50	7.96%		$\Sigma_b^*\pi$	1.66	7.33%
		total	31.39			total	22.66	
$\Sigma_b(6095)$	$ J = \frac{1}{2}^-, 0\rangle$	$\Lambda_b\pi$	6.00	100.0%	$ ^4P_{1/2}\rangle$	$\Lambda_b\pi$	4.00	28.15%
		$\Sigma_b\pi$	\dots	\dots		$\Sigma_b\pi$	9.50	66.85%
		$\Sigma_b^*\pi$	\dots	\dots		$\Sigma_b^*\pi$	0.71	5.0%
		total	6.00			total	14.21	
$\Sigma_b(6096)$	$ J = \frac{3}{2}^-, 2\rangle$	$\Lambda_b\pi$	35.18	90.07%	$ ^2P_{3/2}\rangle$	$\Lambda_b\pi$	29.31	74.60%
		$\Sigma_b\pi$	3.25	8.32%		$\Sigma_b\pi$	4.81	12.24%
		$\Sigma_b^*\pi$	0.63	1.61%		$\Sigma_b^*\pi$	5.17	13.16%
		total	39.06			total	39.29	
$\Sigma_b(6087)$	$ J = \frac{3}{2}^-, 1\rangle$	$\Lambda_b\pi$	\dots	\dots	$ ^4P_{3/2}\rangle$	$\Lambda_b\pi$	5.38	20.46%
		$\Sigma_b\pi$	1.47	5.28%		$\Sigma_b\pi$	0.20	0.76%
		$\Sigma_b^*\pi$	26.39	94.72%		$\Sigma_b^*\pi$	20.71	78.78%
		total	27.86			total	26.29	
$\Sigma_b(6084)$	$ J = \frac{5}{2}^-, 2\rangle$	$\Lambda_b\pi$	31.38	81.85%	$ ^4P_{5/2}\rangle$	$\Lambda_b\pi$	31.38	81.85%
		$\Sigma_b\pi$	1.09	2.84%		$\Sigma_b\pi$	1.09	2.84%
		$\Sigma_b^*\pi$	5.77	15.05%		$\Sigma_b^*\pi$	5.77	15.05%
		total	38.34			total	38.34	

as a function of the mixing angle ϕ are shown in Fig. 1. It is found that the mixed state $|\Sigma_b P_{\frac{1}{2}}^-\rangle_1$ has a width of $\sim 20 - 30$ MeV, its decays are dominated by the $\Sigma_b\pi$ channel, the decay properties are less sensitive to the mixing angle. If the mixing angle is less than the angle $\phi \simeq 35^\circ$ obtained in the heavy quark limit, there is a small decay rate into the $\Lambda_b\pi$ channel. The other mixed state $|\Sigma_b P_{\frac{1}{2}}^-\rangle_2$ has a relatively narrower width, which is sensitive to the mixing angle. If we take the same mixing angle $\phi \simeq 24^\circ$ as that of $\Omega_c(3000)$ [18], except for the main decay mode $\Lambda_b\pi$, the $\Sigma_b\pi$ may play an obvious role in its strong decays. The decay properties of $|\Sigma_b P_{\frac{1}{2}}^-\rangle_1$ and $|\Sigma_b P_{\frac{1}{2}}^-\rangle_2$ are inconsistent with the observations of $\Sigma_b(6097)$.

2. $J^P = 3/2^-$ states

For the $J^P = 3/2^-$ states $|J^P = \frac{3}{2}^-, 1\rangle$ and $|J^P = \frac{3}{2}^-, 2\rangle$ in the j - j coupling scheme, according to Eq. (9) they can relate to the states in the L - S coupling scheme:

$$\left| J^P = \frac{3}{2}^-, 1 \right\rangle = -\sqrt{\frac{1}{6}}|^2P_{3/2}\rangle + \sqrt{\frac{5}{6}}|^4P_{3/2}\rangle, \quad (15)$$

$$\left| J^P = \frac{3}{2}^-, 2 \right\rangle = \sqrt{\frac{5}{6}}|^2P_{3/2}\rangle + \sqrt{\frac{1}{6}}|^4P_{3/2}\rangle. \quad (16)$$

If let $\cos \phi = \sqrt{\frac{5}{6}}$ and $\sin \phi = \sqrt{\frac{1}{6}}$, we get the mixing angle $\phi \simeq 24^\circ$. The decay properties of $|J^P = \frac{3}{2}^-, 1\rangle$ and $|J^P = \frac{3}{2}^-, 2\rangle$ in the Σ_b family are calculated within the chiral quark model. Our results are listed in Table. III as well. From the Table. III, it is found that the $|J^P = \frac{3}{2}^-, 1\rangle$ state has a width of ~ 28 MeV, and dominantly decays into the $\Sigma_b^*\pi$ channel; while the $|J^P = \frac{3}{2}^-, 2\rangle$ state has a relatively broad width of ~ 39 MeV, and dominantly decays into the $\Lambda_b\pi$ channel. Both the decay mode and width indicate that $\Sigma_b(6097)$ is a good candidate of the light spin $j = 2$ state $|J^P = \frac{3}{2}^-, 2\rangle$. The partial width ratio between $\Lambda_b\pi$ and $\Sigma_b\pi$ for $|J^P = \frac{3}{2}^-, 2\rangle$ is predicted to be

$$\frac{\Gamma[\Lambda_b\pi]}{\Gamma[\Sigma_b\pi]} \simeq 12, \quad (17)$$

which can be tested in future experiments. It should be mentioned that the main decay properties, such as the total decay width and dominant decay modes, of the $J^P = 3/2^-$ states $|J^P = \frac{3}{2}^-, 1\rangle$ and $|J^P = \frac{3}{2}^-, 2\rangle$ in the j - j coupling scheme are similar to those of $|^4P_{3/2}\rangle$ and $|^2P_{3/2}\rangle$ in the L - S coupling scheme, respectively (see Table. IV).

Considering the mixing angle of the physical states with $J^P = 3/2^-$ may have some deviations from the j - j couplings, adopting the mixing scheme defined in Eq. (14) we also plot the strong decay width as a function of mixing angle ϕ in Fig. 1 for a reference. It is shown that when ϕ varies in $0 \sim 24^\circ$, the total widths of both $|\Sigma_b P_{\frac{3}{2}}^-\rangle_1$ and $|\Sigma_b P_{\frac{3}{2}}^-\rangle_2$ are less sensitive to the mixing angle, however, the partial width

ratio of $\Gamma[\Lambda_b\pi]/\Gamma[\Sigma_b^*\pi]$ is very sensitive to the mixing angle. The measurements of the ratio $\Gamma[\Lambda_b\pi]/\Gamma[\Sigma_b^*\pi]$ might be helpful to determine the mixing angle.

3. $J^P = 5/2^-$ state

For the $J^P = 5/2^-$ state, the L - S coupling scheme is equal to the j - j coupling scheme, i.e.,

$$\left| J^P = \frac{5^-}{2}, 2 \right\rangle = \left| {}^4P_{5/2} \right\rangle. \quad (18)$$

From Table III, it is seen that both the total decay width and the dominant $\Lambda_b\pi$ decay mode of $|J^P = \frac{5^-}{2}, 2\rangle$ also favor the observations of $\Sigma_b(6097)$. The decay rate of the $J^P = 5/2^-$ state into $\Sigma_b^*\pi$ is sizable as well. The partial width ratio between the two main decay channels $\Lambda_b\pi$ and $\Sigma_b^*\pi$ is predicted to be

$$\frac{\Gamma[\Lambda_b\pi]}{\Gamma[\Sigma_b^*\pi]} \simeq 5. \quad (19)$$

If the $\Sigma_b(6097)$ resonance is the $J^P = 5/2^-$ state indeed, it should be observed in the $\Sigma_b^*\pi$ channel as well.

As a whole, $\Sigma_b(6097)$ seems to favor the $j = 2$ states with $J^P = 3/2^-$ or $J^P = 5/2^-$. To distinguish these two states, more observations in the $\Sigma_b\pi$ and $\Sigma_b^*\pi$ channels are needed in future experiments. If $\Sigma_b(6097)$ corresponds to $|J^P = \frac{3^-}{2}, 2\rangle$, it should be observed in the $\Sigma_b\pi$ channel. On the other hand, if $\Sigma_b(6097)$ is the $J^P = 5/2^-$ state, it should be observed in the $\Sigma_b^*\pi$ channel.

B. $\Xi_b'(6227)^-$ as the P -wave Ξ_b' states

The measured mass of $\Xi_b(6227)$ indicates that it is a good candidate of the λ -mode P -wave excitations [4]. Within the L - S coupling scheme, the strong decay behaviors of the P wave Ξ_b' states have been studied in Ref. [16]. It is found that $\Xi_b(6227)$ favors the $J^P = 3/2^-$ $|\Xi_b' {}^2P_{3/2}\rangle$ state or $J^P = 5/2^-$ $|\Xi_b' {}^4P_{5/2}\rangle$ state. Moreover, due to the theoretical uncertainties and lack of the branching ratio information, the possibilities of the $\Xi_b(6227)$ as other P wave Ξ_b' states cannot be excluded.

In this work, we study the strong decays of the λ -mode P -wave Ξ_b' state in the j - j coupling scheme, and our results are listed in Table IV.

1. $J^P = 1/2^-$ states

It is found that the $J^P = 1/2^-$ state $|J^P = \frac{1^-}{2}, 0\rangle$ has a width of 51 MeV, this state mainly decays into $\Lambda_b K$ and $\Xi_b\pi$ channels. The decay width of $|J^P = \frac{1^-}{2}, 0\rangle$ is about a factor 3 larger than that of $\Xi_b(6227)$. While the other $J^P = 1/2^-$ state $|J^P = \frac{1^-}{2}, 1\rangle$ has a width of 15 MeV, and dominantly decays into $\Xi_b^*\pi$ channel. Although the decay width of $|J^P = \frac{1^-}{2}, 1\rangle$




FIG. 2: Partial and total strong decay widths of the $J^P = 1/2^-$ and $J^P = 3/2^-$ states with $M = 6227$ MeV in the Ξ_b' families as a function of mixing angle ϕ . The solid curves stand for the total widths.

is close to that of $\Xi_b(6227)$, the decay modes are inconsistent with the observations. For a comparison, in Table IV we also list the strong decay properties of the $J^P = 1/2^-$ states $|{}^2P_{1/2}\rangle$ and $|{}^4P_{1/2}\rangle$ which are calculated within the L - S coupling scheme in Ref. [16]. The predictions show obvious differences between the j - j and L - S coupling schemes (see Table IV).

Considering the mixing angle between $|{}^2P_{1/2}\rangle$ and $|{}^4P_{1/2}\rangle$ for the physical states with $J^P = 1/2^-$ may have some deviations from the j - j coupling scheme, the strong decay widths $|\Xi_b' P_{1/2}^1\rangle_1$ and $|\Xi_b' P_{1/2}^1\rangle_2$ as a function of mixing angle ϕ are shown in Fig. 2. It is found that the decay properties of $|\Xi_b' P_{1/2}^1\rangle_2$ are less sensitive to the mixing angle ϕ . However, the decay properties of $|\Xi_b' P_{1/2}^1\rangle_1$ shows some sensitivities to the mixing angle. For example, if one adopts the mixing angle $\phi \simeq 35^\circ$ in the heavy quark limit the $\Lambda_b K$ mode for $|\Xi_b' P_{1/2}^1\rangle_1$ is forbidden, while if $\phi \simeq 15^\circ$ the decay rate into $\Lambda_b K$ is sizable. From the decay properties shown in Fig. 2, one can find that the $\Xi_b'(6227)$ do not favor the mixed states $|\Xi_b' P_{1/2}^1\rangle_1$ and $|\Xi_b' P_{1/2}^1\rangle_2$.

2. $J^P = 3/2^-$ states

For the $J^P = 3/2^-$ state $|J^P = \frac{3^-}{2}, 1\rangle$ in the Ξ_b' family, it has a width of ~ 14 MeV, and dominantly decays into $\Xi_b'(5945)\pi$, the decay mode is inconsistent with the observations of $\Xi_b'(6227)$. While the other $J^P = 3/2^-$ state $|J^P = \frac{3^-}{2}, 2\rangle$ has a width of ~ 24 MeV, and mainly decays into $\Lambda_b K$

TABLE IV: Partial widths (MeV) and branching fractions for the strong decays of the $1P$ -wave states in the j - j coupling scheme compared with that in the L - S coupling scheme taken from Ref. [16] in the Ξ'_b family. The masses of the P -wave bottom baryons are adopted from the quark model predictions in Ref. [4].

State	$ J, j\rangle$	Channel	Γ_i (MeV)	\mathcal{B}_i	$ ^{2S+1}L_J\rangle$	Channel	Γ_i (MeV) [16]	\mathcal{B}_i [16]
$\Xi'_b(6233)$	$ J = \frac{1}{2}^-, 1\rangle$	$\Lambda_b K$	$ ^2P_{1/2}\rangle$	$\Lambda_b K$	12.11	44.77%
		$\Xi_b \pi$		$\Xi_b \pi$	4.77	17.63%
		$\Xi'_b \pi$	13.85	90.76%		$\Xi'_b \pi$	9.23	34.12%
		$\Xi'_b(5945)\pi$	1.41	9.24%		$\Xi'_b(5945)\pi$	0.94	3.48%
		total	15.26			total	27.05	
$\Xi'_b(6227)$	$ J = \frac{1}{2}^-, 0\rangle$	$\Lambda_b K$	38.07	71.71%	$ ^4P_{1/2}\rangle$	$\Lambda_b K$	17.28	53.60%
		$\Xi_b \pi$	15.02	28.29%		$\Xi_b \pi$	10.01	31.05%
		$\Xi'_b \pi$		$\Xi'_b \pi$	4.54	14.08%
		$\Xi'_b(5945)\pi$		$\Xi'_b(5945)\pi$	0.41	1.27%
		total	53.09			total	32.24	
$\Xi'_b(6234)$	$ J = \frac{3}{2}^-, 2\rangle$	$\Lambda_b K$	7.31	26.02%	$ ^2P_{3/2}\rangle$	$\Lambda_b K$	4.14	17.14%
		$\Xi_b \pi$	17.89	63.69%		$\Xi_b \pi$	14.91	61.74%
		$\Xi'_b \pi$	1.60	5.70%		$\Xi'_b \pi$	2.37	9.81%
		$\Xi'_b(5945)\pi$	1.29	4.59%		$\Xi'_b(5945)\pi$	2.73	11.30%
		total	28.09			total	24.15	
$\Xi'_b(6224)$	$ J = \frac{3}{2}^-, 1\rangle$	$\Lambda_b K$	$ ^4P_{3/2}\rangle$	$\Lambda_b K$	0.98	6.19%
		$\Xi_b \pi$		$\Xi_b \pi$	2.67	16.87%
		$\Xi'_b \pi$	0.72	5.02%		$\Xi'_b \pi$	0.10	0.63%
		$\Xi'_b(5945)\pi$	13.61	94.98%		$\Xi'_b(5945)\pi$	12.08	76.31%
		total	14.33			total	15.83	
$\Xi'_b(6226)$	$ J = \frac{5}{2}^-, 2\rangle$	$\Lambda_b K$	4.20	17.22%	$ ^4P_{5/2}\rangle$	$\Lambda_b K$	4.20	17.22%
		$\Xi_b \pi$	16.37	67.12%		$\Xi_b \pi$	16.37	67.12%
		$\Xi'_b \pi$	0.60	2.46%		$\Xi'_b \pi$	0.60	2.46%
		$\Xi'_b(5945)\pi$	3.22	13.20%		$\Xi'_b(5945)\pi$	3.22	13.20%
		total	24.39			total	24.39	

and $\Xi_b \pi$ final states, and the partial width ratio between these two channels is predicted to be

$$\frac{\Gamma[\Lambda_b K]}{\Gamma[\Xi_b \pi]} \simeq 0.36. \quad (20)$$

This ratio predicted by us is obviously smaller than the 3P_0 model prediction $\frac{\Gamma[\Lambda_b K]}{\Gamma[\Xi_b \pi]} \simeq 0.89$ in Ref. [14]. Both the decay modes and width of $|J^P = \frac{3}{2}^-, 2\rangle$ are consistent with the observations of $\Xi_b(6227)$. It should be mentioned that the strong decay properties of the $J^P = 3/2^-$ Ξ'_b states $|J^P = \frac{3}{2}^-, 1\rangle$ and $|J^P = \frac{3}{2}^-, 2\rangle$ in the j - j coupling scheme are similar to those of $|^4P_{3/2}\rangle$ and $|^2P_{3/2}\rangle$ in the L - S coupling scheme, respectively (see Table. IV).

The mixing angle of the physical states with $J^P = 3/2^-$ may lie between the L - S and j - j coupling limit, we show the strong decay widths as a function of mixing angle ϕ from $0^\circ \sim 24^\circ$ in Fig. 2. It can be found that the deviation from the j - j coupling mixing has small influence on the strong decay behaviors, and our conclusions remain. It should be mentioned that the recent QCD sum rule analysis also suggested that $\Xi_b(6227)$ may be a $J^P = 3/2^-$ state [34].

3. $J^P = 5/2^-$ state

For the $|J^P = \frac{5}{2}^-, 2\rangle$ state, the L - S coupling scheme is equal to the j - j coupling scheme. The results indicate that the $\Xi'_b(6227)$ can be assigned as the $|J^P = \frac{5}{2}^-, 2\rangle$ state as well. The partial width ratio between $\Lambda_b K$ and $\Xi_b \pi$ is predicted to be

$$\frac{\Gamma[\Lambda_b K]}{\Gamma[\Xi_b \pi]} \simeq 0.25, \quad (21)$$

which is slightly smaller than that for $|J^P = \frac{3}{2}^-, 2\rangle$. The ratio predicted by us is obviously smaller than the 3P_0 model prediction $\frac{\Gamma[\Lambda_b K]}{\Gamma[\Xi_b \pi]} \simeq 0.94$ in Ref. [14].

As a whole, $\Xi'_b(6227)$ seems to favor the $j = 2$ states with $J^P = 3/2^-$ or $J^P = 5/2^-$. Our conclusion is also consistent with the calculations of the 3P_0 model and QCD sum rule [14, 35]. To further understand the nature of $\Xi'_b(6227)$, more measurements, such as the ratio $\Gamma[\Lambda_b K]/\Gamma[\Xi_b \pi]$, are suggested to be carried out in future experiments.

TABLE V: Partial widths (MeV) and branching fractions for the strong decays of the $1P$ -wave states in the j - j coupling scheme compared with that in the L - S coupling scheme taken from Ref. [16] in the Ω_b family. The masses of the P -wave bottom baryons are adopted from the quark model predictions in Ref. [4].

State [4]	$ J^P, j\rangle$	$\Gamma[\Xi_b K]$ (MeV)	$ ^{2S+1}L_J\rangle$	$\Gamma[\Xi_b K]$ (MeV) [16]
$\Omega_b(6339)$	$ J = \frac{1}{2}^-, 1\rangle$...	$ ^2P_{1/2}\rangle$	49
$\Omega_b(6330)$	$ J = \frac{1}{2}^-, 0\rangle$	143	$ ^4P_{1/2}\rangle$	95
$\Omega_b(6340)$	$ J = \frac{3}{2}^-, 2\rangle$	2.2	$ ^2P_{3/2}\rangle$	1.8
$\Omega_b(6331)$	$ J = \frac{3}{2}^-, 1\rangle$...	$ ^4P_{3/2}\rangle$	0.22
$\Omega_b(6334)$	$ J = \frac{5}{2}^-, 2\rangle$	1.6	$ ^4P_{5/2}\rangle$	1.6




FIG. 3: Partial decay widths of the $\Xi_b K$ channel for the $J^P = 1/2^-$ and $J^P = 3/2^-$ Ω_b states as a function of mixing angle ϕ . The resonance mass is taken the quark model prediction $M = 6330$ MeV.

C. The P -wave Ω_b states

For the $1P$ -wave Ω_b states, there are no signals from experiments. By taking the predicted masses from RQM [4], their strong decays in j - j coupling scheme are calculated and shown in Table V. For the $J^P = 1/2^-$ state $|J^P = \frac{1}{2}^-, 0\rangle$, a large decay width of 148 MeV is obtained, its decays are governed by the $\Xi_b K$ mode. The decay mode $\Xi_b K$ for the $j = 1$ states $|J^P = \frac{1}{2}^-, 1\rangle$ and $|J^P = \frac{3}{2}^-, 1\rangle$ are forbidden, thus, these states should be very narrow states. The two $j = 2$ states $|J^P = \frac{3}{2}^-, 2\rangle$ and $|J^P = \frac{5}{2}^-, 2\rangle$ are predicted to be very narrow states with a width of $\sim 1 - 2$ MeV, which might be found in future experiments. The strong decay properties of the $1P$ -wave Ω_b states were also studied in Ref. [15, 34]. Our main results are consistent with the predictions in Ref. [15]. For a comparison, the results calculated from the L - S coupling scheme [16] are also listed in Table. V. It is found that the strong decay properties of $|J = \frac{1}{2}^-, 0\rangle$ and $|J = \frac{3}{2}^-, 2\rangle$ are similar to those of $|^4P_{1/2}\rangle$ and $|^2P_{3/2}\rangle$ in the L - S coupling scheme, respectively. However, the decay properties of $|J = \frac{1}{2}^-, 1\rangle$ and $|J = \frac{3}{2}^-, 1\rangle$ are very different from those in the L - S coupling scheme.

The physical states with $J^P = 1/2^-$ as mixed states between $|^2P_{1/2}\rangle$ and $|^4P_{1/2}\rangle$, the mixing angle may not be the ideal angle $\phi = 35^\circ$ predicted in the heavy quark limit. Taking the

mixing scheme as defined in Eq. (14), the strong decay widths of the $J^P = 1/2^-$ states as a function of the mixing angle ϕ are presented in Fig. 3. When the mixing angle lies in $0^\circ \sim 35^\circ$, the width of $|\Omega_b P_{\frac{1}{2}}^-\rangle_1$ is relatively small, while $|\Omega_b P_{\frac{1}{2}}^-\rangle_2$ is a broad state with a width of $\sim 120 \pm 20$ MeV. For the $J^P = 3/2^-$ states, when the mixing angle varies from 0° to 24° , the $|\Omega_b P_{\frac{3}{2}}^-\rangle_1$ state is a narrow state with a width of ~ 1 MeV, while the partial width of $|\Omega_b P_{\frac{3}{2}}^-\rangle_2$ into $\Xi_b K$ channel may be less than 0.2 MeV. The $J^P = 3/2^-$ states might be found in the $\Xi_b K$ channel in future experiments.

V. SUMMARY

In the j - j coupling scheme, the strong decays of the low-lying P -wave singly bottom heavy baryons belonging to $\mathbf{6}_F$ are studied within the chiral quark model. Our results show that the newly observed resonances $\Sigma_b(6097)^\pm$ can be assigned as the light spin $j = 2$ states with spin-parity numbers $J^P = 3/2^-$ or $J^P = 5/2^-$. If $\Sigma_b(6097)$ corresponds to $|J^P = \frac{3}{2}^-, 2\rangle$, it might be observed in the $\Sigma_b \pi$ channel, while if $\Sigma_b(6097)$ is the $J^P = 5/2^-$ state, it might be observed in the $\Sigma_b^* \pi$ channel.

The newly observed resonance $\Xi_b(6227)^-$ favors the light spin $j = 2$ states with spin-parity numbers $J^P = 3/2^-$ or $J^P = 5/2^-$ in the Ξ_b' family. $\Xi_b(6227)^-$ may be the strange partner of $\Sigma_b(6097)^\pm$. The $J^P = 3/2^-$ state $|J^P = \frac{3}{2}^-, 2\rangle$ and the $J^P = 5/2^-$ state $|J^P = \frac{5}{2}^-, 2\rangle$ in the Ξ_b' family have similar strong decay properties. In order to identify them, angular distributions of their decays in either strong decay modes or radiative transitions should be needed [16].

Considering the heavy quark symmetry, the $J^P = 1/2^-$ P -wave Ω_b state $|J^P = \frac{1}{2}^-, 0\rangle$ might be a broad state with a width of ~ 150 MeV. The two $j = 2$ states $|J^P = \frac{3}{2}^-, 2\rangle$ and $|J^P = \frac{5}{2}^-, 2\rangle$ are predicted to be very narrow states with a width of $\sim 1 - 2$ MeV, which might be found in future experiments. The $\Xi_b K$ decay mode for the $j = 1$ states $|J^P = \frac{1}{2}^-, 1\rangle$ and $|J^P = \frac{3}{2}^-, 1\rangle$ are forbidden in the heavy quark limit, thus these two states should be very narrow states. To looking for the missing P -wave Ω_b states, the $\Xi_b K$ decay mode is worth to observing.

Acknowledgments

No.U1832173, and No. 11705056.

This work is supported, in part, by the National Natural Science Foundation of China under Grants No.11775078,

-
- [1] R. Aaij *et al.* [LHCb Collaboration], Observation of a new Ξ_b^- resonance, Phys. Rev. Lett. **121**, 072002 (2018).
- [2] R. Aaij *et al.* [LHCb Collaboration], Observation of two resonances in the $\Lambda_b^0\pi^\pm$ systems and precise measurement of Σ_b^\pm and $\Sigma_b^{*\pm}$ properties, arXiv:1809.07752 [hep-ex].
- [3] S. Capstick and W. Roberts, Quark models of baryon masses and decays, Prog. Part. Nucl. Phys. **45**, S241 (2000).
- [4] D. Ebert, R. N. Faustov and V. O. Galkin, Spectroscopy and Regge trajectories of heavy baryons in the relativistic quark-diquark picture, Phys. Rev. D **84**, 014025 (2011).
- [5] D. Ebert, R. N. Faustov and V. O. Galkin, Masses of excited heavy baryons in the relativistic quark model, Phys. Lett. B **659**, 612 (2008).
- [6] D. Ebert, R. N. Faustov and V. O. Galkin, Masses of heavy baryons in the relativistic quark model, Phys. Rev. D **72**, 034026 (2005).
- [7] T. Yoshida, E. Hiyama, A. Hosaka, M. Oka and K. Sadato, Spectrum of heavy baryons in the quark model, Phys. Rev. D **92**, 114029 (2015).
- [8] W. Roberts and M. Pervin, Heavy baryons in a quark model, Int. J. Mod. Phys. A **23**, 2817 (2008).
- [9] A. Valcarce, H. Garcilazo and J. Vijande, Towards an understanding of heavy baryon spectroscopy, Eur. Phys. J. A **37**, 217 (2008).
- [10] M. Karliner, B. Keren-Zur, H. J. Lipkin and J. L. Rosner, The Quark Model and b Baryons, Annals Phys. **324**, 2 (2009).
- [11] B. Chen, K. W. Wei and A. Zhang, Assignments of Λ_Q and Ξ_Q baryons in the heavy quark-light diquark picture, Eur. Phys. J. A **51**, 82 (2015).
- [12] Q. Mao, H. X. Chen, W. Chen, A. Hosaka, X. Liu and S. L. Zhu, QCD sum rule calculation for P-wave bottom baryons, Phys. Rev. D **92**, 114007 (2015).
- [13] Z. G. Wang, Analysis of the $1/2^-$ and $3/2^-$ heavy and doubly heavy baryon states with QCD sum rules, Eur. Phys. J. A **47**, 81 (2011).
- [14] B. Chen, K. W. Wei, X. Liu and A. Zhang, Role of newly discovered $\Xi_b(6227)^-$ for constructing excited bottom baryon family, Phys. Rev. D **98**, 031502 (2018).
- [15] B. Chen and X. Liu, Assigning the newly reported $\Sigma_b(6097)$ as a P -wave excited state and predicting its partners, arXiv:1810.00389 [hep-ph].
- [16] K. L. Wang, Y. X. Yao, X. H. Zhong and Q. Zhao, Strong and radiative decays of the low-lying S - and P -wave singly heavy baryons, Phys. Rev. D **96**, 116016 (2017).
- [17] R. Aaij *et al.* [LHCb Collaboration], Observation of five new narrow Ω_c^0 states decaying to $\Xi_c^+K^-$, Phys. Rev. Lett. **118**, 182001 (2017).
- [18] K. L. Wang, L. Y. Xiao, X. H. Zhong and Q. Zhao, Understanding the newly observed Ω_c states through their decays, Phys. Rev. D **95**, 116010 (2017).
- [19] H. Y. Cheng, Charmed baryons circa 2015, Front. Phys. **10**, 101406 (2015).
- [20] A. Manohar and H. Georgi, Chiral quarks and the nonrelativistic quark Model, Nucl. Phys. B **234**, 189 (1984).
- [21] X. H. Zhong and Q. Zhao, Strong decays of newly observed D_{sJ} states in a constituent quark model with effective Lagrangians, Phys. Rev. D **81**, 014031 (2010).
- [22] X. H. Zhong and Q. Zhao, Strong decays of heavy-light mesons in a chiral quark model, Phys. Rev. D **78**, 014029 (2008).
- [23] X. H. Zhong, Strong decays of the newly observed $D(2550)$, $D(2600)$, $D(2750)$, and $D(2760)$, Phys. Rev. D **82**, 114014 (2010).
- [24] L. Y. Xiao and X. H. Zhong, Strong decays of higher excited heavy-light mesons in a chiral quark model, Phys. Rev. D **90**, 074029 (2014).
- [25] X. H. Zhong and Q. Zhao, Charmed baryon strong decays in a chiral quark model, Phys. Rev. D **77**, 074008 (2008).
- [26] L. H. Liu, L. Y. Xiao and X. H. Zhong, Charm-strange baryon strong decays in a chiral quark model, Phys. Rev. D **86**, 034024 (2012).
- [27] L. Y. Xiao and X. H. Zhong, Ξ baryon strong decays in a chiral quark model, Phys. Rev. D **87**, 094002 (2013).
- [28] L. Y. Xiao, K. L. Wang, Q. F. Lü, X. H. Zhong and S. L. Zhu, Strong and radiative decays of the doubly charmed baryons, Phys. Rev. D **96**, 094005 (2017).
- [29] H. Nagahiro, S. Yasui, A. Hosaka, M. Oka and H. Noumi, Structure of charmed baryons studied by pionic decays, Phys. Rev. D **95**, 014023 (2017).
- [30] Y. X. Yao, K. L. Wang and X. H. Zhong, Strong and radiative decays of the low-lying D -wave singly heavy baryons, arXiv:1803.00364 [hep-ph].
- [31] R. Koniuk and N. Isgur, Baryon Decays in a Quark Model with Chromodynamics, Phys. Rev. D **21**, 1868 (1980) Erratum: [Phys. Rev. D **23**, 818 (1981)].
- [32] S. Godfrey and N. Isgur, Mesons in a Relativized Quark Model with Chromodynamics, Phys. Rev. D **32**, 189 (1985).
- [33] M. Tanabashi *et al.* [Particle Data Group], Review of Particle Physics, Phys. Rev. D **98**, 030001 (2018).
- [34] S. S. Agaev, K. Azizi and H. Sundu, Decay widths of the excited Ω_b baryons, Phys. Rev. D **96**, 094011 (2017).
- [35] T. M. Aliev, K. Azizi, Y. Sarac and H. Sundu, Structure of the $\Xi_b(6227)^-$ Resonance, arXiv:1808.08032 [hep-ph].