

Recursion schemes, discrete differential equations and characterization of polynomial time computation

Olivier Bournez ^{*1}, Arnaud Durand ^{†2}, and Sabrina Ouazzani ^{‡3}

¹*Ecole Polytechnique, LIX, 91128 Palaiseau Cedex, France*

²*Université Paris Diderot, IMJ-PRG, CNRS UMR 7586, Case
7012, 75205 Paris cedex 13, France*

³*LIX, 91128 Palaiseau Cedex, France and LACL, Université
Paris-Est Créteil, 61 avenue du Général de Gaulle, 94010 Créteil,
France*

October 5, 2018

Abstract

This paper studies the expressive and computational power of discrete Ordinary Differential Equations (ODEs). It presents a new framework using discrete ODEs as a central tool for computation and provides several implicit characterizations of complexity and computability classes.

The proposed framework presents an original point of view on complexity and computability classes. It also unifies in an elegant setting various constructions that have been proposed for characterizing these classes. This includes Cobham's and Bellantoni and Cook's definition of polynomial time and later extensions on the approach, as well as recent characterizations of computability and complexity by classes of ordinary differential equations. It also helps understanding the relationships between analog computations and classical discrete models of computation theory.

At a more technical point of view, this paper points out the fundamental role of linear (discrete) ordinary differential equations and classical ODE tools such as changes of variables to capture computability and complexity measures, or as a tool for programming various algorithms.

^{*}Email: bournez@lix.polytechnique.fr. Supported by RACAF Project from Agence Nationale de la Recherche and Labex Digicosme Project ACDC.

[†]Email: durand@math.univ-paris-diderot.fr

[‡]Email: sabrina@lix.polytechnique.fr. Supported by Labex Digicosme Project ACDC.

Contents

1	Introduction	3
1.1	Related works	4
1.1.1	Analog computations:	4
1.2	Classical complexity theory:	4
1.3	Structure of the paper	5
2	Discrete differentiability	5
2.1	Discrete derivation, integration and exponentiation	5
2.2	Discrete Ordinary Differential Equations (ODE)	7
3	Programming with discrete ODE	8
4	Bounded schemes in computation theory	11
4.1	Computability theory and bounded schemes	11
4.2	Complexity theory and bounded schemes	13
5	Computability and Discrete ODEs	15
5.1	About positive and negative integers and encodings	15
5.2	Recursive and subrecursive classes of functions	15
5.3	Subrecursive functions and discrete ODEs	16
5.4	Elementary functions, Grzegorzcyk hierarchy and linear discrete ODEs	16
5.5	Computability and discrete ODEs	19
5.5.1	By adding a minimization operator	19
5.5.2	By programming minimization	19
6	Restricted recursion and integration schemes	20
6.1	General theory	22
6.2	Fundamental alternative view	24
6.3	Length-ODEs	25
6.4	Linear length-ODEs	26
7	A characterization of polynomial time	28
7.1	Register machines	28
7.2	A characterization of polynomial time	30
8	Further works	32
A	Discrete Calculus	37
A.1	Some basic statements	37
A.2	The discrete integral and primitive	38
A.3	Integration by part	40
A.4	Derivative of a composition	40
A.5	Falling power	40
A.6	Exponential	41

A.7	Falling Exponential	41
A.8	Solving some particular ODEs	41
A.9	Solving affine ODEs	42
A.10	Derivative of some particular functions	45
B	Review of some results and models	46
B.1	Random access machines	46

1 Introduction

Since the beginning of its foundations, classification of problems, by various models of computation, either by their complexity or by their computability properties, is a thriving field of computer science. Nowadays, classical (digital) computer science problems also deal with continuous data coming from different areas and modeling involves the use of tools like numerical analysis, probability theory or differential equations. Thus new characterizations related to these fields have been proposed. On a dual way, the quest for new type of computers recently led to revisit the power of some models for analog machines based on differential equations, and to compare them to modern digital models. In both contexts, when discussing the related issues, one has to overcome the fact that today's (digital) computers are by essence discrete machines while the objects under study are continuous and naturally correspond to Ordinary Differential Equations (ODEs).

We consider here an original approach in between the two worlds: discrete oriented computation with differential equations.

Indeed, ODE appear to be a natural way of expressing properties and are intensively used, in particular in applied science. The theory of classical (continuous) ODEs is rather very well understood, broadly studied and taught with a plethora literature, see e.g. [1, 4, 13]. We are here interested here in a discrete more recent counterpart of classical continuous ODEs: discrete ODEs, whose theory is sometimes called *discrete calculus*, or *finite calculus*. See for e.g. [19, 18, 22, 26] or our brief presentation in Appendix A.

In this article, we prove that various classes of complexity and computability theory can be very elegantly and simply defined using discrete ordinary differential equations. We also demonstrate through this discussion how some techniques from analog world such as changes of variables can be used to solve efficiently some (classical, digital) problems.

As far as we know, this is the first time that computations with discrete ODEs and their related complexity aspects have been considered. By contrast, characterizations have been recently obtained with classical (continuous) ODEs of various classes of functions, mostly in the framework of computable analysis. The hardness of solving continuous ODEs has been intensively discussed: for example [25] establishes some basis of the complexity aspects of ODEs and more recent work like [24] or [14] establishes links between complexity or effective aspects of such differential equations. Hence, the computational power of

continuous ODEs is much more understood.

We believe that investigating the expressive power of discrete ODE, can really help to better understand complexity of computation for both the discrete and continuous settings. Indeed, on one hand, as a consequence, our work relates classical (discrete) complexity classes to analog computations, i.e. computations over the reals, as analog computation have been related in various ways to continuous ordinary differential equations, and as discrete ordinary differential equations provide clear hints about their continuous counterparts. But on the other hand, it also opens a new perspective on classical discrete computations, i.e. computation that deals with bits, words, or integers. In this discrete setting, our work falls under the scope of so-called implicit complexity, i.e. characterization complexity measures in a machine independent way. Combining these two approaches, it helps to clearly point out which aspects of the statements are related to continuous computations versus discrete computations.

This original work point out the fundamental role of linear (discrete) ordinary differential equations in computability and complexity theory: when considered in general, this provides a characterization of elementary functions. When considered with suitable (length related) changes of variables, this provides a characterization of polynomial time. This work also opens a way to revisit seminal results such as Cobham's [12] and Bellantoni and Cook's [2] definition of polynomial time as syntactic constraints imposing only linear discrete ODEs.

1.1 Related works

1.1.1 Analog computations:

In the context of analog computations there have been several results relating classical complexity to various classes of continuous ODEs. In particular, a serie of papers has been devoted to study various classes of the so-called \mathbb{R} -recursive functions, after their introduction in [31]. At the complexity level, characterizations of complexity classes such as **PTIME** and **NPTIME** using \mathbb{R} -recursive algebra have been obtained [33], motivated in particular by the idea of transferring classical questions from complexity theory to the context of real and complex analysis [30, 33, 32]. More recently, it has been proved that polynomial differential equations can be considered as a very simple and elegant model in which computable functions over the reals and polynomial time computable functions over the reals can be defined without any reference to concepts from discrete computation theory [5, 35]. Refer to [7, 6] for an up to date survey about various works on analog computations, in particular in a computation theory perspective.

1.2 Classical complexity theory:

Implicit complexity has been developed in many ways to provide machine independent characterizations of various computability and complexity classes in the discrete setting. This includes characterizations of complexity classes based

on lambda calculus (e.g. [28]), finite model theory and descriptive complexity (e.g. [16]), on function algebra (e.g. [12, 2, 29, 36]), or yet one combining the latter two approaches (e.g. [20, 37]). This approach has also been proved useful to measure the expressive power of various formalisms with wide applications in database and constraint theory and programming languages (see [10, 15, 21] for more complete references).

1.3 Structure of the paper

In Section 2 a short introduction to discrete differentiability is given followed in Section 3 by an illustration, through examples, of the programming ability of discrete ODE. First formal definitions of discrete ODE are given in Section 5 together with characterizations of primitive recursion and elementary functions in this context. Section 6 introduces the notion (and basic theory) of length-ODE which is central in the characterization of **FPTIME** (Section 7). Section 8 discusses some extensions of the results. Due to lack of space, an important Appendix is given.

2 Discrete differentiability

In this section, we review some basic notions of discrete calculus to help intuition in the rest of the paper (see Appendix A for a more complete review).

2.1 Discrete derivation, integration and exponentiation

Discrete derivatives are usually intended to concern functions over the integers of type $\mathbf{f} : \mathbb{N}^p \rightarrow \mathbb{Z}^d$, but the statements and concepts considered in our discussions are also valid more generally for functions of type $\mathbf{f} : \mathbb{Z}^p \rightarrow \mathbb{Z}^d$, for some integers p, d , or even functions $\mathbf{f} : \mathbb{R}^p \rightarrow \mathbb{R}^d$. The basic idea is to consider the following concept of derivative.

Definition 1 (Discrete Derivative) *The discrete derivative of $\mathbf{f}(x)$ is defined as $\Delta\mathbf{f}(x) = \mathbf{f}(x+1) - \mathbf{f}(x)$. We will also write \mathbf{f}' for $\Delta\mathbf{f}(x)$ to help to understand statements with respect to their classical continuous counterparts.*

Several results from classical derivatives generalize to these settings: this includes linearity of derivation $(a \cdot f(x) + b \cdot g(x))' = a \cdot f'(x) + b \cdot g'(x)$, formulas for products and division such as $(f(x) \cdot g(x))' = f'(x) \cdot g(x+1) + f(x) \cdot g'(x)$. A fundamental concept is the following:

Definition 2 (Discrete Integral) *Given some function $\mathbf{f}(x)$, we write $\int_a^b \mathbf{f}(x)\delta x$ as a synonym for*

$$\int_a^b \mathbf{f}(x)\delta x = \sum_{x=a}^{x=b-1} \mathbf{f}(x)$$

with the convention that it values 0 when $a = b$ and $\int_a^b \mathbf{f}(x)\delta x = -\int_b^a \mathbf{f}(x)\delta x$ when $a > b$.

The telescope formula yields the so-called Fundamental Theorem of Finite Calculus:

Theorem 1 (Fundamental Theorem of Finite Calculus) *Let $\mathbf{F}(x)$ be some function. Then,*

$$\int_a^b \mathbf{F}'(x)\delta x = \mathbf{F}(b) - \mathbf{F}(a).$$

As for classical functions, any function has a set of primitives defined up to some additive constant, and techniques such as integration by part can be used.

Lemma 1 (Derivation of an integral with parameters) *Consider*

$$\mathbf{F}(x) = \int_{a(x)}^{b(x)} \mathbf{f}(x, t)\delta t.$$

Then

$$\mathbf{F}'(x) = \int_{a(x)}^{b(x)} \frac{\partial \mathbf{f}}{\partial x}(x, t)\delta t + \int_0^{-a'(x)} \mathbf{f}(x+1, a(x+1)+t)\delta t + \int_0^{b'(x)} \mathbf{f}(x+1, b(x)+t)\delta t$$

In particular, when $a(x) = a$ and $b(x) = b$ are constant functions,

$$\mathbf{F}'(x) = \int_a^b \frac{\partial \mathbf{f}}{\partial x}(x, t)\delta t$$

A classical concept in discrete calculus is the one of falling power defined as $x^{\underline{m}} = x \cdot (x-1) \cdot (x-2) \cdots (x-(m-1))$. This notion is motivated by a derivative formula similar to the classical one for powers in the continuous setting. In a similar spirit, we introduce the following concept. This seems not standard (as far as the authors know) but of clear interest.

Definition 3 (Falling exponential) *Given some function $\mathbf{U}(x)$, the expression \mathbf{U} to the falling exponential x , denoted by $\overline{2}^{\mathbf{U}(x)}$, stands for*

$$\overline{2}^{\mathbf{U}(x)} = (1 + \mathbf{U}'(x-1)) \cdots (1 + \mathbf{U}'(1)) \cdot (1 + \mathbf{U}'(0)) = \prod_{t=0}^{t=x-1} (1 + \mathbf{U}'(t)).$$

with the convention that $\prod_0^0 = \mathbf{id}$, where \mathbf{id} is the identity (sometimes denoted 1 hereafter)

This is motivated by the remark that for all $x \in \mathbb{Z}$, $2^x = \overline{2}^x$, and:

Theorem 2 (Derivative of a falling exponential) *The discrete derivative of a falling exponential is given by*

$$\left(\overline{2}^{\mathbf{U}(x)}\right)' = \mathbf{U}'(x) \cdot \overline{2}^{\mathbf{U}(x)}.$$

2.2 Discrete Ordinary Differential Equations (ODE)

We will focus in this article on discrete Ordinary Differential Equations (ODE) on functions with several variables, that is to say for example on equations of the (possibly vectorial) form:

$$\frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial x} = \mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}). \quad (1)$$

As expected $\frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial x}$ stands for the derivative of functions $f(x, \mathbf{y})$ considered as a function of x , when \mathbf{y} is fixed. When some initial value is given, this is called an *Initial Value Problem (IVP)*, also called a *Cauchy Problem*. That is to say, we are given a problem of type:

$$\begin{aligned} \frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial x} &= \mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) \\ \mathbf{f}(0, \mathbf{y}) &= \mathbf{g}(\mathbf{y}) \end{aligned} \quad (2)$$

with functions \mathbf{g}, \mathbf{h} of suitable dimensions and domains: Our aim here is to discuss total functions whose domain and range is either of the form $\mathcal{D} = \mathbb{N}$, \mathbb{Z} , or possibly a finite product $\mathcal{D} = \mathcal{D}_1 \times \cdots \times \mathcal{D}_k$ where each $\mathcal{D}_i = \mathbb{N}, \mathbb{Z}$. By considering that $\mathbb{N} \subset \mathbb{Z}$, we assume that the range is always \mathbb{Z}^d for some d . The concept of solution for such ODEs is as expected: assume $h : \mathbb{Z}^d \times \mathbb{N} \times \mathbb{Z}^p \rightarrow \mathbb{Z}$ (or $h : \mathbb{Z}^d \times \mathbb{Z} \times \mathbb{Z}^p \rightarrow \mathbb{Z}$), a solution over \mathcal{D} is a function $f : \mathcal{D} \times \mathbb{Z}^p \rightarrow \mathbb{Z}^d$ that satisfies the equations for all x, \mathbf{y} .

We will only consider well-defined ODEs such as above in this article (but variants with partially defined function could be considered as well). Observe that an IVP of the form (1) always admits a (necessarily unique) solution over \mathbb{N} since f can be defined inductively with $\mathbf{f}(0, \mathbf{y}) = \mathbf{g}(\mathbf{y})$ and $\mathbf{f}(x + 1, \mathbf{y}) = \mathbf{f}(x, \mathbf{y}) + \mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y})$.

Remark 1 Notice that this is not necessarily true over \mathbb{Z} : As an example, consider $f'(x) = -f(x) + 1$, $f(0) = 0$. By definition of $f'(x)$, we must have $f(x + 1) = 1$ for all x , but if $x = -1$, $f(0) = 1 \neq 0$.

Remark 2 (Sign function) It is very instructive to realize that the solution of this IVP over \mathbb{N} is the sign $\mathbf{sg}_{\mathbb{N}}(x)$ function defined by $\mathbf{sg}_{\mathbb{N}}(x) = 1$ if $x > 0$ and $\mathbf{sg}_{\mathbb{N}}(x) = 0$ in the other case.

Affine (also called linear) ordinary differential equations will play a very important role in what follows, i.e. discrete ordinary differential equations of the form $\mathbf{f}'(x) = \mathbf{A}(x) \cdot \mathbf{f}(x) + \mathbf{B}(x)$.

Remark 3 Recall that the solution of $f'(x) = a(x)f(x) + b(x)$ for classical continuous derivatives turns out to be given by (This is usually obtained by the variation of parameter method - see Appendix A for a short review of the method and for other classical ODE):

$$f(x) = f(0)e^{\int_0^x a(t)dt} + \int_0^x b(u)e^{\int_u^x a(t)dt} du. \quad (3)$$

This generalizes to discrete ordinary differential equations, and this works even vectorially:

Lemma 2 (Solution of ODE $\mathbf{f}'(x, \mathbf{y}) = \mathbf{A}(x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(x, \mathbf{y})$) For matrices \mathbf{A} and vectors \mathbf{B} and \mathbf{G} , the solution of equation $\mathbf{f}'(x, \mathbf{y}) = \mathbf{A}(x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(x, \mathbf{y})$ with initial conditions $\mathbf{f}(0, \mathbf{y}) = \mathbf{G}(\mathbf{y})$ is

$$\mathbf{f}(x, \mathbf{y}) = \left(\sqrt{2}^{\int_0^x \mathbf{A}(t, \mathbf{y}) \delta t} \right) \cdot \mathbf{G}(\mathbf{y}) + \int_0^x \left(\sqrt{2}^{\int_{u+1}^x \mathbf{A}(t, \mathbf{y}) \delta t} \right) \cdot \mathbf{B}(u, \mathbf{y}) \delta u. \quad (4)$$

3 Programming with discrete ODE

In this section, we show that several algorithms can actually be naturally solved using discrete ODEs, or viewed as discrete ODEs: Basically, for now, we suppose that composition of functions, constant and the following basic functions can be used freely as functions from \mathbb{Z} to \mathbb{Z} :

- arithmetic operations: $+$, $-$, \times
- $\ell(x)$ returns the length in binary of $x \in \mathbb{N}$.
- $\mathbf{sg}(x) : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $\mathbf{sg}(x) = 1$ if $x > 0$ and $\mathbf{sg}(x) = 0$ in the other case. $\mathbf{sg}_{\mathbb{N}}(x) : \mathbb{N} \rightarrow \mathbb{Z}$ defined by $\mathbf{sg}_{\mathbb{N}}(x) = 1$ if $x > 0$ and $\mathbf{sg}_{\mathbb{N}}(x) = 0$ in the other case.

Recall that $\mathbf{sg}_{\mathbb{N}}(x)$ is the solution over \mathbb{N} of some IVP and hence is very natural in this context. From these basic functions, for readability, one may define useful functions as synonyms:

- $\bar{\mathbf{sg}}(x)$ stands for $\bar{\mathbf{sg}}(x) = (1 - \mathbf{sg}(x)) \times (1 - \mathbf{sg}(-x))$: it tests if $x = 0$ for $x \in \mathbb{Z}$. $\bar{\mathbf{sg}}_{\mathbb{N}}(x)$ stands for $\bar{\mathbf{sg}}_{\mathbb{N}}(x) = 1 - \mathbf{sg}_{\mathbb{N}}(x)$: it tests if $x = 0$ for $x \in \mathbb{N}$.
- $\mathbf{if}(x, y, z)$ stands for $\mathbf{if}(x, y, z) = y + \bar{\mathbf{sg}}(x) \cdot (z - y)$ and $\mathbf{if}_{\mathbb{N}}(x, y, z)$ stands for $\mathbf{if}_{\mathbb{N}}(x, y, z) = y + \bar{\mathbf{sg}}_{\mathbb{N}}(x) \cdot (z - y)$. We have for both versions (The point is that the first considers $x \in \mathbb{Z}$ while the second assumes $x \in \mathbb{N}$):

$$\mathbf{if}(x, y, z) = \begin{cases} y & \text{if } x = 0 \\ z & \text{otherwise} \end{cases} \quad \mathbf{if}_{\mathbb{N}}(x, y, z) = \begin{cases} y & \text{if } x = 0 \\ z & \text{otherwise} \end{cases}$$

- We will extensively use these functions below: $\mathbf{if}(x < x', y, z)$ will be a synonym for $\mathbf{if}(\mathbf{sg}(x - x' - 1), y, z)$. $\mathbf{if}(x = x', y, z)$ will be a synonym for $\mathbf{if}(\bar{\mathbf{sg}}(x - x'), y, z)$.

Example 1 (Computing the minimum of a function:) The minimum of a function $\min f : x \mapsto \min\{f(y) : 0 \leq y \leq x\}$ is given by $F(x, x)$ where F can be computed recursively by

$$\begin{aligned} F(0) &= f(0) \\ F(t+1, x) &= \mathbf{if}(F(t, x) < f(x), F(t, x), f(x)) \end{aligned}$$

This can be interpreted as a discrete ordinary differential equation:

$$\frac{\partial F(t, x)}{\partial t} = H(F(t, x), f(x), t, x) = \begin{cases} 0 & \text{if } F(t, x) < f(x) \\ -F(t, x) + f(x) & \text{if } F(t, x) \geq f(x). \end{cases}$$

The value $\min f(x) = F(x, x)$ can then be computed using the solution of the above discrete ODE. In integral form, we have:

$$F(x, y) = F(0) + \int_0^x H(F(t, y), t, y) \delta t.$$

Remark 4 We also see through this example that such an integral (equivalently discrete ODE) can always be considered as a (recursive) algorithm: compute the integral from its definition as a sum to compute the function. Notice that this algorithm is not polynomial as this basically takes time x to compute $\min f$, i.e. not polynomial with respect to the usual convention for measuring complexity based on the binary length of arguments.

Example 2 (Computing the integer part and divisions, Length-ODE:)

Suppose that we want to compute $\lfloor \sqrt{x} \rfloor = \max\{y \leq x : y \cdot y \leq x\}$ and $\lfloor \frac{x}{y} \rfloor = \max\{z \leq x : z \cdot y = x\}$. It can be done by the following uniform method. Let f, h be some functions with h being non decreasing. We compute some_h with $\text{some}_h(x) = y$ s.t. $|f(x) - h(y)|$ is minimal. When $h(x) = x^2$ and $f(x) = x$, it holds that: $\lfloor \sqrt{x} \rfloor = \text{if}(\text{some}_h(x)^2 \leq x, \text{some}_h(x), \text{some}_h(x) - 1)$.

The function some_h can be computed as a solution of an ODE as in the preceding example. However, there is a more efficient way to do it based on what one usually does with classical ordinary differential equations: performing a change of variable so that it becomes logarithmic in x . Indeed, we write $\text{some}_h(x) = G(\ell(x), x)$ for some function $G(t, x)$ defined by:

$$\begin{aligned} G(0, x) &= x \\ G(t + 1, x) &= \text{if}(h(G(t, x)) = f(x), G(t, x), \\ &\quad \text{if}(h(G(t, x)) > f(x), G(t, x) + 2^{\ell(x)-t-1}, G(t, x) - 2^{\ell(x)-t-1})) \end{aligned}$$

Or, if one prefers, $G(t, x)$ is solution of

$$\frac{\partial G(t, x)}{\partial t} = E(G(t, x), t, x) = \begin{cases} +2^{\ell(x)-t-1} & \text{if } h(G(t, x)) > g(x) \\ 0 & \text{if } h(G(t, x)) = g(x) \\ -2^{\ell(x)-t-1} & \text{if } h(G(t, x)) < g(x) \end{cases}$$

This is indeed a differential equation whose solution is converging fast (in polynomial time) to what we want. Reformulating what we just did, we wrote $\text{some}_h(x) = G(\ell(x), x)$ using the solution of the above discrete ODE, i.e. the solution of $G(T, y) = x + \int_0^T E(G(t, y), t, y) \delta t$. This provides a polynomial time algorithm to solve our problems using a new parameter $t = \ell(x)$ logarithmic in x . Such techniques will be at the heart of the coming results.

Notice that the theory of ODEs also provides very natural alternative ways to compute various quantities. This is very clear when considering numeric functions such as \tan , \sin , etc.

Example 3 (Computing \tan with discrete ODEs, iterative algorithms)

As an illustration, suppose you want to compute $\tan(x_0)$ for say $x_0 = 72$. One way to do it is to observe that $\tan(x)' = (1 + \tan(x)\tan(x+1))$. From fundamental theorem of finite calculus we can hence write:

$$\tan(x_0) = 0 + \int_0^{x_0} \tan(x)\delta x \tag{5}$$

$$= 0 + \tan(1) \cdot \int_0^{x_0} (1 + \tan(x)\tan(x+1))\delta x \tag{6}$$

Inspired from previous remarks, the point is that Equation (5) can be interpreted as an algorithm: it provides a way to compute $\tan(x_0)$ as an integral (or if you prefer as a sum).

Thinking about what means this integral, discrete ODE (19), also encoded by (6), can also be interpreted as $\tan(x+1) - \tan(x) = \tan(1) \cdot [1 + \tan(x)\tan(x+1)]$ that is to say $\tan(x+1) = f(\tan(x))$ where $f(X) = \frac{X + \tan(1)}{1 - \tan(1)X}$. Hence, this is suggesting a way to compute $\tan(72)$ by a method close to express that $\tan(x_0) = f^{[x_0]}(0)$. That is to say Equations (5) and (6) can be interpreted as providing a way to compute $\tan(72)$ using an iterative algorithm: they basically encode some recursive way of computing \tan .

Of course, a similar principle would hold for \sin , or \cos using discrete ODEs obtained above, and for many other functions starting from expression of their derivative.

Remark 5 Given x_0 , (even if we put aside how to deal with involved real quantities) a point is that computing $\tan(x_0)$ using this method can not be considered as polynomial time, as the (usual) convention is that time complexity is measured in term of the length of x_0 , and not on x_0 .

Could we do the same computation faster using a change of variables? This is at the heart of the coming constructions and discussions.

Example 4 (Computing suffixes with discrete ODEs)

Discrete ODEs turns out to be very natural in many other contexts, in particular non numerical ones, where they would probably not be expected. We illustrate the discussion by a way to compute fast (in polynomial time) the suffix function: The suffix function, $\text{suffix}(x, y)$ takes as input two integers x and y and outputs the $\ell(y) = t$ least significant bits of the binary decomposition of x . We describe below a way to compute a suffix working over a parameter t , that is logarithmic in x . Consider the following amazing algorithm that can be interpreted as a fix-point definition of the function: $\text{suffix}(x, y) = F(\ell(x), y)$ where

$$F(T, y) = x + \int_0^T \text{if}(\ell(F(t, x)) = 1, 0, -2^{\ell(F(t, x)) - 1})\delta t.$$

4 Bounded schemes in computation theory

After this teaser, the rest of this article aims at discussing which problems can be solved using discrete ordinary differential equations, and with which complexity. Before doing so, we need to review some basic concepts and results from computation theory that we will be needed in the rest of this article and that have been obtained at this date.

4.1 Computability theory and bounded schemes

Classical recursion theory deals with functions over integers, that is to say with functions $\mathbf{f} : \mathbb{N}^p \rightarrow \mathbb{N}^d$ for some positive integers p, d .

It is well known that all main classes of classical recursion theory can be characterized as closures of a set of basic functions by a finite number of basic rules to build new functions: See e.g. [36, 34, 11]:

Theorem 3 (Total Recursive functions) *A total function over the integers is computable if and only if it belongs to the smallest set of functions that contains constant function $\mathbf{0}$, the projection functions π_i^p , the function successor \mathbf{s} , that is closed under composition, primitive recursion and safe minimization.*

In this statement, $\mathbf{0}$, π_i^p and \mathbf{s} are respectively the functions from, $\mathbb{N} \rightarrow \mathbb{N}$, $\mathbb{N}^p \rightarrow \mathbb{N}$ and $\mathbb{N} \rightarrow \mathbb{N}$ defined as $n \mapsto 0$, $(n_1, \dots, n_p) \mapsto n_i$, and $n \mapsto n + 1$.

We also recall here the basic definitions used in the above statement:

Definition 4 (Primitive recursion) *Given functions $g : \mathbb{N}^p \rightarrow \mathbb{N}$ and $h : \mathbb{N}^{p+2} \rightarrow \mathbb{N}$, function $f = \text{REC}(g, h)$ defined by primitive recursion from g and h is the function $\mathbb{N}^{p+1} \rightarrow \mathbb{N}$ satisfying*

$$\begin{aligned} f(0, \mathbf{y}) &= g(\mathbf{y}) \\ f(x + 1, \mathbf{y}) &= h(f(x, \mathbf{y}), x, \mathbf{y}). \end{aligned}$$

Definition 5 (Primitive recursive functions) *A function over the integers is primitive recursive if and only if it belongs to the smallest set of functions that contains constant function $\mathbf{0}$, the projection functions π_i^p , the functions successor \mathbf{s} , that is closed under composition and primitive recursion.*

Primitive recursive functions have been stratified into various subclasses. We recall here the Grzegorzcyk hierarchy in the rest of this subsection.

Definition 6 (Bounded sum) *Given functions $g(\mathbf{y}) : \mathbb{N}^p \rightarrow \mathbb{N}$,*

- *function $f = \text{BSUM}(g) : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$ is defined by $f : (x, \mathbf{y}) \mapsto \sum_{z \leq x} g(z, \mathbf{y})$.*
- *function $f = \text{BSUM}_{<}(g) : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$ is defined by $f : (x, \mathbf{y}) \mapsto \sum_{z < x} g(z, \mathbf{y})$ for $x \neq 0$, and 0 for $x = 0$.*

Definition 7 (Bounded product) Given functions $g : \mathbb{N}^p \rightarrow \mathbb{N}$,

- function $f = \text{BPROD}(g) : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$ is defined by $f : (x, \mathbf{y}) \mapsto \prod_{z \leq x} g(z, \mathbf{y})$.
- function $f = \text{BPROD}_{<}(g)$ is defined by $f : (x, \mathbf{y}) \mapsto \prod_{z < x} g(z, \mathbf{y})$ for $x \neq 0$, and 1 for $x = 0$.

We have

$$\begin{aligned} \text{BSUM}(g)(x, \mathbf{y}) &= \text{BSUM}_{<}(g)(x, \mathbf{y}) + g(x, \mathbf{y}) \\ \text{BPROD}(g)(x, \mathbf{y}) &= \text{BPROD}_{<}(g)(x, \mathbf{y}) \cdot g(x, \mathbf{y}) \end{aligned}$$

Definition 8 (Elementary functions) A function over the integers is elementary if and only if it belongs to the smallest set of functions that contains constant function $\mathbf{0}$, the projection functions π_i^p , the functions successor \mathbf{s} , addition $+$, limited subtraction $\ominus : (n_1, n_2) \mapsto \max(0, n_1 - n_2)$, and that is closed under composition, bounded sum BSUM and bounded product BPROD .

We denote by \mathcal{E} the class of elementary functions.

Class \mathcal{E} contains many classical functions. In particular:

Lemma 3 ([36, Lemma 2.5, page 6]) $(x, y) \mapsto \lfloor x/y \rfloor$ is in \mathcal{E} .

Lemma 4 ([36]) $(x, y) \mapsto x \cdot y$ is in \mathcal{E} .

The following normal form is also well-known. We consider safe minimization instead of classical minimization as we focus in this article only on total functions.

Definition 9 ((Safe) Minimization) Given function $g : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$, such that for all x there exists \mathbf{y} with $g(x, \mathbf{y}) = 0$, function $f = \text{SMIN}(g)$ defined by (safe) minimization from g is the (total) function $\mathbb{N}^p \rightarrow \mathbb{N}$ satisfying $\text{SMIN}(g) : \mathbf{y} \mapsto \min\{x; g(x, \mathbf{y}) = 0\}$.

Theorem 4 (Normal form for computable functions [23, 36]) Any total recursive function f can be written as $f = g(\text{SMIN}(h))$ for some elementary functions g and h .

Consider the family of functions E_n defined by induction as follows. When f is a function, $f^{[d]}$ denotes its d -th iterate: $f^{[0]}(\mathbf{x}) = x$, $f^{[d+1]}(\mathbf{x}) = f(f^{[d]}(\mathbf{x}))$:

$$\begin{aligned} \mathbf{E}_0(x) &= s(x) = x + 1, \\ \mathbf{E}_1(x, y) &= x + y, \\ \mathbf{E}_2(x, y) &= (x + 1) \cdot (y + 1), \\ \mathbf{E}_3(x) &= 2^x, \\ \mathbf{E}_{n+1}(x) &= \mathbf{E}_n^{[x]}(1) \text{ for } n \geq 3. \end{aligned}$$

Definition 10 (Bounded recursion) Given functions $g(\mathbf{y}) : \mathbb{N}^p \rightarrow \mathbb{N}$ and $h(f, x, \mathbf{y}) : \mathbb{N}^{p+2} \rightarrow \mathbb{N}$ and $i(x, \mathbf{y}) : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$, the function $f = \text{BR}(g, h)$ defined by bounded recursion from g and h is defined as the function $\mathbb{N}^{p+1} \rightarrow \mathbb{N}$ verifying

$$\begin{aligned} f(0, \mathbf{y}) &= g(\mathbf{y}) \\ f(x+1, \mathbf{y}) &= h(f(x, \mathbf{y}), x, \mathbf{y}) \end{aligned}$$

under the condition that:

$$f(x, \mathbf{y}) \leq i(x, \mathbf{y}).$$

Definition 11 (Grzegorzcyk hierarchy (see [36])) Class \mathcal{E}^0 denotes the class that contains the constant function $\mathbf{0}$, the projection functions π_i^p , the successor function \mathbf{s} , and that is closed under composition and bounded recursion.

Class \mathcal{E}^n for $n \geq 1$ is defined similarly except that functions \max and \mathbf{E}_n are added to the list of initial functions.

Theorem 5 ([34, 8]) Let $n \geq 3$. A function is in class \mathcal{E}_n iff it belongs to the smallest set of functions that contains constant function $\mathbf{0}$, the projection functions π_i^p , the functions successor \mathbf{s} , addition $+$, subtraction \ominus , and the function \mathbf{E}_n and that is closed under composition, bounded sum and bounded product.

The above proposition means that closure under bounded recursion is equivalent to using both closure under bounded sum and closure under bounded product. Indeed, as explained in chapter 1 of [36] (see Theorem 3.1 for details), bounded recursion can be expressed as a minimization of bounded sums and bounded products, itself being expressed as a bounded sum of bounded products.

The following facts are known:

Theorem 6 ([36, 34, 11])

$$\begin{aligned} \mathcal{E}_3 &= \mathcal{E} \subsetneq \mathcal{PR} \\ \mathcal{E}_n &\subsetneq \mathcal{E}_{n+1} \text{ for } n \geq 3 \\ \mathcal{PR} &= \bigcup_i \mathcal{E}_i \end{aligned}$$

4.2 Complexity theory and bounded schemes

We suppose the reader familiar with the well-known complexity classes **P**TIME (polynomial time), **N**PTIME or (non-deterministic polynomial time) or **P**SPACE (polynomial space). We denote by **F**PTIME (resp. **F**PSPACE) the class of functions, $f : \mathbb{N}^k \rightarrow \mathbb{N}$ with $k \in \mathbb{N}$, computable in polynomial time (resp. polynomial space) on deterministic Turing machines. Note that if **F**PTIME is closed by composition, it is not the case of **F**PSPACE since the size of the output can be exponentially larger than the size of the input.

It turns out that the main complexity classes have also been characterized algebraically, by restricted form of recursion scheme. A foundational result in that spirit is due to Cobham, who gave in [12] a characterization of function computable in polynomial time. The idea is to consider schemes similar to primitive recursion, but with restricting the number of induction steps.

Let $\mathbf{0}(\cdot)$ and $\mathbf{1}(\cdot)$ be the successor functions defined by $\mathbf{0}(x) = 2.x$ and $\mathbf{1}(x) = 2.x + 1$.

Definition 12 (Bounded recursion on notations) *A function f is defined by bounded recursion scheme on notations from g, h_0, h_1, k , denoted by $f = \text{BRN}(g, h_0, h_1)$, if*

$$\begin{aligned} f(\mathbf{0}, \mathbf{y}) &= g(\mathbf{y}) \\ f(\mathbf{0}(x), \mathbf{y}) &= h_0(f(x, \mathbf{y}), x, \mathbf{y}) \text{ for } x \neq 0 \\ f(\mathbf{1}(x), \mathbf{y}) &= h_1(f(x, \mathbf{y}), x, \mathbf{y}) \end{aligned}$$

under the condition that:

$$f(x, \mathbf{y}) \leq k(x, \mathbf{y})$$

for all x, \mathbf{y} .

Based on this scheme, Cobham proposed the following class of functions:

Definition 13 (\mathcal{F}_p) *The class \mathcal{F}_p is the smallest class of primitive recursive functions containing $\mathbf{0}$, the projections π_i^p , the successor functions $\mathbf{0}(x) = 2.x$ and $\mathbf{1}(x) = 2.x + 1$, the function $\#$ defined by $x\#y = 2^{\ell(x) \times \ell(y)}$ and closed by composition and by bounded recursion scheme on notations.*

This class turns out to be a characterization of polynomial time:

Theorem 7 ([12], see [10] for a proof) $\mathcal{F}_p = \mathbf{FPTIME}$.

Cobham's result opened the way to various characterizations of complexity classes, or various ways to control recursion schemes. This includes the famous characterization of **PTIME** from Bellantoni and Cook in [2] and by Leivant in [27]. Refer to [10, 11] for monographies presenting a whole serie of results in that spirit.

The task to capture **FPSPACE** is less easy since the principle of such characterizations is to use classes of functions closed by composition. However, for function with a reasonable output size some characterizations have been obtained. Let us denote by $\mathcal{F}_{\mathbf{PSPACE}}$, the class of functions of polynomial growth i.e. of functions $f : \mathbb{N}^k \rightarrow \mathbb{N}$, such that, for all $\mathbf{x} \in \mathbb{N}^k$, $\ell(f(\mathbf{x})) = O(\max_{1 \leq i \leq k} \ell(x_i))$. The following then holds:

Theorem 8 ([38],[11, Theorem 6.3.16]) *A function over the integers is in $\mathcal{F}_{\mathbf{PSPACE}}$ if and only if it belongs to the smallest set of functions that contains the constant function $\mathbf{0}$, the projection functions π_i^p , the functions successor \mathbf{s} , function $\#$ that is closed under composition and bounded recursion.*

5 Computability and Discrete ODEs

Before coming back to efficient algorithms and complexity theory, we consider functions defined by ODE under the prism of computability. This part is clearly inspired by ideas from [8, 9], but adapted here for our framework of discrete ODEs that we believe to provide simpler explanations of statements of these papers.

5.1 About positive and negative integers and encodings

As classical computability is mainly dealing with functions over the natural integers, i.e. over \mathbb{N} , while schemes with discrete ODEs naturally deals with functions over the integers, i.e. over \mathbb{Z} , we need to fix some conventions to be able to compare classes over the integers. Notice that this is very natural in our framework to consider functions that may take negative values.

Definition 14 (Representation of integers) *The set \mathbb{Z} of integers can be encoded by the set $\{0, 1\} \times \mathbb{N}$: couple (s, n) with $s \in \{0, 1\}$, $n \in \mathbb{N}$ encodes $(-1)^s n$. Notice that 0 corresponds both to $(0, 0)$ and $(1, 0)$. To avoid confusion, we will denote by $\tilde{\mathbb{Z}}$ the set $\{0, 1\} \times \mathbb{N}$.*

We will only deal with classes \mathcal{C} of functions over either the natural integer \mathbb{N} or integers \mathbb{Z} . We basically use the same convention for functions : let $f : \mathbb{N}^k \times \mathbb{Z}^h \rightarrow \mathbb{N}^r \times \mathbb{Z}^s$, we denote by $\tilde{f} : \mathbb{N}^k \times \tilde{\mathbb{Z}}^h \rightarrow \mathbb{N}^r \times \tilde{\mathbb{Z}}^s$ the function equivalent to f with above representation.

Note that, if x, y and z are such that $x = y + z$ then $\tilde{x} = \tilde{y} \tilde{+} \tilde{z}$ and $\tilde{+}$ is primitive recursive. The same holds for multiplication and subtraction.

5.2 Recursive and subrecursive classes of functions

A first key remark is that at a computability level, many schemes can actually be seen as particular natural types of ODEs.

The coming discussion is clearly inspired from considerations in [8, 9] for classical continuous ODEs. Here we focus on discrete ODEs: We believe that the settings here in terms of discrete ODEs instead of classical continuous ODEs is really simpler and helps to understand the constructions in the series of related papers that have been published later on for characterizing various classes of the so-called \mathbb{R} -recursive functions (see previous discussions on related work for references). Our settings in particular avoid discussions related to how to deal with noise in computations, as we are living in a world where computations are exact, compared to the above mentioned settings. Furthermore, we believe it clearly helps the intuition of many of the constructions done in all these references.

5.3 Subrecursive functions and discrete ODEs

First, the purpose of this subsection is to observe that primitive recursion is basically a discrete ODE schemata:

Definition 15 ((Scalar) Discrete ODE schemata) *Given $g : \mathbb{N}^p \rightarrow \mathbb{N}$ and $h : \mathbb{Z} \times \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$, we say that f is defined by discrete ODE solving from g and h , denoted by $f = \text{ODE}(g, h)$, if $f : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ corresponds to the (necessarily unique) solution of Initial Value Problem*

$$\begin{aligned} \frac{\partial f(x, \mathbf{y})}{\partial x} &= h(f(x, \mathbf{y}), x, \mathbf{y}) \\ f(0, \mathbf{y}) &= g(\mathbf{y}). \end{aligned} \quad (7)$$

Remark 6 *To be more general, we could take $g : \mathbb{N}^p \rightarrow \mathbb{Z}$. However, this would be of no use in the context of this paper.*

Lemma 5 (Primitive recursion vs Discrete ODEs) *1. Consider g and h as in Definition 15 and $f = \text{ODE}(g, h)$. Then f is primitive recursive when g and h are. When $f : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$, then f is primitive recursive under the same conditions*

2. Consider g and h as in Definition 4. Then $f = \text{REC}(g, h)$ corresponds also to $f = \text{ODE}(g, \bar{h})$ where $\bar{h} : \mathbb{N}^{p+2} \rightarrow \mathbb{Z}$ is defined by

$$\bar{h}(f(x, \mathbf{y}), x, \mathbf{y}) = h(f(x, \mathbf{y}), x, \mathbf{y}) - f(x, \mathbf{y}).$$

Proof: For statement 1., applying the ODE schemata on primitive recursive functions \tilde{h} and g , it holds that: $f(0, \mathbf{y}) = g(\mathbf{y})$ and $\tilde{f}(x+1, \mathbf{y}) = \tilde{h}(\tilde{f}(x, \mathbf{y}), x, \mathbf{y}) + \tilde{f}(x, \mathbf{y})$, where addition is redefined to apply to elements of \mathbb{Z} . This is easily seen to be primitive recursive. When $f : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$, one can extract $f(x, \mathbf{y})$ from $\tilde{f}(x, \mathbf{y})$ by a primitive recursive function.

For statement 2., remark that $h(f(x, \mathbf{y}), x, \mathbf{y}) - f(x, \mathbf{y}) = f(x+1, \mathbf{y}) - f(x, \mathbf{y}) = \frac{\partial f(x, \mathbf{y})}{\partial x}$. \square

Lemma 5 combined with Definition 4 provides the following important characterization of primitive recursive functions in terms of discrete ODEs.

Theorem 9 (A discrete ODE characterization of primitive recursive functions)

The set of primitive recursive functions \mathcal{PR} is the intersection with $\mathbb{N}^{\mathbb{N}}$ of the smallest set of functions that contains the zero functions $\mathbf{0}$, the projection functions π_i^p , the addition and subtraction functions $+$ and $-$, and that is closed under composition and discrete ODE schemata.

5.4 Elementary functions, Grzegorzcyk hierarchy and linear discrete ODEs

Actually, this is even possible to be more precise, and provide a characterization of the various subrecursive classes introduced up to now. This part is clearly inspired from ideas from [8, 9], adapted here for discrete ODEs.

This is very natural to restrict to linear ODEs. This provides natural ways to talk about elementary functions and levels of the Grzegorzczk hierarchy.

Definition 16 ((Scalar) Linear ODE schemata) *Given $g : \mathbb{N}^p \rightarrow \mathbb{N}$, $a : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ and $b : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$, we say that f is obtained by linear ODE solving from g, a and b , denoted by $f = \text{LI}(g, a, b)$, if $f : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ corresponds to the (necessarily unique) solution of Initial Value Problem*

$$\begin{aligned} \frac{\partial f(x, \mathbf{y})}{\partial x} &= a(x, \mathbf{y}) \cdot f(x, \mathbf{y}) + b(x, \mathbf{y}) \\ f(0, \mathbf{y}) &= g(\mathbf{y}). \end{aligned} \quad (8)$$

First observe that bounded sums and products are of this specific form:

Lemma 6 (Bounded sum) *Let $k : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$ be given. Then $f = \text{BSUM}_{<}(k)$ is the unique solution of initial value problem*

$$\begin{aligned} \frac{\partial f(x, \mathbf{y})}{\partial x} &= k(x, \mathbf{y}) \\ f(0, \mathbf{y}) &= 0 \end{aligned}$$

Lemma 7 (Bounded product) *Let $k : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$ be given. Then $f = \text{BPROD}_{<}(k)$ is the unique solution of initial value problem*

$$\begin{aligned} \frac{\partial f(x, \mathbf{y})}{\partial x} &= f(x, \mathbf{y}) \cdot (k(x, \mathbf{y}) - 1) \\ f(0, \mathbf{y}) &= 1 \end{aligned}$$

In the context of Ordinary Differential Equations, this is very natural not to restrict to scalar functions, and the following makes a clear natural sense.

Definition 17 (Linear ODE schemata) *Given a vector $\mathbf{G} = (G_i)_{1 \leq i \leq k}$ matrix $\mathbf{A} = (A_{i,j})_{1 \leq i, j \leq k}$, $\mathbf{B} = (B_i)_{1 \leq i \leq k}$ whose coefficients corresponds to functions $g_i : \mathbb{N}^p \rightarrow \mathbb{N}^k$, and $a_{i,j} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ and $b_{i,j} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ respectively, we say that \mathbf{f} is obtained by linear ODE solving from g, \mathbf{A} and \mathbf{B} , denoted by $\mathbf{f} = \text{LI}(\mathbf{G}, \mathbf{A}, \mathbf{B})$, if $f : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^k$ corresponds to the (necessarily unique) solution of Initial Value Problem*

$$\begin{aligned} \frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial x} &= \mathbf{A}(x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(x, \mathbf{y}) \\ \mathbf{f}(0, \mathbf{y}) &= \mathbf{G}(\mathbf{y}). \end{aligned} \quad (9)$$

One key observation behind the coming characterizations is the following:

Lemma 8 (Elementary vs Linear ODEs) *Consider \mathbf{G}, \mathbf{A} and \mathbf{B} as in Definition 17. Then $\mathbf{f} = \text{LI}(\mathbf{G}, \mathbf{A}, \mathbf{B})$ is elementary when \mathbf{G}, \mathbf{A} and \mathbf{B} are.*

Proof: We do the proof in the scalar case, writing a, b, g for $\mathbf{A}, \mathbf{B}, \mathbf{G}$. The general (vectorial) case follows from similar arguments. By Lemma 2, it follows that:

$$f(x, \mathbf{y}) = \left(\prod_{t=0}^{t=x-1} (1 + a(t, \mathbf{y})) \right) \cdot g(\mathbf{y}) + b(x-1, \mathbf{y}) + \sum_{u=0}^{x-2} \left(\prod_{t=u+1}^{x-1} (1 + a(t, \mathbf{y})) \right) \cdot b(u, \mathbf{y}).$$

Clearly, $\prod_{t=0}^{t=x-1} (1 + a(t, \mathbf{y})) = \text{BPROD}_{<}(1 + a(t, \mathbf{y}))(x, \mathbf{y})$. Similarly,

$$p(u, x, \mathbf{y}) =_{\text{def}} \prod_{t=u+1}^{x-1} (1 + a(t, \mathbf{y})) = \frac{\text{BPROD}_{<}(1 + a(t, \mathbf{y}))(x, \mathbf{y})}{\text{BPROD}_{<}(1 + a(t, \mathbf{y}))(u + 1, \mathbf{y})}$$

As the function $(x, y) \mapsto \lfloor x/y \rfloor$ is elementary from Lemma 3, we get that $p(x, \mathbf{y})$ is elementary.

As multiplication is elementary, it follows that

$$\sum_{u=0}^{x-2} p(x, \mathbf{y}) b(u, \mathbf{y}) = \text{BSUM}_{<}(p(u, x, \mathbf{y}) b(u, \mathbf{y}))(x - 2, \mathbf{y})$$

is also elementary, and \tilde{f} is elementary using closure by composition and multiplication. \square

We get the following elegant characterization of the Elementary functions in terms of Linear ODEs.

Theorem 10 (A discrete ODE characterization of elementary functions)

The set of elementary functions \mathcal{E} is the intersection with $\mathbb{N}^{\mathbb{N}}$ of the smallest set of functions that contains the zero functions $\mathbf{0}$, the projection functions π_i^p , the successor function \mathbf{s} , addition $+$, subtraction $-$, and that is closed under composition and discrete linear ODE schemata (respectively: scalar discrete linear ODE schemata) LI.

Proof:

\square

Inspired by bounded recursion, this also makes sense to consider the following (as expected, we write $\mathbf{u} \leq \mathbf{v}$ if it holds componentwise):

Definition 18 (Bounded discrete ODE schemata) *Given $\mathbf{g}(\mathbf{y}) : \mathbb{N}^p \rightarrow \mathbb{N}^k$ and $\mathbf{h}(\mathbf{f}, x, \mathbf{y}) : \mathbb{Z}^k \times \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^k$, and $\mathbf{i}(x, \mathbf{y}) : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^k$, we say that \mathbf{f} is defined by bounded discrete ODE solving from \mathbf{g}, \mathbf{h} and \mathbf{i} , denoted by $\mathbf{f} = \text{boundedlinODE}(\mathbf{g}, \mathbf{h}, \mathbf{i})$, if $\mathbf{f} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^k$ corresponds to the (necessarily unique) solution of Initial Value Problem*

$$\begin{aligned} \frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial x} &= \mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) \\ \mathbf{f}(0, \mathbf{y}) &= \mathbf{g}(\mathbf{y}) \end{aligned}$$

under the condition that:

$$\mathbf{f}(x, \mathbf{y}) \leq \mathbf{i}(x, \mathbf{y})$$

Lemma 9 (Primitive recursion vs Discrete ODEs) 1. Consider \mathbf{g} , \mathbf{h} , \mathbf{i} as in Definition 18. Then $\tilde{\mathbf{f}} = \text{boundedlinODE}(\mathbf{g}, \tilde{\mathbf{h}}, \mathbf{i})$ is in \mathcal{E}_n when \mathbf{g} and \mathbf{h} and \mathbf{i} are, and $n \geq 3$.

2. Consider \mathbf{g} , \mathbf{h} , \mathbf{i} as in Definition 10. Then $\mathbf{f} = \text{BR}(\mathbf{g}, \mathbf{h}, \mathbf{i})$ corresponds also to $\mathbf{f} = \text{boundedlinODE}(\mathbf{g}, \bar{\mathbf{h}}, \mathbf{i})$ where $\bar{\mathbf{h}} : \mathbb{N}^{p+2} \rightarrow \mathbb{Z}^k$ is defined by $\bar{\mathbf{h}}(\mathbf{f}, x, \mathbf{y}) = \mathbf{h}(\mathbf{f}, x, \mathbf{y}) - \mathbf{f}$.

Proof: For statement 1., this follows from exactly the same proof as for Lemma 8.

Second item can be proved by observing that $\frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial x} = \bar{\mathbf{h}}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) = \mathbf{f}(x+1, \mathbf{y}) - \mathbf{f}(x, \mathbf{y})$ which is equal to $\mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) - \mathbf{f}(x, \mathbf{y})$ by Definition 18. \square

This provide the following elegant characterizations of the levels of the Grzegorzcyk hierarchy in terms of bounded linear ODEs.

Theorem 11 (A discrete ODE characterization of \mathcal{E}_n for $n \geq 3$) For all $n \geq 3$, the set of functions in \mathcal{E}_n is the smallest set of functions that contains \mathbf{E}_n , constant function $\mathbf{0}$, the projection functions π_i^p , the functions successor \mathbf{s} , and that is closed under composition and boundedlinODE.

Proof: Using Theorem 5, this follows from Lemmas 6, 7 and 9. \square

5.5 Computability and discrete ODEs

If we want to talk about computable functions, and not only about subrecursive functions, a first method is to add directly minimization to considered operators.

5.5.1 By adding a minimization operator

Theorem 12 (A discrete ODE characterization of total recursive functions)

The set of total recursive functions \mathcal{E} is the intersection with $\mathbb{N}^{\mathbb{N}}$ of the smallest set of functions that contains the zero functions $\mathbf{0}$, the projection functions π_i^p , the successor function \mathbf{s} , addition $+$, subtraction $-$, and that is closed under composition and discrete (even linear) ODE schemata LI, and safe minimization.

Proof: One direction follows from Theorem 10 (characterization of elementary functions) and Theorem 4 (normal form theorem) in one direction. And from a clear generalization of previous arguments in the other direction. \square

5.5.2 By programming minimization

But actually, minimization can be programmed using discrete ODEs *in some sense*. Indeed, minimization can be programmed in the following sense.

Theorem 13 (Programming Minimization) Consider a function $g : \mathbb{N}^{p+1} \rightarrow \mathbb{N}$. Then the solution of initial value problem

$$\begin{aligned} f(0, \mathbf{y}) &= 0 \\ \frac{\partial f(x, \mathbf{y})}{\partial x} &= \text{if}(g(f(x, \mathbf{y})), 1, 0) \end{aligned}$$

is such that for all \mathbf{y} , $f(x, \mathbf{y})$ is eventually a constant $k = k(\mathbf{y})$ when x increases if and only if there is some x with $g(x, \mathbf{y}) = 0$. This constant $k(\mathbf{y})$ corresponds to $\text{SMIN}(g)(\mathbf{y})$ for all \mathbf{y} .

This leads to the following natural concept: The idea is that $\text{SMIN}(g)$ is computable in the following sense considering $h_1(x, \mathbf{y}) = f(x, \mathbf{y})$ and $h_2(x, \mathbf{y}) = \text{s}\bar{g}_{\mathbb{N}}(g(f(x, \mathbf{y})))$.

Definition 19 (Discrete ODEs as a computational model) We say that a total function $f : \mathbb{N}^p \rightarrow \mathbb{N}$ is ODE computable if there exist some function $h_1, h_2 : \mathbb{N}^{p+1} \rightarrow \mathbb{N}^2$ in the smallest set of functions that contains the zero functions $\mathbf{0}$, the projection functions π_i^p , the successor function \mathbf{s} , and that is closed under composition and discrete ODE schemata such that: for all \mathbf{y} ,

- there exists some $T = T(\mathbf{y})$ with $h_2(T, \mathbf{y}) \neq 0$;
- $f(\mathbf{y}) = h_1(T, \mathbf{y})$ where T is the smallest such T .

The following is then easy to establish:

Theorem 14 (Discrete ODE computability and classical computability are equivalent) A total function f is ODE computable if and only if it is total recursive.

6 Restricted recursion and integration schemes

In order to talk about complexity instead of computability, we need to put some restrictions on integrations schemes.

Remark 7 Observe that this is necessary. Indeed, the solution of a polynomial ordinary differential equation (ODE) can grow very very fast.

Indeed:

$$\begin{aligned} (\bar{2}^x)' &= \bar{2}^x \\ \left(\bar{2}^{\bar{2}^x}\right)' &= \bar{2}^x \cdot \bar{2}^{\bar{2}^x} \\ \left(\bar{2}^{\bar{2}^{\bar{2}^x}}\right)' &= \bar{2}^x \cdot \bar{2}^{\bar{2}^x} \cdot \bar{2}^{\bar{2}^{\bar{2}^x}} \\ &\vdots \end{aligned}$$

and so on, is solution of degree 2 polynomial ODE:

$$\begin{aligned} y_1' &= y_1 \\ y_2' &= y_1 \cdot y_2 \\ y_3' &= y_2 \cdot y_3 \\ &\vdots \end{aligned}$$

with initial condition $y_1(0) = y_2(0) = y_3(0) = \dots = 1$. That means that if we consider a two general integration scheme, then we get such towers of exponentials. Clearly, such a function is not polynomial time computable, as only writing its value in binary cannot be done in polynomial time.

We propose to introduce the following variation on the notion of derivation: derivation along some function $\mathcal{L}(x, \mathbf{y})$.

Definition 20 (\mathcal{L} -ODE) Let $\mathcal{L} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$. We write

$$\frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial \mathcal{L}} = \frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial \mathcal{L}(x, \mathbf{y})} = \mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}), \quad (10)$$

as a formal synonym for

$$\mathbf{f}(x + 1, \mathbf{y}) = \mathbf{f}(x, \mathbf{y}) + (\mathcal{L}(x + 1, \mathbf{y}) - \mathcal{L}(x, \mathbf{y})) \cdot \mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}).$$

Remark 8 This is motivated by the fact that the latter expression is similar to classical formula for classical continuous ODEs:

$$\frac{\delta f(x, \mathbf{y})}{\delta x} = \frac{\delta \mathcal{L}(x, \mathbf{y})}{\delta x} \cdot \frac{\delta f(x, \mathbf{y})}{\delta \mathcal{L}(x, \mathbf{y})}.$$

This will allow us to simulate suitable change of variables using this analogy. We will talk about \mathcal{L} -IVP when some initial condition is added. An important special case is when $\mathcal{L}(x, \mathbf{y})$ corresponds to the length $\mathcal{L}(x, \mathbf{y}) = \ell(x)$ function: we will call this special case length-ODEs.

Example 5 (Example 2 continued) The trick used in Example 2 can be read as using a new parameter $t = \ell(x)$ logarithmic in x , using relation

$$\frac{\partial \text{some}_h(x)}{\partial \ell(x)} = E(\text{some}_h(x), \ell(x), x)$$

Example 6 (Function $2^{\ell(x)}$ and $2^{\ell(x)^2}$) To compute function $f : x \mapsto 2^{\ell(x)}$, a method would consist in computing $f(x)$ using the fact $f(x) = \int_0^x f'(t) \delta t$ which would a priori requires time x ; But a more efficient method consists in stating that $f(x) = F(\ell(x))$ where $F(t) = 2^t$ is solution of IVP $F'(t) = F(t)$, $F(0) = 1$. This is a fast (polynomial) algorithm to solve our problem. Once again, we have

used a change of variable in order to compute faster. Thinking about what we have just done, we have basically observed the fact that

$$(2^{\ell(x+1)})' = \ell(x)' \cdot 2^{\ell(x)} \text{ that is to say } \frac{\partial 2^{\ell(x)}}{\partial \ell(x)} = \ell(x)' \cdot 2^{\ell(x)}$$

This is what led us to consider change of variable $t = \ell(x)$ and what led to above more efficient algorithm, considering $F(t)$ instead of $f(x)$, with similarities with the relation for continuous derivative $\frac{\delta f(x)}{\delta t} = \frac{\delta t}{\delta x} \cdot \frac{\delta f(x)}{\delta t}$.

Suppose now that we want to compute function $f : x \mapsto 2^{\ell(x)^2}$. We can use the same principle, observing that

$$(2^{\ell(x+1)^2})' = (\ell(x+1)^2 - \ell(x)^2) \cdot 2^{\ell(x)^2}$$

that is to say

$$\frac{\partial 2^{\ell(x)^2}}{\partial \mathcal{L}} = (\ell(x)^2)' \cdot 2^{\ell(x)^2} \text{ considering } \mathcal{L}(x) = \ell(x)^2$$

and then noticing that f is consequently computed fast (in polynomial time) as $F(\ell(x)^2)$.

Example 7 $f(x, y) = 2^{\ell(x) \cdot \ell(y)}$ is the solution of the following length-IVP:

$$\begin{aligned} f(0, y) &= 2^{|y|} \\ \frac{\partial f(x, y)}{\partial \ell} &= f(x, y) \cdot (2^{\ell(y)} - 1), \end{aligned}$$

since $2^{\ell(x+1) \cdot \ell(y)} = 2^{\ell(x) \cdot \ell(y)} + \ell(x)' \cdot 2^{\ell(x) \cdot \ell(y)} \cdot (2^{\ell(y)} - 1)$.

6.1 General theory

The following result though simple, illustrate one key property of the \mathcal{L} -ODE scheme under a computational point of view: it's dependence on the number of distinct values of function \mathcal{L} .

Definition 21 ($\text{Jump}_{\mathcal{L}}$) Let $\mathcal{L}(x, \mathbf{y})$ be some function. Fixing \mathbf{y} , we write $\text{Jump}_{\mathcal{L}}(x, \mathbf{y}) = \{0 \leq i \leq x - 1 \mid \mathcal{L}(i + 1, \mathbf{y}) \neq \mathcal{L}(i, \mathbf{y})\}$ (that is to say the set of points where \mathcal{L} has a value that changes) and $\alpha : [0..|\text{Jump}_{\mathcal{L}}(x, \mathbf{y})| - 1] \rightarrow \text{Jump}_{\mathcal{L}}(x, \mathbf{y})$ for an increasing function enumerating these points: If $i_0 < i_1 < i_2 < \dots < i_{\text{card}(\text{Jump}_{\mathcal{L}}(x, \mathbf{y})) - 1}$ denote all elements of $\text{Jump}_{\mathcal{L}}(x, \mathbf{y})$, then $\alpha(j) = i_j \in \text{Jump}_{\mathcal{L}}(x, \mathbf{y})$.

Lemma 10 (Fundamental Observation) Let $k \in \mathbb{N}$, $f : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^d$ and $\mathcal{L} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ be some functions. Assume that (10) holds. Then:

$$\mathbf{f}(x, \mathbf{y}) = \mathbf{f}(0, \mathbf{y}) + \int_0^{\text{card}(\text{Jump}_{\mathcal{L}}(x, \mathbf{y}))} \Delta \mathcal{L}(\alpha(u), \mathbf{y}) \cdot \mathbf{h}(\mathbf{f}(\alpha(u), \mathbf{y}), \alpha(u), \mathbf{y}) \delta u$$

Proof:[Proof of Lemma 10] By definition, we have

$$\mathbf{f}(x+1, \mathbf{y}) = \mathbf{f}(x, \mathbf{y}) + (\mathcal{L}(x+1, \mathbf{y}) - \mathcal{L}(x, \mathbf{y})) \cdot \mathbf{h}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}).$$

Hence,

- as soon as $i \notin \text{Jump}_{\mathcal{L}}(x, \mathbf{y})$, then $\mathbf{f}(i+1, \mathbf{y}) = \mathbf{f}(i, \mathbf{y})$, since $\mathcal{L}(i+1, \mathbf{y}) - \mathcal{L}(i, \mathbf{y}) = 0$. In other words,

$$\Delta \mathbf{f}(i, \mathbf{y}) = 0.$$

- as soon as $i \in \text{Jump}_{\mathcal{L}}(x, \mathbf{y})$, say $i = i_j$, then

$$\Delta \mathbf{f}(i_j, \mathbf{y}) = (\mathcal{L}(i_j+1, \mathbf{y}) - \mathcal{L}(i_j, \mathbf{y})) \cdot \mathbf{h}(\mathbf{f}(i_j, \mathbf{y}), i_j, \mathbf{y})$$

I.e.

$$\Delta \mathbf{f}(i_j, \mathbf{y}) = \Delta \mathcal{L}(i_j, \mathbf{y}) \cdot \mathbf{h}(\mathbf{f}(i_j, \mathbf{y}), i_j, \mathbf{y})$$

Now

$$\begin{aligned} \mathbf{f}(x, \mathbf{y}) &= \mathbf{f}(0, \mathbf{y}) + \int_0^x \Delta \mathbf{f}(t, \mathbf{y}) \delta t \\ &= \mathbf{f}(0, \mathbf{y}) + \sum_{t=0}^{x-1} \Delta \mathbf{f}(t, \mathbf{y}) \\ &= \mathbf{f}(0, \mathbf{y}) + \sum_{i_j \in \text{Jump}_{\mathcal{L}}(x, \mathbf{y})} \Delta \mathbf{f}(i_j, \mathbf{y}) \\ &= \mathbf{f}(0, \mathbf{y}) + \sum_{i_j \in \text{Jump}_{\mathcal{L}}(x, \mathbf{y})} \Delta \mathcal{L}(i_j, \mathbf{y}) \cdot \mathbf{h}(\mathbf{f}(i_j, \mathbf{y}), i_j, \mathbf{y}) \\ &= \mathbf{f}(0, \mathbf{y}) + \sum_{j=0}^{\text{card}(\text{Jump}_{\mathcal{L}}(x, \mathbf{y})) - 1} \Delta \mathcal{L}(\alpha(j), \mathbf{y}) \cdot \mathbf{h}(\mathbf{f}(\alpha(j), \mathbf{y}), \alpha(j), \mathbf{y}) \\ &= \mathbf{f}(0, \mathbf{y}) + \int_0^{\text{card}(\text{Jump}_{\mathcal{L}}(x, \mathbf{y}))} \Delta \mathcal{L}(\alpha(u), \mathbf{y}) \cdot \mathbf{h}(\mathbf{f}(\alpha(u), \mathbf{y}), \alpha(u), \mathbf{y}) \delta u \end{aligned}$$

which corresponds to the expression. \square

The proof of the Lemma is based on (and illustrates) some fundamental aspect of \mathcal{L} -ODE from their definition: for fixed \mathbf{y} , the value of $\mathbf{f}(x, \mathbf{y})$ only changes when the value of $\mathcal{L}(x, \mathbf{y})$ changes. This implies that the value of $\mathbf{f}(x, \mathbf{y})$ must then depend on \mathbf{y} and $\mathcal{L}(x, \mathbf{y})$. We formalize this in the following definition.

Definition 22 (\mathcal{L} -expressiveness) Let $k \in \mathbb{N}$, $f : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^d$ and $\mathcal{L} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ be some functions. We say that $\mathbf{f}(x, \mathbf{y})$ is \mathcal{L} -expressible if there exists some function $\mathbf{g} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^d$ such that $\mathbf{f}(x, \mathbf{y}) = \mathbf{g}(\mathcal{L}(x, \mathbf{y}), \mathbf{y})$.

Corollary 1 *Let $k \in \mathbb{N}$, $f : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^d$ and $\mathcal{L} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ be some functions as above. Then $\mathbf{f}(x, \mathbf{y})$ is \mathcal{L} -expressible.*

Let's make a pause to ponder. From the above results, if \mathcal{L} is chosen such that $|\text{Jump}_{\mathcal{L}}(x, \mathbf{y})| = |\{0 \leq i \leq x - 1 \mid \mathcal{L}(i + 1, \mathbf{y}) = \mathcal{L}(i, \mathbf{y})\}| \leq P(\ell(x), \ell(\mathbf{y}))$ for some polynomial P then, the number of distinct values of $\mathbf{f}(x', \mathbf{y})$ with $x' \leq x$ that are necessary to compute $\mathbf{f}(x, \mathbf{y})$ is polynomial in $\ell(x)$ and $\ell(\mathbf{y})$. Hence, at least in terms of the number of steps (not necessarily in terms of the size of the intermediate objects), $\mathbf{f}(x, \mathbf{y})$ can be computed fast.

6.2 Fundamental alternative view

If previous hypotheses hold, there is then an alternative view to understand the integral, by using a change of variable, and by building a discrete ODE that mimics the computation of the integral. Basically, we are using the fact that we can consider some parameter t corresponding to $\mathcal{L}(x, \mathbf{y})$. Indeed:

Lemma 11 (Fundamental alternative view) *Let $k \in \mathbb{N}$, $f : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}^d$, $\mathcal{L} : \mathbb{N}^{p+1} \rightarrow \mathbb{Z}$ be some functions and assume that (10) holds. Then $\mathbf{f}(x, \mathbf{y})$ is given by $\mathbf{f}(x, \mathbf{y}) = \mathbf{F}(\mathcal{L}(x, \mathbf{y}), \mathbf{y})$ where \mathbf{F} is the solution of initial value problem*

$$\begin{aligned} \frac{\partial \mathbf{F}(t, \mathbf{y})}{\partial t} &= \Delta \mathcal{L}(t, \mathbf{y}) \cdot \mathbf{h}(\mathbf{F}(t, \mathbf{y}), t, \mathbf{y}) \\ \mathbf{F}(0, \mathbf{y}) &= \mathbf{f}(\mathcal{L}(0, \mathbf{x}), \mathbf{y}). \end{aligned}$$

We will say in that case the IVP is converging “in time $\mathcal{L}(x, \mathbf{y})$ ” to $\mathbf{f}(x, \mathbf{y})$. Conversely, if there is such a function \mathbf{F} , then a discrete ODE of the type of (10) can easily be derived.

Example 8 *The previous discussion about the complexity of computing $x \mapsto 2^{\ell(x)}$ and $x \mapsto 2^{\ell(x)^2}$ is a concrete applications of all these remarks.*

Example 9 *Let us consider an example, where $\mathcal{L}(x)$ is not $\ell(x)$ (or a power of it): Suppose we want to compute $f : x \mapsto 2^{\lfloor \sqrt{x} \rfloor}$: Consider $\mathcal{L}(x) = \lfloor \sqrt{x} \rfloor$. We have*

$$\frac{\partial f(x)}{\partial \mathcal{L}(x)} = \mathcal{L}'(x) \cdot f(x) = (\lfloor \sqrt{x+1} \rfloor - \lfloor \sqrt{x} \rfloor) \cdot f(x).$$

One may think that the number $|\text{Jump}_{\mathcal{L}}(x)|$ of \mathcal{L} , i.e. the number of jumps of factor $(\lfloor \sqrt{x+1} \rfloor - \lfloor \sqrt{x} \rfloor)$ is hard to predict, but the point is to look at the method we devised to compute $\lfloor \sqrt{x} \rfloor$ in Example 2: It is basically expressing $\lfloor \sqrt{x} \rfloor$ as some function G of $\text{some}_h(x)$: We wrote $\lfloor \sqrt{x} \rfloor = G(\text{some}_h(x))$ for some function G . Consequently, we could also consider variable $\mathcal{L}_2(x) = \text{some}_h(x)$, and see from expressions that the number of jumps $|\text{Jump}_{\mathcal{L}}(x)|$ of previous \mathcal{L} is actually related to the $|\text{Jump}_{\mathcal{L}_2}(x)|$ of this new $\mathcal{L}_2(x)$. We also have

$$\frac{\partial f(x)}{\partial \mathcal{L}_2(x)} = \mathcal{L}'_2(x) \cdot f(x) = (G(\text{some}_h(x+1)) - G(\text{some}_h(x))) \cdot f(x).$$

Observing that $\text{some}_h(x)$ is in turn computed in “time” $\ell(x)$ using the method of Example 2, the number of jumps for all these $\mathcal{L}(x)$ are always polynomials, and we are guarantee that all these expressions lead to fast (polynomial) algorithms.

Remark 9 This method clearly extends to more general functions: Generalizing the above reasoning, we can compute fast functions of type $x \mapsto g(\lfloor \sqrt{x} \rfloor)$ as soon as we have a fast ODE computing g . Similarly, $\lfloor \sqrt{x} \rfloor$ can be replaced by anything that can be computed fast basically using similar techniques.

s

6.3 Length-ODEs

An important and natural case is the special case where $\mathcal{L}(x, \mathbf{y})$ is the usual one variable length function $\mathcal{L}(x, \mathbf{y}) = \ell(x)$. We will of course write $\frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial \ell}$ in that case for $\frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial \mathcal{L}}$.

We can adapt the Lemma above to this special case of a what we will call length-ODE. Namely:

Corollary 2 (First view) Let $\mathcal{L} : \mathbb{N} \rightarrow \mathbb{N}$ be defined by $\mathcal{L}(x) = \ell(x)$ for all integer x and f satisfies the hypothesis of Lemma 10. Then,

$$\mathbf{f}(x, \mathbf{y}) = \mathbf{f}(0, \mathbf{y}) + \int_0^{\ell(x)} \mathbf{h}(\mathbf{f}(2^u - 1, \mathbf{y}), 2^u - 1, \mathbf{y}) \delta u$$

Or, equivalently:

$$\mathbf{f}(x, \mathbf{y}) = \mathbf{f}(0, \mathbf{y}) + \sum_{i=0}^{\ell(x)-1} \mathbf{h}(\mathbf{f}(2^i - 1, \mathbf{y}), 2^i - 1, \mathbf{y})$$

Proof: Immediate consequence of Lemma 10. Function α is such that $\alpha(i) = 2^i - 1$. \square

Corollary 3 (Alternative view) Let $\mathcal{L} : \mathbb{N} \rightarrow \mathbb{N}$ be defined by $\mathcal{L}(x) = \ell(x)$ for all integer x and f satisfies the hypothesis of Lemma 10. Then $\mathbf{f}(x, \mathbf{y})$ is given by $\mathbf{f}(x, \mathbf{y}) = \mathbf{F}(\ell(x), \mathbf{y})$ where \mathbf{F} is the solution of initial value problem

$$\frac{\partial \mathbf{F}(t, \mathbf{y})}{\partial t} = \mathbf{h}(\mathbf{F}(t, \mathbf{y}), t, \mathbf{y}) \text{ with } \mathbf{F}(0, \mathbf{y}) = \mathbf{f}(0, \mathbf{y})$$

In other words, for $\mathcal{L}(x) = \ell(x)$, this offers us also two ways to present a length-ODE for a function $f(x, \mathbf{y})$: either by considering equation of the type of (10) or by considering $\mathbf{f}(x, \mathbf{y}) = \mathbf{F}(\ell(x), \mathbf{y})$ where \mathbf{F} given by an equation of the form:

$$\frac{\partial \mathbf{F}(t, \mathbf{y})}{\partial t} = \mathbf{h}(\mathbf{F}(t, \mathbf{y}), t, \mathbf{y}) \tag{11}$$

with $\mathbf{F}(0, \mathbf{y}) = \mathbf{f}(0, \mathbf{y})$. As before, the idea is that t is a parameter logarithmic in x , namely $t = \ell(x)$.

Our purpose now is to discuss which kind of problems can be solved efficiently using similar techniques: it turns out to be exactly all of **FPTIME**. It will be made clear from the incoming discussion and results.

6.4 Linear length-ODEs

Remark 10 *In all previous reasoning, we considered that a function over the integers is polynomial time computable if it is in the length of all its arguments, as this is the usual convention. When not explicitly stated, this is our convention. As usual, we also say that some vectorial function (respectively: matrix) is polynomial time computable if all its components are. We will need sometimes to consider also polynomial dependency directly in some of the variables and not on their length: This happens in the next fundamental lemma.*

We write $\|\cdot\|$ for the sup norm: given some matrix $\mathbf{A} = (A_{i,j})_{1 \leq i \leq n, 1 \leq j \leq m}$, $\|\mathbf{A}\| = \max_{i,j} A_{i,j}$.

Lemma 12 (Fundamental observation) *Consider ODE*

$$\mathbf{f}'(x, \mathbf{y}) = \mathbf{A}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}). \quad (12)$$

Assume:

1. *Initial condition $\mathbf{G}(\mathbf{y}) =^{def} \mathbf{f}(0, \mathbf{y})$, as well as Matrix \mathbf{A} and vector \mathbf{B} are polynomial time computable.*
2. *$\ell(\|\mathbf{A}(\mathbf{f}, x, \mathbf{y})\|) \leq \ell(\|\mathbf{f}\|) + p_{\mathbf{A}}(x, \ell(\mathbf{y}))$ for some polynomial $p_{\mathbf{A}}$*
3. *$\ell(\|\mathbf{B}(\mathbf{f}, x, \mathbf{y})\|) \leq \ell(\|\mathbf{f}\|) + p_{\mathbf{B}}(x, \ell(\mathbf{y}))$ for some polynomial $p_{\mathbf{B}}$*

Then its solution $\mathbf{f}(x, \mathbf{y})$ is polynomial time computable in x and the length of \mathbf{y} .

Proof: We know by Lemma 2 that we must have:

$$\mathbf{f}(x, \mathbf{y}) = \left(\overline{2}^{\int_0^x \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y}) \delta t} \right) \cdot \mathbf{G}(\mathbf{y}) + \int_0^x \left(\overline{2}^{\int_{u+1}^x \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y}) \delta t} \right) \cdot \mathbf{B}(\mathbf{f}(u, \mathbf{y}), u, \mathbf{y}) \delta u. \quad (13)$$

The key point is that Equation (13) provides a (recursive) algorithm to compute $\mathbf{f}(x, \mathbf{y})$ for all x . To see it, it may help to see that this can also be expressed as

$$\mathbf{f}(x, \mathbf{y}) = \sum_{u=-1}^{x-1} \left(\prod_{t=u+1}^{x-1} (1 + \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y})) \right) \cdot \mathbf{B}(\mathbf{f}(u, \mathbf{y}), u, \mathbf{y}). \quad (14)$$

with the conventions that $\prod_x^{x-1} \kappa(x) = 1$ and $\mathbf{B}(\cdot, -1, \mathbf{y}) = \mathbf{G}(\mathbf{y})$.

Clearly the number of arithmetic operations to evaluate $\mathbf{f}(x, \mathbf{y})$ by this method is polynomial in x : basically we have to sum $x + 1$ terms, each of them involving at most $x - 1$ multiplications. This can be done in the requested complexity if we are sure that the size of the involved quantities remains polynomial in x and the length of \mathbf{y} .

Since the length of $\mathbf{B}(\mathbf{f}(u, \mathbf{y}), u, \mathbf{y})$ and of $\mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y})$ is at most polynomial in the length of $\mathbf{f}(t, \mathbf{y})$ we only need to be convinced that the size of $\mathbf{f}(x, \mathbf{y})$ remains polynomial. But it holds, as from (12) we get

$$\mathbf{f}(x + 1, \mathbf{y}) = (1 + \mathbf{A}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y})) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y})$$

and hence

$$\begin{aligned} \ell(\mathbf{f}(x + 1, \mathbf{y})) &\leq \max(\ell((1 + \mathbf{A}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}))) + \ell(\mathbf{f}(x, \mathbf{y})), \ell(\mathbf{B}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}))) \\ &\leq \ell(\mathbf{f}(x, \mathbf{y})) + p_{\mathbf{f}}(x, \ell(\mathbf{y})) + 1 \end{aligned}$$

for polynomial $p_{\mathbf{f}}$, that we may assume without loss of generality to be increasing in its first argument. It follows from an easy induction that we must have

$$\ell(\mathbf{f}(x, \mathbf{y})) \leq \ell(G(\mathbf{y})) + x \cdot p_{\mathbf{f}}(x, \ell(\mathbf{y})).$$

□

We now go to specific forms of linear ODEs.

Definition 23 A *sg-polynomial expression* $P(x_1, \dots, x_h)$ is a expression built-on $+$, $-$, \times (often denoted \cdot) and $\mathbf{sg}(\cdot)$ functions over a set of variables $V = \{x_1, \dots, x_h\}$ and integer constants. The degree $\deg(x, P)$ of a term $x \in V$ in P is defined inductively as follows:

- $\deg(x, x) = 1$ and for $x' \in X \cup \mathbb{Z}$ such that $x' \neq x$, $\deg(x, x') = 0$
- $\deg(x, P + Q) = \max\{\deg(x, P), \deg(x, Q)\}$
- $\deg(x, P \times Q) = \deg(x, P) + \deg(x, Q)$
- $\deg(x, \mathbf{sg}(P)) = 0$

A *sg-polynomial expression* P is *essentially constant in x* if $\deg(x, P) = 0$.

Compared to the classical notion of degree in polynomial expression, all subterms that are within the scope of a sign function contributes for 0 to the degree. A vectorial function (resp. a matrix or a vector) is said to be a *sg-polynomial expression* if all its coordinates (resp. coefficients) are. it is said to be *essentially constant* if all its coefficients are.

A (possibly vectorial) *sg-polynomial expression* $\mathbf{g}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y})$ is said to be *essentially linear in $\mathbf{f}(x, \mathbf{y})$* if it is of the form

$$\mathbf{g}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) = \mathbf{A}[\mathbf{f}(x, \mathbf{y}), \mathbf{h}(x, \mathbf{y}), x, \mathbf{y}] \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}[\mathbf{f}(x, \mathbf{y}), \mathbf{h}(x, \mathbf{y}), x, \mathbf{y}]$$

where \mathbf{A} and \mathbf{B} are *sg-polynomial expressions essentially constant in $\mathbf{f}(x, \mathbf{y})$* .

Example 10 The expression $P(x, y, z) = x \cdot \text{sg}((x^2 - z) \cdot y) + y^3$ is linear in x , essentially constant in z and not linear in y . The expression $P(x, 2^{\ell(y)}, z) = \text{sg}(x^2 - z) \cdot z^2 + 2^{\ell(y)}$ is essentially constant in x , essentially linear in $2^{\ell(y)}$ (but not essentially constant) and not essentially linear in z . The expression: $\text{if}(x, y, z) = y + \overline{\text{sg}}(x) \cdot (z - y) = y + (1 - \text{sg}(x)) \cdot (z - y)$ is essentially constant in x and linear in y and z .

Definition 24 Function \mathbf{f} is linear \mathcal{L} -ODE definable (from \mathbf{u} and \mathbf{g}) if it corresponds to the solution of \mathcal{L} -IVP

$$\begin{aligned} \frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial \mathcal{L}} &= \mathbf{u}(\mathbf{f}(x, \mathbf{y}), \mathbf{h}(x, \mathbf{y}), x, \mathbf{y}) \\ \mathbf{f}(0, \mathbf{y}) &= \mathbf{g}(\mathbf{y}) \end{aligned} \quad (15)$$

where \mathbf{u} is essentially linear in $\mathbf{f}(x, \mathbf{y})$. When $\mathcal{L}(x, \mathbf{y}) = \ell(x)$, such a system is called linear length-ODE.

The previous statements lead to the following:

Lemma 13 (Fundamental Observation for linear \mathcal{L} -ODE) Assume that $\mathbf{f}(x, \mathbf{y})$ is solution of (15). Then $\mathbf{f}(x, \mathbf{y})$ can be computed in polynomial time under the following conditions:

1. $\mathbf{f}(0, \mathbf{y}) = \mathbf{g}(\mathbf{y})$ is computable in polynomial-time.
2. function \mathbf{h} is computable in polynomial time.
3. there exist $c \in \mathbb{N}$, such that, for each \mathbf{y} , $|\text{Jump}_{\mathcal{L}}(\mathbf{y})| \leq \ell(x)^c$

Proof: Thanks to condition above, we can replace parameter x and derivation in $\mathcal{L}(x, \mathbf{y})$ by a parameter $t \leq \ell(x)^c$ and derivation in t by Lemma 11.

This leads to an ODE of the form:

$$\mathbf{f}'(x, \mathbf{y}) = \overline{\mathbf{A}}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \overline{\mathbf{B}}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}).$$

by setting

$$\begin{aligned} \overline{\mathbf{A}}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) &= \mathbf{A}(\mathbf{f}(x, \mathbf{y}), \mathbf{h}(x, \mathbf{y}), x, \mathbf{y}) \\ \overline{\mathbf{B}}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) &= \mathbf{B}(\mathbf{f}(x, \mathbf{y}), \mathbf{h}(x, \mathbf{y}), x, \mathbf{y}) \end{aligned}$$

But then Lemma 12 applies, and we get precisely the conclusion, observing that the fact that the corresponding matrix $\overline{\mathbf{A}}$ and vector $\overline{\mathbf{B}}$ are essentially constant in $\mathbf{f}(x, \mathbf{y})$ guarantees hypotheses of Lemma 12. \square

7 A characterization of polynomial time

7.1 Register machines

A register machine program (a.k.a. goto program) is a finite sequence of ordered labeled instructions acting on a finite set of registers of one of the following type:

- increment the j th register R_j by the value of k th register R_k and go the next instruction:

$$R_j := R_j + R_k$$

- decrement the j th register R_j by the value of k th register R_k and go the next instruction:

$$R_j := R_j - R_k$$

- set the j th register R_j to integer k , for $\ell \in \{0, 1\}$ and go the next instruction:

$$R_j := k$$

- if register j is equal to 0, go to instruction p else go to next instruction.

$$\text{if } R_j = 0 \text{ goto } p$$

- halt the program: halt

In the following, since coping with negative numbers on classical models of computation can be done through simple encodings, we will not restrict ourself to non-negative numbers.

Definition 25 Let $t : \mathbb{N} \rightarrow \mathbb{N}$. A function $f : \mathbb{N}^p \rightarrow \mathbb{Z}$ is computable in time t by a register machine M with k registers if:

- when starting in initial configuration with registers $R_1, \dots, R_{\min(p,k)}$ set to $x_1, \dots, x_{\min(p,k)}$ and all other registers to 0 and
- starting on the first instruction (of label 0),

Machine M ends its computation after at most $t(\ell(\mathbf{x}))$ instructions where $\ell(\mathbf{x}) = \ell(x_1) + \dots + \ell(x_p)$ and with register R_0 containing $f(x_1, \dots, x_p)$.

A function is computable in polynomial time by M if there exists $c \in \mathbb{N}$ such that $t(\ell(\mathbf{x})) \leq \ell(\mathbf{x})^c$ for all $\mathbf{x} = (x_1, \dots, x_p)$.

The definition of register machines might look rudimentary however, the following is easy (but tedious) to prove for any reasonable encoding of integer by Turing machines.

Theorem 15 A function f from $\mathbb{N}^p \rightarrow \mathbb{Z}$ is computable in polynomial time on Turing machines iff it is computable in polynomial time on register machines.

7.2 A characterization of polynomial time

The above result shows that function defined by linear length-ODE from function computable in polynomial time, are indeed polynomial time. We are now ready to introduce a recursion scheme based on solving linear differential equation to capture polynomial time.

Remark 11 *Since the function we define take their values in \mathbb{N} and have output in \mathbb{Z} , composition is an issue. Instead of considering restrictions of these function with output in \mathbb{N} (which is always possible, even by syntactically expressible constraints), we simply admit that composition may not be defined in some cases.*

Definition 26 *Let \mathbb{DL} be the smallest subset of functions, that contains $\mathbf{0}$, $\mathbf{1}$, projections π_i^p , the length function $\ell(x)$, the addition function $x+y$, the subtraction function $x-y$, the multiplication function $x \times y$ (often denoted $x \cdot y$), the sign function $\text{sg}(x)$ and closed under composition (when defined) and linear length-ODE scheme.*

Remark 12 *As our results will show, the definition of \mathbb{DL} would remain the same by considering closure under any kind of \mathcal{L} -ODE with \mathcal{L} satisfying the hypothesis of Lemma 13.*

Example 11 *A number of natural functions are in \mathbb{DL} . the following result is immediate by inspection of the example from Section 3 and 6. Functions $2^{\ell(x)}$, $2^{\ell(x) \cdot \ell(y)}$, $\text{if}(x, y, z)$, $\text{suffix}(x, y)$, $\lfloor \sqrt{x} \rfloor$, $\lfloor \frac{x}{y} \rfloor$, $2^{\lfloor \sqrt{x} \rfloor}$ all belong to \mathbb{DL} .*

Theorem 16 $\mathbb{DL} = \mathbf{FPTIME}$

Proof: The inclusion $\mathbb{DL} \subseteq \mathbf{FPTIME}$ is a consequence of the fundamental observation proved in Lemma 13, on the fact that arithmetic operations that are allowed can be computed in polynomial time and that \mathbf{FPTIME} is closed under composition of functions.

We now prove that $\mathbf{FPTIME} \subseteq \mathbb{DL}$. Let $f : \mathbb{N}^p \rightarrow \mathbb{N}$ be computable in polynomial time and M a k registers machine that compute f in time $\ell(\mathbf{x})^c$ for some $c \in \mathbb{N}$. We first describe the computation of M by simultaneous recursion scheme on length for functions $R_0(t, \mathbf{x}), \dots, R_k(t, \mathbf{x})$ and $\text{inst}(t, \mathbf{x})$ that give, respectively, the values of each register and the label of the current instruction at time $\ell(t)$.

We start with an informal description of the characterization. Initializations of the functions are given by: $R_0(0, \mathbf{x}) = 0, R_1(0, \mathbf{x}) = x_1, \dots, R_p(0, \mathbf{x}) = x_p, R_{p+1}(0, \mathbf{x}) = \dots = R_k(0, \mathbf{x}) = 0$ et $\text{inst}(0, \mathbf{x}) = 0$. Let $m \in \mathbb{N}$ be the number of instructions of M and let $l \leq m$. Recall that, for a function f , $\frac{\partial f}{\partial L}(t, \mathbf{x})$ represents a manner to describe $f(t+1, \mathbf{x})$ from $f(t, \mathbf{x})$ when $L(t+1) = L(t) + 1$. We denote by, $\text{next}_l^f, \text{next}_l^h, h \leq k$, the evolution of the instruction function and of register R_h after applying instruction l at any such instant t . They are defined as follows:

- If instruction of label l if of the type $R_j := R_j + R_k$, then:
 - $\text{next}_l^I = 1$ since $\text{inst}(t + 1, \mathbf{x}) = \text{inst}(t, \mathbf{x}) + 1$
 - $\text{next}_l^j = R_k(t, \mathbf{x})$ since $R_j(t + 1, \mathbf{x}) = R_j(t, \mathbf{x}) + R_k(t, \mathbf{x})$
 - $\text{next}_l^h = 0$ since $R_h(t, \mathbf{x})$ does not change for $h \neq j$
- If instruction of label l if of the type $R_j := R_j - R_k$, then:
 - $\text{next}_l^I = 1$ since $\text{inst}(t + 1, \mathbf{x}) = \text{inst}(t, \mathbf{x}) + 1$
 - $\text{next}_l^j = -R_k(t, \mathbf{x})$ since $R_j(t + 1, \mathbf{x}) = R_j(t, \mathbf{x}) - R_k(t, \mathbf{x})$
 - $\text{next}_l^h = 0$ since $R_h(t, \mathbf{x})$ does not change for $h \neq j$
- If instruction of label l if of the type $R_j := \ell$, for $\ell \in \{0, 1\}$ then:
 - $\text{next}_l^I = 1$ since $\text{inst}(t + 1, \mathbf{x}) = \text{inst}(t, \mathbf{x}) + 1$
 - $\text{next}_l^j = \ell - R_j(t, \mathbf{x})$ since $R_j(t + 1, \mathbf{x}) = \ell$
 - $\text{next}_l^h = 0$ since $R_h(t, \mathbf{x})$ does not change for $h \neq j$
- If instruction of label l if of the type if $R_j = 0$ goto p , then:
 - $\text{next}_l^I = \text{if}(R_j(t, \mathbf{x}), p - \text{inst}(t, \mathbf{x}), 1)$ since, in case $R_j(t, \mathbf{x}) = 0$ instruction number goes from $\text{inst}(t, \mathbf{x})$ to p .
 - $\text{next}_l^h = 0$
- If instruction of label l if of the type **Halt**, then:
 - $\text{next}_l^I = 0$ since the machine stays in the same instruction when halting
 - $\text{next}_l^h = 0$.

The definition of function inst by derivation on length is now given by (we use a more readable "by case" presentation):

$$\frac{\partial \text{inst}}{\partial \ell}(t, \mathbf{x}) = \text{case} \begin{cases} \text{inst}(t, \mathbf{x}) = 1 & \text{next}_1^I \\ \text{inst}(t, \mathbf{x}) = 2 & \text{next}_2^I \\ \vdots \\ \text{inst}(t, \mathbf{x}) = m & \text{next}_m^I \end{cases}$$

Expanded as an arithmetic expression, this give:

$$\frac{\partial \text{inst}}{\partial \ell}(t, \mathbf{x}) = \sum_{l=0}^m \left(\prod_{i=0}^{l-1} \text{sg}(\text{inst}(t, \mathbf{x}) - i) \right) \cdot \bar{\text{sg}}(\text{inst}(t, \mathbf{x}) - l) \cdot \text{next}_l^I$$

Note that each next_l^I is an expression in terms of $\text{inst}(t, \mathbf{x})$ and, in some cases, in $\text{sg}(R_j(t, \mathbf{x}))$, too (for a conditional statement). Similarly, for each $j \leq k$:

$$\frac{\partial R_j}{\partial \ell}(t, \mathbf{x}) = \sum_{l=0}^m \left(\prod_{i=0}^{l-1} \text{sg}(\text{inst}(t, \mathbf{x}) - i) \right) \cdot \text{sg}(\text{inst}(t, \mathbf{x}) - l) \cdot \text{next}_l^j$$

It is easily seen that, in each of these expressions above, there is at most one occurrence of $\text{inst}(t, \mathbf{x})$ and $R_j(t, \mathbf{x})$ that is not under the scope of an essentially constant function (i.e. the sign functions). Hence, the expressions are of the prescribed form.

We know M works in time $\ell(\mathbf{x})^c$ for some fixed $c \in \mathbb{N}$. Both functions $\ell(\mathbf{x}) = \ell(x_1) + \dots + \ell(x_p)$ and $B(\mathbf{x}) = 2^{\ell(\mathbf{x}) \cdot \ell(\mathbf{x})}$ are in \mathbb{DL} . It is easily seen that : $\ell(\mathbf{x})^c \leq B^{(c)}(\ell(\mathbf{x}))$ where $B^{(c)}$ is the c -fold composition of function B .

We can conclude by setting $f(\mathbf{x}) = R_0(B^{(c)}(\max(\mathbf{x})), \mathbf{x})$. \square

The following normal form theorem can also be obtained (Compared to Definition 15, no function \mathbf{h} is allowed on the right hand side):

Definition 27 (Normal linear \mathcal{L} -ODE (NL-ODE)) *Functions \mathbf{f} are definable by a normal linear \mathcal{L} -ODE if it corresponds to the solution of \mathcal{L} -ODE $\frac{\partial \mathbf{f}(x, \mathbf{y})}{\partial \mathcal{E}} = \mathbf{u}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y})$ where \mathbf{u} is essentially linear in $\mathbf{f}(x, \mathbf{y})$.*

Definition 28 (SLL) *A function $\mathbf{f} : \mathbb{N}^k \rightarrow \mathbb{N}^k$ is in SLL if there exists $\mathbf{g} : \mathbb{N}^{k+1} \rightarrow \mathbb{N}$ and $h : \mathbb{N}^k \rightarrow \mathbb{N}$ such that:*

- g is solution of a normal linear length-ODE $\frac{\partial \mathbf{g}(x, \mathbf{y})}{\partial \ell(x)} = \mathbf{u}(\mathbf{g}(x, \mathbf{y}), x, \mathbf{y})$
- h is the solution of a single linear length-ODE
- and, for all $\mathbf{y} \in \mathbb{N}^k$: for some integer c .

From the proof of Theorem 16 the result below can be easily obtained. It expresses that composition need to be used only once as exemplified in the above definition.

Theorem 17 SLL = FPTIME

Proof: In the proof of Theorem 16, the definition of each function $\text{inst}, R_0, \dots, R_k$ are done through a linear system of SLL-ODE that uses only the basic arithmetic and sign functions. Composition is used only to bound the computation by $B^{(c)}(\max(\mathbf{x}))$, whose definition can be obtained through a simple length-ODE. \square

8 Further works

Previous ideas can be extended to provide a characterization of **FPSPACE** by considering random access machines (RAM) instead of register machines (see Appendix B.1 for definitions) with specific instructions sets. Depending on the set of basic operations allowed in the RAM model, polynomial time computation relates to very different complexity classes as witnessed by the following statements (see formal statement and proof of Theorem 31 in appendix):

1. A function $f : \mathbb{N}^k \rightarrow \mathbb{Z}$ is computable in polynomial time, i.e. is in **FPTIME**, iff it is computable in polynomial time on a $\{+, -\}$ -RAM with unit cost.
2. A function $f : \mathbb{N}^k \rightarrow \mathbb{Z}$ is computable in **FPSPACE** iff it is computable in polynomial time on a $\{+, -, \times, \div\}$ -RAM with unit cost.

Second item follows from the following arguments: It as been proved in [17], that a function f is in **FPSPACE** iff it is the difference of two functions $f_1, f_2 : \mathbb{N} \rightarrow \mathbb{N}$ in $\#\mathbf{PSPACE}$, the class of functions that counts the number of accepting computations of a non deterministic polynomial space Turing machine. It follows from the result from [3], that a function is computable in polynomial time on a $\{+, -, \times, \div\}$ -RAM if and only if it belongs to $\#\mathbf{PSPACE}$.

Using random access machines (RAM) instead of register machines, **FPSPACE** can be shown to correspond to functions of type $f(\mathbf{y}) = g_1(h(\mathbf{y}), \mathbf{y})$ where \mathbf{g} is defined as a specific class of polynomial length-ODE with substitutions, and conversely.

We leave this characterization for future work, as we believe that this statement can be improved to an even simpler statement, and scheme.

References

- [1] V. I. Arnold. *Ordinary Differential Equations*. MIT Press, 1978.
- [2] S. Bellantoni and S. Cook. A new recursion-theoretic characterization of the poly-time functions. *Computational Complexity*, 2:97–110, 1992.
- [3] Alberto Bertoni, Giancarlo Mauri, and Nicoletta Sabadini. A Characterization of the Class of Functions Computable in Polynomial Time on Random Access Machines. *STOC*, pages 168–176, 1981.
- [4] G. Birkhoff and G.-C. Rota. *Ordinary Differential Equations*. John Wiley & Sons, 4th edition, 1989.
- [5] O. Bournez, D. S. Graça, and A. Pouly. Polynomial Time corresponds to Solutions of Polynomial Ordinary Differential Equations of Polynomial Length. *Journal of the ACM*, 64(6):38:1–38:76, 2017. doi:10.1145/3127496.
- [6] O. Bournez and A. Pouly. A Survey on Analog Models of Computation. Technical report, May 2018. arXiv:1805.05729.
- [7] Olivier Bournez and Amaury Pouly. *Handbook of Computability and Complexity in Analysis*, chapter A Survey on Analog Models of Computation. Springer. To appear (arXiv version in [6]), 2018.
- [8] Manuel L. Campagnolo. *Computational Complexity of Real Valued Recursive Functions and Analog Circuits*. PhD thesis, Universidade Técnica de Lisboa, 2001.

- [9] Manuel L. Campagnolo, Christopher Moore, and José Félix Costa. An analog characterization of the Grzegorzczak hierarchy. *Journal of Complexity*, 18(4):977–1000, 2002.
- [10] P. Clote. Computational models and function algebras. In Edward R. Griffor, editor, *Handbook of Computability Theory*, pages 589–681. North-Holland, Amsterdam, 1998.
- [11] Peter Clote and Evangelos Kranakis. *Boolean functions and computation models*. Springer Science & Business Media, 2013.
- [12] A. Cobham. The intrinsic computational difficulty of functions. In Y. Bar-Hillel, editor, *Proceedings of the International Conference on Logic, Methodology, and Philosophy of Science*, pages 24–30. North-Holland, Amsterdam, 1962.
- [13] E. A. Coddington and N. Levinson. *Theory of Ordinary Differential Equations*. Mc-Graw-Hill, 1955.
- [14] Pieter Collins and Daniel S Graça. Effective computability of solutions of ordinary differential equations the thousand monkeys approach. *Electronic Notes in Theoretical Computer Science*, 221:103–114, 2008.
- [15] Heinz-Dieter Ebbinghaus and Jörg Flum. *Finite Model Theory*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1995.
- [16] R. Fagin. Generalized first-order spectra and polynomial-time recognizable sets. In R. M. Karp, editor, *Complexity in Computer Computations*, pages 43–73. American Mathematics Society, Providence R.I., 1974.
- [17] Matthias Galota and Heribert Vollmer. Functions computable in polynomial space. *Information and Computation*, 198(1):56–70, April 2005.
- [18] David Gleich. Finite calculus: A tutorial for solving nasty sums. *Stanford University*, 2005.
- [19] Ronald L Graham, Donald E Knuth, Oren Patashnik, and Stanley Liu. Concrete mathematics: a foundation for computer science. *Computers in Physics*, 3(5):106–107, 1989.
- [20] Y. Gurevich. Algebras of feasible functions. In *Twenty Fourth Symposium on Foundations of Computer Science*, pages 210–214. IEEE Computer Society Press, 1983.
- [21] N. Immerman. *Descriptive Complexity*. Springer, 1999.
- [22] FA Izadi, N Aliev, and G Bagirov. *Discrete Calculus by Analogy*. Bentham Science Publishers, 2009.
- [23] L. Kalmár. Egyzzerü példa eldönthetetlen aritmetikai problémára. *Mate és Fizikai Lapok*, 50:1–23, 1943.

- [24] A. Kawamura. Lipschitz continuous ordinary differential equations are polynomial-space complete. In *2009 24th Annual IEEE Conference on Computational Complexity*, pages 149–160. IEEE, 2009.
- [25] Ker-I Ko. On the computational complexity of ordinary differential equations. *Information and Control*, 58(1-3):157–194, July/August/September 1983.
- [26] Gustavo Lau. Discrete calculus. URL: <http://www.acm.ciens.ucv.ve/main/entrenamiento/material/DiscreteCalculus.pdf>.
- [27] D. Leivant. Intrinsic theories and computational complexity. In *LCC'94*, number 960 in *Lecture Notes in Computer Science*, pages 177–194, 1995.
- [28] D. Leivant and J-Y Marion. Lambda calculus characterizations of poly-time. *Fundamenta Informatica*, 19(1,2):167,184, September 1993.
- [29] Daniel Leivant and Jean-Yves Marion. Ramified recurrence and computational complexity II: substitution and poly-space. In L. Pacholski and J. Tiuryn, editors, *Computer Science Logic, 8th Workshop, CSL'94*, volume 933 of *Lecture Notes in Computer Science*, pages 369–380, Kazimierz, Poland, 1995. Springer.
- [30] Bruno Loff, José Félix Costa, and Jerzy Mycka. The new promise of analog computation. In *Computability in Europe 2007: Computation and Logic in the Real World.*, 2007.
- [31] Cristopher Moore. Recursion theory on the reals and continuous-time computation. *Theoretical Computer Science*, 162(1):23–44, 5 August 1996.
- [32] Jerzy Mycka and José Félix Costa. What lies beyond the mountains? Computational systems beyond the Turing limit. *European Association for Theoretical Computer Science Bulletin*, 85:181–189, February 2005.
- [33] Jerzy Mycka and José Félix Costa. The $P \neq NP$ conjecture in the context of real and complex analysis. *Journal of Complexity*, 22(2):287–303, 2006.
- [34] P. Odifreddi. *Classical Recursion Theory*, volume 125 of *Studies in Logic and the foundations of mathematics*. North-Holland, April 1992.
- [35] Amaury Pouly. *Continuous models of computation: from computability to complexity*. PhD thesis, Ecole Polytechnique and Unidersidade Do Algarve, Defended on July 6, 2015. 2015. <https://pastel.archives-ouvertes.fr/tel-01223284>, Prix de Thèse de l'Ecole Polytechnique 2016, Ackermann Award 2017.
- [36] H.E. Rose. *Subrecursion*. Oxford university press, 1984.
- [37] V. Sazonov. Polynomial computability and recursivity in finite domains. *Elektronische Informationsverarbeitung und Kybernetik*, 7:319–323, 1980.

- [38] David B Thompson. Subrecursiveness: Machine-independent notions of computability in restricted time and storage. *Mathematical Systems Theory*, 6(1-2):3–15, 1972.
- [39] Jan van Leeuwen, editor. *Handbook of Theoretical Computer Science (Vol. A): Algorithms and Complexity*. MIT Press, Cambridge, MA, USA, 1990.

A Discrete Calculus

The following text is based on [18, 22, 26]: We do so using intentionally some notations from continuous ODEs in order to help understanding to people familiar to classical continuous theory. We provide proofs for most of the statements, but some of the proofs are not repeated here, as they just follow from easy computations, or as they are classical and can be found in these references.

Discrete ODEs are basically usually intended to concern functions over the integers of type $\mathbf{f} : \mathbb{N}^p \rightarrow \mathbb{Z}^q$, but its statements and concepts considered in this section are also valid more generally for functions of type $\mathbf{f} : \mathbb{Z}^p \rightarrow \mathbb{Z}^q$, for some integers p, q , or even functions $\mathbf{f} : \mathbb{R}^p \rightarrow \mathbb{R}^q$.

The basic idea is to consider the following concept of derivative.

Definition 29 (Discrete Derivative) *The discrete derivative of $\mathbf{f}(x)$ is defined as $\Delta\mathbf{f}(x) = \mathbf{f}(x+1) - \mathbf{f}(x)$. We will also write in this article \mathbf{f}' for $\Delta\mathbf{f}(x)$ to help to understand statements with respect to their classical continuous counterparts.*

Remark 13 *The previous concept corresponds to the right derivative. We can write $\mathbf{f}'_r(x)$ or $\Delta^+\mathbf{f}(x)$ to emphasise that fact. A left derivative version could also be considered: This would correspond to $\Delta^-\mathbf{f}(x) = \mathbf{f}(x-1) - \mathbf{f}(x)$, that we will sometimes write $\mathbf{f}'_l(x)$.*

In the rest of this section, we will only talk about above right derivative, but all results could easily be adapted to deal with left derivative. Actually, left and right derivatives are related by the following observation:

Lemma 14 (Left vs Right Derivative) *When $x, x+1, x-1$ fall in the domain of \mathbf{f} , we always have:*

$$\begin{aligned}\mathbf{f}'_l(x+1) &= -\mathbf{f}'_r(x) \\ \mathbf{f}'_l(x) &= -\mathbf{f}'_r(x-1)\end{aligned}$$

A.1 Some basic statements

Theorem 18 (Linearity) *For any functions \mathbf{f} and \mathbf{g} , and constant \mathbf{c} :*

$$\begin{aligned}(\mathbf{f}(x) + \mathbf{g}(x))' &= \mathbf{f}'(x) + \mathbf{g}'(x) \\ (\mathbf{c} \cdot \mathbf{f}(x))' &= \mathbf{c} \cdot \mathbf{f}'(x)\end{aligned}$$

Theorem 19 (Inverse) *Consider $x \neq 0$.*

$$\left(\frac{1}{x}\right)' = -\frac{1}{x} \cdot \frac{1}{x+1}$$

Theorem 20 (Division) Assume $g(x) \neq 0$, and $g(x+1) \neq 0$.

$$\left(\frac{f}{g}\right)' = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)g(x+1)}$$

Theorem 21 (Product)

$$\begin{aligned} (\mathbf{f}(x)\mathbf{g}(x))' &= \mathbf{f}'(x)\mathbf{g}(x+1) + \mathbf{f}(x)\mathbf{g}'(x) \\ &= \mathbf{f}(x+1)\mathbf{g}'(x) + \mathbf{f}'(x)\mathbf{g}(x) \end{aligned}$$

A.2 The discrete integral and primitive

Definition 30 (Discrete Integral) Given some function $\mathbf{f}(x)$, we write $\int_a^b \mathbf{f}(x)\delta x$ as a synonym for

•

$$\int_a^b \mathbf{f}(x)\delta x = \sum_{x=a}^{x=b-1} \mathbf{f}(x)$$

when $a < b$, (pay attention to the fact that the bound is $b-1$ on right, and b on left)

• 0 when $a = b$,

• and, when $a > b$:

$$\int_a^b \mathbf{f}(x)\delta x = - \int_b^a \mathbf{f}(x)\delta x$$

The following holds from a basic computation (from telescope formula):

Theorem 22 (Fundamental Theorem of Finite Calculus) Let $\mathbf{F}(x)$ be some function. Then,

$$\int_a^b \mathbf{F}'(x)\delta x = \mathbf{F}(b) - \mathbf{F}(a).$$

As a consequence:

Definition 31 (Discrete Primitive) Let $\mathbf{f}(x)$ be some function, and \mathbf{C} some constant (of suitable dimension if \mathbf{f} is vectorial). Then the function

$$\mathbf{F}(x) = \mathbf{C} + \sum_{x=0}^{x-1} \mathbf{f}(x)$$

is such that $\mathbf{F}'(x) = \mathbf{f}(x)$ and $\mathbf{F}(0) = \mathbf{C}$. As expected, \mathbf{F} is called a primitive of $\mathbf{f}(x)$.

Corollary 4 Let $\mathbf{f}(x)$ be some function, and $\mathbf{F}(x)$ its primitive.

$$\mathbf{F}(b) - \mathbf{F}(a) = \int_a^b \mathbf{F}'(x) \delta x = \sum_{x=a}^{x=b-1} \mathbf{f}(x)$$

And:

$$\sum_{x=a}^{x=b} \mathbf{f}(x) = \mathbf{F}(b+1) - \mathbf{F}(a)$$

Remark 14 Recall that for classical continuous derivative if $F(x) = \int_{a(x)}^{b(x)} f(x, t) dt$ then

$$F'(x) = f(x, b(x))b'(x) - f(x, a(x))a'(x) + \int_{a(x)}^{b(x)} \frac{\partial f}{\partial x}(x, t) dt$$

This generalizes to the following:

Lemma 15 (Derivation of an integral with parameters) Consider

$$\mathbf{F}(x) = \int_{a(x)}^{b(x)} \mathbf{f}(x, t) \delta t.$$

Then

$$\mathbf{F}'(x) = \int_{a(x)}^{b(x)} \frac{\partial \mathbf{f}}{\partial x}(x, t) \delta t + \int_0^{-a'(x)} \mathbf{f}(x+1, a(x+1)+t) \delta t + \int_0^{b'(x)} \mathbf{f}(x+1, b(x)+t) \delta t$$

In particular, when $a(x) = a$ and $b(x) = b$ are constant functions,

$$\mathbf{F}'(x) = \int_a^b \frac{\partial \mathbf{f}}{\partial x}(x, t) \delta t$$

Proof:

$$\begin{aligned} \mathbf{F}(x+1) - \mathbf{F}(x) &= \sum_{t=a(x+1)}^{b(x+1)-1} \mathbf{f}(x+1, t) - \sum_{t=a(x)}^{b(x)-1} \mathbf{f}(x, t) \\ &= \sum_{t=a(x)}^{b(x)-1} (\mathbf{f}(x+1, t) - \mathbf{f}(x, t)) + \sum_{t=a(x+1)}^{t=a(x)-1} \mathbf{f}(x+1, t) + \sum_{t=b(x)}^{b(x+1)-1} \mathbf{f}(x+1, t) \\ &= \sum_{t=a(x)}^{b(x)-1} \frac{\partial \mathbf{f}}{\partial x}(x, t) + \sum_{t=a(x+1)}^{t=a(x)-1} \mathbf{f}(x+1, t) + \sum_{t=b(x)}^{b(x+1)-1} \mathbf{f}(x+1, t) \\ &= \sum_{t=a(x)}^{b(x)-1} \frac{\partial \mathbf{f}}{\partial x}(x, t) + \sum_{t=0}^{t=-a(x+1)+a(x)-1} \mathbf{f}(x+1, a(x+1)+t) \\ &\quad + \sum_{t=0}^{b(x+1)-b(x)-1} \mathbf{f}(x+1, b(x)+t) \end{aligned}$$

□

A.3 Integration by part

Theorem 23 (Integration by part)

$$\int_a^b \mathbf{u}(x)\mathbf{v}'(x)\delta x = [\mathbf{u}(x)\mathbf{v}(x)]_a^b - \int_a^b \mathbf{u}'(x)\mathbf{v}(x+1)\delta x$$

where $[\mathbf{u}(x)\mathbf{v}(x)]_a^b$ stands for $\mathbf{u}(b)\mathbf{v}(b) - \mathbf{u}(a)\mathbf{v}(a)$

Proof: Write $(\mathbf{u}(x)\mathbf{v}(x))' = \mathbf{u}(x)\mathbf{v}'(x) + \mathbf{u}'(x)\mathbf{v}(x+1)$, and hence $\mathbf{u}(x)\mathbf{v}'(x) = (\mathbf{u}(x)\mathbf{v}(x))' - \mathbf{u}'(x)\mathbf{v}(x+1)$. Then integrate. □

A.4 Derivative of a composition

The following can be established:

Theorem 24 (Derivative of $f \circ g$)

$$\mathbf{f}(g(x))' = \int_0^{g'(x)} \mathbf{f}'(g(x) + k)\delta k$$

Proof: Write

$$\begin{aligned} \mathbf{f}(g(x+1)) - \mathbf{f}(g(x)) &= \int_{g(x)}^{g(x+1)} \mathbf{f}'(t)\delta t \\ &= \int_0^{g(x+1)-g(x)} \mathbf{f}'(g(x) + k)\delta k \\ &= \int_0^{g'(x)} \mathbf{f}'(g(x) + k)\delta k \end{aligned} \quad (16)$$

□

A.5 Falling power

With analogy with the concept of derivative of a power, this is traditional to define (m stands for some natural integer).

Definition 32 (Falling power) *The expression x to the m falling is denoted by $x^{\underline{m}}$ (sometimes denoted by $(x)_m$) stands for*

$$x^{\underline{m}} = x \cdot (x-1) \cdot (x-2) \cdots (x-(m-1)).$$

This is motivated by the following observation:

Theorem 25 (Derivative of a falling power) *The discrete derivative of a falling power having exponent m is m times the next lowest falling power: That is*

$$(x^{\underline{m}})' = m \cdot x^{\underline{m-1}}$$

A.6 Exponential

Theorem 26 (Exponential c^x) *Let c be some positive constant. We have*

$$(c^x)' = (c - 1) \cdot c^x.$$

In particular

$$(2^x)' = 2^x.$$

More generally,

Theorem 27 (Exponential $c^{f(x)}$) *Let c be some positive constant. We have*
 $(c^{f(x)})' = (c^{f'(x)} - 1) \cdot c^{f(x)}.$

A.7 Falling Exponential

In a spirit similar to the falling power above, we propose to introduce the following concept. This seems not standard (as far as the authors know, but this seems to be of clear interest).

We assume $x \in \mathbb{N}$ in the following discussions.

Definition 33 (Falling exponential) *Given some function $\mathbf{U}(x)$, the expression \mathbf{U} to the falling exponential x , denoted by*

$$\bar{2}^{\mathbf{U}(x)} = (1 + \mathbf{U}'(x-1)) \cdots (1 + \mathbf{U}'(1)) \cdot (1 + \mathbf{U}'(0)) = \prod_{t=0}^{t=x-1} (1 + \mathbf{U}'(t)).$$

with the convention that $\prod_0^0 = \mathbf{id}$, where \mathbf{id} is the identity (e.g. 1 for the scalar case).

This is motivated by the following two observations:

Lemma 16 *For all $x \in \mathbb{Z}$, $2^x = \bar{2}^x$*

Theorem 28 (Derivative of a falling exponential) *The discrete derivative of a falling exponential is given by*

$$\left(\bar{2}^{\mathbf{U}(x)}\right)' = \mathbf{U}'(x) \cdot \bar{2}^{\mathbf{U}(x)}$$

In particular, we can easily build towers of exponentials using polynomial ordinary differential equations (ODEs):

A.8 Solving some particular ODEs

We will here consider the discrete variants of some particular (linear) ODEs.

Remark 15 *We assume implicitly in all this section that $x \in \mathbb{N}$, i.e. we discuss solutions of the ODEs over the domain \mathbb{N} .*

Remark 16 Recall that the solution of $\mathbf{f}'(x) = \mathbf{b}(x)$, $\mathbf{f}(0) = 0$ for classical continuous derivatives is given by

$$\mathbf{f}(x) = \int_0^x \mathbf{b}(t) dt.$$

Here we have something very similar:

Lemma 17 (Solution of ODE $\mathbf{f}'(x) = \mathbf{b}(x)$) The solution of $\mathbf{f}'(x) = \mathbf{b}(x)$, $\mathbf{f}(0) = \mathbf{0}$ is

$$\int_0^x \mathbf{b}(t) \delta t$$

Proof: Consider $\mathbf{f}(x) = \int_0^x \mathbf{b}(t) \delta t = \sum_{t=0}^{t=x-1} \mathbf{b}(t)$. For $x = 0$, we have $\mathbf{f}(x) = 0$. For $x > 0$, we have $\mathbf{f}'(x) = \mathbf{f}(x+1) - \mathbf{f}(x) = \mathbf{b}(x)$. \square

Remark 17 Recall that the solution of $f'(x) = a(x)f(x)$, $f(0) = 1$ (respectively: or more generally for the vectorial constant case $\mathbf{f}'(x) = \mathbf{A} \cdot \mathbf{f}(x)$, $\mathbf{f}(0) = \mathbf{1}$) for classical continuous derivatives is given by

$$f(x) = e^{\int_0^x a(t) dt}$$

(resp. $\mathbf{f}(x) = e^{t\mathbf{A}}$). Something very similar holds in the discrete setting:

Lemma 18 (Solution of ODE $\mathbf{f}'(x) = \mathbf{A}(x) \cdot \mathbf{f}(x)$) The solution of $\mathbf{f}'(x) = \mathbf{A}(x) \cdot \mathbf{f}(x)$ is

$$\bar{\int}_0^x \mathbf{A}(t) \delta t \cdot \mathbf{f}(0).$$

Notice that

$$\bar{\int}_0^x \mathbf{A}(t) \delta t = (1 + \mathbf{A}(x-1)) \cdots (1 + \mathbf{A}(1)) \cdot (1 + \mathbf{A}(0)) = \prod_{t=0}^{t=x-1} (1 + \mathbf{A}(t)).$$

Proof: Consider $\mathbf{f}(x) = \bar{\int}_0^x \mathbf{A}(t) \delta t \cdot \mathbf{f}(0)$. This values $\mathbf{f}(0)$ in 0. For $x > 0$, we have $\mathbf{f}(x+1) = (1 + \mathbf{A}(x)) \cdot \mathbf{f}(x)$ and hence $\mathbf{f}(x+1) - \mathbf{f}(x) = \mathbf{A}(x) \cdot \mathbf{f}(x)$. \square

A.9 Solving affine ODEs

We now go to affine (also called linear) ODE $\mathbf{f}'(x) = \mathbf{A}(x) \cdot \mathbf{f}(x) + \mathbf{B}(x)$. This affine ODEs play a key role in this article.

Remark 18 The solutions of $f'(x) = a(x) \cdot f(x) + b(x)$ for classical continuous derivatives are

$$f(x) = f(0)e^{\int_0^x a(t) dt} + \int_0^x b(u)e^{\int_u^x a(t) dt} du \quad (17)$$

Remark 19 It is usually obtained by variation of parameter method: we search $f(x)$ of the form $f(x) = f_1(x)k(x)$ where $f_1(x)$ is solution of $f_1'(x) = a(x)f_1(x)$ (from above discussions, we hence have $f_1(x) = f_1(0)e^{\int_0^x a(t)dt}$). Indeed, the trick is then that must have $f_1'(x)k(x) + f_1(x)k'(x) = a(x)f_1(x)k(x) + b(x)$: Factors of $k(x)$ cancels, and we get $f_1(x)k'(x) = b(x)$.

Multiplying by $e^{-\int_0^x a(t)dt}$ both sides, we get $e^{-\int_0^x a(t)dt}f_1(x)k'(x) = e^{-\int_0^x a(t)dt}b(x)$ which simplifies to $f_1(0)k'(x) = e^{-\int_0^x a(t)dt}b(x)$, equation in $f_1(0)k(x)$ than can be solved by a simple integral:

$$f_1(0)k(x) = f_1(0)k(0) + \int_0^x b(u)e^{-\int_0^u a(t)dt}du,$$

and then reporting the expression of $f_1(0)k(x)$

$$\begin{aligned} f(x) &= f_1(x)k(x) \\ &= f_1(0)e^{\int_0^x a(t)dt}k(x) \\ &= f_1(0)k(0)e^{\int_0^x a(t)dt} + e^{\int_0^x a(t)dt} \int_0^x b(u)e^{-\int_0^u a(t)dt}du \\ &= f_1(0)k(0)e^{\int_0^x a(t)dt} + \int_0^x b(u)e^{\int_u^x a(t)dt}du. \end{aligned}$$

Considering value in 0, we realize that $f_1(0)k(0)$ is actually $f(0)$ and obtain the above solution.

Remark 20 The solution (17) is the sum of a solution to $f'(x) = a(x)f(x)$, i.e. of the ODE with the non-linear term, and of a solution that values 0 in 0.

Remark 21 This extends for the vectorial case for classical continuous derivatives. This is usually obtained using the concept of resolvent: resolvent $\mathbf{R}(\mathbf{x}, \mathbf{x}_0)$ is by definition such that solutions of $\mathbf{f}'(x) = \mathbf{A}(x) \cdot \mathbf{f}(x)$ with $\mathbf{f}(0) = \mathbf{y}_0$ correspond to $\mathbf{f}(x) = \mathbf{R}(x, x_0) \cdot \mathbf{y}_0$.

In the case where $\mathbf{A}(x) = \mathbf{A}$ is constant, the resolvent is given by $\mathbf{R}(x, x_0) = e^{(x-x_0) \cdot \mathbf{A}}$.

The solutions of $\mathbf{f}'(x) = \mathbf{A}(x) \cdot \mathbf{f}(x) + \mathbf{B}(x)$ for classical continuous derivatives are then given by

$$\begin{aligned} \mathbf{f}(x) &= \mathbf{R}(x, 0) \cdot \mathbf{f}(0) + \mathbf{R}(x, 0) \cdot \int_0^x \mathbf{R}(0, u) \cdot \mathbf{b}(u)du \\ &= \mathbf{R}(x, 0) \cdot \mathbf{f}(0) + \int_0^x \mathbf{R}(x, u) \cdot \mathbf{b}(u)du \end{aligned} \quad (18)$$

in the general case.

In the discrete case, something similar holds. It is detailed below in the context of functions with several variables to be used, as it is, later.

Lemma 19 (Solution of ODE $\mathbf{f}'(x, \mathbf{y}) = \mathbf{A}(x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(x, \mathbf{y})$) For matrices \mathbf{A} and vectors \mathbf{B} and \mathbf{G} , the solution of equation $\mathbf{f}'(x, \mathbf{y}) = \mathbf{A}(x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(x, \mathbf{y})$ with initial conditions $\mathbf{f}(0, \mathbf{y}) = \mathbf{G}(\mathbf{y})$ is

$$\left(\overline{2}^{\int_0^x \mathbf{A}(t, \mathbf{y}) \delta t} \right) \cdot \mathbf{G}(\mathbf{y}) + \int_0^x \left(\overline{2}^{\int_{u+1}^x \mathbf{A}(t, \mathbf{y}) \delta t} \right) \cdot \mathbf{B}(u, \mathbf{y}) \delta u.$$

Notice that this can also be written:

$$\left(\prod_{t=0}^{t=x-1} (1 + \mathbf{A}(t, \mathbf{y})) \right) \cdot \mathbf{G}(\mathbf{y}) + \mathbf{B}(x-1, \mathbf{y}) + \sum_{u=0}^{x-2} \left(\prod_{t=u+1}^{x-1} (1 + \mathbf{A}(t, \mathbf{y})) \right) \cdot \mathbf{B}(u, \mathbf{y}).$$

This can also be expressed by simpler expression:

$$\sum_{u=-1}^{x-1} \left(\prod_{t=u+1}^{x-1} (1 + \mathbf{A}(t, \mathbf{y})) \right) \cdot \mathbf{B}(u, \mathbf{y}).$$

with the (not so usual) conventions that $\prod_x^{x-1} \kappa(x) = 1$ and $\mathbf{B}(-1, \mathbf{y}) = \mathbf{G}(\mathbf{y})$.

Before getting to the proof, we start by another Lemma:

Lemma 20 (Solution of $\mathbf{H}'(x) = \mathbf{A}(x) \cdot \mathbf{H}(x) + \mathbf{B}(x)$ with $\mathbf{H}(x) = 0$) *The solution of $\mathbf{H}'(x) = \mathbf{A}(x) \cdot \mathbf{H}(x) + \mathbf{B}(x)$ with $\mathbf{H}(x) = 0$ is given by*

$$\mathbf{H}(x) = \int_0^x \left(\bar{2}^{\int_{u+1}^x \mathbf{A}(t) \delta t} \right) \cdot \mathbf{B}(u) \delta u.$$

Proof: Consider above expression. We then have.

$$\begin{aligned} \mathbf{H}(0) &= 0 \text{ and from Lemma 15} \\ \mathbf{H}'(x) &= \int_0^x \mathbf{A}(x) \cdot \left(\bar{2}^{\int_{u+1}^x \mathbf{A}(t) \delta t} \right) \cdot \mathbf{B}(u) \delta u + \bar{2}^{\int_{x+1}^x \mathbf{A}(t) \delta t} \cdot \mathbf{B}(x) \\ &= \mathbf{A}(x) \cdot \mathbf{H}(x) + \mathbf{B}(x). \end{aligned}$$

□

We can now go to the proof of Lemma 2.

Proof: From linearity of derivation, we must have $\mathbf{f}_1(x, \mathbf{y}) = \mathbf{f}(x, \mathbf{y}) - \mathbf{H}(x, \mathbf{y})$ solution of $\mathbf{f}'_1(x, \mathbf{y}) = \mathbf{A}(x, \mathbf{y}) \cdot \mathbf{f}_1(x, \mathbf{y})$, where $\mathbf{H}(x, \mathbf{y})$ satisfies $\mathbf{H}(0, \mathbf{y}) = 0$ and $\mathbf{H}'(x, \mathbf{y}) = \mathbf{A}(x, \mathbf{y}) \cdot \mathbf{H}(x, \mathbf{y}) + \mathbf{B}(x, \mathbf{y})$. The solution of latter equation is given by Lemma 20.

A general solution f_1 of former equation is (see above)

$$\mathbf{f}_1(x, \mathbf{y}) = \bar{2}^{\int_0^x \mathbf{A}(t, \mathbf{y}) \delta t} \cdot \mathbf{f}_1(0, \mathbf{y}).$$

This leads to the above expression. □

Remark 22 *Fomula (4) can also be expressed by simpler expression:*

$$\sum_{u=-1}^{x-1} \left(\prod_{t=u+1}^{x-1} (1 + \mathbf{A}(t, \mathbf{y})) \right) \cdot \mathbf{B}(u, \mathbf{y}).$$

with the (not so usual) conventions that $\prod_x^{x-1} \kappa(x) = 1$ and $\mathbf{B}(-1, \mathbf{y}) = \mathbf{G}(\mathbf{y})$.

Exactly the same (first) proof shows that the following generalization holds:

Lemma 21 (Solution of ODE $\mathbf{f}'(x, \mathbf{y}) = \mathbf{A}(\mathbf{f}(x, \mathbf{y}), x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(f(x, \mathbf{y}), x, \mathbf{y})$)
For matrices \mathbf{A} and vectors \mathbf{B} and \mathbf{G} , the solution of equation $\mathbf{f}'(x, \mathbf{y}) = \mathbf{A}(f(x, \mathbf{y}), x, \mathbf{y}) \cdot \mathbf{f}(x, \mathbf{y}) + \mathbf{B}(f(x, \mathbf{y}), x, \mathbf{y})$ with initial conditions $\mathbf{f}(0, \mathbf{y}) = \mathbf{G}(\mathbf{y})$ satisfies

$$\mathbf{f}(x, \mathbf{y}) = \left(\overline{2}^{\int_0^x \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y}) \delta t} \right) \cdot \mathbf{G}(\mathbf{y}) + \int_0^x \left(\overline{2}^{\int_{u+1}^x \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y}) \delta t} \right) \cdot \mathbf{B}(\mathbf{f}(u, \mathbf{y}), u, \mathbf{y}) \delta u.$$

In an analog way to above, this can also be written:

$$\mathbf{f}(x, \mathbf{y}) = \left(\prod_{t=0}^{t=x-1} (1 + \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y})) \right) \cdot \mathbf{G}(\mathbf{y}) + \mathbf{B}(x-1, \mathbf{y}) + \sum_{u=0}^{x-2} \left(\prod_{t=u+1}^{x-1} (1 + \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y})) \right) \cdot \mathbf{B}(\mathbf{f}(u, \mathbf{y}), u, \mathbf{y}).$$

or as

$$\mathbf{f}(x, \mathbf{y}) = \sum_{u=-1}^{x-1} \left(\prod_{t=u+1}^{x-1} (1 + \mathbf{A}(\mathbf{f}(t, \mathbf{y}), t, \mathbf{y})) \right) \cdot \mathbf{B}(\mathbf{f}(u, \mathbf{y}), u, \mathbf{y}).$$

with the conventions that $\prod_x^{x-1} \kappa(x) = 1$ and $\mathbf{B}(\cdot, -1, \mathbf{y}) = \mathbf{G}(\mathbf{y})$.

A.10 Derivative of some particular functions

We now provide some other examples of functions with their derivative.

Theorem 29 (sin, cos) *We have:*

$$\begin{aligned} \sin(x)' &= 2 \cdot \sin\left(\frac{1}{2}\right) \cdot \cos\left(x + \frac{1}{2}\right) \\ \cos(x)' &= -2 \cdot \sin\left(\frac{1}{2}\right) \cdot \sin\left(x + \frac{1}{2}\right) \end{aligned}$$

Theorem 30 (tan) *Whenever $\cos(x) \neq 0$ and $\cos(x+1) \neq 0$, we have:*

$$\begin{aligned} \tan(x)' &= \frac{1}{2} \sin\left(\frac{1}{2}\right) \cos\left(\frac{1}{2}\right) \frac{1}{\cos(x) \cos(x+1)} \\ &= \tan(1) \cdot (1 + \tan(x) \tan(x+1)) \end{aligned}$$

B Review of some results and models

B.1 Random access machines

In the following we consider the random access machine with unit cost as computation model. Let Op be a set of arithmetic operations. A Op -RAM is the collection of a **potentially infinite** set of registers (R_i) where $i \in \mathbb{N}^*$ and two special registers A, B . A program is a finite sequence of ordered labeled instructions, I_0, \dots, I_r acting on registers of one of the following type:

1. $A := l, B := l, l \in \mathbb{N}$
2. $A := A \odot B$ or $B := A \odot B$, for $\odot \in Op$
3. $B := A, A := B$
4. $A := R_A$ meaning that A receive the content of the register whose address is in A , provided A is non negative (indirect addressing).
5. $R_A := B$ meaning that the register whose address is in (non negative) A receives the content of register B .
6. If $A = B$ then goto I_i else goto I_j (pour tout $i, j \leq r$).
7. halt

Definition 34 Let $t : \mathbb{N} \rightarrow \mathbb{N}$. A function $f : \mathbb{N}^p \rightarrow \mathbb{Z}$ is computable in time t by a RAM machine M if:

- when starting in initial configuration with registers R_1, \dots, R_p set to x_1, \dots, x_p and all other registers to 0 and
- starting on the first instruction (of label 0),

machine M ends its computation after at most $t(\ell(x))$ instructions where $\ell(x) = \ell(x_1) + \dots + \ell(x_p)$ and with register A containing $f(x_1, \dots, x_p)$.

A function is computable in polynomial time by M if there exists $c \in \mathbb{N}$ such that $t(\ell(x)) \leq \ell(x)^c$ for all $x = (x_1, \dots, x_p)$.

Depending on the set of basic operations allowed in the RAM model, polynomial time computation relates to very different complexity classes as witnessed by the following result.

Theorem 31 1. A function $f : \mathbb{N}^k \rightarrow \mathbb{Z}$ is computable in polynomial time, i.e. is in **FPTIME**, iff it is computable in polynomial time on a $\{+, -\}$ -RAM with unit cost.

2. A function $f : \mathbb{N}^k \rightarrow \mathbb{Z}$ is computable in **FPSPACE** iff it is computable in polynomial time on a $\{+, -, \times, \div\}$ -RAM with unit cost.

Proof: Let $f : \mathbb{N}^k \rightarrow \mathbb{Z}$, let $f_1, f_2 : \mathbb{N}^k \rightarrow \mathbb{N}$ defined by $f_1 = \max\{0, f\}$, $f_2 = \max\{0, -f\}$. Remark that $f = f_1 - f_2$. The following is easily seen, through a reasonable representation of integers (see, e.g., Definition 14) and a straightforward simulation of arithmetic operations: function f is computable in polynomial time on a $\{+, -\}$ -RAM (resp. $\{+, -, \times, \div\}$ -RAM) if and only if f_1 and f_2 are computable in polynomial time on a $\{+, -\}$ -RAM (resp. $\{+, -, \times, \div\}$ -RAM). From that, the first item follows easily by classical simulation between machine models [39].

It as been proved in [17], that a function f is in **FPSPACE** iff it is the difference of two functions $f_1, f_2 : \mathbb{N} \rightarrow \mathbb{N}$ in **#PSPACE**, the class of functions that counts the number of accepting computations of a non deterministic polynomial space Turing machine. We conclude using the early remarks of the proof and the result from [3], that a function is computable in polynomial time on a $\{+, -, \times, \div\}$ -RAM if and only if it belongs to **#PSPACE**. \square