

Nonreciprocal Directional Dichroism Induced by Quantum Metric Dipole

Yang Gao and Di Xiao

Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA

(Dated: March 21, 2022)

We identify the quantum metric dipole as the geometric origin of the nonreciprocal directional dichroism which describes the change in the refractive index upon reversing the light propagation direction. Specifically, we find that the static limit of the nonreciprocal directional dichroism corresponds to a quadrupolar transport current from the quantum metric dipole, in response to a quadrupolar electric field. Moreover, at finite frequency, we demonstrate that the steepest slope of the averaged quantum metric dipole determines a peak. Finally, we illustrate both features in a low-energy model.

Spatially dispersive optical effects can often yield incisive information on the structural and electronic properties of matter. A well-known example is natural optical activity in noncentrosymmetric materials [1]. If time-reversal symmetry is also broken, spatial dispersion can give rise to another phenomenon known as nonreciprocal directional dichroism (NDD) [2], referring to the difference in refractive index between counter-propagating lights. Due to its dependence on both broken time-reversal and inversion symmetry, NDD provides a powerful probe of the dynamical coupling between electricity and magnetism in matter [3–11].

To date, the microscopic understanding of NDD has been dominated by molecular theories of electromagnetic multipoles [12, 13]. Despite its wide adoption in the literature, this formalism cannot access physics associated with the geometric structure of Bloch states in the momentum space, whose importance has been made increasingly clear in recent years [11, 14, 15]. In particular, it has been shown that optical activity and the gyrotropic magnetic effect are connected to the Berry curvature and orbital magnetic moment, revealing their geometric origin [16–18].

In this Letter, we identify the quantum metric as the geometric origin of NDD. Specifically, NDD is connected to the first order moment of quantum metric, which is hence referred to as the quantum metric dipole. Using the semiclassical transport theory, we show that the integration of the quantum metric dipole over the Fermi surface yields a static current driven by a quadrupolar electric field. Such current is the DC counterpart of NDD and exists in metals with broken time-reversal and inversion symmetry. Moreover, the quantum metric can be interpreted as the quadrupole moment of the Bloch state. Our result thus provides an interesting dual to the role of orbital magnetic moment in optical activity: the dependence of the refractive index on light helicity is determined by dipole of orbital magnetic moment [17, 18], while that on propagation direction is determined by dipole of electric quadrupole moment.

We then use the linear response theory to show that NDD at finite frequency is also determined by the quantum metric dipole. We find that the steepest slope of

the averaged quantum metric dipole can give rise to a peak in the differential refractive index between counter-propagating lights. Finally, we illustrate both the static quadrupolar current and the peak structure of NDD in a low-energy minimal model. Our result shows that NDD can be used to probe the geometric structure of Bloch states, and also opens the door to band structure engineering of NDD.

Phenomenological theory of NDD.—We first give a brief account of the phenomenological theory of NDD in terms of the optical conductivity [19], along with some symmetry considerations. In medium, the light propagation is characterized by the refractive index n , which can be solved from the Maxwell equations. The wave equation for the electric field reads

$$\nabla \times (\nabla \times \mathbf{E}) = -\mu_0 \frac{\partial \mathbf{J}}{\partial t} - \frac{1}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2}, \quad (1)$$

where μ_0 is the vacuum permeability and c is the speed of light. Let us consider a monochromatic light polarized along the x -direction and propagating along the z -direction. The current \mathbf{J} is induced by the electric field through the conductivity tensor $\sigma_{ij}(\omega, \mathbf{q})$, $J_i(\omega, \mathbf{q}) = \sigma_{ij}(\omega, \mathbf{q}) E_j(\omega, \mathbf{q})$, with ω and \mathbf{q} being the frequency and wave vector of the light. If the spatial dispersion is weak, we can expand $\sigma_{ij}(\omega, \mathbf{q})$ in powers of \mathbf{q} ,

$$\sigma_{ij}(\omega, \mathbf{q}) = \sigma_{ij}(\omega, 0) + \sigma_{ijk}(\omega, 0) q_k + \dots \quad (2)$$

The derivation of n can be simplified by exerting the following symmetry constraints. First, we assume the mirror- z symmetry is broken but mirror- x symmetry is present. This forbids the existence of $\sigma_{xy}(\omega, \mathbf{q})$. We further assume that the system is rotationally invariant about the z -axis, so that $\sigma_{xxz} = \sigma_{yyz}$. By inserting the electric field profile $\mathbf{E} \propto e^{i\omega n_z z/c - i\omega t}$ in Eq. (1) and using the above symmetry assumptions, we obtain $n_z^2 = (n_0 + i\kappa_0)^2 + ic\mu_0 n_z \sigma_{xxz}$ with $n_0 + i\kappa_0 = (1 + i\mu_0 c^2 \sigma_{xx}/\omega)^{1/2}$. The solution reads

$$n_z = ic\mu_0 \sigma_{xxz}/2 + \sqrt{(n_0 + i\kappa_0)^2 - c^2 \mu_0^2 \sigma_{xxz}^2/4}. \quad (3)$$

If we reverse the propagation direction, the first term in Eq. (3) flips sign. Therefore, the difference in the refrac-




FIG. 1. The electric quadrupolar current. The electric field in red arrows has the quadrupolar profile. The resulting quadrupolar current is in purple arrows.

tive index is $\Delta n = n_z - n_{-z} = ic\mu_0\sigma_{xxz}(\omega, 0)$. Alternatively, we can write [20]

$$\Delta n = \frac{1}{2}ic\mu_0[\sigma_{xxz}(\omega, 0) - \sigma_{xxz}^*(-\omega^*, 0)]. \quad (4)$$

Here $\omega \rightarrow \omega + i\eta$ has a small imaginary part.

The symmetry requirement for NDD can be also seen by noting that Δn implies a current $J_x \propto \partial_z E_x$. Because J_x and $\partial_z E_x$ transform in the opposite way under time reversal, inversion, and mirror- z operations, Δn vanishes identically if any of the above symmetries is present.

Electric quadrupolar current.—To reveal the geometric origin of NDD, we first consider its static counterpart, i.e. a current driven by a spatially varying but static electric field in metals with the above symmetry constraints. As $\nabla \times \mathbf{E} = 0$, the electric field must be quadrupolar as shown in Fig. 1. We will adopt the semiclassical transport theory since the band geometry enters naturally in the semiclassical equations of motion. Below we sketch the derivation and leave the details in the Supplementary Materials [20]. Our starting point is the equations of motion in crystals under slowly changing electric field

$$\dot{\mathbf{r}} = \frac{1}{\hbar}\partial_{\mathbf{k}}\tilde{\varepsilon}_m - \dot{\mathbf{k}} \times \boldsymbol{\Omega}_m - \boldsymbol{\Omega}_{\mathbf{k}\mathbf{r},m} \cdot \dot{\mathbf{r}}, \quad (5)$$

$$\hbar\dot{\mathbf{k}} = -e\mathbf{E} + \boldsymbol{\Omega}_{\mathbf{r}\mathbf{k},m} \cdot \hbar\dot{\mathbf{k}}, \quad (6)$$

where $\boldsymbol{\Omega}_m = -2\text{Im}\langle\partial_{\mathbf{k}}u_m|\times|\partial_{\mathbf{k}}u_m\rangle$ is the momentum-space Berry curvature with $|u_m\rangle$ being the periodic part of Bloch functions in the m -th band, and $(\boldsymbol{\Omega}_{\mathbf{k}\mathbf{r},m})_{ij} = -2\text{Im}\langle\partial_{k_i}u_m|\partial_{r_j}u_m\rangle$ is the mixed Berry curvature with $(\boldsymbol{\Omega}_{\mathbf{k}\mathbf{r},m})_{ij} = -(\boldsymbol{\Omega}_{\mathbf{r}\mathbf{k},m})_{ji}$. The appearance of $\boldsymbol{\Omega}_{\mathbf{k}\mathbf{r},m}$ is due to the modification of the Bloch function by the inhomogeneous electric field.

The quantity of interest here is the band energy given by $\tilde{\varepsilon}_m = \varepsilon_m + \delta\varepsilon$, where ε_m is the unperturbed band energy and $\delta\varepsilon = \frac{1}{2}e(\partial_i E_j)g_{ij,m}$ is the correction to ε_m [20]. Here $g_{ij,m}$ is the quantum metric [21, 22]

$$g_{ij,m} = \text{Re}\langle\partial_{k_i}u_m|\partial_{k_j}u_m\rangle - A_{i,m}A_{j,m}, \quad (7)$$

where $A_{i,n} = \langle u_n|i\partial_{k_i}u_n\rangle$ is the intraband Berry connection. Geometrically, $g_{ij,m}$ measures the distance between neighboring Bloch states [21, 22]. As the quantum metric enters in the energy correction by coupling to $\partial_i E_j$, it can be viewed as the electric quadrupole of Bloch states, consistent with the definition of the electric quadrupole in electromagnetic theory.

Since the system considered here breaks both time-reversal and inversion symmetry, a linear magnetoelectric coupling is allowed. Consequently, a spatially varying electric field can induce a spatially varying magnetization, which gives rise to a magnetization current. It is well established that the magnetization current should be discounted and the transport current reads (Eq. (4) of Ref. [23])

$$\begin{aligned} \mathbf{J}^{\text{tr}} &= -e \sum_m \int \frac{d\mathbf{k}}{8\pi^3} D\dot{\mathbf{r}} f_m \\ &- \nabla_{\mathbf{r}} \times \frac{e}{\hbar} \sum_m \int \frac{d\mathbf{k}}{8\pi^3} k_B T \boldsymbol{\Omega}_m \log\left(1 + e^{\frac{\mu - \varepsilon_m}{k_B T}}\right), \end{aligned} \quad (8)$$

where f_m is the Fermi function, and $D = 1 + \text{Tr} \boldsymbol{\Omega}_{\mathbf{k}\mathbf{r},m}$ is modified density of states [24].

To evaluate Eq. (8) at the order $\partial\mathbf{E}$, we also need the response of the Berry curvature to the electric field, which reads $\boldsymbol{\Omega}'_m = 2e\nabla_{\mathbf{k}} \times \text{Re} \sum_{n \neq m} \mathbf{A}_{nm} (\mathbf{A}_{nm} \cdot \mathbf{E}) / \omega_{mn}$ [25], where $\omega_{mn} = \varepsilon_m - \varepsilon_n$ and $\mathbf{A}_{nm} = \langle u_m|i\partial_{\mathbf{k}}u_n\rangle$ is the interband Berry connection.

By plugging Eq. (5) and $\boldsymbol{\Omega}'_m$ into Eq. (8), we can obtain the transport current. We find that only the term containing $\partial_{\mathbf{k}}\delta\varepsilon$ remains and all other contributions cancel. We leave the detail in the Supplementary Materials [20]. The final result reads

$$\mathbf{J}_x^{\text{tr}} = -2e \int \frac{d\mathbf{k}}{8\pi^3} f_m \partial_{k_x} \delta\varepsilon = \gamma_{xxz} \partial_z E_x, \quad (9)$$

where

$$\gamma_{xxz} = \frac{e^2}{\hbar} \sum_m \int \frac{d\mathbf{k}}{8\pi^3} G_{xxz,m} f'_m, \quad (10)$$

Here $G_{ijk,m} = v_{i,m}g_{jk,m}$ with $v_{i,m}$ being the band velocity is the quantum metric dipole, defined similarly to the Berry curvature dipole [26]. The factor 2 in Eq. (9) appears because in the static case $\partial_z E_x = \partial_x E_z$. $f'_m = \partial f_m / \partial \varepsilon$. Since the quantum metric is the Bloch state quadrupole, $G_{ijk,m}$ can also be viewed as the momentum space dipole of the electric quadrupole.

Equation (9) is an intrinsic current independent of the transport relaxation time. It is also a Fermi surface effect and is hence important in metals and semiconductors. Since this current is in response to the variation of the electric field not the electric itself, it persists even if the net electric field across the whole sample is zero.

In addition to the electric quadrupole obtained here, Bloch states also have orbital magnetic moments that

couple to the magnetic field [15]. According to Ref. [17, 18], by substituting the electric quadrupole (i.e. quantum metric) in Eq. (10) with the orbital magnetic moment, we obtain the geometric part of the natural optical activity. Thus the dipoles of the quantum metric and orbital magnetic moment complete a full understanding of the geometric origin for distinct spatially dispersive optical phenomena in materials.

NDD at finite frequency.—We now reveal the geometric origin of NDD at finite frequency. For this purpose, we need to express Δn in Eq. (4) in terms of Bloch functions. We start from the standard Kubo formula of the optical conductivity

$$\sigma_{xx}(\omega, \mathbf{q}) = -\frac{e^2}{i\omega} \sum_{m,n} \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{(f_{m\mathbf{k}-\mathbf{q}} - f_{n\mathbf{k}})M_{ij}}{\varepsilon_{m\mathbf{k}-\mathbf{q}} - \varepsilon_{n\mathbf{k}} + \hbar\omega + i\eta}, \quad (11)$$

where $M_{ij} = \langle u_{m\mathbf{k}-\mathbf{q}} | \hat{v}_i | u_{n\mathbf{k}} \rangle \langle u_{n\mathbf{k}} | \hat{v}_j | u_{m\mathbf{k}-\mathbf{q}} \rangle$ and \hat{v} is the velocity operator. It is obvious that the photon wave vector \mathbf{q} shifts the crystal momentum and connects neighboring Bloch states in momentum space which are linked by the quantum metric. As such, the quantum metric should appear naturally in NDD.

To explicitly demonstrate the role of quantum metric, we expand $\sigma_{xx}(\omega, \mathbf{q})$ in Eq. (11) to the linear order of \mathbf{q} . The general expression is presented in the Supplemental Materials [20]. The result can be simplified by considering the optical transition between two bands. In this case we obtain [20]

$$\Delta n = -\frac{e^2 c \mu_0}{2\hbar} \sum_{m,n=c,v} \int \frac{d\mathbf{k}}{8\pi^3} \left[A(G_{zxx,m} f'_m + G_{zxx,n} f'_n) + B(G_{zxx,m} + G_{zxx,n}) + C(G_{xxz,m} + G_{xxz,n}) \right]. \quad (12)$$

In Eq. (12) we have explicitly separated the geometric contribution from the spectral contribution. The coefficients A , B and C only depend on the spectrum:

$$\begin{aligned} A &= \frac{\omega_{mn}^2}{\omega_{mn}^2 - \omega^2} + i\pi\omega\delta(\omega + \omega_{mn}), \\ B &= -\frac{2\Delta f_{mn}\omega_{mn}^3}{(\omega_{mn}^2 - \omega^2)^2} + i\pi\frac{\Delta f_{mn}\omega_{mn}^2}{\omega} \frac{d}{d\omega}\delta(\omega + \omega_{mn}), \\ C &= \frac{2\Delta f_{mn}\omega_{mn}}{\omega_{mn}^2 - \omega^2} - 2i\pi\Delta f_{mn}\delta(\omega + \omega_{mn}), \end{aligned}$$

where $\Delta f_{mn} = f_m - f_n$. The indices m and n run between the conduction (c) and valence (v) band. We note that in insulators the expression for $\text{Re } \Delta n$ is consistent with $\text{Im } \sigma_{xxz}$ in Ref. [27].

The quantum metric dipole G fully conforms to the symmetry requirement for NDD. g_{jk} is even under time reversal (T) and inversion (I) operations. Due to the appearance of the velocity, G_{ijk} is odd under T or I and even under combined TI . Moreover, both G_{xxz} and




FIG. 2. Energy spectrum near each valley. v' causes the conduction band bottom and the valence band top to shift oppositely along k_z , creating indirect band gaps. The energy difference at $k_z = 0$ yields the band gap parameter Δ_i .

G_{zxx} are odd under mirror- z symmetry. Therefore, Δn will vanish identically if T or I or mirror- z symmetry is present. Such behaviour is consistent with the previous symmetry analysis.

We note that it is tempting to take the $\omega \rightarrow 0$ limit of σ_{xxz} derived from Eq. (11) and compare with the semiclassical result. However, as we have already shown above, there are localized current contributions to σ_{xxz} that need to properly discounted. The details will be discussed elsewhere.

Armed with the insight that NDD has its geometrical origin in the quantum metric dipole, next we exam the peak structure of Δn as a function of ω . We will focus on $\text{Im } \Delta n$ which can be measured by light absorbance. In Eq. (12), the imaginary parts of the first and third term contain delta functions and hence experience a peak when the averaged quantum metric dipole reaches a maxima, e.g., when the joint density of state experiences the Van Hove singularity. However, the second term in Eq. (12) also contributes a peak of a different origin. It contains the derivative of the delta function and can be rewritten as

$$\frac{i\pi c \mu_0}{2\omega} \frac{d}{d\lambda} \langle G_{zxx,m} + G_{zxx,n} \rangle \Big|_{\lambda=-\omega}, \quad (13)$$

where $\langle x \rangle(\lambda) = \sum_{m,n=c,v} \int (d\mathbf{k}/8\pi^3) x \Delta f_{mn} \omega_{mn}^2 \delta(\omega_{mn} - \lambda)$ is the average of x for a fixed parameter λ . Equation (13) shows that $\text{Im } \Delta n$ will exhibit a peak when the averaged quantum metric dipole has the steepest slope. Due to the Kramers-Kronig relation, $\text{Re } \Delta n$ should also have a peak at the same frequency.

Low energy model.—To demonstrate the geometrical features of NDD, we consider the following Hamiltonian

$$\hat{H} = v'_i k_z + v\tau_i k_x \sigma_x + v k_y \sigma_y + \left(\Delta_i + \frac{k_z^2}{2m_i} \right) \sigma_z, \quad (14)$$

where i is the valley index, and for each valley $\tau_i = \pm 1$, $2\Delta_i$ is the band gap, v is the in-plane Fermi velocity,




FIG. 3. Sketch of the formation of a net quantum metric dipole. Streamline arrows and different colors show the direction and magnitude of the quantum metric dipole field, respectively. Although arrows are symmetric, the color map clearly has a dipolar structure, generating a net quantum metric dipole.

and $m_i > 0$ is the effective mass along the z -direction. The first term with v'_i introduces an anisotropy in the z -direction and hence effectively breaks the mirror- z symmetry for each valley. The energy spectrum around each valley is shown in Fig. 2. For fixed k_z , the corresponding 2D Hamiltonian is a gapped Dirac model that can be realized in layered two-dimensional magnets such as MnPX₃ ($X = \text{S, Se}$) monolayers [28, 29]. The k_z dependence can be introduced by stacking these 2D magnets along the z -direction and the v' term will appear when the stacking order breaks the mirror- z symmetry.

Since the valleys contribute additively, we shall limit our discussion to just one valley. The low energy model has a net quantum metric dipole in each valley. For illustration purpose, we consider the dipoles of g_{xx} for the valence band, which form a vector $(G_{xxx}, G_{yxx}, G_{zxx})$. After integrating this vector over k_x , we find that G_{xxx} vanishes and the remaining components form a vector in the k_y - k_z plane. In Fig. 3 we sketch such vector field using streamline arrows and color map to represent its direction and magnitude, respectively. We observe that the direction distribution is symmetric. However, the magnitude distribution is only symmetric about $k_y = 0$ axis. About $k_z = 0$ axis, it has a clear dipolar structure with a stronger hot spot in the lower-half plane due to v'_i .

We now quantitatively demonstrate the influence of the quantum metric dipole on Δn for a two-valley system. For each valley, we calculate the averaged quantum metric dipole $\langle G \rangle = \langle G_{zxx,m} + G_{zxx,n} \rangle$ with the chemical potential in the band gap. We find that $\langle G \rangle = \Delta^2 S(\omega/\Delta) \sqrt{mv'^2/2\Delta} \sqrt{\omega/(2\Delta) - 1}$ with $S(\omega/\Delta)$ being a dimensionless structure factor [20]. Clearly, the magnitude of $\langle G \rangle$ is proportional to the strength of the mirror- z symmetry breaking term, v' . Moreover, it has the steepest slope at $\omega = 2\Delta$. Consequently, according to Eq. (13),




FIG. 4. Δn (Panel a) and the electric quadrupolar current (Panel b) in the low-energy model. The parameters are chosen as follows: $\Delta_1 = \Delta$, $\Delta_2 = 2\Delta$, $v'_1 = -v'_2 = v'$, $\frac{1}{2}m(v')^2 = 0.95\Delta$. The system has a small global band gap 0.1Δ . In Panel a, $\langle G \rangle$ is in units of Δ^2 . In Panel b, γ_{xxz} is in units of $e^2/(16\pi^2\hbar)$.

Δn has a term proportional to $1/\sqrt{\omega/(2\Delta) - 1}$ and hence has a peak at $\omega = 2\Delta$ (the band edge).

In Fig. 4a we plot Δn and G as a function of ω with the Fermi energy lies in the band gap. Clearly there are two peaks. This is because according to above analysis, for each valley, the slope of $\langle G \rangle$ has a maxima at $\omega = 2\Delta_i$. Since $\Delta_1 \neq \Delta_2$, there will be two peaks corresponding to the optical transition from each valley. If the model has T or I symmetry, then $\Delta_2 = \Delta_1$ and $v'_1 = -v'_2$. The two peaks will appear at the same position with opposite values. Therefore, they will cancel each other and Δn vanishes.

This low-energy model also allows the static electric quadrupolar current. In Fig. 4b, we calculate the relevant conductivity γ_{xxz} at zero temperature. We find that $\gamma_{xxz} = 0$ when μ lies in the global band gap. As μ reaches the lower band bottom but still falls inside the gap of the other valley, γ_{xxz} increases with the density of states. When μ further increases, the other valley will also contribute but with opposite signs. Hence γ_{xxz} starts decreasing. When μ is large enough, γ_{xxz} approaches 0, as the quantum metric dipole decays faster than the increasing density of states.

In summary, we have identified the quantum metric dipole as the geometric origin of NDD. The static counterpart of NDD is a current driven by a quadrupolar electric field, obtained by integrating the quantum metric dipole over the Fermi surface. Moreover, the steepest slope of the averaged quantum metric dipole yields a peak in NDD, which could be useful for band structure engineering of NDD.

We acknowledge useful discussions with Kenneth Burch, Nuh Gedik, Ajit Srivastava, Haidan Wen, Xiaodong Xu and Qi Zhang. This work is supported by the Department of Energy, Basic Energy Sciences, Grant No. DE-SC0012509.

-
- [1] L. Landau, E. Lifshitz, and L. Pitaevskii, *Electrodynamics of Continuous Media* (Pergamon Press, New York, 1984).
- [2] R. Fuchs, *Phil. Mag.* **11**, 647 (1965).
- [3] J. Goulon, A. Rogalev, C. Goulon-Ginet, G. Benayoun, and L. Paolasini, *Phys. Rev. Lett.* **85**, 4385 (2000).
- [4] M. Kubota, T. Arima, Y. Kaneko, J. He, X. Yu, and Y. Tokura, *Phys. Rev. Lett.* **92**, 137401 (2004).
- [5] T. Arima, *J. Phys. Condens. Matter* **20**, 434211 (2008).
- [6] I. Kézsmárki, N. Kida, H. Murakawa, S. Bordács, Y. Onose, and Y. Tokura, *Phys. Rev. Lett.* **106**, 057403 (2011).
- [7] Y. Takahashi, R. Shimano, Y. Kaneko, H. Murakawa, and Y. Tokura, *Nat. Phys* **8**, 121 (2011).
- [8] Y. Okamura, F. Kagawa, M. Mochizuki, M. Kubota, S. Seki, S. Ishiwata, M. Kawasaki, Y. Onose, and Y. Tokura, *Nat. Commun.* **4**, 2391 (2013).
- [9] I. Kézsmárki, D. Szaller, S. Bordács, V. Kocsis, Y. Tokunaga, Y. Taguchi, H. Murakawa, Y. Tokura, H. Engelkamp, T. Rößm, and U. Nagel, *Nat. Commun.* **5**, 3203 (2014).
- [10] S. Toyoda, N. Abe, S. Kimura, Y. Matsuda, T. Nomura, A. Ikeda, S. Takeyama, and T. Arima, *Phys. Rev. Lett.* **115**, 267207 (2015).
- [11] Y. Tokura and N. Nagaosa, *Nat. Commun.* **9**, 3740 (2018).
- [12] E. B. Graham and R. E. Raab, *Phil. Mag. B* **66**, 269 (1992).
- [13] L. D. Barron, *Molecular Light Scattering and Optical Activity* (Cambridge University Press, Cambridge, 2004).
- [14] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, *Rev. Mod. Phys* **82**, 1539 (2010).
- [15] D. Xiao, M.-C. Chang, and Q. Niu, *Rev. Mod. Phys.* **82**, 1959 (2010).
- [16] S. Zhong, J. Orenstein, and J. E. Moore, *Phys. Rev. Lett.* **115**, 117403 (2015).
- [17] J. Ma and D. A. Pesin, *Phys. Rev. B* **92**, 235205 (2015).
- [18] S. Zhong, J. E. Moore, and I. Souza, *Phys. Rev. Lett.* **116**, 077201 (2016).
- [19] R. M. Hornreich and S. Shtrikman, *Phys. Rev.* **171**, 1065 (1968).
- [20] Supplemental materials..
- [21] J. P. Provost and G. Vallee, *Comm. Math. Phys.* **76**, 289 (1980).
- [22] R. Resta, *Eur. Phys. J. B* **79**, 121 (2011).
- [23] D. Xiao, Y. Yao, Z. Fang, and Q. Niu, *Phys. Rev. Lett.* **97**, 026603 (2006).
- [24] D. Xiao, J. Shi, and Q. Niu, *Phys. Rev. Lett.* **95**, 137204 (2005).
- [25] Y. Gao, S. A. Yang, and Q. Niu, *Phys. Rev. Lett.* **112**, 166601 (2014).
- [26] I. Sodemann and L. Fu, *Phys. Rev. Lett.* **115**, 216806 (2015).
- [27] A. Malashevich and I. Souza, *Phys. Rev. B* **82**, 245118 (2010).
- [28] X. Li, T. Cao, Q. Niu, J. Shi, and J. Feng, *Proc. Natl. Acad. Sci. U. S. A.* **110**, 3738 (2013).
- [29] N. Sivadas, S. Okamoto, and D. Xiao, *Phys. Rev. Lett.* **117**, 267203 (2016).