Counting homomorphisms in plain exponential time

Amineh Dadsetan[∗] and Andrei A. Bulatov†‡

October 9, 2018

Abstract

In the counting Graph Homomorphism problem $(\#$ GRAPHHOM) the question is: Given graphs G, H , find the number of homomorphisms from G to H. This problem is generally $#P$ -complete, moreover, Cygan et al. proved that unless the ETH is false there is no algorithm that solves this problem in time $O(|V(H)|^{o(|V(G)|)})$. This, however, does not rule out the possibility that faster algorithms exist for restricted problems of this kind. Wahlström proved that $\#\text{GraphHom}$ can be solved in plain exponential time, that is, in time $k^{|V(G)|} \text{poly}(|V(H)|, |V(G)|)$ provided H has clique width k . We generalize this result to a larger class of graphs, and also identify several other graph classes that admit a plain exponential algorithm for $\#$ GRAPHHOM.

1 Introduction

The Exponential Time Hypothesis (ETH) [\[12\]](#page-31-0) essentially suggests that the Satisfiability problem does not admit an algorithm that is significantly faster than the straightforward brute force algorithm. The ETH has been widely used to obtain (conditional) lower bounds on the complexity of various problems, see [\[14\]](#page-31-1) for a fairly recent survey. It however does not forbid nontrivial algorithms for many other hard problems.

One of such problems is the Graph Homomorphism problem (Graph-HOM for short). A homomorphism from a graph G to a graph H is a mapping $\varphi: V(G) \to V(H)$ such that for any edge $ab \in E(G)$ the pair $\varphi(a)\varphi(b)$ is an edge of H . GRAPHHOM asks, given graphs G and H , whether or not there exists a homomorphism from G to H [\[10\]](#page-31-2). In the counting version of this problem, denoted $\#\text{GraphHOM}$, the goal is to find the number of homomorphisms from G to H . These two problems can be solved just by checking all possible mappings from a given graph G to a given graph H, which takes time $O^*(|V(H)|^{|V(G)|})$,

[∗]amineh.dadsetan@gmail.com

[†]abulatov@sfu.ca

[‡]This work was supported by an NSERC Discovery grant

where O^* denotes asymptotics up to a polynomial factor. Assuming the ETH Cygan et al. [\[4\]](#page-30-0) proved that the general GRAPHHOM and therefore $\#\text{Graph}$ -HOM cannot be solved in time $|V(H)|^{o(|V(G)|)}$. Related hardness results have also been obtained in [\[16,](#page-31-3) [3\]](#page-30-1).

In spite of this result, there are several ways to restrict GraphHom which sometimes result in a problem admitting a faster algorithm. For graph classes $\mathcal{G}, \mathcal{H}, \mathsf{GRAPHHom}(\mathcal{G}, \mathcal{H})$ denotes the problem GRAPHHOM in which the input graphs G, H belong to \mathcal{G}, \mathcal{H} , respectively. $\#\text{GRAPHIOM}$ can be restricted in the same way. Both problems have received much attention in their own rights and as a special case of a more general Constraint Satisfaction Problem, and much is known about their computational complexity. In particular, it is known when GRAPHHOM(–, H) [\[11\]](#page-31-4) and $\#\text{GraphHom}(-, \mathcal{H})$ [\[5\]](#page-30-2) are solvable in polynomial time and when they are NP- or #P-complete. Symbol − here means that an input graph is not restricted. Similarly, it is known when $\text{GRAPHHom}(\mathcal{G}, -)$ [\[9\]](#page-31-5) and $\#\text{GraphHom}(\mathcal{G},-)$ are solvable in polynomial time.

Here we are interested in such restrictions that give rise to problems solvable still in exponential time but much faster than brute force. Specifically, GRAPHHOM(G, H) or $\#$ GRAPHHOM(G, H) is said to be solvable in *plain exponential* time if there is a solution algorithm running in time $O^*(c^{|V(G)|+|V(H)|}),$ where c is a constant. If the problem $\#\text{GRAPHHom}(-, \mathcal{H})$ is solvable in plain exponential time, the class H is said to be a *plain exponential class*.

The most well known plain exponential class of graphs is K , the class of all cliques. Note that $\#\text{GraphHom}(-, \mathcal{K})$ is equivalent to the $\#\text{Graph Coulon}$ -ING problem, in which the problem is, given a graph G and a number k , to find the number of k -colourings of G . A fairly straightforward dynamic programming algorithm solves this problem in time $O^*(3^{|V(G)|})$; we outline this algorithm in Example [6.](#page-3-0) A more sophisticated algorithm [\[13\]](#page-31-6) solves this problem in time $O^*(2^{|V(G)|})$. If H is a class of graphs of tree width k then $\#\text{GraphHom}(-, \mathcal{H})$ is solvable in time $O^*(k+3)^{|V(G)|}$, see, [\[8\]](#page-31-7). For the class \mathcal{D}_c of graphs of degree at most c the problems $\#\text{GraphHom}(\mathcal{D}_c)$ and $\#\text{GraphHom}(\mathcal{D}_c)$ can be solved in time $O^*(c^{|V(G)|})$ by a minor modification of the brute force enumer-ation algorithm. Finally, Wahlström [\[17\]](#page-31-8) obtained probably the most general result so far on plain exponential graph classes, proving that if $\mathcal H$ only contains graphs of clique width k then $\#\text{GraphHom}(-, \mathcal{H})$ can be solved in time $O^*((2k+1)^{|V(G)|+|V(H)|})$. The algorithm from [\[17\]](#page-31-8) is also dynamic programming and uses the representation of (labeled) graphs of bounded clique width through a sequence of operations such as disjoint union, connecting vertices with certain labels, and relabelling vertices (k*-expressions*).

In this paper we further expand the class of graphs for which plain exponential counting algorithms are possible by adding one more operation to the construction of graphs of bounded clique width. The new class of graphs includes families of graphs of unbounded clique width, for instance, hypercubes, and therefore is strictly larger than the class of graphs of bounded clique width. By means of this new set of operations one can define a new graph 'width' measure that we call *extended clique width*. Graphs of extended clique width at most k can also be represented by *extended* k-*expressions*. Let \mathcal{X}_k denote the class of graph of extended clique width at most k.

We then show that given an arbitrary graph G , a graph H of extended clique width k, and an extended k-expression Φ representing H, the number hom (G, H) of homomorphisms from G to H can be found in time $O^*((2k)^{|G|})$. Similar to [\[17\]](#page-31-8), the algorithm is dynamic programming and iteratively computes numbers hom (G', H') , where G' is an induced subgraph of G and H' is a graph represented by a subexpression of Φ . Clearly, as one cannot assume that an extended k -expression representing H is known in advance, this algorithm alone does not guarantee that \mathcal{X}_k is plain exponential. However, we also show that given a graph H of extended width at most k , an extended k -expression representing H can be found in time $O^*((2k)^{|H|})$. Combined with the previous result we thus obtain the following

Theorem 1. For any fixed k the class \mathcal{X}_k is plain exponential.

Apart from graphs of bounded extended clique width we identify two less general plain exponential classes of graphs. The first one consists of subdivisions of cliques: Let S be a class of graphs, then $\mathcal{K}(S)$ denotes the class of graphs H obtained as follows. Take $H' \in \mathcal{S}$, a clique on vertices $\{v_1, \ldots, v_n\}$, and for any edge $v_i v_j$ of the clique, $i \neq j$, replace this edge with a copy of H', that is, connect v_i, v_j to all vertices of H' and include all the edges of H' .

Theorem 2. For any plain exponential class S of graphs, the class $K(S)$ is also *plain exponential.*

The second class consists of well studied Kneser graphs: \mathcal{KG}_k is the class of graphs, whose vertices are the k -element subsets of a certain set, and two vertices are connected if and only if the corresponding subsets are disjoint.

Theorem 3. For every k the class \mathcal{KG}_k is plain exponential.

2 (Extended) Clique width

2.1 Homomorphisms, plain exponential time

As always we denote the vertex set of a graph G by $V(G)$, and its edge set by $E(G)$. A *homomorphism* of a graph G to a graph H is a mapping $\varphi: V(G) \to$ $V(H)$ such that $\varphi(u)\varphi(v) \in E(H)$ for any $uv \in E(G)$. The *Counting Graph Homomorphism* problem $\#\text{GRAPHHOM}$ is defined as follows: given graphs G, H , find the number of homomorphisms from G to H . Its decision version $$ does there exist a homomorphism from G to H — is denoted by GRAPHHOM. Graph homomorphisms and the related combinatorial problems have been ex-tensively studied [\[10\]](#page-31-2). If H is allowed only from a class $\mathcal H$ of graphs, the resulting counting and decision problems are denoted $\#\text{GraphHom}(-, \mathcal{H})$ and $GRAPHHOM(-, \mathcal{H})$, respectively.

We will be concerned with the complexity and the best running time of algorithms for $\#\text{GraphHom}(-, \mathcal{H})$. In particular, we say that a class H of graphs is *plain exponential* if there is an algorithm that solves the problem $\#\text{GraphHom}(-, \mathcal{H})$ in *plain exponential time*: there exists a constant c such that on input $G, H, H \in \mathcal{H}$, the algorithm runs in time $O^*(c^{|V(G)|+|V(H)|}),$ where O^* means asymptotics up to a factor polynomial in $|V(G)|, |V(H)|$. Note that we will always assume that G and H are connected, since otherwise the existence or the number of homomorphisms from G to H can be deduced from those of their connected components.

Example 4. (*H*-COLOURING.) If H consists of just one graph, H , the problems $\#\text{GraphHom}(-, \mathcal{H})$, GRAPHHOM(-, H) are known as $\#H\text{-}\text{ColoURING}$ and H -COLOURING, respectively. The $#H$ -COLOURING problem is solvable in polynomial time if H is a complete graph with all loops present, or is a complete bipartite graph $[5]$. The H-COLOURING problem is solvable in polynomial time if H contains a loop or is bipartite [\[11\]](#page-31-4). Otherwise these problems are $\#P$ - and NP-complete, respectively. Since the brute force algorithm for this problems runs in $O(|V(H)|^{|V(G)|})$ time, #H-COLOURING and H-COLOURING are always solvable in plain exponential time. Also, by inspecting the solution algorithms from [\[5,](#page-30-2) [11\]](#page-31-4) these results can be slightly generalized: $\#\text{GRAPHHOM}(-, \mathcal{H})$ is solvable in polynomial time whenever every graph from $\mathcal H$ is a complete graph with all loops, or a complete bipartite graph. Similarly $\text{GRAPHHOM}(-, \mathcal{H})$ is polynomial time solvable if every graph from H contains a loop or is bipartite.

Example 5. (Graphs of bounded degree.) As is mentioned in the introduction, if the degrees of graphs from H are bounded by a number c, the (improved) brute force algorithm solves $\#\text{GraphHom}(-, \mathcal{H})$, GRAPHHOM $(-, \mathcal{H})$. Let G, H be input graphs, $H \in \mathcal{H}$. We assume G is connected; otherwise the procedure below has to be performed for each connected component, and the results multiplied. Order the vertices v_1, \ldots, v_n of G in such a way that each vertex except for the first one is adjacent to one of the preceding vertices. Then the brute force algorithm is organized as follows: Assign images to v_1, \ldots, v_n in turn. There are |H| possibilities to map v_1 , but then if v_i is adjacent to v_j , $j < i$, the image of v_j is fixed, and therefore there are at most c possibilities for the image of v_i . Thus, the algorithm runs in $O^*(c^n)$. This approach also allows H to have bounded number of vertices of high degree.

Example 6. (Graphs of bounded clique width.) Let \mathcal{C}_k denote the class of all graphs of clique width at most k (to be defined in Section [2.2\)](#page-4-0). Then #GRAPHHOM(-, \mathcal{C}_k), GRAPHHOM(-, \mathcal{C}_k) can be solved in time $O^*((2k+1)^{|V(G)|+|V(H)|})$, implying that \mathcal{C}_k is plain exponential [\[17\]](#page-31-8).

Here we briefly describe the simple algorithm solving $\#\text{GraphHom}(-, \mathcal{K})$, where K is the class of cliques. Given a graph G and a number s (or, equivalently, the clique K_s) the solution algorithm maintains an array $N(S, \ell)$ for $S \subseteq V$ and $\ell \leq s$, which contains the number of homomorphisms from the subgraph of G induced by S to an ℓ -element clique. To compute each $N(S, \ell)$ we go over all subsets $S' \subseteq S$, consider the vertices from S' to be mapped to the ℓ -th vertex of the ℓ -clique. Then there are $N(S-S', \ell-1)$ ways to map the remaining vertices, and $N(S, \ell)$ is the sum of all numbers like this. It is not hard to see that the running time of this algorithm is $O^*(3^{|V(G)|})$. It can be improved to run in time $O[*](2^{|G|})$ [\[13\]](#page-31-6), and some further improvements are possible in certain cases [\[6\]](#page-31-9).

We will often deal with vertex labeled graphs. It will be convenient to represent labels on vertices of a graph G as a *label function* $\pi: V(G) \to [k]$ ([k] = $\{1,\ldots,k\}$, in which case we say that G is k-labeled. Graph $G = (V, E)$ equipped with a label function π will be denoted by $\mathbb{G} = (V, E, \pi)$. The k-labeled graph G is then called a k-labelling of G. Let $\mathbb{G}_1 = (V_1, E_1, \pi_1)$ and $\mathbb{G}_2 = (V_2, E_2, \pi_2)$ are k-labeled graph. A mapping $\varphi: V_1 \to V_2$ is a *homomorphism* of k-labeled graph \mathbb{G}_1 to k-labeled graph \mathbb{G}_2 if it is a homomorphism of graph $G_1 = (V_1, E_1)$ to $G_2 = (V_2, E_2)$ respecting the labelling, that is, $\pi_2(\varphi(v)) = \pi_1(v)$ for every $v \in V_1$.

The following notation will also be useful. Let again $\mathbb{G}_1, \mathbb{G}_2$ be k-labeled graphs, such that V_1, V_2 are disjoint. Then $\mathbb{G}_1 \bigoplus \mathbb{G}_2 = (V_1 \uplus V_2, E_1 \uplus E_2, \pi_1 \uplus \pi_2),$ where

$$
\pi_1 \uplus \pi_2(v) = \begin{cases} \pi_1(v), & \text{if } v \in V_1, \\ \pi_2(v), & \text{if } v \in V_2. \end{cases}
$$

Finally, the subgraph of a graph $G = (V, E)$ induced by a set $S \subseteq V$ is denoted by G[S]. For a k-labeled graph $\mathbb{G} = (V, E, \pi)$, by $\mathbb{G}[S]$ we denote the k-labeled subgraph induced by $S \subseteq V$. Note that the labelling function of $\mathbb{G}[S]$ is $\pi|_S$, i.e., the restriction of π on the set S.

2.2 Clique width and k-expressions

The simplest way to introduce clique width of a graph is through k-expressions.

Definition 7. *The following operators are defined on* k*-labeled graphs.*

- \cdot_i *: Construct a graph with one vertex, which is labeled* $i \in [k]$ *.*
- \bullet $\rho_{i\rightarrow j}(\mathbb{G})$: Relabel all vertices with label $i \in [k]$ of a k-labeled graph \mathbb{G} to *label* $j \in [k]$ *.*
- $\eta_{ij}(\mathbb{G})$, for $i \neq j$: Add edges from every vertex labeled i to every vertex *labeled* j *in* G*, i.e. add edges* uv *for any vertices* u, v *where* u *has label* i *and* v *has label* j*.*
- $\mathbb{G}_1 \bigoplus \mathbb{G}_2$: The disjoint union of k-labeled graphs \mathbb{G}_1 and \mathbb{G}_2 .

A k-expression *is any (properly formed) formula using the above operators.*

Every k*-expression represents a* k*-labeled graph. We say that a graph* G = (V, E) *is represented by k-expression* Φ *, if there exists a k-labelling* π *of the vertices of* G *such that* Φ *represents* $\mathbb{G} = (V, E, \pi)$ *. A graph has* clique width k *if it is represented by a* k*-expression. The class of all graphs of clique width* k *is denoted by* \mathcal{C}_k *.*

Wahlström in [\[17\]](#page-31-8) used k-expressions of graphs to show that \mathcal{C}_k is plain exponential. However, k -expressions suitable for his plain exponential algorithm must satisfy an extra condition. Let Φ be a k-expression representing a k-labeled graph \mathbb{G} . Note that any subformula of Φ represents a subgraph of \mathbb{G} . We say that k-expression Φ is *safe* if for every its subexpression $\Phi_1 \bigoplus \Phi_2$ such that Φ_1, Φ_2 represent graphs $\mathbb{G}_1, \mathbb{G}_2$, respectively, the graph \mathbb{G}_1 equals $\mathbb{G}[V(\mathbb{G}_i)]$ for $i = 1, 2$. In other words all edges of G between vertices of \mathbb{G}_i , $i = 1, 2$, are already edges of \mathbb{G}_i .

Lemma 8 ([\[17\]](#page-31-8)). *(1) Every graph of clique width* k *can be represented by a safe* k*-expression.*

(2) A safe k*-expression for a graph of clique width* k *can be found in plain exponential time.*

2.3 Extended k-expressions

In this section we introduce a more general version of k -expressions, and accordingly a more general version of clique width.

Fix a natural k. By \overrightarrow{n} we denote a vector $(n_1, \ldots, n_k) \in ([k] \cup \{0\})^k$. For such a vector \vec{n} , let

$$
\mathcal{L}(\overrightarrow{n}) = \{ (i_1, j_1, i_2, j_2) \mid i_1, i_2 \in [k], 1 \le j_1 \le n_{i_1}, 1 \le j_2 \le n_{i_2} \}.
$$

New k -expressions require two more operators on k -labeled graphs. The first one does not have analogues in k-expressions.

Definition 9. Let $\overrightarrow{n} = (n_1, \ldots, n_k) \in ([k] \cup \{0\})^k$, $\sigma : [k] \rightarrow [k]$, and $\mathcal{S} \subseteq \mathcal{L}(\overrightarrow{n})$. *Also, let* S *be a symmetric set, that is, if* $(i_1, j_1, i_2, j_2) \in S$ *then* $(i_2, j_2, i_1, j_1) \in$ S. Operator $\beta_{\vec{n},\sigma,S}$ *transforms* k-labeled graph $\mathbb{G}_1 = (V_1, E_1, \pi_1)$ to a k-labeled *graph* $\mathbb{G}_2 = (V_2, E_2, \pi_2)$ *as follows:*

- $V_2 = \bigcup_{i=1}^k C_i$ *, where* $C_i = \{a_j | j \in \{0, ..., n_i\}$ *,* $a \in V_1$ *and* $\pi_1(a) = i\}$ *. The vertices of* $a_0, a \in V_1$, are called original vertices of $\mathbb{G}_2 = \beta_{\overrightarrow{n}, \sigma, \mathcal{S}}(\mathbb{G}_1)$ and *are identified with their corresponding vertices from* V1*;*
- $(a_j, b_{j'}) \in E_2$ *if and only if* $(a, b) \in E_1$ *, and* $(\pi_1(a), j, \pi_1(b), j') \in S$ *or* $j = j' = 0;$

•
$$
\pi_2(a_j) = \begin{cases} \pi_1(a), & \text{if } j = 0, \\ \sigma(\pi_1(a)), & \text{otherwise.} \end{cases}
$$

We also refer to this operator as the beta operator*.*

The second operator combines disjoint union with a sequence of adding edges operators.

Definition 10. Let $\mathcal{T} \subseteq [k] \times [k]$. Operator $\eta_{\mathcal{T}}$ takes two k-labeled graphs as input and produces a k-labeled graph as output. For k-labeled graphs \mathbb{G}_1 =

 (V_1, E_1, π_1) *, and* $\mathbb{G}_2 = (V_2, E_2, \pi_2)$ *,* V_1, V_2 *disjoint the k-labeled graph* $\eta_{\mathcal{T}}(\mathbb{G}_1, \mathbb{G}_2)$ (V, E, π) *, is defined as follows:*

$$
V = V_1 \cup V_2
$$

\n
$$
E = E_1 \cup E_2 \cup \{(a, b) \mid a \in V_1, b \in V_2, \pi_1(a) = i, \pi_2(b) = j, (i, j) \in \mathcal{T}\}
$$

\n
$$
\pi = \begin{cases} \pi_1(a), & \text{if } a \in V_1, \\ \pi_2(a), & \text{if } a \in V_2. \end{cases}
$$

We also refer to this operator as the connect operator*.*

An *extended* k*-expression* is a (properly formed) expression that involves operators \cdot_i ($i \in [k]$), $\rho_{i \to j}$ ($i, j \in [k]$), $\beta_{\overrightarrow{n}, \sigma, \mathcal{S}}$, and $\eta_{\mathcal{T}}$, where $\overrightarrow{n}, \sigma, \mathcal{S}, \mathcal{T}$ are as in Definitions [9,](#page-5-0) [10.](#page-5-1) Similar to k -expressions, extended k -expressions represent k labeled graphs, as well as usual graphs. Next we explore what kind of graphs and k-labeled graphs can be represented by extended k-expressions.

Note that if \mathbb{G}_1 and \mathbb{G}_2 are two isomorphic k-labeled graphs, and \mathbb{G}_1 has an extended k-expression Φ , then Φ is an extended k-expression for \mathbb{G}_2 as well.

As is easily seen, the connect operator can be expressed through disjoint union and adding edges. However, we will need properties similar to the safety of k-expressions. Unfortunately, the beta operator does not allow an equally clean and easy definition of safety, as in the case or k expressions, and we use the connect operator instead.

Let $\mathbb{G} = \eta_{\mathbb{F}}(\mathbb{G}_1, \mathbb{G}_2)$. It is straightforward from the definition that $\mathbb{G}[V(\mathbb{G}_1)]$ is equal to \mathbb{G}_1 and $\mathbb{G}[V(\mathbb{G}_2)]$ is equal to \mathbb{G}_2 , that is, $\eta_{\mathcal{T}}$ does not add edges inside $\mathbb{G}_1, \mathbb{G}_2$. Similarly, if $\mathbb{G} = \beta_{\vec{\pi}, \mathbb{S}, \sigma}(\mathbb{G}_1)$, then again $\mathbb{G}[V(\mathbb{G}_1)]$ is equal to \mathbb{G}_1 . Similar to k-expressions we say that an extended k-expression Φ is *safe* if for every its subexpressions $\eta_{\mathcal{T}}(\Phi_1, \Phi_2)$ and $\beta_{\vec{n},\sigma,S}(\mathbb{G}_1)$ such that Φ_1, Φ_2 represent graphs $\mathbb{G}_1, \mathbb{G}_2$, respectively, it holds $\mathbb{G}_1 = \mathbb{G}[V(\mathbb{G}_i)]$ for $i = 1, 2$. The following property is straightforward.

Lemma 11. *Any extended* k *expression is safe.*

For an extended k-expression Φ , size(Φ) denotes the total number of operands, connect operators, beta operators, and the maximal subsequences of relabelling operators of Φ.

A graph $G = (V, E)$ is said to have *extended clique width* k if there is a klabelling π of G and an extended k-expression Φ that represents $\mathbb{G} = (V, E, \pi)$. If such a π exists we also say that Φ represents G. The class of all graphs of extended clique width is denoted by \mathcal{X}_k .

We complete this section showing that \mathcal{C}_k is a subset of \mathcal{X}_k .

Proposition 12. *Any graph* G *that can be represented by a* k*-expression, can also be represented by an extended* k*-expression.*

Proof. We start with a piece of terminology. For a sequence Ψ of operators of the form $\rho_{i\to j}$ and η_{ij} , consider the k-labeled graph $\mathbb{G} = \Psi(\mathbb{G}_1 \bigoplus \mathbb{G}_2)$ for some $\mathbb{G}_1, \mathbb{G}_2$. Expression Ψ is said to *connect* a vertex a to a vertex b, if ab is an edge

of \mathbb{G} , but not of $\mathbb{G}_1 \bigoplus \mathbb{G}_2$, that is, if there is an operator η_{ij} in Ψ that connects a to b. Expression Ψ is said to relabel a vertex a with label i to label j, if a has label *i* in $\mathbb{G}_1 \bigoplus \mathbb{G}_2$ and label *j* in \mathbb{G} . Also, $\hat{\Psi}$ denotes the sequence of operators that is obtained from Ψ by removing all the η_{ij} operators..

Let Φ be a k-expression representing graph G. By Lemma [8](#page-5-2) Φ can be assumed safe. We proceed by induction on the structure of Φ . If G is a graph with one vertex, there is nothing to prove. If G has more than one vertex, we can write Φ as $\Psi(\mathbb{G}_1 \bigoplus \mathbb{G}_2)$ where, \mathbb{G}_1 and \mathbb{G}_2 are represented by some subexpressions of Φ and Ψ is a sequence of operators of the form $\rho_{i\to j}$ and η_{ij} . Let $\mathbb{G}_1 = (V_1, E_1, \pi_1)$ and $\mathbb{G}_2 = (V_2, E_2, \pi_2)$.

As is easily seen, for any $a \in V_1$ and $b \in V_2$, if $ab \in E(G)$, then there is operator η_{st} in Ψ that connects a to b. Conversely, if some operator η_{st} in Ψ connects vertex $a \in V_1$ to a vertex $b \in V_2$ where, $\pi_1(a) = i$ and $\pi_2(b) = j$, then, Ψ connects every vertex $x \in V_1$ to every vertex $y \in V_2$, with $\pi_1(x) = i$ and $\pi_2(y) = j$, because the vertices from $\mathbb{G}_1 \oplus \mathbb{G}_2$ with the same label, remain with the same label, after applying any operator. So, there is a set $\mathcal T$ of pairs (i, j) such that Ψ connects every vertex $x \in V_1$ to every vertex $y \in V_2$ with $\pi_1(x) = i$ and $\pi_2(y) = j$.

Therefore, the set of edges between \mathbb{G}_1 and \mathbb{G}_2 which are added by Ψ is the same as those which are added by $\eta_{\mathcal{T}}$. Also, $\mathbb{G}[V_1] = \mathbb{G}_1$ and $\mathbb{G}[V_2] = \mathbb{G}_2$, as Φ is safe. Thus, $\eta_{\mathcal{T}}(\mathbb{G}_1, \mathbb{G}_2)$ and $\Psi(\mathbb{G}_1 \bigoplus \mathbb{G}_2)$ have the same set of vertices and same set of edges. Moreover, Ψ relabels any vertices same way as Ψ does. Therefore $\Psi(\mathbb{G}_1 \bigoplus \mathbb{G}_2)$ and $\hat{\Psi}\eta_{\mathcal{T}}(\mathbb{G}_1, \mathbb{G}_2)$ represent the same k-labeled graph. By the induction hypothesis the result follows. 口

Corollary 13. *Every graph that has clique width* k *also has extended clique width* k*.*

Next we show that not all graphs of extended clique width k also have clique width k. More precisely, we present a class of graph of extended clique width 2 that does not have bounded clique width.

2.4 Graph class of bounded extended but not regular clique width

An *n*-dimensional hypercube, denoted HC_n is a graph whose vertices are *n*-bit binary vectors, and two vertices are adjacent if the Hamming distance between them is exactly 1. Let $\mathcal{HC} = \{HC_n \mid n \in \mathbb{N}\}\.$ We first show that each hypercube is represented by an extended 2-expression, and therefore has extended width 2.

Extended 2-expressions for hypercubes are constructed by induction on the dimensionality of the hypercube. The base cases of induction are HC_0 and HC_1 . An extended 2-expression for HC_0 is \cdot_1 , and an extended 2-expression for HC_1 is $\eta_{\{(1,2)\}}(\cdot_1, \cdot_2)$.

Suppose that for $m < n$ the graph HC_m has an extended 2-expression. Let Φ be an extended 2-expression for HC_n. Let $\vec{n} = (1, 1)$, let $\sigma : [2] \rightarrow 2$ be the function given by $\sigma(1) \mapsto 2, \sigma(2) \mapsto 1$, and let S be $\{(1, 1, 1, 1), (2, 1, 2, 1), (1, 1, 2, 1),$ $(2, 1, 1, 1), (1, 0, 2, 1), (2, 1, 1, 0), (2, 0, 1, 1), (1, 1, 2, 0)$. Then it is not hard to see that $\beta_{\overrightarrow{n},\sigma,\mathcal{S}}\Phi$ is an extended 2-expression for HC_{n+1}.

Lemma 14. HC *does not have bounded clique width.*

Proof. Let k be a constant, let n be a sufficiently large number, let V be a set of cardinality n, and, for the sake of contradiction, let Φ_0 be a k-expression for HC_n. We define a finite sequence of k-expressions $\{\Phi_i\}$ which starts with k -expression Φ_0 .

For $i \geq 0$, the $(i + 1)$ -th element of the sequence is defined from the *i*-th element of the sequence as follows: if Φ_i represents a graph with more than 1 vertices, it has the form $\Psi_i(\Phi_{i,l} \bigoplus \Phi_{i,r})$, where Ψ_i is a sequence of recolouring and renaming operators and $\Phi_{i,l}$ and $\Phi_{i,r}$ are two k-expressions. Then, let Φ_{i+1} be the k-expression from $\{\Phi_{i,l}, \Phi_{i,r}\}\$ that represents the graph with the greatest number of vertices. Thus, if G_i is the graph represented by Φ_i and G_{i+1} the graph represented by Φ_{i+1} , then $|V(G_{i+1})| \geq |V(G_i)|/2$.

Let G_i be the first graph in the sequence such that $|V(G_i)|$ is less than *n*. As G_i is the first such graph, $|V(G_{i-1})| \geq n$. By the observation above, $|V(G_i)| \ge n/2$. By the Pigeonhole principle, for some label in [k], say 1, there are more than $n/2k$ vertices of G_i labeled with it. Since n is sufficiently large, there are at least three vertices that are labeled 1. Let us denote these vertices a, b, c. Since G_i contains fewer than n vertices and the degree of a in G equals n, there are at least two vertices outside G_i adjacent to a. Denote them d and e. Now, at some point in Φ vertices d, e are connected to a. However, as b, c have the same label as a, the operator $\eta_{i\ell}$ that connects a to d and a to e, also adds edges bd, be, cd, ce. So, the subgraph of HC_n induced by $\{a, b, c, d, e\}$ contains a complete bipartite graph $K_{3,2}$.

Let us view a, b, c, d, e as n-bit vectors. Then d differs from from each of a, b, c in one bit, but the position of that bit is different, say, it is 1,2 and 3, respectively. This means that a, b, c are equal in all the remaining positions. Since e also differs from each of a, b, c in one position, these must be the same positions, and $d = e$. \Box

2.5 Finding an extended k -expression for a given graph

Next we show how to find an extended k-expression for a given graph G if it has one, in time $O^*(k^{|V(G)|})$.

One of the ingredients of our algorithm is the problem of deciding whether two k-labeled graphs are isomorphic. k-labeled graphs $\mathbb{G} = (V_1, E_1, \pi_1), \mathbb{H} =$ (V_2, E_2, π_2) are isomorphic if there exists an isomorphism φ from the graph $G = (V_1, E_1)$ to $H = (V_2, E_2)$ such that $\pi_1(a) = \pi_2(\varphi(a))$ for $a \in V_1$. We show that this problem can be reduced to the regular Graph Isomorphism problem and use the celebrated result by Babai [\[1\]](#page-30-3) that there is an algorithm that, given graphs G and H , decides whether there exists an isomorphism between G and *H* in time $O(2^{\log(|V(G)|)^{O(1)}})$.

Proposition 15. *There is an algorithm that decides if* k*-labeled graphs* G, H *are isomorphic and runs in time* $O^*(2^{\log((k+2)n)^{O(1)}})$ *.*

Before proving Proposition [15,](#page-9-0) we introduce a gadget construction, see, Fig [1,](#page-9-1) which we borrow from [\[7\]](#page-31-10) with minor modifications. Let us denote the gadget in Fig [1](#page-9-1) by D. It is proved in [\[7\]](#page-31-10) that for each homomorphism $\varphi : D \to D$ and $i \in [5]$, $\phi(z) = z$ and $\varphi(z) \neq \varphi(x_i)$.

Figure 1: The graph D

We join q more such gadgets D in a row to construct a larger gadget T_q that has total $q+1$ copy of gadget D. We label the left-most vertex of the *i*-th block in the chain by z_{i-1} except for the first block, see, Fig [2.](#page-9-2) Each automorphism of T_q preserves the order on z's which means for each $i \in [q]$ and isomorphism $\varphi: T_q \to T_q$, $\varphi(z_i) = z_i$. This property is also proved in [\[7\]](#page-31-10).

Figure 2: Gadget T_q

For each $i \in [q]$, we replace z_i in T_q with a copy of K_{n+3} , a clique on $n+3$ vertices, and connect every vertex of K_{n+3} to all the neighbours of z_i , its special vertex, in the next subsequent block that is $(i+1)$ th block. Note that only one special vertex of the $(i + 1)$ th copy of K_{n+3} , that is called z_i , is connected to some vertices of previous block. Denote the new graph by $T_{q,n}$, see Fig [3](#page-9-3)

Figure 3: Gadget $T_{a,n}$

Let $\mathbb{G} = (V_1, E_1, \pi_1), \mathbb{H} = (V_2, E_2, \pi_2)$ be k-labeled graphs, and let $G =$ $(V_1, E_1), H = (V_2, E_2)$. Now we construct an instance G', H' of GRAPH Iso-MORPHISM. Let $A_q = \{a_1, ..., a_q\}$. Then the graph G' consists of a copy of G, a copy of $T_{q,n}$, and a copy of vertices from A_q with the following additional edges: for each $i \in [q]$ the vertex z_i from the $(i + 1)$ th block of $T_{q,n}$ is adjacent to the vertex a_i . Also, we add edges from G to vertices from A_q : for a vertex $b \in V_1$ with $\pi_1(b) = i$, we add an edge $\{b, a_i\}$, see, Fig [4.](#page-10-0)

Figure 4: Constructing G'

Graph H' is constructed in a similar way. Let $T'_{q,n+3}$ be a copy of $T_{q,n}$ such that for each $i \in [q], z_i$ is renamed to z'_i in $T'_{q,n}$. Also, let A'_q be the set of vertices $\{a'_1, ..., a'_q\}$. Graph H' consists of a copy of H, a copy of $T'_{q,n}$, and a copy of vertices from A'_q with the following additional edges: for each $i \in [q]$, the vertex z_i' from the $(i+1)$ th block of $T'_{q,n}$ is adjacent to the vertex a_i' . Also, we add edges from H to vertices from A'_q : for a vertex $c \in V_2$ with $\pi_2(b) = i$, we add an edge $\{c, a'_i\}.$

We need several properties of this construction. Some of them are proved in [\[7\]](#page-31-10) and the rest are proved here.

Lemma 16. (1) Any isomorphism φ from G' to H' maps $T_{q,n}$ to $T'_{q,n}$.

- (2) Any isomorphism φ from G' to H' bijectively maps $T_{q,n}$ to $T'_{q,n}$ so that *the for each* $i \in [q]$ *,* z_i *is mapped to* z'_i *.*
- (3) Any isomorphism φ from G' to H' maps vertices from A_q to vertices from A'_q so that a_i is mapped to a'_i for each $i \in [q]$.
- (4) *Any isomorphism* φ *from* G' *to* H' *maps* G *to* H *.*

Proof. $(1),(2)$ are proved in [\[7\]](#page-31-10).

(3) By items (1),(2) the restriction of φ on $T_{q,n}$ is a bijective and surjective mapping from $T_{q,n}$ to $T'_{q,n}$, such that $\varphi(z_i) = z'_i$ for each $i \in [q]$. This also means that each vertex of $T'_{q,n}$ is the image of some vertex from $T_{q,n}$ under φ .

Thus, for each $i \in [q]$, $\varphi(a_i)$ cannot be from $T'_{q,n}$ as otherwise φ would not be a bijection. On the other hand, for each $i \in [q]$, since φ is an isomorphism and (z_i, a_i) is an edge of G', $\varphi(a_i)$ is adjacent to $\varphi(z_i) = z'_i$. Now, for each $i \in [q]$, the adjacent vertices of z_i' are either in $T'_{q,n}$ or it is a_i' . As $\varphi(a_i)$ cannot be in $T'_{q,n}$ and has to be adjacent to z'_i , thus $\varphi(\alpha_i) = \alpha'_i$.

(4) By items (1)–(3) the restriction of φ on $T_{q,n}$ and A_q is a bijective and surjective mapping from $T_{q,n}$ to $T'_{q,n}$, and from A_q to from A'_q . Thus each vertex from $T'_{q,n}$ or A'_q is the image of some vertex in G' that is also in $T_{q,n}$ or A_q , under the φ . Hence the image of each vertex of G is a vertex from H, under φ . \Box

Now we can prove Proposition [15.](#page-9-0)

Proof of Proposition [15.](#page-9-0) Let G′ and H′ be two graphs constructed from G and H as described above. First we show that there is a isomorphism from \mathbb{G} to \mathbb{H} if and only if there is an isomorphism from G' to H' .

Let φ be an isomorphism from G to H. We show that its natural extension φ' mapping $T_{q,n}$ to $T'_{q,n}$ and vertices from A_q to vertices from A'_q is an isomorphism from G' to H' . The fact that it maps edges of G' to edges of H' is non-trivial only for edges of G' from G to vertices of A_q . Consider an edge from a vertex $b \in V_1$ with $\pi_1(b) = i$ to the vertex a_i . We have $\pi_2(\varphi'(b)) = i$ as the φ' is an extension of an isomorphism from $\mathbb G$ to $\mathbb H$. Also $\varphi'(a_i) = a'_i$ means that φ' maps (a_i, b) to an edge of H' .

For the reverse direction, let φ' be an isomorphism from G' to H' . We show that its restriction on $V(G)$ is an isomorphism from G to H. Since by Lemma 16(4) φ' maps G to H, it is enough to check that φ' preserves the labelling. In order to do this consider a vertex $b \in V_1$ with $\pi_1(b) = i$. Then $\pi_2(\varphi(b)) = i$ as well, as otherwise $(\varphi'(b), a'_i)$ would not be an edge of H'.

Therefore, there is a polynomial time reduction the Isomorphism problem for k -labeled graphs to GRAPH ISOMORPHISM such that the number of vertices only grows by a constant factor $(k+2)$. Now by [\[1\]](#page-30-3) there is an algorithm that solves the instance (G', H') in time $O(2^{\log((k+2)n)^{O(1)}})$. The result follows. \Box

The following theorem is the main result of this section.

Theorem 17. *If graph* G *has extended clique width* k*, then an extended* k*expression for* G *can be found in time* $O^*((4k+4)^{|V(G)|})$ *.*

Proof. We create an array N of size $(k + 1)^n$ whose entries $N(\mathbb{G}')$ are labeled with a k-labelling \mathbb{G}' of a subgraph G' of G . For any entry $N(\mathbb{G}')$ the k-labeled graph G′ either has an extended k-expression or it does not. The goal is to set the value of each entry $N(\mathbb{G}')$ to some extended k-expression for \mathbb{G}' if it has one and to "no" otherwise.

Now we consider more detailed possibilities for each \mathbb{G}' . There are four cases. Case 1 takes place if \mathbb{G}' has an extended k-expression that ends with a Beta operator; Case 2 takes place if \mathbb{G}' it has an extended k-expression that ends with a connect operator; Case 3 takes place if \mathbb{G}' has an extended k-expression that

ends with a sequence of relabelling operators; and, finally, Case 4 takes place if \mathbb{G}' does not have an extended k-expression.

All one-element k-labeled graphs are obviously represented by an extended k-expression. Let us suppose the values of each entry $N(\mathbb{G}')$, where \mathbb{G}' contains at most $n-1$ vertices is set correctly. Then, we want to set the correct values for entries of the array whose associated k -labeled graph has exactly n vertices. We use the dynamic programming approach that consists of two phases. In Phase 1, for each entry $N(\mathbb{G}')$ such that \mathbb{G}' has n vertices, we check if \mathbb{G}' satisfies the conditions of Case 1. Then for each k-labeled graph like this that does not satisfy the conditions of Case 1 we check if it falls in Case 2. In Phase 2, by relabelling \mathbb{G}' for which $N(\mathbb{G}')$ is assigned a value, we find a new extended kexpression for G′ that do not satisfy the conditions of Cases 1 and 2, but satisfy the conditions of Case 3. In the end, for each \mathbb{G}' that belongs to none of Cases 1, 2, or 3, we set the value $N(\mathbb{G}')$ to "no" because it does not have an extended k-expression. In the rest of this proof, for a k-labeled graph \mathbb{G}' , we show how to check if it satisfies the conditions each of Cases 1 and 2.

Let $\mathbb{G}' = (V', E', \pi')$ be a k-labeled graph with $|V'| = n$ and it has an extended k-expression that ends with a beta operator. Then there is an induced subgraph \mathbb{G}_1' of \mathbb{G}' such that the result of application of a beta operator to \mathbb{G}_1' is isomorphic to \mathbb{G}' and \mathbb{G}'_1 has an extended k-expression. Thus, there exist $\sigma : [k] \to [k], \overrightarrow{n} \in (\{0\} \cup [k])^k, \mathcal{S} \subseteq \mathcal{L}(\overrightarrow{n})$, and a set $V'_1 \subset V'$, such that

- (A) $\mathbb{G}'_2 = (V'_2, E'_2, \pi'_2) = \beta_{\overrightarrow{\pi}, \mathbb{S}, \sigma}(\mathbb{G}'[V'_1])$ is isomorphic to \mathbb{G} , and
- (B) $\mathbb{G}'[V_1']$ has an extended k-expression.

Conversely, if there exist $V'_1 \subset V'$, $\sigma : [k] \to [k]$, $\overrightarrow{n} \in (\{0\} \cup [k])^k$, $\mathcal{S} \subseteq \mathcal{L}(\overrightarrow{n})$ satisfying conditions (A),(B), then \mathbb{G}'_2 has an extended k-expression that ends with a beta operator. As \mathbb{G}'_2 and \mathbb{G}' are isomorphic, \mathbb{G}' has an extended kexpression that ends with a beta operator as well. Thus, the sufficient and necessary conditions for \mathbb{G}' to have an extended k-expression that ends with a Beta operator, is that there exist $V'_1 \subset V', \sigma : [k] \to [k], \vec{n} \in (\{0\} \cup [k])^k, \mathcal{S} \in$ $\mathcal{L}(\overrightarrow{n})$ satisfying (A),(B).

The algorithm now searches through all possible selection of V'_1 , σ , S , to check if conditions (A) , (B) satisfied for any of them. Let us evaluate the running time of this procedure. Checking condition (A) takes time $O(2^{\log((k+2)n)^{O(1)}})$ by Proposition [15,](#page-9-0) while condition (B) can be verified by looking up the existing entry $N(\mathbb{G}'[V_1'])$ in $O(1)$ time. There are 2^n choices for V_1' and k^k choices for σ . Vector \overrightarrow{n} can be chosen in $(k+1)^k$ ways, and so $\mathcal{L}(\overrightarrow{n})$ has at most $(k+1)^2(k)^2$ elements. Thus, S can be chosen in at most $2^{(k+1)^2(k)^2}$ ways. Thus, the total running time of filling up $N(\mathbb{G}')$ in this case is upper bounded by

$$
2^{|V(G)|} \times k^k \times (k+1)^2(k)^2 \times 2^{(k+1)^2(k)^2} \times O(2^{\log((k+2)n)^{O(1)}}) = O^*(2^{2|V(G)|}).
$$

Now let us suppose that $\mathbb{G}' = (V', E', \pi')$ has an extended k-expression that ends with a connect operator. Then due to the safety of extended k expressions there exist two induced subgraphs \mathbb{G}_1' and \mathbb{G}_2' of \mathbb{G}' such that, first, they both are represented by extended k-expressions, and, second, there is $\mathcal{T} \subseteq$ $[k]^2$, such that $\eta_{\mathcal{T}}(\mathbb{G}'_1, \mathbb{G}'_2)$ is identical to \mathbb{G}' . Thus to find an extended kexpression for \mathbb{G}' it suffices to go through all partitions of V' into sets V'_1 and V'_2 and for each partition check the following two conditions. First, check if $\mathbb{G}_1' = \mathbb{G}'[V_1']$ and $\mathbb{G}_2' = \mathbb{G}'[V_2']$ have an extended k-expression by looking up the entries $N(\mathbb{G}'_1), N(\mathbb{G}'_2)$. Second, check if there is $\mathcal{T} \subseteq [k]^2$ such that $\eta_{\mathcal{T}}(\mathbb{G}'_1, \mathbb{G}'_2)$ is identical to G. Since there are at most $2^{|V(G)|}$ ways to partition V' into V'_1 and V'_2 , takes time $O(2^{|V(G)|})$ to check if \mathbb{G}' falls into Case 2.

So far we have registered an extended k -expression for every \mathbb{G}' that satisfies the conditions of Case 1 or Case 2. Now, start Phase 2 and check whether any of the remaining k-labeled graphs G′ satisfies the conditions of Case 3. In order to do that we go through all k-labeled graphs \mathbb{G}' with n vertices and such that $N(\mathbb{G}')$ contains an extended k-expression Φ , that is initially for all \mathbb{G}' that fall into Cases 1,2. Then we consider every possible single relabelling ρ_{ij} in turn. If $\rho_{ij}(\mathbb{G}')$ is a k-labeled graph such that $N(\mathbb{G}')$ does not have an extended kexpression, then we set $N(\rho_{ij}(\mathbb{G}')) = \rho_{ij}(\Phi)$. We repeat this process for each k-labeled graph G′ , until no new entries can be filled. The time required for Phase 2 in total, for all \mathbb{G}' , not only those with n vertices is bounded by number of all k -labellings of all subgraphs of G times the number of possible operators ρ_{ij} . As is easily seen, the time required for Phase 2 in total is

$$
(k+1)^{|V(G)|} \times k^2 = O^*((k+1)^{|V(G)|})
$$

Time complexity: The array we construct has $(k+1)^{|V(G)|}$ entries. The time required to complete Phase 1 for all the entries is bounded by $O^*(4^{|V(G)|} \times$ $(k+1)^{|V(G)|}$. The time to complete Phase 2 for all entries is bounded by $O^*((k+1)^{|V(G)|})$. Thus the total running time is $O^*((4k+4)^{|V(G)|})$. \Box

3 Counting homomorphism to labeled graphs given an extended k-expression

In this section we prove our main result. Let $\mathsf{hom}(G, H)$ denote the number of homomorphisms from a graph G to a graph H .

Theorem 18. *Let* G *and* H *be two graphs, and let* k*-labeled graph* H *be a* k*labelling of graph* H. Given an extended k-expression Φ for \mathbb{H} , hom (G, H) can *be found in time* $O^*((2k+1)^{|V(G)|})$

The following notation and terminology will be used throughout this section. The number of homomorphism from graph G to graph H is denoted by $hom(G, H)$. Also, $Hom(G, H)$ denotes the set of all homomorphisms from G to H. Let $X \subseteq V(G)$, and let $\chi : X \to [k]$ be a label function. A mapping φ from X to k-labeled graph $\mathbb{H} = (V, E, \pi)$ is said to be consistent with χ if for every $x \in X$ it holds $\pi(\varphi(x)) = \chi(x)$. Let hom $_{\chi}(G, \mathbb{H})$, HOM $_{\chi}(G, \mathbb{H})$, map $_{\chi}(G, \mathbb{H})$, and $\mathsf{MAP}_\chi(G, \mathbb{H})$, denote the number of homomorphisms from $G[X]$ to $\mathbb H$ consistent with χ , the set of all homomorphisms from $G[X]$ to $\mathbb H$ consistent with χ , the number of all mappings from $G[X]$ to $\mathbb H$ consistent with χ , and the set of all mappings from $G[X]$ to H consistent with χ , respectively.

Let Φ be an extended k-expression for a k-labelling $\mathbb H$ of the graph H. We proceed by induction on the structure of Φ . More precisely, our algorithm will compute entries $\textsf{hom}(\mathbb{G}[X], \mathbb{H}')$, where $X \subseteq V(G)$ and \mathbb{G} is a k-labelling of G , and \mathbb{H}' is the k-labeled graph represented by a subformula of Φ . Operator \cdot_i creating a graph \mathbb{H}' with a single vertex labeled i gives the base case of induction. In this case $\textsf{hom}(\mathbb{G}[X], \mathbb{H}') = 1$ if all vertices of X are labeled i and $\mathbb{G}[X]$ has no edges; otherwise hom $(\mathbb{G}[X], \mathbb{H}') = 0$. Finally, after computing the numbers hom(\mathbb{G}, \mathbb{H}) for all the k-labellings \mathbb{G} of G we complete using the following observation.

Observation 19. *Let* G *and* H *be graphs, and let* k*-labeled graph* H *be a* k*labelling of* H*. Then*

$$
\hom(G,H)=\sum_{\chi: V(G)\to [k]}\hom_\chi(G,\mathbb{H})
$$

It therefore suffices to show how to compute $\text{hom}(\mathbb{G}[X], \mathbb{H}')$, where $\mathbb G$ is an arbitrary k-labelling of $G, X \subseteq V(G)$, and \mathbb{H}' is represented by a subformula Φ' of Φ , provided hom($\mathbb{G}[Y], \mathbb{H}'$) is known for all $Y \subseteq X$ and \mathbb{H}'' represented by a subformula Φ'' of Φ' with $\Phi'' \neq \Phi'$. We consider 3 cases depending on the last operator of Φ' .

3.1 Relabelling Operator

Let Ψ be a string of relabelling operators, applied on a k-labeled graph \mathbb{G} . We say that Ψ relabels i to j, if the application of Ψ on a vertex labeled with i gives a vertex labeled with j.

Let $\mathbb{H}' = (V', E', \pi')$ be a k-labeled graph, and let $\mathbb{H} = \Psi(\mathbb{H}') = (V, E, \pi)$ be the result of application of Ψ to \mathbb{H}' . Let $X \subseteq V(G)$ and let $\chi : X \to [k]$ be any k-label function. Also, let $\varphi \in HOM_{\chi}(G, \mathbb{H})$. Then there is a function $\chi': X \to [k]$ such that $\varphi \in \text{HOM}_{\chi'}(G, \mathbb{H}').$ Let call χ' the *consistent labelling* of φ . To count the elements of $\text{HOM}_\chi(G, \mathbb{H})$, we partition this set into sets of homomorphisms that share the same consistent labelling, and find the number of elements of these smaller sets.

Let $\mathcal{D}(\chi)$ denote the set of all of functions $\chi': X \to [k]$ that satisfy $\chi = \sigma(\chi')$. We show that $\mathcal{D}(\chi)$ is the set of consistent labellings of all homomorphisms from $\mathsf{HOM}_\chi(G, \mathbb{H}').$

Lemma 20. *Let* χ , G , \mathbb{H} , and \mathbb{H}' be as above. Then,

$$
\text{hom}_{\chi}(G,\mathbb{H}) = \sum_{\chi' \in \mathcal{D}(\chi)} |\text{HOM}_{\chi'}(G,\mathbb{H}')|.
$$
 (1)

Proof. Let $\varphi \in \text{HOM}_\chi(G, \mathbb{H})$. Also, let $\chi' : X \to [k]$, be the consistent labelling of φ . As is easily seen, χ' satisfies $\chi = \sigma(\chi')$. Now, let $\chi' \in \mathcal{D}(\chi)$, then the set $\text{HOM}_{\chi'}(G, \mathbb{H}')$ is the set of all elements of $\text{HOM}_{\chi}(G, \mathbb{H})$ that have χ' as their consistent labelling. Note that for two different $\chi', \chi'' \in \mathcal{D}(\chi)$ the sets $\text{HOM}_{\chi'}(G,\mathbb{H}')$ and $\text{HOM}_{\chi''}(G,\mathbb{H}')$ are disjoint. Thus,

$$
\mathsf{HOM}_\chi(G, \mathbb{H}) = \bigcup_{\chi' \in \mathcal{D}(\chi)} \mathsf{HOM}_{\chi'}(G, \mathbb{H}')
$$

The result follows.

We now have an algorithm computing all the required numbers of homomorphisms.

Corollary 21. Let $Y \subseteq V(G)$ and $\gamma: Y \rightarrow [k]$. There is an algorithm that *given* $\textsf{hom}_{\zeta}(G, \mathbb{H}')$ *for all functions* ζ *from subsets of* $V(G)$ *to* [k] *as input, finds* $\mathsf{hom}_{\gamma}(G,\mathbb{H})$ *in time* $O^*(2^{|V(G)|}).$

Proof. First for a fixed $X \in V(G)$ and a fixed function $\chi : X \to [k]$ we calculate the upper bound on the size of $\mathcal{D}(\chi)$. For a function $\chi' \in \mathbb{D}$, we can count those functions by considering their possible values on each element of X . Since the codomain of χ' is [k] and size of X is bounded by $|V(G)|$, there are at most $k^{|V(G)|}$ possibilities for χ' . By Lemma [22](#page-16-0) hom_{γ} (G, \mathbb{H}) can be found by summing over $O^*(k^{|V(G)|})$ known values. That can be done in time $O^*(k^{|V(G)|})$. \Box

3.2 Connect Operators

Let $\mathbb{H}' = (V', E', \pi')$ and $\mathbb{H}'' = (V'', E'', \pi'')$ be k-labeled graphs, let \mathcal{T} be a subset of $[k] \times [k]$, and let $\mathbb{H} = \eta \tau(\mathbb{H}', \mathbb{H}'')$ be a k-labeled graph. Let $X \subseteq V(G)$, and let $\chi: X \to [k]$ be any k-label function.

To count elements of $\text{HOM}_\chi(G, \mathbb{H})$, one can partition them into smaller subsets of elements and count the elements in each subset. For any $\varphi \in$ HOM_X(G, H), let $\varphi' = \varphi|_{X'}$ and $\varphi'' = \varphi|_{X''}$, where X' and X'' are preimages of $V(\mathbb{H}')$ and $V(\mathbb{H}'')$ under φ , respectively. Then φ' and φ'' are homomorphisms from $G[X']$ to \mathbb{H}' and from $G[X'']$ to \mathbb{H}' , respectively. Thus, there are unique k-label functions $\chi': X' \to V(\mathbb{H}')$ and $\chi'': X'' \to V(\mathbb{H}'')$ with which φ' and φ'' are consistent. Let us call the pair of k-label functions (χ', χ'') , the *consistent pair* of φ . The idea is to partition $\text{HOM}_\chi(G, \mathbb{H})$ into smaller subsets such that for each of them all the homomorphisms φ in that subset share the same consistent pair.

Let $\text{All}(\chi)$ be the set of all pairs of k-label functions (χ', χ'') , $\chi' : X' \to$ $V(\mathbb{H}'), \chi'' : X'' \to V(\mathbb{H}'')$, where X' and X'' are disjoint subsets of X and:

- (a.1) $X' \cup X'' = X$; and also $\chi|_{X'} = \chi'$ and $\chi|_{X''} = \chi''$.
- (a.2) For any $x' \in X'$, such that $\chi'(x) = i$ and for any $x'' \in X''$ such that $\chi''(x'') = j$, if $x'x'' \in E(G)$ then, $(i, j) \in \mathcal{T}$.

We show that $All(\chi)$ is the set of consistent pairs of homomorphisms from HOM_x(G, H). Also for every pair $(\chi', \chi'') \in All(\chi)$ we count the number of $\varphi \in \text{HOM}_\chi(G, \mathbb{H})$, such that (χ', χ'') is the consistent pair of φ .

 \Box

Lemma 22. *The number* hom_x (G, \mathbb{H}) *satisfies the following equality*

$$
\mathsf{hom}_\chi(G,\mathbb{H})=\sum_{(\chi',\chi'')\in\mathsf{All}(\chi)}\mathsf{hom}_{\chi'}(G,\mathbb{H}')\mathsf{hom}_{\chi''}(G,\mathbb{H}'').\tag{2}
$$

Proof. First, observe that the right hand side of [\(2\)](#page-16-1) counts the elements of the set

$$
R = \bigcup_{(\chi',\chi'')\in\mathrm{All}(\chi)} \mathrm{HOM}_{\chi'}(G,\mathbb{H}')\times\mathrm{HOM}_{\chi''}(G,\mathbb{H}'').
$$

Also, the left hand side equals the cardinality of the set $\text{HOM}_\chi(G, \mathbb{H})$ by defi-nition. So, [\(2\)](#page-16-1) can be proved by constructing a bijection between $\text{HOM}_\chi(G, \mathbb{H})$ and R.

Let $\varphi \in \text{HOM}_X(G, \mathbb{H})$, and let X' and X'' be preimages of $V(\mathbb{H}')$ and $V(\mathbb{H}'')$ under φ , respectively. Consider the mapping $\tau : \varphi \mapsto (\varphi|_{X'}, \varphi|_{X''})$.

To show that τ is a bijection between $\text{HOM}_\chi(G, \mathbb{H})$ and R, one needs to show that, first, for any $\varphi \in \text{HOM}_\chi(G, \mathbb{H}), \tau(\varphi)$ is an element of R; and, second, that τ is surjective and injective.

Let $\varphi' = \varphi|_{X'}$ and $\varphi'' = \varphi|_{X''}$. As is easily seen, φ' and φ'' are elements of $\text{HOM}(G, \mathbb{H}')$ and $\text{HOM}(G, \mathbb{H}'')$, respectively. Also, there is a unique pair of functions $\chi': X' \to [k]$ and $\chi'': X'' \to [k]$ such that φ' and φ'' are consistent with χ' and χ'' , respectively, i.e., $\varphi' \in \text{HOM}_{\chi'}(G, \mathbb{H}')$ and $\varphi'' \in \text{HOM}_{\chi''}(G, \mathbb{H}'')$. Since $\varphi' = \varphi|_{X'}$ and $\varphi'' = \varphi|_{X''}$ are consistent with χ' and χ'' , respectively, $\chi|_{X'} = \chi'$ and $\chi|_{X''} = \chi''$. This means that (χ', χ'') satisfies Property (a.1).

Let $x' \in X'$ and $x'' \in X''$ be such that $\chi'(x') = i$ and $\chi''(x'') = j$ and also, $x'x'' \in E(G)$. Then, $\chi(x') = \chi'(x') = i$ and $\chi(x'') = \chi(x'') = j$. Since $\varphi(x')\varphi(x'')$ is an edge of $\mathbb{H} = \eta_{\mathcal{T}}(\mathbb{H}', \mathbb{H}'')$, then $(i, j) \in \mathcal{T}$. Thus (χ', χ'') satisfies Properties (a.1) and (a.2), therefore $(\chi', \chi'') \in All(\chi)$. This means $(\varphi', \varphi'') \in$ $\text{HOM}_{\chi'}(G,\mathbb{H}') \times \text{HOM}_{\chi''}(G,\mathbb{H}'') \subseteq R$. In other words, we have shown that for any $\varphi \in \text{HOM}_\chi(G, \mathbb{H})$, the consistent pair of φ belongs to $\text{All}(\chi)$.

For two nonidentical elements φ and ψ of HOM_x(G, H), $\tau(\varphi)$ is clearly not equal to $\tau(\psi)$, so φ is injective.

Finally, we show that τ is surjective. Let $\chi': X' \to [k]$ and $\chi'': X'' \to [k]$ be any two functions such that $(\chi', \chi'') \in All(\chi)$. Let φ' and φ'' be two homomorphisms in HOM_{X'} (G, \mathbb{H}') and HOM_{X'}' (G, \mathbb{H}'') , respectively, i.e., $(\varphi', \varphi'') \in R$. Let

$$
\varphi(x) = \begin{cases} \varphi'(x), & x \in X', \\ \varphi''(x), & x \in X''. \end{cases}
$$

By Property (a.1), $\chi|_{X'} = \chi'$ and $\chi|_{X''} = \chi''$. Then, φ is consistent with χ , i.e. $\varphi \in \mathsf{MAP}_\chi(G, \mathbb{H}).$

Let $x' \in X'$ and $x'' \in X''$ be such that $x'x'' \in E(G)$. To show that φ is a homomorphism, observe that $\varphi(x') = \varphi'(x')$ and $\varphi(x'') = \varphi''(x'')$. Thus, $\pi'(\varphi'(x')) = \chi'(x') = i$ and $\pi''(\varphi''(x'')) = \chi''(x'') = j$. Since $x'x'' \in E(G)$ by Property (a.2), $(i, j) \in \mathcal{T}$ and thus $\varphi'(x')\varphi''(x'') \in E(\mathbb{H})$. Therefore, φ is a homomorphism.

For an arbitrary element (χ', χ'') of $\text{All}(\chi)$ and an arbitrary element (φ', φ'') of $\text{HOM}_{\chi'}(G,\mathbb{H}') \times \text{HOM}_{\chi''}(G,\mathbb{H}'') \subseteq R$, there is φ from $\text{HOM}_{\chi}(G,\mathbb{H})$ such that $\tau(\varphi) = (\varphi', \varphi'')$ proving τ is surjective.

The bijection τ between $\text{HOM}_\chi(G, \mathbb{H})$ and R, proves [\(2\)](#page-16-1).

□

Lemma 23. Let X be a subset of $V(G)$ and let χ be a function $\chi: X \to [k]$. *There is an algorithm that given* $hom_{\zeta}(G, \mathbb{H}')$ *and* $hom_{\zeta}(G, \mathbb{H}'')$ *for all functions* ζ *from a subset of* $V(G)$ *to* $[k]$ *as input, finds* hom $_{\chi}(G, \mathbb{H})$ *in time* $O^*(2^{|V(G)|})$ *.*

Proof. By Lemma [22](#page-16-0) hom_{χ} (G, \mathbb{H}) can be found by summing over $O^*(2^{|V(G)|})$ known values. \Box

3.3 Beta operators

In this part, we show how to make a recursive step in the case when the last operator of Φ′ is a beta operator. Before explaining this step, we need several definitions.

A *retraction* is a homomorphism ψ from a graph G_2 to its subgraph G_1 such that $\psi(v) = v$ for each vertex v of G_1 . In this case the subgraph G_1 is called a *retract* of G_2 . A retraction from a k-labeled graph $\mathbb{G}_2 = (V_2, E_2, \pi_2)$ to a klabeled graph $\mathbb{G}_1 = (V_1, E_1, \pi_1)$ is defined to be a retraction from $G_2 = (V_2, E_2)$ to $G_1 = (V_1, E_1)$ preserving the label function π_2 , that is, $\pi_2(v) = \pi_1(\psi(v))$ for all $v \in V_2$.

It will be convenient for us to subdivide operator $\beta_{\vec{n},\sigma,S}$ into two steps: the first one is expansion of the original graph using $\vec{\pi}$ and $\vec{\mathcal{S}}$, and the second is relabelling of some vertices of the resulting graph using σ . More specifically, let $\mathbb{H} = (V, E, \pi)$ be a k-labeled graph, $\overrightarrow{n} \subseteq ([k] \cup \{0\})^k$, $S \subseteq \mathcal{L}(\overrightarrow{n})$, and $\sigma : [k] \to [k]$. Then $\mathbb{H}' = (V', E', \pi') = \alpha_{\overrightarrow{n}, \mathcal{S}}(\mathbb{H})$ is given by

- $V' = \bigcup_{i=1}^{k} C_i$, where $C_i = \{a_j | j \in \{0, ..., n_i\}, a \in V \text{ and } \pi_1(a) = i\}.$ The vertices $a_0, a \in V$, are called original vertices of $\mathbb{H}' = \alpha_{\overrightarrow{n}, S}(\mathbb{H})$ and are identified with their corresponding vertices from V ;
- $(a_j, b_{j'}) \in E'$ if and only if $(a, b) \in E$, and $(\pi(a), j, \pi(b), j') \in S$ or $j =$ $j' = 0;$
- $\pi'(a_j) = \pi(a)$.

Then $\mathbb{H}'' = (V'', E'', \pi'') = \beta_{\vec{\pi}, \sigma, \mathcal{S}}(\mathbb{H})$, that is, $V'' = V'$, $E'' = E'$, and

$$
\pi''(a_j) = \begin{cases} \pi(a), & \text{if } j = 0, \\ \sigma(\pi(a)), & \text{otherwise.} \end{cases}
$$

As is easily seen, $\mathbb H$ is an induced subgraph of $\mathbb H'$, and a retract. Indeed, the mapping μ that maps every $o_j \in V(\mathbb{H}')$ to o (recall that o_j is a 'copy' of some $o \in V(\mathbb{H})$ is a retraction.

The objective is to find a method to express the number of homomorphisms from induced subgraphs of G to \mathbb{H}'' given those from induced subgraphs of G to H.

Lemma 24. Let $Y \subseteq V(G)$ and let γ be a function $Y \to [k]$. There is an *algorithm that given* $hom_{\mathcal{C}}(G, \mathbb{H})$ *for all functions* ζ *from a subset of* $V(G)$ *to* [k] *as input, finds* hom_{γ}(\tilde{G} , \mathbb{H}'') *in time* $O((2(k+1))^{2|V(G)|})$ *.*

We break this down into two steps. The main result of Step I, which is summarized in Lemma [25,](#page-19-0) finds an equality for the number of homomorphisms from *G* to H'. Then, the main result of Step II, Lemma [26,](#page-20-0) finds the number of homomorphisms from induced subgraphs of G to \mathbb{H}'' given those for G and \mathbb{H}' .

Step 1

Let $\mathbb{H}' = (V', E', \pi') = \alpha_{\overrightarrow{n}, \mathcal{S}}(\mathbb{H})$ and $\mathbb{H} = (V, E, \pi)$. Let Y be a subset of $V(G)$ and γ a function $Y \to [k]$. Also, set

$$
\mathbb{W}(\gamma) = {\omega \mid \omega : Y \to [k] \cup \{0\} \text{ and } \forall a \in Y, 0 \le \omega(a) \le n_{\gamma(a)}\}.
$$

For $\omega \in \mathbb{W}(\gamma)$, let

$$
\mathsf{HOM}_{\gamma}(G, \mathbb{H}', \omega) = \{ \varphi \mid \varphi \in \mathsf{HOM}_{\gamma}(G, \mathbb{H}') \text{ and } \forall a \in Y, \exists o \in V(\mathbb{H}) \text{ such that } \varphi(a) = o_{\omega(a)} \},
$$

and

$$
\mathsf{MAP}_{\gamma}(G, \mathbb{H}', \omega) = \{ \varphi \mid \varphi \in \mathsf{MAP}_{\gamma}(G, \mathbb{H}') \text{ and } \forall a \in Y, \exists o \in V(\mathbb{H}) \text{ such that } \varphi(a) = o_{\omega(a)} \}.
$$

For the rest of Step I, let X' and X'' be two disjoint subsets of $V(G)$ and let $\chi': X' \to [k]$ and $\chi'': X'' \to [k]$ be arbitrary functions. Also let $X = X' \cup X''$ and let $\chi = \chi' \uplus \chi''$.

Let $\mathsf{HOM}_{\chi',\chi''}(G,\mathbb H')$ denote the set of all elements of $\mathsf{HOM}_\chi(G,\mathbb H')$ that map a vertex a from $V(G)$ to an original vertex of \mathbb{H}' (recall that any vertex of $\mathbb H$ is called an original vertex of \mathbb{H}') if and only if $a \in X'$. The set $\text{HOM}_{\chi',\chi''}(G,\mathbb{H}')$ is an intermediate set that is computed in Step I. In Step 2, this set is used to calculate another value.

For any $\varphi \in \text{HOM}_{\chi',\chi''}(G,\mathbb{H}'),$ there is a unique $\omega \in \mathbb{W}(\chi)$ such that φ is also an element of $\text{HOM}_\chi(G, \mathbb{H}', \omega)$. Let us call ω the *consistent function* of φ . Similar to the method used in Section [3.2,](#page-15-0) one can partition $\text{HOM}_{\chi',\chi''}(G,\mathbb{H'})$ into smaller subsets and count the elements in each smaller subsets. We partition $\text{HOM}_{\chi',\chi''}(G,\mathbb{H}')$ into sets of homomorphisms that all share the same consistent function ω for some $\omega \in \mathbb{W}(\chi)$. As is easily seen, $\mathsf{HOM}_\chi(G, \mathbb{H}', \omega) \cap$ $\mathsf{HOM}_{\chi',\chi''}(G,\mathbb H')$ is such a subset.

Let $\mathcal{B}(\chi', \chi'')$ be the set of all $\omega \in \mathbb{W}(\chi)$ such that ω satisfies the following properties:

(b.1) $\omega \in \mathbb{W}(\chi)$ and $\omega(x) = 0$ iff $x \in X'$.

(b.2) For every $a, b \in X$ such that at least one of them is not an element of X', and $ab \in E(G)$ it holds that $(\chi(a), \omega(a), \chi(b), \omega(b)) \in S$.

We show that, the number of elements in $HOM_{\chi}(G, \mathbb{H}')$ such that ω is their consistent function is the same for any $\omega \in \mathcal{B}(\chi', \chi'')$ and it is zero otherwise.

Lemma 25. Let G , H , H' , χ' , and χ'' be defined as above, then

$$
|\text{HOM}_{\chi',\chi''}(G,\mathbb{H}')|=|\mathcal{B}(\chi',\chi'')|\times \text{hom}_\chi(G,\mathbb{H}).
$$

Proof. One can easily observe that if $\omega \in \mathbb{W}(\chi)$ does not satisfy Property (b.1), then the number of $\varphi \in HOM_{\chi',\chi''}(G,\mathbb{H}')$ that have ω as their consistent function is zero. Otherwise, if $\omega \in \mathbb{W}(\chi)$ satisfies Property (b.1), then $\mathsf{HOM}_\chi(G,\mathbb{H}',\omega)$ is the set of all $\varphi \in \text{HOM}_{\chi',\chi''}(G,\mathbb{H})$ that have ω as their consistent function. So,

$$
\mathsf{HOM}_{X',X''}(G,\mathbb{H}')=\bigcup_{\substack{\omega\in\mathbb{W}(\chi)\\ \omega(x)=0\text{ iff }x\in X'}}\mathsf{HOM}_\chi(G,\mathbb{H}',\omega).
$$

Based on this, to prove the Lemma [25,](#page-19-0) it suffices to prove that if $\omega \in \mathbb{W}(\chi)$ and it satisfies Property (b.1) but it does not satisfy Property (b.2), then the size of $\text{HOM}_\chi(G, \mathbb{H}', \omega)$ is zero, and otherwise, if $\omega \in \mathbb{W}(\chi)$ satisfies Properties (b.1) and (b.2), then the size of $\text{HOM}_{\chi}(G, \mathbb{H}', \omega)$ is equal to the size of $\text{HOM}_{\chi}(G, \mathbb{H})$.

Let ω be a function that satisfies Property (b.1) but ω does not satisfy Property (b.2), i.e., ω is not an element of $\mathcal{B}(\chi', \chi'')$, and let φ be an element of $\text{HOM}_\chi(G, \mathbb{H}', \omega)$. Since ω does not satisfy Property (b.2), there is $a, b \in \mathbb{X}$ such that at least one of them is not in X' and $ab \in E(G)$ and $(\chi(a), \omega(a), \chi(b), \omega(b)) = (\pi'(\varphi(a)), \omega(a), \pi'(\varphi(b)), \omega(b)) \notin S$. So, $(\varphi(a), \varphi(b))$ is not an edge in \mathbb{H}' and φ is not a homomorphism. Thus $|\text{HOM}_\chi(G, \mathbb{H}', \omega)| = 0$.

Now, for a fixed $\omega \in \mathbb{W}(\chi)$ such that ω satisfies Properties (b.1) and (b.2), i.e., $\omega \in \mathcal{B}(\chi', \chi'')$, one can define a bijection between the set $\mathsf{HOM}_\chi(G, \mathbb{H}', \omega)$ and the set $\text{HOM}_\chi(G, \mathbb{H})$, as follows. Suppose that φ is an element of $HOM_\chi(G, \mathbb{H})$. The mapping $\tau_{\varphi}: a \mapsto \varphi(a)_{\omega(a)}$ is an element of $\mathsf{MAP}_\chi(G, \mathbb{H}, \omega)$. We claim that mapping $\tau : \varphi \mapsto \tau_{\varphi}$ is a bijection between $\mathsf{HOM}_\chi(G, \mathbb{H})$ and $\mathsf{HOM}_\chi(G, \mathbb{H}', \omega)$. In so doing, one should prove that first: τ_{φ} is in $\text{HOM}_{\chi}(G, \mathbb{H}', \omega)$; and second, $\tau: HOM_\chi(G, \mathbb{H}) \to HOM_\chi(G, \mathbb{H}', \omega)$ is injective and surjective.

For any $a, b \in X'$ and $ab \in E(G)$, as ω satisfies Property $(b.1), \omega(a) = \omega(b) =$ 0. Thus, $\tau_{\varphi}(a) = \varphi(a)_{\omega(a)} = \varphi(a)_{0} = \varphi(a)$ and $\tau_{\varphi}(b) = \varphi(b)_{\omega(b)} = \varphi(b)_{0} = \varphi(b)$. Since φ is a homomorphism, $\tau_{\varphi}(a)\tau_{\varphi}(b) = \varphi(a)\varphi(b) \in E(\mathbb{H}) \subseteq E(\mathbb{H}^{\prime})$. Thus the image of ab under τ_{φ} is an edge.

For any $a, b \in X$, such that at least one of a and b is not an element of X' , and $ab \in E(G)$, by definition $\tau_{\varphi}(a) = \varphi(a)_{\omega(a)}$ and $\tau_{\varphi}(b) = \varphi(b)_{\omega(b)}$. Moreover, since $\pi(\varphi(a)) = \chi(a)$ and $\pi(\varphi(b)) = \chi(b)$, as ω satisfies Property (b.2), $(\chi(a), \omega(a), \chi(b), \omega(b)) = (\pi(\varphi(a)), \omega(a), \pi(\varphi(b)), \omega(b)) \in S$. Consequently, by the definition of operator $\alpha_{\vec{\pi},\mathcal{S}}$, as $\varphi(a)\varphi(b) \in E(\mathbb{H})$, then $\varphi(a)_{\omega(a)}\varphi(b)_{\omega(b)}$ is an edge of \mathbb{H}' . Thus for any $\varphi \in \text{HOM}_\chi(G, \mathbb{H})$, τ_φ is a homomorphism. Since $\tau_{\varphi} \in \text{MAP}_{\chi}(G, \mathbb{H}', \omega)$ for every $\varphi \in \text{HOM}_{\chi}(G, \mathbb{H})$, and τ_{φ} is a homomorphism, τ_{φ} is an element of the set $\mathsf{HOM}_\chi(G, \mathbb{H}', \omega)$.

Second step is to prove that τ is injective. Consider $\varphi \neq \psi$ two different elements of $\text{HOM}_\chi(G, \mathbb{H})$. There is an element a in their domain such that $\varphi(a) \neq \psi(a)$. Since $\tau_{\varphi}(a) = \varphi(a)_{\omega(a)} \neq \tau_{\psi}(a) = \psi(a)_{\omega(a)}$ we obtain $\tau_{\varphi} \neq \tau_{\psi}$.

Finally, we should prove that τ is surjective. For any homomorphism $\varphi' \in$ $\mathsf{HOM}_\chi(G, \mathbb{H}', \omega), \mu(\varphi')$ is an element of $\mathsf{HOM}_\chi(G, \mathbb{H})$ (recall that μ is the natural retraction from \mathbb{H}' to \mathbb{H} .). As is easy to see, $\tau_{\mu(\varphi')} = \varphi'$. Thus τ is surjective.

The bijection φ between $\text{HOM}_\chi(G, \mathbb{H})$ and $\text{HOM}_\chi(G, \mathbb{H}', \omega)$ proves that the cardinalities of these two sets are equal. 口

Step II

Recall that $\mathbb{H}'' = \beta_{\overrightarrow{n},\sigma,S}(\mathbb{H})$ and $\mathbb{H}' = \alpha_{\overrightarrow{n},S}(\mathbb{H})$. Let X be a subset of $V(G)$ and let χ be a k-label function from X to [k]. Also, let $\varphi \in \text{HOM}_\chi(G, \mathbb{H}^{\prime\prime})$, and let X' and X'' be the preimages of $V(\mathbb{H})$ and $V(\mathbb{H}') \setminus V(\mathbb{H})$ under φ , respectively. Then, there is a pair of functions $\chi': X' \to [k]$ and $\chi'': X'' \to [k]$ such that $\varphi \in \text{HOM}_{\chi',\chi''}(G,\mathbb{H}').$ Let us call (χ',χ'') the *consistent partition* of φ . To count the elements of the set $\text{HOM}_\chi(G, \mathbb{H}^{\prime\prime})$, we partition this set into smaller sets of homomorphisms that share the same consistent partition, and count the number of elements in the smaller sets.

Let $\mathcal{C}(\chi)$ denote the set of all pairs of functions $(\chi', \chi''), \chi' : X' \to [k], \chi''$: $X'' \to [k], X', X''$ disjoint, that satisfy the following conditions:

- $(c.1)$ $X' \cup X'' = X$, and
- (c.2) $\chi|_{X'} = \chi'$ and $\chi|_{X''} = \sigma(\chi'')$.

We show that $\mathcal{C}(\chi)$ is the set of consistent partitions of the homomorphisms in $\mathsf{HOM}_\chi(G, \mathbb{H}'')$.

Lemma 26. Let χ , G, \mathbb{H}' , and \mathbb{H}'' be defined as above. Then,

$$
\text{hom}_{\chi}(G,\mathbb{H}'') = \sum_{(\chi',\chi'')\in\mathcal{C}(\chi)} |\text{HOM}_{\chi',\chi''}(G,\mathbb{H}')|.
$$
 (3)

Proof. Let $\varphi \in \text{HOM}_X(G, \mathbb{H}^{\prime\prime})$, and let X' and X'' be the preimages of $V(\mathbb{H})$ and $V(\mathbb{H}^{\prime\prime}) \setminus V(\mathbb{H})$ under φ , respectively. Also let $(\chi', \chi''), \chi : X' \to [k], \chi''$: $X'' \to [k]$, be the consistent partition of φ . As is easily seen, (χ', χ'') satisfies Properties (c.1) and (c.2).

Now, let $(\chi', \chi'') \in C(\chi)$, then the set $\text{HOM}_{\chi', \chi''}(G, \mathbb{H}')$ is the set of all elements of $\text{HOM}_\chi(G, \mathbb{H}^{\prime\prime})$ that have (χ', χ'') as their consistent partition. Thus

$$
\mathsf{HOM}_{\chi}(G,\mathbb{H}'') = \bigcup_{(\chi',\chi'')\in\mathcal{C}(\chi)} \mathsf{HOM}_{\chi',\chi''}(G,\mathbb{H}').\tag{4}
$$

Equation [\(3\)](#page-20-1) follows from [\(4\)](#page-20-2).

 \Box

Putting things together

We are now in a position to prove Lemma [24](#page-18-0) and Theorem [18.](#page-13-0)

Proof of Lemma [24.](#page-18-0) By Lemmas [25](#page-19-0) and [26,](#page-20-0)

$$
\begin{aligned} \hom_\gamma(G,\mathbb{H}'')&=\sum_{(\gamma',\gamma'')\in\mathcal{C}(\gamma)}|\mathrm{HOM}_{\gamma',\gamma''}(G,\mathbb{H}')|\\ &=\sum_{(\gamma',\gamma'')\in\mathcal{C}(\gamma)}|\mathcal{B}(\gamma',\gamma'')|\times\hom_{\gamma'\uplus\gamma''}(G,\mathbb{H}),\end{aligned}
$$

where, the first equality follows from Lemma [26](#page-20-0) and the second equality follows from Lemma [25.](#page-19-0)

The cardinality of $\mathcal{C}(\gamma)$ is bounded by $2^{|V(G)|}$. Also, for any $(\gamma', \gamma'') \in \mathcal{C}(\gamma)$ the cardinality of $\mathcal{B}(\gamma', \gamma'')$ is bounded by $(2k)^{V(G)}$ and thus that cardinality is computable in time $O((2k)^{V(G)})$ by brute force. Therefore, the time complexity of calculating

$$
\sum_{(\gamma',\gamma'')\in\mathcal{C}(\gamma)} |\mathcal{B}(\gamma',\gamma'')|\times\hom_{\gamma'\uplus\gamma''}(G,\mathbb{H})
$$

is $O^*((4k)^{|V(G)|})$ and so is the time complexity of computing $\mathsf{hom}_{\gamma}(G, \mathbb{H}^{\prime\prime}).$

Proposition 27. *Let* G *be a graph,* H *a* k*-labeled graph, and let* Φ *be an extended* k -expression for \mathbb{H} . Then, there is an algorithm that calculates $\text{hom}_{\chi}(G, \mathbb{H})$ for *all* k-label functions $\chi : X \to [k]$ *, where* $X \subseteq V(G)$ *, in total time*

$$
\text{size}(\Phi) \times (2(k+1))^{2|V(G)|},
$$

given numbers $\text{hom}_{\chi'}(G, \mathbb{H}')$ *for all label functions* χ' *and k-labeled graphs represented by subexpressions of* Φ*.*

Proof. The base case of induction is when $size(\Phi) = 1$ which takes place when H is a labeled vertex. In this case, there is a brute force algorithm for computing hom_x(G, H), with running time of $O(|V(G)|) \times (k+1)^{|V(G)|}$ that is less than $\mathsf{size}(\Phi)(2(k+1))^{2|V(G)|}.$

If $size(\Phi) > 1$, as Φ is an extended k-expression, there are three possibilities:

- $\Phi = \beta_{\overrightarrow{n},\sigma,\mathcal{S}}(\Phi')$, where Φ' is another extended k-expression;
- $\Phi = \eta_{\mathcal{T}}(\Phi', \Phi'')$, where Φ' and Φ'' are two extended k-expressions;
- $\Phi = \Psi(\Phi')$, where λ is a sequence of relabelling operators and Φ' is another extended k-expression that does not begin with a relabelling operator.

The k-labeled graphs represented by Φ', Φ'' are denoted $\mathbb{H}', \mathbb{H}''$. Let start with the first case. Let $\mathbb{H} = \beta_{\vec{\pi},\sigma,S}(\mathbb{H}')$. Since size(Φ') is less than size(Φ), by induction hypothesis, there is an algorithm that calculates $\textsf{hom}_{\chi}(G, \mathbb{H}')$ for all k-label functions $\chi: X \to [k]$, where $X \subseteq V(G)$, in total time size $(\Phi') \times (2(k +$ $(1))^{2|V(G)|}.$

Suppose that for all k-label functions $\chi: X \to [k]$ such that $X \subseteq V(G)$ the value of $\hom_\chi(G, \mathbb{H}')$ is known. Then by Lemma [24,](#page-18-0) for any k-label function $\chi: X \to [k]$, where $X \subseteq V(G)$, the value $\text{hom}_{\chi}(G, \mathbb{H})$ can be found in time $(2k)^{|V(G)|}$ and, since there are $(k+1)^{V(G)}$ of those functions, $\textsf{hom}_{\chi}(G,\mathbb{H}')$ can also be found for all k-label functions $\chi : X \to [k]$, where $X \subseteq V(G)$, in time $(k+1)^{|V(G)|} \times (2(k+1)^{|V(G)|}) \leq (2(k+1))^{2|V(G)|}.$

So, hom_x(G, H) for all k-label functions $\chi: X \to [k]$, where $X \subseteq V(G)$, can be found in total time $O((\mathsf{size}(\Phi')+1) \times (2(k+1))^{|V(G)|})$, that is, bounded by size $(\Phi)\times (2(k+1))^{2|V(G)|}.$

The proof for the second and third cases is similar with Lemma [22](#page-16-0) and Lemma [20](#page-14-0) in place of Lemma [24.](#page-18-0) 口

Proof of Theorem [18.](#page-13-0) By Observation [19,](#page-14-1) $hom(G, H)$ equals the sum of $hom_v(G, \mathbb{H})$ over all k-label functions $\chi : V(G) \to [k]$. By Proposition [27,](#page-21-0) the time required to compute $\textsf{hom}_\chi(G,\mathbb{H})$ for all k-label functions $\chi: V(G) \to [k]$ is $O(\textsf{size}(\Phi) \times (2(k+1))^{2|V(G)|})$. Also, the time for summing up over $k^{V(G)}$ numbers is $O(k^{V(G)})$. Thus the total running time can be bounded by

$$
O(\text{size}(\Phi) \times (2(k+1))^{2|V(G)|}) + O(k^{V(G)}) = O(\text{size}(\Phi) \times (2(k+1))^{2|V(G)|}).
$$

4 Other examples of plain exponential classes

In this section we provide two plain exponential classes of graphs. We start with subdivisions of cliques.

4.1 Subdivided Cliques

The *subdivision* of an edge uv by a graph H is a graph with vertex set $V(H)$ ∪ $\{u, v\}$ and edge set $E(H) \cup \bigcup \{ut, vt\}$. The subdivision of a graph G by $t\in V(H)$

a graph H is the graph obtained by replacing every edge uv of G with its subdivision by a copy of H (a disjoint copy for each edge). Let $\mathcal{K}(\mathcal{H})$ denote the class of subdivisions of cliques by graphs from a class H .

The following theorem is the main result of this section.

Theorem 28. Let H be a plain exponential class of graphs. Then $K(H)$ is also *plain exponential.*

More precisely, if $\#\text{GRAPHHom}(-, \mathcal{H})$ *can be solved in time* $O^*(c^{|V(G)|+|V(H)|}),$ *c constant, for any given graphs* G *and* $H \in \mathcal{H}$, *then* $\#\text{GRAPHHom}(-, \mathcal{K}(\mathcal{H}))$ *can be solved in time* $O^*(c_1^{2(|V(G)|+|V(H)|})$ *, where* $c_1 = \max(c, 2)$ *.*

The following theorem from [\[13\]](#page-31-6) is our main technical tool. This theorem gives an upper bound on time complexity of the generic problem of summing over the partitions of n elements into a given number of weighted subsets. We translate our counting problem as summation of this kind and use Theorem [29](#page-23-0) to prove an upper bound for time complexity of our counting problem.

Theorem 29 ([\[13\]](#page-31-6)). *Let* M *be a* n*-element set and* f *a function from the power set of* M *to the set of real numbers. Also, suppose that for any subset* $X \subseteq M$, *the value of* $f(X)$ *can be obtained in time* $O(1)$ *. Let*

$$
\operatorname{par}(f) = \sum_{P} \prod_{i=1}^{n} f(P_i) \tag{5}
$$

where, $P = (P_1, ..., P_n)$ *runs through all ordered partitions of* M *into* k disjoint *subsets of* M *(*k *fixed). There is an algorithm that calculate* par(f) *in time* $O^*(2^{|M|}).$

With specific choices of function f this class of partition problems expresses various counting problems. For instance, to compute the number of n -colourings of a given graph G we let $f(S) = 1$ if the vertex subset S is an independent set and 0 otherwise. As is easily seen, for any n-partition P of $V(G)$ such that for each $i \in [n]$, P_i is an independent set, P can be associated with a proper ncolouring of G and vice versa. Also $\text{par}(f)$ is equal to the number of *n*-partitions of $V(G)$ that P_i is an independent set for each $i \in [n]$. Thus, the number of *n*-colouring of G equals $\text{par}(f)$.

Here, we count the number of H -colorings of a graph G by expressing this number as $\textsf{par}(f)$ for some function f. Without loss of generality, we can assume that G is a connected graph as otherwise, the number of homomorphisms from G to H is the product of the numbers of homomorphisms from the connected components of G to H .

Let H be the subdivision of K_n by a graph $U \in \mathcal{H}$. We denote the set of all vertices in H that are originally from $V(K_n)$ by $H_A = \{v_1, ..., v_n\}$ and the set of the remaining vertices of H by $H_B = \bigcup_{i,j \in [n], i \neq j} \{V(U_{ij})\}$, where U_{ij} is the copy of U subdividing the edge $v_i v_j$. Let (A, B) be a 2-partition of $V(G)$. A homomorphism φ from G to H is said to be *consistent* with (A, B) if the image of A under φ is a subset of H_A and the image of B under φ is a subset of H_B .

Let C_B be the set of all connected components of $G[B]$. Also let $\mathbf{C} \in C_B$ and let $N(\mathbf{C})$ be the set of all elements of A that are adjacent to at least one vertex from C. Unless $A = \emptyset$, the set $N(C)$ is not empty as otherwise it contradicts the assumption that G is connected. Then we fix an arbitrary vertex of $N(\mathbf{C})$ and refer to it as $s_{\mathbf{C}}$.

Let φ be an H-colouring of G. Then it is easy to see that vertices from G that are mapped by φ to vertices of H_A should be an independent set. Thus we use brute force to consider all 2-partitions (A, B) of $V(G)$, where A is an independent set and we count the number of homomorphisms from G to H that are consistent with each (A, B) and then sum those numbers up to get the total number of homomorphisms from G to H .

Let $C \in C_B$. Since any homomorphism φ consistent with (A, B) should map C to H_B and the image of C under φ should be connected, the image of C under φ is a subset of one of the copies of U. Also, the image under φ of all vertices in A adjacent to C is a set of at most two elements as otherwise it is not possible for image of C to be adjacent to the image of all elements of $N(C)$.

Thus, one can associate φ with a relation denoted $\varphi' \subseteq (A \cup C_B) \times V(H_A)$ that contains pairs $(a, \varphi(a)), a \in A$, and (C, b) , where $C \in C_B$ and $b \in \varphi(N(C))$. Relation φ' defined this way may not be a function because it may relate a component C to more than one element of $V(H_A)$. To fix this, we introduce two copies of C denoted \mathbf{C}^0 and \mathbf{C}^1 . Since the image of $N(\mathbf{C})$ under φ contains at most 2 elements, φ' relates each copy of C to one of those elements of the image of $N(\mathbf{C})$ under φ . Now, let $S = A \cup \bigcup_{\mathbf{C} \in C_B} {\{ \mathbf{C}^0, \mathbf{C}^1 \}}$, i.e., S includes the vertices of A along with two copies of elements of C_B . Thus we can associate homomorphisms from G to H that are consistent with (A, B) with a relation from S to $V(H_A)$.

In order to convert φ' into a mapping, for any φ we require the associated φ' to associate \mathbb{C}^1 and $s_{\mathbb{C}}$ with the same element of $V(H_A)$ and \mathbb{C}^0 with the other element if the image of $N(\mathbf{C})$ contains two vertices. Restricted this way φ' is a mapping $\varphi' : S \to V(H_A)$.

As any mapping from S to $V(H_A)$ naturally corresponds to a *n*-partition of S, we can associate homomorphisms from $V(G)$ to $V(H)$ that are consistent with (A, B) with *n*-partitions of S. For an *n*-partition $P = (P_1, \ldots, P_n)$ of S, we define *mappings compatible* with P as follows. A mapping $\varphi : V(G) \to V(H)$ is compatible with P if it satisfies the following properties:

- (A) φ maps elements of $P_i \cap A$ to v_i for each $i \in [n]$.
- (B) For every $C \in C_B$, such that $C^0 \in P_i$ and $C^1 \in P_j$, for $i \neq j$, the mapping $\varphi|_{\mathbf{C}}$ is a homomorphism that maps \mathbf{C} to U_{ij} .
- (C) For every $\mathbf{C} \in C_B$ such that $\mathbf{C}^0, \mathbf{C}^1 \in P_i$, $\varphi|_{\mathbf{C}}$ is a homomorphism that maps C to U_{ij} , where $j \in [n] - \{i\}$ is arbitrary.

We define function f from the power set of S to natural numbers as follows. First, f is set to 0 for subsets $X \subseteq S$ such that if X is a class of a partition P, say, $X = P_i$, then there are no homomorphisms compatible with P. So suppose P has at least one homomorphism compatible with it. Then for any $C \in C_B$, the class of P that contains $a \in N(\mathbb{C})$ also contains one of \mathbb{C}^0 or \mathbb{C}^1 . So, for $X \subseteq S$,

 $f(X) = 0$ if there is $a \in X$ and $\mathbf{C} \in C_B$ such that $a \in N(\mathbf{C})$ but neither of \mathbb{C}^0 or \mathbb{C}^1 is in the set X.

Also, if P contains at least one of \mathbb{C}^0 or \mathbb{C}^1 in one of its classes for some $\mathbf{C} \in C_B$, then there should be at least one of vertex from $N(\mathbf{C})$ in the same subset as otherwise P would have no compatible homomorphisms. Thus for $X \subseteq S$,

 $f(X) = 0$ if for some $C \in C_B$, $C^1 \in X$ and/or $C^0 \in X$ but none of the elements of $N(\mathbf{C})$ is in X.

Recall that the n-partition compatible with a homomorphism is supposed to have \mathbb{C}^1 and $s_{\mathbb{C}}$ in the same class of P. Therefore, for $X \subseteq S$,

 $f(X) = 0$ if for some $\mathbf{C} \in C_B$, $\mathbf{C}^1 \in X$ and $s_{\mathbf{C}} \notin X$.

Now we need to set the value of f for subsets that can be a class in a partition with compatible homomorphisms. For $X \subseteq S$,

• if none of the above is true for X, $f(X) = x \cdot y$, where

$$
x = \prod_{\mathbf{C}: \mathbf{C}^0, \mathbf{C}^1 \in X} (n-1) \text{hom}(\mathbf{C}, U), \text{ and}
$$

$$
y = \prod_{\mathbf{C}: \mathbf{C}^1 \in X, \mathbf{C}^0 \notin X} \text{hom}(\mathbf{C}, U).
$$

We defer an explanation of this equality to the proof of the following

Lemma 30. *The number of homomorphisms from* G *to* H *that are consistent with* (A, B) *equals*

$$
\sum_{P} \prod_{i=1}^{n} f(P_i),\tag{6}
$$

where $P = (P_1, ..., P_n)$ *runs through all n-partitions of* S.

Proof. To prove this lemma, we prove that the *n*-partitions of S induce a partition on the set of homomorphisms from G to H that are consistent with (A, B) , by grouping all homomorphism according to their compatible n-partitions. Then, we show that for each *n*-partition P of S , the number of homomorphisms compatible with P equals $\prod_{i=1}^{n} f(P_i)$. These two claims prove the lemma.

Let P be a fixed n-partition of S such that $\prod_{i=1}^{n} f(P_i) \neq 0$, and let φ be a mapping compatible with P. As is easily seen, φ is a mapping from G to H that is consistent with (A, B) . Since A is an independent set, $\varphi|_A$ is a homomorphism for any mapping φ . By Properties (B) and (C), for any connected component $\mathbf{C} \in C_B$, $\varphi|_{V(\mathbf{C})}$ is a homomorphism hence $\varphi|_B$ is a homomorphism as well. Now let $a \in A$ and $b \in V(\mathbf{C})$, where $\mathbf{C} \in C_B$. Since $\prod_{i=1}^n f(P_i) \neq 0$ the possible cases are: (1) $a, \mathbf{C}^0, \mathbf{C}^1 \in P_i$ for some $i \in [n]$; (2) $a, \mathbf{C}^1 \in P_i$ and $\mathbf{C}^0 \in P_j$ such that $i \neq j$, if $a = s_{\mathbf{C}}$, and $\mathbf{C}^1 \in P_i$ and $a, \mathbf{C}^0 \in P_j$ otherwise. In case (1) by Property 3 $\varphi(\mathbf{C})$ is in graph U_{ij} for some $0 \leq j \leq n$, and by Property (A), $\varphi(a)$ is v_i , and since there is an edge between any vertex of U_{ij} and v_i , $\varphi(a)\varphi(b)$ is an edge of H . The argument in case (2) goes in the similar way. So, any mapping compatible with partition P is a homomorphism that is consistent with (A, B) .

Now, we prove that for any homomorphism φ from G to H that is consistent with (A, B) , there is exactly one *n*-partition P of S such that φ is compatible with P. Let us fix a homomorphism φ from G to H that is consistent with (A, B) . By Properties (A) – (C) of the compatible homomorphism, an *n*-partition P with which φ is compatible should satisfy the following properties

(a) For any $a \in A$, $a \in P_i$ if and only if $\varphi(a) = v_i$.

- (b) For any $C \in C_B$ such that the image of $N(C)$ under φ is $\{v_i, v_j\}$ for $1 \leq i \neq j \leq n$, and $\varphi(s_{\mathbf{C}}) = v_i$ it holds $\mathbf{C}^0 \in P_j$ and $\mathbf{C}^1 \in P_i$. Note that \mathbf{C}^0 and \mathbf{C}^1 are not interchangeable because $s_{\mathbf{C}}$ and \mathbf{C}^1 should be included in the same class of P.
- (c) For any $\mathbf{C} \in C_B$ such that the image of $N(\mathbf{C})$ under φ is $\{v_i\}$, it holds that $\mathbf{C}^0, \mathbf{C}^1 \in P_i$.

As is easily seen, there is at least one n-partition that satisfies all of the above properties. On the other hand, no two n-partitions of S can satisfy all the above properties because the properties specify for each of elements of S which subset of the compatible partition to go, uniquely. Thus, there is exactly one *n*-partition of S that is compatible with φ .

Let P be an n-partition of S and \hat{P} be the set of all homomorphisms compatible with P. We show that $|\hat{P}| = \prod_{i=1}^{n}$ $\prod_{i=1} f(P_i)$. For any connected component $\mathbf{C} \in C_B$, such that $\mathbf{C}^0 \in P_i$ and $\mathbf{C}^1, s_{\mathbf{C}} \in P_j$ for $i \neq j$, hom (\mathbf{C}, U) is a contributing factor to $f(P_j)$. Note that $\text{hom}(\mathbf{C}, U)$ does not contribute anything to $f(P_i)$. Also, for any connected component $\mathbf{C} \in C_B$, such that $\mathbf{C}^0, \mathbf{C}^1 \in P_i$ for some $i \in [n]$, the value $f(P_i)$ contains a factor $(n-1)$ hom (C, U) . Thus,

$$
\prod_{i=1}^{n} f(P_i) = \prod_{\substack{\mathbf{C}: \mathbf{C}^0, \mathbf{C}^1 \in P_i \\ \text{for some } i \in [n]}} (n-1) \text{hom}(\mathbf{C}, U) \times \prod_{\substack{\mathbf{C}: \mathbf{C}^1 \in P_i, \mathbf{C}^0 \notin P_i \\ \text{for some } i \in [n]}} \text{hom}(\mathbf{C}, U). \tag{7}
$$

On the other hand by Property (a), $\varphi|_A$ is the same for every $\varphi \in \hat{P}$. Let $\mathbf{C} \in C_B$. Let $\varphi_1, \varphi_2 \in \hat{P}$ be two homomorphisms such that $\varphi_1|_{V(\mathbf{C})}$ and $\varphi_2|_{V(\mathbf{C})}$ are not identical. Then, let $\varphi: V(G) \to V(H)$ be defined as follows:

$$
\varphi(v) = \begin{cases} \varphi_1(v), & \text{if } v \in V(G) \setminus V(\mathbf{C}), \\ \varphi_2(v), & \text{if } v \in V(\mathbf{C}). \end{cases}
$$

Mapping φ is a homomorphism as $\varphi_1|_A$ and $\varphi_2|_A$ are the same and also C is disconnected from the $V(G) \setminus (V(\mathbf{C}) \cup A)$. Thus, the number of compatible homomorphisms of partition P can be obtained by multiplying the number of all different mappings we can get by restricting those homomorphisms to a component $\mathbf{C} \in C_B$, over all $\mathbf{C} \in C_B$. Therefore,

$$
|\hat{P}| = \prod_{\mathbf{C} \in C_B} \left| \{ \varphi|_{V(\mathbf{C})}: \varphi \in \hat{P} \} \right|
$$

=
$$
\prod_{\substack{\mathbf{C}: \mathbf{C}^0, \mathbf{C}^1 \in P_i \\ \text{for some } i \in [n]}} \left| \{ \varphi|_{V(\mathbf{C})}: \varphi \in \hat{P} \} \right| \times \prod_{\substack{\mathbf{C}: \mathbf{C}^1 \in P_i, \mathbf{C}^0 \notin P_i \\ \text{for some } i \in [n]}} \left| \{ \varphi|_{V(\mathbf{C})}: \varphi \in \hat{P} \} \right|.
$$

Let again $\mathbf{C} \in C_B$, such that $\mathbf{C}^0 \in P_i$ and $\mathbf{C}^1 \in P_j$ for $i \neq j$. By Property (a) every $\varphi \in \hat{P}$ maps $P_i \cap A$ to v_i and since φ is homomorphism, it maps C to U_{ij} . Therefore,

$$
\left| \{ \varphi|_{V(\mathbf{C})}: \varphi \in \hat{P} \} \right| = \hom(\mathbf{C}, U).
$$

Now let $\mathbf{C} \in C_B$ such that $\mathbf{C}^0, \mathbf{C}^1 \in P_i$. By Property (C), the image of \mathbf{C} under φ can be any of U_{ij} where $j \in [n] - \{i\}$. Thus,

$$
\left| \{ \varphi|_{V(\mathbf{C})}: \varphi \in \hat{P} \} \right| = (n-1)\text{hom}(\mathbf{C},U).
$$

Therefore,

$$
|\hat{P}| = \prod_{\substack{\mathbf{C}: \mathbf{C}^0, \mathbf{C}^1 \in P_i \\ \text{for some } i \in [n]}} (n-1) \text{hom}(\mathbf{C}, U) \times \prod_{\substack{\mathbf{C}: \mathbf{C}^1 \in p, \mathbf{C}^0 \notin P_i \\ \text{for some } i \in [n]}} \text{hom}(\mathbf{C}, U) \tag{8}
$$

By [\(8\)](#page-27-0) and [\(7\)](#page-26-0), $|\hat{P}| = \prod_{n=1}^{\infty}$ $\prod_{i=1} f(P_i).$

Hence, the number of homomorphisms from G to H that are consistent \prod^n with (A, B) is equal to Σ $\prod_{i=1} f(P_i)$, where $P = (P_1, ..., P_n)$ runs through all P n-partitions of S. \Box

Now, we are in a position to prove Theorem [28](#page-22-0)

Proof of Theorem [28.](#page-22-0) We brute force over all partitions of set $V(G)$ into two sets A and B, and for each partition (A, B) of $V(G)$, we calculate function f with respect to (A, B) and then we find the par(f). Let us fix a partition (A, B) of $V(G)$. The time needed to calculate function f with respect to (A, B) , over all subsets of S is bounded by $O^*(c^{|V(G)|})$ as the time to calculate $\mathsf{hom}(\mathbb{C}, U)$ is bounded by $O(c^{|V(C)|})$ and there are at most $|V(G)|$ of them. By Lemma [30](#page-25-0) the number of homomorphisms from G to H that are consistent with (A, B) can be represented as $\text{par}(f)$ for the corresponding f. By Theorem [29,](#page-23-0) the total time to calculate $\textsf{par}(f)$ is bounded by $O^*(2^{|V(G)|})$. Thus, the time complexity to calculate the number of homomorphisms from G to H that are consistent with (A, B) is bounded by $O^*(c_1^{|V(G)|}), c_1 = \max(c, 2)$.

The value $\textsf{hom}(G, H)$ equals the sum of numbers of homomorphisms from G to H that are consistent with (A, B) over all 2-partitions (A, B) of $V(G)$. Since there are $2^{V(G)}$ different partitions (A, B) and for any such partition the time complexity of counting the number of homomorphisms from G to H is bounded by $O^*(c_1^{|V(G)|})$, there is a algorithm that computes hom (G, H) in time $O^*(c_1^{2|V(G)|}).$

By Theorem 5.4 from [\[15\]](#page-31-11) it follows that $\mathcal{K}(\mathcal{H})$ does not always have bounded clique width; for instance when H contains only one graph that is a single vertex.

4.2 Kneser Graphs

Kneser graphs give another example of a plain exponential class of graphs.

The *Kneser graph* $KG_{n,k}$ is the graph whose vertex set is the set of k-element subsets of a set of n elements, and two vertices are adjacent if and only if the two corresponding sets are disjoint. By \mathcal{KG}_k we denote the class of all Kneser graphs for a fixed k . The main result of this section is the following

Theorem 31. For any k , \mathcal{KG}_k is plain exponential.

More precisely, there is an algorithm that given a graph G *and* $n \in \mathbb{N}$ *finds* $\mathsf{hom}(G,\mathrm{KG}_{n,k})$ *in time* $O^*(2^{k|V(G)|}).$

Remark 32. *The running time of our algorithm is not plain exponential if* k *is not a constant. Thus, Theorem [31](#page-28-0) does not guarantee that the class of all Kneser graphs irrespective of the parameter* k *is plain exponential. Proving or disproving it remains an open problem.*

We use the following fact.

Theorem 33 ([\[13\]](#page-31-6)). *There is an algorithm that given* k *and a graph* G *counts the number of k-colouring of graph* G *in time* $O^*(2^{|V(G)|})$ *.*

Proof of Theorem [31.](#page-28-0) Let $G^{(k)}$ denote the graph obtained by replacing each of its vertices with a clique of size k and replacing each of its edges with a complete bipartite graph on $k+k$ vertices. For $a \in V(G)$ let $\psi(a)$ denote the set of vertices of the clique replacing v in $G^{(k)}$.

First, we introduce a many to one correspondence between elements of $\mathsf{HOM}(G^{(k)}, K_n)$ and $\mathsf{HOM}(G,\mathrm{KG}_{n,k})$. Let $\tau: \mathsf{HOM}(G^{(k)}, K_n) \to \mathsf{HOM}(G,\mathrm{KG}_{n,k})$ be defined by setting $\tau(\varphi): V(G) \to \mathrm{KG}_{n,k}$ to be the mapping $v \mapsto {\varphi(u)}|u \in$ $\psi(v)$. Notice that the cardinality of $\{\varphi(u) | u \in \psi(v)\}$ equals k because $G^{(k)}[\psi(v)]$ is a k-clique and φ is a homomorphism from $G^{(k)}$ to K_n . Therefore $\tau(\varphi)(v)$ is always a vertex of $KG_{n,k}$.

Next, we prove that for $\varphi \in \text{HOM}(G^{(k)}, K_n), \varphi^* = \tau(\varphi)$ is a homomorphism. Let $ab \in E(G)$ then $\varphi^*(a) = {\varphi(u) | u \in \psi(a)}$ and $\varphi^*(b) = {\varphi(u) | u \in \psi(b)}$. Since $G^{(k)}[\psi(a) \cup \psi(b)]$ is a clique, and φ is a homomorphism $\varphi^*(a)$ and $\varphi^*(b)$ are two disjoint set thus $\varphi^*(a)\varphi^*(b)$ is an edge of KG_{n,k}. Therefore $\varphi^* \in$ $HOM(G, KG_{n,k}).$

Next, we prove that for any element σ of HOM($G, KG_{n,k}$), $\tau(\varphi) = \sigma$ for exactly $(k!)^{|V(G)|}$ homomorphisms $\varphi \in \text{HOM}(G^{(k)}, K_n)$. Take the set of all $\varphi \in \text{MAP}(G^{(k)}, K_n)$ such that for any $v \in V(G)$ the image of $\psi(v)$ under φ equals $\sigma(v)$. First, there are $(k!)^{|V(G)|}$ of them as for a fixed σ and a fixed $v \in V(G)$, the image of any $G[\psi(v)]$ under such a mapping, has k! different possibilities. Second, observe that for any $\varphi \in \mathsf{MAP}(G^{(k)}, K_n)$ such that for any $v \in V(G)$ the image of $\psi(v)$ under φ equals $\sigma(v)$, φ is a homomorphism, we have $\tau(\varphi) = \sigma$. Hence, φ is a surjective $(k!)^{|V(G)|}$ to 1 mapping from $\textsf{HOM}(G^{(k)}, K_n)$ to $\text{HOM}(G, \text{KG}_{n,k})$. Therefore

$$
|\text{HOM}(G, \text{KG}_{n,k})| = \frac{\text{HOM}(G^{(k)}, K_n)}{(k!)^{|V(G)|}}.
$$

Since there is an algorithm that computes $\mathsf{hom}(G^{(k)}, K_n)$ in time $O^*(2^{k|V(G)|}),$ there is an algorithm that computes $\text{hom}(G, \text{KG}_{n,k})$ in the same time. \Box

Next we show that \mathcal{KG}_k does not have bounded clique width.

Theorem 34. *The class* \mathcal{KG}_2 *of Kneser graphs does not have bounded clique width.*

The next theorem is the main ingredient.

Theorem 35 ([\[2\]](#page-30-4)). Let \mathcal{H} be a class of graphs with bounded clique width. Then, *the class of complements of graphs from* H *is also of bounded clique width.*

Proposition 36. *The class of complements of graphs of* \mathcal{KG}_2 *does not have bounded clique width.*

Proof. Let k be a constant, let n be a sufficiently large number, let V be a set of cardinality n, and, for the sake of contradiction, let Φ_0 be a k-expression for the complement of a Kneser graph $KG_{n,2}$ whose vertices are 2-element subsets of V. In particular, this means that each vertex of $KG_{n,2}$ receives one of the k labels. We define a finite sequence of k-expressions $\{\Phi_i\}$ which starts with k -expression Φ_0 .

For $i \geq 0$, the $(i + 1)$ -th element of the sequence is defined from the *i*-th element of the sequence as follows: if Φ_i represents a graph with more than 1 vertices, it has the form $\Psi_i(\Phi_{i,l} \bigoplus \Phi_{i,r})$, where Ψ_i is a sequence of recolouring and connecting operators and $\Phi_{i,l}$ and $\Phi_{i,r}$ are two k-expressions. Then, let Φ_{i+1} be the k-expression from $\{\Phi_{i,l}, \Phi_{i,r}\}\$ that represents the graph with greatest number of vertices. Thus, if G_i is the graph represented by Φ_i and G_{i+1} the graph represented by Φ_{i+1} , then $|V(G_i)| \geq |V(G_{i-1})|/2$.

Let graph G be a subgraph of G_0 . For $v \in V$, let *count* of v in graph G, denoted by $c(v)$, be the number of vertices from G of which v is an element. Note that each vertex of G_0 is a 2-elements set of elements in V. Let G_i be the first graph in the sequence such that the number of vertices whose count in G_i is $n-1$ is less than $c_0 = (n-1)/k$. There are two cases: first, in G_i the count of every vertex is less than $n-1$, and second, there is at least one vertex in V whose count in G_i is $n-1$. Also note that the count of any $v \in V$ in G_0 or any of its subgraphs, is less than n .

Let A denote the set of all elements of V whose count is $n-1$ in G_{i-1} . Then, $|A| \geq c_0$. For each element of A there are $n-1$ different vertices of G_{i-1} which contain it. Thus the sum of $c(v)$ over all $v \in A$ is $(n-1)|A|$ which is number of all vertices of G_{i-1} that have one of their elements in A and the other element in $V \setminus A$, plus twice the number of vertices of G_{i-1} that have both elements in A. Thus the number of vertices of G_{i-1} that have at least one element in A equals $(n-1)|A| - {A \choose 2}$. Observe that $|V(G_{i-1})|$ is greater than the number of those vertices of G_{i-1} that contain an element from A. As $\frac{n-1}{k} \leq |A| \leq n$,

$$
|V(G_{i-1})| \ge (n-1)|A| - \binom{|A|}{2} = |A| \left(n - 1 - \frac{|A| - 1}{2} \right) \ge \frac{(n-1)^2}{2k}.
$$

Therefore

$$
|V(G_i)| \ge |V(G_{i-1})| \setminus 2 \ge \frac{(n-1)^2}{4k}.
$$

Let v_1 be an element of V whose count in G_i is the largest among all elements of V . By the Pigeonhole principle,

$$
c(v_1) \ge \frac{|V(G_i)|}{n} \ge \frac{2(n-1)^2}{4nk}.
$$

Let $G_i(v)$ be the set of all vertices of G_i that have v_1 as an element. Again, by the Pigeonhole principle, there are at least $c(v_1)/k$ vertices of $G_i(v_1)$ that are labeled with the same label. So, for sufficiently large n there are at least 3 vertices in G_i that all have v_1 as an element and are labeled with the same label in G_i . Now, let $\{v_1, v_2\}$, $\{v_1, v_3\}$, and $\{v_1, v_4\}$ be different elements of $G_i(v_1)$ that are labeled with the same label.

Now, let us consider the case when the count of each vertex of V is less than $n-1$ in G_i . In this case there is $v_5 \in V$ such that $\{v_2, v_5\} \notin V(G_i)$. Clearly, $v_5 \neq v_3$ or $v_5 \neq v_4$. Without loss of generality, assume $v_5 \neq v_3$.

In the case when there are vertices with count $n-1$ in G_i , let $v_1 \in V$ be an element whose count in G_i is $n-1$. Then, by the Pigeonhole principle, there are $(n-1)/k$ elements of $G_i(v_1)$ that have the same label. Since there are at least $(n-1)/k$ elements of $G_i(v_1)$ with the same label, and there are at most c_0 elements of V with count $n-1$ in G_i , there are $\{v_1, v_2\}$, $\{v_1, v_3\}$, $\{v_1, v_4\}$ ∈ $V(G_i)$, such that $\{v_1, v_2\}$, $\{v_1, v_3\}$, and $\{v_1, v_4\}$ are labeled the same label in G_i and also count of v_2 is less than $n-1$ in G_i . Therefore, there is $v_5 \in V$ such that $\{v_2, v_5\} \notin V(G_i)$ and (say) $v_5 \neq v_3$.

Thus, in both cases, there are $v_1, v_2, v_3, v_5 \in V$ such that $\{v_1, v_2\}$ and $\{v_1, v_3\}$ are from $V(G_i)$, and also are labeled with the same label, and $\{v_2, v_5\} \notin V(G_i)$, and $v_5 \neq v_3$. Since G_i is represented by a subexpression of Φ_0 and $\{v_2, v_5\} \notin G_i$, the vertex $\{v_2, v_5\}$ should connect to $\{v_1, v_2\}$ at some later point. Since $\{v_1, v_2\}$ and $\{v_1, v_3\}$ both have the same label, any operator η_{ij} that connects $\{v_2, v_5\}$ to $\{v_1, v_2\}$, also connects $\{v_2, v_5\}$ to $\{v_1, v_3\}$ that is not an edge of G_0 . A contradiction. П

References

- [1] László Babai. Graph isomorphism in quasipolynomial time [extended abstract]. In *STOC*, pages 684–697, 2016.
- [2] Alexandre Blanch´e, Konrad K. Dabrowski, Matthew Johnson, Vadim V. Lozin, Daniël Paulusma, and Viktor Zamaraev. Clique-width for graph classes closed under complementation. In *MFCS*, pages 73:1–73:14, 2017.
- [3] Jianer Chen, Xiuzhen Huang, Iyad A. Kanj, and Ge Xia. Strong computational lower bounds via parameterized complexity. *J. Comput. Syst. Sci.*, 72(8):1346–1367, 2006.
- [4] Marek Cygan, Fedor V. Fomin, Alexander Golovnev, Alexander S. Kulikov, Ivan Mihajlin, Jakub Pachocki, and Arkadiusz Socala. Tight bounds for graph homomorphism and subgraph isomorphism. In *SODA*, pages 1643– 1649, 2016.
- [5] Martin E. Dyer and Catherine S. Greenhill. The complexity of counting graph homomorphisms. *Random Struct. Algorithms*, 17(3-4):260–289, 2000.
- [6] Fedor V. Fomin, Serge Gaspers, and Saket Saurabh. Improved exact algorithms for counting 3- and 4-colorings. In *COCOON*, pages 65–74, 2007.
- [7] Fedor V. Fomin, Alexander Golovnev, Alexander S. Kulikov, and Ivan Mihajlin. Lower bounds for the graph homomorphism problem. In *ICALP*, pages 481–493, 2015.
- [8] Fedor V. Fomin, Pinar Heggernes, and Dieter Kratsch. Exact algorithms for graph homomorphisms. *Theory Comput. Syst.*, 41(2):381–393, 2007.
- [9] Martin Grohe. The complexity of homomorphism and constraint satisfaction problems seen from the other side. *J. ACM*, 54(1):1:1–1:24, 2007.
- [10] Pavol Hell and Jaroslav Nešetřil. *Graphs and homomorphisms*, volume 28 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, 2004.
- [11] Pavol Hell and Jaroslav Nešetřil. On the complexity of H-coloring. *Journal of Combinatorial Theory, Ser.B*, 48:92–110, 1990.
- [12] Russell Impagliazzo and Ramamohan Paturi. Complexity of k-sat. In *CCC*, pages 237–240, 1999.
- [13] Mikko Koivisto. An $o^*(2^n)$ algorithm for graph coloring and other partitioning problems via inclusion–exclusion. In *FOCS*, pages 583–590, 2006.
- [14] Daniel Lokshtanov, Dániel Marx, and Saket Saurabh. Lower bounds based on the exponential time hypothesis. *Bulletin of the EATCS*, 105:41–72, 2011.
- [15] Johann A. Makowsky and Udi Rotics. On the clique-width of graphs with few p4's. *International Journal of Foundations of Computer Science*, 10(03):329–348, 1999.
- [16] Patrick Traxler. The time complexity of constraint satisfaction. In *IWPEC*, pages 190–201, 2008.
- [17] Magnus Wahlström. New plain-exponential time classes for graph homomorphism. In *CSR*, pages 346–355, 2009.