Classifying k-Edge Colouring for H-free Graphs*

Esther Galby¹, Paloma T. Lima², Daniël Paulusma³, and Bernard Ries¹

Abstract. A graph is H-free if it does not contain an induced subgraph isomorphic to H. For every integer k and every graph H, we determine the computational complexity of k-EDGE COLOURING for H-free graphs.

1 Introduction

A graph G = (V, E) is k-edge colourable for some integer k if there exists a mapping $c : E \to \{1, ..., k\}$ such that $c(e) \neq c(f)$ for any two edges e and f of G that have a common end-vertex. The chromatic index of G is the smallest integer k such that G is k-edge colourable. Vizing proved the following classical result.

Theorem 1 ([20]). The chromatic index of a graph G with maximum degree Δ is either Δ or $\Delta + 1$.

The EDGE COLOURING problem is to decide if a given graph G is k-edge colourable for some given integer k. Note that (G, k) is a yes-instance if G has maximum degree at most k-1 by Theorem 1 and that (G, k) is a no-instance if G has maximum degree at least k+1. If k is fixed, that is, k is not part of the input, then we denote the problem by k-EDGE COLOURING. It is trivial to solve this problem for k=2. However, the problem is NP-complete if $k \geq 3$, as shown by Holyer for k=3 and by Leven and Galil for $k \geq 4$.

Theorem 2 ([10,14]). For $k \geq 3$, k-Edge Colouring is NP-complete even for k-regular graphs.

Due to the above hardness results we may wish to restrict the input to some special graph class. A natural property of a graph class is to be closed under vertex deletion. Such graph classes are called hereditary and they form the focus of our paper. To give an example, bipartite graphs form a hereditary graph class, and it is well-known that they have chromatic index Δ . Hence, EDGE COLOURING is polynomial-time solvable for bipartite graphs, which are perfect and triangle-free. In contrast, Cai and Ellis [4] proved that for every $k \geq 3$, k-EDGE COLOURING is NP-complete for k-regular comparability graphs, which are also perfect. They also proved the following two results, the first one of which shows that EDGE COLOURING is NP-complete for triangle-free graphs (the graph C_s denotes the cycle on s vertices).

Theorem 3 ([4]). Let $k \geq 3$ and $s \geq 3$. Then k-EDGE COLOURING is NP-complete for k-regular C_s -free graphs.

Theorem 4 ([4]). Let $k \geq 3$ be an odd integer. Then k-EDGE COLOURING is NP-complete for k-regular line graphs of bipartite graphs.

It is also known that EDGE COLOURING is polynomial-time solvable for chordless graphs [16], series-parallel graphs [11], split-indifference graphs [19] and for graphs of treewidth at most k for any constant k [1].

It is not difficult to see that a graph class \mathcal{G} is hereditary if and only if it can be characterized by a set $\mathcal{F}_{\mathcal{G}}$ of forbidden induced subgraphs (see, for example, [12]). Malyshev determined the complexity of 3-EDGE COLOURING for every hereditary graph class \mathcal{G} , for which $\mathcal{F}_{\mathcal{G}}$ consists of graphs that each have at most five vertices, except perhaps two graphs that may contain six vertices [17]. Malyshev performed a

Department of Informatics, University of Fribourg, Switzerland, {esther.galby,bernard.ries}@unifr.ch
Department of Informatics, University of Bergen, Norway, paloma.lima@uib.no
Department of Computer Science, Durham University, UK, daniel.paulusma@durham.ac.uk

^{*} The second author was supported by the Research Council of Norway via the project CLASSIS. The third author was supported by the Leverhulme Trust (RPG-2016-258).

similar complexity study for EDGE COLOURING for graph classes defined by a family of forbidden (but not necessarily induced) graphs with at most seven vertices and at most six edges [18].

We focus on the case where $\mathcal{F}_{\mathcal{G}}$ consists of a single graph H. A graph G is H-free if G does not contain an induced subgraph isomorphic to H. We obtain the following dichotomy for H-free graphs.

Theorem 5. Let $k \geq 3$ be an integer and H be a graph. If H is a linear forest, then k-EDGE COLOURING is polynomial-time solvable for H-free graphs. Otherwise k-EDGE COLOURING is NP-complete even for k-regular H-free graphs.

We obtain Theorem 5 by combining Theorems 3 and 4 with two new results. In particular, we will prove a hardness result for k-regular claw-free graphs for even integers k (as Theorem 4 is only valid when k is odd).

2 Preliminaries

The graphs C_n , P_n and K_n denote the path, cycle and complete graph on n vertices, respectively. A set I is an *independent set* of a graph G if all vertices of I are pairwise nonadjacent in G. A graph G is *bipartite* if its vertex set can be partitioned into two independent sets A and B. If there exists an edge between every vertex of A and every vertex of B, then G is *complete bipartite*. The *claw* $K_{1,3}$ is the complete bipartite graph with |A| = 1 and |B| = 3.

Let G_1 and G_2 be two vertex-disjoint graphs. The *join* operation \times adds an edge between every vertex of G_1 and every vertex of G_2 . The *disjoint union* operation + merges G_1 and G_2 into one graph without adding any new edges, that is, $G_1 + G_2 = (V(G_1) \cup V(G_2), E(G_1) \cup E(G_2))$. We write rG to denote the disjoint union of r copies of a graph G.

A forest is a graph with no cycles. A linear forest is a forest of maximum degree at most 2, or equivalently, a disjoint union of one or more paths. A graph G is a cograph if G can be generated from K_1 by a sequence of join and disjoint union operations. A graph is a cograph if and only if it is P_4 -free (see, for example, [3]). The following well-known lemma follows from this equivalence and the definition of a cograph.

Lemma 1. Every connected P_4 -free graph on at least two vertices has a spanning complete bipartite subgraph.

Let G = (V, E) be a graph. For a subset $S \subseteq V$, the graph $G[S] = (S, \{uv \in E \mid u, v \in S\})$ denotes the subgraph of G induced by S. We say that S is connected if G[S] is connected. Recall that a graph G is H-free for some graph H if G does not contain H as an induced subgraph. A subset $D \subset V(G)$ is dominating if every vertex of $V(G) \setminus D$ is adjacent to least one vertex of D. We will need the following result of Camby and Schaudt.

Theorem 6 ([5]). Let $t \ge 4$ and G be a connected P_t -free graph. Let X be any minimum connected dominating set of G. Then G[X] is either P_{t-2} -free or isomorphic to P_{t-2} .

Let G = (V, E) be some graph. The degree of a vertex $u \in V$ is equal to the size of its neighbourhood $N(u) = \{v \mid uv \in E\}$. The graph G is r-regular if every vertex of G has degree r. The line graph of G is the graph L(G), which has vertex set E and an edge between two distinct vertices e and f if and only if e and f have a common end-vertex in G.

3 The Proof of Theorem 5

To prove our dichtomy, we first consider the case where the forbidden induced subgraph H is a claw. As line graphs are claw-free, we only need to deal with the case where the number of colours k is even due to Theorem 4. For proving this case we need another result of Cai and Ellis, which we will use as a lemma. Let c be a k-edge colouring of a graph G = (V, E). Then a vertex $u \in V$ misses colour i if none of the edges incident to u is coloured i.

Lemma 2 ([4]). For even $k \ge 2$, the complete graph K_k has a k-edge colouring with the property that $V(K_k)$ can be partitioned into sets $\{u_i, u_i'\}$ $(1 \le i \le \frac{k}{2})$, such that for $i = 1, \ldots, \frac{k}{2}$, vertices u_i and u_i' miss the same colour, which is not missed by any of the other vertices.

We use Lemma 2 to prove the following result, which solves the case where k is even and $H = K_{1,3}$.

Lemma 3. Let $k \ge 4$ be an even integer. Then k-EDGE COLOURING is NP-complete for k-regular claw-free graphs.

Proof. Recall that k-EDGE COLOURING for k-regular graphs is NP-complete for every integer $k \geq 4$ due to Theorem 2. Consider an instance (G,k) of k-EDGE COLOURING, where G is k-regular for some even integer $k = 2\ell \geq 4$. From G we construct a graph G' as follows. First we replace every vertex v in G by the gadget H(v) shown in Figure 1. Next we connect the different gadgets in the following way. Every gadget H(v) has exactly k pendant edges, which are incident with vertices $v_1, \ldots, v_\ell, v_{\ell+1}, \ldots, v_{2\ell}$, respectively. As G is k-regular, every vertex has k neighbours in G. Hence, we can identify each edge uv of G with a unique edge uhv_i in G', which is a pendant edge of both H(u) and H(v). It is readily seen that G' is k-regular and claw-free.




Fig. 1. The gadget H(v) where $K_i(v)$ is a complete graph of size 2ℓ for i = 1, 2. Note that edges inside $K_1(v)$ and $K_2(v)$ are not drawn.

First suppose that G is k-edge colourable. Let c be a k-edge colouring of G. Consider a vertex $v \in V(G)$. For every neighbour u of v in G, we colour the pendant edge in H(v) corresponding to the edge uv with colour c(uv). As c assigned different colours to the edges incident to v, the 2ℓ pendant edges of H(v) will receive pairwise distinct colours, which we denote by $x_1, \ldots, x_\ell, y_1, \ldots, y_\ell$. By Lemma 2, we can colour the edges of $K_1(v)$ in such a way that for $i = 1, \ldots, \ell$, v_i and v_i' miss colour x_i . For $i = 1, \ldots, \ell$, we can therefore assign colour x_i to edge $v_i'w$. Similarly, we may assume that for $i = 1, \ldots, \ell$, $v_{\ell+i}$ and $v_{\ell+i}'$ miss colour y_i . For $i = 1, \ldots, \ell$, we can therefore assign colour y_i to edge $v_{\ell+i}'w$. Recall that the colours $x_1, \ldots, x_\ell, y_1, \ldots, y_\ell$ are all different. Hence, doing this procedure for each vertex of G yields a k-edge colouring c' of G'.

Now suppose that G' is k-edge colourable. Let c' be a k-edge colouring of G'. Consider some $v \in V(G)$. Denote the pendant edges of H(v) by e_i for $i = 1, \ldots, 2\ell$, where e_i is incident to v_i (and to some vertex u_h in a gadget H(u) for each neighbour u of v in G). Suppose that c' gave colour x to an edge wv_i' for some $1 \le i \le \ell$, say to wv_1' , but not to any edge e_i for $i = 1, \ldots, \ell$. Note that wv_2', \ldots, wv_ℓ' cannot be coloured x. As every vertex of G' has degree $k = 2\ell$, every v_i with $1 \le i \le \ell$ and every v_j' with $1 \le i \le \ell$ is incident to some edge coloured x. As x is neither the colour of e_1, \ldots, e_ℓ nor the colour of wv_2', \ldots, wv_ℓ' , the complete graph $K_1(v) - v_1'$ contains a perfect matching all of whose edges have colour x. However, $K_1(v) - v_1'$ has odd size $2\ell - 1$. Hence, this is not possible. We conclude that each of the (pairwise distinct) colours of wv_1', \ldots, wv_ℓ' , which we denote by x_1, \ldots, x_ℓ , is the colour of an edge e_i for some $1 \le i \le \ell$.

Let y_1, \ldots, y_ℓ be the (pairwise distinct) colours of $wv'_{\ell+1}, \ldots, wv'_{2\ell}$, respectively. By the same arguments as above, we find that each of those colours is also the colour of a pendant edge of H(v) that is incident to a vertex $v_{\ell+i}$ for some $1 \le i \le \ell$. Note that $x_1, \ldots, x_\ell, y_1, \ldots, y_\ell$ are 2ℓ pairwise distinct colours, as they are colours of edges incident to the same vertex, namely vertex w. Hence, we can define a k-colouring c of G by setting $c(uv) = c'(u_h v_i)$ for every edge $uv \in E(G)$ with corresponding edge $u_h v_i \in E(G')$.

We note that the graph G' in the proof of Lemma 3 is not a line graph, as the gadget H(v) is not a line graph: the vertices v'_1, v'_2, v_1, w form a diamond and by adding the pendant edge incident to v_1 and the edge $wv'_{\ell+1}$ we obtain an induced subgraph of H(v) that is not a line graph.

To handle the case where the forbidden induced subgraph H is a path, we make the following observation.

Observation 1 If a graph G of maximum degree k has a dominating set of size at most p, then G has at most p(k+1) vertices.

We use Observation 1 in the proof of the following lemma.

Lemma 4. Let $k \geq 3$ and $t \geq 1$. Then k-Edge Colouring is constant-time solvable for P_t -free graphs.

Proof. Let G be a P_t -free graph with maximum degree Δ . If $\Delta \leq k-1$, then G is k-edge colourable by Theorem 1. If $\Delta \geq k+1$, then G is not k-edge colourable. Hence, we may assume that $\Delta = k$. We may assume without loss of generality that G is connected. We claim that G has at most f(k,t) vertices for some function f that only depends on k and t. As we assume that k and t are constants, this means that we can check in constant time if G is k-edge colourable.

We prove the above claim by induction on t. First suppose t = 4 (and observe that if the claim holds for t = 4, it also holds for $t \le 3$). As G is connected, G has a dominating set of size 2 due to Lemma 1. Hence, by Observation 1, G has at most f(k,2) = 2(k+1) vertices. Now suppose $t \ge 5$. Let X be an arbitrary minimum connected dominating set of G. By Theorem 6, G[X] is either P_{t-2} -free or isomorphic to P_{t-2} . In the first case we use the induction hypothesis to conclude that G[X] has at most f(k, t-2) vertices. Hence, G has at most f(k, t-2)(k+1) vertices by Observation 1. In the second case, we find that G has at most (t-2)(k+1) vertices. We set $f(k,t) = \max\{f(k,t-2)(k+1), (t-2)(k+1)\}$.

We are now ready to prove Theorem 5, which we restate below.

Theorem 5. (restated) Let $k \geq 3$ be an integer and H be a graph. If H is a linear forest, then k-EDGE COLOURING is polynomial-time solvable for H-free graphs. Otherwise k-EDGE COLOURING is NP-complete even for k-regular H-free graphs.

Proof. First suppose that H contains a cycle C_s for some $s \geq 3$. Then the class of H-free graphs is a superclass of the class of C_s -free graphs. This means that we can apply Theorem 3. From now on assume that H contains no cycle, so H is a forest. Suppose that H contains a vertex of degree at least 3. Then the class of H-free graphs is a superclass of the class of $K_{1,3}$ -free graphs, which in turn forms a superclass of the class of line graphs. Hence, if k is odd, then we apply Theorem 4, and if k is even, then we apply Lemma 3. From now on assume that H contains no cycle and no vertex of degree at least 3. Then H is a linear forest, say with ℓ connected components. Let $t = \ell |V(H)|$. Then the class of H-free graphs is contained in the class of H-free graphs. Hence we may apply Lemma 4. This completes the proof of Theorem 5.

4 Conclusions

We gave a complete complexity classification of k-EDGE COLOURING for H-free graphs, showing a dichotomy between constant-time solvable cases and NP-complete cases. We saw that this depends on H being a linear forest or not. It would be interesting to prove a dichotomy result for EDGE COLOURING restricted to H-free graphs. Note that due to Theorem 5 we only need to consider the case where H is a linear forest. However, even determining the complexity for small linear forests H, such as the cases where $H = 2P_2$ and $H = P_4$,

turns out to be a difficult problem. In fact, the computational complexity of EDGE COLOURING for split graphs, or equivalently, $(2P_2, C_4, C_5)$ -free graphs [8] and for P_4 -free graphs has yet to be settled, despite the efforts towards solving the problem for these graph classes [6,7,15].

On a side note, a graph is k-edge colourable if and only if its line graph is k-vertex colourable. In contrast to the situation for EDGE COLOURING, the computational complexity of VERTEX COLOURING has been fully classified for H-free graphs [13]. However, the complexity status of k-VERTEX COLOURING is still open for P_t -free graphs if k = 3 and $t \ge 8$ after Bonomo et al. [2] proved polynomial-time solvability for k = 3 and t = 7; see the survey [9] for further background information.

Acknowledgements

The present work was done when the second author was visiting the University of Fribourg funded by a scholarship of the University of Fribourg.

References

- 1. H. L. Bodlaender. Polynomial algorithms for graph isomorphism and chromatic index on partial k-trees. *Journal of Algorithms*, 11(4):631–643, 1990.
- 2. F. Bonomo, M. Chudnovsky, P. Maceli, O. Schaudt, M. Stein, and M. Zhong. Three-coloring and list three-coloring of graphs without induced paths on seven vertices. *Combinatorica*, 38:779–801, 2018.
- 3. A. Brandstädt, V. B. Le, and J. P. Spinrad. *Graph Classes: A Survey*. SIAM Monographs on Discrete Mathematics and Applications. Society for Industrial and Applied Mathematics (SIAM), 1999.
- 4. L. Cai and J. A. Ellis. NP-completeness of edge-colouring some restricted graphs. *Discrete Applied Mathematics*, 30(1):15–27, 1991.
- 5. E. Camby and O. Schaudt. A new characterization of P_k -free graphs. Algorithmica, 75(1), 2016.
- B.-L. Chen, H.-L. Fu, and M. T. Ko. Total chromatic number and chromatic index of split graphs. Journal of Combinatorial Mathematics and Combinatorial Computing, 17:137–146, 1995.
- S. M. de Almeida, C. P. de Mello, and A. Morgana. Edge-coloring of split graphs. Ars Combinatoria, 119:363–375, 2015.
- 8. S. Földes and P. L. Hammer. Split graphs. Congressus Numerantium, XIX:311-315, 1977.
- 9. P. A. Golovach, M. Johnson, D. Paulusma, and J. Song. A survey on the computational complexity of colouring graphs with forbidden subgraphs. *Journal of Graph Theory*, 84(4):331–363, 2017.
- 10. I. Holyer. The NP-completeness of edge-coloring. SIAM Journal on Computing, 10(4):718–720, 1981.
- 11. D. S. Johnson. The NP-completeness column: An ongoing guide. J. Algorithms, 6(3):434-451, 1985.
- 12. S. Kitaev and V. V. Lozin. Words and Graphs. Monographs in Theoretical Computer Science. An EATCS Series. Springer, 2015.
- 13. D. Král', J. Kratochvíl, Zs. Tuza, and G. J. Woeginger. Complexity of coloring graphs without forbidden induced subgraphs. *Proc. WG 2001, LNCS*, 2204:254–262, 2001.
- 14. D. Leven and Z. Galil. NP completeness of finding the chromatic index of regular graphs. *Journal of Algorithms*, 4(1):35–44, 1983.
- 15. A. R. C. Lima, G. Garcia, L. Zatesko, and S. M. de Almeida. On the chromatic index of cographs and join graphs. *Eletronic Notes in Discrete Mathematics*, 50:433–438, 2015.
- R. C. S. Machado, C. M. H. de Figueiredo, and N. Trotignon. Edge-colouring and total-colouring chordless graphs. Discrete Mathematics, 313(14):1547–1552, 2013.
- 17. D. S. Malyshev. The complexity of the edge 3-colorability problem for graphs without two induced fragments each on at most six vertices. Sib. elektr. matem. izv., 11:811–822, 2014.
- 18. D. S. Malyshev. Complexity classification of the edge coloring problem for a family of graph classes. *Discrete Mathematics and Applications*, 27:97–101, 2017.
- C. Ortiz, N. Maculan, and J. L. Szwarcfiter. Characterizing and edge-colouring split-indifference graphs. Discrete Applied Mathematics, 82(1-3):209-217, 1998.
- 20. V. G. Vizing. On an estimate of the chromatic class of a p-graph. Diskret. Analiz., 3:25–30, 1964.