

POLYNOMIALS OVER STRICT SEMIRINGS DO NOT ADMIT UNIQUE FACTORIZATION

ALEXANDER AGUDELO

ABSTRACT. In this short note, we prove the claim of the title.

INTRODUCTION

It is well-known that the polynomial algebra over the tropical semifield does not satisfy unique factorization. For example,

$$(T + 1)(T^2 + 1) = T^3 + T^2 + T + 1 = (T + 1)^3$$

are two different factorizations of the same polynomial. Note that this failure of unique factorization should not be confused with the unique factorization property for *tropical (polynomial) functions*, which holds since polynomials that define the same function are identified; also cf. the comments at the end of section 2 in [3].

It is clear that the above example works for polynomial algebras over any idempotent semiring. What is not so clear, and to our knowledge not mentioned in the literature yet, is that unique factorization fails for polynomial algebras over any strict semiring.

Examples for which unique factorization fails are the polynomial algebra over the next semirings:

- Any idempotent semirings.
- The natural numbers \mathbb{N} .
- The non-negative real numbers $\mathbb{R}_{\geq 0}$.
- The ambient semiring $(\mathbb{T}^{\text{hyp}})^+ = \mathbb{N}[\mathbb{R}_{>0}]$ of the tropical hyperfield \mathbb{T}^{hyp} ; cf. [2, Ex. 5.7.4] for more details on \mathbb{T}^{hyp} .
- $S_0 = \mathbb{N}[x, y]/(xy \sim 1, x + y \sim 1)$. This semiring is interesting since there is a semiring morphism from S_0 to R if and only if there is an element $a \in R^\times$ such that $a + a^{-1} = 1$, and this last property leads to different cases in our proof.

THE THEOREM

A (*commutative*) *semiring* (with 0 and 1) is a set R with two constants 0 and 1 and two binary operations $+$ and \cdot such that both $(R, +, 0)$ and $(R, \cdot, 1)$ are commutative monoids and such that $0 \cdot a = 0$ and $a(b + c) = ab + ac$ for all $a, b, c \in R$. A semiring R is *strict* if $a + b = 0$ implies $a = b = 0$ for all $a, b \in R$. It is *without zero divisors* if $ab = 0$ implies $a = 0$ or $b = 0$ for all $a, b \in R$. An element in R is a *unit* if it has an inverse element in the multiplicative monoid of R . The set of all units of R will be denoted by R^\times . A non-zero non-unit element of R is *irreducible* if it is not a product of two non-units. The semiring R admits the *unique factorization property* if it is without zero divisor and every non-zero non-unit element has a unique factorization into irreducible elements, up to permutations and multiplication by units. The *polynomial algebra* over R is the semiring $R[T]$ of formal sums $\sum_{i \in \mathbb{N}} a_i T^i$, where there is n such that $a_n = 0$ for all $i > n$, together with the usual addition and multiplication of polynomials.

Our main theorem is the following.

Theorem 1. *Polynomial algebras over strict semirings without zero divisors do not satisfy the unique factorization property.*

THE PROOF

Let R be a strict semiring without zero divisors, we know that $R[T]$ is without zero divisor and the units of $R[T]$ are the units of R (cf. [1, Ex. 4.23]). This means, as usual, two irreducible polynomials of different degree can not differ by a unit. Thus if we find one polynomial that admits two different factorization into irreducible polynomials of different degree we are done.

Note that the polynomial $T^5 + T^4 + T^3 + T^2 + T + 1$ admits the following two factorization

$$(T + 1)(T^4 + T^2 + 1) = (T^3 + 1)(T^2 + T + 1).$$

The next lemmas tell us when the polynomial above are irreducible.

Lemma 2. *For all $a \in R^\times$ and $n \geq 1$ the polynomial $T^n + a$ is irreducible.*

Proof. It is obvious that $T + a$ is irreducible. For $n \geq 2$ assume that $T^n + a$ is reducible, then there are $1 \leq m < n$ and $c_0, \dots, c_{m-1}, d_0, \dots, d_{n-m-1} \in R$ such that

$$T^n + a = (T^m + c_{m-1}T^{m-1} + \dots + c_0)(T^{n-m} + d_{n-m-1}T^{n-m-1} + \dots + d_0).$$

Therefore $c_0d_0 = a$ and $\sum_{i=1}^k c_i d_{k-i} = 0$ for all $1 \leq k \leq n-1$. In particular, d_0 is a unit. For $k = m$, we get $d_0 + \sum_{i=1}^{m-1} c_i d_{k-i} = 0$. Since R is strict, $d_0 = 0$, which is a contradiction. \square

Lemma 3. *Either $T^2 + T + 1$ is irreducible or there is $a \in R^\times$ such that $a + a^{-1} = 1$. In the latter case, $T^2 + T + 1 = (T + a)(T + a^{-1})$.*

Proof. Suppose that $T^2 + T + 1$ is reducible. Then there are $c, d \in R$ such that

$$T^2 + T + 1 = (T + c)(T + d).$$

This implies $cd = 1$ and $c + d = 1$. \square

Lemma 4. *Either $T^4 + T^2 + 1$ is irreducible or there is $a \in R^\times$ such that $a + a^{-1} = 1$. In the latter case, $T^4 + T^2 + 1 = (T^2 + a)(T^2 + a^{-1})$.*

Proof. Suppose that $T^4 + T^2 + 1$ is reducible. Now we have two possibilities, $T^4 + T^2 + 1$ factors as the product of one linear and one cubic polynomial or as the product of two quadratics polynomials.

In the first case, there are $c_0, d_0, d_1, d_2 \in R$ such that

$$T^4 + T^2 + 1 = (T + c_0)(T^3 + d_2T^2 + d_1T + d_0).$$

Therefore $c_0d_0 = 1$ and $c_0 + d_2 = 0$, which is impossible since R is strict.

In the second case, there are $c_0, c_1, d_0, d_1 \in R$ such that

$$T^4 + T^2 + 1 = (T^2 + c_1T + c_0)(T^2 + d_1T + d_0).$$

Therefore $c_0d_0 = 1$, $c_0d_1 + c_1d_0 = 0$, $c_1d_1 + c_0 + d_0 = 1$ and $c_1 + d_1 = 0$. Since R is strict, we get $c_1 = d_1 = 0$, $c_0d_0 = 1$ and $c_0 + d_0 = 1$. \square

We are ready to finish the proof of Theorem 1.

If there is no $a \in R^\times$ such that $a + a^{-1} = 1$, then Lemmas 2, 3 and 4 show that all factors in

$$(T + 1)(T^4 + T^2 + 1) = (T^3 + 1)(T^2 + T + 1)$$

are irreducible. Otherwise, we have

$$(T + 1)(T^2 + a)(T^2 + a^{-1}) = (T^3 + 1)(T + a)(T + a^{-1}).$$

Lemma 2 implies that all factors in the above expression are irreducible. In both cases we have found two different factorization of $T^5 + T^4 + T^3 + T^2 + T + 1$ into irreducible polynomials of different degree. \square

REFERENCES

- [1] Jonathan S. Golan. *Semirings and their applications*. Kluwer Academic Publishers, Dordrecht, 1999. Updated and expanded version of it The theory of semirings, with applications to mathematics and theoretical computer science [Longman Sci. Tech., Harlow, 1992; MR1163371 (93b:16085)].
- [2] Oliver Lorscheid. Blueprints and tropical scheme theory. Lecture notes, <http://oliver.impa.br/2018-Blueprints/versions/lecturenotes180521.pdf>, version from May 21, 2018.
- [3] David Speyer and Bernd Sturmfels. Tropical mathematics. *Math. Mag.*, 82(3):163–173, 2009.
E-mail address: alagudel@impa.br

INSTITUTO NACIONAL DE MATEMÁTICA PURA E APLICADA, RIO DE JANEIRO, BRAZIL