ZERO ENTROPY AUTOMORPHISMS OF COMPACT KÄHLER MANIFOLDS AND DYNAMICAL FILTRATIONS

TIEN-CUONG DINH, HSUEH-YUNG LIN, KEIJI OGUISO, AND DE-QI ZHANG

In memory of Jean-Pierre Demailly

ABSTRACT. We study zero entropy automorphisms of a compact Kähler manifold X. Our goal is to bring to light some new structures of the action on the cohomology of X, in terms of the so-called dynamical filtrations on $H^{1,1}(X,\mathbb{R})$. Based on these filtrations, we obtain the first general upper bound on the polynomial growth of the iterations $(g^m)^* \circ H^2(X,\mathbb{C})$ where g is a zero entropy automorphism, in terms of dim X only.

We also give an upper bound for the (essential) derived length $\ell_{ess}(G, X)$ for every zero entropy subgroup G, again in terms of the dimension of X only. We propose a conjectural upper bound for the essential nilpotency class $c_{ess}(G, X)$ of a zero entropy subgroup G.

Finally, we construct examples showing that our upper bound of the polynomial growth (as well as the conjectural upper bound of $c_{ess}(G, X)$) are optimal.

CONTENTS

| 1. | Introduction | 1 |
|------------|--|----|
| 2. | Essential derived lengths and nilpotency classes | Ę |
| 3. | Dynamical filtrations and consequences | 8 |
| 4. | Explicit examples | 18 |
| References | | 21 |

1

1. INTRODUCTION

Let X be a compact Kähler manifold and let $\operatorname{Aut}(X)$ be the group of holomorphic automorphisms of X. Recall that for every $g \in \operatorname{Aut}(X)$, according to theorems of Gromov and Yomdin, the topological entropy of g is equal to the logarithm of the spectral radius of its action on the cohomology ring of X, see [6, 13, 28]. Groups with positive entropy elements have been intensively studied during the last decades using techniques and ideas from complex dynamics and algebraic geometry, see *e.g.* [3, 7, 21] and the references therein. Despite these developments, much less is known about zero entropy elements (see *e.g.* [3, §3]), which will be the main focus of this paper.

One of the aims of this paper is to excavate some hidden structures of zero entropy group actions on the cohomology of X, in terms of various dynamical filtrations on $H^{1,1}(X,\mathbb{R})$.

Introduction

²⁰¹⁰ Mathematics Subject Classification. 14J50, 32M05, 32H50, 37B40.

Key words and phrases. Automorphisms of compact Kähler manifolds, zero entropy automorphisms, dynamical filtrations, polynomial growth, derived length, nilpotency class.

We defer the discussion on these dynamical filtrations to the later part of the introduction, and start with the consequences.

1.1. Polynomial growths of pullback actions.

As a consequence of Gromov–Yomdin's theorem, an automorphism $g \in \operatorname{Aut}(X)$ of a compact Kähler manifold X has zero entropy if and only if the pullback $g^* : H^{\bullet}(X, \mathbb{C}) \circlearrowleft$ is quasi-unipotent (see e.g. Proposition 2.6(1)). In this case, the iterations $(g^m)^* :$ $H^{\bullet}(X, \mathbb{C}) \circlearrowright \text{ of } g^*$ have polynomial growth when m tends to infinity. The study of the polynomial growth of $(g^m)^* : H^2(X, \mathbb{C}) \circlearrowright$ was initiated by Lo Bianco [18]: he proved that the growth satisfies

$$\|(g^m)^*: H^2(X, \mathbb{C}) \circlearrowleft \| =_{m \to \infty} O(m^4).$$

when dim X = 3. With the help of dynamical filtrations, we generalize Lo Bianco's theorem to compact Kähler manifolds of arbitrary dimension.

Theorem 1.1. Let X be a compact Kähler manifold of dimension $n \ge 1$. Let g be any automorphism of zero entropy of X. Then the action of g^m on $H^{1,1}(X, \mathbb{C})$ satisfies

$$||(g^m)^* : H^{1,1}(X, \mathbb{C}) \circlearrowleft || = O(m^{2(n-1)})$$

when m tends to infinity. This estimation is optimal in terms of $n = \dim X$.

More generally, the action of g^m on each factor $H^{p,q}(X,\mathbb{C})$ of the Hodge decomposition with $0 \leq p, q \leq n$, satisfies

$$||(g^m)^* : H^{p,q}(X, \mathbb{C}) \circlearrowleft || = O(m^{(p'+q')(n-1)})$$

when m tends to infinity; here $p' := \min(p, n-p)$ and $q' := \min(q, n-q)$.

According to [3, §3.2], Theorem 1.1 appears to be the first general result on the growth of actions on $H^2(X, \mathbb{C})$ induced by zero entropy automorphisms. Note that

$$||(g^m)^* : H^{p,q}(X,\mathbb{R}) \circlearrowleft || = O(m^{k-1})$$

if and only if the Jordan blocks of $g^* : H^{p,q}(X, \mathbb{R}) \bigcirc$ in its Jordan form are of size $k \times k$ or smaller.

1.2. Derived lengths of zero entropy subgroups.

Given a group H, its p-th derived subgroups $H^{(p)}$ are defined inductively by

$$H^{(0)} := H$$
 and $H^{(i+1)} := [H^{(i)}, H^{(i)}],$

where $[H^{(i)}, H^{(i)}]$ is the subgroup of H generated by $[g, h] = ghg^{-1}h^{-1}$ $(g, h \in H^{(i)})$. By definition, $H^{(p)} = \{1\}$ for some integer $p \ge 0$ if and only if H is solvable. We call the minimum of such p the *derived length* of H (when H is solvable) and denote it by $\ell(H)$. If H is not solvable, we set $\ell(H) := \infty$.

A subgroup $H \subset \operatorname{GL}(N, \mathbb{R})$ is said to be *unipotent* if 1 is the only eigenvalue of every element of H. It is known that a subgroup H of $\operatorname{GL}(N, \mathbb{R})$ is unipotent if and only if His conjugate to a subgroup consisting of upper triangular matrices with all entries on the diagonal being 1 [15, §17.5]. Note that in this discussion, we can replace \mathbb{R} by \mathbb{C} as we have the natural inclusions $\operatorname{GL}(N, \mathbb{R}) \subseteq \operatorname{GL}(N, \mathbb{C}) \subseteq \operatorname{GL}(2N, \mathbb{R})$.

Since unipotent subgroups are nilpotent and hence solvable, every unipotent subgroup H has derived length $\ell(H) < \infty$.

We say that a subgroup $G \subset \operatorname{Aut}(X)$ has zero entropy, and call it a zero entropy subgroup, if all its elements have zero entropy. For every subgroup $G \subset \operatorname{Aut}(X)$, we define

$$G_0 := G \cap \operatorname{Aut}^0(X),$$

where $\operatorname{Aut}^{0}(X)$ is the identity component of $\operatorname{Aut}(X)$. Note that $\operatorname{Aut}^{0}(X)$ is a connected complex Lie group of finite dimension by Bochner–Montgomery's theorem [1]. For every zero entropy subgroup $G \subset \operatorname{Aut}(X)$ one can show that there exists a finite index subgroup $G' \subset G$ such that the action of G'/G'_{0} on $H^{p,p}(X,\mathbb{R})$ realizes G'/G'_{0} as a unipotent subgroup of $\operatorname{GL}(H^{p,p}(X,\mathbb{R}))$ for every $1 \leq p \leq \dim X - 1$. Moreover, the invariant

$$\ell_{\rm ess}(G, X) := \ell(G'/G'_0),$$

does not depend on the choice of G' (see Proposition 2.6). We call $\ell_{\text{ess}}(G, X)$ the essential derived length of the subgroup $G \subset \text{Aut}(X)$.

We will prove the following in §3, based on dynamical filtrations.

Theorem 1.2. Let X be a compact Kähler manifold of dimension $n \ge 1$. Then every subgroup $G \subset Aut(X)$ of zero entropy satisfies

$$\ell_{\rm ess}(G, X) \le n - 1.$$

1.3. Dynamical filtrations.

In this subsection, we present one example of dynamical filtration, the one which underlies the estimate in Theorem 1.1.

Let X be a compact Kähler manifold of dimension $n \ge 1$. For simplicity, if L and M are two cohomology classes, we denote by $L \cdot M$ or LM their cup-product. We also identify $H^0(X, \mathbb{R})$ and $H^{2n}(X, \mathbb{R})$ with \mathbb{R} in the canonical way, where $n = \dim X$. So classes in these groups are identified with real numbers. The *nef cone* Nef $(X) \subset H^{1,1}(X, \mathbb{R})$ is defined as the closure of the Kähler cone.

Let $G \subset \operatorname{Aut}(X)$ be a subgroup such that the induced action $G \circlearrowleft H^{1,1}(X)$ is unipotent. For instance, G can be the cyclic group generated by some finite power of a zero entropy automorphism. Based on a Lie–Kolchin type theorem [16, Theorem 1.1], there exists a sequence

 $M_1, \ldots, M_n \in H^{1,1}(X, \mathbb{R})$

such that each $L_i := M_1 \cdots M_i \neq 0 \in H^{i,i}(X, \mathbb{R})$ (with $L_0 := 1 \in H^0(X, \mathbb{R})$) is G-invariant and satisfies

$$L_i \in \overline{L_{i-1} \cdot \operatorname{Nef}(X)}.$$

Following [29] and [6], we call $M_1, \ldots, M_n \in H^{1,1}(X, \mathbb{R})$ a quasi-nef sequence for $G \oslash X$. For $\beta \in H^{i,i}(X, \mathbb{R})$, we write $\beta \equiv 0$ if $\beta \cdot H^{1,1}(X, \mathbb{R})^{n-i} = 0$. Define

$$F_i := \left\{ \alpha \in H^{1,1}(X, \mathbb{R}) \mid L_i \alpha \equiv 0 \right\}$$

and

$$F'_i := \left\{ \alpha \in F_i \mid L_{i-1} \alpha^2 \equiv 0 \right\}.$$

Based on the mixed Hodge–Riemann relations [9, 12] and the positivity of nef classes, we will get:

Theorem 1.3. The subsets F_i and F'_i form a filtration

$$0 = F_0 \subset F'_1 \subset F_1 \subset \cdots \subset F'_{n-1} \subset F_{n-1} \subset F'_n = H^{1,1}(X,\mathbb{R})$$

of G-stable vector spaces, where $n = \dim X$. Moreover,

- (1) dim $F'_i/F_{i-1} \leq 1$; and the following assertions are equivalent:
 - (a) dim $F'_i/F_{i-1} = 1$;
 - (b) $F'_i = F_{i-1} \oplus \mathbb{R} \cdot M_i;$
 - (c) $L_{i-1}M_i^2 = 0.$
- (2) For every $g \in G$, there exists a unique strictly decreasing sequence of integers

$$n-1 \ge s_1 > \dots > s_r \ge 1$$

such that for every Kähler class $\omega \in H^{1,1}(X, \mathbb{R})$, we have

$$(g^* - \operatorname{Id})^{2j-1}(\omega) \in F_{s_j} \setminus F'_{s_j} \quad and \quad (g^* - \operatorname{Id})^{2j}(\omega) \in F'_{s_j} \setminus F_{s_j-1}$$

for $1 \le j \le r$, and $(g^* - \operatorname{Id})^{2r+1}(\omega) = 0$.

The upper bound estimate in Theorem 1.1 for $(g^m)^* : H^{1,1}(X, \mathbb{R}) \circlearrowleft$ (which is the essential part of the statement) follows from Theorem 1.3. The proof of Theorem 1.2 is based on a more refined dynamical filtration; the reader is referred to §3 for details.

See [17] for other applications of dynamical filtrations.

1.4. Nilpotency classes: a conjecture and some consequences.

The descending central series of a group H is defined by

$$\Gamma_0 H := H$$
 and $\Gamma_{i+1} H := [\Gamma_i H, H] = [H, \Gamma_i H].$

By definition, $\Gamma_p H = \{1\}$ for some integer $p \ge 0$ if and only if H is nilpotent. We call the minimum of such p the *nilpotency class* of H (when H is nilpotent) and denote it by c(H). If H is not nilpotent, we set $c(H) := \infty$.

Let $G \subset \operatorname{Aut}(X)$ be a zero entropy subgroup. There exists a finite index subgroup $G' \subset G$ such that the action of G'/G'_0 on $H^{p,p}(X,\mathbb{R})$ realizes G'/G'_0 as a unipotent subgroup of $\operatorname{GL}(H^{p,p}(X,\mathbb{R}))$ for every $1 \leq p \leq \dim X - 1$. Similar to the definition of essential derived length, we define the essential nilpotency class

$$c_{\rm ess}(G,X) := c(G'/G'_0)$$

of $G \subset \operatorname{Aut}(X)$ to be the nilpotency class of G'/G'_0 ; according to Proposition 2.6, it does not depend on the choice of G'.

For any group H, the inclusion $H^{(i)} \subset \Gamma_i H$ implies $\ell(H) \leq c(H)$. We even have $H^{(i)} = 0$ for the *i*-th derived group, provided $2^i > c(H)$ (see *e.g.* [24, §5.1.12, Proof]), which implies

(1.1)
$$\ell(H) \le \lfloor \log_2(c(H)) \rfloor + 1$$

if H is non-trivial.

We believe that the upper bound in Theorem 1.2 is not optimal and propose the following more precise conjecture involving the Kodaira dimension $\kappa(X)$. In view of (1.1), this conjecture would improve Theorem 1.2 in a significant way: the right hand side of the inequality there would be replaced by $\lfloor \log_2(n-1) \rfloor + 1$.

Conjecture 1.4. Let X be a compact Kähler manifold of dimension $n \ge 1$. For every subgroup $G \subset \operatorname{Aut}(X)$ of zero entropy, we have

$$c_{\rm ess}(G, X) \le n - \max\{\kappa(X), 1\}.$$

We will construct some concrete examples in §4 showing the optimality of the upper bounds in Theorem 1.1 and Conjecture 1.4.

Remark 1.5. In [11], Epstein and Thurston studied upper bounds for the derived length of a solvable or nilpotent *connected* Lie group acting continuously on a Hausdorff space. Their examples in [11, Remark 1.5] may justify the necessity to quotient-away the continuous part G_0 of G in Conjecture 1.4. See also [4, §6.2] for relevant results.

Terminology and Notation.

In this paper, we work in the category of analytic spaces. All manifolds are assumed to be connected. By Zariski closures, we also mean *analytic* Zariski closures unless otherwise specified. Given two sequences (x_m) and (y_m) in $\mathbb{R}_{\geq 0}$ or more generally in a salient convex closed cone, the relation $x_m \leq y_m$ means $Cy_m - x_m$ belongs to this cone for some constant C > 0 independent of m, and $x_m \simeq y_m$ means $x_m \leq y_m$ and $y_m \leq x_m$.

When a group N acts on a space V preserving some structure on V, we denote by $N|_V$ the image of N in the group of automorphisms of V. For instance, if G is a group acting on a complex manifold X, then $G|_X$ is the image of G in Aut(X), and $G|_{H^p(X,\mathbb{Z})}$ is the image of G in $GL(H^p(X,\mathbb{Z}))$ for the pullback action. For a normal subgroup $N_1 \leq N$, we set $(N/N_1)|_V = (N|_V)/(N_1|_V)$.

We say that a property holds for *very general* (resp. *general*) parameters or points if it holds for all parameters or points outside a countable (resp. finite) union of proper closed analytic subvarieties of the space of parameters or points.

Acknowledgments.

The authors are supported by NUS and MOE grants R-146-000-248-114 and MOE-T2EP20120-0010; Taiwan Ministry of Education Yushan Young Scholar Fellowship (NTU-110VV006), and Taiwan Ministry of Science and Technology (110-2628-M-002-006-); a JSPS Grant-in-Aid (A) 20H00111 and an NCTS Scholar Program; and an ARF of NUS, respectively. We would like to thank Fei Hu and Jun-Muk Hwang for helpful discussions; Serge Cantat for various comments and for bringing the references [4, 11] to our attention; the referees for very detailed and constructive suggestions; and KIAS, National Center for Theoretical Sciences (NCTS) in Taipei, NUS and the University of Tokyo for the hospitality and support during the preparation of this paper.

2. Essential derived lengths and nilpotency classes

2.1. Unipotent subgroups.

Lemma 2.1 below should be well known, but we give a proof due to the lack of reference.

Lemma 2.1. Let V be a real vector space of finite dimension. Let Γ be a subgroup of $\operatorname{GL}(V)$. Assume that there is an integer $N \geq 1$ such that g^N is unipotent for every $g \in \Gamma$. Then there is a finite-index subgroup Γ' of Γ which is a unipotent subgroup of $\operatorname{GL}(V)$.

Proof. Let Γ denote the algebraic Zariski closure of Γ in $\operatorname{GL}(V_{\mathbb{C}})$, where $V_{\mathbb{C}} := V \otimes_{\mathbb{R}} \mathbb{C}$. This is a complex linear algebraic group. By hypothesis, for $g \in \Gamma$, all eigenvalues of g are N-th roots of unity. It follows that the coefficients of the characteristic polynomial of g belong to a finite set. These coefficients can be seen as polynomial functions in the matrix coefficients of $g \in \operatorname{GL}(V_{\mathbb{C}})$. We deduce that the same property holds for all $g \in \overline{\Gamma}$. Let H be the component of the identity of $\overline{\Gamma}$. By continuity, the characteristic polynomial of $g \in H$ is constantly equal to $(t-1)^{\dim V}$, as this is the case for the identity element. Thus 1 is the only eigenvalue for all $g \in H$.

Define $\Gamma' := \Gamma \cap H$; Γ' is a unipotent subgroup of GL(V) by [15, §17.5]. Since $\overline{\Gamma}$ is algebraic, it has a finite number of connected components. It follows that H is a finite-index subgroup of $\overline{\Gamma}$ and Γ' is a finite-index subgroup of Γ .

Theorem 2.2. Let $V_{\mathbb{Z}}$ be a free \mathbb{Z} -module of finite rank and let $\rho : \Gamma \to \operatorname{GL}(V_{\mathbb{Z}})$ be a group homomorphism such that $\operatorname{Ker}(\rho)$ is finite and all elements of $\rho(\Gamma)$ are unipotent. Then

- (1) (Mal'cev's theorem) Γ is countable, finitely generated, and admits only countably many subgroups. Moreover, all subgroups of Γ are finitely generated as well.
- (2) There is a finite-index normal subgroup Γ' of Γ such that $\Gamma' \cap \text{Ker}(\rho) = \{1\}$.

Proof. (1) By the assumption, $\rho(\Gamma)$ is a unipotent, hence solvable, subgroup of $\operatorname{GL}(V_{\mathbb{Z}})$ by [15, §17.5]. Thus the result follows from a theorem of Mal'cev [25, p.26, Corollary 1].

(2) By Zorn's lemma, there exists a maximal normal subgroup $\Gamma' \subset \Gamma$ such that $\Gamma' \cap \operatorname{Ker}(\rho) = \{1\}$. It is enough to show that Γ' is of finite-index in Γ . Assume by contradiction that the group $K := \Gamma/\Gamma'$ is infinite. Denote by $\pi : \Gamma \to K$ the canonical homomorphism. If L is a normal subgroup of K, then $\pi^{-1}(L)$ is a normal subgroup of Γ containing Γ' . To complete the proof of the lemma, i.e. to get a contradiction, it is enough to construct such a group L with $L \neq \{1\}$ and $L \cap \pi(\operatorname{Ker}(\rho)) = \{1\}$.

Since $\rho(\Gamma)$ is a unipotent subgroup of $\operatorname{GL}(V_{\mathbb{Z}})$, we have a finite sequence

$$\Gamma = \Gamma_0 \vartriangleright \Gamma_1 \vartriangleright \cdots \vartriangleright \Gamma_{m+1} \subset \ker(\rho) \quad \text{with} \quad \Gamma_{j+1} := [\Gamma, \Gamma_j]$$

As remarked above, all such subgroups Γ_j of Γ are finitely generated. Since K is infinite and $\ker(\rho)$ is finite, there is a j such that $\pi(\Gamma_j)$ is infinite but $\pi(\Gamma_{j+1})$ is finite. So $\pi(\Gamma_j)/\pi(\Gamma_{j+1})$ is an infinite abelian group which is finitely generated. Thus, there is an element $a \in \pi(\Gamma_j)$ which is of infinite order. Clearly, a^m does not belong to $\pi(\operatorname{Ker}(\rho))$ for $m \neq 0$ because $\ker(\rho)$ is finite. We need the following:

Claim 2.3. For any $k \in K$, there is an integer l > 0 such that $ka^{ml}k^{-1} = a^{ml}$ for every m.

We prove the claim. By construction, $\pi(\Gamma_j)/\pi(\Gamma_{j+1})$ is contained in the center of $K/\pi(\Gamma_{j+1})$. Therefore, for every $p \geq 1$, there is some $b_p = b_p(k) \in \pi(\Gamma_{j+1})$ such that $ka^pk^{-1} = b_pa^p$. Since $\pi(\Gamma_{j+1})$ is finite, there are p < q such that $b_p = b_q$. We then have for l := q - p

$$ka^{l}k^{-1} = (ka^{p}k^{-1})^{-1}(ka^{q}k^{-1}) = a^{l}$$

This implies the claim for m = 1 and then for every m.

We resume the proof of Theorem 2.2. Since K is finitely generated, the claim provides an $l \ge 1$ such that $ka^{l}k^{-1} = a^{l}$ for every $k \in K$. It follows that the group generated by a^{l} is normal in K. Since it has a trivial intersection with $\pi(\text{Ker}(\rho))$, we get a contradiction. This ends the proof of Theorem 2.2.

2.2. Essential derived lengths and nilpotency classes.

Lemma 2.4 is well known; we give a proof for completeness.

Lemma 2.4. Let H be a unipotent subgroup of $GL(N, \mathbb{R})$ (or $GL(N, \mathbb{C})$), and let H' be a finite-index subgroup of H. Then we have $\ell(H') = \ell(H)$ and c(H') = c(H).

Proof. By [15, §17.5], H can be regarded as a subgroup of the group of unipotent upper triangular matrices in $\operatorname{GL}(N, \mathbb{C})$ for a suitable N. Let $\overline{H} \subseteq \operatorname{GL}(N, \mathbb{C})$ be the algebraic Zariski closure of H. Then \overline{H} is still a unipotent subgroup of $\operatorname{GL}(N, \mathbb{C})$ and hence connected [15, Exercise 15.5.6].

Let $H \supseteq H^{(1)} \supseteq H^{(2)} \supseteq \cdots$ be the derived series of H. Then, $\overline{H}^{(i)}$ is equal to $\overline{H^{(i)}}$, the algebraic Zariski closure of $H^{(i)}$, see *e.g.* [20, Proof of Lemma 2.1(2)]. It follows that the derived length of H is equal to that of \overline{H} .

Since H' is of finite-index in H, so is the group $\overline{H'}$ in the group \overline{H} . Finally, since the latter two groups are both unipotent subgroups of $\operatorname{GL}(N, \mathbb{C})$, they are connected, and hence equal. Thus, we have $\ell(H) = \ell(\overline{H}) = \ell(\overline{H'}) = \ell(H')$. By replacing the derived series by the central series, the same argument shows c(H') = c(H).

Lemma 2.5. Let X be a compact Kähler manifold of dimension n. Let $G \subset Aut(X)$ be a subgroup of zero entropy. Then there is a finite-index subgroup G' of G satisfying the following properties for every $1 \le p \le n-1$.

(1) The kernels of the canonical representations

$$\rho_p: G' \to \operatorname{GL}(H^{2p}(X,\mathbb{R})) \quad and \quad \rho_{p,p}: G' \to \operatorname{GL}(H^{p,p}(X,\mathbb{R}))$$

are both equal to G'_0 .

(2) The images of both ρ_p and $\rho_{p,p}$ are unipotent subgroups of $GL(H^{2p}(X,\mathbb{R}))$ and $GL(H^{p,p}(X,\mathbb{R}))$, respectively.

Proof. It is enough to prove the statement for a fixed p because we can then deduce the lemma using a simple induction on p.

The argument of this paragraph is well-known. Let g be an automorphism of X of zero entropy. For every $m \in \mathbb{Z}$, g^m also has zero entropy. Therefore, by Gromov–Yomdin's theorem, all eigenvalues of $\rho_p(g^m)$ are of modulus less than or equal to 1. Since $\rho_p(g^m)$ preserves the image of $H^{2p}(X,\mathbb{Z})$ in $H^{2p}(X,\mathbb{R})$, we deduce that the characteristic polynomial of $\rho_p(g^m)$ belongs to a finite family of polynomials, as their coefficients are bounded integers. If λ is an eigenvalue of $\rho_p(g)$ then λ^m is an eigenvalue of $\rho_p(g^m)$ and hence belongs to a finite set which is independent of m. Thus, there is an integer $N \geq 1$ such that $\lambda^N = 1$ for all eigenvalues of $\rho_p(g)$.

According to Lemma 2.1, replacing G by a finite-index subgroup, we have that $\rho_p(G)$ contains only unipotent elements of $GL(H^{2p}(X,\mathbb{R}))$. So we have the property (2) in the lemma for both ρ_p and $\rho_{p,p}$.

The kernel K of $\rho_{p,p} : G \to \operatorname{GL}(H^{p,p}(X,\mathbb{R}))$ is the set of $g \in G$ whose action on $H^{p,p}(X,\mathbb{R})$ is trivial. In particular, K preserves the p-th power of a Kähler class. By a

generalized version of Fujiki–Lieberman's theorem [8, Theorem 2.1], the quotient K/G_0 is a finite group (for p = 1, we can refer to the original theorem of Fujiki and Lieberman). It follows that $\rho_p(K) \subset \operatorname{GL}(H^{2p}(X,\mathbb{R}))$ is finite and unipotent, thus trivial. This implies $K = \ker(\rho_p)$.

Finally, we apply Theorem 2.2 to $\Gamma := G/G_0$ and $V_{\mathbb{Z}} = H^{2p}(X, \mathbb{Z})/\text{torsion}$. According to this theorem, we can replace G by a finite-index subgroup G' and assume that $G'_0 = \ker(\rho_{p|G'}) = \ker(\rho_{p,p|G'})$. This implies the property (1) for both $\rho_{p,p}$ and ρ_p .

As immediate consequences of Lemmas 2.4 and 2.5, we now have the following.

Proposition 2.6. Let X be a compact Kähler manifold of dimension $n \ge 1$ and let $G \subset Aut(X)$ be a zero entropy subgroup. We have the following assertions.

- (1) G admits a finite-index subgroup G' such that for any $1 \le p \le n-1$, the natural map $G'/G'_0 \to G'|_{H^{2p}(X,\mathbb{R})}$ (resp. $G'/G'_0 \to G'|_{H^{p,p}(X,\mathbb{R})}$) embeds G'/G'_0 as a unipotent subgroup of $\operatorname{GL}(H^{2p}(X,\mathbb{R}))$ (resp. $\operatorname{GL}(H^{p,p}(X,\mathbb{R}))$).
- (2) For every finite-index subgroup G'' of G such that $G''/G''_0 \to \operatorname{GL}(H^{p,p}(X,\mathbb{R}))$ is an isomorphism onto a unipotent subgroup for some $1 \le p \le n-1$, the derived length $\ell(G''/G''_0)$ and the nilpotency class $c(G''/G''_0)$ of G''/G''_0 do not depend on the choice of G'', nor on p.

Proof. (1) is contained in Lemma 2.5. (2) follows from Lemma 2.4, as it implies $\ell(G'/G'_0) = \ell(G''/G''_0)$ and $c(G'/G'_0) = c(G''/G''_0)$.

3. Dynamical filtrations and consequences

3.1. Quasi-nef sequences.

Let $\mathcal{K}^i(X)$ denote the closure of the convex cone generated by the classes of smooth strictly positive closed (i, i)-forms in $H^{i,i}(X, \mathbb{R})$. This is a salient (that is, no line contained) convex closed cone with non-empty interior and is $\operatorname{Aut}(X)$ -invariant. In particular, $\mathcal{K}^1(X) = \operatorname{Nef}(X)$ is the nef cone.

For any class $L \in \mathcal{K}^i(X) \setminus \{0\}$ with $0 \le i \le n-1$, denote by Nef(L) the closure of the cone $L \cdot \operatorname{Nef}(X)$ in $H^{i+1,i+1}(X,\mathbb{R})$.

Remark 3.1. The image of a salient closed convex cone under a linear map is not necessarily closed, see e.g. [19, Remark 2.5]. Here is another example suggested by the referee. The image of

$$C = \left\{ \left. (x,y,z) \in \mathbb{R}^3 \right. \middle| \left. (y-z)^2 + x^2 \le z^2, y \ge 0, z \ge 0 \right. \right\} \subset \mathbb{R}^3$$

under the projection $\mathbb{R}^3 \to \mathbb{R}^2$ to the *xy*-plane is the open upper half-plane union $\{0\}$. For later use, observe that $C + \{x = y = 0\}$ is not closed in \mathbb{R}^3 .

Define also for $0 \le i \le n$

$$\mathcal{N}^{i}(X) := \left\{ \beta \in H^{i,i}(X, \mathbb{R}) \mid \beta \equiv 0 \right\},\$$

where we recall that $\beta \equiv 0$ means $\beta \cdot H^{1,1}(X,\mathbb{R})^{n-i} = 0$. This is a linear subspace of $H^{i,i}(X,\mathbb{R})$ preserved by Aut(X). Clearly $\mathcal{N}^n(X) = 0$ and $\mathcal{N}^0(X) = 0$. We also have $\mathcal{N}^i(X) \cdot M \subset \mathcal{N}^{i+1}(X)$ for all $M \in H^{1,1}(X,\mathbb{R})$. We have $\mathcal{N}^{n-1}(X) = 0$ by Poincaré duality, and hence $\mathcal{N}^1(X) = 0$ by hard Lefschetz's theorem (see e.g. [27, Theorem 6.25]).

Lemma 3.2. Let $L \in \mathcal{K}^i(X) \setminus \{0\}$ with $0 \le i \le n-1$. Then:

- (1) Nef(L) is a salient convex closed cone contained in the vector space $L \cdot H^{1,1}(X, \mathbb{R})$ and in the cone $\mathcal{K}^{i+1}(X)$.
- (2) $\mathcal{K}^{i+1}(X) \cap \mathcal{N}^{i+1}(X) = \{0\} \text{ and } \operatorname{Nef}(L) \cap \mathcal{N}^{i+1}(X) = \{0\}.$ In particular, $\mathcal{K}^{i+1}(X) + \mathcal{N}^{i+1}(X)$ and $\operatorname{Nef}(L) + \mathcal{N}^{i+1}(X)$ are closed in $H^{i+1,i+1}(X, \mathbb{R}).$
- (3) Let M_1, \ldots, M_p be in $H^{1,1}(X, \mathbb{R})$ such that LM_1, \ldots, LM_p are in Nef(L). Then $LM_1 \cdots M_p$ belongs to $\mathcal{K}^{i+p}(X)$. In particular, we have $LM_1 \cdots M_p = 0$ if and only if $LM_1 \cdots M_p c_1 \cdots c_{n-i-p} = 0$ for some Kähler classes c_1, \ldots, c_{n-i-p} .
- (4) Let M_1, \ldots, M_p be in $H^{1,1}(X, \mathbb{R})$ such that LM_1, \ldots, LM_p are in Nef $(L) + \mathcal{N}^{i+1}(X)$. Then $LM_1 \cdots M_p$ belongs to $\mathcal{K}^{i+p}(X) + \mathcal{N}^{i+p}(X)$.

Proof. (1) By definition, Nef(L) is convex and closed. Clearly, $L \cdot \text{Nef}(X)$ is contained in $L \cdot H^{1,1}(X, \mathbb{R})$ and in $\mathcal{K}^{i+1}(X)$ which are both closed. It follows that Nef(L) is also contained in $L \cdot H^{1,1}(X, \mathbb{R})$ and $\mathcal{K}^{i+1}(X)$. Since $\mathcal{K}^{i+1}(X)$ is salient, Nef(L) satisfies the same property.

(2) Since each class $\alpha \in \mathcal{K}^{i+1}(X) \setminus \{0\}$ can be represented by a non-zero positive closed current, we have $\alpha \cdot c_1 \cdots c_{n-i-1} \neq 0$ for all Kähler classes c_1, \ldots, c_{n-i-1} . Thus $\mathcal{K}^{i+1}(X) \cap \mathcal{N}^{i+1}(X) = \{0\}$ and $\operatorname{Nef}(L) \cap \mathcal{N}^{i+1}(X) = \{0\}$. The second assertion is then a direct consequence.

(3) For simplicity, we consider p = 2 as the general case can be obtained in the same way. Since LM_1, LM_2 are in Nef(L), there are nef classes $N_{1,k}, N_{2,\ell}$ such that $LN_{1,k}$ converges to LM_1 and $LN_{2,\ell}$ converges to LM_2 . We deduce that

$$LM_1M_2 = \lim_{k \to \infty} LN_{1,k}M_2 = \lim_{k \to \infty} LM_2N_{1,k} = \lim_{k \to \infty} LN_{2,\ell_k}N_{1,k},$$

where we choose l_k large enough so that $||LN_{2,\ell_k}N_{1,k} - LM_2N_{1,k}|| \leq 1/k$. It is now clear that LM_1M_2 belongs to $\mathcal{K}^{i+2}(X)$. The last assertion is true, as we just observed in (2), for every class in $\mathcal{K}^{i+2}(X)$ and in particular for the class LM_1M_2 .

(4) As above, we assume p = 2 for simplicity. By hypothesis, there are nef classes $N_{1,k}, N_{2,\ell}$ and classes $N_1^{(0)}, N_2^{(0)} \in \mathcal{N}^{i+1}(X)$ such that

$$LM_1 = \left(\lim_{k \to \infty} LN_{1,k}\right) + N_1^{(0)}$$
 and $LM_2 = \left(\lim_{\ell \to \infty} LN_{2,\ell}\right) + N_2^{(0)}.$

For suitable l_k going to infinity fast enough, we have

$$LM_1M_2 = \left(\lim_{k \to \infty} LN_{1,k}M_2\right) + N_1^{(0)}M_2 = \lim_{k \to \infty} \left(LN_{1,k}N_{2,l_k} + N^{(k)}\right)$$

with $N^{(k)} := N_{1,k}N_2^{(0)} + N_1^{(0)}M_2$. Since $LN_{1,k}N_{2,\ell} \in \mathcal{K}^{i+2}(X)$ and $N^{(k)} \in \mathcal{N}^{i+2}(X)$, using (2), we deduce from the above identities that $LM_1M_2 \in \mathcal{K}^{i+2}(X) + \mathcal{N}^{i+2}(X)$.

Let $G \subset \operatorname{Aut}(X)$ be a subgroup. Until the end of §3.3, we assume that the action $G \circlearrowleft H^{\bullet}(X, \mathbb{R})$ is unipotent; see Lemma 2.5.

Lemma 3.3. There is a sequence of classes $L_i \in \mathcal{K}^i(X) \setminus \{0\}$ for $0 \leq i \leq n$ such that:

- (1) $L_0 = 1$ and $L_{i+1} \in \operatorname{Nef}(L_i)$ for every $0 \le i \le n-1$. In particular, there is a class $M_{i+1} \in H^{1,1}(X, \mathbb{R})$ such that $L_{i+1} = L_i M_{i+1}$; and
- (2) L_i is fixed by G for every $0 \le i \le n$.

Note that in Lemma 3.3, for the reason similar to Remark 3.1, we cannot assume M_{i+1} to be nef.

Proof. We construct the sequence by induction. Assume that L_0, \ldots, L_i are already constructed and satisfy the above properties (1) and (2). We see that $\operatorname{Nef}(L_i) \neq 0$ is invariant by G. Since $G \circ H^{\bullet}(X, \mathbb{R})$ is unipotent, it factors through a solvable group action. Therefore, by a version of Lie-Kolchin's theorem for cone [16, Theorem 1.1], there is a ray in $\operatorname{Nef}(L_i)$ which is preserved by G. Choose L_{i+1} in this ray. Since 1 is the only eigenvalue of $g^*: H^{i+1,i+1}(X, \mathbb{R}) \circ$ for any $g \in G$, we deduce that L_{i+1} is fixed by G. This completes the proof of the lemma.

The sequence $M_1, \ldots, M_n \in H^{1,1}(X, \mathbb{R})$ as in Lemma 3.3 is called a *quasi-nef* sequence.

3.2. Dynamical filtrations.

Let $M_1, \ldots, M_n \in H^{1,1}(X, \mathbb{R})$ be a quasi-nef sequence constructed with respect to the *G*-action (see Lemma 3.3). Consider the following *G*-stable linear subspaces of $H^{1,1}(X, \mathbb{R})$. Set $F_n := H^{1,1}(X, \mathbb{R})$ and define for $0 \le i \le n-1$

(3.1)
$$F_{i} := \left\{ M \in H^{1,1}(X, \mathbb{R}) \mid L_{i}M \cdot H^{1,1}(X, \mathbb{R})^{n-i-1} = 0 \right\} \\ = (L_{i} \smile \bullet)^{-1} \left(\mathcal{N}^{i+1}(X) \right).$$

They form an increasing filtration

$$\cdots \subset F_i \subset F_{i+1} \subset \cdots$$
.

By Poincaré duality, in the definition of F_i we can replace n - i - 1 by n - i - 2 (when $i \leq n-2$) without changing this space. We have $F_0 = \mathcal{N}^1(X) = 0$. We also see that F_{n-1} is the hyperplane

$$\{ M \in H^{1,1}(X,\mathbb{R}) \mid L_{n-1}M = 0 \} \subset H^{1,1}(X,\mathbb{R}).$$

Consider Kähler classes c_1, \ldots, c_{n-i-2} and define the quadratic form Q_i on $H^{1,1}(X, \mathbb{R})$ by

(3.2)
$$Q_i(M, M') := L_i M M' c_1 \cdots c_{n-i-2} \text{ for } M, M' \in H^{1,1}(X, \mathbb{R}).$$

Here, the right hand side is a real number since $H^{n,n}(X,\mathbb{R})$ is canonically identified with \mathbb{R} . By the definition of F_i , this quadratic form induces a quadratic form on $H^{1,1}(X,\mathbb{R})/F_i$ that we still denote by Q_i . Consider another Kähler class c_{n-i-1} and the primitive subspace¹

$$P_i := \{ M \in H^{1,1}(X, \mathbb{R}) \mid L_i M c_1 \cdots c_{n-i-1} = 0 \}.$$

Since L_i belongs to $\mathcal{K}^i(X) \setminus \{0\}$, we have $L_i c_1 \cdots c_{n-i-1} \neq 0$. Therefore, by Poincaré duality, P_i is a hyperplane of $H^{1,1}(X,\mathbb{R})$. It contains F_i , hence P_i/F_i is a hyperplane of $H^{1,1}(X,\mathbb{R})/F_i$.

Lemma 3.4. The quadratic form Q_i is negative semi-definite on P_i , hence also on P_i/F_i . Moreover, Q_i is negative semi-definite on P_{i+1}/F_i , and hence on F_{i+1}/F_i .

¹This is a generalization of the classical primitive subspace $\operatorname{Ker}(c^{n-1} \smile \bullet) \subset H^{1,1}(X, \mathbb{R})$ for some Kähler class c.

We have seen that

$$P_{i+1} = \left\{ M \in H^{1,1}(X, \mathbb{R}) \mid L_{i+1}Mc_1 \cdots c_{n-i-2} = 0 \right\}.$$

is a hyperplane of $H^{1,1}(X,\mathbb{R})$. Recall that L_{i+1} can be approximated by classes $\{L_i c'_k\}_{k\geq 1}$ with c'_k Kähler. Therefore, we can approximate P_{i+1} by

$$P'_{k} := \left\{ M \in H^{1,1}(X, \mathbb{R}) \mid L_{i}Mc'_{k}c_{1}\cdots c_{n-i-2} = 0 \right\}.$$

Finally, since Q_i is negative semi-definite on P'_k , by continuity, it is negative semi-definite on P_{i+1} as well. This implies the second assertion of the lemma because F_{i+1} is contained in P_{i+1} .

Lemma 3.5. Let M and M' be classes in $H^{1,1}(X,\mathbb{R})$ such that

$$Q_i(M, M') = Q_i(M', M') = 0.$$

Assume also that $L_i M \in \operatorname{Nef}(L_i) \setminus \{0\}$. Then there is a vector $(a, b) \in \mathbb{R}^2 \setminus \{0\}$, unique up to a multiplicative constant, such that

$$L_i(aM+bM')c_1\cdots c_{n-i-2}=0.$$

If the subspace spanned by M and M' is a plane, then $Vect(M, M') \cap P$ is the real line spanned by aM + bM'.

Proof. Since the classes c_k are Kähler, the hypothesis on L_iM implies that

$$L_i M c_1 \cdots c_{n-i-2} c_{n-i-1} \neq 0$$
 and $L_i M c_1 \cdots c_{n-i-2} \neq 0$

Therefore, we have $M \notin P$ and (a, b) is unique up to a multiplicative constant. The lemma is clear when M and M' are collinear. Assume now that M and M' are not collinear. Since $\Pi := \mathsf{Vect}(M, M')$ is a plane, $\Pi \cap P$ is a line. Let N := aM + bM' be a vector which spans this real line. It is enough to show that this vector satisfies the identity in the lemma.

By Poincaré duality, we only have to show that $Q_i(N, N') = 0$ for every $N' \in H^{1,1}(X, \mathbb{R})$. Since $N \in \Pi \cap P$, Lemma 3.4 implies $Q_i(N, N) \leq 0$. We also have

$$Q_i(N,N) = a^2 Q_i(M,M) \ge 0,$$

because $L_i M^2 \in \mathcal{K}^{i+2}(X)$ by Lemma 3.2(3) and the hypothesis $L_i M \in \operatorname{Nef}(L_i)$. We deduce that $Q_i(N, N) = 0$, and that $Q_i(M, M) = 0$ when $a \neq 0$. In any case, we have $Q_i(N, N') = 0$ for $N' \in \Pi$ because $Q_i(M, M') = 0$.

Note that $H^{1,1}(X,\mathbb{R}) = \Pi + P$. Now, it suffices to check the identity $Q_i(N,N') = 0$ for $N' \in P$. Since $Q_i(N,N) = 0$, using Lemma 3.4 and Cauchy-Schwarz's inequality, we have for $N' \in P$

$$Q_i(N, N')^2 \le Q_i(N, N) \cdot Q_i(N', N') = 0.$$

The lemma follows.

For $1 \leq i \leq n$, define F'_i as the vector subspace of $F_i \subset H^{1,1}(X, \mathbb{R})$ spanned by the cone $C_i := \left\{ M \in F_i \mid L_{i-1}M \in \operatorname{Nef}(L_{i-1}) + \mathcal{N}^i(X) \right\}.$

We can deduce from Lemma 3.7 below that either $C_i = F_{i-1}$ or C_i is a closed half-space of F'_i having F_{i-1} as the boundary. We have

$$F_{i-1} \subset C_i \subset F'_i \subset F_i$$

since $L_{i-1}F_{i-1} \subset \mathcal{N}^i(X)$. Both C_i and F'_i are G-invariant. In particular, we have a G-invariant filtration

$$(3.3) \qquad \cdots \subset F'_{i-1} \subset F_i \subset F_i \subset \cdots$$

We also have $F'_n = H^{1,1}(X, \mathbb{R})$ since $F_n = H^{1,1}(X, \mathbb{R})$; thus C_n contains the nef cone Nef(X) of X.

Lemma 3.6. For $1 \le i \le n$, we have $C_i \cap (-C_i) = F_{i-1}$.

Proof. Clearly, F_{i-1} is contained in $C_i \cap (-C_i)$. For $M \in C_i$, there are nef classes N_k and a class $N^{(0)} \in \mathcal{N}^i(X)$ such that

$$L_{i-1}M = \lim_{k \to \infty} L_{i-1}N_k + N^{(0)}$$

In particular, if c_1, \ldots, c_{n-i} are Kähler classes, then $L_{i-1}Mc_1 \cdots c_{n-i} \ge 0$. Assume now that $M \in C_i \cap (-C_i)$. We obtain that $L_{i-1}Mc_1 \cdots c_{n-i} = 0$ for every Kähler classes c_1, \ldots, c_{n-i} . Thus, $L_{i-1}MH^{1,1}(X, \mathbb{R})^{n-i} = 0$ and $M \in F_{i-1}$.

Recall that $L_i = L_{i-1}M_i$ by the definition of M_i . Note also that dim $F'_n/F_{n-1} = 1$ because F_{n-1} is a hyperplane of $F'_n = H^{1,1}(X, \mathbb{R})$. Since $M_n \notin F_{n-1}$, we see that $F'_n = F_{n-1} \oplus \mathbb{R} \cdot M_n$.

Lemma 3.7. Let $1 \le i \le n$. We have the following properties.

- (1) dim $F'_i/F_{i-1} \le 1$ and $F'_i = \{ \alpha \in F_i \mid L_{i-1}\alpha^2 \equiv 0 \}.$
- (2) The following assertions are equivalent:
 - (a) dim $F'_i/F_{i-1} = 1;$

(b)
$$F'_i = F_{i-1} \oplus \mathbb{R} \cdot M_i$$
;

- (c) $L_{i-1}M_i^2 \equiv 0;$
- (d) $L_{i-1}M_i^2 = 0;$
- (e) C_i is the closed half-space of F'_i containing M_i and having F_{i-1} as the boundary.
- (3) For every $\alpha \in F'_i$ (resp. $\alpha \in C_i \setminus F_{i-1}$), $L_{i-1}\alpha$ is proportional (resp. positively proportional) to L_i modulo $\mathcal{N}^i(X)$.

Proof. Lemma 3.7 for i = n follows from

$$F'_n = F_{n-1} \oplus \mathbb{R} \cdot M_n = H^{1,1}(X, \mathbb{R}).$$

From now on, we assume that $i \leq n-1$.

(1) Let Q_{i-1} be the quadratic form defined using $L_{i-1}c_1 \cdots c_{n-i-1}$ for some Kähler classes c_j . By the second assertion of Lemma 3.4, if $M' \in F_i$ then $Q_{i-1}(M', M') \leq 0$. If $M' \in C_i$, we have $Q_{i-1}(M', M') \geq 0$ according to Lemma 3.2(4). Therefore, we have $Q_{i-1}(M', M') = 0$ for $M' \in C_i$ and hence for $M' \in F'_i$ as well because C_i spans F'_i . This proves

$$F'_i \subset \left\{ \alpha \in F_i \mid L_{i-1} \alpha^2 \equiv 0 \right\}.$$

The other inclusion and dim $F'_i/F_{i-1} \leq 1$ follow easily from the following claim.

Claim. Let M' be any class in F_i satisfying $Q_{i-1}(M', M') = 0$ for all Kähler classes c_1, \ldots, c_{n-i-1} . Then M' is proportional to M_i modulo F_{i-1} .

Proof of Claim. Assume to the contrary that M' is not proportional to M_i modulo F_{i-1} . By the definitions of F_i and M_i , we have $Q_{i-1}(M_i, M') = 0$; we also have $Q_{i-1}(M', M') = 0$ by hypothesis. According to Lemma 3.5, there is a vector $(a, b) \in \mathbb{R}^2 \setminus \{0\}$, unique up to a multiplicative constant, such that

(A)
$$L_{i-1}(aM_i + bM')c_1 \cdots c_{n-i-1} = 0.$$

Also since M' and M_i span a plane, by the second assertion of Lemma 3.5, this (a, b) is the unique (up to a multiplicative constant) solution of the equation

(B)
$$L_{i-1}(aM_i + bM')c_1 \cdots c_{n-i-1}c_{n-i} = 0.$$

So Equation (B) is equivalent to Equation (A) which can be obtained from (B) by removing the factor c_{n-i} . Since this property holds for arbitrary Kähler classes c_1, \ldots, c_{n-i} , Equation (B) is equivalent to any equation obtained from (B) by removing a factor c_j :

(A')
$$L_{i-1}(aM_i + bM')c_1 \cdots c_{j-1}c_{j+1} \dots c_{n-i-1}c_{n-i} = 0.$$

We conclude that (A) and (A') are equivalent. In other words, Equation (A) remains equivalent if we replace a factor c_j by any other Kähler class. Therefore, if (a, b) is as above, after replacing one by one the factors c_j with arbitrary Kähler classes, we have

 $L_{i-1}(aM_i + bM')c_1 \cdots c_{n-i-1} = 0$

for all Kähler classes c_1, \ldots, c_{n-i-1} . Since Kähler classes span $H^{1,1}(X, \mathbb{R})$, we obtain

$$L_{i-1}(aM_i + bM')H^{1,1}(X, \mathbb{R})^{n-i-1} = 0$$

Hence M' has to be proportional to M_i modulo F_{i-1} .

(2) The above claim implies that (a) implies (b). That (b) implies (c) follows from Part (1) of the lemma. As $L_{i-1}M_i = L_i \in \operatorname{Nef}(L_{i-1}) \subset \mathcal{K}^i(X)$, it follows from Lemma 3.2(3) that (c) is equivalent to (d). We show that (c) implies (a). Since dim $F'_i/F_{i-1} \leq 1$, it suffices to show that $F_{i-1} \neq F'_i$. Assume that $L_{i-1}M_i^2 \equiv 0$, then $M_i \in F_i$ because $L_iM_i = L_{i-1}M_i^2 \equiv 0$, and therefore $M_i \in F'_i$ by Part (1) of the lemma. As $L_{i-1}M_i = L_i \neq 0$, we have $M_i \notin F_{i-1}$. Hence $F_{i-1} \neq F'_i$ and (c) implies (a). It is clear that (e) implies (a). Finally, assuming (a), (b), (c), it is clear that $M_i \in C_i$, $M_i \notin F_{i-1}$ and hence (e) is true, thanks to Lemma 3.6.

(3) Recall that $L_i = L_{i-1}M_i$. The assertion is clear when $F'_i = C_i = F_{i-1}$. Otherwise, this is a consequence of (b) and (d) in Part (2).

3.3. Polynomial growth of the pullback action: proofs of Theorems 1.1 and 1.3.

Let $M_1, \ldots, M_n \in H^{1,1}(X, \mathbb{R})$ be a quasi-nef sequence with respect to the *G*-action. Let F_i and F'_i be the filtrations constructed previously.

Let $g \in G$ and let ω be a Kähler class. We have the expansion:

(3.4)
$$(g^m)^*(\omega) = ((g^* - \operatorname{Id}) + \operatorname{Id})^m(\omega) = \sum_{0 \le j \le m} \binom{m}{j} \omega_j \text{ with } \omega_j := (g^* - \operatorname{Id})^j(\omega).$$

The following proposition combined with Lemma 3.7 proves Theorem 1.3.

Proposition 3.8. Recall that $g^* : H^{1,1}(X) \circlearrowleft$ is unipotent. Assume that $g^* \neq \text{Id}$. Then there exist an integer $1 \le r \le n-1$ and a unique strictly decreasing sequence of integers

$$n-1 \ge s_1 > \dots > s_r \ge 1$$

depending on g but independent of the Kähler class ω , such that $\omega_{2j-1} \in F_{s_j} \setminus F'_{s_j}$ and $\omega_{2j} \in F'_{s_j} \setminus F_{s_j-1}$ for $1 \leq j \leq r$, and $\omega_{2r+1} = 0$. Moreover, we have that dim $F'_{s_j}/F_{s_j-1} = 1$, ω_{2j} is positively proportional to M_{s_j} modulo F_{s_j-1} , and $F'_{s_j} = F_{s_j-1} \oplus \mathbb{R} \cdot M_{s_j}$.

For later use (after Lemma 3.12), in order to recall that the s_j in Proposition 3.8 depends on $g \in \text{Aut}(X)$, we will sometimes use the notation $s_j(g) := s_j$. We will also use the notation $\omega_1(g) := \omega_1$ in the expansion (3.4) from time to time.

Before proving Proposition 3.8, we first deduce the upper bounds stated in Theorem 1.1.

Proof of Theorem 1.1 (for the optimality, see §4.1). Observe that it is enough to prove this theorem for any positive iterate of g instead of g. Since g has zero entropy, up to replacing g by some positive iterate, we can assume by Lemma 2.5 that $g^* : H^{1,1}(X) \circlearrowleft$ is unipotent. We apply the above study for $G \subset \operatorname{Aut}(X)$ the cyclic group generated by g. We deduce from Proposition 3.8 that for $m \geq 2r$,

$$(g^m)^*(\omega) = \sum_{0 \le j \le 2r} \binom{m}{j} \omega_j.$$

Since $r \leq n-1$, we obtain that $||(g^m)^*(\omega)|| \lesssim m^{2(n-1)}$ and hence $||(g^m)^*(\omega^p)|| \lesssim m^{2p(n-1)}$. Since ω^p is in the interior of the cone $\mathcal{K}^p(X)$, it follows that (see also [5, Proposition 5.8])

 $\|(g^m)^*: H^{p,p}(X,\mathbb{R}) \circlearrowleft \| \lesssim m^{2p(n-1)}.$

Since $(g^m)^* : H^{p,p}(X, \mathbb{R}) \circlearrowleft$ is dual to $(g^{-m})^* : H^{n-p,n-p}(X, \mathbb{R}) \circlearrowright$, we deduce that $\|(g^m)^* : H^{p,p}(X, \mathbb{R}) \circlearrowright \| \lesssim m^{2(n-p)(n-1)}.$

Thus,

$$\|(g^m)^*: H^{p,p}(X,\mathbb{R}) \circlearrowleft \| \lesssim m^{2p'(n-1)}$$

with $p' = \min(p, n-p)$. Finally, we obtain the estimate given in Theorem 1.1 and conclude its proof by using:

$$||(g^m)^* : H^{p,q}(X,\mathbb{R}) \circlearrowleft ||^2 \le C ||(g^m)^* : H^{p,p}(X,\mathbb{R}) \circlearrowright |||(g^m)^* : H^{q,q}(X,\mathbb{R}) \circlearrowright ||$$

for some constant C > 0 independent of g and m, see the proof of [5, Proposition 5.8]. \Box

Remark 3.9.

(1) A well-known result says that if $||(g^m)^* : H^{1,1}(X, \mathbb{R}) \circ || \sim m^k$ when $m \gg 1$, then k = 2r is always an *even* integer. Indeed, if k is odd, then for a general Kähler class ω , we see from the expansion (3.4) that the Kähler classes $(g^m)^*(\omega)$ and $(g^{-m})^*(\omega)$ would be asymptotically opposite. This contradicts the property that the cone Nef(X) is salient.

(2) For the other values of (p,q), the estimate can be slightly improved. Indeed, this is clear when r < n-1. When r = n-1, we have $s_i = n-i$ for every *i* and we can take $L_1 = M_1 = \omega_{2r}$. Then, by Lemma 3.7 and Proposition 3.8, we get $\omega_{2r}^2 \equiv 0$ and $\omega_{2r}\omega_{2r-1} \equiv 0$. Thus the following holds whenever $p \geq 2$ and regardless of the value of *r*:

$$||(g^m)^* : H^{p,p}(X,\mathbb{R}) \circlearrowleft || = O(m^{2p(n-1)-2}).$$

Proof of Proposition 3.8. First observe that $g^* = \text{Id on } F'_i/F_{i-1}$ because g^* is unipotent and dim $F'_i/F_{i-1} \leq 1$ according to Lemma 3.7. In particular, if $\omega_k \in F'_i \setminus F_{i-1}$ then $\omega_p \notin F'_i$ for p < k and $\omega_{k+1} \in F_{i-1}$. This remark will be used several times in the proof.

Starting from a Kähler class ω , we first construct by induction the sequence s_j which verifies the properties in Proposition 3.8 except its independence of ω . We will show later that r and s_j do not depend on the choice of ω .

We have $\omega_0 = \omega \in F'_n = H^{1,1}(X, \mathbb{R})$ and for simplicity we set $s_0 := n$. Suppose that $j \geq 1$ and the sequence (s_k) is already constructed for k up to j-1, so that $\omega_{2i-1} \in F_{s_i} \setminus F'_{s_i}$ and $\omega_{2i} \in F'_{s_i} \setminus F_{s_i-1}$ for $i = 1, \ldots, j-1$. For i = j-1 (when j > 1), this gives

$$\omega_{2j-3} \in F_{s_{j-1}} \setminus F'_{s_{j-1}} \quad \text{and} \quad \omega_{2j-2} \in F'_{s_{j-1}} \setminus F_{s_{j-1}-1}$$

If $\omega_{2j-1} = 0$ (in particular, when $s_{j-1} = 1$) we end the construction and define r := j - 1. Assume now that $\omega_{2j-1} \neq 0$, so $s_{j-1} > 1$. It is enough to show the existence of $1 \leq s_j < s_{j-1}$ such that $\omega_{2j-1} \in F_{s_j} \setminus F'_{s_j}$ and $\omega_{2j} \in F'_{s_j} \setminus F_{s_j-1}$.

Let $s_j \geq 1$ be the integer such that $\omega_{2j-1} \in F_{s_j} \setminus F_{s_j-1}$. By Lemma 3.7, the space $F'_{s_{j-1}}/F_{s_{j-1}-1}$ is of dimension 0 or 1. Therefore, since $\omega_{2j-1} = (g^* - \operatorname{Id})(\omega_{2j-2})$ and ω_{2j-2} belongs to $F'_{s_{j-1}} \setminus F_{s_{j-1}-1}$, we necessarily have $\omega_{2j-1} \in F_{s_{j-1}-1}$. It follows that $s_j < s_{j-1}$.

Let $k \geq 2j-1$ be the largest integer such that ω_k belongs to $F_{s_j} \setminus F_{s_j-1}$. So we have $L_{s_j-1}\omega_k \cdot H^{1,1}(X,\mathbb{R})^{n-s_j} \neq 0$. We only need to show that $\omega_k \in F'_{s_j}$ and k = 2j. Note that these properties imply that $\omega_{2j-1} \notin F'_{s_j}$ thanks to the remark at the beginning of the proof.

Observe that modulo F_{s_i-1} we have

$$(g^m)^*(\omega) = \omega_0 + \binom{m}{1}\omega_1 + \dots + \binom{m}{k}\omega_k$$

Therefore, there are classes $N^{(m)}$ in $L_{s_i-1}F_{s_i-1} \subset \mathcal{N}^{s_j}(X)$ such that

$$L_{s_j-1}\omega_k = \lim_{m \to \infty} \left((k!)m^{-k}L_{s_j-1}(g^m)^*(\omega) + N^{(m)} \right).$$

Lemma 3.2(2) implies that $L_{s_j-1}\omega_k$ is in Nef $(L_{s_j-1}) + \mathcal{N}^{s_j}(X)$. Since $\omega_k \in F_{s_j}$, we conclude that ω_k belongs to $C_{s_j} \subset F'_{s_j}$. As $\omega_k \notin F_{s_j-1}$, by Lemma 3.7(3), ω_k is positively proportional to M_{s_j} in F'_{s_j}/F_{s_j-1} .

Since $\omega_k \in F'_{s_j} \setminus F_{s_j-1}$, Lemma 3.6 implies that $-\omega_k$ doesn't belong to C_{s_j} . Since $(g^{-m})^* = (\mathrm{Id} - (g^*)^{-1} (g^* - \mathrm{Id}))^m$, we have

$$(g^{-m})^*(\omega) = \omega_0 - \binom{m}{1}(g^{-1})^*(\omega_1) + \dots + (-1)^k \binom{m}{k}(g^{-k})^*(\omega_k) \in H^{1,1}(X,\mathbb{R})/F_{s_j-1}$$

which implies

$$(g^{k-m})^*(\omega) = (g^k)^*(\omega_0) - \binom{m}{1}(g^{k-1})^*(\omega_1) + \dots + (-1)^k\binom{m}{k}\omega_k \in H^{1,1}(X,\mathbb{R})/F_{s_j-1}.$$

Hence for some classes $N^{[m]}$ in $L_{s_j-1}F_{s_j-1} \subset \mathcal{N}^{s_j}(X)$,

$$(-1)^{k} L_{s_{j}-1}\omega_{k} = \lim_{m \to \infty} \left((k!)m^{-k} L_{s_{j}-1}(g^{k-m})^{*}(\omega) + N^{[m]} \right).$$

It follows as above that $(-1)^k \omega_k$ belongs to C_{s_j} . Thus, k is even and hence $k \geq 2j$.

To show that k = 2j, we assume by contradiction that $k \ge 2j+2$. Observe that, for $q \ge 2j-1$ we have $\omega_q \in F_{s_j}$ by the definition of s_j . Therefore $L_{s_j-1}M_{s_j}\omega_q = L_{s_j}\omega_q \in \mathcal{N}^{s_j+1}(X)$,

and hence $L_{s_j-1}\omega_k\omega_q \in \mathcal{N}^{s_j+1}(X)$ (where we recall that $L_{s_j-1}\omega_k$ is positively proportional to $L_{s_j-1}M_{s_j}$ modulo $\mathcal{N}^{s_j}(X)$). In particular, this holds for $q \geq k-2$. We also observe that by the definition of k, we have $\omega_q \in F_{s_j-1}$ whenever $q \geq k+1$, so $L_{s_j-1}\omega_q \in \mathcal{N}^{s_j}(X)$. Thus, we have (using the expansion (3.4))

$$\lim_{m \to \infty} \left((k-1)!^2 m^{-2k+2} L_{s_j-1}(g^m)^*(\omega^2) \right) = L_{s_j-1} \omega_{k-1}^2 \mod \mathcal{N}^{s_j+1}(X).$$

When we multiply by Kähler classes, the left hand side gives a non-negative number while the right hand side gives a non-positive number due to Lemma 3.4 (because $\omega_{k-1} \in F_{s_j}$). We then obtain that $L_{s_j-1}\omega_{k-1}^2 = 0$ modulo $\mathcal{N}^{s_j+1}(X)$. This contradicts Lemma 3.7(1) and the fact that $\omega_{k-1} \in F_{s_j} \setminus F'_{s_j}$, the latter due to $\omega_k \in F'_{s_j} \setminus F_{s_j-1}$ and the remark at the beginning of the proof. Thus k = 2j as desired. Finally, the fact that ω_k is positively proportional to M_{s_j} in F'_{s_j}/F_{s_j-1} together with Lemma 3.7(2) shows that s_j satisfies the "moreover part" of Proposition 3.8.

We prove now that the sequence s_j doesn't depend on the choice of ω . Let ω' be another Kähler class. We have $\omega \leq \omega' \leq \omega$ which implies that

$$L_i(g^m)^*(\omega) \lesssim L_i(g^m)^*(\omega') \lesssim L_i(g^m)^*(\omega)$$
.

It follows from the polynomial growth (when $m \to \infty$) of the expansion

$$L_i c_1 \cdots c_{n-i-1} (g^m)^*(\omega) = \sum_{0 \le j \le 2r} \binom{m}{j} L_i c_1 \cdots c_{n-i-1} \omega_j \qquad (c_1, \ldots, c_{n-i-1} \text{ K\"ahler})$$

and the similar one for ω' that k is the smallest integer such that $L_i\omega_k \in \mathcal{N}^{i+1}(X)$ if and only if k is the smallest integer such that $L_i\omega'_k \in \mathcal{N}^{i+1}(X)$. So for any integer k, $\omega_k \in F_i$ if and only if $\omega'_k \in F_i$, which shows that s_j is independent of ω .

Finally, the uniqueness of the sequence (s_k) is clear from the property $\omega_{2j-1} \in F_{s_j} \setminus F'_{s_j}$ and (3.3). This completes the proof of Proposition 3.8 and also that of Theorem 1.3.

3.4. Essential derived length: proof of Theorem 1.2.

Thanks to Lemmas 2.4 and 2.5, in order to prove Theorem 1.2, we may and will replace G by a suitable finite-index subgroup and assume that the properties (1) and (2) in Lemma 2.5 hold for G instead of G'.

From now on, we fix some classes L_i with $0 \le i \le n$ and M_i with $1 \le i \le n$ as in Lemma 3.3. Define the group H_i by

$$H_i := \{g \in G : g^* = \operatorname{Id} \text{ on } L_i \cdot H^{1,1}(X, \mathbb{R}) / \mathcal{N}^{i+1}(X) \}$$
$$= \{g \in G : L_i \cdot (g^* - \operatorname{Id}) H^{1,1}(X, \mathbb{R}) \subset \mathcal{N}^{i+1}(X) \},\$$

where the quotient in the first display is understood as the quotient of $L_i \cdot H^{1,1}(X, \mathbb{R})$ by its intersection with $\mathcal{N}^{i+1}(X)$. Equivalently,

(3.5)
$$H_i = \{g \in G : (g^* - \mathrm{Id})H^{1,1}(X, \mathbb{R}) \subset F_i \} \\ = \{g \in G : g^* = \mathrm{Id} \text{ on } H^{1,1}(X, \mathbb{R})/F_i \}.$$

Since $F_{i+1} \supset F_i$, the sequence H_i is increasing, H_i is normal in G and hence in H_j for $i \leq j$. Moreover, G/H_i acts faithfully on $L_i \cdot H^{1,1}(X, \mathbb{R})/\mathcal{N}^{i+1}(X)$. Also $H_0 = G_0$, because Lemma 2.5 holds for G instead of G'. So if $H_0 = G$ or equivalently if G acts trivially on $H^{1,1}(X, \mathbb{R})$, then Theorem 1.2 holds obviously.

Lemma 3.10. Assume that $H_0 \neq G$. Then there is an integer $0 \leq l \leq n-2$ such that $H_l \neq G$ and $H_{l+1} = G$.

Proof. Since G acts trivially on $H^{n,n}(X,\mathbb{R})$, $H_{n-1} = G$. The lemma follows easily.

For $1 \leq j \leq n-1$, we define

$$W_j := \left\{ \begin{array}{l} \alpha \in F_j \\ L_{j-1}(M_j\beta + \alpha^2)c_1 \cdots c_{n-j-1} \ge 0 \text{ for all Kähler classes } c_i \end{array} \right\}$$

Note that we obtain the same space if we assume that β is Kähler, because when the inequality in the definition of W_j holds for β it holds for all β' such that $\beta' - \beta$ is nef. We call W_j a primitive root space².

Lemma 3.11. For $1 \le j \le n-1$, W_j is a G-invariant vector space containing F'_j .

Proof. It is clear that W_j is invariant by G and $W_j \supset F'_j$, see Lemma 3.7. Moreover, W_j is stable under scalar multiplications, so we only need to show that $W_j + W_j \subset W_j$. Fix Kähler classes c_1, \ldots, c_{n-j-1} and let Q_{j-1} be the quadratic form (3.2) defined using $L_{j-1}c_1 \cdots c_{n-j-1}$. By Lemma 3.4, Q_{j-1} is negative semi-definite on F_j , so the function $F_j \ni v \mapsto Q_{j-1}(v, v)$ is concave. In particular, for every $\alpha_1, \alpha_2 \in W_j$,

$$Q_{j-1}(\alpha_1 + \alpha_2, \alpha_1 + \alpha_2) \geq \frac{1}{2} (Q_{j-1}(2\alpha_1, 2\alpha_1) + Q_{j-1}(2\alpha_2, 2\alpha_2))$$

= 2(Q_{j-1}(\alpha_1, \alpha_1) + Q_{j-1}(\alpha_2, \alpha_2)).

It follows that $\alpha_1 + \alpha_2 \in W_j$.

Lemma 3.12. Let ω be a Kähler class. Then, the class

$$\omega_{2j-1} = (g^* - \operatorname{Id})^{2j-1}(\omega)$$

belongs to W_{s_j} for every $1 \leq j \leq r$.

Proof. We have for every Kähler classes c_i

$$m^{-4j+2}L_{s_j-1}(g^m)^*(\omega^2)c_1\cdots c_{n-s_j-1} \ge 0.$$

By Proposition 3.8, $L_{s_j-1}\omega_p H^{1,1}(X,\mathbb{R})^{n-s_j} = 0$ whenever $p \ge 2j+1$. Observe also that

$$L_{s_j-1}\omega_{2j}\omega_q H^{1,1}(X,\mathbb{R})^{n-s_j-1} = L_{s_j}\omega_q H^{1,1}(X,\mathbb{R})^{n-s_j-1} = 0$$

whenever $q \ge 2j - 1$ because ω_{2j} is proportional to M_{s_j} modulo F_{s_j-1} by Proposition 3.8. Therefore, using the expansion (3.4), we get from the last inequality

$$m^{-4j+2}L_{s_j-1}\left(2\binom{m}{2j}\binom{m}{2j-2}\omega_{2j}\omega_{2j-2} + \binom{m}{2j-1}^2\omega_{2j-1}^2\right)c_1\cdots c_{n-s_j-1} + o(1) \ge 0.$$

Multiplying the last line by (2j)!(2j-1)! and taking $m \to \infty$ give

$$L_{s_j-1}((4j-2)\omega_{2j}\omega_{2j-2}+(2j)\omega_{2j-1}^2)c_1\cdots c_{n-s_j-1} \ge 0.$$

By Proposition 3.8, ω_{2j} is positively proportional to M_{s_j} modulo F_{s_j-1} . Therefore, the last inequality implies that $\omega_{2j-1} \in W_{s_j}$.

 $^{^{2}\}alpha$ can be regarded as a square root of a class which is bounded below by $-M_{j}\beta$ in some sense.

End of the proof of Theorem 1.2. We can assume that $H_0 \neq G$ as in Lemma 3.10. Take an integer l as in that lemma. Since $l \leq n-2$, it is enough to show that $\ell(G/G_0) \leq l+1$.

Since $H_0 = G_0$ and $H_{l+1} = G$, in order to complete the proof, it is enough to show that H_{i+1}/H_i is commutative. For this purpose, we can assume that $H_{i+1} \neq H_i$. Observe also that H_{i+1}/H_i acts faithfully on $L_i H^{1,1}(X, \mathbb{R})/\mathcal{N}^{i+1}(X)$ and on $H^{1,1}(X, \mathbb{R})/F_i$, see (3.5). We continue to use the expansion (3.4). By (3.5) and Proposition 3.8, h belongs to H_i if and only if $\omega_1(h) \in F_i$ for every Kähler class ω , or equivalently, $s_1(h) \leq i$. Therefore, for any $g \in H_{i+1} \setminus H_i$, we have $s_1(g) = i + 1$ and, since $\omega_1(g) \notin F'_{i+1}$ by Proposition 3.8, H_{i+1}/H_i acts faithfully on $H^{1,1}(X, \mathbb{R})/F'_{i+1}$.

The following lemma is obvious.

Lemma 3.13. Let V be a vector space over an arbitrary field. Let Γ be a subset of GL(V). Assume there is a vector subspace W of V invariant by Γ such that Γ acts trivially on W and on V/W. Then Γ is a commutative set.

By Lemmas 3.12 and Lemma 3.13, it is enough to show that g acts trivially on W_{i+1}/F'_{i+1} . Assume by contradiction that the last property is wrong. Then since

$$(g^* - \mathrm{Id})^2(H^{1,1}(X,\mathbb{R})) \subset F'_{s_1(g)} = F'_{i+1},$$

there are classes α and M' in $W_{i+1} \setminus F'_{i+1}$ such that $(g^m)^*(\alpha) = \alpha + mM'$ modulo F'_{i+1} . Choose a class β as in the definition of W_{i+1} . As we observed above, we can assume that β is a Kähler class ω . Using that definition, the expansion (3.4), Lemma 3.4, and Proposition 3.8, we have for every Kähler classes c_j

$$0 \le -L_i(g^m)^*(\alpha^2)c_1 \cdots c_{n-i-2} \le L_i M_{i+1}(g^m)^*(\omega)c_1 \cdots c_{n-i-2} = L_i M_{i+1}\omega_0 c_1 \cdots c_{n-i-2}.$$

Here, since $L_i M_{i+1} = L_{i+1}$ and $\omega_j \in F_{i+1}$ for $j \geq 1$ (because $s_1 = i + 1$), these ω_j do not contribute for last equality. Taking $m \to \infty$, we see that $L_i M'^2 c_1 \cdots c_{n-i-2} = 0$. We have used here Lemma 3.7(3) which implies that $L_i W_{i+1} F'_{i+1} \subset \mathcal{N}^{i+2}(X)$. So $L_i M'^2 = 0$ modulo $\mathcal{N}^{i+2}(X)$. According to Lemma 3.7, we have $M' \in F'_{i+1}$ which is a contradiction. This completes the proof of Theorem 1.2.

4. Explicit examples

Let $n \geq 1$ be an integer and let $E = \mathbb{C}/\Lambda$ be an elliptic curve.

4.1. Torus examples for the optimality of the upper bound in Theorem 1.1.

Let $X := E^n = \mathbb{C}^n / \Lambda^n$ with coordinates z_1, \ldots, z_n . Define the automorphism $g : X \to X$ by

$$g(z_1,\ldots,z_n) = (z_1,z_1+z_2,\ldots,z_{n-1}+z_n).$$

With respect to the bases dz_1, \ldots, dz_n and $d\overline{z}_1, \ldots, d\overline{z}_n$ of $H^{1,0}(X)$ and $H^{0,1}(X)$ respectively, both $g^* : H^{1,0}(X) \to H^{1,0}(X)$ and $g^* : H^{0,1}(X) \to H^{0,1}(X)$ correspond to the Jordan matrix

$$J_n = \begin{bmatrix} 1 & 1 & & \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ & & & & 1 \end{bmatrix}.$$

Since $g^* : H^{1,1}(X) \to H^{1,1}(X)$ is isomorphic to the tensor product of $g^* : H^{1,0}(X) \to H^{1,0}(X)$ with $g^* : H^{0,1}(X) \to H^{0,1}(X)$, the corresponding matrix of $g^* : H^{1,1}(X) \to H^{1,1}(X)$ is the Kronecker product (see e.g. [14])

$$J_n \otimes J_n = \begin{bmatrix} J_n & J_n & & \\ & \ddots & \ddots & \\ & & \ddots & J_n \\ & & & J_n \end{bmatrix}$$

In particular, $g^*: H^{1,1}(X) \to H^{1,1}(X)$ is unipotent. Set $N_n := J_n - \mathrm{Id}_n$,

$$N := \begin{bmatrix} N_n & 0 & & \\ & \ddots & \ddots & \\ & & \ddots & 0 \\ & & & N_n \end{bmatrix} \quad \text{and} \quad J := \begin{bmatrix} 0 & J_n & & \\ & \ddots & \ddots & \\ & & \ddots & J_n \\ & & & 0 \end{bmatrix}$$

The linear map $(g^* - \mathrm{Id}) : H^{1,1}(X) \to H^{1,1}(X)$ is represented by the matrix N + J. We have $N^k = J^k = 0$ for every $k \ge n$ and

$$N^{n-1} = \begin{bmatrix} N_n^{n-1} & 0 & & \\ & \ddots & \ddots & \\ & & \ddots & 0 \\ & & & N_n^{n-1} \end{bmatrix} \quad \text{and} \quad J^{n-1} = \begin{bmatrix} 0 & \dots & 0 & J_n^{n-1} \\ & \ddots & & 0 \\ & & & \ddots & \vdots \\ & & & & 0 \end{bmatrix}.$$

Note that the multiplication of the first row of N^{n-1} with the last column of J^{n-1} is not zero. Therefore, since NJ = JN, we have

$$(N+J)^{2n-2} = \sum_{k=0}^{2n-2} {\binom{2n-2}{k}} N^k J^{2n-2-k} = {\binom{2n-2}{n-1}} N^{n-1} J^{n-1} \neq 0.$$

This example shows that

$$||(g^m)^*: H^2(X, \mathbb{C}) \circlearrowleft || \sim_{m \to \infty} Cm^{2(n-1)}$$

for some C > 0, thus the upper bound of the polynomial growth in Theorem 1.1 is optimal in terms of n.

Remark 4.1. More generally, we have

$$||(g^m)^*: H^{p,q}(X,\mathbb{R}) \circlearrowleft || \sim_{m \to \infty} Cm^{p(n-p)+q(n-q)}.$$

Thus the estimate in Theorem 1.1 is optimal for $p, q \in \{0, 1, n-1, n\}$. We omit the proof.

4.2. Some explicit examples related to Conjecture 1.4.

4.2.1. Torus examples.

Denote by $U(n, \mathbb{Z})$ the group of $n \times n$ upper triangular matrices whose entries are integers and whose diagonal entries are 1. Let E_{ij} be the $n \times n$ -matrix whose (i, j)-entry is 1 and other entries are 0. It is well-known and easy to see directly that $U(n, \mathbb{Z})$ is generated by

$$\tau_{ij} := I_n + E_{ij} \quad \text{with} \quad 1 \le i < j \le n \,,$$

and $U(n, \mathbb{Z})$ has nilpotency class n - 1.

The group $G := U(n, \mathbb{Z})$ acts on $X = E^n$, and the induced action on $H^2(E^n, \mathbb{Z}) = \wedge^2 H^1(E^n, \mathbb{Z})$ is faithful and unipotent. Hence, $G \subset \operatorname{Aut}(X)$ is a zero entropy subgroup. We have $G \cap \operatorname{Aut}^0(X) = \{1\}$ because $U(n, \mathbb{Z})$ acts on $H^2(E^n, \mathbb{Z})$ faithfully. Thus, by Lemma 2.4, we have

$$c_{\text{ess}}(G, X) = c(U(n, \mathbb{Z})) = n - 1 = \dim X - 1.$$

4.2.2. Some examples with positive Kodaira dimension.

Let $n \geq 2$ and κ be integers such that $1 \leq \kappa \leq n-1$. Fix a smooth projective variety B of dimension κ having an ample canonical divisor K_B (hence $\kappa(B) = \dim B = \kappa$) such that there is a surjective morphism $\rho: B \to E$. One may find such a B as a general member of a very ample linear system of $V := E^{\kappa+1}$ and ρ the projection to the first factor of $E^{\kappa+1}$. Indeed, by the adjunction formula, $K_B = (K_V + B)|_B \sim B|_B$ is the restriction of an ample divisor, whence it is ample.

Consider the product $X := E^{n-\kappa} \times B$. This is a smooth projective variety of dimension $n \ge 2$. By [26, Theorem 15.1],

$$\kappa(X) = \kappa(B) = \dim B = \kappa.$$

The group $U(n - \kappa, \mathbb{Z})$ acts faithfully on X by

$$U(n-\kappa,\mathbb{Z}) = U(n-\kappa,\mathbb{Z}) \times \{\mathrm{Id}_B\} \subset \mathrm{Aut}((E,0)^{n-\kappa}) \times \mathrm{Aut}(B) \subset \mathrm{Aut}(E^{n-\kappa} \times B).$$

For each *i* with $1 \leq i \leq n - \kappa$, we define $\rho_i \in \operatorname{Aut}(X)$ by

$$E^{n-\kappa} \times B \ni (z_1, \dots, z_i, \dots, z_{n-\kappa}, w) \mapsto (z_1, \dots, z_i + \rho(w), \dots, z_{n-\kappa}, w) \in E^{n-\kappa} \times B.$$

Then $\rho_i \in \operatorname{Aut}(X/B)$ with respect to the canonical projection onto B. Denote by A the abelian subgroup of $\operatorname{Aut}(X)$ generated by the ρ_i for $1 \leq i \leq n - \kappa$. Finally, define G to be the subgroup of $\operatorname{Aut}(X)$ generated by A and $U(n - \kappa, \mathbb{Z})$.

Since A (resp. $U(n-\kappa,\mathbb{Z})$) acts on each fiber $E^{n-\kappa}$ as a translation (resp. zero entropy) group, by the relative dynamical degree formula in [10, Theorem 1.1], the action of G on X is of zero entropy. This can also be obtained using the definition of entropy and the fact that for $g \in G$ the Lipschitz norm of g^m is bounded by a polynomial in m.

Proposition 4.2. Let X be as above. Then, $G_0 = \{1\}$ and

$$c_{\rm ess}(G, X) = \dim X - \kappa(X).$$

Proof. Using Blanchard's lemma [2, Cor. 2.3], for $X = E^{n-\kappa} \times B$, we obtain the following natural isomorphisms

$$\operatorname{Aut}^{0}(X) \cong \operatorname{Aut}^{0}(E^{n-\kappa}) \times \operatorname{Aut}^{0}(B) \cong E^{n-\kappa} \times {\operatorname{Id}}_{B} \cong E^{n-\kappa}.$$

Here, we also use the assumption that B is of general type and hence Aut(B) is finite, see [26, Cor. 14.3]; it follows that $Aut^{0}(B)$ is trivial.

Let $t \in B$ be a very general point. Then $\rho(t) \in (E, 0)$ is a non-torsion point, i.e., an infinite order element of the abelian group (E, 0). Then, the action of G on

$$q^{-1}(t) = E^{n-\kappa} \times \{t\} \simeq E^{n-\kappa}$$

is faithful and is given, for $z \in E^{n-\kappa}$, by

$$z \mapsto Az + \rho(t)b$$
 with $A \in U(n - \kappa, \mathbb{Z})$ and $b \in \mathbb{Z}^{n-\kappa}$

If $t' \in B$ is another very general point, the intersection between $\rho(t)\mathbb{Z}^{n-\kappa}$ and $\rho(t')\mathbb{Z}^{n-\kappa}$ is trivial, i.e., equal to $\{0\}$. Therefore, we see that the identity is the only element in Gwhich belongs to $\operatorname{Aut}^0(X)$. Thus, we have $G_0 = \{1\}$ and therefore $c_{\operatorname{ess}}(G, X) = c(G)$, provided that there exists an embedding $G \hookrightarrow \operatorname{GL}(V)$ of G as a unipotent subgroup for some finite dimensional vector space V. Indeed, by the definition of $c_{\operatorname{ess}}(G, X)$, there exists a finite index subgroup $G' \subset G$ such that $c_{\operatorname{ess}}(G, X) = c(G'/G'_0)$. As $G_0 = \{1\}$, we have $G'_0 = \{1\}$. Thus

$$c_{\rm ess}(G, X) = c(G'/G'_0) = c(G') = c(G),$$

where the last equality follows from Lemma 2.4.

It remains to find such an embedding $G \hookrightarrow \operatorname{GL}(V)$ and show that $c(G) = n - \kappa$. This is now a purely group theoretical problem and we don't need to use the action on cohomology. As G acts faithfully on $q^{-1}(t)$, from the description of the action of G on $q^{-1}(t)$ we may identify G with the unipotent affine transformation subgroup

$$P := \mathbb{Z}^{n-\kappa} \rtimes U(n-\kappa,\mathbb{Z}) \cong U(n-\kappa+1,\mathbb{Z}) \subset \mathrm{GL}(n-\kappa+1,\mathbb{R}).$$

Here, the last isomorphism is the natural one given by

$$\mathbb{Z}^{n-\kappa} \rtimes U(n-\kappa,\mathbb{Z}) \ni f(x) = Ax + b \leftrightarrow \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix} \in U(n-\kappa+1,\mathbb{Z}).$$

It also follows that $c(G) = n - \kappa$. This proves Proposition 4.2.

4.2.3. Some rationally connected examples.

Finally, we construct examples with Kodaira dimension $-\infty$. Let $n \ge 2$. Let E_{ω} be an elliptic curve with period $\omega = (-1 + \sqrt{-3})/2$ a primitive third root of unity. Let

$$\overline{X}_n := E_\omega^n / \langle -\omega I_n \rangle,$$

 $\pi: E_{\omega}^n \to \overline{X}_n$ the quotient map, and $X_n \to \overline{X}_n$ the blow-up along the maximal ideals of all singular points of \overline{X}_n . Then X_n is a smooth projective variety and the action of $G := U(n, \mathbb{Z})$ on E_{ω}^n descends to a faithful *biholomorphic* action on X_n . As the *G*-action on E_{ω}^n has zero entropy, so do the *G*-action on X_n . We verify that

$$c_{\rm ess}(G, X_n) = n - 1.$$

Furthermore, if $n \in \{2, 3, 4, 5\}$, then X_n is rationally connected (see *e.g.* [22, Proof of Corollary 4.6] and also [23]), and hence $\kappa(X_n) = -\infty$.

In view of the above examples we ask the following.

Question 4.3. Let $n \ge 6$. Can we construct an *n*-dimensional smooth projective variety X of Kodaira dimension $-\infty$ (e.g. a rational, or rationally connected variety) which admits a zero entropy subgroup G such that $c_{ess}(G, X) = n - 1$?

References

- S. Bochner and D. Montgomery, Locally compact groups of differentiable transformations, Ann. of Math. (2) 47 (1946), 639–653. 3
- [2] M. Brion, On automorphism groups of fiber bundles, Publ. Mat. Urug. 12 (2011), 39-66. 20
- [3] S. Cantat, Automorphisms and dynamics: a list of open problems, In: Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures, pages 619–634. World Sci. Publ., Hackensack, NJ, 2018. 1, 2

- [4] S. Cantat and J. Xie, Algebraic actions of discrete groups: the p-adic method, Acta Math., 220(2):239–295, 2018.
- [5] T.-C. Dinh, Suites d'applications méromorphes multivaluées et courants laminaires, J. Geom. Anal. 15(2) (2005), 207-227. 14
- [6] T.-C. Dinh, Tits alternative for automorphism groups of compact Kähler manifolds, Acta Math. Vietnamatica, Volume 37 (special volume VIASM), Number 4 (2012), 513–529. 1, 3
- [7] T.-C. Dinh, Pluripotential theory and complex dynamics in higher dimension, In: Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited lectures, pages 1561–1581. World Sci. Publ., Hackensack, NJ, 2018. 1
- [8] T.-C. Dinh, F. Hu and D.-Q. Zhang, Compact Kähler manifolds admitting large solvable groups of automorphisms, Adv. Math. 281 (2015), 333–352. 8
- [9] T.-C. Dinh and V.-A. Nguyen, The mixed Hodge-Riemann bilinear relations for compact Kähler manifolds, *Geom. Funct. Anal.* 16 (2006), no. 4, 838–849. 3, 11
- [10] T.-C. Dinh, V.-A. Nguyen and T.-T. Truong, On the dynamical degrees of meromorphic maps preserving a fibration, *Comm. Contemporary. Math.* 14, No. 6 (2012), 18 pp. 20
- [11] D. B. A. Epstein and W. P. Thurston, Transformation groups and natural bundles, Proc. London Math. Soc. (3), 38(2):219–236, 1979. 5
- [12] M. Gromov, Convex sets and Kähler manifolds, Advances in differential geometry and topology, 1–38, World Sci. Publ., Teaneck, NJ, 1990. 3, 11
- [13] M. Gromov, On the entropy of holomorphic maps, Enseign. Math. 49 (2003), 217–235, Manuscript (1977). 1
- [14] R. A. Horn and C.R. Johnson, Topics in Matrix Analysis, Cambridge University Press, 1991. 19
- [15] J. E. Humphreys, Linear Algebraic Groups, Graduate Texts in Mathematics, Volume 21, Springer, 1975. 2, 6, 7
- [16] J. Keum, K. Oguiso and D.-Q. Zhang, Conjecture of Tits type for complex varieties and theorem of Lie-Kolchin type for a cone, *Math. Res. Lett.* 16 (2009), no. 1, 133–148. 3, 10
- [17] H.-Y. Lin, K. Oguiso and D.-Q. Zhang, Polynomial log-volume growth in slow dynamics and the GK-dimensions of twisted homogeneous coordinate rings, arXiv:2104.03423. 4
- [18] F. Lo Bianco, On the cohomological action of automorphisms of compact Kähler threefolds, Bull. Soc. Math. France 147 (2019), no. 3, 469-514. 2
- [19] E. Looijenga, Discrete automorphism groups of convex cones of finite type, Compos. Math. 147 (2014), no. 11, 1939-1962. 8
- [20] K. Oguiso, Tits alternative in hypekärhler manifolds, Math. Res. Lett. 13 (2006), 307–316. 7
- [21] K. Oguiso. Some aspects of explicit birational geometry inspired by complex dynamics. In Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. II, pages 695–721. Kyung Moon Sa, Seoul, 2014. 1
- [22] K. Oguiso, Pisot units, Salem numbers and higher dimensional projective manifolds with primitive automorphisms of positive entropy, *International Mathematics Research Notices*, (2019) no. 5, 1373– 1400. 21
- [23] K. Oguiso and T.-T. Truong, Explicit examples of rational and Calabi-Yau threefolds with primitive automorphisms of positive entropy, *Kodaira Centennial issue of the Journal of Mathematical Sciences*, the University of Tokyo 22 (2015), 361–385. 21
- [24] D. Robinson, A course in the theory of groups, Second edition. Graduate Texts in Mathematics, 80. Springer-Verlag, New York, 1996. xviii+499 pp. 4
- [25] D. Segal, Polycyclic groups, volume 82 of Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 1983. 6
- [26] K. Ueno, Classification theory of algebraic varieties and compact complex spaces, Lecture Notes in Mathematics 439, Springer, 1975. 20
- [27] C. Voisin, Hodge theory and complex algebraic geometry I, Cambridge Studies in Advanced Mathematics 76, Cambridge University Press, 2007. 8
- [28] Y. Yomdin, Volume growth and entropy, Isr. J. Math. 57 (1987), 285-300. 1
- [29] D.-Q. Zhang, A theorem of Tits type for compact Kähler manifolds, Invent. Math. 176 (2009), no. 3, 449-459. 3

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE *Email address*: matdtc@nus.edu.sg

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN *Email address*: hsuehyunglin@ntu.edu.tw

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, JAPAN; NATIONAL CENTER FOR THEORETICAL SCIENCES, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN

Email address: oguiso@g.ecc.u-tokyo.ac.jp

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE *Email address:* matzdq@nus.edu.sg