

# Testing the standard model of cosmology with the SKA: the cosmic radio dipole

C. A. P. Bengaly<sup>1\*</sup>, T. M. Siewert<sup>2</sup>, D. J. Schwarz<sup>2</sup>, R. Maartens<sup>1,3</sup>

<sup>1</sup>*Department of Physics & Astronomy, University of the Western Cape, Cape Town 7535, South Africa*

<sup>2</sup>*Fakultät für Physik, Universität Bielefeld, Postfach 100131, 33501 Bielefeld, Germany*

<sup>3</sup>*Institute of Cosmology & Gravitation, University of Portsmouth, Portsmouth PO1 3FX, United Kingdom*

13 August 2022

## ABSTRACT

The dipole anisotropy seen in the cosmic microwave background radiation is interpreted as due to our peculiar motion. The Cosmological Principle implies that this cosmic dipole signal should also be present, with the same direction, in the large-scale distribution of matter. Measurement of the cosmic matter dipole constitutes a key test of the standard cosmological model. Current measurements of this dipole are barely above the expected noise and unable to provide a robust test. Upcoming radio continuum surveys with the SKA should be able to detect the dipole at high signal to noise. We simulate number count maps for SKA survey specifications in Phases 1 and 2, including all relevant effects. Nonlinear effects from local large-scale structure contaminate the cosmic (kinematic) dipole signal, and we find that removal of radio sources at low redshift ( $z \lesssim 0.5$ ) leads to significantly improved constraints. We forecast that the SKA could determine the dipole direction in Galactic coordinates with an error of  $\sim (1^\circ, 3^\circ) - (5^\circ, 5^\circ)$ , depending on the sensitivity. The predicted errors on the relative speed are  $\sim 6 - 10\%$ . These measurements would significantly reduce the present uncertainty on the direction of the radio dipole, and thus enable the first critical test of consistency between the matter and CMB dipoles.

**Key words:** Cosmology: observations; Cosmology: theory; (cosmology:) large-scale structure of the Universe;

## 1 INTRODUCTION

The Cosmological Principle underlying the standard model of the Universe requires that the CMB and the matter distribution should be isotropic on large scales, after we remove the cosmic kinematic dipole due to the motion of the Solar System relative to the cosmic rest-frame. In particular, this implies that the matter distribution should have a kinematic dipole with the same direction as that of the CMB. This constitutes a critical test of the foundations of the standard cosmological model (Ellis & Baldwin 1984; Schwarz et al. 2015).

Since the CMB dipole establishes the cosmic rest-frame of the early Universe at last photon scattering and the observed distribution of matter is probing the late Universe (at redshifts of order unity), the prediction that the early and late cosmic rest-frames should coincide is still awaiting accurate observational verification.

The CMB dipole was measured with high accuracy by the *Planck* collaboration (Akrami et al. 2018):

$$v = 369.82 \pm 0.11 \text{ km/s} \\ \text{towards } (264.021 \pm 0.011, 48.253 \pm 0.005)^\circ, \quad (1)$$


where the direction is given in Galactic coordinates.

This dipole is expected to be dominated by the kinematic contribution, which is  $O(10^2)$  larger than the intrinsic fluctuations in the standard  $\Lambda$ CDM model. Since the cosmic variance of the dipole is very large, significant non-kinematic contributions remain possible and need to be tested by other means. Probing the dipole of the matter distribution in addition to that of the CMB will help to tighten constraints on putative non-kinematic contributions.

The extragalactic radio sky offers an excellent opportunity to perform an independent test of the Cosmological Principle. The radio continuum dipole is expected to be dominated by the kinematic dipole. This is not the case for galaxy surveys at visible or infrared wavebands: the number counts in wide area surveys in those wavebands are dominated by objects at redshifts well below unity, so that the large-scale structure dominates over the kinematic signal. By contrast, radio continuum surveys have median redshifts above one, which suppresses the effect of local large-scale structure. Another advantage is that radio waves are not subject to extinction and thus the sky area that can be reliably observed by radio surveys exceeds that of optical and infrared surveys.

The largest available wide-area radio continuum surveys include the NRAO VLA Sky Survey (NVSS) (Condon et al. 1998) and the Tiered GMRT Sky Survey (TGSS) (Intema et al. 2016).

\* E-mail: carlosap87@gmail.com



**Figure 1.** Angular number density (left), clustering bias (centre), and magnification bias (right) as a function of redshift, for flux thresholds 1, 5, 10 and  $20 \mu\text{Jy}$ .

Measurements performed using these and other radio continuum catalogues have found that the radio dipole is compatible with the CMB dipole direction, but the dipole amplitude is 2–5 times larger than the signal observed in the CMB (Blake & Wall 2002; Singal 2011; Gibelyou & Huterer 2012; Rubart & Schwarz 2013; Tiwari et al. 2014; Tiwari & Jain 2015; Tiwari & Nusser 2016; Colin et al. 2017; Bengaly et al. 2018b). However, no significant evidence for anomalous anisotropy in galaxy counts was reported at lower redshift ranges, such as those probed by visible and infrared catalogues (Itoh et al. 2010; Gibelyou & Huterer 2012; Yoon et al. 2014; Alonso et al. 2015a; Yoon & Huterer 2015; Javanmardi & Kroupa 2017; Bengaly et al. 2017, 2018a; Rameez et al. 2018).

Measurement of the continuum radio dipole is one of the High-Priority Science Objectives of the SKA,<sup>1</sup> which can be extracted from the same type of survey that will allow us to measure or constrain primordial non-Gaussianity on the largest angular scales (Maartens et al. 2015). Here we focus on the prospects of measuring the dipole direction and amplitude by means of the SKA.

We produce mock catalogues for SKA continuum surveys that include the effects of shot noise, large-scale structure and the kinematic dipole, and then we forecast the errors on SKA measurements of the radio dipole. This provides the details that underpin the results that we presented in the SKA1 Cosmology Red Book (Bacon et al. 2018). Here we include all the effects from large-scale structure, which were not included in the SKA Science Book. In addition, we extend the Red Book results to an alternative ‘optimistic’ flux threshold for SKA1, as well as to ‘optimistic’ and ‘realistic’ flux thresholds for SKA2. This analysis also updates and extends previous forecasts presented in Schwarz et al. (2015) (see also Crawford 2009; Itoh et al. 2010; Yoon & Huterer 2015).

## 2 ANALYSIS

### 2.1 Survey specifications

Radio continuum surveys measure the integrated source flux in a frequency band, with rms noise  $S_{\text{rms}}$ . For the proposed SKA1 sur-

vey (Bacon et al. 2018):

$$\text{SKA1 : } f_{\text{sky}} \approx 0.5, \quad 350 < \nu < 1050 \text{ MHz}, \quad (2)$$

$$S_{\text{rms}} \sim 1 \mu\text{Jy}, \quad 0 < z \lesssim 5, \quad (3)$$

where  $f_{\text{sky}}$  is the fraction of sky observed,  $\nu$  is the observed radio frequency and  $z$  is the redshift. Specifications for SKA2 have not been formalised, but the rms noise is expected to be  $\sim 10$  times smaller and the sky coverage to increase from  $\sim 20,000$  to  $\sim 30,000 \text{ deg}^2$ :

$$\text{SKA2 : } f_{\text{sky}} \approx 0.7, \quad S_{\text{rms}} \sim 0.1 \mu\text{Jy}, \quad 0 < z \lesssim 6. \quad (4)$$

We assume that SKA2 has the same frequency range as SKA1.

For the flux threshold, we adopt optimistic and realistic values as follows:

$$\text{SKA1 : } \quad \text{optimistic: } S > 10 \mu\text{Jy}, \quad \text{realistic: } S > 20 \mu\text{Jy}, \quad (5)$$

$$\text{SKA2 : } \quad \text{optimistic: } S > 1 \mu\text{Jy}, \quad \text{realistic: } S > 5 \mu\text{Jy}. \quad (6)$$

In the SKA1 Red Book,  $S > 22.4 \mu\text{Jy}$  is used (Bacon et al. 2018), which corresponds to the realistic case for SKA1.

### 2.2 Simulated data


In order to forecast the SKA dipole constraints, we generate 500 mock catalogues of radio number count maps. This is done in two steps:

- Compute the theoretical angular power spectrum  $C_\ell$  using CAMB SOURCES (Challinor & Lewis 2011), which includes the effects on the observed number counts of redshift space distortions and lensing magnification.
- Input the theoretical  $C_\ell$  and the redshift distribution of sources  $n(z)$  to the lognormal code FLASK (Xavier et al. 2016), to generate mock SKA number count maps.

Computation of the theoretical  $C_\ell$  requires as inputs the clustering bias  $b(z)$ , the redshift distribution of radio sources  $n(z)$ , and the magnification bias  $s(z)$  (which determines the effect of lensing magnification on number counts). These quantities can be estimated by using the SKA Simulated Skies ( $S^3$ ) data-base (Wilman et al. 2008). We use the code<sup>2</sup> of Alonso et al. (2015b), which produces a semi-analytic fit to the  $S^3$  luminosity function (details about the code are provided in their Appendix B). The results are shown in Fig. 1.

<sup>1</sup> <https://astronomers.skatelescope.org/wp-content/uploads/2016/12/SKA-TEL-SKO-0000122.pdf>

<sup>2</sup> <http://intensitymapping.physics.ox.ac.uk/codes.html>



**Figure 2.** Examples of SKA number count realisations for  $S > 1$  (upper left), 5 (upper right), 10 (lower left), and  $20\mu\text{Jy}$  (lower right).

For the fiducial cosmological model, we use the *Planck* 2015 best-fit flat  $\Lambda\text{CDM}$  parameters (Ade et al. 2016). We compute  $C_\ell$  in 6 redshift bins:

$$0 < z < .5, .5 < z < 1, 1 < z < 2, 2 < z < 3, 3 < z < 4, 4 < z < 5, \quad (7)$$

using a top-hat window function. For each bin, we use in CAMB SOURCES the clustering and magnification biases shown in Fig. 1. The nonlinear matter power spectrum is included using an updated HALOFIT code (Takahashi et al. 2012).

The  $C_\ell$  in the 6 redshift bins, together with the source redshift distribution  $n(z)$ , are separately input into the lognormal code FLASK to generate 6 simulated sky maps. The mock catalogues from these number count maps are given in the HEALPIX (Górski et al. 2015) grid resolution  $N_{\text{side}} = 64$  for each redshift bin.

Then we stack the 6 maps together to obtain an SKA mock number count map for each flux threshold, as one would observe in an actual radio continuum catalogue. In this way, we ensure that not only the shot noise, but also the expected clustering is carefully taken into account, including all theoretical effects (the kinematic dipole is also included, as explained below). Examples of these realisations at each flux threshold are given in Fig. 2.

### 2.3 Fiducial kinematic dipole

Following Bengaly et al. (2018b), the effect of the kinematic dipole is input in the mock maps through a dipole modulation of the number counts. To first order in perturbations, a boost of the observer leads to

$$N_{\text{obs}}(\mathbf{n}, > S) = N_{\text{rest}}(\mathbf{n}, > S) \left[ 1 + A \mathbf{n} \cdot \hat{\beta} \right], \quad (8)$$

where  $N$  is the source count per solid angle about the unit direction  $\mathbf{n}$  and above a flux threshold  $S$ , and  $v = \beta c$  is the velocity of the

Solar System observer relative to the CMB rest-frame. The boost amplitude is (Ellis & Baldwin 1984):

$$A = [2 + x(1 + \alpha)]\beta. \quad (9)$$

The number count contrast is defined by

$$\delta(\mathbf{n}, > S) = \frac{N(\mathbf{n}, > S) - \bar{N}(> S)}{\bar{N}(> S)}, \quad (10)$$

where  $\bar{N}$  is the average number of sources per solid angle. Then (8) gives

$$\delta_{\text{obs}}(\mathbf{n}, > S) = \delta_{\text{rest}}(\mathbf{n}, > S) + A \cos \theta, \quad \cos \theta = \mathbf{n} \cdot \hat{\beta}. \quad (11)$$

The expression (9) assumes that the flux density and counts are given by power laws:

$$S \propto \nu^{-\alpha}, \quad \bar{N}(> S) \propto S^{-x}. \quad (12)$$


The fiducial radio kinematic dipole amplitude is computed from (9) using the SKA1 Red Book values (Bacon et al. 2018),

$$x = 1, \quad \alpha = 0.76, \quad (13)$$


for all flux thresholds, while  $\beta$  and  $\hat{\beta}$  are taken from the CMB dipole values in (1). Then the fiducial dipole amplitude is

$$A_{\text{kin}} = 4.62 \times 10^{-3}. \quad (14)$$

At low redshifts, the kinematic dipole is dominated by the dipole induced by the nonlinear influence of local large-scale structure. The local structure dipole is a contaminant to the cosmic kinematic signal, whose effects decay with redshift. We found that an effective way to suppress this contamination is to excise all sources in the first redshift bin  $z < 0.5$ . The consequent increase in shot noise is not significant: unlike current surveys, SKA surveys are



**Figure 3.** Dipole directions from (15) for the flux limits indicated, based on 500 simulations each, in Galactic coordinates and stereographic projection. Dots show the CMB dipole (white), and the kinematic dipole with (red) and without (blue) local structure.



**Figure 4.** Dipole amplitude corresponding to Fig. 3. *Left:* Histogram for 4 flux thresholds, including all sources ( $z > 0$ ). The fiducial value (14) is the black line. *Right:*  $S > 20 \mu\text{Jy}$  case, with and without  $z < 0.5$  sources.

not dominated by shot noise until  $\ell > 500$  (see Fig. 2 in Pant et al. 2018). The removal of low- $z$  sources should be feasible by cross-correlating radio continuum data with optical, infra-red or future 21cm data, as discussed in the SKA1 Red Book (Bacon et al. 2018).

## 2.4 Estimator

Linear estimators of the dipole suffer from bias, as discussed in Rubart & Schwarz (2013). Quadratic estimators that are free from this problem can be constructed. A straightforward choice of such an estimator for the dipole in pixel space is to vary the monopole and dipole of number counts so as to find

$$\min \sum_p \frac{[N_p(\mathbf{n}, > S) - \bar{N}(> S) (1 + A \cos \theta_p)]^2}{\bar{N}(> S) (1 + A \cos \theta_p)}. \quad (15)$$

The sum is taken over all unmasked pixels and the monopole  $\bar{N}(> S)$  is an estimate of the all-sky average.

Comparing (15) with (8), we have approximated  $N_{\text{rest}}(\mathbf{n}, > S)$  by  $\bar{N}(> S)$ : in other words, the expression (15) neglects clustering and effectively assumes a Poisson distribution. In principle one could model deviations from such a Poisson distribution due to

large-scale structure by including the full covariance matrix, using the theoretically expected angular two-point correlation function.<sup>3</sup> We leave the inclusion of clustering for a future study.

Instead, here we suppress the effect of large-scale structure on the estimator by using rather large pixels and downgrading each simulated map to  $N_{\text{side}} = 16$ . Then we minimise the estimator for each simulation on a three-dimensional grid of dimension  $49152 \times 6 \times 20$ . The dipole direction is probed along the pixel centres defined for  $N_{\text{side}} = 64$ . The mean distance between neighbouring pixel centres at this resolution is  $0.92^\circ$ . Using the HEALPix ‘ring’ scheme, we effectively reduce the two angular dimensions to a one-dimensional index that probes the full sphere. The other two grid directions are the monopole and dipole amplitudes, which vary for the dipole amplitude in steps of  $5 \times 10^{-4}$ .

<sup>3</sup> Note that we do not assign any physical meaning to the minimum of the estimator in this work, i.e., we are not calculating p-values.

**Table 1.** Averaged dipole direction (Galactic coordinates) and amplitude for 500 simulations at 4 flux thresholds, with and without  $z < 0.5$  sources.

Sample	$S >$ ( $\mu\text{Jy}$ )	$l$ (deg)	$b$ (deg)	$A$ ( $10^{-3}$ )
full	1.0	$263.89 \pm 3.06$	$47.19 \pm 6.57$	$4.67 \pm 0.49$
$z \geq 0.5$	1.0	$264.62 \pm 1.2$	$47.45 \pm 3.53$	$4.65 \pm 0.30$
full	5.0	$263.29 \pm 6.71$	$46.99 \pm 7.65$	$4.70 \pm 0.63$
$z \geq 0.5$	5.0	$264.67 \pm 2.09$	$47.39 \pm 3.57$	$4.72 \pm 0.36$
full	10.0	$263.42 \pm 5.93$	$46.89 \pm 7.52$	$4.79 \pm 0.63$
$z \geq 0.5$	10.0	$264.71 \pm 2.05$	$47.22 \pm 3.87$	$4.75 \pm 0.40$
full	20.0	$260.11 \pm 20.33$	$44.63 \pm 15.66$	$5.20 \pm 1.23$
$z \geq 0.5$	20.0	$264.89 \pm 5.54$	$46.65 \pm 5.07$	$4.79 \pm 0.46$
fiducial		264.02	48.25	4.62

### 3 RESULTS

The results of 500 simulations for each flux threshold are shown in Fig. 3 for the dipole direction and Fig. 4 for the dipole amplitude. The numerical values are summarised in Table 1.

The central values and spread shown are calculated via the arithmetic mean and standard deviation from the simulations for each sample, which are straightforward for the latitude  $b$  and amplitude  $A$ . For the longitude  $l$ , we need to account for the fact that close to the pole, a small shift can lead to a large difference in  $l$ , whereas close to the equator  $l$  is a good measure. We therefore use a weighted mean and standard deviation, defined as follows:

$$\bar{l} = \frac{1}{k} \sum_i w_i l_i, \quad w_i = \frac{k \cos b_i}{\sum_i \cos b_i}, \quad (16)$$

$$\sigma_l^2 = \frac{1}{k-1} \sum_i w_i (l_i - \bar{l})^2. \quad (17)$$

Here the sums are over  $i = 1, \dots, k$ , where  $k (= 500)$  is the number of realisations.

The dipole amplitudes are consistent among the four flux thresholds, but increase slightly with increasing flux limits. Our results recover the assumption of a kinematic radio dipole with expected amplitude given by (14). We find the best agreement for the lowest flux threshold and without local structure:

$$A = (4.65 \pm 0.30) \times 10^{-3}, \quad \text{for } S > 1 \mu\text{Jy} \text{ and } z \geq 0.5. \quad (18)$$

The contribution from local structure increases with the flux limit, becoming significant for the highest flux limit.

The recovered dipole directions are also in agreement with the CMB dipole position, given in (1), for all flux thresholds. For the lowest flux sample, and without local structure, we measure the dipole direction with degree-level accuracy:

$$(l, b) = (264.62 \pm 1.2, 47.45 \pm 3.53)^\circ, \quad (19)$$

for  $S > 1 \mu\text{Jy}$  and  $z \geq 0.5$ .

When we remove local structure, the error on the dipole position drops significantly and the directions become more consistent with the CMB dipole direction. As is evident in Fig. 3, the spread of the individual dipole directions is larger for higher flux limits. The simulations without local structure (blue) are still concentrated around the CMB dipole direction (white), while the full simulations (red) begin to spread over the northern galactic hemisphere.

In order to have a comparison with the estimator (15), we present forecasts using an alternative estimator in Appendix A.

### 4 CONCLUSIONS

The standard model of cosmology is highly successful in accounting for current observations of the CMB and large-scale structure. Nevertheless, it remains of crucial importance to also test the theoretical consistency of the model, using current and upcoming observations. A fundamental test is to probe the consistency between the kinematic dipoles measured in the CMB and in the large-scale structure. A mis-match between these dipoles could indicate either a violation of the Cosmological Principle on the large scales where it is required to hold, or a sign of new physical features in the primordial Universe.

The CMB kinematic dipole has been measured with exquisite accuracy by *Planck*. Measurement of this dipole in galaxy surveys faces formidable problems. The survey needs to cover a significant fraction of the sky with low shot noise, in order to make a detection of the dipole. In addition, the survey must reach redshifts significantly above  $z \sim 1$ , in order to suppress the strong contamination of the kinematic dipole by local structure.

These requirements are in principle met by wide radio continuum surveys. The existing all-sky NVSS and TGSS are unable to deliver the signal-to-noise needed for a robust measurement of the kinematic dipole direction and amplitude. We have investigated the capacity of next-generation SKA surveys to make a measurement that allows for a consistency test. By simulating SKA number count maps that include clustering, the fiducial kinematic dipole and shot noise, we have shown that SKA1 should be able to make a measurement with  $\sim 8 - 10\%$  accuracy on the amplitude and  $\sim (2^\circ, 4^\circ) - (5^\circ, 5^\circ)$  accuracy on the direction. Although this is well behind *Planck* precision, checking the consistency of the two directions within degree level would be a remarkable test. SKA2 delivers increased precision, but not by a significant margin: in other words, this critical consistency will already be feasible in Phase 1 and does not have to wait for Phase 2.

A critical aspect of our constraints is the significant improvement that we achieved by excising sources at  $z < 0.5$  in order to suppress the contamination from local structure. Cross-correlating the SKA map with current and upcoming optical/infrared and low- $z$  21cm data should allow for this excision to be implemented effectively.

We did not model any systematic instrumental effects here; some of them are addressed in the SKA1 Science Red Book and are shown to not be show-stoppers. A future study should test if there is any unexpected cross-talk between possible systematics and large-scale structure that could undermine accurate measurement of the kinematic dipole.

**Acknowledgements**

We thank Song Chen for comments and Mario Santos for very useful discussions. CB and RM acknowledge support from the South African SKA Project and the National Research Foundation of South Africa (Grant No. 75415). RM was also supported by the UK Science & Technology Facilities Council (Grant No. ST/N000668/1). TS and DJS gratefully acknowledge support from the Deutsche Forschungsgemeinschaft (DFG) within the Research Training Group 1620 “Models of Gravity”.

**Table A1.** As in Table 1, but using the alternative estimator (A1).

Sample	$S >$ ( $\mu\text{Jy}$ )	$l$ (deg)	$b$ (deg)	$A$ ( $10^{-3}$ )
full	1.0	$262.97 \pm 19.92$	$48.14 \pm 10.76$	$5.64 \pm 0.36$
$z \geq 0.5$	1.0	$261.56 \pm 13.69$	$48.14 \pm 7.72$	$5.46 \pm 0.20$
full	5.0	$259.50 \pm 32.34$	$48.14 \pm 14.84$	$5.72 \pm 0.43$
$z \geq 0.5$	5.0	$258.75 \pm 22.11$	$51.26 \pm 10.22$	$5.50 \pm 0.23$
full	10.0	$258.75 \pm 32.86$	$48.14 \pm 15.69$	$5.76 \pm 0.41$
$z \geq 0.5$	10.0	$258.33 \pm 24.65$	$49.70 \pm 10.35$	$5.56 \pm 0.26$
full	20.0	$258.67 \pm 48.42$	$43.41 \pm 20.27$	$6.38 \pm 0.82$
$z \geq 0.5$	20.0	$259.84 \pm 28.14$	$49.70 \pm 12.33$	$5.74 \pm 0.32$

**Table A2.** Clustering and shot noise dipole amplitudes, and SNR (A3).

Sample	$S >$ ( $\mu\text{Jy}$ )	$A_{\text{LSS}}$ ( $10^{-3}$ )	$A_{\text{PN}}$ ( $10^{-3}$ )	SNR
full	1.0	1.389	0.107	3.446
$z \geq 0.5$	1.0	1.005	0.120	4.748
full	5.0	1.949	0.237	2.397
$z \geq 0.5$	5.0	1.294	0.245	3.574
full	10.0	2.052	0.314	2.274
$z \geq 0.5$	10.0	1.415	0.349	3.238
full	20.0	3.270	0.448	1.436
$z \geq 0.5$	20.0	1.723	0.482	2.641

**APPENDIX A: ALTERNATIVE ESTIMATOR**

For comparison, we evaluate the constraints using an alternative hemispherical comparison estimator, following Bengaly et al. (2018b) (see also Bengaly et al. 2017, 2018a):

$$\Delta(\theta) \equiv \frac{\sigma_i^U(\theta) - \sigma_i^D(\theta)}{\sigma} = A \cos \theta, \quad (\text{A1})$$

where

$$\sigma_i^J = \frac{N_i^J}{2\pi(f_{\text{sky}})_i^J}, \quad \sigma = \frac{N_{\text{total}}}{4\pi f_{\text{sky}}}. \quad (\text{A2})$$

Here  $i = 1, \dots, 12288$  labels the hemisphere decomposition, corresponding to  $N_{\text{side}} = 32$  grid resolution<sup>4</sup>, and  $J = U, D$  identifies the ‘up’ and ‘down’ hemispheres in this pixelisation scheme.

For each hemisphere,  $N_i^J$  is the number of sources,  $(f_{\text{sky}})_i^J$  is the observed sky fraction, and  $\sigma_i^J$  is the source density, with  $N_i^U + N_i^D = N_{\text{total}}$ .  $\theta$  is the angle between the  $i$ -pixel centre and the observer’s motion, so that the maximum  $\Delta(\theta)$  value provides the kinematic dipole amplitude, and the pixel centre position where it occurs is regarded as the dipole direction.

The results are shown in Fig. A1 and Table A1. It is apparent that the recovered dipole amplitudes are larger than the predicted value (14), and significantly larger than the values recovered via the quadratic estimator (15). This shows that the linear estimator (A1) is biased (see Rubart & Schwarz 2013 for a discussion of different linear estimators that show similar effects).

Despite the bias, this estimator is clearly detecting the signal and recovers the fiducial direction, however with a larger spread of values, as is clear from Fig. A1. The advantage of linear estimators is that they are less compute-intensive and easy to implement. We thus can use them to obtain fast order-of-magnitude estimates of the individual contributions to the cosmic radio dipole.

<sup>4</sup> Note that  $N_{\text{side}} = 16$  was chosen in Bengaly et al. (2017, 2018a,b).




Figure A1. Same as Fig. 3, but using the alternative estimator (A1).

### Signal to Noise estimate

We can make an estimate of the signal-to-noise ratio (SNR) for detection of the kinematic dipole, using the expression in Bengaly et al. (2018b) (see also Itoh et al. 2010; Baleisis et al. 1998):

$$\text{SNR} = \frac{A_{\text{kin}}}{\sqrt{A_{\text{LSS}}^2 + A_{\text{PN}}^2}}. \quad (\text{A3})$$

Here  $A_{\text{kin}}$  is given by (14),  $A_{\text{LSS}}$  is the dipole contribution from large-scale structure, and  $A_{\text{PN}}$  is the dipole contribution from Poisson noise.

The last two quantities are estimated using (A1) on 500 catalogues, constructed as follows:

- $A_{\text{LSS}}$  : LSS-only simulations, no PN,
- $A_{\text{PN}}$  : PN-only simulations, no LSS,

where the PN-only maps are homogeneous (no clustering). The median values of the dipole amplitudes are regarded as  $A_{\text{LSS}}$  and  $A_{\text{PN}}$ , respectively. We perform this procedure for the full sample, as well as the sub-sample with  $z < 0.5$  sources excised. This gives a rough estimate of how well the kinematic signal can be detected with the SKA survey specifications.

The results are presented in Table A2. It is evident that  $A_{\text{LSS}}$  is almost half the expected kinematic dipole amplitude (14) for the highest flux limits, but this number significantly decreases when the local sources are removed. Although the shot noise contribution  $A_{\text{PN}}$  slightly increases after removal of  $z < 0.5$  sources, there is still a significant gain in SNR.

Note that the SKA SNR is much larger than that of the existing surveys NVSS and TGSS, for which  $\text{SNR} \approx 1$  (Bengaly et al. 2018b).

### REFERENCES

- P. A. R. Ade, *et al.* [Planck Collaboration], “Planck 2015 results. XIII. Cosmological parameters,” *Astron. Astrophys.* **594**, A13 (2016) [arXiv:1502.01589].
- Y. Akrami *et al.* [Planck Collaboration], “Planck 2018 results. I. Overview and the cosmological legacy of Planck,” arXiv:1807.06205.
- D. Alonso, A. I. Salvador, F. J. Sánchez, M. Bilicki, J. García-Bellido and E. Sánchez, *Mon. Not. Roy. Astron. Soc.* **449** (2015) no.1, 670 [arXiv:1412.5151].
- D. Alonso, P. Bull, P. G. Ferreira, R. Maartens and M. Santos, “Ultra large-scale cosmology in next-generation experiments with single tracers,” *Astrophys. J.* **814**, no. 2, 145 (2015) [arXiv:1505.07596].
- D. J. Bacon, *et al.* [SKA Cosmology Working Group], “Cosmology with SKA1 – 2018 Red Book,” to appear.
- A. Baleisis, O. Lahav, A. J. Loan and J. V. Wall, “Searching for large scale structure in deep radio surveys,” *Mon. Not. Roy. Astron. Soc.* **297**, 545 (1998) [astro-ph/9709205].
- C. A. P. Bengaly, A. Bernui, J. S. Alcaniz, H. S. Xavier and C. P. Novaes, “Is there evidence for anomalous dipole anisotropy in the large-scale structure?,” *Mon. Not. Roy. Astron. Soc.* **464**, no. 1, 768 (2017) [arXiv:1606.06751].
- C. A. P. Bengaly, C. P. Novaes, H. S. Xavier, M. Bilicki, A. Bernui and J. S. Alcaniz, “The dipole anisotropy of WISE x SuperCOSMOS number counts,” *Mon. Not. Roy. Astron. Soc.* **475**, no. 1, L106 (2018) [arXiv:1707.08091].
- C. A. P. Bengaly, R. Maartens and M. G. Santos, “Probing the Cosmological Principle in the counts of radio galaxies at different frequencies,” *JCAP* **1804**, no. 04, 031 (2018) [arXiv:1710.08804].
- C. Blake and J. Wall, “Detection of the velocity dipole in the radio galaxies of the NRAO VLA Sky Survey,” *Nature* **416**, 150 (2002) [astro-ph/0203385].
- A. Challinor and A. Lewis, “The linear power spectrum of observed source number counts,” *Phys. Rev. D* **84**, 043516 (2011) [arXiv:1105.5292].
- J. Colin, R. Mohayaee, M. Rameez and S. Sarkar, “High redshift radio galaxies and divergence from the CMB dipole,” *Mon. Not. Roy. Astron. Soc.* **471**, no. 1, 1045 (2017) [arXiv:1703.09376].
- J. J. Condon, W. D. Cotton, E. W. Greisen, Q. F. Yin, R. A. Perley, G. B. Taylor and J. J. Broderick, “The NRAO VLA Sky survey,” *Astron. J.* **115**, 1693 (1998).
- F. Crawford, “Detecting the Cosmic Dipole Anisotropy in Large-Scale Ra-

- dio Surveys,” *Astrophys. J.* **692**, 887 (2009) [arXiv:0810.4520].
- G. F. R. Ellis and J. E. Baldwin, “On the expected anisotropy of radio source counts,” *Mon. Not. Roy. Astron. Soc.* **206**, 377 (1984)
- S. Ghosh, P. Jain, G. Kashyap, R. Kothari, S. Nadkarni-Ghosh and P. Tiwari, “Probing statistical isotropy of cosmological radio sources using SKA,” *J. Astrophys. Astron.* **37**, 25 (2016) [arXiv:1610.08176].
- C. Gibelyou and D. Huterer, “Dipoles in the Sky,” *Mon. Not. Roy. Astron. Soc.* **427**, 1994 (2012) [arXiv:1205.6476].
- K. M. Gorski, E. Hivon, A. J. Banday, B. D. Wandelt, F. K. Hansen, M. Reinecke and M. Bartelman, “HEALPix - A Framework for high resolution discretization, and fast analysis of data distributed on the sphere,” *Astrophys. J.* **622**, 759 (2005) [astro-ph/0409513].
- H. T. Intema, P. Jagannathan, K. P. Mooley and D. A. Frail, “The GMRT 150 MHz All-sky Radio Survey: First Alternative Data Release TGSS ADR1,” *Astron. Astrophys.* **598**, A78 (2017) [arXiv:1603.04368].
- Y. Itoh, K. Yahata and M. Takada, “A dipole anisotropy of galaxy distribution: Does the CMB rest-frame exist in the local universe?,” *Phys. Rev. D* **82**, 043530 (2010) [arXiv:0912.1460].
- B. Javanmardi and P. Kroupa, “Anisotropy in the all-sky distribution of galaxy morphological types,” *Astron. Astrophys.* **597**, A120 (2017) [arXiv:1609.06719].
- R. Maartens *et al.* [SKA Cosmology SWG Collaboration], “Overview of Cosmology with the SKA,” *PoS AASKA* **14**, 016 (2015) [arXiv:1501.04076]
- N. Pant, A. Rotti, C. A. P. Bengaly and R. Maartens, “Measuring our velocity from fluctuations in number counts,” arXiv:1808.09743.
- M. Rameez, R. Mohayaee, S. Sarkar and J. Colin, “The dipole anisotropy of ALLWISE galaxies,” *Mon. Not. Roy. Astron. Soc.* **477**, 1772 (2018) [arXiv:1712.03444].
- M. Rubart and D. J. Schwarz, “Cosmic radio dipole from NVSS and WENSS,” *Astron. Astrophys.* **555**, A117 (2013) [arXiv:1301.5559].
- D. J. Schwarz *et al.*, “Testing foundations of modern cosmology with SKA all-sky surveys,” *PoS AASKA* **14**, 032 (2015) [arXiv:1501.03820].
- A. K. Singal, “Large peculiar motion of the solar system from the dipole anisotropy in sky brightness due to distant radio sources,” *Astrophys. J.* **742**, L23 (2011) [arXiv:1110.6260].
- R. Takahashi, M. Sato, T. Nishimichi, A. Taruya and M. Oguri, “Revising the Halofit Model for the Nonlinear Matter Power Spectrum,” *Astrophys. J.* **761**, 152 (2012) [arXiv:1208.2701].
- P. Tiwari, R. Kothari, A. Naskar, S. Nadkarni-Ghosh and P. Jain, “Dipole anisotropy in sky brightness and source count distribution in radio NVSS data,” *Astropart. Phys.* **61**, 1 (2014) [arXiv:1307.1947].
- P. Tiwari and P. Jain, “Dipole Anisotropy in Integrated Linearly Polarized Flux Density in NVSS Data,” *Mon. Not. Roy. Astron. Soc.* **447**, 2658 (2015) [arXiv:1308.3970].
- P. Tiwari and A. Nusser, “Revisiting the NVSS number count dipole,” *JCAP* **1603**, 062 (2016) [arXiv:1509.02532].
- R. J. Wilman *et al.*, “A semi-empirical simulation of the extragalactic radio continuum sky for next generation radio telescopes,” *Mon. Not. Roy. Astron. Soc.* **388**, 1335 (2008) [arXiv:0805.3413].
- H. S. Xavier, F. B. Abdalla and B. Joachimi, “Improving lognormal models for cosmological fields,” *Mon. Not. Roy. Astron. Soc.* **459**, no. 4, 3693 (2016) [arXiv:1602.08503].
- M. Yoon, D. Huterer, C. Gibelyou, A. Kovács and I. Szapudi, “Dipolar modulation in number counts of WISE-2MASS sources,” *Mon. Not. Roy. Astron. Soc.* **445**, L60 (2014) [arXiv:1406.1187].
- M. Yoon and D. Huterer, “Kinematic dipole detection with galaxy surveys: forecasts and requirements,” *Astrophys. J.* **813**, no. 1, L18 (2015) [arXiv:1509.05374].

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.