

Mixed H/He bursts in SAX J1748.9–2021 during the spectral change of its 2015 outburst

Z. Li^{1,2,3}, V. De Falco^{4,2}, M. Falanga^{2,5}, E. Bozzo⁶, L. Kuiper⁷, J. Poutanen^{8,9,10}, A. Cumming¹¹, D. K. Galloway^{12,13}, and S. Zhang¹⁴

¹ Department of Physics, Xiangtan University, Xiangtan, 411105, P. R. China
e-mail: lizhaosheng@xtu.edu.cn

² International Space Science Institute (ISSI), Hallerstrasse 6, 3012 Bern, Switzerland

³ Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics / Laboratory for High-Energy Physics, University of Bern, Switzerland

⁴ Departement Physik, Universität Basel, Klingelbergstrasse 82, 4056 Basel, Switzerland

⁵ International Space Science Institute Beijing, No.1 Nanertiao, Zhongguancun, Haidian District, 100190 Beijing, China

⁶ Department of Astronomy, University of Geneva, chemin d'écogia 16, 1290, Versoix, Switzerland

⁷ SRON, Netherlands Institute for Space Research, Sorbonnelaan 2, 3584 CA, Utrecht, The Netherlands

⁸ Tuorla Observatory, Department of Physics and Astronomy, FI-20014 University of Turku, Finland

⁹ Space Research Institute of the Russian Academy of Sciences, Profsoyuznaya str. 84/32, 117997 Moscow, Russia

¹⁰ Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden

¹¹ Department of Physics and McGill Space Institute, McGill University, 3600 Rue University, Montreal QC, Canada H3A 2T8

¹² School of Physics and Astronomy, Monash University, VIC 3800, Australia

¹³ Monash Centre for Astrophysics, Monash University, VIC 3800, Australia

¹⁴ Laboratory for Particle Astrophysics, Institute of High Energy Physics, Beijing 100049, P. R. China

Accepted for publication in A&A on 12/10/2018

ABSTRACT

SAX J1748.9–2021 is a transiently accreting X-ray millisecond pulsar. It is also known as an X-ray burster source discovered by *Beppo-SAX*. We analysed the persistent emission and type-I X-ray burst properties during its 2015 outburst. The source varied from hard to soft state within half day. We modeled the broad-band spectra of the persistent emission in the (1 – 250) keV energy band for both spectral states using the quasi-simultaneous *INTEGRAL* and *Swift* data. The broad-band spectra are well fitted by an absorbed thermal Comptonization model, *COMPBS*, in a slab geometry. The best-fits for the two states indicate significantly different plasma temperature of 18 and 5 keV and the Thomson optical depth of 3 and 4, respectively. In total, 56 type-I X-ray bursts were observed during the 2015 outburst, of which 26 detected by *INTEGRAL* in the hard state, 25 by *XMM-Newton* in the soft state, and 5 by *Swift* in both states. As the object transitioned from the hard to the soft state, the recurrence time for X-ray bursts decreased from ≈ 2 to ≈ 1 hr. The relation between the recurrence time, Δt_{rec} , and the local mass accretion rate per unit area onto the compact object, \dot{m} , is fitted by a power-law model, and yielded as best fit at $\Delta t_{\text{rec}} \sim \langle \dot{m} \rangle^{-1.02 \pm 0.03}$ using all X-ray bursts. In both cases, the observed recurrence times are consistent with the mixed hydrogen/helium bursts. We also discuss the effects of type-I X-ray bursts prior to the hard to soft transition.

Key words. pulsars: individual SAX J1748.9–2021 – stars: neutron – X-ray: binaries – X-ray: bursts

1. Introduction

Neutron stars (NSs), in low mass X-ray binaries (LMXBs), accrete matter from their companion stars, forming an accretion disk (Frank et al. 2002). The accreted matter on the NS surface can trigger thermonuclear flashes, called type-I X-ray bursts (see e.g., Lewin et al. 1993). The burst spectra are described by a blackbody with peak temperatures reaching $kT_{\text{bb}} \approx 3$ keV, and with a gradual softening due to the cooling of the NS photosphere (see Lewin et al. 1993; Strohmayer & Bildsten 2006; Galloway et al. 2008, for a review). The burst peak luminosity can reach the Eddington limit, $L_{\text{Edd}} \approx 2 \times 10^{38}$ erg s⁻¹, and the total burst energy release is of the order $\sim 10^{39-42}$ erg, depending on the burst type (e.g., Galloway et al. 2008).

During type-I X-ray bursts, the injected soft X-ray photons may affect the NS hot inner accretion flow or the surrounding hot corona. Indeed, a deficit has been observed at 30 – 50 keV

photons in IGR J17473–2721, Aql X-1, 4U 1636–536, and GS 1826–238, implying that the soft X-ray photons, emitted during type-I X-ray bursts, cool down the electrons producing hard X-ray emission (Chen et al. 2012, 2013; Ji et al. 2013, 2014a, 2015). The hard persistent X-ray component is thought to be derived from Comptonization in the hot corona (Chen et al. 2012, 2013). A similar study has also been performed during the hard state of the source 4U 1728–34, however, during the occurrence of type-I X-ray bursts the persistent hard X-ray deficit was not detected by *Rossi X-ray Timing Explorer*, due to the limitation of the response in hard X-ray band (Ji et al. 2014b), but confirmed by *INTEGRAL* (Kajava et al. 2017). The persistent emission level above 40 keV was about a factor of three lower compared with the persistent emission without burst activities (Kajava et al. 2017). Also in this case the interpretation indicates that the soft radiation injection into the accretion disk or corona may induce spectral variability during type-I X-ray bursts.

It is believed that with varying mass accretion rate the spectral changes of NSs in LMXBs follow an atoll or Z-track on an X-ray color-color diagram (CCD) (Hasinger & van der Klis 1989; van der Klis 2006). The accretion rate increases from the hard (island) spectral state, where the soft emission is much reduced and the spectrum is dominated by a power-law-like spectrum up to the energies of 200 keV, to the soft (banana) spectral state, where most of the energy is emitted below ~ 20 keV (e.g., Barret 2001; Gierliński & Done 2002). The spectral change has been studied during type-I X-ray bursts (Falanga et al. 2006), while we study SAX J1748.9–2021, as an example to investigate if the large number of type-I X-ray bursts during its 2015 February – June outburst can trigger a spectral state change, i.e., from a persistent hard to a soft spectral state. In this paper, we report on the *INTEGRAL* and *Swift* observations of SAX J1748.9–2021. The properties of the largest set of X-ray bursts as well as the correlation of all observed type-I X-ray bursts with the source in the soft and hard state are investigated.

1.1. The source SAX J1748.9–2021

The X-ray source SAX J1748.9–2021 was discovered by *BeppoSAX* during an outburst in 1998 (in 't Zand et al. 1998). The source has been identified to be located within the globular cluster NGC 6440 at a distance of ≈ 8.2 kpc (in 't Zand et al. 1999; Valenti et al. 2007, and references therein). The detection of a type-I X-ray burst associated to SAX J1748.9–2021 confirmed the source as a NS hosted in a LMXB system (in't Zand et al. 2001). SAX J1748.9–2021 has also been observed during its quiescent state with *Chandra* that identified its optical counterpart candidate (in't Zand et al. 2001, and references therein). Altamirano et al. (2008) suggested that the companion star might be a main-sequence or a slightly evolved star with a mass ranging between $(0.85 - 1.1) M_{\odot}$. Recently, the companion star has been confirmed by the Hubble Space Telescope, and it turns out to be a main-sequence star with a mass of $(0.70 - 0.83) M_{\odot}$, a radius of $(0.88 \pm 0.02) R_{\odot}$, and a surface temperature of (5250 ± 80) K (Cadelano et al. 2017). This source has been observed in outburst in 1998, 2001, 2005, 2009 – 2010, and 2015 and classified also as an atoll source (in 't Zand et al. 1998; in't Zand et al. 2001; Markwardt & Swank 2005; Suzuki et al. 2009; Patruno et al. 2009; Bozzo et al. 2015; Sanna et al. 2016; Pintore et al. 2016). During both 2001 and 2005 outbursts the source showed intermittent pulsations at 442 Hz on timescales of hundreds of seconds (Gavriil et al. 2007; Altamirano et al. 2008), from which it was possible to infer the orbital period of ≈ 8.76 hr and the magnetic field of $B \gtrsim 1.3 \times 10^8$ G (Altamirano et al. 2008). Therefore, this source has been classified as an accreting millisecond X-ray pulsar (AMXP; see Poutanen 2006; Patruno & Watts 2012, for reviews).

2. Observations and data

2.1. INTEGRAL

The data were obtained with *INTEGRAL* (Winkler et al. 2003) from 2015 February 16 to April 20, i.e., from satellite revolution 1508 to 1532 for a total effective exposure time of ~ 856 ks. These revolutions include a public Target of Opportunity (ToO) observation for ~ 100 ks during the revolution 1511.

We analyzed data from the coded mask imager IBIS/ISGRI (Ubertini et al. 2003; Lebrun et al. 2003), covering the 20 – 250 keV energy band, and from the two JEM-X monitors covering the 4 – 22 keV energy range (Lund et al. 2003). The observa-

tion in revolution 1511 was performed in the hexagonal dithering mode, which permits to keep always the source within both the field-of-view of IBIS/ISGRI and JEM-X. For all other revolutions, we considered all the pointings for which the source was located at a maximum off-set angle with respect to the satellite aim point of $\lesssim 12^{\circ}$ for IBIS/ISGRI and $\lesssim 2^{\circ}5$ for JEM-X in order to minimize calibration uncertainties. The data reduction was performed using the standard OFFLINE SCIENCE ANALYSIS (OSA) version 10.2 distributed by the ISDC (Courvoisier et al. 2003). The algorithms used for the spatial and spectral analysis of ISGRI and JEM-X are described in Goldwurm et al. (2003) and Lund et al. (2003), respectively.

In Fig. 1, we show part of the ISGRI field of view (significance map) centered on the position of SAX J1748.9–2021. The source is clearly detected in the mosaic, and we estimated a detection significance of $\sim 126 \sigma$ in the 20 – 100 keV energy range. The best determined position is $\alpha_{J2000} = 17^{\text{h}}48^{\text{m}}52^{\text{s}}.20$ and $\delta_{J2000} = -20^{\circ}21'32''.62$ with an associated uncertainty of $0'.4$ at 90% confidence level (Gros et al. 2003), consistent with the best determined position (Pooley et al. 2002).




Fig. 1. The IBIS/ISGRI sky image centered on SAX J1748.9–2021 in the 20 – 100 keV range of the ~ 100 ks ToO observation. The size of each pixel is $3'$.

Using the ISGRI data in the energy band 15 – 300 keV, starting from revolution 1508, we performed a timing analysis to search for the pulsations. We applied the same timing techniques as used for the ISGRI data in the past (see e.g., Kuiper et al. 2003; Falanga et al. 2005a, 2007, 2011, 2012; De Falco et al.

2017b,a), adopting the ephemeris reported in Sanna et al. (2016). No pulsed emission has been detected above 15 keV, which is in line with the very low pulsed fraction of $\sim 2\%$ as detected by *XMM-Newton* above 10 keV (Sanna et al. 2016).

We extracted the IBIS/ISGRI and JEM-X light curve of SAX J1748.9–2021 at a time scale of typically $\sim 2-3$ ks, i.e. the duration of a pointing (science window). To search for type-I X-ray bursts, we used the JEM-X data also at a time resolution of 2 seconds, as discussed in Sect. 5. The JEM-X and ISGRI spectra were extracted for the hard and soft state quasi-contemporaneous with *Swift* observations (see Sect. 4).

2.2. *Swift*

Swift/XRT (Burrows et al. 2005) detected SAX J1748.9–2021 in outburst after a 2 ks observation on 2015 February 18 (Bozzo et al. 2015). The XRT monitoring campaign covered the 2015 outburst from February 25 to July 30 comprising a total of 40 pointings and the effective exposure time of ~ 43 ks.

We processed the *Swift*/XRT data by using standard procedures (Burrows et al. 2005) and the calibration files version 20160113. The *Swift*/XRT data were taken both in window-timing (WT) and photon-counting (PC) modes (processed with the XRTPIPELINE v.0.13.2). Filtering and screening criteria were applied by using FTOOLS contained in the HEASOFT software package (v6.19)¹. We extracted source and background light curves and spectra by selecting event grades of 0–2 and 0–12 for the WT and PC mode, respectively. We used the latest spectral redistribution matrices in the HEASARC calibration database. Ancillary response files, accounting for different extraction regions, vignetting and PSF corrections, were generated by using the XRTMARKF task. When required, we corrected PC data for pile-up, and used the XRTLCCORR task to account for this correction in the background-subtracted light curve.

2.3. *XMM-Newton*

SAX J1748.9–2021 was observed with *XMM-Newton* (Jansen et al. 2001) on March 4, 2015 (Obs.ID. 0748391301) continuously for about ~ 100 ks with the first results reported in Pintore et al. (2016). We reanalysed the *XMM-Newton* observation to obtain each burst flux and its fluence. Following the standard procedure, the TIMING data from European Photon Imaging Camera (EPIC)-pn (Strüder et al. 2001) were reduced and the last 5 ks of the data were ignored because of flaring background. The data were filtered with FLAG=0 and PATTERN ≤ 4 . After the extraction of the lightcurve, for each burst, we located its start and end time at which the **count rate** increased and decreased to 10% of its peak value above the persistent intensity level, respectively. The spectrum of each burst was obtained in the range RAWX=[31:43] without the column RAWX=37 to reduce the pile-up effect. The persistent spectrum prior to a burst was regarded as its background. Each burst spectrum was fitted by the model TBABS*BBODYRAD, and the burst bolometric flux was calculated using the relation $F = 1.076 \times 10^{-11} (kT_{bb}/1 \text{ keV})^4 K_{bb} \text{ ergs cm}^{-2} \text{ s}^{-1}$ (Galloway et al. 2008), where kT_{bb} and K_{bb} are the blackbody temperature and normalization, respectively. The *XMM-Newton* data reduction and analysis was similar to the procedure described in Pintore et al. (2016). All reported fluxes are unabsorbed.

3. Outburst light curve

We report in Fig. 2 the outburst profile of SAX J1748.9–2021 from February 16 to May 2, 2015. SAX J1748.9–2021 exhibited significant spectral variation during the outburst, passing from hard to soft state within ≈ 0.5 d around MJD 57079.5. The hard state lasted for the first ≈ 10 d, while the soft state lasted about 50 d, until the source returned to quiescence. Most AMXPs that underwent an outburst for a few weeks to months showed a common outburst profile, i.e., the light curve decays exponentially until it reaches a break, after which the flux drops linearly to the quiescence level without exhibiting strong spectral variation (e.g., Gierliński et al. 2002; Gierliński & Poutanen 2005; Falanga et al. 2005b,a, 2007; Ibragimov & Poutanen 2009; Falanga et al. 2011, 2012; De Falco et al. 2017b,a). However, this source is known to be an atoll source with significant flux variations between hard and soft state and showing intermittent pulsations whose origin is unclear (see e.g., Patruno et al. 2009).

4. Spectral analysis

The spectral analysis was carried out using XSPEC version 12.6 (Arnaud 1996). We studied the broad-band X-ray spectrum divided in the hard and soft state, see Fig. 2. For the hard state we consider the INTEGRAL/JEM-X (3.5–22) keV and ISGRI (20–250) keV ToO data between MJD 57076.52–57077.82, and the quasi-simultaneous *Swift*/XRT/WT (1–10) keV data starting on MJD 57078.43 for 1 ks (obs. ID 00033646003). For the soft state we consider the INTEGRAL/JEM-X (4–22) keV and ISGRI (20–50) keV public data between MJD 57079.83–57084.03 and one simultaneous *Swift*/XRT/WT (1–10) keV data for a total of 2 ks (obs ID. 00033646004). We did not merge together the additional two close-by *Swift* pointings ID. 00033646006 and ID. 00033646005, as the roll angle between the three pointings was different and the statistical quality of the first pointing we used was sufficient to perform an accurate broad-band fit. We verified a posteriori that performing the same broad-band fit with the other two *Swift* pointings would have resulted in similar values of the model parameters (to within the uncertainties). For all spectra the bursts intervals have been removed from the data set. For each instrument, a multiplication factor was included in the fit to take into account the uncertainty in the cross-calibration of the instruments. For all fits the factor was fixed at 1 for the INTEGRAL/ISGRI data. All uncertainties in the spectral parameters are given at a 1σ confidence level for a single parameter.

We fit all the combined XRT/JEM-X/ISGRI average spectra separately for the hard and soft spectral state, respectively, using first a simple and phenomenological cutoff power-law model and afterwards a physical motivated thermal Comptonization model, COMPPS, in the slab geometry (Poutanen & Svensson 1996). The later model has been used previously to fit AMXPs broad-band spectra (e.g., Gierliński et al. 2002; Gierliński & Poutanen 2005; Falanga et al. 2005b,a, 2007; Ibragimov & Poutanen 2009; Falanga et al. 2011, 2012; De Falco et al. 2017b,a). The main parameters are the absorption column density, N_H , the Thomson optical depth, τ_T , across the slab, the electron temperature, kT_e , the temperature, kT_{seed} , of the soft-seed thermal photons (assumed to be injected from the bottom of the slab), the apparent area of the seed photons, A_{seed} , and $\cos \theta$ that we fixed at 0.5, where θ is the inclination angle between the slab normal and the observer line of sight. The best-fit parameters of all models are reported in Table 1. At variance with Pintore et al. (2016) we could not find any evidence of the DISKBB component from the residuals of our best fit comptonization models (see also Fig. 3).

¹ <http://heasarc.gsfc.nasa.gov/docs/software.html>.




Fig. 2. SAX J1748.9–2021 light curve during the 2015 outburst. The vertical dashed lines, indicate the ~ 100 ks *INTEGRAL* ToO time interval. The black, green, and red vertical lines indicate the times of the detected type-I X-ray bursts with *INTEGRAL*, *Swift* and *XMM-Newton*, respectively. In the bottom panel the inset shows the hard ISGRI light curve zoomed in at the spectral change time interval. The count rate drops to zero within 0.5 day. The hard and soft state data points used in the spectral analysis are color coded with light blue and pink, respectively (see Sect. 4).

We note that the column density between the two states changes from $N_{\text{H}} \sim 0.5 \times 10^{22} \text{ cm}^{-2}$ to $N_{\text{H}} \sim 0.8 \times 10^{22} \text{ cm}^{-2}$, as well as the plasma temperature, kT_{e} , and the optical depth, τ_{T} (see Table 1). It is remarkable how this spectral change from a hard to a soft state occurred within a half day (see Fig. 2).

Table 1. Best fit parameters determined for the hard and soft spectral state of SAX J1748.9–2021.

	Hard state		Soft state	
	CUTOFFPL	COMPSS	CUTOFFPL	COMPSS
N_{H} (10^{22} cm^{-2})	0.57 ± 0.03	0.52 ± 0.08	0.90 ± 0.06	0.8 ± 0.1
Γ	1.37 ± 0.03	–	1.1 ± 0.1	–
E_{cut}	40 ± 2	–	6.7 ± 1.4	–
kT_{e} (keV)	–	18.1 ± 0.8	–	5.0 ± 0.5
kT_{seed} (keV)	–	0.39 ± 0.06	–	0.28 ± 0.15
τ_{T}	–	2.9 ± 0.1	–	3.8 ± 0.5
A_{seed}	–	5000^{+1450}_{-1300}	–	10300^{+6000}_{-6000}
F_{bol} ($10^{-9} \text{ cm}^{-2} \text{ s}^{-1}$) ^a	3.1 ± 0.2	3.1 ± 0.2	6.1 ± 0.2	6.1 ± 0.2
$\chi^2_{\text{red}}/\text{dof}$	1.05/516	1.06/515	0.98/579	0.98/578

Notes. ^(a) Unabsorbed flux in the 0.1 – 250 keV and 0.1 – 50 keV energy range for the hard and soft states, respectively. For the CUTOFFPL model, the parameters are the photon index, Γ , the cutoff energy, E_{cut} . For the COMPSS model, the main parameters are the Thomson optical depth, τ_{T} , across the slab, the electron temperature, kT_{e} , the temperature, kT_{seed} , of the soft-seed thermal photons and the apparent area of the seed photons, A_{seed} .

5. The type-I X-ray bursts

SAX J1748.9–2021 is also known to be a burster source, as during its 2015 outburst 56 type-I X-ray bursts have been detected (see Fig. 2), of which 26 found by *INTEGRAL*/JEM-X, 5 by *Swift*, and for completeness we added also 25 bursts observed by *XMM-Newton* and reported by Pintore et al. (2016). These bursts occurred during both the hard and soft states. However, we analyzed in more detail the bursts detected during the continuous *INTEGRAL* ToO and *XMM-Newton* observations, where the bursts were separated by two distinct time intervals of around $\Delta t_{\text{rec,hard}} \sim 2$ hr (hard state, *INTEGRAL* data) and $\Delta t_{\text{rec,soft}} \sim 1$ hr (soft state, *XMM-Newton* data). All bursts lasted on average ≈ 100 s and showed a rise time of $\sim 2 - 3$ s. The burst decay profile could be well fitted with an exponential function and the correspondingly derived e-folding time is $\tau_{\text{fit}} = 23 \pm 3$ s, that is in agreement with the value reported by Pintore et al. (2016), estimated to be 25 ± 1 s (typo in their article, private communication). We note, that all the bursts outside these two observations also showed comparable τ_{fit} values and similar burst profiles.

The averaged flux in the 0.1 – 50 keV and 0.1 – 250 keV energy range of the persistent emission during the soft and hard states, respectively, was $F_{\text{pers,soft}} \approx (5.48 - 6.13) \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$ for *XMM-Newton* bursts and $F_{\text{pers,hard}} \approx (2.88 - 3.35) \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$ when *INTEGRAL*/JEM-X detected the bursts. The corresponding luminosities are $L_{\text{pers,soft}} \approx (4.4 - 4.9) \times 10^{37} \text{ erg}$




Fig. 3. The unfolded broad-band spectrum of SAX J1748.9–2021 showing the hard and soft spectral states. The data points are from the *Swift*/XRT (black crosses), *INTEGRAL*/JEM-X (red crosses), and *INTEGRAL*/ISGRI (green crosses). The best fit is obtained with the COMPPS model, represented by a solid blue line. The lower panels show the residuals between the data and the model. The *Swift*/XRT spectra have been rebinned for a better visualization.




Fig. 4. Recurrence time versus local mass accretion rate per unit area onto the compact object, where the circles and triangles represent the type-I X-ray bursts detected by *XMM-Newton* and *INTEGRAL*/JEM-X, respectively. The dashed red lines represent the best-fit power-law model for all bursts $\Delta t_{\text{rec}} \sim \langle \dot{m} \rangle^{-1.02 \pm 0.03}$. The recurrence time decreased from ~ 2 hr to ~ 1 hr as the persistent spectrum showed the transition from the hard to the soft state.

s^{-1} , i.e. (12–13)% L_{Edd} , and $L_{\text{pers,hard}} \approx (2.32 - 2.70) \times 10^{37} \text{ erg s}^{-1}$, i.e. (6–7)% L_{Edd} , where $L_{\text{Edd}} = 3.8 \times 10^{38} \text{ erg cm}^{-2} \text{ s}^{-1}$ is the Eddington luminosity adopted by Kuulkers et al. (2003). We have used as distance to the source $d = 8.2$ kpc.

The local accretion rate per unit area onto the compact object is $\dot{m} = L_{\text{pers}}(1+z)(4\pi R^2(GM/R))^{-1}$, i.e., $\dot{m}_{\text{soft}} \approx 2.67 \times 10^4 \text{ g cm}^{-2} \text{ s}^{-1}$ and $\dot{m}_{\text{hard}} \approx (1.30 - 1.51) \times 10^4 \text{ g cm}^{-2} \text{ s}^{-1}$, where the gravitational redshift is $1+z = 1.31$ for a canonical NS with a mass $M = 1.4M_{\odot}$ and a radius of $R = 10$ km. The observed recurrence time increases with \dot{m} as $\Delta t_{\text{rec}} \sim \langle \dot{m} \rangle^{-1.02 \pm 0.03}$ (see Fig. 4).

For the bursts in the soft state, the total burst fluence is $f_{\text{b,soft}} \approx 0.35 \times 10^{-6} \text{ erg cm}^{-2}$, obtained from the burst flux multiplied by the corresponding burst duration. For the bursts in the hard state, the peak flux is $F_{\text{peak,hard}} \approx 1.6 \times 10^{-8} \text{ erg cm}^{-2} \text{ s}^{-1}$, that multiplied by $\tau_{\text{fit}} \approx 23$ s give an estimation of the total burst fluence of $f_{\text{b,hard}} \approx 0.37 \times 10^{-6} \text{ erg cm}^{-2}$. We note, that these values are close to the directly measured total fluence per burst.

We can then estimate the ignition depth at the onset of the burst with the equation $y_{\text{ign}} = 4\pi f_{\text{b}} d^2 (1+z) (4\pi R^2 Q_{\text{nuc}})^{-1}$, where the nuclear energy generated for solar composition (assuming $\bar{X} = 0.7$) is $Q_{\text{nuc}} \approx 1.31 + 6.95\bar{X} - 1.92\bar{X}^2 \text{ MeV/nucleon} \approx 4.98 \text{ MeV/nucleon}$ (Goodwin et al. submitted). Note that the considered values Q_{nuc} include losses owing to neutrino emission as determined by using KEPLER. Therefore we obtained $y_{\text{ign,soft}} \approx 0.59 \times 10^8 \text{ g cm}^{-2}$ and $y_{\text{ign,hard}} \approx 0.63 \times 10^8 \text{ g cm}^{-2}$ for the soft and hard state, respectively.

Once the ignition depth is known, the recurrence time between the bursts can be calculated by using the equation $\Delta t_{\text{rec}} = (y_{\text{ign}}/\dot{m})(1+z)$. For the soft state, we obtain $\Delta t_{\text{rec,soft}} \sim 0.81$ hr, and for the hard state we obtain $\Delta t_{\text{rec,hard}} \sim 1.6$ hr in the mixed hydrogen/helium burning case. For the pure helium burning, the predicted recurrence time is much longer. Compared with the observed recurrence time and \dot{m} , these results confirm that the burst type is mixed hydrogen/helium with solar abundances for both the hard and soft state (see Fig. 4). In addition this can be additionally validated by the α value (see e.g., Strohmayer & Bildsten 2006). We computed the ratio of the integrated persistent flux to the burst fluence, $\alpha = \tau_{\text{rec}} F_{\text{per}}/f_{\text{b}}$, which are 64 ± 10 for the soft bursts and 60 ± 6 for the hard bursts. The α -values are consistent with the mixed hydrogen/helium bursts from SAX J1748.9–2021 reported by Galloway et al. (2008).

If the anisotropy of the persistent and burst emission, i.e., with a factor of ξ_{p} and ξ_{b} respectively, are considered, it will introduce several consequences (see He & Keek 2016, and references therein). The local accretion rate (\dot{m}) should be multiplied by the factor ξ_{p} , however, it will not effect the obtained power-law index of the $\Delta t_{\text{rec}} - \langle \dot{m} \rangle$ relation if the accretion disk shape did not change dramatically from the soft to hard transition (in both cases, \dot{m} is multiplied by the same factor). The predicted recurrence time is proportional to the observed value, $f_{\text{b}}/F_{\text{pers}}$, so it should be changed with a factor of $\xi_{\text{b}}/\xi_{\text{p}}$. The intrinsic α parameter can be obtained from the observed value divided by a factor of $\xi_{\text{b}}/\xi_{\text{p}}$ (Fujimoto 1988). The $\xi_{\text{b}}/\xi_{\text{p}}$ factor is estimated in the range 1–1.5 for the low inclination system of SAX J1748.9–2021 (He & Keek 2016). These modifications still support the mixed hydrogen/helium bursts occurred in SAX J1748.9–2021.

6. Discussions and Conclusions

SAX J1748.9–2021 has shown 17 bursts during the 1998 outburst detected by *Beppo-SAX*, 16 bursts during the 2001 outburst, 2 bursts during the 2005 outburst, and 13 bursts during the 2009–2010 outburst, all detected by *RXTE* (Galloway et al. 2008). The total number of X-ray bursts was enlarged by the 56 bursts detected during the 2015 outburst. The bursts in soft and hard states are similar in terms of their e-folding time, the peak flux, and the total fluence. Compared with the calculated recurrence time, we conclude that all bursts are generated from mixed hydrogen/helium burning on the NS surface. However, this source appeared to alternate between helium or mixed hydrogen/helium bursts independently of the persistent flux (Galloway et al. 2008), while during the 2015 outburst, studied in detail in this work, the mixed hydrogen/helium bursts follow the

theoretical predicted recurrence time as function of the mass accretion rate $\Delta t_{\text{rec}} \sim \langle \dot{m} \rangle^{-1.02 \pm 0.03}$ (see Fig. 4).

We have studied the spectral behaviour of SAX J1748.9–2021 during its 2015 outburst by using the available *INTEGRAL*, *Swift*, and *XMM-Newton* data. The source shows a spectral change from hard to soft state within ~ 0.5 day around MJD 57079.5 (see Fig. 2). Usually, the hard to soft transitions occurred in the rise phase of the outbursts in LMXBs (see Remillard & McClintock 2006, for reviews). The hard to soft transitions in LMXBs were explained in different ways, i.e., the increasing of mass accretion rate (Esin et al. 1997), the shrinking of the corona size (Homan et al. 2001), the competition between inner halo and outer standard thin accretion disk (Smith et al. 2002), and non-stationary accretion (Yu & Yan 2009). The duration of the outbursts' rise phase in LMXBs can be regarded as the timescale of the hard to soft transition, which has a mean value of 5 days and a minimum value of ~ 1 day (Yan & Yu 2015). Compared with the large sample of outbursts in 36 LMXBs (Yan & Yu 2015), the rapid spectral transition in SAX J1748.9–2021 during the 2015 outburst had the shortest duration. We suggest that the production of soft X-ray photons from type-I X-ray bursts may have accelerated the hard-to-soft spectral transition.

Before the transition, the persistent emission in the 20 – 100 keV range maintained at a high level in ~ 5 days (see the bottom panel in Fig. 2). Considering the recurrence time of ~ 2 hrs in the hard state, we expect that a total number of type-I bursts of ~ 60 should have occurred in SAX J1748.9–2021, and so ~ 35 of these are probably missed because the source was outside the field of view of JEM-X. With the averaged energy release, 3×10^{39} ergs, of type-I X-ray bursts, the total energy of 1.8×10^{41} ergs was emitted in the hard state. The X-ray bursts in SAX J1748.9–2021 may play two roles in the hard-to-soft transition by ejecting a bunch of soft X-ray photons and dragging the accretion flow inward to enhance the persistent X-ray emission (Walker 1992). In both cases, the produced soft photons pass through the corona and cool down the electrons there, which can reduce the corona size and shorten the duration of the hard-to-soft transition. Moreover, we found the source EXO 1745-248 showing a similar behavior, especially considering the two outbursts at MJD 51776 and MJD 52459, respectively. The first outburst was more energetic than the second one. However, the hard to soft transition duration in the first outburst, i.e., ~ 4 days, was smaller than the second one, i.e., ~ 5 days (Yan & Yu 2015). Meanwhile, dozens of X-ray bursts happened before the hard to state transition in the first outburst (Galloway et al. 2008). Considering that more total energy released in the first outburst, we suggested that X-ray bursts may also accelerate the hard to soft spectral transition during the first outburst in EXO 1748-248.

Acknowledgements. We appreciate the suggestions from the referee, which improve the paper. Z.L. thank S.S. Weng for helpful discussions. Z.L. was supported by the Swiss Government Excellence Scholarships for Foreign Scholars. Z.L. thanks the International Space Science Institute and University of Bern for the hospitality. This work was supported by National Natural Science Foundation of China (Grant No. 11703021) and Hunan Provincial Natural Science Foundation of China (Grant Nos. 2017JJ3310 and 2018JJ3495). V.D.F. and M.F. acknowledge the Swiss National Science Foundation project 200021_149865, who financed this research project. V.D.F. acknowledges the International Space Science Institute in Bern for their support. J.P. was supported by the grant 14.W03.31.0021 of the Ministry of Education and Science of the Russian Federation.

References

Altamirano, D., Casella, P., Patruno, A., Wijnands, R., & van der Klis, M. 2008, *ApJ*, 674, L45

- Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 17
- Barret, D. 2001, *Advances in Space Research*, 28, 307
- Bozzo, E., Kuulkers, E., & Ferrigno, C. 2015, *The Astronomer's Telegram*, 7106
- Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, *Space Sci. Rev.*, 120, 165
- Cadelano, M., Pallanca, C., Ferraro, F. R., et al. 2017, *ApJ*, 844, 53
- Chen, Y.-P., Zhang, S., Zhang, S.-N., et al. 2013, *ApJ*, 777, L9
- Chen, Y.-P., Zhang, S., Zhang, S.-N., Li, J., & Wang, J.-M. 2012, *ApJ*, 752, L34
- Courvoisier, T. J.-L., Walter, R., Beckmann, V., et al. 2003, *A&A*, 411, L53
- De Falco, V., Kuiper, L., Bozzo, E., et al. 2017a, *A&A*, 603, A16
- De Falco, V., Kuiper, L., Bozzo, E., et al. 2017b, *A&A*, 599, A88
- Esin, A. A., McClintock, J. E., & Narayan, R. 1997, *ApJ*, 489, 865
- Falanga, M., Bonnet-Bidaud, J. M., Poutanen, J., et al. 2005a, *A&A*, 436, 647
- Falanga, M., Götz, D., Goldoni, P., et al. 2006, *A&A*, 458, 21
- Falanga, M., Kuiper, L., Poutanen, J., et al. 2005b, *A&A*, 444, 15
- Falanga, M., Kuiper, L., Poutanen, J., et al. 2011, *A&A*, 529, A68
- Falanga, M., Kuiper, L., Poutanen, J., et al. 2012, *A&A*, 545, A26
- Falanga, M., Poutanen, J., Bonning, E. W., et al. 2007, *A&A*, 464, 1069
- Frank, J., King, A., & Raine, D. J. 2002, *Accretion Power in Astrophysics: Third Edition*, 398
- Fujimoto, M. Y. 1988, *ApJ*, 324, 995
- Galloway, D. K., Muno, M. P., Hartman, J. M., Psaltis, D., & Chakrabarty, D. 2008, *ApJS*, 179, 360
- Gavriil, F. P., Strohmayer, T. E., Swank, J. H., & Markwardt, C. B. 2007, *ApJ*, 669, L29
- Gierliński, M. & Done, C. 2002, *MNRAS*, 331, L47
- Gierliński, M., Done, C., & Barret, D. 2002, *MNRAS*, 331, 141
- Gierliński, M. & Poutanen, J. 2005, *MNRAS*, 359, 1261
- Goldwurm, A., David, P., Foschini, L., et al. 2003, *A&A*, 411, L223
- Gros, A., Goldwurm, A., Cadolle-Bel, M., et al. 2003, *A&A*, 411, L179
- Hasinger, G. & van der Klis, M. 1989, *A&A*, 225, 79
- He, C.-C. & Keek, L. 2016, *ApJ*, 819, 47
- Homan, J., Wijnands, R., van der Klis, M., et al. 2001, *ApJS*, 132, 377
- Ibragimov, A. & Poutanen, J. 2009, *MNRAS*, 400, 492
- in 't Zand, J., Heise, J., Bazzano, A., et al. 1998, *IAU Circ.*, 6997
- in 't Zand, J. J. M., Verbunt, F., Strohmayer, T. E., et al. 1999, *A&A*, 345, 100
- in 't Zand, J. J. M., van Kerkwijk, M. H., Pooley, D., et al. 2001, *ApJ*, 563, L41
- Jansen, F., Lumb, D., Altieri, B., et al. 2001, *A&A*, 365, L1
- Ji, L., Zhang, S., Chen, Y., et al. 2013, *MNRAS*, 432, 2773
- Ji, L., Zhang, S., Chen, Y., et al. 2015, *ApJ*, 806, 89
- Ji, L., Zhang, S., Chen, Y., et al. 2014a, *ApJ*, 782, 40
- Ji, L., Zhang, S., Chen, Y.-P., et al. 2014b, *A&A*, 564, A20
- Kajava, J. J. E., Sánchez-Fernández, C., Kuulkers, E., & Poutanen, J. 2017, *A&A*, 599, A89
- Kuiper, L., Hermsen, W., Walter, R., & Foschini, L. 2003, *A&A*, 411, L31
- Kuulkers, E., den Hartog, P. R., in 't Zand, J. J. M., et al. 2003, *A&A*, 399, 663
- Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, *A&A*, 411, L141
- Lewin, W. H. G., van Paradijs, J., & Taam, R. E. 1993, *Space Sci. Rev.*, 62, 223
- Lund, N., Budtz-Jørgensen, C., Westergaard, N. J., et al. 2003, *A&A*, 411, L231
- Markwardt, C. B. & Swank, J. H. 2005, *The Astronomer's Telegram*, 495
- Patruno, A., Altamirano, D., Hessels, J. W. T., et al. 2009, *ApJ*, 690, 1856
- Patruno, A. & Watts, A. L. 2012, *ArXiv e-prints*, 1206.2727
- Pintore, F., Sanna, A., Di Salvo, T., et al. 2016, *MNRAS*, 457, 2988
- Pooley, D., Lewin, W. H. G., Verbunt, F., et al. 2002, *ApJ*, 573, 184
- Poutanen, J. 2006, *Advances in Space Research*, 38, 2697
- Poutanen, J. & Svensson, R. 1996, *ApJ*, 470, 249
- Remillard, R. A. & McClintock, J. E. 2006, *ARA&A*, 44, 49
- Sanna, A., Burderi, L., Riggio, A., et al. 2016, *MNRAS*, 459, 1340
- Smith, D. M., Heindl, W. A., & Swank, J. H. 2002, *ApJ*, 569, 362
- Strohmayer, T. & Bildsten, L. 2006, *New views of thermonuclear bursts*, ed. W. H. G. Lewin & M. van der Klis, 113–156
- Strüder, L., Briel, U., Dennerl, K., et al. 2001, *A&A*, 365, L18
- Suzuki, M., Negoro, H., Nakahira, S., et al. 2009, *The Astronomer's Telegram*, 2360
- Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131
- Valenti, E., Ferraro, F. R., & Origlia, L. 2007, *AJ*, 133, 1287
- van der Klis, M. 2006, *Rapid X-ray Variability*, ed. W. H. G. Lewin & M. van der Klis, 39–112
- Walker, M. A. 1992, *ApJ*, 385, 642
- Winkler, C., Courvoisier, T. J.-L., Di Cocco, G., et al. 2003, *A&A*, 411, L1
- Yan, Z. & Yu, W. 2015, *ApJ*, 805, 87
- Yu, W. & Yan, Z. 2009, *ApJ*, 701, 1940