

# The Contact Process on Random Graphs and Galton-Watson Trees

Xiangying Huang and Rick Durrett  
Dept. of Math, Duke University

June 20, 2022

## Abstract

The key to our investigation is an improved (and in a sense sharp) understanding of the survival time of the contact process on star graphs. Using these results, we show that for the contact process on Galton-Watson trees, when the offspring distribution (i) is subexponential the critical value for local survival  $\lambda_2 = 0$  and (ii) when it is geometric( $p$ ) we have  $\lambda_2 \leq C_p$ , where the  $C_p$  are much smaller than previous estimates. We also study the critical value  $\lambda_c(n)$  for “prolonged persistence” on graphs with  $n$  vertices generated by the configuration model. In the case of power law and stretched exponential distributions where it is known  $\lambda_c(n) \rightarrow 0$  we give estimates on the rate of convergence. Physicists tell us that  $\lambda_c(n) \sim 1/\Lambda(n)$  where  $\Lambda(n)$  is the maximum eigenvalue of the adjacency matrix. Our results suggest that this is not correct.

## 1 Introduction

Harris [13] introduced the contact process on  $\mathbb{Z}^d$  in 1974. The state at time  $t$  is  $\xi_t \subset \mathbb{Z}^d$ . It is often thought of as a model for the spread of species. In this case  $\xi_t$  is the set of occupied sites, and sites in  $\xi_t^c$  are vacant. However, it can also be viewed as a spatial SIS epidemic model. In this case  $\xi_t$  is the set of infected sites, and sites in  $\xi_t^c$  are susceptible. Both interpretations are common in the literature, so the reader will see both here.

In the contact process on a graph  $G$ , occupied sites become vacant at rate 1, and give birth onto vacant neighbors at rate  $\lambda$ . Let  $\xi_t^0$  be the process starting from only the origin occupied and let  $\xi_t^1$  be the process starting from all sites occupied. Harris introduced the critical value

$$\lambda_c = \inf\{\lambda : P(\xi_t^0 \neq \emptyset \text{ for all } t) > 0\},$$

and proved that  $0 < \lambda_c < \infty$ . He also showed that for  $\lambda > \lambda_c$ ,  $\xi_t^1$  converges to a limit that is a nontrivial stationary distribution. A rich theory has been developed for the contact process on  $\mathbb{Z}^d$ . See Liggett’s 1999 book [17] for a summary of much of what is known.

Pemantle [28] was the first to study the contact process on the tree  $\mathbb{T}^d$  in which each vertex has degree  $d + 1$ . Here, and in what follows, we assume  $d \geq 2$  since  $\mathbb{T}_1 = \mathbb{Z}$ . Let 0 be

the root of the tree and let  $P_0$  be the probability measure for the process starting from only the root occupied. Pemantle found that the contact process on  $\mathbb{T}^d$  had two critical values.

$$\begin{aligned}\lambda_1 &= \inf\{\lambda : P_0(\xi_t \neq \emptyset \text{ for all } t) > 0\}, \\ \lambda_2 &= \inf\{\lambda : \liminf_{t \rightarrow \infty} P_0(0 \in \xi_t) > 0\}.\end{aligned}$$

By deriving bounds on the critical values, he showed that  $\lambda_1 < \lambda_2$  when  $d \geq 3$ . Liggett [16] improved the bounds to settle the case  $d = 2$ . At about the same time, Stacey [32] gave a proof that  $\lambda_1 < \lambda_2$  that did not rely on bounds. The stationary distributions and limiting behavior of the contact process on trees is an interesting subject that has been extensively studied. See Liggett’s book [17] for an account of the results.

## 1.1 Results for star graphs

Let  $G_k$  be a star graph with center 0 and leaves  $1, 2, \dots, k$  and let  $\xi_t$  be set of vertices infected in the contact process at time  $t$ . Write the state  $\xi_t$  as  $(i, j)$  where  $i$  is the number of infected leaves and  $j = 1$  if the center is infected and  $j = 0$  otherwise. We write  $P_{i,j}$  for the law of the process starting from  $(i, j)$ . Pemantle [28] was the first to study the persistence time of the contact process on stars. See his Section 4. He did his analysis on the “ladder graph”  $\{0, \dots, n\} \times \{0, 1\}$  so he ended up with a very approximate superharmonic function  $W(\xi)$ . Let  $i$  be the number of infected leaves, and let  $I(\xi) = 1$  if the root is infected and  $= 0$  otherwise.

$$W(\xi) = e^{-\lambda i/10} \left( 1 - I(\xi) \frac{(e^{\lambda/10} - 1)}{\lambda} \right).$$

To make the connection change Pemantle’s  $n$  (the number of leaves) to our  $k$  and note that his birth rate  $\lambda = \alpha/\sqrt{n}$ . Pemantle has an interesting heuristic discussion on pages 2015–2016 that explains why this form is reasonable. However the 10’s that are thrown in to make it is easier to prove it is superharmonic ruin its accuracy.

Here, following the approach in [4], we will reduce to a discrete time one dimensional chain, we will only look at times when  $j = 1$ . When the state is  $(i, 0)$  with  $i > 0$ , the next event will occur after exponential time with mean  $1/(i\lambda + i)$ . The probability that it will be the reinfection of the center is  $\lambda/(\lambda + 1)$ . The probability it will be the healing of a leaf is  $1/(\lambda + 1)$ . Thus, the number of leaf infections  $N$  that will be lost while the center is healthy has a shifted geometric distribution with success probability  $\lambda/(\lambda + 1)$ , i.e.,

$$P(N = j) = \left( \frac{1}{\lambda + 1} \right)^j \cdot \frac{\lambda}{\lambda + 1} \quad \text{for } j \geq 0.$$

Note that

$$EN = \frac{\lambda + 1}{\lambda} - 1 = \frac{1}{\lambda}.$$

The next step is to modify the chain so that the infection rate is 0 when the number of infected leaves is at least

$$L = pk \quad \text{where } p = \lambda/(1 + 2\lambda). \tag{1}$$

(The reader will see the reasons that underlie this choice later.) Note that the number of infected leaves is always  $\leq pk$  and the number of uninfected leaves is  $\geq (1-p)k$ . Thus if we look at the embedded discrete time process for the contact process on the star and only look at times when the center is infected, the process dominates  $Y_n$  where

$$\begin{array}{ll} \text{jump} & \text{with prob} \\ Y_n \rightarrow Y_n - 1 & pk/D \\ Y_n \rightarrow \min\{Y_n + 1, pk\} & \lambda(1-p)k/D \\ Y_n \rightarrow Y_n - N & 1/D \end{array}$$

Here  $N$  is independent of  $Y_n$  and the denominator

$$D = pk + \lambda(1-p)k + 1 \leq k + \lambda k + 1 \leq (2 + \lambda)k. \quad (2)$$

The fact that  $Y_n$  is a reflecting random walk will simplify computations. We will use the process to lower bound survival times. Before the infection on the star graph goes extinct it will spend most of its time near  $pk$ , (i) this does not lose much compared to the more accurate birth and death chain, which uses the actual number of infected leaves not just a bound, and (ii) we make only a small error when we return to continuous time by assuming that jumps happen at the maximum rate. In [4] it is shown, see Lemma 2.2 on page 2339, that

**Lemma 1.1.** *Suppose  $\lambda \leq 1$  and  $\lambda^2 k \geq 50$ . Let  $L_0 = \lambda k/4$  and  $S_0 = \frac{1}{4L_0} \exp(k\lambda^2/80)$ . Then*

$$P_{L,i} \left( \inf_{t \leq S_0} |\xi_t| \leq 0.4L_0 \right) \leq 7e^{-\lambda^2 L_0/80}.$$

In contrast our Lemma 2.4 will show that if  $L = \lambda k/(1 + 2\lambda)$  and  $b = \epsilon L$

$$P_{L,1} \left( \inf_{t \leq S} \xi_t \leq bL \right) \leq (3 + \lambda)(1 + \lambda/2)^{-\epsilon L} \quad \text{where} \quad S = \frac{1}{(2 + \lambda)2k} (1 + \lambda/2)^{L(1-2\epsilon)}. \quad (3)$$

Part of the improvement comes from simply replacing  $L_0$  by  $L$  and 0.4 by  $\epsilon$ , but the most important change is to construct a more accurate superharmonic function. If one is proving that a critical value is 0, as [28] and [4] were, then it is not harmful to be off by a large constant factor, but if we are trying to get a good positive upper bound we need to be accurate.

The more precise lower bound we developed here was useful in a summer 2018 research project with four Duke undergrads to prove that on the  $(1, n)$  tree, in which vertices of degree 1 and  $n$  alternate, we have

**Theorem 1.** *Theorem 5 in [14]. On the  $(1, n)$  tree, as  $n \rightarrow \infty$  the critical value*

$$\lambda_2 \sim \sqrt{0.5(\log n)/n}.$$

In that paper the following upper bound on the survival time is proved. Let  $K = \lambda k/(\lambda + 1)$ . Let  $T_{0,0}$  be the time until the state is 0, 0, i.e., infection dies out.

**Theorem 2.** *Lemma 4 in [14]. Let  $K = \lambda n / (n + 1)$ . For any  $\epsilon > 0$ , the contact process on star graph has*

$$E_{K,1}T_{0,0} \leq (\log n)e^{(1+\epsilon)\lambda^2 n}.$$

when  $n$  is sufficiently large.

If  $\lambda^2 n \rightarrow \infty$  then the  $\log n$  prefactor can be absorbed by changing  $\epsilon$  however it is important if  $\lambda = O(1/\sqrt{n})$ , since in this case the exponential is  $O(1)$ .

In contrast, the lower bound time  $T$  from (3), ignoring the prefactor, is

$$(1 + \lambda/2)^{L(1-2\epsilon)} \approx \exp\left((1 - 2\epsilon)\frac{\lambda^2 k}{2(1 + 2\lambda)}\right).$$

If  $\lambda$  is small then term in the exponential is about 1/2 the one in Theorem 2. Strictly speaking these results are not sharp (on the exponential scale) but a factor of 2 is much better than the factor of 80 that appears in [4]. It is not clear which result gives the right answer. The result in Theorem 2 is proved by looking at the first time the center becomes healthy and then all of the leaves become healthy before the center is reinfected. At first sight this bound seems crazy, but the calculations above show that it is fairly accurate. We have not been able to finding a good subharmonic function for  $Y_n$  to find a better upper bound so we leave it to a clever reader to determine the nature of the large deviation event that wipes out the infection on the star.

## 1.2 Galton-Watson trees

Given an offspring distribution  $p_k$ , we construct a Galton-Watson trees as follows. Starting with the root, each individual has  $k$  children with probability  $p_k$ . Pemantle has shown that

**Theorem 3.** *Theorem 3.2 in [28]. There are constants  $c_2$  and  $c_3$  so that if  $\mu$  is the mean of the offspring distribution, then for any  $k > 1$ , if we let  $r_k = \max\{2, c_2 \log(1/kp_k)/\mu\}$*

$$\lambda_2 < c_3 \sqrt{r_k \log r_k \log(k)/k}.$$

If the offspring distribution in the Galton-Watson tree is a stretched exponential  $p_k = c_\gamma \exp(-k^\gamma)$  with  $\gamma < 1$  then  $\log(1/kp_k) \sim k^\gamma$  and hence  $\lambda_2 = 0$ .

Given this result, it is natural to ask about the critical values  $\lambda_1$  and  $\lambda_2$  when degrees have a geometric distribution.  $p_k = (1 - p)^{k-1}p$  for  $k \geq 1$ . The most interesting problem is to prove  $\lambda_1 > 0$ . Here, we prove upper bounds.

**Theorem 4.**  $\lambda_1 \leq p/(1 - p)$ .

*Proof.* Modify the contact process so that births from a site can only occur on sites further from the root. Each vertex  $x$  will be occupied at most once. If  $x$  is occupied then it will give birth with probability  $\lambda/(\lambda + 1)$  onto each neighbor  $y$ . The birth events are not independent but that is not important. If we let  $Z_n$  be the number of sites at distance  $n$  that are ever occupied,  $Z_n$  is a branching process in which the offspring distribution has mean  $\lambda/((\lambda + 1) \cdot p)$  which is  $> 1$  if  $\lambda > p/(1 - p)$ .  $\square$

When  $p_k = (1-p)^{k-1}p$ ,  $\log(1/kp_k) \sim c_p k$ , so Theorem 3 gives a finite upper bound on  $\lambda_2$ . It is difficult to trace through all the calculations to get an explicit lower bound. However, Pemantle uses  $e^{-1}/5 = 0.0735$  as the lower bound for the probability of long time survival starting with only the center of a large degree star graph occupied, while Lemma 2.5 gives  $1 - 3k^{-1/3}$  when the degree is  $k$ . This probability  $e^{-1}/5$  appears cubed near the end of his proof, so we think that his bound is much worse than the following:

**Theorem 5.** *If  $p_k = 2^{-k}$  for  $k \geq 1$ , then  $\lambda_2 \leq 2.5$ .*

This result is proved by combining our new estimates for the contact process on stars with the mysterious Lemma 2.4 in Pemantle’s paper [28] (see Lemma 3.3 below).

The proof works for a general geometric  $p_k = (1-p)^{k-1}p$ ,  $k \geq 1$ . We cannot get a nice formula for the upper bound as a function of  $p$  but the upper bounds can easily be computed numerically and graphed. These upper bounds are only interesting for small  $p$ . A Galton-Watson tree with  $p_0 = 0$  and  $p_1 < 1$  contains a copy of  $\mathbb{Z}$  (start with a vertex with two children and follow their descendants) so using Liggett’s bound on  $\lambda_c(\mathbb{Z})$  proved in [15] we conclude  $\lambda_2 \leq 2$  for all  $0 < p < 1$ .




Figure 1: Upper bounds on  $\lambda_2$  (solid line) and  $\lambda_1$  (dotted line) as a function of  $p$  for the geometric degree distribution. The graph is computed by using (10).

In addition, the proof of Theorem 5 yields an improvement of Pemantle’s result for stretched exponential distributions. We say that  $p_k$  is subexponential if

$$\limsup_{k \rightarrow \infty} (1/k) \log p_k = 0.$$

**Theorem 6.** *If the offspring distribution  $p_k$  for a Galton-Watson tree is subexponential and has mean  $\mu > 1$  then  $\lambda_2 = 0$ .*

Note that  $\lambda_2 = 0$  implies  $\lambda_1 = 0$ .

We conjecture that the result in Theorem 6 is sharp. To have a concrete problem, we consider only the special case of geometric distributions

**Conjecture 1.** *Suppose that degree distribution has  $P(d(x) \geq k) = (1 - p)^k$  for  $k \geq K$ . Then  $\lambda_2 > 0$ .*

When the degree distribution is bounded, i.e.,  $d(x) \leq D$ , there is a trivial lower bound on the critical value:  $\lambda_1 \geq 1/D$ . When the degree distribution is unbounded, lower bounds become much more difficult than upper bounds. One can prove upper bounds by giving a recipe that guarantees survival of the process. For lower bounds one must consider everything that can possibly happen. Menard and Singh [18] have recently proved  $\lambda_1 > 0$  for the contact process on random geometric graphs (RGG). A random geometric graph is constructed as follows. Vertices are points of a Poisson process on  $\mathbb{R}^d$  with  $d \geq 2$ . Two vertices  $v \neq w$  are connected if  $\|v - w\| < R$ . For percolation properties of RGG's see the book by Penrose [30].

They prove their results by studying “cumulative merging processes.” The result of applying such a procedure is a partition of the vertices of a weighted graph, which has the property that if  $C, C'$  are clusters and  $C \neq C'$  then

$$d(C, C') > \min\{r(C), r(C')\}^\alpha \tag{4}$$

where  $r(C)$  is the sum of the weights in the cluster. Theorem 4.1 in their paper shows that if choose  $\Delta > 0$ , let

$$r(x) = d(x)1_{\{d(x) \geq \Delta\}}$$

and show (4) holds for some  $\alpha \geq 5/2$  then  $\lambda_1 > 0$ . The key to the proof is to use partition to define a “time delayed branching Markov chain” that controls the growth of the contact process. While the proof is ingenious, it cannot work on Galton-Watson trees since in that setting useful partitions satisfying (4) do not exist when  $\alpha > 1$ . If a vertex has large enough degree, it will with positive probability belong to an infinite component.

### 1.3 Finite graphs

Consider the contact process on  $\{-n, \dots, n\}$  starting from all sites occupied and let  $\tau_n = \inf\{t : \xi_t = \emptyset\}$ . Combining results of Durrett and Liu [10] and Durrett and Schonmann [11] gives the following results

(i) If  $\lambda < \lambda_c$  then there is a constant  $\gamma_1(\lambda)$  so that

$$\tau_n / \log n \rightarrow \gamma_1(\lambda) \quad \text{in probability.}$$

(ii) If  $\lambda > \lambda_c$  then there is a constant  $\gamma_2(\lambda)$  so that

$$(\log \tau_n) / n \rightarrow \gamma_2(\lambda) \quad \text{in probability.}$$

(iii) When  $\lambda > \lambda_c$  there is “metastability:”

$$\tau_n / E\tau_n \Rightarrow \text{exponential}(1).$$

where  $\Rightarrow$  means convergence in distribution. Intuitively, the process on the interval stays exponentially long in a state that looks like the stationary distribution for the process on  $\mathbb{Z}$ , and then suddenly dies out.

Results on  $\mathbb{Z}^d$  with  $d > 1$  had to wait for the work of Bezuidenhout and Grimmett [1], who showed that in  $d > 1$  the contact process dies out at the critical value and in doing so introduced a block construction that can be used to study the supercritical process. Mountford [19] proved the metastability result in 1993 and that  $(\log \tau_n)/n^d \rightarrow \gamma(\lambda)$  in 1999, see [20].

Stacey [31] studied the contact process on a tree truncated at height  $\ell$ ,  $\mathbb{T}_\ell^d$ . To be precise, the root has degree  $d$ , vertices at distance  $0 < k < \ell$  from the root have degree  $d + 1$ , while those at distance  $\ell$  have degree 1. Cranston, Mountford, Mourrat, and Valesin improved Stacey's result to establish that the time to extinction starting from all sites occupied  $\tau_\ell^d$  satisfies

**Theorem 7.** [6] (a) For any  $0 < \lambda < \lambda_2(\mathbb{T}^d)$  there is a  $c \in (0, \infty)$  so that as  $\ell \rightarrow \infty$

$$\tau_\ell^d / \log |\mathbb{T}_\ell^d| \rightarrow c \quad \text{in probability.}$$

(b) For any  $\lambda_2(\mathbb{T}^d) < \lambda < \infty$  there is a  $c \in (0, \infty)$  so that as  $\ell \rightarrow \infty$

$$\log(\tau_\ell^d) / |\mathbb{T}_\ell^d| \rightarrow c \quad \text{in probability.}$$

Moreover  $\tau_\ell^d / E\tau_\ell^d$  converges to a mean one exponential.

When a tree is truncated at a finite distance, a positive fraction of the sites are on the boundary. A more natural finite version of a tree is a random regular graph in which all vertices have degree  $d + 1$ . In this case there is no boundary and the graph has the same distribution viewed from any point. If there are  $n$  vertices, the graph looks like  $\mathbb{T}_d$  in neighborhoods of a point that have  $\leq n^{1/3}$  vertices. Mourrat and Valesin have shown for a random regular graph, the time to extinction starting from all sites occupied  $\tau_n$  satisfies:

**Theorem 8.** [23] (a) For any  $0 < \lambda < \lambda_1(\mathbb{T}^d)$  there is a  $C < \infty$  so that as  $n \rightarrow \infty$

$$P(\tau_n < C \log n) \rightarrow 1,$$

(b) For any  $\lambda_1(\mathbb{T}^d) < \lambda < \infty$  there is a  $c > 0$  so that as  $n \rightarrow \infty$

$$P(\tau_n > e^{cn}) \rightarrow c.$$

Notice that the threshold in the second result comes at  $\lambda_1$ , while the one in Stacey's result comes at  $\lambda_2$ . The difference is that when  $\lambda \in (\lambda_1, \lambda_2)$  on the infinite tree the origin is in the middle of linearly growing vacant region. On the truncated tree the system dies out when the vacant region is large enough. However, on the random regular graph the occupied sites will later return to the origin. Durrett and Jung [9] investigated the qualitative differences between  $\lambda \in (\lambda_1, \lambda_2)$  and  $\lambda > \lambda_2$  on the small world graph.

To construct a random graph  $G_n$  on the vertex set  $\{1, 2, \dots, n\}$  having a specified degree distribution, we use the *configuration model*. Let  $d_1, \dots, d_n$  be independent and have the distribution  $P(d_i = k) = p_k$ . In order to have a valid degree sequence, we condition on the event  $E_n = \{d_1 + \dots + d_n \text{ is even}\}$ . Since  $P(E_n) \rightarrow 1/2$  as  $n \rightarrow \infty$ , the conditioning will have a little effect on the distribution of  $d_i$ 's. Having chosen the degree sequence  $(d_1, d_2, \dots, d_n)$ ,

we attach  $d_i$  half-edges to the vertex  $i$ , and then pair these half-edges at random. This procedure may produce a graph with self-loops or parallel edges, but we will ignore that problem for the moment.

In the early 2000’s physicists studied the contact process on a random graphs with a power-law degree distribution, i.e., the degree of each vertex is  $k$  with probability

$$p_k \sim Ck^{-\alpha} \quad \text{as } k \rightarrow \infty.$$

Pastor-Satorras and Vespignani [25, 26, 27] have made an extensive study of this model using mean-field methods. Their nonrigorous computations suggest the following conjectures about  $\lambda_c$  the threshold for “prolonged persistence” of the contact process, and the critical exponent  $\beta$ , that controls the rate at which the equilibrium density of occupied sites  $\rho(\lambda)$  goes to 0, i.e.,  $\rho(\lambda) \sim C(\lambda - \lambda_c)^\beta$ .

- If  $\alpha \leq 3$ , then  $\lambda_c = 0$ . If  $\alpha < 3$  then  $\beta = 1/(3 - \alpha)$ .
- If  $3 < \alpha \leq 4$ , then  $\lambda_c > 0$  and  $\beta = 1/(\alpha - 3) > 1$ .
- If  $\alpha > 4$ , then  $\lambda_c > 0$  and  $\beta = 1$ .

See also Section V of [24]. The values of  $\beta$  quoted above are given in formula (29) of [24].

Chatterjee and Durrett [4] showed in 2009 that  $\lambda_c > 0$  not correct when  $\alpha > 3$  and  $P(d_i \leq 2) = 0$ . The last condition guarantees that the graph is connected and that random walks on the graph have good mixing properties. They only proved survival for time  $\exp(O(n^{1-\epsilon}))$  but they obtained bounds on the critical exponent  $\beta$ .

In 2013 Mountford, Mourrat, Valesin, and Yao [22] extended the results of [4] to include  $2 < \alpha \leq 3$  and proved upper and lower bounds that had the same dependence on  $\lambda$  but different constants

$$\begin{cases} \lambda^{1/(3-\alpha)} & 2 < \alpha \leq 5/2 \\ \lambda^{2\alpha-3} \log^{2-\alpha}(1/\lambda) & 5/2 < \alpha \leq 3 \\ \lambda^{2\alpha-3} \log^{4-2\alpha}(1/\lambda) & 3 < \alpha \end{cases}$$

The result for  $2 < \alpha \leq 5/2$  agrees with the mean-field calculations quoted above but that formula is claimed to hold for  $2 < \alpha < 3$ . Figure 2 gives a visual comparison of the mean-field and rigorous results for critical exponents. For more about why the change occurs at  $5/2$  see the next section and [22]. Three years later, Mountford, Mourrat, Valesin, and Yao [21] showed the survival time  $\geq e^{cn}$ .

## 1.4 Critical value asymptotics when $\lambda_c = 0$

While the results cited above show that the mean-field calculations are not correct, physicists have never said they were wrong. Indeed, a 2010 paper Castellano and Pastor-Satorras [2] claims they knew the right answer all along. “Already in 2003, Wang et al [33] argued that the SIS epidemic threshold on any graph is set by the largest eigenvalue of the adjacency matrix,  $\Lambda$

$$\lambda_c(n) = 1/\Lambda(n). \tag{5}$$






Figure 2: Mean field critical exponents (solid line) versus rigorous results (dashed line) as  $\alpha$  varies from 2 to 4.5.

Two years earlier Pemantle and Stacey [29] proved that  $1/\Lambda(n)$  is the critical value of branching random walk on the graph. This should not be surprising since the number of paths of length  $n$  is  $\approx \Lambda^n$ . However, it is far from obvious why this should also be the critical value for the contact process, which is often much larger

The first question that needs to be addressed before (5) can become a theorem is the definition of  $\lambda_c$ . According to page 942 of the 2015 survey paper in *Reviews of Modern Physics* [24] “Above the epidemic threshold, the activity must be *endemic*, so that the average time to absorption is  $O(e^{cn})$ .” To make it clear that they wanted to insist on this standard we note that the discussion continued with

“Chatterjee and Durrett proved that in graphs with power law degree distribution  $ET > \exp(O(N^{1-\delta}))$  for any  $\delta > 0$ . This result pointed to a vanishing threshold but still left the possibility for nonendemic long-lived metastable states.”

Survival for time  $e^{cn}$  is certainly the gold standard for prolonged persistence, but following the footsteps of Ganesh, Masoulié, and Towsley [12], we will accept survival for time  $\exp(O(n^\epsilon))$  for some  $\epsilon > 0$  as evidence that  $\lambda > \lambda_c$ .

The proofs of (5) in [33] and [3] do not provide a lower bound on survival time. They let  $n \rightarrow \infty$  to obtain a nonlinear dynamical system (NLDS). To explain, note that if we let  $p_{i,t}$  be the probability  $i$  is infected at time  $t$  and let  $\zeta_{j,t}$  be the probability  $j$  does not receive infection at time  $t$  then

$$\zeta_{i,t} = \prod_{j:j \sim i} (1 - \beta p_{j,t-1})$$

$$1 - p_{i,t} = (1 - p_{i,t-1})\zeta_{i,t} + \delta p_{i,t-1}\zeta_{i,t}$$

Then they argue that if  $\lambda > \Lambda^{-1}$  then one of the eigenvalues of the linearization of the NLDS around 0 is  $> 1$ , see the Appendix of [3]. It is not clear what the last conclusion implies

in terms of persistence. Wang et al [33] use (5) to conclude that the critical value for the contact process on a star graph with  $n$  leaves is  $1/\sqrt{n}$ .

The results discussed in Section 1.1 show that the survival time on the star graph increase dramatically when  $\lambda$  changes from  $O(1/\sqrt{n})$  to  $\gg 1/\sqrt{n}$ . However, the claim that critical value on a star graph is  $1/\Lambda(n)$  is contradicted by Theorem 2 which shows that if  $\lambda = \alpha/n^{1/2}$  then for large  $n$

$$E_{K,1}T_{0,0} \leq e^{2\alpha} \log n$$

where  $K = \lambda n/(\lambda + 1)$ . It is not hard to show that the time needed to go from  $n$  to  $K$  is  $O(\log n)$ . Thus the survival time is  $O(\log n)$  which is much smaller than the  $O(e^{cn})$  that [24] demands. Since the results in Section 1.1. show that the survival time is  $\exp(O(\lambda^2 n))$ , we would have to take  $\lambda > 0$  independent of  $n$  for the contact process on the start to survive for this long

Returning to the implications of (5) for the contact process, the maximum eigenvalue of the adjacency matrix of a random graph is trivially  $\geq d_{max}^{1/2}$  (generate paths by going back and forth between a vertex with degree  $d_{max}$  and its neighbors). Using results of Chung, Lu, and Vu [5] for the maximum eigenvalue for random graphs the authors of [2] concluded that the critical value for power law random graphs satisfies

$$\lambda_c \sim \begin{cases} \langle d \rangle / \langle d^2 \rangle & 2 < \alpha < 5/2 \\ 1/\sqrt{d_{max}} & 5/2 < \alpha \end{cases}$$

where  $d_{max}$  is the maximum degree in the graph, and  $\langle d \rangle$ ,  $\langle d^2 \rangle$  are the average values of  $d(x)$  and  $d(x)^2$  for the graph. More concretely

$$\lambda_c(n) \sim \begin{cases} n^{(\alpha-3)/(\alpha-1)} & 2 < \alpha < 5/2 \\ n^{-1/2(\alpha-1)} & 5/2 < \alpha \end{cases}$$

Using our results we can prove an upper bound on  $\lambda_c$  that supports this prediction when  $\alpha > 3$ . Here  $a = \alpha - 1$ .

**Theorem 9.** *Suppose that the degree distribution has*

$$P(d(x) \geq k) = 3^a k^{-a} \quad \text{for } k \geq 3.$$

*We assume  $a > 2$  so that  $Ed(x)^2 < \infty$ . Let  $\lambda = n^{-(1-2\eta)/2a}$  and  $\eta > 0$ . If we start from all 1's then there is an  $\epsilon > 0$  so that the system survives for time  $\exp(O(n^\epsilon))$  with high probability.*

Combining this result with the fact that  $1/\Lambda$  gives the critical value for branching random walk and hence a lower bound on the critical value for the contact process we have

$$\lambda_c(n) = n^{-(1+o(1))/2a}. \tag{6}$$

Next we consider the stretched exponential

$$P(d(x) \geq k) = \exp(-x^{1/b} + 3^{1/b}) \quad \text{for } k \geq 3.$$

where  $b > 1$ . In this case, the maximum degree vertex on a graph with  $n$  vertices is  $\sim \log^b n$ , so the maximum eigenvalue  $\Lambda \sim \log^{b/2} n$  and the formula in (5) predicts that  $\lambda_c \approx \log^{-b/2} n$  but our rigorous upper bound is larger.

**Theorem 10.** *Suppose  $\lambda_n = \log^{(1-\eta)(1-b)/2} n$ . If we start from all 1's then for any  $\epsilon > 0$  the system survives for time  $\exp(O(n^{1-\epsilon}))$  with high probability.*

We believe that the last result gives the right answer.

**Conjecture 2.** *Suppose  $\lambda_n = \log^{-a/2} n$  where  $a > b - 1$ . If we start from all 1's then for any  $\epsilon > 0$  the system dies out by time  $\exp(O(n^\epsilon))$  with high probability.*

The remainder of the paper is devoted to proofs. Section 2 gives our results for the star graph. Section 3 proves our results for Galton-Watson trees. Section 4 gives the asymptotics for  $\lambda_c(n)$ .

## 2 Results for the star graph

Recall that we set

$$L = pk \quad \text{where} \quad p = \lambda/(1 + 2\lambda).$$

**Lemma 2.1.** *Let  $e^\theta = 1/(1 + \lambda/2)$ . If  $k$  is large enough  $e^{\theta Y_n}$  is a supermartingale while  $Y_n \in (0, pk)$ .*

*Proof.* We begin by noting that

$$\begin{aligned} E(\exp(\theta Y_{n+1}) - \exp(\theta Y_n) | Y_n = y) &= e^{\theta y} (e^\theta - 1) \lambda (1 - p) k / D \\ &+ e^{\theta y} (e^{-\theta} - 1) p k / D + \frac{e^{\theta y}}{D} \left[ \sum_{j=0}^{\infty} \left( \frac{e^{-\theta}}{1 + \lambda} \right)^j \left( \frac{\lambda}{1 + \lambda} \right) - 1 \right]. \end{aligned} \quad (7)$$

The term in square brackets is

$$\frac{1}{1 - e^{-\theta}/(1 + \lambda)} \cdot \frac{\lambda}{1 + \lambda} - 1 = \frac{\lambda}{1 + \lambda - e^{-\theta}} - 1 = \frac{e^{-\theta} - 1}{1 + \lambda - e^{-\theta}} \geq 0.$$

Note that this implies we must take  $e^{-\theta} < 1 + \lambda$ .

The first two terms are

$$\frac{e^{\theta y} k}{D} ((e^\theta - 1) \lambda (1 - p) + (e^{-\theta} - 1) p),$$

so we begin by solving

$$(e^\theta - 1) \lambda (1 - p) + (e^{-\theta} - 1) p = 0.$$

Rearranging and setting  $x = e^\theta$  we want

$$x^2 \lambda (1 - p) - [\lambda (1 - p) + p] x + p = 0.$$

Factoring we have

$$(\lambda (1 - p) x - p)(x - 1) = 0.$$

Since  $p = \lambda/(1 + 2\lambda)$  the smaller root is

$$\frac{p}{\lambda(1-p)} = \frac{\lambda/(1+2\lambda)}{\lambda(1+\lambda)/(1+2\lambda)} = \frac{1}{1+\lambda}.$$

We let  $e^\theta = 1/(1 + \lambda/2) \in (1/(1 + \lambda), 1)$  so that there is a  $\delta > 0$  with

$$e^\theta \lambda(1-p) + e^{-\theta} p = [\lambda(1-p) + p] - \delta$$

and hence

$$(e^\theta - 1)\lambda(1-p)k + (e^{-\theta} - 1)pk + \frac{e^{-\theta} - 1}{1 + \lambda - e^{-\theta}} = -\delta k + \frac{e^{-\theta} - 1}{1 + \lambda - e^{-\theta}}.$$

From this we see that if  $k$  is large enough  $e^{\theta Y_n}$  is a supermartingale while  $Y_n \in (0, pk)$ . Note that when  $Y_n$  is small the number of infected leaves may become 0 before the center is reinfected but in this case the number of lost infections  $N$  is truncated.  $\square$

Let  $T_\ell^- = \inf\{n : Y_n \leq \ell\}$  and let  $T_m^+ = \inf\{n : Y_n \geq m\}$ .

**Lemma 2.2.** *Let  $a, b \in (0, L)$ . If  $b < a$  then*

$$P_a(T_b^- < T_L^+) \leq (1 + \lambda/2)^{b-a}.$$

*Proof.* To estimate the hitting probability let  $\phi(x) = \exp(\theta x)$  where we take  $e^\theta = 1/(1 + \lambda/2)$  and note that if  $\tau = T_b^- \wedge T_L^+$  then  $\phi(Y(t \wedge \tau))$  is a supermartingale. Let  $q = P_a(T_b^- < T_L^+)$ . Using the optional stopping theorem we have

$$q\phi(Y_b^-) + (1-q)\phi(Y_L^+) \leq \phi(a).$$

It is possible that  $Y_b^- < b$ . Note that since  $\theta < 0$ , we have  $\phi(x) \geq \phi(b)$  for  $x \leq b$ . Hence,

$$q\phi(b) + (1-q)\phi(L) \leq \phi(a).$$

Dropping the second term on the left,  $q \leq \phi(a)/\phi(b) = (1 + \lambda/2)^{b-a}$ , which completes the proof.  $\square$

**Lemma 2.3.** *If  $R_L = \inf\{n > T_{L-1}^- : Y_n = L\}$  and  $b \in [0, L)$  then*

$$P_L(T_b^- < R_L) \leq (2 + \lambda)(1 + \lambda/2)^{b-L}.$$

**Remark.** Here, and in later lemmas, the computation of explicit constants is somewhat annoying. However, when we consider asymptotics for critical values,  $\lambda$  will go to 0, so we will need to know how the constants depend on  $\lambda$ .

*Proof.* To compute the left-hand side we break things down according to the first jump. The definition of  $R_L$  allows us to ignore the attempted upward jumps that do nothing. Recall that  $L = pk$ . The jump is to  $L - 1$  with probability  $pk/(pk + 1)$  and to  $L - j$  with probability  $\frac{\lambda}{(1+\lambda)^{j+1}} \cdot \frac{1}{1+pk}$ . In the first case the probability of returning to  $L$  before going below  $b$  is

$$\leq (1 + \lambda/2)^{b-(L-1)} = (1 + \lambda/2) \cdot (1 + \lambda/2)^{b-L}.$$

In the second we have to sum over the possible values of  $L - j$ . Using Lemma 2.2

$$\begin{aligned}
&\leq (1 + \lambda/2)^{b-L} \sum_{j=1}^{\infty} \frac{\lambda}{(1 + \lambda)^{j+1}} (1 + \lambda/2)^j + \frac{\lambda}{1 + \lambda} P_L(T_b^- < R_L) \\
&\leq (1 + \lambda/2)^{b-L} \frac{\lambda}{\lambda + 1} \cdot \sum_{j=0}^{\infty} \left( \frac{1 + \lambda/2}{1 + \lambda} \right)^j + \frac{\lambda}{1 + \lambda} P_L(T_b^- < R_L) \\
&= 2(1 + \lambda/2)^{b-L} + \frac{\lambda}{1 + \lambda} P_L(T_b^- < R_L).
\end{aligned}$$

Noting that  $\max\{2, 1 + \lambda/2\} \leq 2(1 + \lambda/2) - \delta$  for some small  $\delta < \lambda$ , we have the following relation,

$$P_L(T_b^- < R_L) \leq \frac{\lambda}{(1 + \lambda)(1 + pk)} P_L(T_b^- < R_L) + (2 + \lambda - \delta)(1 + \lambda/2)^{b-L}.$$

Hence for  $k$  sufficiently large, we have

$$P_L(T_b^- < R_L) \leq (2 + \lambda)(1 + \lambda/2)^{b-L}.$$

□

**Lemma 2.4.** Let  $b = \epsilon L$  and  $S = \frac{1}{(2+\lambda)2k} (1 + \lambda/2)^{L(1-2\epsilon)}$

$$P_{L,1} \left( \inf_{t \leq S} \xi_t \leq b \right) \leq (3 + \lambda)(1 + \lambda/2)^{-L\epsilon}.$$

*Proof.* Let  $M = (1 + \lambda/2)^{L(1-2\epsilon)}$ . By Lemma 2.3 the probability that the chain fails to return  $M$  times to  $L$  before going below  $\epsilon L$  is

$$\leq (2 + \lambda)(1 + \lambda/2)^{-L\epsilon}.$$

Using Chebyshev's inequality on the sum  $S_M$  of  $M$  exponentials with mean 1 (and hence variance 1),

$$P(S_M < M/2) \leq 4/M.$$

When the number of infected leaves is  $\leq L$  maximum jump rate is  $D \leq (2 + \lambda)k$  so

$$P \left( \frac{S_M}{(2 + \lambda)k} \leq \frac{(1 + \lambda/2)^{L(1-2\epsilon)}}{2(2 + \lambda)k} \right) \leq 4(1 + \lambda/2)^{-L(1-2\epsilon)}.$$

Adding up the error probabilities completes the proof. □

Up to this point we have shown that if a star has  $L$  infected leaves it will remain infected for a long time. To make this useful, we need estimates about what happens when the star starts with only the center infected. Let  $T_{0,0}$  be the first time the star is healthy. We use the pair  $(n, i)$  to denote the state of the star graph, where  $n$  is the number of infected leaves and  $i$  indicates the state of the center ( $i = 1$  means the center is infected).

**Lemma 2.5.** *Let  $\lambda > 0$  be fixed and  $K = \lambda k^{1/3}$ . Then for large  $k$*

$$\begin{aligned} P_{0,1}(T_K^+ > T_{0,0}) &\leq 2\lambda k^{-1/3}, \\ P_{K,1}(T_{0,0} < T_L^+) &\leq k^{-1/3}, \\ E_{0,1}(T_L^+ | T_L^+ < T_{0,0}) &\leq 2/\lambda. \end{aligned}$$

*Proof.* Clearly

$$P_{0,1}(T_K^+ < T_{0,0}) \geq \prod_{j=0}^{K-1} \frac{(k-j)\lambda}{1 + (k-j)\lambda + j}$$

so subtracting the last inequality from  $1 = \prod_{j=0}^{K-1} 1$  and using Lemma 3.4.3 from [8]

$$P_{0,1}(T_K^+ > T_{0,0}) \leq \sum_{j=0}^{K-1} \frac{1+j}{(k-j)\lambda} \leq \frac{\lambda^2 k^{2/3}}{(k - \lambda k^{1/3})\lambda} \leq 2\lambda k^{-1/3}.$$

For the second result we use the supermartingale  $e^{\theta Y_n}$  from Lemma 2.1. If  $q = P_{K,1}(T_{0,0} < T_L^+)$ , using optional stopping theorem we have

$$q \cdot 1 + (1-q)e^{\theta L} \leq e^{\theta K}.$$

Dropping the second term on the left,

$$q \leq e^{\theta K} = (1 + \lambda/2)^{-K} \leq k^{-1/3}.$$

To bound the time we return to continuous time

jump	at rate
$Y_t \rightarrow Y_t - 1$	$pk$
$Y_t \rightarrow \min\{Y_t + 1, pk\}$	$\lambda(1-p)k$
$Y_t \rightarrow Y_t - N$	$1$

Before time  $V_L = T_{0,0} \wedge T_L^+$  the drift of  $Y_t$  is at least

$$\mu = \lambda(1-p)k - pk - 1/\lambda = \lambda pk - 1/\lambda \tag{8}$$

so  $Y_t - \mu t$  is a submartingale. Stopping this martingale at the bounded stopping time  $V_L \wedge t$

$$EY(V_L \wedge t) - \mu E(V_L \wedge t) \geq EY_0 \geq 0.$$

Since  $EY(V_L \wedge t) \leq L$ , it follows that

$$E(V_L \wedge t) \leq \frac{L}{\mu} = \frac{pk}{\lambda pk - 1/\lambda},$$

where  $p = \lambda/(1 + 2\lambda)$ , so if  $\lambda$  is fixed and  $k$  is large

$$E(V_L \wedge t) \leq 2/\lambda$$

which completes the proof. □

Combining Lemmas 2.4 and 2.5 we have the following

**Lemma 2.6.** *Let  $A_t$  denote the number of infected leaves at time  $t$  and take  $S$  as in Lemma 2.4. Define  $G = \{\inf_{k^{2/3} \leq t \leq S} |A_t| \geq \epsilon L\}$ . If  $\lambda > 0$  is fixed*

$$P_{0,1}(G) \geq 1 - C_\lambda k^{-1/3} \quad (9)$$

for some constant  $C_\lambda$ .

When  $G$  occurs, we say the star at 0 is good.

### 3 Proofs of results for Galton-Watson trees

In the previous section we developed estimates for the contact process on stars. The next step is to obtain estimates on the probability of “pushing an infection from one star to another.” When  $\lambda > 0$  is fixed we have to be careful not to lose too much.

**Lemma 3.1.** *Let  $v_0, v_1, \dots, v_r$  be a path in the graph and suppose that  $v_0$  is infected at time 0. Then there is a  $\gamma > 0$  so that the probability that  $v_r$  will become infected by time  $2r$  is*

$$\geq \left( \frac{\lambda}{\lambda + 1} \right)^r (1 - \exp(-\gamma r)).$$

If  $\epsilon > 0$  and we let  $\hat{\lambda} = (1 - \epsilon)\lambda/(\lambda + 1)$  then for large  $r$  this probability is  $\geq \hat{\lambda}^r$ .

*Proof.* The probability that  $v_{i-1}$  infects  $v_i$  before it is cured is  $\lambda/(1 + \lambda)$ . When this transfer of infection occurs the amount of time is  $t_i$  exponential with rate  $1 + \lambda$ . By large deviations for the exponential distribution  $P(t_1 + \dots + t_r > 2r) \leq e^{-\gamma r}$  for some  $\gamma > 0$ .  $\square$

**Lemma 3.2.** *Run the contact process on a graph consisting for a star of size  $k$  to which there has been added a single chain  $v_1, \dots, v_r$  of length  $r$  where  $v_1$  is a neighbor of 0, the center of the star. Suppose that at time 0 there are  $L$  infected leaves. For large  $r$  the probability  $v_r$  will not be infected before time  $T = m(2r + 1)$  is*

$$\leq (1 - \hat{\lambda}^r)^m.$$

*Proof.* Consider a sequence of times  $t_i = (2r + 1)i$  for  $i \geq l$ . 0 may not be infected at time  $t_i$  but since the star at 0 is good the number of infected neighbors is  $\geq \epsilon L$  and it will with high probability be infected by time  $t_i + 1$ . By Lemma 3.1 the probability  $v_r$  is successfully infected in  $[t_i, t_{i+1})$  is  $\geq (c\hat{\lambda})^r$  when 0 is good, even if we condition on the events up to time  $t_i$ . The desired result follows.  $\square$

**Remark.** Due to the way the proof is done, if we condition on 0 being good then success on two different chains are independent events.

To prepare for the proof of the main results we need the next lemma. Let  $\varphi(x) = \sum_{n=0}^{\infty} p_n x^n$  be the generating function of the Galton-Watson tree. We will apply the result to

$$f(t) = P(0 \in \xi_t) \geq p_k P(0 \in \xi_t | 0 \text{ has at least } k \text{ children}).$$

**Lemma 3.3.** *Let  $H$  be any nondecreasing function on the nonnegative reals with  $H(x) \geq x$  when  $x \in [0, x_0]$ . If  $f$  satisfies (i)  $\inf_{0 \leq t \leq L} f(t) > 0$  and (ii)  $f(t) \geq H(\inf_{0 \leq s \leq t-L} f(s))$  for  $t \geq L$  some  $L > 0$  then  $\liminf_{t \rightarrow \infty} f(t) > 0$ .*

*Proof.* For any  $t_0$  and  $\epsilon > 0$ , (ii) implies that there is a decreasing sequence  $t_i$  with  $t_{i+1} \leq t_i - L$  and  $t_k < L$  for some  $k$

$$f(t_i) \geq H(f(t_{i+1})) - \epsilon 2^{-i}.$$

If  $f(t_i) < x_0$  for all  $1 \leq i \leq k$  then

$$f(t_i) \geq f(t_{i+1}) - \epsilon 2^{-i}$$

and summing gives  $f(t_0) > f(t_k) - \epsilon$  which gives the desired result. Suppose now that  $j$  is the smallest index with  $f(t_j) > x_0$ . If  $j = 0$  we have  $f(t_0) > x_0$ . If  $j = 1$  we have  $f(t_0) \geq H(x_0)$ . If  $j \geq 2$  we have

$$f(t_0) \geq f(t_{j-1}) - \epsilon \geq H(x_0) - \epsilon$$

so in all cases we get the desired conclusion.  $\square$

**Proof for  $p_n = 2^{-n}$ ,  $n \geq 1$ .** The mean of the offspring distribution 2. Let  $Z_r$  be the number of vertices at distance  $r$  from 0 and let  $v_r^1, \dots, v_r^J$  be those that have exactly  $k$  children, where  $J$  is a random variable that represents the number of such vertices.

Since the root has degree  $k$  and  $p_k = 2^{-k}$  if we set  $r = k$

$$EJ \geq k\mu^{r-1}p_k = k/2,$$

where  $\mu = 2$  is the mean offspring number.

If we condition on the value of  $W = Z_r/(k\mu^{r-1})$  and let  $\bar{J} = (J|W)$  then

$$\bar{J} = \text{Binomial}(k2^{r-1}W, 2^{-k}).$$

Let  $M$  be the random number of vertices among  $v_r^1, \dots, v_r^J$  that are infected before time

$$S = \frac{1}{2k(2 + \lambda)}(1 + \lambda/2)^{L(1-2\epsilon)}$$

defined in Lemma 2.4. By Lemma 3.2 the probability a given vertex will not become infected is (recall  $r = k$ )

$$\leq (1 - \hat{\lambda}^k)^{\lceil S/(2k+1) \rceil} \leq \exp(-(2k(2k+1)(2+\lambda))^{-1} \hat{\lambda}^k (1 + \lambda/2)^{(1-2\epsilon)\lambda k/(1+2\lambda)})$$

When  $\lambda = 2.5$

$$\frac{\lambda}{\lambda + 1} (1 + \lambda/2)^{\lambda/(1+2\lambda)} = 1.0014$$

and it follows that if  $k$  is large

$$EM \geq (1 - \delta)EJ.$$

The remark after Lemma 3.2 implies that if we condition on the value of  $W$  and let  $\bar{M} = (M|W)$  then  $\bar{M} \geq \text{Binomial}(k2^{r-1}W, 2^{-k}(1 - \delta))$ .



Let

$$H_1(t) = P(v_r^i \in \xi(t - S) \text{ for some } 1 \leq i \leq J)$$

and

$$H_2(t) = P(0 \in \xi(t) | v_r^i \in \xi(t - S) \text{ for some } 1 \leq i \leq J),$$

so that  $f(t) \geq H_1(t)H_2(t)$ .

Fix  $t > 2S$  and let

$$\chi(t) = \inf\{f(s) : s \leq t - S\}.$$

For simplicity in notation we will write  $\chi(t)$  as  $\chi$ .

Ignore all but the first infection of each  $v_r^i$  by its parent, any of these will evolve independently from the time  $s < S$  it is first infected, and will be infected at time  $t - S$  with probability at least  $\chi$ . Thus given  $M$  the number of infected at time  $t - S$  will dominate  $N = \text{binomial}(M, \chi)$ . If we let  $\bar{N} = \text{binomial}(\bar{M}, \chi)$  and take a small  $\delta > 0$ , then by Lemma 2.3 in [28] we see that there exists a  $\varepsilon > 0$  such that

$$P(\bar{N} \geq 1) \geq (1 - \delta)\chi EM \wedge \varepsilon$$

Therefore  $H_1(t) \geq (1 - \delta)\chi EM \wedge \varepsilon$  when  $t > 2S$ .

Finally, if some  $v_r^i$  is infected at time  $t - S$  then the probability of finding 0 infected at time  $t$  is bounded below by  $\rho_1\rho_2$  where

- $\rho_1$  is the probability that the contact process starting with only  $v_r^i$  infected at time  $t - S$  infects 0 at some time  $s$  with  $t - S \leq s \leq t$ . By Lemmas 2.5, 2.6, and 3.2,  $\rho_1 \geq 1 - \delta$ .
- $\rho_2$  is the probability 0 is infected at time  $t$  given the infection of 0 at such a time  $s$ . By Lemma 2.5 and 2.6, at time  $t - \epsilon$  there should be at least  $\epsilon L$  infected leaves. Hence 0 is infected at  $t$  with probability at least  $(1 - e^{-\lambda\epsilon^2 L})e^{-\epsilon}$ .

So  $\rho_2 \geq 1 - \delta$  when we pick  $\epsilon$  sufficiently small. Thus

$$f(t) \geq \begin{cases} \chi(t)EM(1 - \delta)^3 \wedge \varepsilon & t > 2S, \\ \inf_{0 \leq s \leq 2S} f(s) & S \leq t \leq 2S. \end{cases}$$

We can take  $\varepsilon < \inf_{0 \leq s \leq 2S} f(s)$  so that  $f(t) \geq \chi(t)EM(1 - \delta)^3 \wedge \varepsilon$  for all  $t \geq S$ . The result now follows from Lemma 3.3 with  $L = S$  and  $H(x) = (1 - \delta)^3(EM)x \wedge \varepsilon$ .

**Proof for**  $p_n = (1 - p)^{n-1}p$ . The mean of the offspring distribution is  $1/p$ . Let  $Z_r$  be the number of vertices at distance  $r$  from 0 and let  $v_r^1, \dots, v_r^J$  be those that have exactly  $k$  children. Since the root has degree  $k$  and  $p_k = (1 - p)^{k-1}p$

$$EJ \geq k(1/p)^{r-1}(1 - p)^{k-1}p.$$

In this case we want to pick  $r$  so that  $(1/p)^r(1 - p)^k \approx 1$ . Ignoring the fact that  $r$  and  $k$  must be integers this means

$$r/k = \log(1 - p)/\log p.$$

Let  $M$  be the random number of vertices among  $v_r^1, \dots, v_r^J$  that are infected before time  $S$ . By Lemma 3.2 the probability a given vertex will not become infected is

$$\leq (1 - \hat{\lambda}^r)^{\lceil S/(2r+1) \rceil} \leq \exp(-(2k(2r+1)(2+\lambda))^{-1} \hat{\lambda}^r (1 + \lambda/2)^{(1-2\epsilon)\lambda k/(1+2\lambda)}).$$

We will get a lower bound on  $\lambda_2$  if

$$\left(\frac{\lambda}{\lambda+1}\right)^{r/k} \cdot (1 + \lambda/2)^{\lambda/(1+2\lambda)} > 1. \quad (10)$$

If we want to graph the bound as a function of  $p$  it is better to work backwards. Given  $\lambda$  the second factor is  $> 1$  so we can easily find the value of  $r/k$  that makes this 1. Having done this we can easily compute the value of  $p$  for which  $\lambda$  gives the upper bound on  $\lambda_2$ .

**Proof for subexponential distributions.** We suppose that the mean of the offspring distribution is  $\mu > 1$ . If  $p_k$  is subexponential, i.e.,

$$\limsup_{k \rightarrow \infty} (1/k) \log p_k = 0,$$

then for any  $\delta$  there is a  $k$  with  $p_k \geq (1 - \delta)^k$  then we can take

$$\frac{r}{k} = -\frac{\log(1 - \delta)}{\log \mu}.$$

Given any  $\lambda > 0$ , (10) will hold if  $\delta$  is small enough, so  $\lambda_2 = 0$ .

## 4 Asymptotics for $\lambda_c$

We begin with some general computations and then consider our two examples: power laws and stretched exponential.

**Survival on star graph.** Our first step is to adapt Lemma 2.4 to the situation in which  $\lambda \rightarrow 0$ . For reasons that will become clear when we prove Lemma 4.3 we have to modify the definition of  $p$ :

$$p = (1 - \epsilon) \frac{\lambda}{1 + \lambda}, \quad L = pk, \quad \text{and} \quad b = \epsilon L.$$

Defining  $Y_n$  as before

**Lemma 4.1.** *Let  $\epsilon > 0$ . If  $\lambda/(1 + 2\lambda) < \epsilon$  then  $(1 + \lambda/2)^{-Y_n}$  is a supermartingale.*

*Proof.*  $(1 - p) = (1 + \lambda\epsilon)/(\lambda + 1)$  so we have

$$\frac{p}{\lambda(1 - p)} = \frac{1 - \epsilon}{1 + \lambda\epsilon}.$$

The right-hand side is  $< 1/(1 + \lambda)$  when

$$1 + \lambda - \epsilon - \epsilon\lambda < 1 + \lambda\epsilon,$$

which holds if  $\lambda/(1 + 2\lambda) < \epsilon$ , so the desired result follows from the proof of Lemma 2.1.  $\square$

**Lemma 4.2.** *Let  $\epsilon > 0$  be fixed. If  $k \rightarrow \infty$  and  $\lambda \rightarrow 0$  then for large  $k$*

$$P_{L,1} \left( \inf_{t \leq T} \xi_t \leq b \right) \leq 4 \exp(-(1-3\epsilon)\lambda^2 k/4),$$

where  $T = \exp((1-4\epsilon)\lambda^2 k/4)$ .

*Proof.* It follows from Lemma 2.4 that if  $S = (1/2k(2+\lambda))(1+\lambda/2)^{L(1-2\epsilon)}$  then

$$P_{L,1} \left( \inf_{t \leq S} \xi_t \leq b \right) \leq (3+\lambda)(1+\lambda/2)^{-L(1-2\epsilon)},$$

but now

$$(1-\epsilon)L = (1-\epsilon)^2 \lambda k / (\lambda+1) > (1-2\epsilon) \lambda k / (\lambda+1).$$

Expanding  $\log(1+x) = x - x^2/2 + x^3/3 - \dots$  and noting that if  $x < 1$  then the right-hand side is an alternating series with decreasing terms

$$\begin{aligned} (1+\lambda/2)^{-(1-2\epsilon)\lambda k/(1+\lambda)} &= \exp \left( -(1-2\epsilon) \frac{\lambda k}{1+\lambda} \log(1+\lambda/2) \right) \\ &\leq \exp \left( -(1-2\epsilon) \frac{\lambda k}{1+\lambda} \left[ \frac{\lambda}{2} - \frac{\lambda^2}{8} \right] \right) \\ &\leq \exp(-(1-3\epsilon)\lambda^2 k/4), \end{aligned}$$

when  $\lambda$  is small. To convert the formula for  $S$  we note that

$$\begin{aligned} (1/2k(2+\lambda))(1+\lambda/2)^{L(1-2\epsilon)} &= \frac{1}{2k(2+\lambda)} \exp \left( \frac{(1-\epsilon)(1-2\epsilon)\lambda k}{(\lambda+1)} \cdot \log(1+\lambda/2) \right) \\ &\geq \frac{1}{6k} \exp \left( \frac{(1-3\epsilon)\lambda k}{(\lambda+1)} \cdot \left[ \frac{\lambda}{2} - \frac{\lambda^2}{8} \right] \right) \\ &\geq \exp((1-4\epsilon)\lambda^2 k/4), \end{aligned}$$

when  $\lambda$  is small, which completes the proof.  $\square$

**Push.** Now we work with the configuration model. Let  $p_k = P(d(x) = k)$  and suppose that

- (i)  $\sum_k k^2 p_k < \infty$ ,
- (ii)  $P(d(x) = k) = 0$  for  $k \leq 2$ .

The first assumption implies that the size biased degree distribution  $q_{j-1} = j p_j / E d(x)$  has finite mean  $\nu$ . The second implies that the diameter of our graph  $\sim (\log n) / \log \nu$ . See Lemma 3.4.1 in [7]. Hence Lemma 3.1 implies that if  $v_0, \dots, v_r$  is a path in the graph and  $v_0$  is infected at time 0, then the probability  $v_r$  will be infected by time  $2r$  is, for large  $r$ ,  $\geq (\lambda/2)^r$ . Let

$$\kappa = n^{3\nu \log(2/\lambda)}.$$

If  $n$  is large then the distance between any two vertices is  $\leq 2\nu \log n$ . Thus the probability that one star can transfer its infection to another before time  $2r\kappa$  is

$$\begin{aligned} &\geq 1 - (1 - (\lambda/2)^{2\nu \log n})^\kappa = 1 - (1 - n^{-2\nu \log(2/\lambda)})^\kappa \\ &\geq 1 - \exp(-n^{\nu \log(2/\lambda)}). \end{aligned} \tag{11}$$

**Ignition on star graph.** We have more work to do this time. The proof of Lemma 2.5 requires that  $K = \lambda k^{2/3} \rightarrow \infty$ , and we need the new definition of  $L$  in part (iii).

Recall that  $T_m^+ = \inf\{n : Y_n \geq m\}$ .

**Lemma 4.3.** *Let  $K = \lambda k / \sqrt{\log k}$ . If  $\lambda \rightarrow 0$  and  $\lambda^2 k \rightarrow \infty$  then for large  $k$*

$$\begin{aligned} (i) P_{0,1}(T_K^+ > T_{0,0}) &\leq 5/\sqrt{\log k}, \\ (ii) P_{K,1}(T_{0,0} < T_L^+) &\leq \exp(-\lambda^2 k / 2\sqrt{\log k}), \\ (iii) E_{0,1} \min\{T_{0,0}, T_L^+\} &\leq 2/\epsilon. \end{aligned}$$

*Proof.* Let  $p_0(t)$  be the probability a leaf is infected at time  $t$  when there are no infected leaves at time 0 and the central vertex has been infected for all  $s \leq t$ .  $p_0(0) = 0$  and

$$\frac{dp_0(t)}{dt} = -p_0(t) + \lambda(1 - p_0(t)) = \lambda - (\lambda + 1)p_0(t).$$

Solving gives

$$p_0(t) = \frac{\lambda}{\lambda + 1}(1 - e^{-(\lambda+1)t}).$$

As  $t \rightarrow 0$

$$\frac{1 - e^{-(\lambda+1)t}}{(\lambda + 1)t} \rightarrow 1,$$

so if  $t$  is small  $p_0(t) \geq \lambda t / 2$ .

Taking  $t = 4/\sqrt{\log k}$  it follows that the number of infected leaves at time  $t$  dominates  $B = \text{Binomial}(k, 2\lambda/\sqrt{\log k})$

$$P_{0,1}(T_K^+ < T_{0,0}) \geq P(B > K) \exp(-4/\sqrt{\log k}).$$

The second factor is the probability that the center stays infected until time  $4/\sqrt{\log k}$ , and

$$\exp(-4/\sqrt{\log k}) \geq 1 - 4/\sqrt{\log k}.$$

$B$  has mean  $2\lambda k / \sqrt{\log k}$  and variance  $\leq 2\lambda k / \sqrt{\log k}$  so Chebyshev's inequality implies

$$P(B < \lambda k / \sqrt{\log k}) \leq \frac{2\lambda k / \sqrt{\log k}}{(\lambda k / \sqrt{\log k})^2} \leq \frac{2\sqrt{\log k}}{\lambda k} \leq 1/\sqrt{\log k}$$

if  $k$  is large.

For (ii) we use the supermartingale from Lemma 4.1, which is the same as the one from Lemma 2.1, and simplify formulas as in the proof of Lemma 4.2. If  $q = P_{K,1}(T_{0,0} < T_L^+)$  then for  $\lambda$  small optional stopping theorem gives

$$q \leq (1 + \lambda/2)^{-\lambda k / \sqrt{\log k}} \leq \exp(-\lambda^2 k / 2\sqrt{\log k}).$$

For (iii), we follow the argument in Lemma 2.5. We return to continuous time and note that by (8) the drift is

$$\leq \mu = \lambda(1 - p)k - pk - 1/\lambda$$

so  $Y_t - \mu t$  is a submartingale before time  $V_L = T_{0,0} \wedge T_L^+$ . Using the optional stopping theorem as before we conclude

$$E(V_L) \leq \frac{L}{\mu} = \frac{(1-\epsilon)\lambda k}{1+\lambda} \cdot \frac{1}{\lambda(1-p)k - pk - 1/\lambda}.$$

Recalling the definition of  $p$

$$\begin{aligned} \lambda(1-p)k - pk &= \lambda \left[ k - \frac{(1-\epsilon)\lambda k}{1+\lambda} \right] - \frac{(1-\epsilon)\lambda k}{1+\lambda} - \frac{1}{\lambda} \\ &= \lambda \left[ \frac{\epsilon\lambda k}{1+\lambda} \right] + \frac{\epsilon\lambda k}{1+\lambda} - \frac{1}{\lambda}. \end{aligned}$$

The first term is much smaller than the second so multiplying by  $\lambda/\lambda$

$$\frac{L}{\mu} \sim \frac{\lambda^2 k}{\epsilon\lambda^2 k - (1+\lambda)} \sim 1/\epsilon,$$

since  $\lambda k^2 \rightarrow \infty$ . □

## 4.1 Power law graphs

Suppose  $P(d(x) \geq m) = 3^a m^{-a}$  for  $m \geq 3$ , where  $a > 2$  so that  $Ed(x)^2 < \infty$ . In this case, the maximum degree vertex on a graph with  $n$  vertices is  $\sim n^{1/a}$ , so the maximum eigenvalue  $\Lambda \sim n^{1/2a}$  and the formula in (5) predicts that  $\lambda_c \approx n^{-1/2a}$ . To prove an upper bound on  $\lambda_c$  that is close to this, we suppose that  $\lambda_0 = n^{-(1-2\eta)/2a}$ .

If  $d(x) \geq k = n^{(1-\eta)/a}$  we call the vertex  $x$  a star.

$$P(d(x) \geq n^{(1-\eta)/a}) = 3^a n^{-(1-\eta)}$$

so if  $n$  is large there are  $\geq n^\eta$  stars with high probability. Now  $\lambda_0^2 k = n^{\eta/a}$ . By the estimate in the Lemma 4.2, each individual star survives for time

$$\geq \exp((1-4\eta)n^{\eta/a}/4). \tag{12}$$

with probability  $\geq 1 - 7\exp(-(1-3\epsilon)n^{\eta/a}/4)$ . The time

$$2r\kappa \leq (4\nu \log n) \exp(O(\log^2 n))$$

so (11) implies that with high probability the chosen star will transfer its infection to its target by time  $2r\kappa$  and we conclude that with high probability no lit star will die out during the process.

Combining these estimates shows that if  $n$  is large then the number of infected stars  $Y_k$  at time  $2r\kappa k$  dominates a discrete time random walk that goes up by 1 with probability  $p > e/(e+1)$  and down by 1 with probability  $1-p$ . Let  $M \geq n^\eta$  be the number of stars. Recalling that  $((1-p)/p)^x$  is harmonic function for a simple random walk that jumps up 1 with probability  $p$ , and down 1 with probability  $1-p$  random walk, we see that  $\exp(-Y_k)$  is a supermartingale while  $Y_k \in (0, M)$ , so

$$P_{0.9M}(T_0 < T_M) \leq e^{-0.9M}.$$

Since each cycle takes at least  $0.1M(2r\kappa)$  units of time, we have survival for time  $\exp(O(n^\epsilon))$  for some  $\epsilon > 0$ .

## 4.2 Stretched exponential

Suppose  $P(d(x) \geq m) = \exp(-m^{1/b} + 3^{1/b})$  for  $m \geq 3$ , where  $b > 1$ . In this case, the maximum degree vertex on a graph with  $n$  vertices is  $\sim \log^b n$ , so the maximum eigenvalue  $\Lambda \sim \log^{b/2} n$  and the formula in (5) predicts that  $\lambda_c \approx \log^{-b/2} n$ .

If  $d(x) \geq k = \eta^b \log^b n$  we call the vertex  $x$  a star.

$$P(d(x) \geq \eta^b \log^b n) = \exp(3^{1/b})n^{-\eta},$$

so if  $n$  is large then the number of stars is  $\geq n^{1-\eta}$  with high probability.

To see what value to take for  $\lambda$  in our lower bound, we set the survival time equal to 1 over the probability of a successful push, that is

$$\exp(\lambda^2 \log^b n) = (2/\lambda)^{2\nu \log n},$$

or taking logs and rearranging

$$\frac{\lambda^2}{\log(2/\lambda)} = 2\nu \log^{1-b} n.$$

This means that the best upper bound we can hope to get is  $\lambda_0 = (\log n)^{(1-\eta)(1-b)/2}$  versus the predicted value of  $\log^{-b/2} n$ .

By our choices we have

$$\lambda_0^2 k = \eta^b (\log n)^{1+\eta(b-1)}$$

so Lemma 4.2 implies that the star survives for time

$$\geq \exp((1 - 4\eta)\eta^b (\log n)^{1+\eta(b-1)}/4)$$

with probability  $\geq 1 - 7 \exp(-(1 - 3\eta)\eta^b (\log n)^{1+\eta(b-1)}/4)$ . The time

$$2r\kappa \leq (4\nu \log n) \exp(\log n \cdot O(\log \log n))$$

so (11) implies that with high probability the chosen star will transfer its infection to its target by time  $2r\kappa$  and we conclude that with high probability no lit star will die out during the process. Comparing with random walk as in the previous proof, we have survival for time  $\exp(O(n^{1-\epsilon}))$  for any  $\epsilon > 0$ .

## References

- [1] Bezuidenhout, C. and Grimmett, G. (1990) The critical contact process dies out. *Ann. Probab.* 18, 1462-1482
- [2] Castellano, C., and Pastor-Satorras, R. (2010) Thresholds for epidemic spreading in networks. *Physical Review Letters*. 105, paper 218701
- [3] Chakrabarti, D., Wang, Y., Wang, C., Leskovec, J., and Faloutsos, C. (2008) *ACM Transactions on Information and System Security*. Vol. 10, No. 4, article 13

- [4] Chatterjee, S., and Durrett, R. (2009) Contact processes on random graphs with power law degree distributions have critical value 0. *Ann. Probab.* 37, 2332-2356
- [5] Chung, F., Lu, L., and Vu, Van. (2003) Spectra of random graphs with given degrees. *Proc. Natl. Acad. Sci.* 100, 6313–6318
- [6] Cranston, M., Mountford, T., Mourrat, J.C., and Valesin, D. (2014) The contact process on finite trees revisited. *ALEA Lat. Am. J. Probab. Math. Stat.* 11, 385-408
- [7] Durrett, R. (2007) *Random Graph Dynamics*. Cambridge U. Press
- [8] Durrett, R. (2010) *Probability: Theory and Examples*. Fourth Edition, Cambridge U Press
- [9] Durrett, R., and Jung, R. (2007) Two phase transitions for contact processes on small worlds. *Stoch. Proc. Appl.* 117, 1910–1927
- [10] Durrett, R., and Liu, Xiu-fang. (1988) The contact process on a finite set. *Ann Probab.* 16, 1158–1173
- [11] Durrett, R and Schonmann, R.H.S. (1988) The contact process on a finite set, II. *Ann. Probab.* 16, 1570-1583
- [12] Ganesh, A., Massoulié, I., and Towsley, D. (2005) The effect of network topology on the spread of epidemics. *Proceedings of the 24th Annual Joint Conference of the IEEE Computer and Communication Societies (INFOCOM)*. 2, 1455–1466
- [13] Harris, T.E. (1974) Contact interactions on a lattice. *Ann. Probab.* 2, 969–988
- [14] Jiang, Y., Kassem, R., York, G., Zhao, B., Huang, X., Junge, M., and Durrett, R. The Contact Process on Periodic Trees arXiv:1808.01863
- [15] Liggett, T. M. (1995) Improved upper bounds for the contact process critical value. *Ann. Probab.* 23, 697-723
- [16] Liggett, T.M. (1996) Multiple transition points for the contact process on the binary tree. *Ann. Probab.* 24, 1455–1490
- [17] Liggett, T.M. (1999) *Stochastic Interacting Systems: Contact, Voter and Exclusion Processes*. Springer-Verlag, New York
- [18] Ménard, L., and Singh, A. (2016) Percolation by cumulative merging and phase transition for the contact process on some random graphs. *Ann. Sci. École Norm. Supér.* 49, 1189–1238
- [19] Mountford, T. S. (1993) A metastable result for the finite multidimensional contact process. *Canad. Math. Bull.* 36, 216-226.
- [20] Mountford, T. S. (1999) Existence of a constant for finite system extinction. *J. Statist. Phys.* 96, 1331-1341.

- [21] Mountford, T., Mourrat, J-C, Valesin, D., and Yao, Q. (2016) Exponential extinction time of the contact process on finite graphs. *Stochastic Process. Appl.* 126, 1974-2013
- [22] Mountford, T., Valesin, D., and Yao, Q. (2013) Metastable densities for the contact process on power law random graphs. *Electron. J. Probab.* 18, paper 103
- [23] Mourrat, J-C, and Valesin, D. (2016) Phase transtions of the contact process on random regular graphs. *Electron. J. Probab.* 21, paper 31
- [24] Pastor-Satorras, R., Castellano, C., van Mieghem, and Vespignani, A (2015) Epidemic processes on complex networks. *Reviews of Modern Physics.* 87, 925–979
- [25] Pastor-Satorras, R. and Vespignani, A. (2001a) Epidemic spreading in scale-free networks. *Phys. Rev. Letters.* 86, 3200–3203
- [26] Pastor-Satorras, R. and Vespignani, A. (2001b) Epidemic dynamics and endemic states in complex networks. *Phys. Rev. E.* 63, 066117
- [27] Pastor-Satorras, R. and Vespignani, A. (2002) Epidemic dynamics in finite size scale-free networks. *Phys. Rev. E.* 65, 035108(R)
- [28] Pemantle, R. (1992) The contact process on trees. *Ann. of Probab.* 20, 2089–2116
- [29] Pemantle, R. and Stacey, A. (2001) The branching random walk and contact process on Galton-Watson and nonhomogeneous trees. *Ann. Probab.* 29, 1563-1590
- [30] Penrose, M. (2003) *Random Geometric Grpahs.* Oxford U. Press
- [31] Stacey, A. (2001) The contact process on finite homogeneous trees. *Probab. Theory Related Fields.* 121, 551-576
- [32] Stacey, A. M. (1996) The existence of an intermediate phase for the contact process on trees. *Ann. Probab.* 24, 1711-1726.
- [33] Wang, Y., Chakrabarti, D. Wang, C., and Faloutsos, C. (2003) Epidemic spreading in networks: An eigenvalue viewpoint. *22nd International Symposium on Reliable Systems.* page 25