

Unexpected curves and Togliatti-type surfaces

Justyna Szpond

May 15, 2019

Abstract

The purpose of this note is to establish a direct link between the theory of unexpected hypersurfaces and varieties with defective osculating behavior. We identify unexpected plane curves of degree 4 as sections of a rational surface X_B of degree 7 in \mathbb{P}^3 with its osculating spaces of order 2 which in every point of X_B have dimension lower than expected. We put this result in perspective with earlier examples of surfaces with defective osculating spaces due to Shifrin and Togliatti. Our considerations are rendered by an analysis of Lefschetz Properties of ideals associated with the studied surfaces.

1 Introduction

In the ground-breaking paper [2] the authors introduced the notion of unexpected curves.

Definition 1. We say that a reduced finite set of points $Z \subset \mathbb{P}^2$ admits an unexpected curve of degree $j + 1$ if there is an integer $j > 0$ such that, for a general point P , jP fails to impose the expected number of conditions on the linear system of curves of degree $j + 1$ containing Z . That is, Z admits an unexpected curve of degree $j + 1$ if

$$h^0(\mathcal{J}_{Z+jP}(j+1)) > \max \left\{ h^0(\mathcal{J}_Z(j+1)) - \binom{j+1}{2}, 0 \right\}.$$

This notion has been generalized to hypersurfaces in projective spaces of arbitrary dimension in the sequel paper [9].

One of the first examples of unexpected curves (in fact this example has motivated this path of research) was discovered in [5, Proof of Proposition 7.3]. The recent work [6] shows that it is the unique example of an unexpected quartic. This example is derived from the B_3 root system, see [9, Example 4.6] and [1, section 2].

On the other hand the study of osculating spaces of projective varieties has a long history, going back to Corrado Segre and Gaston Darboux. A modern treatment has been given in [15].

Definition 2. Let $X \subseteq \mathbb{P}^N$ be a smooth, complete, complex variety. Let $m \geq 1$ be an integer. The m -th osculating space to X at P is the linear subspace $\text{Osc}_P^{(m)}(X)$ determined at a point $P \in X$ by the partial derivatives of order $\leq m$ of the coordinate functions, with respect to a system of local parameters for X at P , evaluated at P .

In particular, $\text{Osc}_P^{(1)}(X)$ is just the tangent space of X at P .

A somewhat more formal, equivalent description is the following.

Keywords: fat points, Lefschetz properties, linear systems, postulation problem, SHGH conjecture, Togliatti system, unexpected hypersurfaces.

Mathematics Subject Classification (2010): 14C20 14J26 14N05 13A15 13E10 13F20.

Let X be a smooth, complete, complex variety of dimension n and let L be a line bundle on X . Let $V = H^0(X, L)$. The m -jet bundle of L is the coherent sheaf

$$\mathcal{J}_m(L) = (p_1)_*(p_2^*L \otimes \mathcal{O}_{X \times X}/\mathcal{J}_\Delta^{m+1}),$$

where $\Delta \subseteq X \times X$ is the diagonal and p_1, p_2 are the projection maps

$$\begin{array}{ccc} & X \times X & \\ p_1 \swarrow & & \searrow p_2 \\ X & & X \end{array}$$

This sheaf is locally free of rank $\binom{n+m}{n}$ and its fiber at a point P can be identified with

$$\mathcal{J}_m(L)_P \simeq H^0(X, L \otimes \mathcal{O}_X/\mathfrak{m}_P^{m+1}),$$

where \mathfrak{m}_P is the ideal sheaf of $P \in X$.

Then the m -th osculating space at P is the projectivization $\mathbb{P}(j_{k,P}(V)) \subseteq \mathbb{P}(V)$ of the image of the evaluation map

$$j_{k,P} : V \longrightarrow H^0(X, L \otimes \mathcal{O}_X/\mathfrak{m}_P^{m+1}).$$

This agrees with Definition 2 in case $X \subset \mathbb{P}^N$ and $L = \mathcal{O}_X(1)$.

It is natural to expect that the space $\text{Osc}_P^{(m)}(X)$ has at a general point $P \in X$ the (projective) dimension $\binom{n+m}{n} - 1$. If this dimension is lower than $\binom{n+m}{n} - 1$ at every point, then following Shifrin [16] we say that X is *hypo-osculating of order m* . More precisely, the deficiency of osculating spaces of X is measured by the number of Laplace equations X satisfies.

Definition 3 (Laplace equation). A projective variety $X \subset \mathbb{P}^N$ of dimension n satisfies δ independent Laplace equations of order m , if for a general point $P \in X$ there is

$$\dim \text{Osc}_P^{(m)} X = \min \left\{ \binom{n+m}{n} - 1, N \right\} - \delta.$$

In this note we are interested in surfaces $X \subseteq \mathbb{P}^5$. It is then expected that $\dim \text{Osc}_P^{(2)}(X) = 5$ at a general point $P \in X$. It has been noticed, apparently first by Eugenio Togliatti [17] that there are hypo-osculating surfaces of order 2 in \mathbb{P}^5 .

Our main results are the following Theorems.

Theorem A. *Let $Z = \{P_1, \dots, P_9\} \subseteq \mathbb{P}^2$ be the B_3 configuration of points. Let $f : Y \longrightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at points of Z with the exceptional divisors E_i over P_i for $i = 1, \dots, 9$. Let X_B be the image of Y under the morphism defined by $H^0(Y, f^*\mathcal{O}_{\mathbb{P}^2}(4) - E_1 - \dots - E_9)$. Then*

- a) X_B is a surface in \mathbb{P}^5 of degree 7 with 3 singular points.
- b) In all points of the smooth locus $(X_B)_{sm}$, the surface is hypo-osculating with

$$\dim \text{Osc}_P^{(2)}((X_B)_{sm}) = 4.$$

- c) There is a 1:1 correspondence between unexpected curves admitted by Z and second osculating spaces of X_B at its smooth points.

Moreover, the divisor cut out on X_B by the second osculating space at a general point of X_B is irreducible.

Theorem B. *Let X'_B be the image of the morphism*

$$\eta : \mathbb{P}^2 \ni (a : b : c) \longrightarrow (a^3 : b^3 : c^3 : a(b^2 - c^2) : b(a^2 - c^2) : c(a^2 - b^2)) \in \mathbb{P}^5.$$

Then

- a) X'_B is a smooth surface of degree 9 in \mathbb{P}^5 .
 b) The surface X'_B is hypo-osculating with

$$\dim \text{Osc}_P^{(2)}(X'_B) = 4$$

for all points $P \in X'_B$, except those in the B_3 configuration, i.e., except the images of points P_1, \dots, P_9 .

Part a) of Theorem A is proved with Proposition 8, part b) is Proposition 9 and part c) is an easy consequence of Lemma 17. Theorem B is proved in Propositions 19 and 20.

2 The Shifrin surface

In this section we recall briefly an example of a surface embedded in \mathbb{P}^5 by a *complete* linear system which satisfies a Laplace equation of order 2. Let $X_S \subseteq \mathbb{P}^5$ be the image of the mapping

$$\varphi : \mathbb{P}^1 \times \mathbb{P}^1 \ni ((s : t), (u, v)) \rightarrow (stv : t^2u : s^2v : stu : s^2u : t^2v) \in \mathbb{P}^5.$$

Let u_0, \dots, u_5 be the coordinates in \mathbb{P}^5 . Then the ideal $I(X_S)$ defining X_S is generated by 6 quadrics

$$\begin{aligned} u_3^2 - u_1u_4, & \quad u_2u_3 - u_0u_4, & \quad u_0u_3 - u_4u_5 \\ u_1u_2 - u_4u_5, & \quad u_0u_1 - u_3u_5, & \quad u_0^2 - u_2u_5 \end{aligned}$$

and its minimal free resolution is

$$S/I(X_S) \leftarrow S \leftarrow S(-2)^{\oplus 6} \leftarrow S(-3)^{\oplus 8} \leftarrow S(-4)^{\oplus 3},$$

where $S = \mathbb{C}[u_0, \dots, u_5]$.

Note that the linear system defining φ is of bidegree $(2, 1)$. If we consider this system directly on $\mathbb{P}^1 \times \mathbb{P}^1$, then we have clearly

$$h^0(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(2, 1)) = 6$$

and for every point $P = (P_1, P_2) \in \mathbb{P}^1 \times \mathbb{P}^1$ there exists a divisor with a triple point there, namely this is

$$D_P = 2 \cdot \{P_1\} \times \mathbb{P}^1 + \mathbb{P}^1 \times \{P_2\}.$$

In particular this divisor is reducible and non-reduced for all points P . Thus, even though the existence of D_P is unexpected from the point of view of counting conditions, it is easily explained by the geometry of the surface.

Property 4. All divisors cut out on the Shifrin surface by 2nd osculating spaces are reducible.

This example has a remarkable property that all its second osculating spaces have the same dimension, which is one less than the expected one. For such surfaces Shifrin introduced in [16] the following terminology.

Definition 5. We say that a surface $X \subseteq \mathbb{P}^5$ is *perfectly* hypo-osculating if

$$\dim \text{Osc}_P^{(2)}(X) = 4$$

for every point $P \in X$.

He formulated a conjecture [16, Conjecture 0.6] to the effect that the surface discussed in this part is the unique smooth perfectly hypo-osculating surface in \mathbb{P}^5 . This conjecture was proved Piene and Tai in [14]. Along these lines we propose a new Problem 23, we relax the assumption of perfectly hypo-osculating and replace it by irreducible hypo-osculating divisors.

3 The Togliatti surface

In this section we discuss the classical example of Togliatti. Let $X_T \subset \mathbb{P}^5$ be the closure of the image of the mapping

$$\psi : \mathbb{P}^2 \ni (x : y : z) \dashrightarrow (x^2y : x^2z : y^2x : y^2z : z^2x : z^2y) \in \mathbb{P}^5.$$

Let, as before, u_0, \dots, u_5 be the coordinates in \mathbb{P}^5 . Then the ideal $I(X_T)$ defining X_T is generated by

$$\begin{aligned} u_2u_4 - u_0u_5, & & u_1u_3 - u_0u_5, & & u_3u_4^2 - u_1u_5^2, & & u_0u_4^2 - u_1^2u_5, & & u_3^2u_4 - u_2u_5^2 \\ u_0u_3u_4 - u_1u_2u_5, & & u_0u_3^2 - u_2^2u_5, & & u_1u_2^2 - u_0^2u_3, & & u_1^2u_2 - u_0^2u_4 \end{aligned}$$

and its minimal free resolution is

$$S/I(X_T) \leftarrow S \leftarrow S(-2)^{\oplus 2} \oplus S(-3)^{\oplus 7} \leftarrow S(-4)^{\oplus 19} \leftarrow S(-5)^{\oplus 16} \leftarrow S(-6)^{\oplus 6} \leftarrow S(-7).$$

For a more geometric description let $P_1 = (1 : 0 : 0)$, $P_2 = (0 : 1 : 0)$ and $P_3 = (0 : 0 : 1)$ be the coordinate points in \mathbb{P}^2 and let $Z = \{P_1, P_2, P_3\}$. Identifying, as usual, global sections with homogeneous polynomials, we may take as a basis of the linear system $V = H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(3) \otimes \mathcal{J}_Z)$ the monomials

$$x^2y, x^2z, y^2x, y^2z, z^2x, z^2y, xyz.$$

Let $f : Y \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at P_1, P_2, P_3 with exceptional divisors E_1, E_2, E_3 respectively, and let $H = f^*\mathcal{O}_{\mathbb{P}^2}(1)$. Then the following holds.

Proposition 6. *The linear system*

$$M = 3H - E_1 - E_2 - E_3$$

is very ample and determines an embedding

$$\varphi_M : Y \rightarrow \mathbb{P}^6,$$

which lifts the rational map

$$\tilde{\psi} : \mathbb{P}^2 \ni (x : y : z) \dashrightarrow (x^2y : x^2z : y^2x : y^2z : z^2x : z^2y : xyz) \in \mathbb{P}^6,$$


i.e., the following diagram commutes

$$\begin{array}{ccc} Y & & \\ f \downarrow & \searrow \varphi_M & \\ \mathbb{P}^2 & \dashrightarrow \tilde{\psi} & \mathbb{P}^6 \end{array}$$

Then the Togliatti surface X_T is the image of $\varphi_M(Y)$ under the projection π from the last coordinate point $(0 : 0 : 0 : 0 : 0 : 0 : 1) \in \mathbb{P}^6$. We have

$$\begin{array}{ccc} Y & \xrightarrow{\varphi_M} & \mathbb{P}^6 \\ f \downarrow & \nearrow \tilde{\psi} & \downarrow \pi \\ \mathbb{P}^2 & \dashrightarrow \psi & \mathbb{P}^5 \end{array}$$

Proof. It suffices to note that M is the anti-canonical bundle on Y , which itself is the Del Pezzo surface of degree 6. \square

Figure 1: C_P

The hypo-osculating divisors for the Togliatti surface X_T have an easy geometric interpretation. Let $P = (a : b : c)$ be a general point in \mathbb{P}^2 . We may assume, in particular that a, b, c are all non-zero. Then the polynomial

$$f_P(x, y, z) = (cx - az)(cy - bz)(bx - ay)$$

defines a degree 3 curve (the union of 3 lines) vanishing in the three coordinate points of \mathbb{P}^2 and vanishing at P to order 3. This is visualized in Figure 1.

Since

$$f_P = bc^2 \cdot x^2y - b^2c \cdot x^2z - ac^2 \cdot y^2x + a^2c \cdot y^2z + ab^2 \cdot z^2x - a^2b \cdot z^2y, \quad (1)$$

it corresponds to a divisor D_P in \mathbb{P}^5 . Since $\text{mult}_P D_P = 3$, we can identify $\text{Osc}_P^{(2)} X_T$ with D_P . Thus, similarly as in the case of the Shifrin surface, we have that all divisors D_P are reducible.

Property 7. All divisors cut out on the Togliatti surface by 2nd osculating spaces are reducible.

4 The unexpected quartics and a B_3 -surface in \mathbb{P}^5

In this section, which is the core of this note, we study a hypo-osculating surface in \mathbb{P}^5 whose general osculating space of order 2 is determined by an irreducible divisor. We begin setting up the notation and recalling basic properties proved in [2].

4.1 The ideal of the B_3 -surface

Let Z be the following set of points in \mathbb{P}^2 :

$$\begin{aligned} P_1 &= (1 : 0 : 0), & P_2 &= (0 : 1 : 0), & P_3 &= (0 : 0 : 1), \\ P_4 &= (1 : 1 : 0), & P_5 &= (1 : -1 : 0), & P_6 &= (1 : 0 : 1), \\ P_7 &= (1 : 0 : -1), & P_8 &= (0 : 1 : 1), & P_9 &= (0 : 1 : -1). \end{aligned}$$

These points impose independent conditions on quartics in \mathbb{P}^2 . The vector space $H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(4) \otimes \mathcal{J}(Z))$ is generated by the following six quartics:

$$\begin{aligned} x^2yz, & & xy^2z, & & xyz^2, \\ xy(x^2 - y^2), & & xz(x^2 - z^2), & & yz(y^2 - z^2) \end{aligned} \quad (2)$$

and their set of zeroes is exactly Z (note that the saturated ideal $I(Z)$ has one generator xyz of degree 3 and three generators of degree 4).

Let $f : Y \rightarrow \mathbb{P}^2$ be the blow up at points P_1, \dots, P_9 with exceptional divisors E_1, \dots, E_9 . Let $H = f^*\mathcal{O}_{\mathbb{P}^2}(1)$.

Proposition 8. *The linear system*

$$M = 4H - E_1 - \dots - E_9$$

is base point free and defines a morphism birational onto its image

$$\varphi_M : Y \rightarrow \mathbb{P}^5$$

which is an isomorphism away from the proper transforms of three lines

$$\begin{aligned} L_1 : x = 0 & \quad \text{through} \quad P_2, P_3, P_8, P_9, \\ L_2 : y = 0 & \quad \text{through} \quad P_1, P_3, P_6, P_7, \\ L_3 : z = 0 & \quad \text{through} \quad P_1, P_2, P_4, P_5. \end{aligned}$$

Proof. Let

$$\begin{aligned} N_1 &= H - E_2 - E_3 - E_8 - E_9, \\ N_2 &= H - E_1 - E_3 - E_6 - E_7, \\ N_3 &= H - E_1 - E_2 - E_4 - E_5 \end{aligned}$$

be proper transformations of the lines L_1, L_2, L_3 .

To begin with, we show that M is nef. Note that

$$M.N_i = 0 \quad \text{for} \quad i = 1, 2, 3.$$

We have also $M.E_i = 1$ for $i = 1, \dots, 9$. Let C be a plane curve of degree d passing through the points P_1, \dots, P_9 with multiplicity m_1, \dots, m_9 different from L_1, L_2, L_3 . Let $\tilde{C} = dH - m_1E_1 - \dots - m_9E_9$ be the proper transform of C . Then

$$\begin{aligned} 0 &\leq \tilde{C}.N_1 = d - m_2 - m_3 - m_8 - m_9, \\ 0 &\leq \tilde{C}.N_2 = d - m_1 - m_3 - m_6 - m_7, \\ 0 &\leq \tilde{C}.N_3 = d - m_1 - m_2 - m_4 - m_5. \end{aligned}$$

Hence

$$3d - 2(m_1 + m_2 + m_3) - (m_4 + m_5 + m_6 + m_7 + m_8 + m_9) \geq 0.$$

Since

$$M.\tilde{C} = 4d - \sum_{i=1}^9 m_i,$$

we conclude that $M.\tilde{C} > 0$. So this shows not only that M is nef but also that N_1, N_2, N_3 are the only effective curves on Y orthogonal to M with respect to the intersection form.

The surface Y is an anticanonical surface. That means that $-K_Y$ is an effective divisor. Indeed, we have

$$-K_Y = N_1 + N_2 + N_3 + E_1 + E_2 + E_3.$$

Linear systems on anticanonical surfaces have been studied intensively by Harbourne. In [7, Theorem III.1.(a)] he showed in particular that if Y is an anticanonical surface and M is a nef linear system on Y such that $(-K_Y.M) \geq 2$, then M is non-special, i.e. $h^1(Y, M) = 0$ and M is base point free. These conditions hold in our situation, since

$$(-K_Y.M) = 12 - 9 = 3.$$

In [8, Proposition 3.2] Harbourne showed moreover that on an anticanonical surface a nef class M with $M^2 > 0$ and such that $(-K_Y.M) \geq 3$ defines a birational morphism onto a surface obtained by contracting all curves perpendicular to M . Since $M^2 = 7$ in our case, we conclude that the image of Y under the morphism $\varphi_{|M|}$ is a surface X_B of degree 7, singular at 3 points resulting from contracting the curves N_1, N_2 and N_3 . \square

Working with specific coordinates introduced at the beginning of this section, we can identify X_B as the closure of the image of the rational map

$$\psi : \mathbb{P}^2 \ni (x : y : z) = (x^2yz : xy^2z : xyz^2 : xy(x^2 - y^2) : xz(x^2 - z^2) : yz(y^2 - z^2)) \in \mathbb{P}^5. \quad (3)$$

In particular, the image of

$$\begin{aligned} N_1 & \text{ is } (0 : 0 : 0 : 0 : 0 : 1), \\ N_2 & \text{ is } (0 : 0 : 0 : 0 : 1 : 0), \\ N_3 & \text{ is } (0 : 0 : 0 : 1 : 0 : 0). \end{aligned}$$

Using the explicit description provided above, we conclude that the ideal $I(X_B)$ of the surface X_B in \mathbb{P}^5 is generated by

$$u_1^2 - u_2^2 - u_0u_5, \quad u_0^2 - u_2^2 - u_1u_4, \quad u_2u_3 - u_1u_4 + u_0u_5, \quad u_0u_1u_3 - u_0u_2u_4 + u_1u_2u_5 - u_3u_4u_5$$

and its minimal free resolution is

$$S/I(X_B) \leftarrow S \leftarrow S(-2)^{\oplus 3} \oplus S(-3) \leftarrow S(-4)^{\oplus 6} \leftarrow S(-5)^{\oplus 3}.$$

4.2 Osculating spaces of the B_3 surface

In this section we show that X_B is hypo-osculating.

Proposition 9. *The surface X_B satisfies one Laplace equation.*

Proof. Using the parametrization in (3), we determine second osculating spaces of X_B . Let $P = (a : b : c)$ be a general point. Evaluating second order derivatives of ψ at P we get the Hesse matrix $H_P\psi$.

	xyz^2	$yz(y^2 - z^2)$	xy^2z	x^2yz	$xz(x^2 - z^2)$	$xy(x^2 - y^2)$
∂^2/∂_x^2	0	0	0	$2bc$	$6ac$	$6ab$
∂^2/∂_y^2	0	$6bc$	$2ac$	0	0	$-6ab$
∂^2/∂_z^2	$2ab$	$-6bc$	0	0	$-6ac$	0
$\partial^2/\partial_x\partial_y$	c^2	0	$2bc$	$2ac$	0	$3(a^2 - b^2)$
$\partial^2/\partial_x\partial_z$	$2bc$	0	b^2	$2ab$	$3(a^2 - c^2)$	0
$\partial^2/\partial_y\partial_z$	$2ac$	$3(b^2 - c^2)$	$2ab$	a^2	0	0

It is easy to check either by hand or by a symbolic algebra computer program (we used Singular [3]) that $\det H_P\psi = 0$. This means that X_B satisfies at least 1 Laplace equation of order 2. Computing $\Lambda^i(H_P\psi)$ for $i = 5, 4, 3$ we conclude that

- the rank of $H_P\psi$ drops by at least 2 along the lines $abc = 0$,
- the rank of $H_P\psi$ drops by 3 at the 3 coordinate points,
- the matrix $H_P\psi$ has always rank at least 3.

□

4.3 The B_3 surface and Lefschetz Properties

In [13] Mezzetti, Miró-Roig and Ottaviani described a connection, based on Macaulay-Matlis duality (also known as apolarity), between projective varieties satisfying at least one Laplace equation and homogeneous ideals in a polynomial ring, generated by polynomials of equal degrees, and failing the Weak Lefschetz Property.

Definition 10 (Weak Lefschetz Property). An artinian homogeneous ideal I in a polynomial ring $R = \mathbb{K}[x_0, \dots, x_N]$ (i.e. an ideal such that the Krull dimension of R/I is zero) satisfies the Weak Lefschetz Property (WLP for short), if there exists a linear form $\ell \in (R/I)_1$ such that for all integers d , the multiplication map

$$\cdot \ell : (R/I)_d \ni f \rightarrow f \cdot \ell \in (R/I)_{d+1} \quad (4)$$

has the maximal rank (i.e. the map is either injective or surjective).

Note that since the algebra R/I is finite dimensional over the ground field \mathbb{K} the condition is relevant for only finitely many values of d . If such an ℓ exists, then the maximal rank property holds for a general linear form (i.e. taken from a Zariski open set in the vector space $(R/I)_1$).

There is another notion defined similarly.

Definition 11 (Strong Lefschetz Property). An ideal I as in Definition 10 is said to satisfy the Strong Lefschetz Property (SLP for short), if there exists ℓ as above such that the multiplication map

$$\cdot \ell^k : (R/I)_d \ni f \rightarrow f \cdot \ell^k \in (R/I)_{d+k} \quad (5)$$

has maximal rank for all d and all $k \geq 1$.

An ideal I is said to fail the WLP in degree d if the map (4) is not of maximal rank for all ℓ .

Similarly, an ideal I fails the SLP at range k and degree d if the map (5) is not of maximal rank for all ℓ .

The interest in the WLP stems partly from the fact that this property imposes strong constraints on the Hilbert function of I . In fact, there is a complete classification of Hilbert functions of artinian algebras satisfying the WLP, [10, Proposition 3.5]. Many algebras are expected to have this property but establishing it is often very difficult. On the other hand ideals failing the WLP often exhibit additional peculiar properties of an algebraic or geometric nature or give rise to such ideals via some standard constructions, for example Macaulay-Matlis duality. We recall now briefly what this duality is. To this end let V be a \mathbb{K} -vector space of dimension $N + 1$ and let us identify the polynomial ring R over \mathbb{K} with the symmetric algebra of forms

$$R = \bigoplus_{d \geq 0} \text{Sym}^d V^*.$$

We define the dual algebra

$$\mathcal{D} = \bigoplus_{d \geq 0} \text{Sym}^d V.$$

Let x_0, \dots, x_N and y_0, \dots, y_N be dual bases of R and \mathcal{D} , respectively. There is a natural pairing

$$\text{Sym}^j V^* \otimes \text{Sym}^i V \ni (F, G) \longrightarrow F \cdot G \in \text{Sym}^{i-j} V$$

inducing on \mathcal{D} the structure of a graded R -module. We can think of this pairing as being defined by partial differentiation in the following way

$$F \cdot G = F(\partial/\partial y_0, \dots, \partial/\partial y_N)G.$$

Usually, one does not distinguish between the coordinates in R and \mathcal{D} as in the following Example.

Example 12. Let $F_1 = x^2yz$ and $F_2 = x^3y$. Let $G = (x + y)^4$. Then

$$F_1 \cdot G = \frac{\partial^4}{\partial x^2 \partial y \partial z} G = 0,$$

$$F_2 \cdot G = \frac{\partial^4}{\partial x^3 \partial y} (x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4) = 4.$$

Definition 13 (Macaulay inverse system). Let $I \subset R$ be a homogeneous ideal. The Macaulay inverse system I^{-1} for I is

$$I^{-1} = \{G \in \mathcal{D} : F \cdot G = 0 \text{ for all } F \in I\}.$$

For the convenience of the reader, we recall the following fundamental result of Mezzetti, Miró-Roig and Ottaviani [13, Theorem 3.2].

Theorem 14. For an artinian ideal $I \subset R$ generated by $r \leq \binom{N+d-1}{N-1}$ forms F_1, \dots, F_r of degree d the following conditions are equivalent:

- i) the ideal I fails the WLP in degree $d - 1$;
- ii) The forms F_1, \dots, F_r become linearly dependent when restricted to a general hyperplane H in \mathbb{P}^N ;
- iii) The image X of \mathbb{P}^N under the morphism defined by the inverse system I^{-1} satisfies at least one Laplace equation of order $d - 1$.

The points in Z determine in the dual projective plane a B_3 arrangement of lines, which is visualized in Figure 2. The missing ninth line is the line at infinity, $z = 0$. We consider




Figure 2: $xyz(x^2 - y^2)(y^2 - z^2)(z^2 - x^2)$

the artinian ideal J generated by the fourth powers of linear forms defining the lines in the B_3 arrangement

$$J = \langle x^4, y^4, z^4, (x - y)^4, (x + y)^4, (x - z)^4, (x + z)^4, (y - z)^4, (y + z)^4 \rangle.$$

Theorem 14 does not apply directly to our situation because the number of generators 9 exceeds the upper bound in its assumptions. However, as we shall see in a moment, condition iii) is satisfied with the order $d - 2$ in place of $d - 1$. Thus it is natural to wonder if this property is reflected in the way J fails some of Lefschetz Properties. To this end we compute first (as in Example 12) the Macaulay-Matlis dual of J :

$$J^{-1} = \langle x^2yz, xy^2z, xyz^2, xy(x^2 - y^2), xz(x^2 - z^2), yz(y^2 - z^2) \rangle.$$

Thus $J^{-1} = \langle (I(Z))_4 \rangle$ by (2). Hence the morphism determined by J^{-1} is exactly the morphism ψ defined in (3). By Proposition 9 the image X_B of the inverse system satisfies one Laplace equation of order 2.

It is convenient to write down explicitly all graded pieces of the artinian algebra $A = R/J$. For $i = 0, \dots, 3$ we have $A_i = (R/J)_i = M^i$, where $M = \langle x, y, z \rangle$ is the maximal ideal and

$$A_4 = \langle xy^3, yz^3, xz^3, xyz^2, xy^2z, x^2yz \rangle.$$

All other graded pieces of A are zero.

Thus the only interesting multiplication maps are those going to A_4 . Taking as a basis of A_2 the monomials $x^2, y^2, z^2, xy, xz, yz$ and a general linear form $\ell = ax + by + cz$, we can write down explicitly the matrix of the multiplication map

$$\cdot \ell^2 : A_2 \longrightarrow A_4$$

as follows

	xy^3	xz^3	yz^3	xyz^2	xy^2z	x^2yz
x^2	$-2ab$	$-2ac$	0	0	0	$2bc$
y^2	$2ab$	0	$-2bc$	0	$2ac$	0
z^2	0	$2ac$	$2bc$	$2ab$	0	0
xy	$b^2 - a^2$	0	0	c^2	$2bc$	$2ac$
xz	0	$c^2 - a^2$	0	$2bc$	b^2	$2ab$
yz	0	0	$c^2 - b^2$	$2ac$	$2ab$	a^2

This matrix has rank at most 5 and at least 3 for any value of a, b, c . The stratification of the degeneracy locus of $\cdot \ell^2$ is in fact exactly the same as that of ψ as described in section 4.2. Thus we have recovered [5, Proposition 7.3].

Proposition 15. *The ideal J fails the SLP in range 2 at degree 2.*

Remark 16. The equivalence between the failure of the SLP in range 2 and degree $d-2$ and the existence of a non-trivial Laplace equation for the image of \mathbb{P}^2 under the mapping determined by the inverse system of J has been observed recently also by Di Gennaro and Ilardi in [4, Corollary 25] and in a more general setting by Di Gennaro, Ilardi and Vallès in [5, Theorem 5.1].

4.4 Unexpected quartics, the B_3 surface and BMSS duality

The surface X_B is of special interest, because in contrast to the Shifrin and the Togliatti surfaces, its second osculating space at a general point cuts out an irreducible divisor. Additionally, the BMSS duality (see [9, Section 4]) associates to X_B a companion surface X'_B , which is also a Togliatti-type surface. We describe these phenomena in this section.

To begin with we identify the hypo-osculating divisors.

Lemma 17. *Let $P = (a : b : c)$ be a general point in \mathbb{P}^2 . Then the quartic*

$$\begin{aligned} f_P(x : y : z) = & 3a(b^2 - c^2) \cdot x^2yz + 3b(c^2 - a^2) \cdot xy^2z + 3c(a^2 - b^2) \cdot xyz^2 \\ & + a^3 \cdot yz(y^2 - z^2) - b^3 \cdot xz(x^2 - z^2) + c^3 \cdot xy(x^2 - y^2) \end{aligned} \quad (6)$$

vanishes at all points of Z and has a triple point at P .

Proof. The existence of f_P has been observed indirectly in [5, Proposition 7.3]. This observation has been repeated explicitly in [2]. In fact it has been the starting point of research presented in this article. The explicit equation of f_P was found in [1, Section 2]. \square

It has been also observed in [1] that the equation (6) can be written down as a cubic in $(a : b : c)$ variables. More precisely we have the following.

Lemma 18. *Let $Q = (x : y : z)$ be a general point in \mathbb{P}^2 with coordinates $(a : b : c)$. Then the cubic*

$$\begin{aligned} g_Q(a : b : c) = & yz(y^2 - z^2) \cdot a^3 + xz(z^2 - x^2) \cdot b^3 + xy(x^2 - y^2) \cdot c^3 \\ & + 3x^2yz \cdot a(b^2 - c^2) - 3xy^2z \cdot b(a^2 - c^2) + 3xyz^2 \cdot c(a^2 - b^2) \end{aligned} \quad (7)$$

has a triple point at Q .

Proof. It is a straightforward calculation. \square

The interplay between f_P and g_Q has been studied in a much more general setting in [9, Section 4]. Here we take a slightly different point of view.

Proposition 19. *Let $I' = \langle a^3, b^3, c^3, a(b^2 - c^2), b(a^2 - c^2), c(a^2 - b^2) \rangle$ and let*

$$\eta : \mathbb{P}^2 \ni (a : b : c) \longrightarrow (a^3 : b^3 : c^3 : a(b^2 - c^2) : b(a^2 - c^2) : c(a^2 - b^2)) \in \mathbb{P}^5$$

be the associated rational map. Then

- η is a morphism (i.e. the ideal I' is artinian),
- the image of η is a smooth surface X'_B of degree 9,
- the ideal $I(X'_B)$ has the following minimal free resolution

$$S/I(X'_B) \leftarrow S \leftarrow S(-3)^{\oplus 7} \oplus S(-4)^{\oplus 2} \leftarrow S(-4)^{\oplus 3} \oplus S(-5)^{\oplus 19} \leftarrow S(-6)^{\oplus 21} \leftarrow S(-7)^{\oplus 8} \leftarrow S(-8).$$

Proof. These claims are easy to check by a symbolic algebra program. \square

We call the surface X'_B the (BMSS) *companion surface* of X_B . It enjoys several peculiar properties which are listed in the next proposition.

Proposition 20. *The surface X'_B is hypo-osculating. Moreover, we have*

$$\dim \text{Osc}_Q^{(2)} X'_B = 4$$

for all points $Q \in \mathbb{P}^2$ with the exception of points $Q \in Z$, where $\dim \text{Osc}_{P_i}^{(2)} X'_B = 3$ for $i = 1, \dots, 9$.

Proof. The assertion follows directly from the analysis of the Hesse matrix of η along the same lines as in the proof of Proposition 9. We omit the easy calculations. \square

Since I' has 6 generators, it is unexpected that it contains the element g_Q vanishing to order 3 at a general point Q . Of course, a cubic curve with a triple point splits in a product of lines, so, similarly as in the case of the Shifrin and the Togliatti surfaces we obtain the following.

Property 21. All divisors cut out on the companion B_3 surface by 2nd osculating spaces are reducible.

5 Final remarks

We conclude the note with the following two problems, which we hope to come back to in the near future.

Problem 22. Are there companion varieties for other Togliatti-type varieties?

Note that there is a certain renaissance of interest in Togliatti systems and hypo-osculating varieties stemming partly from their connection to Lefschetz Properties and partly from the new research direction of unexpected hypersurfaces established in [2], see also [12], [4], [11].

It follows immediately from (1) that the companion surface for the Togliatti surface is the surface itself. In other words, the Togliatti surface is BMSS self-dual.

Taking into account Properties 4, 7, 21 and the irreducibility of hypo-osculating divisors on the B_3 surface, it is tempting to ask the following question, which seems more feasible than the Shifrin conjecture mentioned in section 2.

Problem 23. Does there exist a smooth Togliatti-type surface X in \mathbb{P}^5 such that the divisor cut out on X by the second osculating space at a general point is irreducible?

Acknowledgements. I would like to thank Igor Dolgachev for suggesting that I investigate the surface associated to the B_3 root system. I thank heartily Thomas Bauer and Brian Harbourne for very helpful suggestions on the exposition of the manuscript. I thank also Jean Vallès for explaining to me chronological development of the theory and other useful comments. Finally I would like to thank Ragni Piene for clarifying the status of Shiffrin conjecture for me. The writing of this article was finished in Oberwolfach during the workshop "Asymptotic invariants of homogeneous ideals" held in October 2018. It is a pleasure to thank MFO for perfect working conditions.

References

- [1] T. Bauer, G. Malara, T. Szemberg, and J. Szpond. Quartic unexpected curves and surfaces. *Manuscripta Math.*, to appear, doi.org/10.1007/s00229-018-1091-3.
- [2] D. Cook II, B. Harbourne, J. Migliore, and U. Nagel. Line arrangements and configurations of points with an unexpected geometric property. *Compos. Math.*, 154(10):2150–2194, 2018.
- [3] W. Decker, G.-M. Greuel, G. Pfister, and H. Schönemann. SINGULAR 4-1-1 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de>, 2018.
- [4] R. Di Gennaro and G. Ilardi. Laplace equations, Lefschetz properties and line arrangements. *J. Pure Appl. Algebra*, 222(9):2657–2666, 2018.
- [5] R. Di Gennaro, G. Ilardi, and J. Vallès. Singular hypersurfaces characterizing the Lefschetz properties. *J. Lond. Math. Soc. (2)*, 89(1):194–212, 2014.
- [6] L. Farnik, F. Galuppi, L. Sodomaco, and W. Trok. On the unique unexpected quartic in \mathbf{P}^2 . *arXiv:1804.03590*.
- [7] B. Harbourne. Anticanonical rational surfaces. *Trans. Amer. Math. Soc.*, 349(3):1191–1208, 1997.
- [8] B. Harbourne. Birational morphisms of rational surfaces. *J. Algebra*, 190(1):145–162, 1997.
- [9] B. Harbourne, J. Migliore, U. Nagel, and Z. Teitler. Unexpected hypersurfaces and where to find them, *arXiv:1805.10626*.
- [10] T. Harima, J. C. Migliore, U. Nagel, and J. Watanabe. The weak and strong Lefschetz properties for Artinian K -algebras. *J. Algebra*, 262(1):99–126, 2003.
- [11] E. Mezzetti. Osculating behavior of the Kummer surface in \mathbb{P}^5 . *Eur. J. Math.*, 4(1):372–380, 2018.
- [12] E. Mezzetti and R. M. Miró-Roig. Togliatti systems and Galois coverings. *J. Algebra*, 509:263–291, 2018.
- [13] E. Mezzetti, R. M. Miró-Roig, and G. Ottaviani. Laplace equations and the weak Lefschetz property. *Canad. J. Math.*, 65(3):634–654, 2013.
- [14] R. Piene and H.-s. Tai. A characterization of balanced rational normal scrolls in terms of their osculating spaces. In *Enumerative geometry (Sitges, 1987)*, volume 1436 of *Lecture Notes in Math.*, pages 215–224. Springer, Berlin, 1990.
- [15] W. F. Pohl. Differential geometry of higher order. *Topology*, 1:169–211, 1962.
- [16] T. Shiffrin. The osculatory behavior of surfaces in \mathbf{P}^5 . *Pacific J. Math.*, 123(1):227–256, 1986.

- [17] E. Togliatti. Alcune osservazioni sulle superfici razionali che rappresentano equazioni di laplace. *Ann. Mat. Pura Appl.*, 25(4):325–339, 1929.

Justyna Szpond, Department of Mathematics, Pedagogical University of Cracow, Podchorążych 2, PL-30-084 Kraków, Poland.

E-mail address: `szpond@up.krakow.pl`