

C_0 -semigroups of m -isometries on Hilbert spaces

T. Bermúdez, A. Bonilla and H. Zaway *

October 18, 2018

Abstract

Let $\{T(t)\}_{t \geq 0}$ be a C_0 -semigroup on a separable Hilbert space H . We characterize that $T(t)$ is an m -isometry for every t in terms that the mapping $t \in \mathbb{R}^+ \rightarrow \|T(t)x\|^2$ is a polynomial of degree less than m for each $x \in H$. This fact is used to study m -isometric right translation semigroup on weighted L^p -spaces. We characterize the above property in terms of conditions on the infinitesimal generator operator or in terms of the cogenerator operator of $\{T(t)\}_{t \geq 0}$. Moreover, we prove that a non-unitary 2-isometry on a Hilbert space satisfying the kernel condition, that is,

$$T^*T(KerT^*) \subset KerT^* ,$$

then T can be embedded into a C_0 -semigroup if and only if $dim(KerT^*) = \infty$.

1 Introduction

Let H be a complex Hilbert space and $B(H)$ denote the C^* -algebra of all bounded linear operators on H .

A one parameter family $\{T(t)\}_{t \geq 0}$ of bounded linear operators from H into H is a C_0 -semigroup if:

1. $T(0) = I$.
2. $T(s + t) = T(t)T(s)$ for every $t, s \geq 0$.
3. $\lim_{t \rightarrow 0^+} T(t)x = x$ for every $x \in H$, in the strong operator topology.

The linear operator A defined by

$$Ax := \lim_{t \rightarrow 0^+} \frac{T(t)x - x}{t},$$

*The first and second authors were supported by MINECO and FEDER, Project MTM2016-75963-P. The third author was supported in part by Departamento de Análisis Matemático of Universidad de La Laguna and by a grant of Université de Gabès, UNG 933989527.

for every

$$x \in D(A) := \{x \in H : \lim_{t \rightarrow 0^+} \frac{T(t)x - x}{t} \text{ exists} \}$$

is called the *infinitesimal generator* of the semigroup $\{T(t)\}_{t \geq 0}$. It is well-known that A is a closed densely defined linear operator.

If 1 is in the resolvent set of A , $\rho(A)$, then the Caley transform of A defined by $V := (A + I)(A - I)^{-1}$ is a bounded linear operator since $V = I + 2(A - I)^{-1}$, called the *cogenerator* of the C_0 -semigroup $\{T(t)\}_{t \geq 0}$.

Notice that the point 1 could be on the spectrum of V , $\sigma(V)$.

For a positive integer m , an operator $T \in B(H)$ is called an *m-isometry* if

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \|T^k x\|^2 = 0,$$

for any $x \in H$.

We say that T is a *strict m-isometry* if T is an m -isometry but it is not an $(m - 1)$ -isometry.

- Remark 1.1.**
1. For $m \geq 2$, the strict m -isometries are not power bounded. If T is an m -isometry, then $\|T^n x\|^2$ is a polynomial at n of degree at most $m - 1$, for very x , [6, Theorem 2.1]. In particular, $\|T^n\| = O(n)$ for 3-isometries and $\|T^n\| = O(n^{\frac{1}{2}})$ for 2-isometries.
 2. There are no strict m -isometries on finite dimensional spaces for even m . See [2, Proposition 1.23].
 3. If T is an m -isometry, then $\sigma(T) = \overline{\mathbb{D}}$ or $\sigma(T) \subseteq \partial\mathbb{D}$, [2, Lemma 1.2].

The remainder of the paper is organized as follows. In Section 2, we characterize that $T(t)$ is an m -isometry for every t if the mapping $t \in \mathbb{R}^+ \rightarrow \|T(t)x\|^2$ is a polynomial of degree less than m for each $x \in H$. This fact is used in the last section in order to study m -isometric right translation semigroup on weighted L^p -spaces. Also, we characterize the above property in terms of conditions on the generator operator or in terms that the cogenerator operator be an m -isometry. Moreover, we obtain that $\{T(t)\}_{t \geq 0}$ is an m -isometry for all $t > 0$ if and only if $T(t)$ is an m -isometry, for all $t \in [0, t_1]$ with $t_1 > 0$ or on $[t_1, t_2]$ with $0 < t_1 < t_2$.

Section 3 is devoted to embed m -isometries into C_0 -semigroups. That is, given an m -isometry T finds a C_0 -semigroup $\{T(t)\}_{t \geq 0}$ such that $T(1) = T$. Using the model for 2-isometries we conclude that a non-unitary 2-isometry on a Hilbert space such that satisfies the *kernel condition*, that is

$$T^*T(Ker(T^*)) \subset Ker(T^*),$$

can be embedded into a C_0 -semigroup if and only if $\dim(Ker T^*) = \infty$.

Finally, in Section 4, we obtain a characterization of the right translation C_0 -semigroup on some weighted space, to be a semigroup of m -isometries for all $t > 0$.

2 C_0 -semigroups of m -isometries

Recall that any C_0 -semigroup $\{T(t)\}_{t \geq 0}$ have some real quantities associated as:

1. The *spectral bound*, $s(A)$, by setting

$$s(A) := \sup\{Re\lambda \quad : \quad \lambda \in \sigma(A)\}.$$

2. *Growth bound*, w_0 , given by

$$w_0 := \inf\{w \in \mathbb{R} : \exists M_w \geq 1 \quad : \quad \|T(t)\| \leq M_w e^{wt}, \quad \forall t \geq 0\}.$$

The above quantities are related in the following way, $s(A) \leq w_0$ and also

$$w_0 = \frac{1}{t} \log r(T(t)), \tag{1}$$

where $r(T(t))$ denotes the spectral radius of the operator $T(t)$.

The following lemma allow us to define the cogenerator of a C_0 -semigroup of m -isometries.

Lemma 2.1. *Let $\{T(t)\}_{t \geq 0}$ be a C_0 -semigroup on a separable Hilbert space H consisting of m -isometries and A its generator. Then $1 \in \rho(A)$ and therefore the cogenerator V of $\{T(t)\}_{t \geq 0}$ is well-defined.*

Proof. If $T(t)$ is an m -isometry for all t , then the spectral radius of $T(t)$, $r(T(t))$ is 1 for all t . Using equality (1), then $s(A) \leq w_0 = 0$ and then $1 \in \rho(A)$. \square

The following combinatorial result will be necessary for the proof of Theorem 2.1. Denote $\binom{m}{k} = 0$ if $m < k$ or $k < 0$.

Lemma 2.2. *Let m be a positive integer and p, q be integers such that $0 \leq p, q \leq m$.*

1. *If $p + q \neq m$, then*

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \left\{ \sum_{i=0}^p \binom{m-k}{i} \binom{k}{p-i} (-1)^i \right\} \left\{ \sum_{j=0}^q \binom{m-k}{j} \binom{k}{q-j} (-1)^j \right\} = 0$$

2. *If $p + q = m$, then*

$$\sum_{k=0}^m \binom{m}{k} \left\{ \sum_{i=0}^q \binom{m-k}{i} \binom{k}{q-i} (-1)^i \right\}^2 = 2^m \binom{m}{q} = 2^m \binom{m}{p}.$$

Proof. We define the polynomials r and s by $r(x, y) := 2^m(x + y)^m$ and

$$s(x, y) := ((x + 1)(y + 1) - (x - 1)(y - 1))^m.$$

Then

$$\begin{aligned} r(x, y) &= 2^m \sum_{k=0}^m \binom{m}{k} x^k y^{m-k} \\ s(x, y) &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} (x + 1)^k (x - 1)^{m-k} (y + 1)^k (y - 1)^{m-k} \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \sum_{h, \ell=0}^k \binom{k}{\ell} \binom{k}{h} \sum_{i, j=0}^{m-k} \binom{m-k}{i} \binom{m-k}{j} (-1)^{i+j} x^{j+h} y^{i+\ell} \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \left\{ \sum_{i=0}^p \binom{m-k}{m-k-i} \binom{k}{k+i-p} (-1)^i \right\} \\ &\quad \left\{ \sum_{j=0}^q \binom{m-k}{m-k-j} \binom{k}{k+j-q} (-1)^j \right\} x^{j+h} y^{i+\ell} \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \left\{ \sum_{i=0}^p \binom{m-k}{i} \binom{k}{p-i} (-1)^i \right\} \\ &\quad \left\{ \sum_{j=0}^q \binom{m-k}{j} \binom{k}{q-j} (-1)^j \right\} x^{j+h} y^{i+\ell}. \end{aligned}$$

Denote $\widehat{x^p y^q}^f$ the coefficient of the power $x^p y^q$ on the polynomial f . Since $s(x, y) = r(x, y)$, then $\widehat{x^p y^q}^s = \widehat{x^p y^q}^r$ for every p and q . So, if $p + q \neq m$, then $\widehat{x^p y^q}^s = \widehat{x^p y^q}^r = 0$. If $p + q = m$, then

$$\widehat{x^p y^q}^s = \widehat{x^p y^q}^r = \widehat{x^p y^{m-p}}^r = 2^m \binom{m}{p} = 2^m \binom{m}{m-p} = 2^m \binom{m}{q}.$$

By other hand, it is straightforward to verify that, if $p + q = m$, then

$$\begin{aligned} \widehat{x^p y^q}^s &= \sum_{k=0}^m \binom{m}{k} \left\{ \sum_{i=0}^q \binom{m-k}{m-k-i} \binom{k}{k+i-q} (-1)^i \right\}^2 \\ &= \sum_{k=0}^m \binom{m}{k} \left\{ \sum_{i=0}^q \binom{m-k}{i} \binom{k}{q-i} (-1)^i \right\}^2. \end{aligned}$$

This complete the proof. \square

Given a C_0 -semigroup $\{T(t)\}_{t \geq 0}$, we have two different operators associated to $\{T(t)\}_{t \geq 0}$, the infinitesimal generator A and the cogenerator V , if $1 \in \rho(A)$. This two operators will be useful in the next result, where we obtain a natural generalization to C_0 -semigroup of m -isometries of [12, Proposition 2.2]. See also [13, Proposition 2.6] and [21, Theorem 2].

Theorem 2.1. *Let $\{T(t)\}_{t \geq 0}$ be a C_0 -semigroup on a separable Hilbert space H . Then the following conditions are equivalent:*

- (i) $T(t)$ is an m -isometry for every t .
- (ii) The mapping $t \in \mathbb{R}^+ \rightarrow \|T(t)x\|^2$ is a polynomial of degree less than m for each $x \in H$.
- (iii) The operator inequality

$$\sum_{k=0}^m \binom{m}{k} \langle A^{m-k}x, A^kx \rangle = 0 ,$$

for any $x \in D(A^m)$.

- (iv) The cogenerator V of $\{T(t)\}_{t \geq 0}$ exists and is an m -isometry.

Proof. (i) \Leftrightarrow (ii) If $T(t)$ is an m -isometry for every t , then

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \|T(t + k\tau)x\|^2 = 0 , \quad (2)$$

for every $t, \tau > 0$ and $x \in H$. From the assumption on the semigroup, it is clear that the function $t \in \mathbb{R}^+ \rightarrow f(t) := \|T(t)x\|^2$ is continuous. By [15, Theorem 13.7], the function $f(t)$ is a polynomial of degree less than m for each $x \in H$.

Conversely, if the mapping $t \in \mathbb{R}^+ \rightarrow \|T(t)x\|^2$ is a polynomial of degree less than m for each $x \in H$, then $T(t)$ is an m -isometry for every t , [15, page 271].

(ii) \Leftrightarrow (iii) Let $y \in D(A^m)$. The function $t \in \mathbb{R}^+ \rightarrow \|T(t)y\|^2$ has m th-derivative and it is given by

$$\sum_{k=0}^m \binom{m}{k} \langle A^{m-k}T(t)y, A^kT(t)y \rangle . \quad (3)$$

By (ii) and (3) at $t = 0$ we have that

$$\sum_{k=0}^m \binom{m}{k} \langle A^{m-k}x, A^kx \rangle = 0 ,$$

for any $x \in D(A^m)$.

Conversely, if (3) holds on $D(A^m)$, then the m th-derivative of the function $t \in \mathbb{R}^+ \rightarrow \|T(t)x\|^2$ agree with

$$\sum_{k=0}^m \binom{m}{k} \langle A^{m-k}T(t)x, A^kT(t)x \rangle ,$$

for every $t > 0$ and $x \in D(A^m)$. Since $D(A^m)$ is dense by [17, Theorem 2.7], we obtain the result.

(iii) \Leftrightarrow (iv) It will be sufficient to prove that

$$2^m \sum_{k=0}^m \binom{m}{k} \langle A^{m-k}x, A^kx \rangle = \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \langle (A+I)^k(A-I)^{m-k}x, (A+I)^k(A-I)^{m-k}x \rangle \quad (4)$$

for all $x \in D(A^m)$, since (4) is equivalent to

$$\begin{aligned} & 2^m \sum_{k=0}^m \langle A^{m-k}(A-I)^{-m}y, A^k(A-I)^{-m}y \rangle \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \langle (A+I)^k(A-I)^{-k}y, (A+I)^k(A-I)^{-k}y \rangle \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \|V^k y\|^2, \end{aligned}$$

for all $y \in R(A-I)^m$.

Note that the second part of equality (4) is given by

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \sum_{\ell, h=0}^k \binom{k}{\ell} \binom{k}{h} \sum_{i, j=0}^{m-k} \binom{m-k}{i} \binom{m-k}{j} (-1)^{i+j} \langle A^{\ell+i}x, A^{h+j}x \rangle. \quad (5)$$

We denote $\widehat{A}_{i,j}$ the numerical coefficient of $\langle A^i x, A^j x \rangle$. It is straightforward to verify that if $p+q=m$, then

$$\begin{aligned} \widehat{A}_{q,m-q} &= \widehat{A}_{m-q,q} \\ &= \sum_{k=0}^m \binom{m}{k} \left\{ \sum_{i=0}^q \binom{m-k}{m-k-i} \binom{k}{k+i-q} (-1)^i \right\}^2 \\ &= \sum_{k=0}^m \binom{m}{k} \left\{ \sum_{i=0}^q \binom{m-k}{i} \binom{k}{q-i} (-1)^i \right\}^2 = 2^m \binom{m}{q}, \end{aligned}$$

where the last equality is obtained by part (2) of Lemma 2.2.

If $p+q \neq m$, then

$$\begin{aligned} \widehat{A}_{p,q} &= \widehat{A}_{m-p,m-q} \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \left\{ \sum_{i=0}^p \binom{m-k}{m-k-i} \binom{k}{k+i-p} (-1)^i \right\} \\ &\quad \left\{ \sum_{j=0}^q \binom{m-k}{m-k-j} \binom{k}{k+j-q} (-1)^j \right\} \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \left\{ \sum_{i=0}^p \binom{m-k}{i} \binom{k}{p-i} (-1)^i \right\} \left\{ \sum_{j=0}^q \binom{m-k}{j} \binom{k}{q-j} (-1)^j \right\} \\ &= 0, \end{aligned}$$

where the last equality is obtained by part (1) of Lemma 2.2. So we get the result. \square

Corollary 2.1. *Let $\{T(t)\}_{t \geq 0}$ be a C_0 -semigroup on a separable Hilbert space H . Then $T(t)$ is a strict m -isometry for every $t > 0$ if and only if the cogenerator V of $\{T(t)\}_{t \geq 0}$ is a strict m -isometry.*

In the following corollary, we give an example of m -isometric semigroup.

Corollary 2.2. *Let Q be a nilpotent operator of order n on a separable Hilbert space. Then the C_0 -semigroup generated by Q is a strict $(2n - 1)$ -isometric semigroup.*

Proof. Since Q is the generator of $\{T(t)\}_{t \geq 0}$ and $1 \in \rho(Q)$, then the cogenerator is well-defined and given by

$$V := (Q + I)(Q - I)^{-1}.$$

Thus $V = -(Q + I)(I + Q + \cdots + Q^{n-1}) = -I - 2Q(I + Q + \cdots + Q^{n-2})$, that is, the sum of an isometry and a nilpotent operator of order n . By [7, Theorem 2.2], the cogenerator is a strict $(2n - 1)$ -isometry. Then $\{T(t)\}_{t \geq 0}$ is a strict $(2n - 1)$ -isometric semigroup by Corollary 2.1. \square

For a positive integer m , a linear closed operator A defined on a dense set $D(A) \subset H$ is called an m -symmetry if

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \langle A^{m-k}x, A^kx \rangle = 0,$$

for all $x \in D(A^m)$. We say that A is a *strict m -symmetry* if A is an m -symmetry but it is not an $(m - 1)$ -symmetry. There is not strict m -symmetry bounded for even m . See [1, Page 7].

In the next result, we present the connection between the condition (iii) of Theorem 2.1 and the m -symmetric operators.

Corollary 2.3. *Let A be an m -symmetry on a separable Hilbert space. Then the C_0 -semigroup generated by iA is an m -isometric semigroup.*

Proof. If A is an m -symmetry, then

$$\sum_{k=0}^m \binom{m}{k} \langle (iA)^{m-k}x, (iA)^kx \rangle = (-i)^m \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \langle A^{m-k}x, A^kx \rangle = 0,$$

for all $x \in D(A^m)$. Thus the C_0 -semigroup generated by iA is an m -isometric semigroup. The result is completed by Theorem 2.1. \square

Let w be a weighted function defined on \mathbb{T} . It is defined

$$L_w^2(\mathbb{T}) := \{f : \mathbb{T} \rightarrow \mathbb{C} : \int_{\mathbb{T}} |f(z)|^2 w(z) dz < \infty\}.$$

Let $\{T(t)\}_{t \geq 0}$ be a C_0 -group defined on $L_w^2(\mathbb{T})$ with non-constant weighted function w by $T(t)f(z) := f(e^{it}z)$. Then $T(t)$ is an isometry if and only if t is a multiple of 2π . See [8, page 8].

Proposition 2.1. *Let $\{T(t)\}_{t \in \mathbb{R}}$ be a C_0 -group on a Hilbert space H . Then the following conditions are equivalent:*

- (i) $T(t)$ is an m -isometry for every t .
- (ii) $T(t)$ is an m -isometry for t_1 and t_2 where $\frac{t_1}{t_2}$ is irrational.

Proof. For every t , we have that

$$\sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \|T(t + kt_i)x\|^2 = 0 \quad \text{for } i = 1, 2.$$

Montel's Theorem ([15, Theorem 13.5] & [19, Theorem 1.1]) gives that $\|T(t)x\|^2$ is a polynomial of degree less than m for each $x \in H$, since $\frac{t_1}{t_2}$ is irrational. Thus by Theorem 2.1 we have that $T(t)$ is an m -isometry for every t . \square

If T is an m -isometry, then any power T^r is also an m -isometry, [14, Theorem 2.3]. In general, the converse is not true. However, if T^r and T^{r+1} are m -isometries for a positive integer r , then T is an m -isometry (see, [5, Corollary 3.7]). The stability of the class of m -isometry with powers is fundamental to give necessary and sufficient conditions for a C_0 -semigroup $\{T(t)\}_{t \geq 0}$ to be $T(t)$ an m -isometry for all $t \geq 0$.

Theorem 2.2. *Let $\{T(t)\}_{t \geq 0}$ be a C_0 -semigroup on a Hilbert space H . Then the following properties are equivalent:*

- (i) $T(t)$ is an m -isometry for every $t \geq 0$.
- (ii) $T(t)$ is an m -isometry for every $t \in [0, t_1]$ for some $t_1 > 0$.
- (iii) $T(t)$ is an m -isometry on an interval of the form $[t_1, t_2]$ with $t_1 < t_2$.

Proof. The implications (i) \Rightarrow (ii) and (ii) \Rightarrow (iii) are clear.

(ii) \Rightarrow (i). Fixed $t > 0$, there exist $n \in \mathbb{N}$ and $t' \in [0, t_1]$ such that $t = nt'$. Since $T(t')$ is an m -isometry, then any power of $T(t')$ is an m -isometry by [5, Theorem 3.1]. So, $T(t)$ is an m -isometry.

(iii) \Rightarrow (i). Let us prove that $T(t)$ is an m -isometry for every $t \in (0, \frac{t_2 - t_1}{4}]$.

Choose $k := \lceil \frac{t_1}{t} \rceil + 1$. Then kt and $(k+1)t$ belong to (t_1, t_2) . Thus $T(t)^k$ and $T(t)^{k+1}$ are m -isometries. Hence $T(t)$ is an m -isometry by [5, Corollary 3.7]. \square

3 Embedding m -isometries into C_0 -semigroups

We are interested to study: When can we embed an m -isometry into a continuous C_0 -semigroup?, that is, given an m -isometry T , to find a C_0 -semigroup $\{T(t)\}_{t \geq 0}$ such that $T(1) = T$.

Recall that an isometry T on a Hilbert space can be embedded into a C_0 -semigroup if and only if T is unitary or $\text{codim}(R(T)) = \infty$, where $R(T)$ denotes the range of T . In this case, it is also possible to embed T into an isometric C_0 -semigroup [10, Theorem V.1.19].

Note that, if T can be embedded into C_0 -semigroup, then $\dim(\text{Ker}T)$ and $\dim(\text{Ker}T^*)$ are zero or infinite [10, Theorem V.1.7].

Proposition 3.1. (i) All m -isometries in finite dimensional space are embeddable on a C_0 -group.

(ii) All normal m -isometries are embeddable.

(iii) The weighted forward shift m -isometries are not embeddable.

Proof. (i) On a finite-dimensional space an operator can be embeddable if and only if its spectrum does not contain 0 (see, [10, page 166]). Moreover, on finite-dimensional space the spectrum of m -isometries is contained in the unit circle. Hence any m -isometry on finite dimensional space is embeddable in a group.

(ii) If T is a normal m -isometry, then T is invertible and by [10, Theorem V.1.14] T is embeddable.

(iii) Assume that T is a weighted forward shift m -isometry. Then $\dim \text{Ker}(T^*) = 1$, ([3] & [6]). \square

Let M_z be the multiplication operator on the Dirichlet space $D(\mu)$ for some finite non-negative Borel measure on \mathbb{T} defined by

$$D(\mu) := \{f : \mathbb{D} \rightarrow \mathbb{C} \text{ analytic} : \int_{\mathbb{D}} |f'(z)|^2 \varphi_\mu(z) dA(z) < \infty\},$$

where A denotes the normalized Lebesgue area measure on \mathbb{D} and φ_μ is defined by

$$\varphi_\mu(z) := \frac{1}{2\pi} \int_{[0, 2\pi)} \frac{1 - |z|^2}{|e^{it} - z|^2} d\mu(t),$$

for $z \in \mathbb{D}$.

Proposition 3.2. M_z on $D(\mu)$ can not be embedded into C_0 -semigroup.

Proof. By Richter's Theorem [20], M_z is an analytic 2-isometry with $\dim(\text{Ker}T^*) = 1$, then M_z can not be embedded into C_0 -semigroup. \square

Let Y be an infinite dimensional Hilbert space. Denote by ℓ_Y^2 the Hilbert space of all vector sequences $(h_n)_{n=1}^\infty$ such that $\sum_{n \geq 1} \|h_n\|^2 < \infty$ with the standard inner product. If $(W_n)_{n=1}^\infty \subset B(Y)$ is a uniformly bounded sequence of operators, then the operator $S_W \in B(\ell_Y^2)$ defined by

$$S_W(h_1, h_2, \dots) := (0, \overline{W_1 h_1}, \overline{W_2 h_2}, \dots),$$

for any $(h_1, h_2, \dots) \in \ell_Y^2$, is called the *operator valued unilateral forward weighted shifts* with weights $W := (W_n)_{n \geq 1}$.

Following some ideas of [10, Proposition V.1.18], we obtain the next result.

Lemma 3.1. Let S_W be the valued unilateral forward weighted shift operator with weights $W = (W_n)_{n \geq 1}$ on ℓ_Y^2 where Y be an infinite dimensional Hilbert space. Then S_W can be embedded into C_0 -semigroup.

Proof. The operator S_W is unitarily equivalent to the valued unilateral forward weighted shift operator on $\ell_{L^2([0,1],Y)}^2$. Moreover, $\ell_{L^2([0,1],Y)}^2$ can be identified with $L^2(\mathbb{R}^+, Y)$ by

$$(f_1, f_2 \cdots) \rightarrow (s \rightarrow f_n(s-n), \quad s \in [n, n+1)).$$

For $0 < t \leq 1$, define the following family of operators on $L^2(\mathbb{R}^+, Y)$ by

$$(T(t)f)(s) := \begin{cases} 0 & s < t \\ f(s-t) & n-1+t \leq s < n \\ W_n f(s-t) & n \leq s < n+t, \end{cases}$$

that is,

$$(T(t)f)(s) = \begin{cases} 0 & s < t \\ \sum_{n \geq 1} (f(s-t)\chi_{[n-1+t,n)}(s) + W_n f(s-t)\chi_{[n,n+t)}(s)) & s \geq t. \end{cases}$$

In particular, for $t = 1$, we have that

$$(T(1)f)(s) = \begin{cases} 0 & s < 1 \\ \sum_{n \geq 1} W_n f(s-1)\chi_{[n,n+1)}(s) & s \geq 1. \end{cases}$$

Hence $T(1)$ is unitarily equivalent to S_W .

For $t > 1$, define $T(t) := T^{[t]}(1)T(t - [t])$, where $[t]$ denotes the greatest integer less than or equal to t .

Let us prove that $T(t)$ is a C_0 -semigroup. Given any $f \in L^2(\mathbb{R}^+, Y)$, we have that

$$\lim_{t \rightarrow 0^+} T(t)f(s) = \sum_{n \geq 1} \chi_{[n-1,n)} f(s) = f(s),$$

in the strong topology of $L^2(\mathbb{R}^+, Y)$.

Let $t, t' \in [0, 1)$. Then $T(t)T(t')f(s) = T(t)\tilde{f}(s)$, where

$$\tilde{f}(s) := T(t')f(s) = \begin{cases} 0 & s < t' \\ \sum_{n \geq 1} (\chi_{[n-1+t',n)}(s) + W_n \chi_{[n,n+t')}(s)) f(s-t') & s \geq t'. \end{cases}$$

Then

$$\begin{aligned} & T(t)T(t')f(s) \\ &= \begin{cases} 0 & s < t+t' \\ \sum_{m \geq 1} \left(\sum_{n \geq 1} (\chi_{[n-1+t',n)}(s-t) + W_n \chi_{[n,n+t')}(s-t)) \chi_{[m-1+t,m)}(s) + \right. \\ \left. W_m \sum_{n \geq 1} (\chi_{[n-1+t',n)}(s-t) + W_n \chi_{[n,n+t')}(s-t)) \right. \\ \left. \chi_{[m,m+t)}(s) \right) f(s-t-t') & s \geq t+t' \end{cases} \end{aligned}$$

$$= \begin{cases} 0 & s < t + t' \\ \sum_{m \geq 1} \left\{ \sum_{n \geq 1} (\chi_{[n-1+t'+t, n+t]}(s) + W_n \chi_{[n+t, n+t'+t]}(s)) \chi_{[m-1+t, m]}(s) + \right. \\ \left. W_m \sum_{n \geq 1} (\chi_{[n-1+t'+t, n+t]}(s) + W_n \chi_{[n+t, n+t'+t]}(s)) \right. \\ \left. \chi_{[m, m+t]}(s) \right\} f(s - t' - t) & s \geq t + t' \end{cases} \quad (6)$$

If $t' + t < 1$, then (6) is given by

$$\begin{aligned} T(t)T(t')f(s) &= \begin{cases} 0 & s < t' + t \\ \sum_{n \geq 1} (\chi_{[n-1+t'+t, n]}(s) + W_n \chi_{[n, n+t'+t]}(s)) f(s - t' - t) & s \geq t' + t \end{cases} \\ &= T(t + t')f(s). \end{aligned}$$

Denote $t'' := t' + t - [t' + t]$. If $t' + t > 1$, then $t'' = t' + t - 1$ and

$$\begin{aligned} T(t + t')f(s) &= T^{[t+t']}(1)(T(t + t' - [t + t']))f(s) \\ &= T(1)T(t + t' - 1)f(s) = T(1)T(t'')f(s) \\ &= \begin{cases} 0 & s < t'' \\ T(1) \sum_{n \geq 1} (\chi_{[n-1+t'', n]}(s) + W_n \chi_{[n, n+t'']}(s)) f(s - t'') & s \geq t'' \end{cases} \\ &= \begin{cases} 0 & s - t'' < 1 \\ \sum_{m \geq 1} W_m \sum_{n \geq 1} (\chi_{[n-1+t'', n]}(s-1) + W_n \chi_{[n, n+t'']}(s-1)) \\ \chi_{[m, m+1]}(s) f(s - t'' - 1) & s - t'' \geq 1 \end{cases} \\ &= \begin{cases} 0 & s - t'' < 1 \\ \sum_{m \geq 1} W_m \sum_{n \geq 1} (\chi_{[n+t'', n+1]}(s) + W_n \chi_{[n+1, n+t''+1]}(s)) \\ \chi_{[m, m+1]}(s) f(s - t'' - 1) & s \geq 1 + t'' \end{cases} \\ &= \begin{cases} 0 & s < t + t' \\ \sum_{n \geq 1} (W_n \chi_{[n+t'', n+1]}(s) + W_{n+1} W_n \chi_{[n+1, n+t''+1]}(s)) f(s - t' - t) & s \geq t + t'. \end{cases} \end{aligned}$$

By other hand, by (6) we have that

$$T(t)T(t')f(s) =$$

$$\begin{cases} 0 & s < t + t' \\ \sum_{n \geq 1} (W_n \chi_{[n+t'+t-1, n+1)}(s) + W_{n+1} W_n \chi_{[n+1, n+t'+t)}(s)) f(s - t' - t) & s \geq t + t' \end{cases}$$

This completes the proof. \square

We say that an operator $T \in B(H)$ satisfies the *kernel condition* if

$$T^*T(\text{Ker}T^*) \subset \text{Ker}T^* .$$

Corollary 3.1. *A non-unitary 2-isometry on a Hilbert space satisfying the kernel condition can be embedded into C_0 -semigroup if and only if $\dim(\text{Ker}T^*) = \infty$.*

Proof. If T is a non-unitary 2-isometry on a Hilbert space satisfying the kernel condition as consequence of [4, Theorem 3.8], we obtain that $T \cong U \oplus W$ with U unitary and W a operator valued unilateral forward weighted shifts operator in ℓ_M^2 with $\dim M = \dim(\text{Ker}T^*)$. Thus by [10, Theorem V.1.19] and Lemma 3.1, T can be embedded into C_0 -semigroup. \square

By the Wold Decomposition Theorem for 2-isometries (see [16]), all 2-isometries can be decomposed as directed sum of unitary operator and an analytic 2-isometries.

Question 3.1. An analytic 2-isometry on a Hilbert space can be embedded into C_0 -semigroup if and only if $\dim(\text{Ker}T^*) = \infty$?

4 Translation semigroups of m -isometries

In this section, we discuss examples of semigroups of m -isometries.

Definition 4.1. By a *right admissible weighted function* in $(0, \infty)$, we mean a measurable function $\rho : (0, \infty) \rightarrow \mathbb{R}$ satisfying the following conditions:

1. $\rho(\tau) > 0$ for all $\tau \in (0, \infty)$,
2. there exist constants $M \geq 1$ and $\omega \in \mathbb{R}$ such that $\rho(t + \tau) \leq M e^{\omega t} \rho(\tau)$ holds for all $\tau \in (0, \infty)$ and all $t > 0$.

For a right admissible weighted function, we define the *weighted space* $L^2(\mathbb{R}^+, \rho)$ of measurable functions $f : \mathbb{R}^+ \rightarrow \mathbb{C}$ such that

$$\|f\|_{L^2(\mathbb{R}^+, \rho)} = \int_0^\infty |f(s)|^2 \rho(s) ds < \infty .$$

Then the *right translation semigroup* $(S(t))_{t \geq 0}$ given for $t \geq 0$ and $f \in L^2(\mathbb{R}^+, \rho)$ by

$$(S(t)f)(s) := \begin{cases} 0 & \text{if } s \leq t \\ f(s - t) & \text{if } s > t , \end{cases}$$

is a strongly continuous semigroup and straightforward computation shows that for $s, t \geq 0$ and $f \in L^2(\mathbb{R}^+, \rho)$

$$(S^*(t)f)(s) = \frac{\rho(s + t)}{\rho(s)} f(s + t) .$$

Theorem 4.1. *Let $\{S(t)\}_{t \geq 0}$ be the right translation C_0 -semigroup on $L^2(\mathbb{R}^+, \rho)$ with ρ a continuous function. The semigroup $\{S(t)\}_{t \geq 0}$ is an m -isometry for every $t > 0$ if and only if $\rho(s)$ is a polynomial of degree less than m .*

Proof. By definition, $S(t)$ is an m -isometry for every $t > 0$ if

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \|S^k(t)f\|^2 = 0,$$

for all $t \geq 0$ and $f \in L^2(\mathbb{R}^+, \rho)$. That is,

$$\int_0^\infty \left(\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \frac{\rho(s+kt)}{\rho(s)} \right) |f(s)|^2 \rho(s) ds = 0, \quad (7)$$

for all $t \geq 0$ and $f \in L^2(\mathbb{R}^+, \rho)$. Fixed $t \geq 0$, define

$$g(s) := \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \frac{\rho(s+kt)}{\rho(s)}.$$

If $g(s) \neq 0$, we can suppose without loss of generality that, $g(s) > 0$, for some $s \geq 0$. Then by continuity of ρ and $\rho(\tau) > 0$ for all $\tau \geq 0$, we obtain that there exists an interval $I \subset \mathbb{R}^+$, with finite measure, such that there exists $M > 0$ such that $g(s) > M$ for all $s \in I$.

Let $f_1(s) := \frac{1}{\sqrt{\rho(s)}} \chi_I(s) \in L^2(\mathbb{R}^+, \rho)$. Then by (7) we have that

$$0 = \int_0^\infty g(s) |f_1(s)|^2 \rho(s) ds = \int_I g(s) ds > M \mu(I),$$

which is an absurd. So,

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \frac{\rho(s+kt)}{\rho(s)} = 0,$$

for all $s \geq 0$ and $t \geq 0$. Then by [15, Theorem 13.5], the function ρ is a polynomial of degree less than m . \square

Corollary 4.1. *Let $\{S(t)\}_{t \geq 0}$ be the right translation C_0 -semigroup on $L^2(\mathbb{R}^+, \rho)$ with ρ a continuous function. Then $\{S(t)\}_{t \geq 0}$ is a strict m -isometry for every $t > 0$ if and only if $\rho(s)$ is a polynomial of degree $m - 1$.*

Consider the right weighted translation C_0 -semigroup, S_ρ , defined on $L^2(\mathbb{R}^+)$ as

$$(S_\rho(t)f)(s) := \begin{cases} 0 & \text{if } s \leq t \\ \frac{\rho(s)}{\rho(s-t)} f(s-t) & \text{if } s > t. \end{cases}$$

Then $\{S_\rho(t)\}_{t \geq 0}$ is a strongly continuous semigroup if and only if ρ is a right admissible weight [9] and for $s, t \geq 0$ and $f \in L^2(\mathbb{R}^+)$

$$(S_\rho^*(t)f)(s) = \frac{\rho(s+t)}{\rho(s)}f(s+t).$$

We now improve part (2) of [18, Corollary 3.3].

Theorem 4.2. *Let $\{S_\rho(t)\}_{t \geq 0}$ be the right weighted translation C_0 -semigroup on $L^2(\mathbb{R}^+)$ with ρ a continuous function. Then $\{S_\rho(t)\}_{t \geq 0}$ is an m -isometry for every $t > 0$ if and only if $\rho(s)^2$ is a polynomial of degree less than m .*

Proof. Consider $M_\rho : L^2(\mathbb{R}^+, \rho^2) \rightarrow L^2(\mathbb{R}^+)$ defined by $M_\rho f = \rho f$. Then $S_\rho(t) = M_\rho S(t) M_\rho^{-1}$. Thus $\{S_\rho(t)\}_{t \geq 0}$ is an m -isometry for every $t > 0$ on $L^2(\mathbb{R}^+)$ if and only if $\{S(t)\}_{t \geq 0}$ is an m -isometry for every $t > 0$ on $L^2(\mathbb{R}^+, \rho^2)$ if and only if $\rho(s)^2$ is a polynomial of degree less than m . \square

Corollary 4.2. *Let $\{S_\rho(t)\}_{t \geq 0}$ be the right weighted translation C_0 -semigroup on $L^2(\mathbb{R}^+)$ with ρ a continuous function. Then $\{S_\rho(t)\}_{t \geq 0}$ is a 2-isometry for every $t > 0$ if and only if $\rho(s)^2 = as + b$ for some constants a and b .*

At continuation we characterize the weighted spaces where the adjoint of translation operator is an m -isometry.

Definition 4.2. By a *left admissible weighted function* in $(0, \infty)$, we mean a measurable function $w : (0, \infty) \rightarrow \mathbb{R}$ satisfying the following conditions:

1. $w(\tau) > 0$ for all $\tau \in (0, \infty)$,
2. there exist constants $M \geq 1$ and $\alpha \in \mathbb{R}$ such that $w(\tau) \leq Me^{\alpha t} w(t + \tau)$ holds for all $\tau \in (0, \infty)$ and all $t > 0$.

For a left admissible weighted function we define the *weighted space* $L^2(\mathbb{R}^+, w)$ as the measurable functions $f : \mathbb{R}^+ \rightarrow \mathbb{C}$ such that

$$\|f\|_{L^2(\mathbb{R}^+, w)} = \int_0^\infty |f(s)|^2 w(s) ds < \infty.$$

Let $\{T(t)\}_{t \geq 0}$ be the *left shift semigroup* given for $t \geq 0$ and $f \in L^2(\mathbb{R}^+, w)$ by

$$(T(t)f)(s) := f(s+t).$$

Then $\{T(t)\}_{t \geq 0}$ is a strongly continuous semigroup [11]. For $f \in L^2(\mathbb{R}^+, w)$,

$$(T^*(t)f)(s) = \begin{cases} 0 & \text{if } s \leq t \\ \frac{w(s-t)}{w(s)}f(s-t) & \text{if } s > t. \end{cases}$$

Theorem 4.3. *Let $\{T^*(t)\}_{t \geq 0}$ be the adjoint of left weighted translation C_0 -semigroup on $L^2(\mathbb{R}^+, w)$ such that w is a continuous function. Then $\{T^*(t)\}_{t \geq 0}$ is an m -isometry for every $t > 0$ if and only if $w(s) = \frac{1}{p(s)}$ for some polynomial p of degree less than m .*

Proof. $T^*(t)$ is an m -isometry for every $t > 0$ if and only if

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \|T^{*k}(t)f\|^2 = 0,$$

for all $t > 0$ and $f \in L^2(\mathbb{R}^+, w)$. Then

$$\begin{aligned} 0 &= \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} (-1)^{m-k} \int_{kt}^{\infty} \left| \frac{w(s-kt)}{w(s)} f(s-kt) \right|^2 w(s) ds \\ &= \int_0^{\infty} \left(\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \frac{w(u)}{w(u+kt)} \right) |f(u)|^2 w(u) du, \end{aligned}$$

for all $f \in L^2(\mathbb{R}^+, w)$. As in the proof of Theorem 4.1 we get that $\frac{1}{w(s)}$ is a polynomial of degree less than m . \square

Corollary 4.3. *Let $\{T^*(t)\}_{t \geq 0}$ be the adjoint of left weighted translation C_0 -semigroup on $L^2(\mathbb{R}^+, w)$ such that w is a continuous function. Then $\{T^*(t)\}_{t \geq 0}$ is a strict m -isometry for every $t > 0$ if and only if $w(s) = \frac{1}{p(s)}$ for some polynomial p of degree $m - 1$.*

Acknowledgements

The authors thank to Almira and Hernández-Abreu for calling their attention over Montel's theorems.

References

- [1] J. Agler, A disconjugacy theorem for Toeplitz operators. *Amer. J. Math.*, **112** (1990), no. 1, 1-14.
- [2] J. Agler, M. Stankus, m -isometric transformations of Hilbert space. I, *Integral Equations Operator Theory*, **21** (1995), no. 4, 383-429.
- [3] J. M. Almira and A. J. López-Moreno, On solutions of the Fréchet functional equation. *J. Math. Anal. Appl.* **332** (2007), no. 2, 1119-1133.
- [4] A. Anand, S. Chavan, Z. J. Jablonski and J. Stochel, A solution to the Cauchy dual subnormality problem for 2-isometries, arXiv:1702.01264v3.
- [5] T. Bermúdez, C. Díaz and A. Martínón, Powers of m -isometries, *Studia Math.*, **208** (2012), no. 3, 249–255.
- [6] T. Bermúdez, A. Martínón and E. Negrín, Weighted shift operators which are m -isometries, *Integral Equations Operator Theory*, **68** (2010), no. 3, 301–312.

- [7] T. Bermúdez, A. Martínón and J. A. Noda, An isometry plus a nilpotent operator is an m -isometry, *Applications. J. Math. Anal. Appl.*, **407** (2013), no. 2, 505-512.
- [8] I. Chalendar, J. R. Partington, Compactness, differentiability and similarity to isometry of composition semigroups. *Problems and recent methods in operator theory*, 67–73, *Contemp. Math.*, **687**, Amer. Math. Soc., Providence, RI, 2017.
- [9] M.R. Embry and A. Lambert, Weighted translation semigroups, *Rocky Mountain J. of Math.*, **7** (1977), no. 2, 333-344.
- [10] T. Eisner, *Stability of operators and operator semigroups*, *Operator Theory: Advances and Applications*, **209**. Birkhäuser Verlag, Basel, 2010.
- [11] K.-G. Grosse-Erdmann and A. Peris, *Linear Chaos*, Springer, London, 2011.
- [12] E. A. Gallardo-Gutiérrez and J. R. Partington, C_0 -semigroups of 2-isometries and Dirichlet spaces, *Revista Matemática Iberoamericana*, to appear.
- [13] B. Jacob, J. R. Partington, S. Pott, and A. Wynn, β -admissibility of observation and control operators for hypercontractive semigroups, *J. Evolution Equations*, **18** (2018) no. 1, 153-170.
- [14] Z. J. Jabłonski, Complete hyperexpansivity, subnormality and inverted boundedness conditions. *Integral Equations Operator Theory* **44** (2002), no. 3, 316-336.
- [15] M. Kuczma, *Functional equations in a single variable*, PWN-Polish Scientific Publishers, Warszawa, 1968.
- [16] A. Olofsson, A von Neumann-Wold decomposition of two-isometries, *Acta Sci. Math. (Szeged)*, **70** (2004), no. 3-4, 715–726.
- [17] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*. Applied Mathematical Sciences, **44**. Springer-Verlag, New York, 1983.
- [18] G. M. Phatak and V. M. Sholapurkar, Hyperexpansive weighted translation semigroups, arXiv:1803.08623v1.
- [19] D. Popa and I. Rasa, The Fréchet functional equation with application to the stability of certain operators. *J. Approx. Theory*, **164** (2012), no. 1, 138-144.
- [20] S. Richter, A representation theorem of cyclic analytic two-isometries, *Trans. Amer. Math. Soc.*, **328** (1991), no. 1, 325-349.
- [21] E. Rydhe, An Agler-type model theorem for C_0 -semigroups of Hilbert space contractions, *J. Lond. Math. Soc.*, **93** (2016), no. 2, 420-438.

T. Bermúdez

Departamento de Análisis Matemático, Universidad de La Laguna, 38271, La Laguna (Tenerife), Spain.

e-mail: tbermude@ull.es

A. Bonilla

Departamento de Análisis Matemático, Universidad de La Laguna, 38271, La Laguna (Tenerife), Spain.

e-mail: abonilla@ull.es

H. Zaway

Departamento de Análisis Matemático, Universidad de La Laguna, 38271, La Laguna (Tenerife), Spain.

Department of Mathematics, Faculty of Sciences, University of Gabes, 6072, Tunisia.

e-mail: hajer_zaway@live.fr