

FRACTIONAL SUPERSYMMETRIC QUANTUM MECHANICS AND LACUNARY HERMITE POLYNOMIALS

F. BOUZEFFOUR ^{*}, M. GARAYEV [◇]

ABSTRACT. We consider a realization of fractional supersymmetric of quantum mechanics of order r , where the Hamiltonian and supercharges involve reflection operators. It is shown that the Hamiltonian has r -fold degenerate spectrum and the eigenvalues of hermitian supercharges are zeros of the associated Hermite polynomials of Askey and Wimp. Also it is shown that the associated eigenfunctions involve lacunary Hermite polynomials.

1. INTRODUCTION

Supersymmetry relates bosons and fermions on the basis of \mathbb{Z}_2 -graded superalgebras [1, 2], where the fermionic set is realized by matrices of finite dimension or in terms of Grassmann variables. In [3], M. Plyushchay formulated a supersymmetric quantum mechanics (SUSYQM) for one-dimensional systems without introducing independent fermion ("spin-like") degrees of freedom but taking reflection operator R as a grading operator, see also [4, 5, 6]. In this realization, the components of the supersymmetric Hamiltonian and supercharges are realized as differential operators which additionally involve in their structure the reflection operator being a kind of Yang-Dunkl type operator [4, 7, 8]. In simplest supersymmetric systems of such a nature related to the quantum harmonic oscillator, the eigenstates are expressed in terms of Hermite orthogonal polynomials [9]. The fractional supersymmetric quantum mechanics of order r is an extension of the ordinary supersymmetric quantum mechanics for which the \mathbb{Z}_2 -graded superalgebras are replaced by \mathbb{Z}_r -graded superalgebras [10, 11, 6, 12]. The framework of the fractional supersymmetric quantum mechanics has been shown to be quite fruitful. Among many works, we may quote the deformed Heisenberg algebra introduced in connection with parafermionic and parabosonic systems [4, 12], C_λ -extended oscillator algebra developed in the framework of parasupersymmetric quantum mechanics [13], and the generalized Weyl-Heisenberg algebra W_k related to \mathbb{Z}_k -graded supersymmetric quantum mechanics [12]. Note that the construction of fractional supersymmetric quantum mechanics without employment of fermions and parafermions degrees of freedom was started by Plyushchay [3, 4]. In particular, the idea of realization of fractional supersymmetry in the form as it was presented in refs. [12] and [13], was proposed originally in [6], see also [5].

Due to the well-known fact that the case $r = 2$ corresponds to ordinary supersymmetry, it is interesting to study which supersymmetric Hamiltonian involving reflection operators admits exact eigenfunctions which are expressible in terms of classical orthogonal polynomial system.

In this paper, we give essentially an extension of the construction elaborated by Plyushchay in [3, 4] and develop a fractional supersymmetric quantum of order r without parafermionic degrees of freedom. For this purpose, we use a generalization of Klein operator of the form $K^r = 1$, $r \geq 2$, in order to study some mathematical properties of the fractional

supersymmetric quantum mechanics. We also investigate such a possibility of realizing $r = 2$ supersymmetry for the case of fractional supersymmetric quantum mechanics.

The paper is organized as follows. In Section 2, we review the definition of the ordinary supersymmetric algebra and provide some of their realizations. In section 3, we present a realization of the fractional supersymmetric quantum mechanics. In section 4, we construct a basis involving the lacunary Hermite polynomials that diagonalizes simultaneously the Hamiltonian and the supercharge. We show that the eigenvalues of the supercharge are the zeros of the associated Hermite polynomials.

2. ORDINARY SUPERSYMMETRIC QUANTUM MECHANICS

Let us briefly recall the formalism of the usual supersymmetric quantum mechanics. The SUSYQM, introduced by Witten [1], is generated by three operators Q_- , Q_+ and H satisfying

$$(2.1) \quad Q_{\pm}^2 = 0, \quad [Q_{\pm}, H] = 0, \quad \{Q_-, Q_+\} = H.$$

Superalgebra (2.1) corresponds to a supersymmetric system with $N = 2$ supersymmetry generated by two supercharges being mutually conjugated nilpotent operators. The usual construction of Witten's supersymmetric quantum mechanics with the superalgebra (2.1) is performed by introduction of fermion degrees of freedom (realized in a matrix form, or in terms of Grassmann variables) which commute with bosonic degrees of freedom [2]. Another realization of supersymmetric quantum mechanics is the so-called minimally bosonized supersymmetric quantum. Its supercharges Q_{i-} and Q_{i+} have the expression (see, [3])

$$(2.2) \quad Q_{i-} = \frac{1}{\sqrt{2}}\left(\frac{d}{dx} + W(x)\right)\Pi_i \quad \text{and} \quad Q_{i+} = \frac{1}{\sqrt{2}}\Pi_i\left(-\frac{d}{dx} + W(x)\right), \quad i = 0, 1,$$

where $W(x)$ is an odd function $W(-x) = -W(x)$, Π_0 and Π_1 are the orthogonal projection operators defined by

$$(2.3) \quad \Pi_0 = \frac{1}{2}(1 + R), \quad \text{and} \quad \Pi_1 = \frac{1}{2}(1 - R).$$

Here, R is the reflection operator

$$(2.4) \quad (Rf)(x) = f(-x).$$

A straightforward computation shows that

$$(2.5) \quad Q_{i\pm}^2 = 0,$$

and the corresponding Hamiltonian H_0 and H_1 defined by (2.1), have the forms

$$(2.6) \quad H_0 = -\frac{1}{2}\frac{d^2}{dx^2} + \frac{1}{2}W^2 - \frac{1}{2}RW',$$

$$(2.7) \quad H_1 = -\frac{1}{2}\frac{d^2}{dx^2} + \frac{1}{2}W^2 + \frac{1}{2}RW'.$$

In Eq. (2.2), two pairs of mutually conjugated operators correspond to two different $N = 2$ supersymmetric systems with a bosonized supersymmetry. Each of such two systems can be obtained from the corresponding Witten's system (2.1) with independent fermion degrees of freedom via a special non-local unitary transformation and subsequent reduction described in detail in ref. [6, 14]. In addition, if the initial system (2.2) is in the phase of unbroken supersymmetry, as it happens in the case of superoscillator, one of the corresponding resulting system will be described by unbroken bosonized $N = 2$ supersymmetry,

while another system will be described by the broken $N = 2$ bosonized supersymmetry. These two cases are described by Hamiltonian operators (2.14) and (2.15), respectively. In the particular case, when $W(x) = x$, the supercharges Q_{i-} and Q_{i+} are expressed as

$$(2.8) \quad Q_{i-} = a_- \Pi_i, \quad Q_{i+} = \Pi_i a_+,$$

where a_+ and a_- are the standard creation and annihilation operators:

$$(2.9) \quad a_- = \frac{1}{\sqrt{2}} \left(\frac{d}{dx} + x \right), \quad a_+ = \frac{1}{\sqrt{2}} \left(-\frac{d}{dx} + x \right).$$

Recall that the eigenvalue problem corresponding to the harmonic oscillator is the differential equation for $\psi(x)$

$$(2.10) \quad -\frac{1}{2} \frac{d^2 \psi(x)}{dx^2} + \frac{1}{2} x^2 \psi(x) = \lambda \psi(x).$$

The wave functions corresponding to the well-known eigenvalues

$$(2.11) \quad \lambda_n = n + \frac{1}{2}, \quad n = 0, 1, 2, \dots,$$

are given by

$$(2.12) \quad \psi_n(x) = (\sqrt{\pi} n! 2^n)^{-1/2} e^{-x^2/2} H_n(x),$$

where $H_n(x)$ is the Hermite polynomial of order n . It follows that the wave functions $\psi_n(x)$ form an orthonormal basis in the Hilbert space $L^2(\mathbb{R})$ and satisfy the orthogonality relations

$$(2.13) \quad \int_{\mathbb{R}} \psi_n(x) \psi_m(x) dx = \delta_{nm}.$$

From (2.6) and (2.7), we have

$$(2.14) \quad H_0 = -\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} x^2 - \frac{1}{2} R,$$

$$(2.15) \quad H_1 = -\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} x^2 + \frac{1}{2} R.$$

It follows from equations (2.10) and (2.11) that

$$(2.16) \quad H_0 \psi_n(x) = \left(n + \frac{1 - (-1)^n}{2} \right) \psi_n(x),$$

$$(2.17) \quad H_1 \psi_n(x) = \left(n + \frac{1 + (-1)^n}{2} \right) \psi_n(x).$$

The spectrum of H_0 consists only of the even numbers starting with zero. Each level is degenerate except for the ground states which is unique and the spectrum of H_1 consists only of the odd numbers and each level is degenerate. In [5, 15, 16], the author is used a very simple modification of supercharges (2.2) and Hamiltonian defined in (2.6) and (2.7), and showed that one can obtain a quantum system in the phase of the broken supersymmetry (with all the doubled energy levels organized in the equidistant positive spectrum) described by a linear superalgebra of the form $\mathfrak{eqrefadam}$. Moreover, in the same line one can obtain also the systems described by nonlinear supersymmetric algebra; see [15, 5, 16].

Instead of the supercharges Q_{i-} and Q_{i+} defined in (2.8), we can also reformulate the algebra (2.1) by introducing a new hermitean operators Q_i expressed in the form

$$(2.18) \quad Q_i = \frac{1}{\sqrt{2}} (Q_{i-} + Q_{i+}), \quad i = 0, 1.$$

Hermitian supercharges shown in (2.18), represent a half of the corresponding pairs of the bosonized supercharges for the systems (2.14) and (2.15). From equations (2.8) and (2.9) they share the property $Q_i^\dagger = Q_i$ and commute with the supersymmetric Hamiltonian H_i , so it can be diagonalized simultaneously with H_i . The supercharges Q_i act on the wave functions as

$$\begin{aligned} Q_0\psi_{2n}(x) &= \sqrt{n}\psi_{2n-1}(x), & Q_0\psi_{2n+1}(x) &= \sqrt{n+1}\psi_{2n+2}(x), \\ Q_1\psi_{2n}(x) &= \sqrt{n+1/2}\psi_{2n+1}(x), & Q_1\psi_{2n+1}(x) &= \sqrt{n+1/2}\psi_{2n}(x). \end{aligned}$$

Put

$$(2.19) \quad \psi_{n\varepsilon}^{(0)}(x) = \frac{1}{2}(\psi_{2n+1}(x) + \varepsilon\psi_{2n+2}(x)),$$

$$(2.20) \quad \psi_{n\varepsilon}^{(1)}(x) = \frac{1}{2}(\psi_{2n}(x) + \varepsilon\psi_{2n+1}(x)), \quad \varepsilon = \pm 1.$$

It is readily seen that

$$\begin{aligned} Q_0\psi_{n\varepsilon}^{(0)}(x) &= \varepsilon\sqrt{n+1}\psi_{n\varepsilon}^{(0)}(x), \\ Q_1\psi_{n\varepsilon}^{(1)}(x) &= \varepsilon\sqrt{n+1/2}\psi_{n\varepsilon}^{(1)}(x). \end{aligned}$$

Note that the eigenfunctions $\psi_{n\varepsilon}^{(i)}(x)$ of the supercharges Q_i which are defined in equation (2.19) and (2.20) involve the lacunary Hermite polynomials $H_{2n}(x)$ and $H_{2n+1}(x)$.

2.1. Associated Laguerre and Hermite polynomials. The Hermite polynomials satisfy the three terms recurrence relation

$$(2.21) \quad \begin{aligned} 2xH_n(x) &= H_{n+1}(x) + 2(n+1)H_{n-1}(x) \quad (n > 0), \\ H_{-1}(x) &= 0 \quad H_0(x) = 1. \end{aligned}$$

The associated Hermite polynomials $H_n(x, c)$ are defined by the following three terms recurrence relation

$$(2.22) \quad \begin{aligned} 2xH_n(x, c) &= H_{n+1}(x, c) + 2(n+1+c)H_{n-1}(x, c) \quad (n > 0), \\ H_{-1}(x, c) &= 0 \quad H_0(x, c) = 1. \end{aligned}$$

They were introduced by Askey and Wimp in [17], where their weight functions and explicit formulas were also found. When $c > -1$, they satisfy the orthogonality relations

$$(2.23) \quad \int_{-\infty}^{\infty} \frac{H_n(x, c)H_m(x, c)}{|D_{-c}(ix\sqrt{2})|^2} dx = 2^n \sqrt{\pi} \Gamma(n+c+1) \delta_{nm}.$$

The function $D_{-c}(x)$ is a parabolic cylinder function

$$(2.24) \quad D_{2\nu}(2x) = 2^\nu e^{-x^2} \Psi(-\nu, 1/2; 2x^2),$$

and $\Psi(a, c; x)$ is the Tricomi function [18],

$$(2.25) \quad \Psi(a, c; x) = \frac{1}{\Gamma(a)} \int_0^\infty e^{-xt} t^{a-1} (1+t)^{c-a-1} dt, \quad \Re(a) > 0, \Re(x) > 0.$$

Note that the polynomials $H_n(x, c)$ can be expressed in terms of the associated Laguerre polynomials $L_n^\nu(x, c)$ and $\mathcal{L}_n^\nu(x, c)$, which are defined by the three terms recurrence relations [19]

$$(2.26) \quad (2n+2c+\nu+1-x)L_n^\nu(x, c) = (n+c+1)L_{n+1}^\nu(x, c) + (n+c+\nu)L_{n-1}^\nu(x, c)$$

$$(2.27) \quad L_0^\nu(x, c) = 1, \quad L_1^\nu(x, c) = \frac{2c+\nu+1-x}{c+1}$$

and

$$(2.28) \quad (2n + 2c + \nu + 1 - x)\mathcal{L}_n^\nu(x, c) = (n + c + 1)\mathcal{L}_{n+1}^\nu(x, c) + (n + c + \nu)\mathcal{L}_{n-1}^\nu(x, c)$$

$$(2.29) \quad \mathcal{L}_0^\nu(x, c) = 1, \quad \mathcal{L}_1^\nu(x, c) = \frac{c + \nu + 1 - x}{c + 1}$$

and have the orthogonality relations

$$(2.30) \quad \int_0^\infty L_n^\nu(x, c)L_m^\nu(x, c) x^\nu e^{-x} \frac{|\Psi(c, 1 - \nu; xe^{-i\pi})|^{-2}}{\Gamma(c + 1)\Gamma(\nu + c + 1)} dx = \frac{(\nu + c + 1)_n}{(\nu + 1)_n} \delta_{nm}$$

and

$$(2.31) \quad \int_0^\infty \mathcal{L}_n^\nu(x, c)\mathcal{L}_m^\nu(x, c) x^\nu e^{-x} \frac{|\Psi(c, -\nu; xe^{-i\pi})|^{-2}}{\Gamma(c + 1)\Gamma(\nu + c + 1)} dx = \frac{(\nu + c + 1)_n}{(\nu + 1)_n} \delta_{nm}.$$

By [19], we explicitly have

$$(2.32) \quad H_{2n}(x, c) = (-4)^n (1 + c/2)_n L_n^{1/2}(x^2, c/2),$$

$$(2.33) \quad H_{2n+1}(x, c) = 2x(-4)^n (1 + c/2)_n \mathcal{L}_n^{-1/2}(x^2, c/2).$$

For example, the first polynomials are

$$(2.34) \quad H_0(x, c) = 1, \quad H_1(x, c) = 2x,$$

$$(2.35) \quad H_2(x, c) = 4x^2 - 2(c + 1), \quad H_3(x, c) = 8x^3 - 4(2c + 3)x.$$

Furthermore,

$$(2.36) \quad H_n(x, c) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-2)^k (c)_k (n - k)!}{k! (n - 2k)!} H_{n-2k}(x).$$

The associated Laguerre and Hermite polynomials are also birth and death processes polynomials, see Ismail, Letessier and Valent [19].

The lacunary Hermite polynomials $H_{nr+s}(x)$ may be defined by the following generating function [20]:

$$(2.37) \quad G(r, s) = \sum_{n=0}^{\infty} H_{nr+s}(x) \frac{t^{nr+s}}{(nr + s)!},$$

for arbitrary integers $r \geq 1$ and $0 \leq s \leq r - 1$. The specific generating functions are

$$G(1, 0) = e^{-t^2 + 2xt},$$

$$G(2, 0) = e^{-t^2} \cosh(2xt),$$

$$G(2, 1) = e^{-t^2} \sinh(2xt).$$

Note that $G(1, 0)$ is the generating function of the Hermite polynomials, $G(2, 0)$ and $G(2, 1)$ are linear combinations of $G(1, 0)$ (the same function with x changed by $-x$). Physically, these generating functions are very useful to describe the decomposition of ordinary, even and odd coherent states [21]. They also appeared in a number of problems, including for example the treatment of Cauchy problems in partial differential equations [22, 23] and in calculations involving coherent and squeezed states [23].

3. REALIZATION OF FRACTIONAL SUPERSYMMETRIC QUANTUM MECHANICS

Following Khare [10], a fractional supersymmetric quantum mechanics model of arbitrary order r can be developed by generalizing the fundamental equations (2.1) to the forms

$$(3.1) \quad Q_{\pm}^r = 0, \quad [H, Q_{\pm}] = 0, \quad Q_{-}^{\dagger} = Q_{+},$$

$$(3.2) \quad Q_{-}^{r-1}Q_{+} + Q_{-}^{r-2}Q_{+}Q_{-} + \cdots + Q_{-}Q_{+}Q_{-}^{r-2} + Q_{+}Q_{-}^{r-1} = (r-1)Q_{-}^{r-2}H.$$

It was shown in [11] that the parasupercharges Q_{-} and Q_{+} can be represented by $r \times r$ matrices as natural extensions of the $r = 2$ scheme. Note that the fractional supersymmetric quantum mechanics was also developed without an explicit introduction of parafermionic degrees of freedom by Plyushchay [3], see also [4, 26].

We shall now present a construction of fractional supersymmetric quantum mechanics of order r by using a Klein type operator. Let \mathcal{F} be the Fock space of states for the ordinary bosonic oscillator generated by the complete set of orthonormal vectors $\{|n\rangle\}_{n=0}^{\infty}$:

$$(3.3) \quad |n\rangle = \frac{1}{\sqrt{n!}}a_{+}^n|0\rangle, \quad n = 1, 2, \dots,$$

constructed over the vacuum state defined by the relation,

$$a_{-}|0\rangle = 0.$$

The creation and annihilation operators a_{+} and a_{-} may be realized in Fock space as

$$(3.4) \quad a_{-}|n\rangle = \sqrt{n}|n-1\rangle \quad \text{and} \quad a_{+}|n\rangle = \sqrt{n+1}|n+1\rangle,$$

The number operator N , which counts the number of particles, or objects in general in a system, is defined by $N = a_{+}a_{-}$ and is represented as

$$N|n\rangle = n|n\rangle.$$

It satisfies the commutation relations

$$[a_{-}, N] = a_{-}, \quad [a_{+}, N] = -a_{+}.$$

Now, let $r \geq 2$ be a fixed integer. We introduce the generalized Klein operator K defined by the relations (see, [24, 25])

$$(3.5) \quad K^r = 1, \quad a_{-}K = \varepsilon_r K a_{-} \quad a_{+}K = \varepsilon_r^{-1} K a_{+}, \quad K^{\dagger} = K^{-1},$$

where $\varepsilon_r = e^{\frac{2i\pi}{r}}$ is a primitive root of unity. The operator K can be realized by using the number operator N in the form $K = e^{\frac{2i\pi}{r}N}$, and acts on the space \mathcal{F} as

$$(3.6) \quad K|n\rangle = \varepsilon_r^n |n\rangle,$$

and so introduces \mathbb{Z}_r -grading structure on the Fock space \mathcal{F} as

$$(3.7) \quad \mathcal{F} = \bigoplus_{j=0}^{r-1} \mathcal{F}_j,$$

where

$$\mathcal{F}_j = \{|nr + j\rangle : n = 0, 1, \dots\}.$$

For $j = 0, 1, \dots, r-1$, we denote by Π_j , the orthogonal projection from \mathcal{F} onto its subspace \mathcal{F}_j , which can be represented as

$$(3.8) \quad \Pi_j = \frac{1}{r} \sum_{l=0}^{r-1} \varepsilon_r^{-lj} K^l.$$

Equivalently, the operators K^l are expressed in terms of Π_j as follows

$$(3.9) \quad K^l = \sum_{j=0}^{r-1} \varepsilon_r^{-lj} \Pi_j, \quad l = 0, \dots, r-1.$$

It is clear that they form a system of resolution of the identity:

$$(3.10) \quad \Pi_0 + \Pi_1 + \dots + \Pi_{r-1} = 1 \quad \text{and} \quad \Pi_i \Pi_j = \delta_{ij} \Pi_i.$$

The action of Π_j on \mathcal{F} can be taken to be

$$(3.11) \quad \Pi_j |kr + l\rangle = \delta_{jl} |kr + l\rangle.$$

3.1. Supercharges. Following [3, 4, 12], for every fixed integer j such that $0 \leq j \leq r-1$, one can construct two supercharges Q_{j-} and Q_{j+} by means of the orthogonal projections Π_k and the creation and annihilation operators a_+ and a_- introduced before as follows

$$(3.12) \quad Q_{j-} = a_-(1 - \Pi_j) = (1 - \Pi_{j-1})a_-,$$

$$(3.13) \quad Q_{j+} = a_+(1 - \Pi_{j-1}) = (1 - \Pi_j)a_+.$$

Obviously, the operators a_- , a_+ , Q_{j-} and Q_{j+} satisfy the intertwining relations

$$(3.14) \quad \Pi_{j-1}a_- = a_- \Pi_j \quad \text{and} \quad a_+ \Pi_{j-1} = \Pi_j a_+$$

$$(3.15) \quad Q_{j-}a_- = a_- Q_{j+1-} \quad \text{and} \quad a_+ Q_{j+} = Q_{j+1+} a_+.$$

It is easy to see that the supercharges Q_{j-} and Q_{j+} have the Hermitian conjugation relations

$$(3.16) \quad Q_{j-}^\dagger = Q_{j+}.$$

Furthermore, the operators a_- and a_+ can be represented as

$$(3.17) \quad a_- = \frac{1}{r-1} \sum_{j=0}^{r-1} Q_{j-} \quad \text{and} \quad a_+ = \frac{1}{r-1} \sum_{j=0}^{r-1} Q_{j+}.$$

Now, by making use of relations (3.14) and (3.15), we easily obtain

$$Q_{j-}^k = \begin{cases} a_-^k (1 - \Pi_j - \dots - \Pi_{j+k}) & \text{if } k \leq r-1-j \\ a_-^k (1 - \Pi_j - \dots - \Pi_{N-1} - \dots - \Pi_{k+j-r-1}) & \text{if } r-j \leq k \leq r-1. \end{cases}$$

In particular, for $k = r$, we get

$$(3.18) \quad Q_{j-}^r = a_-^r (1 - \Pi_j - \dots - \Pi_{r-1} - \dots - \Pi_{j-1}) = 0.$$

Similarly, we have

$$Q_{j+}^r = 0.$$

Hence, the supercharge operators $Q_{j\pm}$ are nilpotent operators of order r .

3.2. Supersymmetric Hamiltonian of order r . We can use the operators a_- , a_+ and the orthogonal projections $\Pi_0, \Pi_1, \dots, \Pi_{r-1}$ introduced before for obtaining a realization of supersymmetric Hamiltonian.

The first main result, is

Theorem 3.1. *For fixed $j = 0, \dots, r-1$, the hermitian operators Q_{j-} , Q_{j+} and H_j given by*

$$Q_{j+} = a_+(1 - \Pi_{j-1}), \quad Q_{j-} = a_-(1 - \Pi_j),$$

$$H_j = a_+a_- + \sum_{k=0}^{r-1} (1 + k - r/2) \Pi_{r+j-k-1}$$

satisfy the commutation relations

$$Q_{j\pm}^r = 0, \quad [H, Q_{j\pm}] = 0, \quad Q_{j-}^\dagger = Q_{j+},$$

$$Q_{j-}^{r-1}Q_{j+} + Q_{j-}^{r-2}Q_{j+}Q_{j-} + \dots + Q_{j-}Q_{j+}Q_{j-}^{r-2} + Q_{j+}Q_{j-}^{r-1} = (r-1)Q_{j-}^{r-2}H_j.$$

Furthermore, the superoscillator H_j has r -fold degenerate spectrum and acts on the occupation basis as

$$H_j |nr + j + s\rangle = (rn + r/2 + j) |nr + j + s\rangle.$$

Proof. We first summarize the key properties [12]:

$$(3.19) \quad Q_{j+}Q_{j-}^{r-1} = a_+a_-^{r-1}\Pi_{j-2},$$

$$(3.20) \quad Q_{j-}^{r-1}Q_{j+} = a_-^{r-1}a_+\Pi_{j-1},$$

$$(3.21) \quad Q_{j-}^{r-1-k}Q_{j+}Q_{j-}^k = a_-^{r-1-k}a_+a_-^k(\Pi_{j-2} + \Pi_{j-1}), \quad k = 1, \dots, r-2.$$

From the relation $a_-^k a_+ = a_+ a_-^k + k a_-^{k-1}$ and relations (3.19), (3.20) and (3.21), we have

$$(3.22) \quad \sum_{k=0}^{r-1} Q_{j-}^{r-1-k}Q_{j+}Q_{j-}^k = (r-1)Q_{j-}^{r-2}[a_+a_- + (2 - \frac{r}{2})\Pi_{j-2}(r) + (1 - \frac{r}{2})\Pi_{j-1}].$$

On the other hand, we get from (3.10) that

$$(3.23) \quad Q_{j-}^{r-2} \sum_{k=2}^{r-1} (1 + k - \frac{r}{2}) \Pi_{r+j-k-1} = 0.$$

This yields

$$(3.24) \quad \sum_{k=0}^{r-1} Q_{j-}^{r-1-k}Q_{j+}Q_{j-}^k = (r-1)Q_{j-}^{r-2}H_j,$$

where the operator H_j is given explicitly by

$$(3.25) \quad H_j = a_+a_- + \sum_{k=0}^{r-1} (1 + k - \frac{r}{2}) \Pi_{r+j-k-1}.$$

It remains to prove that

$$(3.26) \quad [H, Q_{j-}] = [H, Q_{j+}] = 0.$$

Indeed, from (3.21), we have

$$\begin{aligned}
 Q_{j-}H &= a_-a_+a_-(1 - \Pi_j) + \sum_{k=0}^{r-2} (1 + k - \frac{r}{2}) a_-(1 - \Pi_j)\Pi_{r+j-k-1} \\
 &= a_+a_-a_-(1 - \Pi_j) + \sum_{k=0}^{r-2} (2 + k - \frac{r}{2}) \Pi_{r+j-k-2}(1 - \Pi_{j-1})a_- \\
 &= a_+a_-a_-(1 - \Pi_j) + \sum_{k=1}^{r-1} (1 + k - \frac{r}{2}) \Pi_{r+j-k-1}(1 - \Pi_{j-1})a_- \\
 &= HQ_{j-}.
 \end{aligned}$$

Thus,

$$(3.27) \quad [H_j, Q_{j-}] = 0.$$

Since the orthogonal projection Π_k is hermitian conjugate $\Pi_k^\dagger = \Pi_k$, we easily check that the operators H_j share the property

$$(3.28) \quad H_j^\dagger = H_j.$$

Combining this with (3.27), we get

$$(3.29) \quad [H_j, Q_{j+}] = 0$$

Given representation (3.4), the action of operators Q_{j-} and Q_{j+} reduces to

$$(3.30) \quad Q_{j-}|nr + j + s\rangle = \sqrt{nr + j + s}|nr + j + s - 1\rangle, \quad s = 1, \dots, r - 1,$$

$$(3.31) \quad Q_{j+}|nr + j + s\rangle = \sqrt{nr + j + s + 1}|nr + j + s + 1\rangle, \quad s = 0, \dots, r - 2,$$

$$(3.32) \quad Q_{j-}|nr + j\rangle = 0, \quad Q_{j+}|(n + 1)r + j - 1\rangle = 0$$

and hence,

$$(3.33) \quad H_j | nr + j + s\rangle = (rn + r/2 + j) | nr + j + s\rangle.$$

□

4. SIMULTANEOUS DIAGONALIZATION

Our second result is the following.

Theorem 4.1. *For fixed integer $j = 0, \dots, r - 1$, the operator Q_j defined by*

$$(4.1) \quad Q_j = \frac{1}{\sqrt{2}}(Q_{j-} + Q_{j+}),$$

is self-adjoint and commutes with the oscillator H_j , that is

$$(4.2) \quad Q_j^\dagger = Q_j, \quad [Q_j, H_j] = 0.$$

Furthermore, for every integer n , the operator Q_j leaves the subspace spanned by the states $\{|nr + j\rangle, \dots, |(n + 1)r + j - 1\rangle\}$ invariant and restricted operator has a nondegenerate spectrum which consists from the roots of the associated Hermite polynomials $H_r(x, c)$, with $c = nr + j - 1$.

and this provides the following inverse relation of (4.9):

$$(4.12) \quad |nr + j + k\rangle = \sum_{s=0}^{r-1} \sqrt{w_s} P_k(x_s(r, n, j)) |n, j, s\rangle.$$

The monic orthogonal polynomials $\tilde{P}_k(x)$, related to $P_k(x)$, satisfy the three term recurrence relation

$$(4.13) \quad x\tilde{P}_k(x) = \tilde{P}_{k+1}(x) + (nr + j + k)/2 \tilde{P}_{k-1}(x) \quad k = 0, \dots, r-1,$$

$$(4.14) \quad \tilde{P}_{-1}(x) = 0, \quad \tilde{P}_0(x) = 1.$$

From the three terms recurrence relations (2.22), we see that the polynomials $\tilde{P}_k(x)$ satisfy (4.13) and (4.14) and can be identified with the associated Hermite polynomial $H_k(x, c)$, with $c = nr + j - 1$. Namely, we have

$$(4.15) \quad \tilde{P}_k(x) = 2^{-k} H_k(x, nr + j - 1), \quad k = 0, 1, \dots, r-1.$$

It is well-known from the theory of orthogonal polynomials [19, 27] that the eigenvalues $x_s(r, n, j)$ of the Jacobi matrix A_{nr} coincide with the roots of the characteristic polynomial $\tilde{P}_r(x)$ given by

$$(4.16) \quad \tilde{P}_r(x) = (x - x_0(r, n, j)) \dots (x - x_{r-1}(r, n, j))$$

and that the discrete weights w_s can be expressed by (see, [19])

$$(4.17) \quad w_s = \frac{\xi_{r-1}}{H_{r-1}(x_s(r, n, j), nr + j - 1) H'_r(x_s(r, n, j), nr + j - 1)}, \quad s = 0, 1, \dots, r-1,$$

where the normalization constants are given by

$$(4.18) \quad \xi_{r-1} = 2^r \frac{\Gamma(r + nr + j - 1)}{\Gamma(nr + j)}.$$

□

5. CONCLUSION

We considered a fractional supersymmetric quantum mechanics of finite order having a structure similar to one constructed by Plyushchay [3, 4]. The Hamiltonian and the supercharges involve reflection operator. We showed this system to be exactly solvable and found that its wave functions are expressed in terms of lacunary Hermite polynomials and associated Hermite polynomials.

REFERENCES

- [1] E. Witten, Dynamical Breaking of Supersymmetry, Nucl. Phys. B **188** (1981) 513.
- [2] H. Nicolai, Supersymmetry and Spin Systems, J. Phys. A **9**, 1497 (1976).
- [3] M. S. Plyushchay, Supersymmetry without fermions, arXiv: hep-th/9404081; Deformed Heisenberg algebra, fractional spin fields and supersymmetry without fermions, Ann. Phys. (N.Y.) **245**, 339 (1996).
- [4] M. S. Plyushchay, Deformed Heisenberg algebra with reflection, Nucl. Phys. B **491**, 619 (1997).
- [5] F. Correa and M. S. Plyushchay, Hidden supersymmetry in quantum bosonic systems, Annals Phys. **322**, 2493 (2007).
- [6] J. Gamboa, M. Plyushchay and J. Zanelli, Three aspects of bosonized supersymmetry and linear differential field equation with reflection, Nucl. Phys. B **543**, 447 (1999).
- [7] M. Rösler, Dunkl operators: theory and applications *Orthogonal Polynomials and Special Functions (Lecture Notes in Mathematics)* vol 1817 (Berlin: Springer)(2003) pp 93–135.

- [8] C. F. Dunkl and Y. Xu, (2001) Orthogonal polynomials of several variables *Encyclopedia of Mathematics and Its Applications* vol 81 (Cambridge: Cambridge University Press)
- [9] S. Post, L. Vinet and A. Zhedanov, Supersymmetric Quantum Mechanics with Reflections, *J. Phys. A* **44**, 435301 (2011).
- [10] A. Khare, Parasupersymmetric quantum mechanics of arbitrary order, *J. Phys. A* **25**, L749 (1992).
- [11] F. Cooper, A. Khare and U. Sukhatme, 2001 *Supersymmetry in Quantum Mechanics* (Singapore: World Scientific)
- [12] M. Daoud and M. Kibler, in *Symmetry and Structural Properties of Condensed Matter*, edited by T. Lulek, B. Lulek and A. Wal (World Scientific, Singapore, 2001); in *Proceedings of the Sixth International Wigner Symposium* (Bogazici Univ. Press, Istanbul, Turkey, 2002).
- [13] C. Quesne and N. Vansteenkiste, $C(\lambda)$ extended harmonic oscillator and (para)supersymmetric quantum mechanics, *Phys. Lett. A* **240**, 21 (1998).
- [14] V. Jakubsky, L. M. Nieto and M. S. Plyushchay, The origin of the hidden supersymmetry, *Phys. Lett. B* **692**, 51 (2010).
- [15] M. Plyushchay, Hidden nonlinear supersymmetries in pure parabosonic systems, *Int. J. Mod. Phys. A* **15**, 3679 (2000).
- [16] F. Correa, L. M. Nieto and M. S. Plyushchay, Hidden nonlinear supersymmetry of finite-gap Lamé equation, *Phys. Lett. B* **644**, 94 (2007).
- [17] R. Askey and J. Wimp, Associated Laguerre and Hermite polynomials, *Proc. Roy. Soc. Edinburgh Sect. A* **96** (1984), no. 1–2, 15–37. MR 741641.
- [18] A. Erdélyi, W. Magnus, F. Oberhettinger and G. F. Tricomi, (1953) *Higher Transcendental Functions*, volume 1. New York: McGraw-Hill.
- [19] M. E. H. Ismail, *Classical and quantum orthogonal polynomials in one variable*, *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, November 2005. ISBN 0521782015.
- [20] I. M. Gessel and P. Jayawant, A triple lacunary generating function for Hermite polynomials, *Electronic J. Comb.* **12** (2005) R30.
- [21] M. M. Nieto, (1998) The discovery of squeezed states in 1927 5th Int. Conf. on Squeezed States and Uncertainty Relations (Balatonfüred, 1997) (NASA Conference Publication NASA/CP-1998-206855) ed D Han, J Janszky, Y S Kim and V I Man'ko (Greenbelt: Goddard Space Flight Center) pp 175–80.
- [22] K. A. Penson, K. Górska, A. Horzela, and G. Dattoli. Quasi-relativistic heat equation via Lévy stable distributions: Exact solutions. *Annalen der Physik*, **530**(3):1700374, 2018. Goddard Space Flight Center) pp 175–80.
- [23] D. Babusci, G. Dattoli, K. Górska, and K.A. Penson. Lacunary generating functions for the Laguerre polynomials. *Séminaire Lotharingien de Combinatoire*, **76**:B76b, 2017.
- [24] F. Bouzeffour, M. Garayev, *Supersymmetric Quantum mechanics on the radial lines*, arXiv:1811.02151.
- [25] F. Bouzeffour, W. Jedidi and M. Garayev, Extended Arithmetic functions. Accepted for publication in *The Ramanujan Journal* DOI: 10.1007/s11139-018-0122-8.
- [26] S. Klishevich and M. Plyushchay, “Supersymmetry of parafermions,” *Mod. Phys. Lett. A* **14**, 2739 (1999).
- [27] T. Chihara, *An introduction to orthogonal polynomials*. Dover Books on Mathematics. Dover Publications, reprint edition, 2011.
- [28] D. Foata, Some Hermite polynomial identities and their combinatorics, *Adv. in Appl. Math.* **2** (1981), 250–259.

* DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCES, KING SAUD UNIVERSITY, P. O BOX 2455 RIYADH 11451, SAUDI ARABIA AND COLLEGE OF SCIENCES, UNIVERSITY OF CARTHAGE, B. O. BOX 64 JARZOUNA 7021, BIZERTE, TUNISIA.

\diamond DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCES, KING SAUD UNIVERSITY, P. O BOX 2455 RIYADH 11451, SAUDI ARABIA AND COLLEGE OF SCIENCES
E-mail address: fbouzeffour@ksu.edu.sa, mgarayev@ksu.edu.sa