# Canonicity and normalization for Dependent Type Theory

Thierry Coquand

Computer Science Department, University of Gothenburg

# Introduction

We show canonicity and normalization for dependent type theory with a cumulative sequence of universes  $U_0: U_1 \ldots$  with  $\eta$ -conversion. We give the argument in a constructive set theory  $\operatorname{CZFu}_{<\omega}$ , designed by P. Aczel [2]. We provide a purely algebraic presentation of a canonicity proof, as a way to build new (algebraic) models of type theory. We then present a normalization proof, which is technically more involved, but is based on the same idea. We believe our argument to be a simplification of existing proofs [15, 16, 1, 7], in the sense that we never need to introduce a reduction relation, and the proof theoretic strength of our meta theory is as close as possible to the one of the object theory [2, 9].

Let us expand these two points. If we are only interested in *canonicity*, i.e. to prove that a closed Boolean is convertible to 0 or 1, one argument for simple type theory (as presented e.g. in [19]) consists in defining a "reducibility"<sup>1</sup> predicate by induction on the type. For the type of Boolean, it means exactly to be convertible to 0 or 1, and for function types, it means that it sends a reducible argument to a reducible value. It is then possible to show by induction on the typing relation that any closed term is reducible. In particular, if this term is a Boolean, we obtain canonicity. The problem of extending this argument for a dependent type system with universes is in the definition of what should be the reducibility predicate for universes. It is natural to try an inductive-recursive definition; this was essentially the way it was done in [15], which is an early instance of an inductive-reductive definition. We define when an element of the universe is reducible, and, by induction on this proof, what is the associated reducibility predicate for the type represented by this element. However, there is a difficulty in this approach: it might well be *a priori* that an element is both convertible for instance to the type of Boolean or of a product type, and if this is the case, the previous inductive-recursive definition is ambiguous.

In [15], this problem is solved by considering first a reduction relation, and then showing this reduction relation to be confluent, and defining convertibility as having a commun reduct. This does *not* work however when conversion is defined as a *judgement* (as in [16, 1]). This is an essential difficulty, and a relatively subtle and complex argument is involved in [1, 7] to solve it: one defines first an *untyped* reduction relation and a reducibility *relation*, which is used first to establish a confluence property.

The main point of this paper is that this essential difficulty can be solved, in a seemingly magical way, by considering *proof-relevant* reducibility, that is where reducibility is defined as a *structure* and not only as a *property*. Such an approach is hinted in the reference [16], but [16] still introduces a reduction relation, and also presents a version of type theory with a restricted form of conversion (no conversion under abstraction, and no  $\eta$ -conversion; this restriction is motivated in [17]).

Even for the base type, reducibility is a structure: the reducibility structure of an element t of Boolean type contains either 0 (if t and 0 are convertible) or 1 (if t and 1 are convertible) and this might a priori contains both 0 and 1. Another advantage of our approach, when defining reducibility in a proof-relevant way, is that the required meta-language is weaker than the one used for a reducibility relation (where one has to do proofs by induction on this reducibility relation).

Yet another aspect that was not satisfactory in previous attempts [1, 7] is that it involved essentially a *partial equivalence relation model*. One expects that this would be needed for a type theory with an extensional equality, but not for the present version of type theory. This issue disappears here: we only consider *predicates* (that are proof-relevant).

<sup>&</sup>lt;sup>1</sup>The terminology for this notion seems to vary: in [12], where is was first introduced, it is called "berechenbarkeit", which can be translated by "computable", in [21] it is called "convertibility", and in [19] it is called "reducibility".

A more minor contribution of this paper is its *algebraic* character. For both canonicity and decidability of conversion, one considers first a general model construction and one obtains then the desired result by instantiating this general construction to the special instance of the initial (term) model, using in both cases only the abstract characteristic property of the initial model.

## **1** Informal presentation

We first give an informal presentation of the canonicity proof by first expliciting the rules of type theory and then explaining the reducibility argument,

#### 1.1 Type system

We use conversion as judgements [1]. Note that it is not clear a priori that subject reduction holds.

$$\begin{array}{ccc} \frac{\Gamma \vdash A:U_n}{\Gamma, x:A \vdash} & \frac{\Gamma \vdash}{\Gamma \vdash x:A} (x:A \ in \ \Gamma) \\ \\ \frac{\Gamma \vdash A:U_n}{\Gamma \vdash \Pi(x:A)B:U_n} & \frac{\Gamma, x:A \vdash t:B}{\Gamma \vdash \lambda(x:A)t:\Pi(x:A)B} & \frac{\Gamma \vdash t:\Pi(x:A)B}{\Gamma \vdash t \ u:B(u)} \\ \\ \frac{\Gamma \vdash A:U_n}{\Gamma \vdash A:U_m} (n \leqslant m) & \frac{\Gamma \vdash U_n:U_m}{\Gamma \vdash U_n:U_m} (n < m) & \frac{\Gamma \vdash N_2:U_n}{\Gamma \vdash N_2:U_n} \end{array}$$

The conversion rules are

$$\begin{array}{c} \displaystyle \frac{\Gamma \vdash t:A \quad \Gamma \vdash A \ \mathrm{conv} \ B:U_n}{\Gamma \vdash t:B} & \displaystyle \frac{\Gamma \vdash t \ \mathrm{conv} \ u:A \quad \Gamma \vdash A \ \mathrm{conv} \ B:U_n}{\Gamma \vdash t \ \mathrm{conv} \ u:B} \\ \\ \displaystyle \frac{\Gamma \vdash t:A \quad \Gamma \vdash t \ \mathrm{conv} \ t:A \quad \Gamma \vdash u \ \mathrm{conv} \ v:A \quad \Gamma \vdash u \ v:A \quad v:A \quad \Gamma \vdash u \ v:A \quad v:A \quad \Gamma \vdash u \ v:A \quad v:A \quad v:A \quad \Gamma \vdash u \ v:A \quad v:$$

We consider type theory with  $\eta$ -rules

$$\frac{\Gamma \vdash t: \Pi(x:A)B \quad \Gamma \vdash u: \Pi(x:A)B \quad \Gamma, x:A \vdash t \ x \ \text{conv} \ u \ x:B}{\Gamma \vdash t \ \text{conv} \ u: \Pi(x:A)B}$$

Finally we add  $N_2: U_1$  with the rules

$$\frac{\Gamma \vdash 0: N_2}{\Gamma \vdash 1: N_2} \qquad \frac{\Gamma, x: N_2 \vdash C: U_n \quad \Gamma \vdash a_0: C(0) \quad \Gamma \vdash a_1: C(1)}{\Gamma \vdash \mathsf{brec} \ (\lambda x.C) \ a_0 \ a_1: \Pi(x: N_2)C}$$

with computation rules brec  $(\lambda x.C) a_0 a_1 0$  conv  $a_0 : C(0)$  and brec  $(\lambda x.C) a_0 a_1 1$  conv  $a_1 : C(1)$ .

### 1.2 Reducibility proof

The informal reducibility proof consists in associating to each closed expression a of type theory (treating equally types and terms) an abstract object a' which represents a "proof" that a is reducible. If A is a (closed) type, then A' is a family of sets over the set Term(A) of closed expressions of type A modulo conversion. If a is of type A then a' is an element of the set A'(a).

The metatheory is a (constructive) set theory with a commulative hierarchy of universes  $\mathcal{U}_n$  [2]. This is defined by structural induction on the expression as follows

• (c a)' is c' a a'

- $(\lambda(x : A)b)'$  is the function which takes as arguments a closed expression a of type A and an element a' in A'(a) and produces b'(a, a')
- $(\Pi(x : A)B)'(w)$  for w closed expression of type  $\Pi(x : A)B$  is the set  $\Pi(a : \text{Term}(A))(a' : A'(a))B'(a,a')(w a)$
- $N'_{2}(t)$  is the set  $\{0 \mid t \text{ conv } 0\} \cup \{1 \mid t \text{ conv } 1\}$
- $U'_n(A)$  is the set  $\operatorname{\mathsf{Term}}(A) \to \mathcal{U}_n$

It can then be shown<sup>2</sup> that if a : A then a' is an element of A'(a) and furthermore that if  $a \operatorname{conv} b : A$  then a' = b' in A'(a) = A'(b). In particular, if  $a : N_2$  then a' is 0 or 1 and we get that a is convertible to 0 and 1.

One feature of this argument is that the required meta theory, here constructive set theory, is known to be of similar strength as the corresponding type theory; for a term involving n universes, the meta theory will need n + 1 universes [9]. This is to be contrasted with the arguments in [15, 1, 7] involving induction recursion which is a much stronger principle.

We believe that the mathematical purest way to formulate this argument is an algebraic argument, giving a (generalized) algebraic presentation of type theory. We then use only of the *term* model the fact that it is the *initial* model of type theory. This is what is done in the next section.

# 2 Model and syntax of dependent type theory with universes

### 2.1 Cumulative categories with families

We present a slight variation (for universes) of the notion of "category" with families  $[10]^3$ . A model is given first by a class of *contexts*. If  $\Gamma, \Delta$  are two given contexts we have a set  $\Delta \to \Gamma$  of *substitutions* from  $\Delta$  to  $\Gamma$ . These collections of sets are equipped with operations that satisfy the laws of composition in a category: we have a substitution 1 in  $\Gamma \to \Gamma$  and a composition operator  $\sigma\delta$  in  $\Theta \to \Gamma$  if  $\delta$  is in  $\Theta \to \Delta$  and  $\sigma$  in  $\Delta \to \Gamma$ . Furthermore we should have  $\sigma 1 = 1\sigma = \sigma$  and  $(\sigma\delta)\theta = \sigma(\delta\theta)$  if  $\theta: \Theta_1 \to \Theta$ .

We assume to have a "terminal" context (): for any other context, there is a unique substitution, also written (), in  $\Gamma \to ($ ). In particular we have () $\sigma = ($ ) in  $\Delta \to ($ ) if  $\sigma$  is in  $\Delta \to \Gamma$ .

We write  $|\Gamma|$  the set of substitutions ()  $\rightarrow \Gamma$ .

If  $\Gamma$  is a context we have a cumulative sequence of sets  $\mathsf{Type}_n(\Gamma)$  of  $types \ over \ \Gamma$  at level n (where n is a natural number). If A in  $\mathsf{Type}_n(\Gamma)$  and  $\sigma$  in  $\Delta \to \Gamma$  we should have  $A\sigma$  in  $\mathsf{Type}_n(\Delta)$ . Furthermore A1 = A and  $(A\sigma)\delta = A(\sigma\delta)$ . If A in  $\mathsf{Type}_n(\Gamma)$  we also have a collection  $\mathsf{Elem}(\Gamma, A)$  of elements of type A. If a in  $\mathsf{Elem}(\Gamma, A)$  and  $\sigma$  in  $\Delta \to \Gamma$  we have  $a\sigma$  in  $\mathsf{Elem}(\Delta, A\sigma)$ . Furthermore a1 = a and  $(a\sigma)\delta = a(\sigma\delta)$ . If A is in  $\mathsf{Type}_n()$  we write |A| the set  $\mathsf{Elem}(), A)$ .

We have a *context extension operation*: if A is in  $\mathsf{Type}_n(\Gamma)$  then we can form a new context  $\Gamma.A$ . Furthermore there is a projection  $\mathsf{p}$  in  $\Gamma.A \to \Gamma$  and a special element  $\mathsf{q}$  in  $\mathsf{Elem}(\Gamma.A, A\mathsf{p})$ . If  $\sigma$  is in  $\Delta \to \Gamma$  and A in  $\mathsf{Type}_n(\Gamma)$  and a in  $\mathsf{Elem}(\Delta, A\sigma)$  we have an extension operation  $(\sigma, a)$  in  $\Delta \to \Gamma.A$ . We should have  $\mathsf{p}(\sigma, a) = \sigma$  and  $\mathsf{q}(\sigma, a) = a$  and  $(\sigma, a)\delta = (\sigma\delta, a\delta)$  and  $(\mathsf{p}, \mathsf{q}) = 1$ .

If a is in  $\mathsf{Elem}(\Gamma, A)$  we write  $\langle a \rangle = (1, a)$  in  $\Gamma \to \Gamma.A$ . Thus if B is in  $\mathsf{Type}_n(\Gamma.A)$  and a in  $\mathsf{Elem}(\Gamma, A)$  we have  $B\langle a \rangle$  in  $\mathsf{Type}_n(\Gamma)$ . If furthermore b is in  $\mathsf{Elem}(\Gamma.A, B)$  we have  $b\langle a \rangle$  in  $\mathsf{Elem}(\Gamma, B\langle a \rangle)$ .

A global type of level n is given by a an element C in  $\mathsf{Type}_n()$ . We write simply C instead of C() in  $\mathsf{Type}_n(\Gamma)$  for () in  $\Gamma \to ()$ . Given such a global element C, a global element of type C is given by an element c in  $\mathsf{Elem}((), C)$ . We then write similarly simply c instead of c() in  $\mathsf{Elem}(\Gamma, C)$ .

Models are sometimes presented by giving a class of special maps (fibrations), where a type are modelled by a fibration and elements by a section of this fibration. In our case, the fibrations are the maps  $\mathbf{p}$  in  $\Gamma.A \to \Gamma$ , and the sections of these fibrations correspond exactly to elements in  $\mathsf{Elem}(\Gamma, A)$ . Any element  $a \; \mathsf{Elem}(\Gamma, A)$  defines a section  $\langle a \rangle = (1, a) : \Gamma \to \Gamma.A$  and any such section is of this form.

 $<sup>^{2}</sup>$ We prove this statement by induction on the derivation and consider a more general statement involving a context; we don't provide the details in this informal part since this will be covered in the next section.

 $<sup>^{3}</sup>$ As emphasized in this reference, these models should be more exactly thought of as *generalized algebraic structures* rather than *categories*; e.g. the initial model is defined up to isomorphism and not up to equivalence). This provides a generalized algebraic notion of model of type theory.

#### 2.2 Dependent product types

A category with families has *product types* if we furthermore have one operation  $\Pi A B$  in  $\mathsf{Type}_n(\Gamma)$  for A is in  $\mathsf{Type}_n(\Gamma)$  and B is in  $\mathsf{Type}_n(\Gamma.A)$ . We should have  $(\Pi A B)\sigma = \Pi (A\sigma) (B\sigma^+)$  where  $\sigma^+ = (\sigma \mathsf{p}, \mathsf{q})$ . We have an abstraction operation  $\lambda b$  in  $\mathsf{Elem}(\Gamma, \Pi A B)$  given b in  $\mathsf{Elem}(\Gamma.A, B)$ . We have an application operation such that  $\mathsf{app}(c, a)$  is in  $\mathsf{Elem}(\Gamma, B\langle a \rangle)$  if a is in  $\mathsf{Elem}(\Gamma, A)$  and c is in  $\mathsf{Elem}(\Gamma, \Pi A B)$ . These operations should satisfy the equations

 $\mathsf{app}(\lambda b, a) = b \langle a \rangle \qquad c = \lambda (\mathsf{app} \ (c\mathsf{p}, \mathsf{q})) \qquad (\lambda b) \sigma = \lambda (b\sigma^+) \qquad \mathsf{app}(c, a) \sigma = \mathsf{app}(c\sigma, a\sigma)$ 

where we write  $\sigma^+ = (\sigma \mathbf{p}, \mathbf{q})$ .

#### 2.3 Cumulative universes

We assume to have global elements  $U_n$  in  $\mathsf{Type}_{n+1}(\Gamma)$  such that  $\mathsf{Type}_n(\Gamma) = \mathsf{Elem}(\Gamma, U_n)$ .

#### 2.4 Booleans

Finally we add the global constant  $N_2$  in  $\mathsf{Type}_0(\Gamma)$  and global elements 0 and 1 in  $\mathsf{Elem}(\Gamma, N_2)$ . Given T in  $\mathsf{Type}_n(\Gamma, N_2)$  and  $a_0$  in  $\mathsf{Elem}(\Gamma, T\langle 0\rangle)$  and  $a_1$  in  $\mathsf{Elem}(\Gamma, T\langle 1\rangle)$  we have an operation  $\mathsf{brec}(T, a_0, a_1)$  producing an element in  $\mathsf{Elem}(\Gamma, \Pi \ N_2 \ T)$  satisfying the equations  $\mathsf{app}(\mathsf{brec}(T, a_0, a_1), 0) = a_0$  and  $\mathsf{app}(\mathsf{brec}(T, a_0, a_1), 1) = a_1$ .

Furthermore,  $\operatorname{brec}(T, a_0, a_1)\sigma = \operatorname{brec}(T\sigma^+, a_0\sigma, a_1\sigma)$ .

# 3 Reducibility model

Given a model of type theory M as defined above, we describe how to build a new associated "reducibility" model  $M^*$ . When applied to the initial/term model  $M_0$ , this gives a proof of canonicity which can be seen as a direct generalization of the argument presented in [19] for Gödel system T. As explained in the introduction, the main novelty here is that we consider a proof-relevant notion of reducibility.

A context of  $\mathsf{M}^*$  is given by a context  $\Gamma$  of the model  $\mathsf{M}$  together with a family of sets  $\Gamma'(\rho)$  for  $\rho$  in  $|\Gamma|$ . A substitution in  $\Delta, \Delta' \to^* \Gamma, \Gamma'$  is given by a pair  $\sigma, \sigma'$  with  $\sigma$  in  $\Delta \to \Gamma$  and  $\sigma'$  in  $\Pi(\nu \in |\Delta|)\Delta'(\nu) \to \Gamma'(\sigma\nu)$ .

The identity substitution is the pair  $1^* = 1, 1'$  with  $1'\rho\rho' = \rho'$ . Composition is defined by  $(\sigma, \sigma')(\delta, \delta') = \sigma\delta, (\sigma\delta)'$  with

$$(\sigma\delta)'\alpha\alpha' = \sigma'(\delta\alpha)(\delta'\alpha\alpha')$$

The set  $\mathsf{Type}_n^*(\Gamma, \Gamma')$  is defined to be the set of pairs A, A' where A is in  $\mathsf{Type}_n(\Gamma)$  and  $A'\rho\rho'$  is in  $|A\rho| \to \mathcal{U}_n$ . We define then  $A'(\sigma, \sigma')\nu\nu' = A'(\sigma\nu)(\sigma'\nu\nu')$ .

We define  $\mathsf{Elem}^*(\Gamma, \Gamma')(A, A')$  to be the set of pairs a, a' where a is in  $\mathsf{Elem}(\Gamma, A)$  and  $a'\rho\rho'$  is in  $A'\rho\rho'(a\rho)$  for each  $\rho$  in  $|\Gamma|$  and  $\rho'$  in  $\Gamma'(\rho)$ . We define then  $(a, a')(\sigma, \sigma') = a\sigma, a'(\sigma, \sigma')$  with  $a'(\sigma, \sigma')\nu\nu' = a'(\sigma\nu)(\sigma'\nu\nu')$ .

The extension operation is defined by  $(\Gamma, \Gamma').(A, A') = \Gamma.A, (\Gamma.A)'$  where  $(\Gamma.A)'(\rho, u)$  is the set of pairs  $\rho', u'$  with  $\rho' \in \Gamma'(\rho)$  and u' in  $A'\rho\rho'(u)$ .

We define an element  $\mathbf{p}^* = \mathbf{p}, \mathbf{p}'$  in  $(\Gamma, \Gamma').(A, A') \to \Gamma, \Gamma'$  by taking  $\mathbf{p}'(\rho, u)(\rho', u') = \rho'$ . We have then an element  $\mathbf{q}, \mathbf{q}'$  in  $\mathsf{Elem}^*((\Gamma, \Gamma').(A, A'), (A, A')\mathbf{p}^*)$  defined by  $\mathbf{q}'(\rho, u)(\rho', u') = u'$ .

#### 3.1 Dependent product

We define a new operation  $\Pi^*(A, A')(B, B') = \Pi A B, (\Pi A B)'$  where  $(\Pi A B)' \rho \rho'(w)$  is the set

$$\Pi(u \in |A\rho|)\Pi(u' \in A'\rho\rho'(u))B'(\rho, u)(\rho', u')(\mathsf{app}(w, u))$$

If b, b' is in  $\mathsf{Elem}^*((\Gamma, \Gamma').(A, A'), (B, B'))$  then  $\lambda^*(b, b') = \lambda b, (\lambda b)'$  where  $(\lambda b)'$  is defined by the equation

 $(\lambda b)'\rho\rho' uu' = b'(\rho,u)(\rho',u')$ 

which is in

$$B'(\rho, u)(\rho', u')(\mathsf{app}((\lambda b)\rho, u)) = B'(\rho, u)(\rho', u')(b(\rho, u))$$

We have an application operation  $\operatorname{app}^*((c, c'), (a, a')) = (\operatorname{app}(c, a), \operatorname{app}(c, a)')$  where  $\operatorname{app}(c, a)' \rho \rho' = c' \rho \rho'(a \rho)(a' \rho \rho')$ .

#### 3.2 Universes

We define  $U'_n(A)$  for A in  $|U_n|$  to be the set of functions  $|A| \to \mathcal{U}_n$ . Thus an element A' of  $U'_n(A)$  is a family of sets A'(u) in  $\mathcal{U}_n$  for u in |A|. The universe  $U^*_n$  of  $M^*$  is defined to be the pair  $U_n, U'_n$  and we have  $\mathsf{Elem}^*((\Gamma, \Gamma'), U^*_n) = \mathsf{Type}^*_n(\Gamma, \Gamma')$ .

#### 3.3 Booleans

We define  $N'_2(u)$  for u in  $|N_2|$  to be the set consisting of 0 if u = 0 and of 1 if u = 1. We have  $N'_2$  in  $U'_0(N_2)$ . Note that  $N'_2(u)$  may not be a subsingleton if we have 0 = 1 in the model. We define  $\operatorname{brec}(T, a_0, a_1)'\rho\rho' uu'$  to be  $a'_0\rho\rho'$  if u' = 0 and to be  $a'_1\rho\rho'$  if u' = 1.

#### 3.4 Main result

**Theorem 3.1.** The new collection of context, with the operations  $\rightarrow^*$ ,  $\mathsf{Type}_n^*$ ,  $\mathsf{Elem}^*$  and  $U_n^*$  and  $N_2^*$  define a new model of type theory.

The proof consists in checking that the required equalities hold for the operations we have defined. For instance, we have

$$\mathsf{app}^*(\lambda^*(b,b'),(a,a')) = (\mathsf{app}(\lambda b,a),\mathsf{app}(\lambda b,a)') = (b(1,a),\mathsf{app}(\lambda b,a)')$$

and

$$\mathsf{app}(\lambda b, a)' \rho \rho' = (\lambda b)' \rho \rho'(a\rho)(a'\rho\rho') = b'(\rho, a\rho)(\rho', a'\rho\rho')$$

and

$$(b(1,a))'\rho\rho' = b'(\rho,a\rho)(1'\rho\rho',a'\rho\rho') = b'(\rho,a\rho)(\rho',a'\rho\rho')$$

When checking the equalities, we only use  $\beta$ ,  $\eta$ -conversions at the metalevel.

There are of course strong similarities with the parametricity model presented in [4]. This model can also be seen as a constructive version of the *glueing* technique [14, 20]. Indeed, to give a family of sets over  $|\Gamma|$  is essentially the same as to give a set X and a map  $X \to |\Gamma|$ , which is what happens in the glueing technique [14, 20].

### 4 The term model

There is a canonical notion of morphism between two models. For instance, the first projection  $M^* \to M$  defines a map of models of type theory. As for models of generalized algebraic theories [10], there is an *initial* model unique up to isomorphism. We define the *term* model  $M_0$  of type theory to be this initial model. As for equational theories, this model can be presented by first-order terms (corresponding to each operations) modulo the equations/conversions that have to hold in any model.

**Theorem 4.1.** In the initial model given u in  $|N_2|$  we have u = 0 or u = 1. Furthermore we don't have 0 = 1 in the initial model.

*Proof.* We have a unique map of models  $M_0 \to M_0^*$ . The composition of the first projection with this map has to be the identity function on  $M_0$ . If u is in  $|N_2|$  the image of u by the initial map has hence to be a pair of the form u, u' with u' in  $N'_2(u)$ . It follows that we have u = 0 if u' = 0 and u = 1 if u' = 1. Since 0' = 0 and 1' = 1 we cannot have 0 = 1 in the initial model  $M_0$ .

## 5 Presheaf model

We suppose given an arbitrary model M. We define from this the following category  $\mathcal{C}$  of "telescopes". An object of  $\mathcal{C}$  is a list  $A_1, \ldots, A_n$  with  $A_1$  in Type(),  $A_2$  in Type( $A_1$ ),  $A_3$  in Type( $A_1.A_2$ ) ... To any such object X we can associate a context  $i(X) = A_1...A_n$  of the model M. If A is in Type(i(X)), we define the set Var(X, A) of numbers  $v_k$  such that  $qp^{n-k}$  is in Elem(i(X), A). We may write simply Elem(X, A) instead of Elem(i(X), A). Similarly we may write Type<sub>n</sub>(X) = Elem(X, U<sub>n</sub>) for Type<sub>n</sub>(i(X)). If  $v_k$  is in Var(X, A) we write  $[v_k] = qp^{n-k}$ . If  $Y = B_1, \ldots, B_m$  is an object of  $\mathcal{C}$ , a map  $\sigma : Y \to X$  is given by a list  $u_1, \ldots, u_n$  such that  $u_p$  is in Var(Y,  $A_p([u_1], \ldots, [u_{p-1}])$ ). We then define  $[\sigma] = ([u_1], \ldots, [u_p]) : i(Y) \to i(X)$ . It is direct to define a composition operation such that  $[\sigma\delta] = [\sigma][\delta]$  which gives a category structure on these objects.

We use freely that we can interpret the language of dependent types (with universes) in any presheaf category [13]. A presheaf F is given by a family of sets F(X) indexed by contexts with restriction maps  $F(X) \to F(Y)$ ,  $u \mapsto u\sigma$  if  $\sigma: Y \to X$ , satisfying the equations u1 = u and  $(u\sigma)\delta = u(\sigma\delta)$  if  $\delta: Z \to Y$ . A dependent presheaf G over F is a presheaf over the category of elements of F, so it is given by a family of sets  $G(X, \rho)$  for  $\rho$  in F(X) with restriction maps.

We write  $\mathcal{V}_0, \mathcal{V}_1, \ldots$  the cumulative sequence of presheaf universes, so that  $\mathcal{V}_n(X)$  is the set of  $\mathcal{U}_n$ -valued dependent presheaves on the presheaf represented by X.

 $\mathsf{Type}_n$  defines a presheaf over this category, with  $\mathsf{Type}_n$  subpresheaf of  $\mathsf{Type}_{n+1}$ . We can see  $\mathsf{Elem}$  as a dependent presheaf over  $\mathsf{Type}_n$  since it determines a collection of sets  $\mathsf{Elem}(X, A)$  for A in  $\mathsf{Type}_n(X)$  with restriction maps.

If A is in  $\mathsf{Type}_n(X)$  we let  $\mathsf{Norm}(X, A)$  (resp.  $\mathsf{Neut}(X, A)$ ) be the set of all expressions of type A that are in normal form (resp. neutral). As for Elem, we can see Neut and Norm as dependent types over  $\mathsf{Type}_n$ , and we have

$$\mathsf{Var}(A) \subseteq \mathsf{Neut}(A) \subseteq \mathsf{Norm}(A)$$

We have an evaluation function [e]: Elem(A) if e: Norm(A). If a is in Elem(A) then we let Norm(A)|a (resp. Neut(A)|a) be the subtypes of Norm(A) (resp. Neut(A)) of elements e such that [e] = a.

Each context  $\Gamma$  defines a presheaf  $|\Gamma|$  by letting  $|\Gamma|(X)$  be the set of all substitutions  $i(X) \to \Gamma$ .

Any element A of  $\mathsf{Type}_n(\Gamma)$  defines internally a function  $|\Gamma| \to \mathsf{Type}_n, \ \rho \mapsto A\rho$ .

We have a canonical isomorphism between  $Var(A) \to Type_n$  and  $Elem(A \to U_n)$ . We can then use this isomorphism to build an operation

$$\pi:\Pi(A:\mathsf{Type}_n)(\mathsf{Var}(A)\to\mathsf{Type}_n)\to\mathsf{Type}_n$$

such that  $(\Pi A B)\rho = \pi(A\rho)((\lambda x : Var(A\rho))B(\rho, [x])).$ 

We can also define, given A: Type<sub>n</sub> and F: Var $(A) \to$  Type<sub>n</sub> an operation  $\Lambda Af$ : Elem $(\pi AF)$ , for  $f: \Pi(x: Var(A))$ Elem(F x).

Similarly, we can define an operation

$$\pi: \Pi(A: \operatorname{Norm}(U_n))(\operatorname{Var}([A]) \to \operatorname{Norm}(U_n)) \to \operatorname{Norm}(U_n)$$

such that  $[\pi AF] = \pi[A](\lambda(x : \mathsf{Var}([A]))[F x])$  and given  $A : \mathsf{Norm}(U_n)$  and  $F : \mathsf{Var}([A]) \to \mathsf{Type}_n$  and  $f : \Pi(x : \mathsf{Var}([A]))\mathsf{Elem}(F x)$  an operation  $\Lambda Af : \mathsf{Norm}(\pi[A]F)$  such that  $[\Lambda Af] = \Lambda[A](\lambda(x : \mathsf{Var}([A]))[f x]))$ .

While equality might not be decidable in Var(A) (because we use arbitrary renaming as maps in the base category), the product operation is injective: if  $\pi AF = \pi BG$  in  $Norm(U_n)$  then A = B in  $Norm(U_n)$  and F = G in  $Var([A]) \rightarrow Type_n$ .

### 6 Normalization model

The model is similar to the reducibility model and we only explain the main operations.

As before, a context is a pair  $\Gamma, \Gamma'$  where  $\Gamma$  is a context of M and  $\Gamma'$  is a dependent family over  $|\Gamma|$ .

A type at level *n* over this context consists now of a pair  $A, \overline{A}$  where A is in  $\mathsf{Type}_n(\Gamma)$  and  $\overline{A}\rho\rho'$ in  $U'_n(A\rho)$  for  $\rho$  in  $|\Gamma|$  and  $\rho'$  in  $\Gamma'(\rho)$ . An element of  $U'_n(T)$  for T in  $\mathsf{Type}_n$  consists in a 4-uple  $T', T_0, \alpha, \beta$  where the element  $T_0$  is in  $\mathsf{Norm}(U_n)|T$ , the element T' is in  $\mathsf{Elem}(T) \to \mathcal{V}_n$ , the element  $\beta$  is in  $\Pi(k: \mathsf{Neut}(T))T'([k])$  and  $\alpha$  is in  $\Pi(u: \mathsf{Elem}(T))T'(u) \to \mathsf{Norm}(T)|u$ . An element of this type is a pair  $a, \overline{a}$  where a is in  $\mathsf{Elem}(\Gamma, A)$  and  $\overline{a}\rho\rho'$  is an element of  $T'(a\rho)$  where  $(T', T_0, \alpha, \beta) = \overline{A}\rho\rho'$ .

The intuition behind this definition is that it is a "proof-relevant" way to express the method of reducibility used for proving normalization [11]: a reducibility predicate has to contain all neutral terms and only normalizable terms. The function  $\alpha$  (resp.  $\beta$ ) is closely connected to the "reify" (resp. "reflect") function used in normalization by evaluation [5], but for a "glued" model.

We redefine  $N_2'(t)$  to be the set of elements u in Norm $(N_2)|t$  such that u is 0 or 1 or is neutral. We define  $\alpha_{N_2}t\nu = \nu$  and  $\beta_{N_2}(k) = k$ .

We define  $\alpha_{U_n} T(T', T_0, \alpha_T, \beta_T) = T_0$  and for K neutral  $\beta_{U_n}(K) = (K', K, \alpha, \beta)$  where K'(t) is  $\mathsf{Neut}([K])|t$  and  $\alpha tk = k$  and  $\beta(k) = k$ .

The set  $\mathsf{Type}_n^*(\Gamma, \Gamma')$  is defined to be the set of pairs  $A, \overline{A}$  where A is in  $\mathsf{Type}_n(\Gamma)$  and  $\overline{A}\rho\rho'$  is in  $U'_n(A\rho)$ .

The extension operation is defined by  $(\Gamma, \Gamma').(A, \overline{A}) = \Gamma.A, (\Gamma.A)'$  where  $(\Gamma.A)'(\rho, u)$  is the set of pairs  $\rho', \nu$  with  $\rho' \in \Gamma'(\rho)$  and  $\nu$  in  $\overline{A}\rho\rho'.1(u)$ .

We define a new operation  $\Pi^*(A, \overline{A})(B, \overline{B}) = C, \overline{C}$  where  $C = \Pi A B$  and  $\overline{C}\rho\rho'$  is the tuple

- $C'(w) = \Pi(a : \mathsf{Elem}(A\rho))\Pi(\nu : T'(u))F'u\nu(\mathsf{app}(w, u))$
- $\beta(k)u\nu = \beta_F u\nu(\operatorname{app}(k, \alpha_T u\nu))$
- $\alpha \ w \ \xi = \Lambda T_0 g$  with  $g(x) = \alpha_F[x] \beta_T(x) (\operatorname{app}(w, [x]))(\xi[x] \beta_T(x))$
- $C_0 = \pi T_0 G$  with  $G(x) = F_0[x]\beta_T(x)$

where we write  $(T', T_0, \alpha_T, \beta_T) = \overline{A}\rho\rho'$  in  $U'_n(A\rho)$  and for each u in  $\mathsf{Elem}(A\rho)$  and  $\nu$  in T'(u) we write  $(F'u\nu, F_0u\nu, \alpha_Fu\nu, \beta_Fu\nu) = \overline{B}(\rho, u)(\rho', \nu)$  in  $U'_n(B(\rho, u))$ . We can check  $[C_0] = (\Pi A B)\rho$  and we have  $C', C_0, \alpha, \beta$  is an element in  $U'_n((\Pi A B)\rho)$ .

We define  $\overline{U_n} = U_n, U_n', \alpha_{U_n}, \beta_{U_n}$  and  $\overline{N_2} = N_2, N_2', \alpha_{N_2}, \beta_{N_2}$ .

If we have T in  $\operatorname{Type}_n(\Gamma, N_2)$  and  $a_0$  in  $\operatorname{Elem}(T\langle 0 \rangle)$  and  $a_1$  in  $\operatorname{Elem}(T\langle 1 \rangle)$  and for each  $\rho : |\Gamma|$  and  $\rho' : \Gamma'(\rho)$  and u in  $\operatorname{Elem}(N_2)$  and  $\nu$  in  $N'_2(u)$  an  $\underbrace{\operatorname{element}(T'u\nu, T_0u\nu, \alpha_Tu\nu, \beta_Tu\nu)}_{T'u\nu}$  in  $U'_n(T(\rho, u))$  and  $\overline{a_0}$  in  $T'00(a_0)$  and  $\overline{a_1}$  in  $T'11a_1$  we define  $f = \overline{\operatorname{brec}(T, a_0, a_1)}\rho\rho'$  as follows. We take  $f \ u \ \nu = \overline{a_0}$  if  $\nu = 0$  and  $f \ u \ \nu = \overline{a_1}$  if  $\nu = 1$  and finally  $f \ u \ \nu = \beta_T u\nu(\operatorname{brec}(\Lambda(N_2, g), \alpha_T 00a_0\overline{a_0}, \alpha_T 11a_1\overline{a_1}))(\nu))$  where  $g(x) = T_0[x]\beta_{N_2}(x)$  if  $\nu$  is neutral.

We thus get, starting from an arbitrary model M, a new model M<sup>\*</sup> with a projection map M<sup>\*</sup>  $\rightarrow$  M. As for the canonicity model, if we start from the initial model M<sub>0</sub> we have an initial map M<sub>0</sub>  $\rightarrow$  M<sup>\*</sup><sub>0</sub> which is a section of the projection map. Hence for any *a* in Elem(*A*) we can compute  $\overline{a}$  in *A'*(*a*) where  $(A', A_0, \alpha_A, \beta_A) = \overline{A}$  and we have  $\alpha_A \ a \ \overline{a}$  in Norm(*A*)|*a*.

**Theorem 6.1.** Equality in  $M_0$  is decidable.

*Proof.* If a and b are of type A we can compute  $\overline{A} = (A', A_0, \alpha, \beta)$ . We then have a = b in Elem(A) if, and only if,  $\alpha a \overline{a} = \alpha b \overline{b}$  in Norm(A) since  $u = [\alpha u \overline{u}]$  for any u in Elem(A). The result then follows from the fact that the equality in Norm((), A) is decidable.

We also can prove that  $\Pi$  is one-to-one for conversions, following P. Hancock's argument presented in [16].

### 7 Conclusion

Our argument extends directly to the addition of dependent sum types with surjective pairing, or inductive types such as the type W A B [18].

The proof is very similar to the argument presented in [16], but it covers conversion under abstraction and  $\eta$ -conversion. Instead of set theory, one could formalize the argument in extensional type theory; presheaf models have been already represented elegantly in NuPrl [6]. As we noticed however, the meta theory only uses the form of extensionality ( $\eta$ -conversion) also used in the object theory, and we should be able to express the normalization proof as a program transformation from one type theory to another. The formulation of the presheaf model as a(n extension of) type theory will be similar to the way cubical type theory [8] expresses syntactically a presheaf model over a base category which is a Lawvere theory. This should amount essentially to work in a type theory with a double context, where substitutions for the first context are restricted to be renamings. We leave this as future work, which, if successful, would refute some arguments in [17] for not accepting  $\eta$ -conversion as definitional equality.

# Acknowledgement

This work started as a reading group of the paper [20] together with Simon Huber and Christian Sattler. The discussions we had were essential for this work; in particular Christian Sattler pointed out to me the reference [3]

# References

- A. Abel and G. Scherer. On Irrelevance and Algorithmic Equality in Predicative Type Theory. In Logical Methods in Computer Science, 8(1):1-36, 2012.
- [2] P. Aczel. On Relating Type Theories and Set Theories. Types for proofs and programs, 118, Lecture Notes in Comput. Sci., 1657, 1999.
- [3] Th. Altenkirch, M. Hofmann and Th. Streicher. Reduction-free normalisation for system F. Unpublished note, 1997.
- [4] J.-Ph. Bernardy, P. Jansson, R. Paterson. Parametricity and dependent types. ICFP 2010: 345-356.
- [5] U. Berger and H. Schwichtenberg. An inverse of the evaluation functional for typed lambda-calculus. Proceedings of LICS 1991.
- [6] M. Bickford. Formalizing Category Theory and Presheaf Models of Type Theory in Nuprl. Preprint, https://arxiv.org/abs/1806.06114, 2018.
- [7] Th. Coquand. An algorithm for testing conversion in type theory. In *Logical frameworks*, p. 255-279, Cambridge University Press, 1991.
- [8] C. Cohen, Th. Coquand, S. Hber, A. Mörtberg. Cubical type theory: a constructive interpretation of the univalence axiom. Proceeding of the Types Conference, 2015.
- [9] L. Crosilla and ML. Rathjen Inaccessible set axioms may have little consistency strength. Ann. Pure Appl. Log. 115, 3370 (2002).
- [10] P. Dybjer. Internal Type Theory. in Types for Programs and Proofs, Springer, 1996.
- [11] S. Fortune, D. Leivant, M. O'Donnell. The Expressiveness of Simple and Second-Order Type Structures. Journal of the ACM, Volume 30 Issue 1, p. 151-185, 1983.
- [12] K. Gödel. Über eine bisher noch nicht benützte Erweiterung des finiten Standpunktes. Dialectica, 12, pp. 280-287, 1958.
- [13] M. Hofmann. Syntax and semantics of dependent type theory. In Semantics of Logic of Computation, Cambridge University Press, 1997.
- [14] J. Lambek and P.J. Scott. Introduction to higher order categorical logic. Cambridge studies in advanced mathematics 7, 1986.
- [15] P. Martin-Löf. An intuitionistic theory of types. Preliminary version 1972; published in 25 Years of Type Theory, 1995.

- [16] P. Martin-Löf. An intuitionistic theory of types: predicative part. Logic Colloquium '73 (Bristol, 1973), pp. 73118.
- [17] P. Martin-Löf. About Models for Intuitionistic Type Theories and the Notion of Definitional Equality. Proceedings of the Third Scandinavian Logic Symposium, 1975, Pages 81-109.
- [18] P. Martin-Löf. Constructive mathematics and computer programming. Logic, methodology and philosophy of science, VI (Hannover, 1979), pp. 153–175, Stud. Logic Found. Math., 104, North-Holland, Amsterdam, 1982.
- [19] J.R. Shoenfield. Mathematical Logic. Addison-Wesley, 1967.
- [20] M. Shulman. Univalence for inverse diagrams and homotopy canonicity. Mathematical Structures in Computer Science, 25:05, p. 12031277, 2014.
- [21] W.W. Tait. Intensional interpretations of functionals of finite type, part I. Journal of Symbolic Logic, 32, pp. 198-212, 1967.